RIP-based Performance Guarantee for Low Rank Matrix Recovery via L_{*-F} Minimization

Yan Li and Liping Zhang

Abstract

In the undetermined linear system b = A(X) + s, vector b and operator A are the known measurements and s is the unknown noise. In this paper, we investigate sufficient conditions for exactly reconstructing desired matrix X being low-rank or approximately low-rank. We use the difference of nuclear norm and Frobenius norm (L_{*-F}) as a surrogate for rank function and establish a new nonconvex relaxation of such low rank matrix recovery, called the L_{*-F} minimization, in order to approximate the rank function closer. For such nonconvex and nonsmooth constrained L_{*-F} minimization problems, based on whether the noise level is 0, we give the upper bound estimation of the recovery error respectively. Particularly, in the noise-free case, one sufficient condition for exact recovery is presented. If linear operator A satisfies the restricted isometry property with $\delta_{4r} < \frac{\sqrt{2r-1}+\sqrt{2}(\sqrt{2r+1})}{\sqrt{2r-1}+\sqrt{2}(\sqrt{2r+1})}$, then r-rank matrix X can be exactly recovered without other assumptions. In addition, we also take insights into the regularized L_{*-F} minimization model since such regularized model is more widely used in algorithm design. We provide the recovery error estimation of this regularized L_{*-F} minimization model via RIP tool. To our knowledge, this is the first result on exact reconstruction of low rank matrix via regularized L_{*-F} minimization.

Index Terms

Low rank matrix recovery, nonconvex optimization, nuclear norm, restricted isometry property, Frobenius norm

I. INTRODUCTION

OW rank matrix recovery (LMR) has been a rapidly growing filed of research in machine learning [1][2][3] and computer vision [4][5]. Mathematically, we hope to acquire the low rank matrix X^o satisfying:

$$\boldsymbol{b} = \mathcal{A}(\boldsymbol{X}^o),\tag{1}$$

1

where **b** is a given nonzero vector and $\mathcal{A} : \mathbb{R}^{m \times n} \to \mathbb{R}^{l}$ is a predesigned measurement linear operator. Hence LMR can be formulated as follows:

$$\min_{\mathbf{X}} \operatorname{rank}(\mathbf{X})$$
s.t. $\mathcal{A}(\mathbf{X}) = \mathbf{b}.$
(2)

A particular class of (2) is to utilize a small number of observation entries to reconstruct matrix X, referred as low rank matrix completion, where $\mathcal{A} := \mathcal{P}_{\Omega}$ the projection operator samples entries from the index set Ω . Unfortunately, problem (2) is generally NP-hard and ill-posed [6], a famous convex surrogate function for rank function is the nuclear norm proposed by Fazel et al. [7] and they established a convex optimization problem over the same constraints:

$$\min_{\mathbf{X}} \|\mathbf{X}\|_{*}$$
s.t. $\mathcal{A}(\mathbf{X}) = \mathbf{b}$,
(3)

where $\|\cdot\|_*$ equals the summation of singular values. Note that the relaxation method is conceptually analogous to the relaxation from ℓ_0 norm to ℓ_1 norm in compressing sensing [8]. Many simple and computationally efficient optimization methods [9][10] exist to solve this type of nuclear norm minimization problem. Variants of nuclear norm, including the truncated nuclear norm [11] and weighted nuclear norm [12] were also proposed in the literature and enhance the recovery performance. Under some suitable conditions related to restricted isometry property (RIP), problem (3) can be guaranteed to produce the the minimum-rank solution [13].

In this study, we focus on a non-convex surrogate for rank function, i.e., the difference of nuclear norm and Frobenius norm (L_{*-F}) to solve problem (5). Invoking the definition of Frobenius norm $||\mathbf{X}||_F := \sqrt{\langle \mathbf{X}, \mathbf{X} \rangle}$, direct manipulations yield $||\mathbf{X}||_* - ||\mathbf{X}||_F = ||\sigma(\mathbf{X})||_1 - ||\sigma(\mathbf{X})||_2$. The contour plot of the $|| \cdot ||_1 - || \cdot ||_2$ ($\ell_1 - \ell_2$) metric presents in Figure 1, which implies it can achieve the goal of sparsity and hence we can build a new nonconvex relaxation of the low rank matrix model (2) as follows:

$$\min_{\mathbf{X}\in\mathbb{R}^{m\times n}} \|\mathbf{X}\|_* - \|\mathbf{X}\|_F \quad \text{s.t. } \mathcal{A}(\mathbf{X}) = \mathbf{b}.$$
(4)

This work was supported in part by the National Natural Science Foundation of China under Grant No. 12171271 (Corresponding author: Liping Zhang) Yan Li and Liping Zhang are with the Department of Mathematical Sciences, Tsinghua University, Beijing 100084, China (email:liyan20@mails.tsinghua.edu.cn; lipingzhang@tsinghua.edu.cn)



Fig. 1: From outside to inside, the value on the contour decreases gradually. It is notable that the solutions of $\ell_1 - \ell_2$ approach two axes, when the value is close to 0. Hence $\ell_1 - \ell_2$ norm gets more sparse solutions than ℓ_1 norm.

In practice, since the measurement b is possibly contaminated by unknown noise s, there produces a type of robustly recovering a low rank matrix in the form of

$$\boldsymbol{b} = \mathcal{A}(\boldsymbol{X}^o) + \boldsymbol{s}. \tag{5}$$

Under this case, we can formulate as the following models, one minimizing the same function with (4) executes the robust constraints to complete the low rank matrix recovery:

$$\min_{\boldsymbol{X} \in \mathbb{R}^{m \times n}} \|\boldsymbol{X}\|_* - \|\boldsymbol{X}\|_F \quad \text{s.t.} \ \|\mathcal{A}(\boldsymbol{X}) - \boldsymbol{b}\|_2 \le \epsilon,$$
(6)

where nonnegative parameter ϵ represents the noise level, and the other one is given by a sparsity regularized optimization problem:

$$\min_{\boldsymbol{X}\in\mathbb{R}^{m\times n}}\mathcal{J}(\boldsymbol{X}) := \|\boldsymbol{X}\|_* - \|\boldsymbol{X}\|_F + \frac{1}{2\lambda}\|\mathcal{A}(\boldsymbol{X}) - b\|_2^2,\tag{7}$$

where $\lambda > 0$ is a tradeoff hyperparameter. Both problems (6) and (7) belong to a special case of difference of convex function (DC) programming. For more details on DC programming and its algorithm implementation, see, e.g., [14][15]. In fact, no matter what function is chosen to replace the rank function, it is necessary to consider the recovery conditions and their resultant recovery error estimates. Subsequently, we will provide sufficient conditions to guarantee the robust reconstruction in bound of $\|\cdot\|_F$ or exact reconstruction of the desired low rank matrix X^o through above three minimization problems including noise setting ($\epsilon \neq 0$) and noiseless context ($\epsilon = 0$).

For this purpose, One of the most commonly used tools is the RIP condition. The matrix version of RIP notion as defined in Definition 1, which was first introduced by Candès and Tao [16], is widely used in sparse signal recovery [17][18][19].

Definition 1. For $T \subseteq \{1, \dots, n\}$ and each number r, r-restricted isometry constants of matrix A is the smallest quantity δ_r such that

$$(1 - \delta_r) \| \boldsymbol{x} \|_2^2 \le \| A_T \boldsymbol{x} \|_2^2 \le (1 + \delta_r) \| \boldsymbol{x} \|_2^2$$

for all subsets T with $|T| \leq r$ and all $x \in \mathbb{R}^{|T|}$. The matrix A is said to satisfy the r-RIP with δ_r .

Inspired by this, Candès and Plan [20] introduced the isometry constants of a linear map \mathcal{A} , as defined in Definition 2. The linear map \mathcal{A} is said to satisfy the RIP at rank r if δ_r is bounded by a sufficiently small constant between 0 and 1. As they mentioned, fix $0 \le \delta < 1$ and let \mathcal{A} be a random measurement ensemble obeying the following condition: for any given $\mathbf{X} \in \mathbb{R}^{n_1 \times n_2}$ and any fixed $0 < \hat{t} < 1$, $P(|||\mathcal{A}(\mathbf{X})||_2^2 - ||\mathbf{X}||_F^2| > \hat{t}||\mathbf{X}||_F^2) \le c_1 \exp(-c_2m)$ for fixed constants $c_1, c_2 > 0$. Then if $m \ge d_1nr$, \mathcal{A} satisfies the RIP with isometry constant $\delta_r \le \delta$ with probability exceeding $1 - c_1 e^{-d_2m}$ for fixed constants $n, d_1, d_2 > 0$. There is a rich literature providing a range of theoretical guarantees under which it is possible to recover a matrix based on the assumption that linear map \mathcal{A} satisfies certain RIP conditions. See, e.g., [21][22][23][24]. Many types of linear map, including random sensing designs [25][26], are known to satisfy the RIP with high probability.

Definition 2. For each integer $r = 1, 2, \dots, n$, the isometry constant δ_r of a linear map \mathcal{A} is the smallest quantity such that

$$(1 - \delta_r) \|\boldsymbol{X}\|_F^2 \le \|\mathcal{A}(\boldsymbol{X})\|_2^2 \le (1 + \delta_r) \|\boldsymbol{X}\|_F^2$$
(8)

holds for all r-rank matrices (any matrix of rank no greater than r).

A. Relation to Existing Work

In the present study, some sufficient conditions based on the RIP analysis to guarantee the recovery of desired matrices through the L_{1-2} metric have been provided. Cai [27] gave a stably recovery condition on imposing an additional assumption on the dimension of desired matrix X^o besides the RIP condition. Let X^o satisfy (5) with $||s||_2 \le \epsilon$, if there exists $r \le \min\{m, n\}$ so that $\alpha(r) := (\frac{\sqrt{2r-1}}{\sqrt{r+1}})^2 > 1$ and the linear map \mathcal{A} satisfies $\delta_{2r} + \alpha(r)\delta_{3r} < \alpha(r) - 1$, the error estimation deriving from problem (6) is bounded by $C_1 \| \mathbf{X}^o - \mathbf{X}_r^o \|_F + C_2 \epsilon$, where $C_1, C_2 > 0$ and \mathbf{X}_r^o is the best rank-r approximation matrix of X^{o} . Hence in the noise-free case ($\epsilon = 0$), X^{o} can be recovered exactly by the constrained optimization problem (4), if X^{o} is r-rank matrix. Ma et al. [28] proposed a truncated L_{1-2} metric and gave the theoretical guarantees to recovery the low rank matrix based on the corresponding constrained model. Note that L_{1-2} metric is a special case for the truncated L_{1-2} metric, hence following from [28] it yields that for the r-rank matrix X^o satisfying (5) with $||s||_2 \le \epsilon$, the error estimation deriving from problem (6) is bounded by $C_3\epsilon$, where $C_3 > 0$. Hence in the noise-free case ($\epsilon = 0$), X^o satisfying (1) can be recovered exactly by problem (4). Above observation suggests that they can not provide a robust error estimation when the information about range of $rank(X^{o})$ is missing. There also exist other forms of characterizations for isometry constant of a linear map by replacing the vector ℓ_2 norm with other vector norms, such as $\ell_p(0 quasi norm [29]. Under the$ framework of RIP with p = 1, Guo et al. [30] presented a recovery guarantee through the truncated L_{1-2} minimization and adopting ℓ_1 constraints, naturally, the recovery estimation of L_{1-2} minimization problem can be acquired. These works give the recovery estimation concerning L_{1-2} minimization approach with constraints. In general, problem (4) and problem (6) is not convenient to be solved in numerical implementation, the common choice is to solve a regularized variant (7) so that some algorithms for unconstrained minimization problem such as DCA [14] and PG [31] can be adopted to complete the recovery of the low rank matrix [27][28][32]. Hence it becomes very significant and necessary to develop some theoretical results for problem (7) and explore its relationship with constrained minimization problem. Moreover, since problem (4) is a surrogate optimization problem of NP hard problem (2), it is also necessary to establish the relation of optimal solutions among them. Motivated by above discussion, in this paper our contribution can be summarized as follows:

- We update the recovery theory based on the RIP conditions of linear map \mathcal{A} for constrained optimization problems (4) and (6) to broaden the range of recoverable low rank matrix. Although authors in [27] recently also give recovery theory based on the RIP conditions for these optimization problems, their recovery estimation is built for the desired matrix X^o satisfying $\min\{m, n\} \ge 24$ and ours breaks this restriction. Different from [28], the recovery theory we proposed is still valid when the range of **rank** (X^o) is unclear.
- To the best of our knowledge, we are the first to provide the upper bound estimation of the approximate error for the regularized L_{*-F} minimization problem (7). Actually, the existing theoretical investigation of problem (7) is limited to its induced algorithms and there is no theoretical guarantee of the regularized recovery estimation. We fill the blink of theoretical investigation to characterize its essential performance in robustly recovering the desired matrix X^o from (5).
- The sufficient condition is provided to demonstrate that it is possible to recover the lowest rank solution exactly by minimizing the difference between nuclear norm and Frobenius norm. Moreover, we also briefly discuss the relation between the global minimizers of problems (7) and (4).

B. Notation

In this subsection, we introduce some related notations used throughout this paper. We present vectors by boldface lowcase letters, e.g., a, matrices by boldface capital letters, e.g., A, sets by capital letters, e.g., A, scalars by lowercase letters, e.g., a. For any positive integer d, [d] denotes the index set $\{1, 2, \dots, d\}$. Given $X \in \mathbb{R}^{m \times n}$. Define $t := \min\{m, n\}$. Let $\sigma(X) := (\sigma_1(X), \dots, \sigma_t(X))$ be a vector composed of X's singular values with $\sigma_1(X) \ge \dots \ge \sigma_t(X) \ge 0$. $E \in \mathbb{R}^{t \times t}$ denotes the identity matrix. For an index set $S \subseteq [d]$, let |S| denote its cardinality and S^c denote its complementarity set. Denote X_i as the *i*-th column of X. Denote X_S as X with all but columns indexed by S set to zero vector. For any vector $h \in \mathbb{R}^n$ and any index subset $S \subseteq [n]$, we denote by h_S the vector whose entries $(h_S)_i = h_i$ for $i \in S$ and 0 otherwise. Besides, we denote by $h_{\max(k)}$ the vector h with all but the largest k entries in absolute values set to 0. The inner products of two vectors $x, y \in \mathbb{R}^n$ and two matrices $X, Y \in \mathbb{R}^{m \times n}$ are denoted by $\langle x, y \rangle := x^\top y$ and $\langle X, Y \rangle := \text{Tr}(X^\top Y)$ where Tr is the matrix trace. For any $x \in \mathbb{R}^t$, we define the operator $\mathcal{D} : \mathbb{R}^t \to \mathbb{R}^{m \times n}$ as follows:

$$\mathcal{D}_{ij}(\boldsymbol{x}) = \begin{cases} x_i & \text{if } i = j, \\ 0 & \text{otherwise.} \end{cases}$$

Let $\mathcal{O}(n)$ be the group of $n \times n$ orthogonal matrices. For any given $X \in \mathbb{R}^{m \times n}$, its singular value decomposition (SVD) is $X = U\mathcal{D}(\sigma(X))V^{\top}$ with $U \in \mathcal{O}(m)$ and $V \in \mathcal{O}(n)$. Denote $\lfloor \cdot \rfloor$ and $\lceil \cdot \rceil$ as the symbols for floor function and ceiling function respectively. Let $\operatorname{supp}(x)$ denote the support of x. We say two matrices X and Y in $\mathbb{R}^{m \times n}$ have simultaneous ordered SVD if there exist $U_X, U_Y \in \mathcal{O}^m$ and $V_X, V_Y \in \mathcal{O}^n$ such that $X = U_X \mathcal{D}(\sigma(X))V_X^{\top}$ and $Y = U_Y \mathcal{D}(\sigma(Y))V_Y^{\top}$ with $\{U_{X_1}, U_{X_2}, \cdots, U_{X_m}\} = \{U_{Y_1}, U_{Y_2}, \cdots, U_{Y_m}\}$ and $\{V_{X_1}, V_{X_2}, \cdots, V_{X_n}\} = \{V_{Y_1}, V_{Y_2}, \cdots, V_{Y_n}\}$. That implies matrix Y can be rewritten as $U_X \mathcal{D}(\pi(\sigma(Y)))V_X^{\top}$, where $\pi_1(\sigma(Y)), \pi_2(\sigma(Y)), \cdots, \pi_t(\sigma(Y))$ are some permutation of singular values of Y.

In this section, we first show that it is possible to recover the lowest rank representation by solving a nonconvex optimization problem. And then, we present the recovery performance of L_{1-2} minimization model with constraints under the framework of RIP.

A. Links Between Problem (2) and Problem (4)

The goal of this subsection is to study the links between the rank minimization problem (2) and its nonconvex surrogate minimization (4). In light of the characterization of locally sparse feasible solutions, we show the globally optimal solution of problem (2) must solve the problem (4) globally.

Definition 3. Denote $\mathcal{F} := \{ X \mid \mathcal{A}(X) = b \}$. $X \in \mathcal{F}$ is called locally sparse if $\nexists Y \in \mathcal{F} \setminus \{ X \}$ such that Y and X have a simultaneous ordered SVD and $supp(\pi(\sigma(Y))) \subseteq supp(\sigma(X))$. Denote $\mathcal{F}_L = \{ X \in \mathcal{F} \mid X \text{ is locally sparse} \}$ as the set of locally sparse feasible solutions.

In fact, the locally sparse feasible solution is locally the sparsest feasible solution.

Lemma 4. For any $X \in \mathcal{F}_L$, there exists $\delta_X > 0$ such that for any $Y \in \mathcal{F}$ having simultaneous ordered SVD with X, if $0 < ||Y - X||_F < \delta_X$, we have $\operatorname{supp}(\sigma(X)) \subset \operatorname{supp}(\pi(\sigma(Y)))$.

Proof. Choose $\delta_X = \min_{i \in \text{supp}(\sigma(X))} \{\sigma_i(X)\}$. For any $Y \in \mathcal{F}$ having simultaneous ordered SVD with X such that $0 < \|Y - X\|_F < \delta_X$, we set Z = X - Y, that is $Z = U_X \mathcal{D}(\sigma(X) - \pi(\sigma(Y))) V_X^{\mathsf{T}}$. For brevity, denote z as $\sigma(X) - \pi(\sigma(Y))$, then we get

$$\|\boldsymbol{z}\|_{\infty} \leq \|\boldsymbol{z}\|_2 < \min_{i \in \operatorname{supp}(\sigma(\boldsymbol{X}))} \{\sigma_i(\boldsymbol{X})\}$$

This yields

$$egin{aligned} & m{\sigma}_i(\sigma(m{Y})) \geq \sigma_i(m{X}) - \|m{z}\|_\infty \ & > \sigma_i(m{X}) - \min_{i \in ext{supp}(\sigma(m{X}))} \{\sigma_i(m{X})\} \geq 0 \end{aligned}$$

for any $i \in \text{supp}(\sigma(X))$, which implies $\text{supp}(\sigma(X)) \subseteq \text{supp}(\pi(\sigma(Y)))$. Moreover, since $X \in \mathcal{F}_L$, we have $\text{supp}(\sigma(X)) \neq \text{supp}(\pi(\sigma(Y)))$. Hence we obtain that $\text{supp}(\sigma(X)) \subset \text{supp}(\pi(\sigma(Y)))$. This completes the proof.

The following results show that the optimal solution sets of problem (4) and problem (2) are contained in the locally sparse sets.

Lemma 5. If X^* solves problem (4) globally, then $X^* \in \mathcal{F}_L$.

Proof. If $X^* \notin \mathcal{F}_L$ and $X^* = U_{X^*}\mathcal{D}(\sigma(X^*))V_{X^*}^{\mathsf{T}}$, then there exists $Y^* \in \mathcal{F} \setminus \{X^*\}$ such that $Y^* = U_{X^*}\mathcal{D}(\pi(\sigma(Y^*)))V_{X^*}^{\mathsf{T}}$ and $\sup(\pi(\sigma(Y))) \subseteq \operatorname{supp}(\sigma(X^*))$. Hence we can find a small enough $\epsilon > 0$ such that

$$\sigma(\boldsymbol{X}^*) - \epsilon \boldsymbol{\pi}(\sigma(\boldsymbol{Y}^*)) \ge 0$$

Define $Z^* := U_{X^*} \mathcal{D}(\frac{\sigma(X^*) - \epsilon \pi(\sigma(Y^*))}{1 - \epsilon}) V_{X^*}^{\intercal}$. By directly calculating, it yields

|| \$7.7 * ||

11 37

$$\mathcal{A}(\mathbf{Z}^*) = \frac{1}{1-\epsilon} \mathcal{A}(\mathbf{X}^*) - \frac{\epsilon}{1-\epsilon} \mathcal{A}(\mathbf{Y}^*) = \mathbf{b}$$

and $\|Z^*\|_* = \frac{1}{1-\epsilon} \|X^*\|_* - \frac{\epsilon}{1-\epsilon} \|Y^*\|_*$ due to the non-negativity of $\sigma_i(X^*) - \epsilon \pi_i(\sigma(Y^*))$. Moreover, it follows from $Y^* \neq X^*$ and $\mathcal{A}(Y^*) = \mathcal{A}(X^*)$ that they are linearly independent, and hence Y^* and Z^* are linearly independent. This implies

$$\|X^*\|_F < \epsilon \|Y^*\|_F + (1-\epsilon) \|Z^*\|_F$$

Therefore, it is obvious that

$$\|\boldsymbol{X}^{*}\|_{*} - \|\boldsymbol{X}\|_{F}$$

> $\epsilon(\|\boldsymbol{Y}^{*}\|_{*} - \|\boldsymbol{Y}^{*}\|_{F}) + (1 - \epsilon)(\|\boldsymbol{Z}^{*}\|_{*} - \|\boldsymbol{Z}^{*}\|_{F})$
 $\geq \min\{\|\boldsymbol{Y}^{*}\|_{*} - \|\boldsymbol{Y}^{*}\|_{F}, \|\boldsymbol{Z}^{*}\|_{*} - \|\boldsymbol{Z}^{*}\|_{F}\},$

which contradicts with the optimality of X^* .

Lemma 6. If X^* solves problem (2) globally, then $X^* \in \mathcal{F}_L$.

Proof. If $X^* \notin \mathcal{F}_L$ and $X^* = U_{X^*}\mathcal{D}(\sigma(X^*))V_{X^*}^{\mathsf{T}}$, then there exists $Y^* \in \mathcal{F} \setminus \{X^*\}$ such that $Y^* = U_{X^*}\mathcal{D}(\pi(\sigma(Y^*)))V_{X^*}^{\mathsf{T}}$ and $\sup(\pi(\sigma(Y^*))) \subseteq \sup(\sigma(X^*))$. Since X^* is optimal, $\sup(\pi(\sigma(Y^*))) = \sup(\sigma(X^*))$ and we denote such support set as S.

According to $\mathbf{X}^* \neq \mathbf{Y}^*$, it yields $\sigma(\mathbf{X}^*) \neq \pi(\sigma(\mathbf{Y}^*))$ and hence $\min_{i \in S} \{\frac{\sigma_i(\mathbf{X}^*)}{\pi_i(\sigma(\mathbf{Y}^*))}\} < 1$ or $\min_{i \in S} \{\frac{\pi_i(\sigma(\mathbf{Y}^*))}{\sigma_i(\mathbf{X}^*)}\} < 1$ must true. Without loss of generality, let $\min_{i \in S} \{\frac{\sigma_i(\mathbf{X}^*)}{\pi_i(\sigma(\mathbf{Y}^*))}\} = \frac{\sigma_k(\mathbf{X}^*)}{\pi_k(\sigma(\mathbf{Y}^*))} = r < 1$ for some $k \in S$. Then

$$\boldsymbol{Z}^* := \frac{1}{1-r} \boldsymbol{X}^* - \frac{r}{1-r} \boldsymbol{Y}^* = \boldsymbol{U}_{X^*} \mathcal{D}(\frac{\sigma(\boldsymbol{X}^*) - r\boldsymbol{\pi}(\sigma(\boldsymbol{Y}^*))}{1-r}) \boldsymbol{V}_{X^*}^{\mathsf{T}},$$

which implies $\mathcal{A}(\mathbf{Z}^*) = \mathbf{b}$, that is, $\mathbf{Z}^* \in \mathcal{F}$. Moreover, denote \mathbf{z} as $\frac{\sigma(\mathbf{X}^*) - r\pi(\sigma(\mathbf{Y}^*))}{1-r}$, then $\mathbf{z}_k = 0$ indicates $\operatorname{supp}(\sigma(\mathbf{Z}^*)) \underset{\neq}{\subseteq} \operatorname{supp}(\sigma(\mathbf{X}^*))$, which contradicts with \mathbf{X}^* being optimal solution of problem (2).

Now, we are in the position to present one of our main recovery result.

Theorem 7. If X^* uniquely solves problem (2) with $\operatorname{rank}(X^*) = s$ and $\min_{i \in \operatorname{supp}(\sigma(X))} \sigma_i(X) > \frac{2(\sqrt{s}-1)}{\operatorname{rank}(X)-1} \|X^*\|_F$ for any $X \in \mathcal{F}_L \setminus \{X^*\}$, then X^* also uniquely solves problem (4).

Proof. First, we will show that for $X \in \mathbb{R}^{m \times n}$, $(\operatorname{rank}(X) - 1)/2 \min_{i \in \operatorname{supp}(\sigma(X))} \sigma_i(X) \leq ||X||_* - ||X||_F \leq (\sqrt{\operatorname{rank}(X)} - 1)||X||_F$. This upper bound can be immediately obtained from the Cauchy-Schwartz inequality. Next, we will give the lower bound.

By directly calculating, it yields

$$\begin{split} \|\mathbf{X}\|_{*} - \|\mathbf{X}\|_{F} &= \frac{\|\mathbf{X}\|_{*}^{2} - \|\mathbf{X}\|_{F}^{2}}{\|\mathbf{X}\|_{*} + \|\mathbf{X}\|_{F}} \\ &= \frac{\sum_{i \neq j} \sigma_{i}(\mathbf{X})\sigma_{j}(\mathbf{X})}{\|\mathbf{X}\|_{*} + \|\mathbf{X}\|_{F}} \\ &= \frac{\sum_{i \neq j \in \text{supp}(\sigma(\mathbf{X}))} \sigma_{i}(\mathbf{X})\sigma_{j}(\mathbf{X})}{\|\mathbf{X}\|_{*} + \|\mathbf{X}\|_{F}} \\ &\geq \frac{(\text{rank}(\mathbf{X}) - 1)\|\mathbf{X}\|_{*}\min_{i \in \text{supp}(\sigma(\mathbf{X}))} \sigma_{i}(\mathbf{X})}{\|\mathbf{X}\|_{*} + \|\mathbf{X}\|_{F}} \\ &= \frac{\text{rank}(\mathbf{X}) - 1}{1 + \frac{\|\mathbf{X}\|_{F}}{\|\mathbf{X}\|_{*}}} \min_{i \in \text{supp}(\sigma(\mathbf{X}))} \sigma_{i}(\mathbf{X}) \\ &\geq \frac{\text{rank}(\mathbf{X}) - 1}{2} \min_{i \in \text{supp}(\sigma(\mathbf{X}))} \sigma_{i}(\mathbf{X}). \end{split}$$

Considering above discussion, we can directly obtain

$$egin{aligned} \|m{X}^*\|_* &- \|m{X}^*\|_F \leq (\sqrt{s}-1)\|m{X}^*\|_F \ &< rac{ ext{rank}(m{X})-1}{2}\min_{i\in ext{supp}(\sigma(m{X}))}\sigma_i(m{X}) \ &\leq \|m{X}\|_* - \|m{X}\|_F \end{aligned}$$

for any $X \in \mathcal{F}_L \setminus \{X^*\}$, which makes sense according to Lemma 6. Thus, the desired result follows directly from Lemma 5. \Box

B. Exact Recovery Theory

In this subsection, we obtain some theoretical results to guarantee the robust recovery through the constrained L_{1-2} minimization problem (4) and problem (6). Our main results not only provide the sufficient conditions of stably recovering the desired matrix X^o , but also characterize the recovery errors with these two approaches. Before proceeding, we provide essential preliminaries and related facts which are helpful to derive stable recovery conditions. We begin with the following fundamental properties respect to the function $\|X\|_* - \|X\|_F$.

Lemma 8. Suppose $X \in \mathbb{R}^{m \times n} \setminus \{0\}$ and $\operatorname{rank}(X) = r$. Let $U\mathcal{D}(\sigma(X))V^{\top}$ be the SVD of X, that is, $X = \sum_{i=1}^{t} \sigma_i(X)U_iV_i^{\top}$. Denote $\Lambda := \operatorname{supp}(\sigma(X))$. Then, we have (a) $(t - \sqrt{t})\sigma_t(X) \le \|X\|_* - \|X\|_F \le (\sqrt{t} - 1)\|X\|_F$; (b) $(r - \sqrt{r})\sigma_r(X) \le \|X\|_* - \|X\|_F \le (\sqrt{r} - 1)\|X\|_F$; (c) $\|X\|_* - \|X\|_F = 0$ if and only if r = 1.

Proof. (a) We can easily get the supper bound from the Cauchy-Schwartz inequality, and it suffices to give the lower bound. Define $r_0 = \lfloor \sqrt{t} \rfloor$. We will give

$$\|\boldsymbol{X}\|_{F} = \sqrt{\sum_{i} \sigma_{i}(\boldsymbol{X})^{2}}$$

$$\leq \sum_{i=1}^{r_{0}} \sigma_{i}(\boldsymbol{X}) + (\sqrt{t} - r_{0})\sigma_{r_{0}+1}(\boldsymbol{X}).$$
(9)

To achieve this, let the left side of (9) be ρ_l and the right side be ρ_r . On the one hand, we have

$$\rho_l^2 = \sum_{i=1}^{r_0} \sigma_i(\mathbf{X})^2 + \sum_{i=r_0+1}^t \sigma_i(\mathbf{X})^2$$

$$\leq \sum_{i=1}^{r_0} \sigma_i(\mathbf{X})^2 + (t-r_0)\sigma_{r_0+1}(\mathbf{X})^2$$

and on the other hand by directly calculating, we obtain

$$\begin{split} \rho_r^2 &= \sum_{i=1}^{r_0} \sigma_i(\boldsymbol{X})^2 + 2(\sqrt{t} - r_0)\sigma_{r_0+1}(\boldsymbol{X}) \sum_{i=1}^{r_0} \sigma_i(\boldsymbol{X}) + \sum_{i=1}^{r_0} \sum_{\substack{j=1\\j \neq i}}^{r_0} \sigma_i(\boldsymbol{X})\sigma_j(\boldsymbol{X}) + (\sqrt{t} - r_0)^2\sigma_{r_0+1}(\boldsymbol{X})^2 \\ &\geq \sum_{i=1}^{r_0} \sigma_i(\boldsymbol{X})^2 + 2(\sqrt{t} - r_0)\sigma_{r_0+1}(\boldsymbol{X}) \sum_{i=1}^{r_0} \sigma_{r_0+1}(\boldsymbol{X}) + \sum_{i=1}^{r_0} \sum_{\substack{j=1\\j \neq i}}^{r_0} \sigma_{r_0+1}(\boldsymbol{X})\sigma_{r_0+1}(\boldsymbol{X}) + (\sqrt{t} - r_0)^2\sigma_{r_0+1}(\boldsymbol{X})^2 \\ &= \sum_{i=1}^{r_0} \sigma_i(\boldsymbol{X})^2 + (t - r_0)\sigma_{r_0+1}(\boldsymbol{X})^2, \end{split}$$

and hence $\rho_l \leq \rho_r$, which implies (9). Then it yields that

$$\|\boldsymbol{X}\|_{*} - \|\boldsymbol{X}\|_{F} \ge \|\boldsymbol{X}\|_{*} - \sum_{i=1}^{r_{0}} \sigma_{i}(\boldsymbol{X}) - (\sqrt{t} - r_{0})\sigma_{r_{0}+1}(\boldsymbol{X})$$
$$\ge \sum_{i=r_{0}+1}^{t} \sigma_{i}(\boldsymbol{X}) - (\sqrt{t} - r_{0})\sigma_{r_{0}+1}(\boldsymbol{X})$$
$$\ge (t - \sqrt{t})\sigma_{t}(\boldsymbol{X}).$$

This yields the desired results.

(b) Note that $|\Lambda| = r$. Define $\hat{X} := \sum_{i \in \Lambda} \sigma_i(X) U_i V_i^{\top}$. Obviously,

$$\|m{X}\|_* - \|m{X}\|_F = \|\hat{m{X}}\|_* - \|\hat{m{X}}\|_F,$$

hence we can apply (a) to get the desired results.

(c) If $\|X\|_* - \|X\|_F = 0$, we can obtain $(r - \sqrt{r})\sigma_r(X) = 0$ by employing the relation (a) and hence r = 1. The other direction is easy.

By a simple application of the parallelogram identity, we then have the next lemma, whose proof can be found in [33].

Lemma 9. For all X, X' obeying $\langle X, X' \rangle = 0$, and $\operatorname{rank}(X) \leq r$, $\operatorname{rank}(X') \leq r'$, we have

$$|\langle \mathcal{A}(\boldsymbol{X}), \mathcal{A}(\boldsymbol{X}') \rangle| \leq \delta_{r+r'} \|\boldsymbol{X}\|_F \|\boldsymbol{X}'\|_F,$$

where $\delta_{r+r'}$ is (r+r')-isometry constant defined by Definition 2.

To show the main results, the following lemma is also necessary.

Lemma 10. Fix positive integer r and \hat{t} . Let m_1, m_2, n_1, n_2 be nonnegative integers which satisfy $m_1 + m_2 = n_1 + n_2 = r$. Then for $k \in [\hat{t}]$, it holds

$$\min_{m_1, m_2, n_1, n_2} \max\{m_1 + n_1 + k, m_2 + n_2 + 2k\} = \max\{r + \lceil \frac{3}{2}k \rceil, 2k\}.$$
(10)

Proof. Obviously, it holds

$$\max\{m_1 + n_1 + k, m_2 + n_2 + 2k\} \ge r + \lceil \frac{3}{2}k \rceil$$
(11)

for any $k \in [\hat{t}]$ since $m_1 + n_1 + k + n_1 + n_2 + 2k = 2r + 3k$.

For $k \ge 2r$. Following from $m_1 + n_1 \le 2r$, it is known that $\max\{m_1 + n_1 + k, m_2 + n_2 + 2k\} = m_2 + n_2 + 2k$ and then we have $\min_{m_1,m_2,n_1,n_2} \max\{m_1 + n_1 + k, m_2 + n_2 + 2k\} = 2k$ by setting $m_2 = n_2 = 0$. For k < 2r. It is known that $r + \lceil \frac{3}{2}k \rceil \ge r + \frac{3}{2}k > 2k$. In fact, we can show $\max\{m_1 + n_1 + k, m_2 + n_2 + 2k\} = r + \lceil \frac{3}{2}k \rceil$

by setting an appropriate m_1, m_2, n_1, n_2 . If k is odd, we set

$$m_{1} = \lceil \frac{r}{2} + \frac{k}{4} + \frac{1}{4} \rceil;$$

$$m_{2} = r - \lceil \frac{r}{2} + \frac{k}{4} + \frac{1}{4} \rceil;$$

$$n_{1} = r + \frac{k}{2} + \frac{1}{2} - \lceil \frac{r}{2} + \frac{k}{4} + \frac{1}{4} \rceil;$$

$$n_{2} = \lceil \frac{r}{2} + \frac{k}{4} + \frac{1}{4} \rceil - \frac{k}{2} - \frac{1}{2}.$$

If k is even, we set

$$m_1 = \lceil \frac{r}{2} + \frac{k}{4} \rceil;$$

$$m_2 = r - \lceil \frac{r}{2} + \frac{k}{4} \rceil;$$

$$n_1 = r + \frac{1}{2}k - \lceil \frac{r}{2} + \frac{k}{4} \rceil;$$

$$n_2 = \lceil \frac{r}{2} + \frac{k}{4} \rceil - \frac{k}{2}.$$

This indicates that

$$\min_{m_1, m_2, n_1, n_2} \max\{m_1 + n_1 + k, m_2 + n_2 + 2k\} = r + \lceil \frac{3}{2}k \rceil.$$

Combining the above two cases, we obtain the desired results.

In view of above lemmas, we are ready to give our main results for recovery analysis via constrained L_{1-2} minimization problem. We first present the recovery estimation in the noise-free case, i.e., the vector s = 0 in (5). The proof is based on the block decomposition of a matrix and the technique result of Lemma 10. Denote $U\mathcal{D}(\sigma(\mathbf{X}^o))\mathbf{V}^{\top}$ as SVD of \mathbf{X}^o . For any fixed positive integer $r \leq t$, the best rank-r approximation matrix \mathbf{X}_r^o of \mathbf{X}^o is defined as $U\mathcal{D}(\mathbf{x}^r)\mathbf{V}^{\top}$ where $\mathbf{x}^r \in \mathbb{R}^t$ and $x_i^r = \sigma_i(\mathbf{X}^o)$ for $i \in [r]$ and $x_i^r = 0$ otherwise.

Theorem 11. Let $\mathcal{A} : \mathbb{R}^{m \times n} \to \mathbb{R}^l$ be a linear map and $\boldsymbol{b} \in \mathbb{R}^l$. Set the desired matrix \boldsymbol{X}^o with $\mathcal{A}(\boldsymbol{X}^o) = \boldsymbol{b}$. For a given positive integer $r \leq t$, define $\hat{t} = \min\{m-r, n-r\}$. If there exists positive integer $k \in [\hat{t}] \setminus \{1\}$ such that linear map \mathcal{A} obeys $\delta_{2r+k} < 1$ and

$$\beta := \frac{\sqrt{2}\delta_{\max\{r+\lceil \frac{3}{2}k\rceil, 2k\}}(\sqrt{2r}+1)}{(1-\delta_{2r+k})(\sqrt{k}-1)} < 1,$$
(12)

2

then we obtain that any optimal solution \hat{X}^* of problem (4) satisfies

$$\|\boldsymbol{X}^{o} - \boldsymbol{X}^{*}\|_{F} \le \alpha \|\boldsymbol{X}^{o} - X_{r}^{o}\|_{F}$$

where $\alpha := \frac{2(\sqrt{2r}+1)+2\beta(\sqrt{k}-1)}{(\sqrt{k}-1)(1-\beta)(\sqrt{2r}+1)}$. Moreover, if \mathbf{X}^o is r-rank matrix, then the unique minimizer of problem (4) is exactly \mathbf{X}^o .

Proof. Define $Z := \hat{X}^* - X^o$ and $X_{r_c}^o := U\mathcal{D}(x^{r_c})V^{\top}$ where $x_i^{r_c} = \sigma_i(X^o)$ for $i \in [t] \setminus [r]$ and $x_i^{r_c} = 0$ otherwise. Obviously, $X^o = X_r^o + X_{r_c}^o$. Fix any nonnegative integer m_i, n_i $(i \in [3])$ satisfying $m_1 + m_2 = n_1 + n_2 = r$, $m_1 + m_2 + m_3 = m$ and $n_1 + n_2 + n_3 = n$. Denote

$$r_1 := m_1 + n_1 + k, \quad r_2 := m_2 + n_2 + 2k, \quad r_3 = \max\{r_1, r_2\}.$$

In the sequel, we get a block decomposition of Z with respect to X^{o} as follows:

$$\boldsymbol{U}^{\top} \boldsymbol{Z} \boldsymbol{V} := \begin{pmatrix} \boldsymbol{Z}_{11} & \boldsymbol{Z}_{12} & \boldsymbol{Z}_{13} \\ \boldsymbol{Z}_{21} & \boldsymbol{Z}_{22} & \boldsymbol{Z}_{23} \\ \boldsymbol{Z}_{31} & \boldsymbol{Z}_{32} & \boldsymbol{Z}_{33} \end{pmatrix}$$
(13)

with $Z_{ij} \in \mathbb{R}^{m_j \times n_j}$ $(i, j \in [3])$. Define

$$\boldsymbol{Z}_1 := \boldsymbol{U} \begin{pmatrix} \boldsymbol{Z}_{11} & \boldsymbol{Z}_{12} & \boldsymbol{Z}_{13} \\ \boldsymbol{Z}_{21} & 0 & 0 \\ \boldsymbol{Z}_{31} & 0 & 0 \end{pmatrix} \boldsymbol{V}^{\top}, \quad \boldsymbol{Z}_2 := \boldsymbol{U} \begin{pmatrix} 0 & 0 & 0 \\ 0 & \boldsymbol{Z}_{22} & \boldsymbol{Z}_{23} \\ 0 & \boldsymbol{Z}_{32} & 0 \end{pmatrix} \boldsymbol{V}^{\top}, \quad \boldsymbol{Z}_3 := \boldsymbol{U} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & \boldsymbol{Z}_{33} \end{pmatrix} \boldsymbol{V}^{\top}.$$

Naturally, we can decompose Z as

$$Z = Z_1 + Z_2 + Z_3 = Z^r + Z^{r_c}, (14)$$

where $Z^r := Z_1 + Z_2$, $Z^{r_c} := Z_3$. It is known that $\operatorname{rank}(Z_1) \le m_1 + n_1$, $\operatorname{rank}(Z_2) \le m_2 + n_2$ and $\operatorname{rank}(Z_1 + Z_2) \le 2r$. Also, Z_1 , Z_2 and Z_3 are orthogonal each other. Then, we have

$$\begin{split} \| \boldsymbol{X}^{o} + \boldsymbol{Z} \|_{*} - \| \boldsymbol{X}^{o} + \boldsymbol{Z} \|_{F} \\ = \| \boldsymbol{X}_{r}^{o} + \boldsymbol{X}_{r_{c}}^{o} + \boldsymbol{Z}^{r} + \boldsymbol{Z}^{r_{c}} \|_{*} - \| \boldsymbol{X}^{o} + \boldsymbol{Z}^{r} + \boldsymbol{Z}^{r_{c}} \|_{F} \\ \geq \| \boldsymbol{X}_{r}^{o} + \boldsymbol{Z}^{r_{c}} \|_{*} - \| \boldsymbol{X}_{r_{c}}^{o} \|_{*} - \| \boldsymbol{Z}^{r} \|_{*} - \| \boldsymbol{X}^{o} \|_{F} - \| \boldsymbol{Z}^{r} \|_{F} - \| \boldsymbol{Z}^{r_{c}} \|_{F} \\ = \| \boldsymbol{X}_{r}^{o} \|_{*} + \| \boldsymbol{Z}^{r_{c}} \|_{*} - \| \boldsymbol{X}_{r_{c}}^{o} \|_{*} - \| \boldsymbol{Z}^{r} \|_{*} - \| \boldsymbol{X}^{o} \|_{F} - \| \boldsymbol{Z}^{r} \|_{F} - \| \boldsymbol{Z}^{r_{c}} \|_{F}. \end{split}$$

Here the last equality follows from [13, Lemma 2.3] with $X_r^o Z^{r_c \top} = 0$ and $X_r^o^{\top} Z^{r_c} = 0$. Besides, we can get

$$\|\hat{X}^*\|_* - \|\hat{X}^*\|_F \le \|X^o\|_* - \|X^o\|_F$$

due to the optimality of \hat{X}^* . Hence we obtain

$$\|\boldsymbol{Z}^{r_c}\|_* - \|\boldsymbol{Z}^{r_c}\|_F \le \|\boldsymbol{Z}^r\|_* + \|\boldsymbol{Z}^r\|_F + 2\|\boldsymbol{X}^o_{r_c}\|_*.$$
(15)

Denote $P\mathcal{D}(\sigma(Z_{33}))Q^{\top}$ as the SVD of Z_{33} where $\sigma(Z_{33}) \in \mathbb{R}^{\hat{t}}$ and $P \in \mathbb{R}^{(m-r) \times (m-r)}$, $Q \in \mathbb{R}^{(n-r) \times (n-r)}$ are orthogonal matrices. We begin dividing $[\hat{t}]$ into subsets of size $k(1 < k \leq \hat{t})$, that is

$$[\hat{t}] = T_1 \cup T_2 \cup \cdots \cup T_h,$$

where each T_i contains k indices probably except T_h , and T_1 contains the indices of k largest coefficients of $\sigma(\mathbf{Z}_{33})$, T_2 contains the indices of the next k largest coefficients, and so on. Define

$$\boldsymbol{Z}_{T_i} := \boldsymbol{U} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & \boldsymbol{P} \mathcal{D}(\sigma_{T_i}(\boldsymbol{Z}_{33})) \boldsymbol{Q}^\top \end{pmatrix} \boldsymbol{V}^\top$$
(16)

for $1 \le i \le h$. Hence Z_1, Z_2, Z_{T_i} are all orthogonal to each other and $\operatorname{rank}(Z_{T_i}) \le k$. Then, on the one hand, we obtain

$$\begin{split} &\|\mathcal{A}(\boldsymbol{Z}_{1}+\boldsymbol{Z}_{2}+\boldsymbol{Z}_{T_{1}})\|_{2}^{2} \\ = &\langle \mathcal{A}(\boldsymbol{Z}_{1}+\boldsymbol{Z}_{2}+\boldsymbol{Z}_{T_{1}}), \mathcal{A}(\boldsymbol{Z}_{1}+\boldsymbol{Z}_{2}+\boldsymbol{Z}_{T_{1}}) \rangle \\ = &\langle \mathcal{A}(\boldsymbol{Z}_{1}+\boldsymbol{Z}_{2}+\boldsymbol{Z}_{T_{1}}), \mathcal{A}(\boldsymbol{Z}) \rangle - \langle \mathcal{A}(\boldsymbol{Z}_{1}+\boldsymbol{Z}_{2}+\boldsymbol{Z}_{T_{1}}), \mathcal{A}(\boldsymbol{Z}_{3}-\boldsymbol{Z}_{T_{1}}) \rangle \\ = &- \langle \mathcal{A}(\boldsymbol{Z}_{1}+\boldsymbol{Z}_{2}+\boldsymbol{Z}_{T_{1}}), \mathcal{A}(\boldsymbol{Z}_{3}-\boldsymbol{Z}_{T_{1}}) \rangle. \end{split}$$

On the other hand, direct calculation yields

$$\begin{aligned} &|\langle \mathcal{A}(\mathbf{Z}_{1} + \mathbf{Z}_{2} + \mathbf{Z}_{T_{1}}), \mathcal{A}(\mathbf{Z}_{3} - \mathbf{Z}_{T_{1}})\rangle| \\ &= |\langle \mathcal{A}(\mathbf{Z}_{1}), \mathcal{A}(\mathbf{Z}_{3} - \mathbf{Z}_{T_{1}})\rangle + \langle \mathcal{A}(\mathbf{Z}_{2} + \mathbf{Z}_{T_{1}}), \mathcal{A}(\mathbf{Z}_{3} - \mathbf{Z}_{T_{1}})\rangle \\ &\leq \sum_{i \geq 2} |\langle \mathcal{A}(\mathbf{Z}_{1}), \mathcal{A}(\mathbf{Z}_{T_{i}})\rangle| + \sum_{i \geq 2} |\langle \mathcal{A}(\mathbf{Z}_{2} + \mathbf{Z}_{T_{1}}), \mathcal{A}(\mathbf{Z}_{T_{i}})\rangle| \\ &\leq (\delta_{r_{1}} ||\mathbf{Z}_{1}||_{F} + \delta_{r_{2}} ||\mathbf{Z}_{2} + \mathbf{Z}_{T_{1}}||_{F}) \sum_{i \geq 2} ||\mathbf{Z}_{T_{i}}||_{F} \\ &\leq \sqrt{2} \delta_{r_{3}} ||\mathbf{Z}_{1} + \mathbf{Z}_{2} + \mathbf{Z}_{T_{1}}||_{F} \sum_{i \geq 2} ||\mathbf{Z}_{T_{i}}||_{F}, \end{aligned}$$

where the second inequality comes from Lemma 9 and the third inequality comes from the monotonicity of the *RIP* constant. Easily, invoking Lemma 10, we can find that

$$\begin{split} &|\langle \mathcal{A}(\boldsymbol{Z}_1 + \boldsymbol{Z}_2 + \boldsymbol{Z}_{T_1}), \mathcal{A}(\boldsymbol{Z}_3 - \boldsymbol{Z}_{T_1})\rangle| \\ \leq &\sqrt{2}\delta_{\max\{r + \lceil \frac{3}{2}k \rceil, 2k\}} \|\boldsymbol{Z}_1 + \boldsymbol{Z}_2 + \boldsymbol{Z}_{T_1}\|_F \sum_{i \geq 2} \|\boldsymbol{Z}_{T_i}\|_F \end{split}$$

since the arbitrariness of m_1, n_1, m_2, n_2 .

Hence

$$\begin{aligned} & \|\mathcal{A}(\boldsymbol{Z}_1 + \boldsymbol{Z}_2 + \boldsymbol{Z}_{T_1})\|_2^2 \\ \leq & \sqrt{2}\delta_{\max\{r + \lceil \frac{3}{2}k \rceil, 2k\}} \|\boldsymbol{Z}_1 + \boldsymbol{Z}_2 + \boldsymbol{Z}_{T_1}\|_F \sum_{i \ge 2} \|\boldsymbol{Z}_{T_i}\|_F, \end{aligned}$$

combining with the fact that

$$\|\mathcal{A}(\mathbf{Z}_1 + \mathbf{Z}_2 + \mathbf{Z}_{T_1})\|_2^2 \ge (1 - \delta_{2r+k}) \|\mathbf{Z}_1 + \mathbf{Z}_2 + \mathbf{Z}_{T_1}\|_F^2,$$

we have

$$\|\boldsymbol{Z}_{1} + \boldsymbol{Z}_{2} + \boldsymbol{Z}_{T_{1}}\|_{F} \leq \frac{\sqrt{2\delta_{\max\{r + \lceil \frac{3}{2}k \rceil, 2k\}}}}{1 - \delta_{2r+k}} \sum_{i \geq 2} \|\boldsymbol{Z}_{T_{i}}\|_{F}.$$
(17)

Now we will give an upper bound for the right side of (17). For any $q \in T_i$ with $i \ge 2$, according to Lemma 8 it yields

$$\sigma_q \le \min_{p \in T_{i-1}} \sigma_p(\mathbf{Z}_{33}) \le \frac{\|\mathbf{Z}_{T_{i-1}}\|_* - \|\mathbf{Z}_{T_{i-1}}\|_F}{k - \sqrt{k}},$$

which implies

$$\|\sigma_{T_i}(\mathbf{Z}_{33})\|_2 \le \sqrt{k} \frac{\|\mathbf{Z}_{T_{i-1}}\|_* - \|\mathbf{Z}_{T_{i-1}}\|_F}{k - \sqrt{k}}.$$

By direct calculation, we have

$$\sum_{i=2}^{h} \|\boldsymbol{Z}_{T_{i}}\|_{F} \leq \frac{1}{\sqrt{k}-1} \sum_{i=2}^{h} (\|\boldsymbol{Z}_{T_{i-1}}\|_{*} - \|\boldsymbol{Z}_{T_{i-1}}\|_{F})$$

$$\leq \frac{1}{\sqrt{k}-1} \sum_{i=1}^{h} (\|\boldsymbol{Z}_{T_{i}}\|_{*} - \|\boldsymbol{Z}_{T_{i}}\|_{F})$$

$$\leq \frac{1}{\sqrt{k}-1} (\|\boldsymbol{Z}_{3}\|_{*} - \|\boldsymbol{Z}_{3}\|_{F})$$

$$\leq \frac{\sqrt{2r}+1}{\sqrt{k}-1} \|\boldsymbol{Z}_{1} + \boldsymbol{Z}_{2}\|_{F} + \frac{2}{\sqrt{k}-1} \|\boldsymbol{X}_{r_{c}}^{o}\|_{*}, \qquad (18)$$

where the third inequality holds since

$$\sum_{i=1}^{h} \|\boldsymbol{Z}_{T_i}\|_F \ge \sqrt{\sum_{i=1}^{h} \|\boldsymbol{Z}_{T_i}\|_F^2} = \|\boldsymbol{Z}_3\|_F,$$

and the last inequality comes from (15). Combining with (17), it is easy to verify that

$$\|\boldsymbol{Z}_{1} + \boldsymbol{Z}_{2} + \boldsymbol{Z}_{T_{1}}\|_{F} \leq \frac{2\beta}{(\sqrt{2r} + 1)(1 - \beta)} \|\boldsymbol{X}_{r_{c}}^{o}\|_{*}$$
(19)

since $1 - \beta > 0$ from the assumption. Then we have

$$egin{aligned} \|m{Z}\|_F &\leq \|m{Z}_1 + m{Z}_2 + m{Z}_{T_1}\|_F + \sum_{i=2}^h \|m{Z}_{T_i}\|_F \ &\leq rac{\sqrt{k} + \sqrt{2r}}{\sqrt{k} - 1} \|m{Z}_1 + m{Z}_2 + m{Z}_{T_1}\|_F + rac{2}{\sqrt{k} - 1} \|m{X}^o_{r_c}\|_* \ &\leq lpha \|m{X}^o_{r_c}\|_*, \end{aligned}$$

where the first inequality follows from (18) and the last inequality follows from (19). This indicates

$$\|\hat{\boldsymbol{X}}^* - \boldsymbol{X}^o\|_F \le \alpha \|\boldsymbol{X}^o_{r_c}\|_*.$$

Hence, we get the desired results. The special case $rank(X^o) \le r$ implies $X^o = X_r^o$ and hence $\hat{X}^* = X^o$ is trivial. Thus, we complete the proof.

Naturally, by setting different values of k, we can get different bounds on isometry constant. When choosing k = 2r, we obtain the following RIP condition involved in δ_{4r} .

Corollary 12. Let $\mathcal{A} : \mathbb{R}^{m \times n} \to \mathbb{R}^{l}$ be a linear map and $\boldsymbol{b} \in \mathbb{R}^{l}$. Set the desired matrix \boldsymbol{X}^{o} satisfies $\mathcal{A}(\boldsymbol{X}^{o}) = \boldsymbol{b}$, that implies the vector $\boldsymbol{s} = 0$ in (5). For a given positive integer $r \leq t$, if linear map \mathcal{A} obeys

$$\delta_{4r} < \frac{\sqrt{2r-1}}{\sqrt{2r-1} + \sqrt{2}(\sqrt{2r+1})},\tag{20}$$

then any optimal solution \hat{X}^* of problem (4) satisfies

$$\|\boldsymbol{X}^{o} - \hat{\boldsymbol{X}}^{*}\|_{F} \leq \hat{\alpha} \|\boldsymbol{X}^{o} - X_{r}^{o}\|_{F},$$

where $\hat{\alpha} := \frac{2 + (2\sqrt{2} - 2)\delta_{4r}}{(\sqrt{2r} - 1) - [(\sqrt{2r} - 1) + \sqrt{2}(\sqrt{2r} + 1)]\delta_{4r}}$. Moreover, if \mathbf{X}^o is r-rank matrix, the unique minimizer of problem (4) is exactly \mathbf{X}^o .

Next we shall establish the recovery estimation of constrained L_{*-F} minimization when measurements are contaminated by noise.

Theorem 13. Under the assumptions of Theorem 11 except that the desired matrix X^o satisfies $\mathcal{A}(X^o) + s = b$ with perturbation $||s||_2 \leq \epsilon$, we have that the optimal solution X^{opt} of problem (6) satisfies

$$\|\boldsymbol{X}^{o} - \boldsymbol{X}^{opt}\|_{F} \leq \alpha \|\boldsymbol{X}^{o} - X_{r}^{o}\|_{F} + \bar{\alpha}\epsilon,$$

where

$$\alpha := \frac{2(\sqrt{2r}+1) + 2\beta(\sqrt{k}-1)}{(\sqrt{k}-1)(1-\beta)(\sqrt{2r}+1)},$$

and

$$\bar{\alpha} := \frac{2(\sqrt{k} + \sqrt{2r})\sqrt{1 + \delta_{2r+k}}}{(\sqrt{k} - 1)(1 - \beta)(1 - \delta_{2r+k})}.$$

Moreover, if X^o is r-rank matrix, then it yields

$$\|\boldsymbol{X}^{o} - \boldsymbol{X}^{opt}\|_{F} \leq \bar{\alpha}\epsilon.$$

Proof. Let $X^{opt} = X^o + Z$, then starting from the block decomposition of Z with respect to X^o and the fact

$$\|X^{opt}\|_{*} - \|X^{opt}\|_{F} \le \|X^{o}\|_{*} - \|X^{o}\|_{F},$$

we repeat the arguments in the proof of Theorem 11 and obtain

$$-\sqrt{2}\delta_{\max\{r+\lceil\frac{3}{2}k\rceil,2k\}}(\|\boldsymbol{Z}_{1}+\boldsymbol{Z}_{2}+\boldsymbol{Z}_{T_{1}}\|_{F})\sum_{i\geq 2}^{h}\|\boldsymbol{Z}_{T_{i}}\|_{F}$$
$$\leq \langle \mathcal{A}(\boldsymbol{Z}_{1}+\boldsymbol{Z}_{2}+\boldsymbol{Z}_{T_{1}}),\mathcal{A}(\boldsymbol{Z})\rangle - \|\mathcal{A}(\boldsymbol{Z}_{1}+\boldsymbol{Z}_{2}+\boldsymbol{Z}_{T_{1}})\|_{2}^{2}$$

and

$$\sum_{i=2}^{h} \|\boldsymbol{Z}_{T_{i}}\|_{F} \leq \frac{\sqrt{2r}+1}{\sqrt{k}-1} \|\boldsymbol{Z}_{1}+\boldsymbol{Z}_{2}\|_{F} + \frac{2}{\sqrt{k}-1} \|\boldsymbol{X}_{r_{c}}^{o}\|_{*}.$$
(21)

Easily, we can find

$$-\sqrt{2}\delta_{\max\{r+\lceil\frac{3}{2}k\rceil,2k\}}(\|\boldsymbol{Z}_{1}+\boldsymbol{Z}_{2}+\boldsymbol{Z}_{T_{1}}\|_{F})\sum_{i\geq2}^{n}\|\boldsymbol{Z}_{T_{i}}\|_{F}$$

$$\leq 2\epsilon\sqrt{1+\delta_{2r+k}}\|\boldsymbol{Z}_{1}+\boldsymbol{Z}_{2}+\boldsymbol{Z}_{T_{1}}\|_{F}-(1-\delta_{2r+k})\|\boldsymbol{Z}_{1}+\boldsymbol{Z}_{2}+\boldsymbol{Z}_{T_{1}}\|_{F}^{2}$$

which together with (21) yields

$$(1 - \delta_{2r+k}) \| \mathbf{Z}_1 + \mathbf{Z}_2 + \mathbf{Z}_{T_1} \|_F$$

$$\leq \sqrt{2} \delta_{\max\{r + \lceil \frac{3}{2}k \rceil, 2k\}} \frac{\sqrt{2r} + 1}{\sqrt{k} - 1} \| \mathbf{Z}_1 + \mathbf{Z}_2 \|_F + \frac{2\sqrt{2} \delta_{\max\{r + \lceil \frac{3}{2}k \rceil, 2k\}}}{\sqrt{k} - 1} \| \mathbf{X}_{r_c}^o \|_* + 2\sqrt{1 + \delta_{2r+k}} \epsilon.$$

Since

$$\beta := \frac{\sqrt{2}\delta_{\max\{r + \lceil \frac{3}{2}k \rceil, 2k\}}(\sqrt{2r} + 1)}{(1 - \delta_{2r+k})(\sqrt{k} - 1)} < 1$$

and $\|Z_1 + Z_2\|_F \le \|Z_1 + Z_2 + Z_{T_1}\|_F$, we have

$$\|\boldsymbol{Z}_{1} + \boldsymbol{Z}_{2} + \boldsymbol{Z}_{T_{1}}\|_{F} \leq \frac{2\sqrt{2}\delta_{\max\{r+\lceil\frac{3}{2}k\rceil,2k\}}}{(1-\delta_{2r+k})(\sqrt{k}-1)(1-\beta)} \|\boldsymbol{X}_{r_{c}}^{o}\|_{*} + \frac{2\sqrt{1+\delta_{2r+k}}}{(1-\delta_{2r+k})(1-\beta)}\epsilon.$$
(22)

$$\|\boldsymbol{Z}\|_{F} \leq \|\boldsymbol{Z}_{1} + \boldsymbol{Z}_{2} + \boldsymbol{Z}_{T_{1}}\|_{F} + \sum_{i \geq 2}^{h} \|\boldsymbol{Z}_{T_{i}}\|_{F}$$
$$\leq \frac{\sqrt{k} + \sqrt{2r}}{\sqrt{k} - 1} \|\boldsymbol{Z}_{1} + \boldsymbol{Z}_{2} + \boldsymbol{Z}_{T_{1}}\|_{F} + \frac{2}{\sqrt{k} - 1} \|\boldsymbol{X}_{r_{c}}^{o}\|_{*}$$
$$\leq \alpha \|\boldsymbol{X}_{r_{c}}^{o}\|_{*} + \bar{\alpha}\epsilon.$$

Here the first inequality comes from (21), and the last inequality comes from (22) and the definitions of α and $\bar{\alpha}$. Additionally, from the above discussion, the desired result is easily obtained when X^o is *r*-rank matrix. Thus, we complete the proof. \Box

III. MAIN RESULTS ON UNCONSTRAINED MINIMIZATION

A. Relation Between Problem (4) and Problem (7)

We now show that in some sense, problem (4) can be solved via solving problem (7). We note that the regularization term $\|X\|_* - \|X\|_F$ is nonconvex and nonsmooth, hence the result is nontrivial.

Theorem 14. Assume linear map \mathcal{A} obeys $\delta_1 < 1$. Let $\{\lambda_n\}$ be a decreasing sequence of positive numbers with $\lambda_n \to 0$ and \bar{X}_{λ_n} be the optimal solution of the problem (7) with $\lambda = \lambda_n$. If problem (4) is feasible, then sequence $\{\bar{X}_{\lambda_n}\}$ is bounded and any of its accumulation points is the optimal solution of the problem (4).

Proof. Let \bar{X} be any feasible point of problem (4), then $\mathcal{A}(\bar{X}) = b$. Since \bar{X}_{λ_n} is the optimal solution of problem (7) with respect to $\lambda = \lambda_n$, we have

$$\|\bar{\boldsymbol{X}}_{\lambda_{n}}\|_{*} - \|\bar{\boldsymbol{X}}_{\lambda_{n}}\|_{F} + \frac{1}{2\lambda_{n}}\|\mathcal{A}(\bar{\boldsymbol{X}}_{\lambda_{n}}) - \boldsymbol{b}\|_{2}^{2} \leq \|\bar{\boldsymbol{X}}\|_{*} - \|\bar{\boldsymbol{X}}\|_{F}.$$
(23)

Since $\lambda_n \to 0$ as $n \to \infty$, the sequences $\{\|\bar{X}_{\lambda_n}\|_* - \|\bar{X}_{\lambda_n}\|_F\}$ and $\{\|\mathcal{A}(\bar{X}_{\lambda_n}) - b\|_2\}$ converge to zero and so they are bounded.

Next we will show that the sequence $\{\bar{X}_{\lambda_n}\}$ is bounded and hence it has at least one accumulation point. Denote $U_{\lambda_n} \mathcal{D}(\sigma(\bar{X}_{\lambda_n})) V_{\lambda_n}$ as the SVD of \bar{X}_{λ_n} . Define

$$ar{oldsymbol{X}}_{\lambda_n}^1 := oldsymbol{U}_{\lambda_n} \mathcal{D}(\sigma^1(ar{oldsymbol{X}}_{\lambda_n}))oldsymbol{V}_{\lambda_n}$$

and

$$ar{X}_{\lambda_n}^2 := U_{\lambda_n} \mathcal{D}(\sigma^2(ar{X}_{\lambda_n})) V_{\lambda_n},$$

where $\sigma_i^1(\bar{X}_{\lambda_n}) = \sigma_i(\bar{X}_{\lambda_n})$ for $i \leq 2$, $\sigma_i^2(\bar{X}_{\lambda_n}) = \sigma_i(\bar{X}_{\lambda_n})$ for i > 2 and others are 0. Likewise, let us define matrices

$$oldsymbol{Y}_{\lambda_n}^1 \coloneqq oldsymbol{U}_{\lambda_n} \mathcal{D}(oldsymbol{y}_{\lambda_n}^1) oldsymbol{V}_{\lambda_n}, \hspace{0.2cm} oldsymbol{Y}_{\lambda_n}^2 \coloneqq oldsymbol{U}_{\lambda_n} \mathcal{D}(oldsymbol{y}_{\lambda_n}^2) oldsymbol{V}_{\lambda_n}$$

where $(\boldsymbol{y}_{\lambda_n}^1)_i = \sigma_i(\bar{\boldsymbol{X}}_{\lambda_n})$ for i = 1, $(\boldsymbol{y}_{\lambda_n}^2)_i = \sigma_i(\bar{\boldsymbol{X}}_{\lambda_n})$ for i > 1 and others are 0. Clearly, we have

$$ar{oldsymbol{X}}_{\lambda_n} = ar{oldsymbol{X}}_{\lambda_n}^1 + ar{oldsymbol{X}}_{\lambda_n}^2 = oldsymbol{Y}_{\lambda_n}^1 + oldsymbol{Y}_{\lambda_n}^2.$$

By direct computation, it follows from Lemma 8 (b) that

$$\begin{split} \|\bar{\boldsymbol{X}}_{\lambda_{n}}\|_{*} - \|\bar{\boldsymbol{X}}_{\lambda_{n}}\|_{F} \geq & \|\bar{\boldsymbol{X}}_{\lambda_{n}}^{1}\|_{*} + \|\bar{\boldsymbol{X}}_{\lambda_{n}}^{2}\|_{*} - \|\bar{\boldsymbol{X}}_{\lambda_{n}}^{1}\|_{F} - \|\bar{\boldsymbol{X}}_{\lambda_{n}}^{2}\|_{F} \\ \geq & \|\bar{\boldsymbol{X}}_{\lambda_{n}}^{1}\|_{*} - \|\bar{\boldsymbol{X}}_{\lambda_{n}}^{1}\|_{F} \\ \geq & (2 - \sqrt{2})\sigma_{2}(\bar{\boldsymbol{X}}_{\lambda_{n}}), \end{split}$$

which, together with the boundedness of the sequence $\{\|\bar{X}_{\lambda_n}\|_* - \|\bar{X}_{\lambda_n}\|_F\}$, implies the sequence $\{\|Y_{\lambda_n}^2\|_F\}$ is bounded. Hence, the inequality

$$\|\mathcal{A}(oldsymbol{Y}_{\lambda_n}^2) - oldsymbol{b}\|_2 \leq \|\mathcal{A}\| \|oldsymbol{Y}_{\lambda_n}^2\|_F + \|oldsymbol{b}\|_2$$

yields the sequence $\{\|\mathcal{A}(\mathbf{Y}_{\lambda_n}^2) - \mathbf{b}\|_2\}$ is bounded where $\|\mathcal{A}\|$ is the operator norm of linear map \mathcal{A} . As a result, since $\{\|\mathcal{A}(\bar{\mathbf{X}}_{\lambda_n}) - \mathbf{b}\|_2\}$ is bounded and

$$\|\mathcal{A}(\boldsymbol{Y}_{\lambda_n}^1)\|_2 \leq \|\mathcal{A}(\bar{\boldsymbol{X}}_{\lambda_n}) - b\|_2 + \|\mathcal{A}(\boldsymbol{Y}_{\lambda_n}^2) - \boldsymbol{b}\|_2,$$

we obtain that the sequence $\{\|\mathcal{A}(\mathbf{Y}_{\lambda_n}^1)\|_2\}$ is bounded. Furthermore, since linear map \mathcal{A} obeys $\delta_1 < 1$, it follows from (8) that the sequence $\{\|\mathbf{Y}_{\lambda_n}^1\|_F\}$ is also bounded. Hence, the boundedness of $\{\bar{\mathbf{X}}_{\lambda_n}\}$ can be seen from $\bar{\mathbf{X}}_{\lambda_n} = \mathbf{Y}_{\lambda_n}^1 + \mathbf{Y}_{\lambda_n}^2$ and the boundedness of $\{\|\mathbf{Y}_{\lambda_n}\|_F\}$ and $\{\|\mathbf{Y}_{\lambda_n}^2\|_F\}$.

Besides, the inequality (23) shows

$$\frac{1}{2} \| \mathcal{A}(\bar{\boldsymbol{X}}_{\lambda_n}) - \boldsymbol{b} \|_2^2 \le \lambda_n (\| \bar{\boldsymbol{X}} \|_* - \| \bar{\boldsymbol{X}} \|_F)$$

for any $\lambda_n \to 0$. Let X^* be any accumulation point of the sequence $\{\bar{X}_{\lambda_n}\}$. Then we can derive that $\mathcal{A}(X^*) = \mathbf{b}$, which combines $\|X^*\|_* - \|X^*\|_F \le \|\bar{X}\|_* - \|\bar{X}\|_F$ and the arbitrariness of \bar{X} lead to X^* being the optimal solution of problem (4). This completes the proof.

B. Recovery Error Estimation for Problem (7)

In this subsection, we provide some theoretical investigations to guarantee the robust recovery of the regularized L_{1-2} minimization problem (7). Let us start with some powerfully technical tools used in the proof of our main results. The following lemma states an elementary geometric fact: any point in a ploytope can be represented as a convex combination of sparse vectors.

Lemma 15. [Sparse Representation of a Polytope [21]] For a positive number α and a positive integer s, define the polytope $T(\alpha, s) \subset \mathbb{R}^p$ by

$$T(\alpha, s) = \{ \boldsymbol{v} \in \mathbb{R}^p : \| \boldsymbol{v} \|_{\infty} \le \alpha, \| \boldsymbol{v} \|_1 \le s \alpha \}$$

For any $\boldsymbol{v} \in \mathbb{R}^p$, define the set of sparse vectors $\mathcal{U}(\alpha, s, \boldsymbol{v}) \subset \mathbb{R}^p$ by

$$\mathcal{U}(\alpha, s, \boldsymbol{v}) = \{ \boldsymbol{u} \in \mathbb{R}^p : \operatorname{supp}(\boldsymbol{u}) \subseteq \operatorname{supp}(\boldsymbol{v}), \|\boldsymbol{u}\|_0 \le s, \|\boldsymbol{u}\|_1 = \|\boldsymbol{v}\|_1, \|\boldsymbol{u}\|_\infty \le \alpha \}.$$
(24)

Then $v \in T(\alpha, s)$ if and only if v is in the convex hull of $\mathcal{U}(\alpha, s, v)$. In particularly, any $v \in T(\alpha, s)$ can be expressed as

$$oldsymbol{v} = \sum_{i=1}^N \lambda_i oldsymbol{u}_i$$

for some positive integer N, where $u_i \in \mathcal{U}(\alpha, s, v)$ and

$$\sum_{i=1}^{N} \lambda_i = 1 \quad \text{with} \quad 0 \le \lambda_i \le 1.$$
(25)

The following lemma established in [34] is also necessary which give an inequality between the sums of the α th power of two sequences of nonnegative numbers based on the inequality of their sums.

Lemma 16. Suppose $m \ge r$, $a_1 \ge a_2 \ge \cdots \ge a_m \ge 0$, and $\sum_{i=1}^r a_i \ge \sum_{i=r+1}^m a_i$, then for all $\alpha \ge 1$,

$$\sum_{j=r+1}^{m} a_j^{\alpha} \le \sum_{i=1}^{r} a_i^{\alpha}.$$

More generally, suppose $a_1 \ge a_2 \ge \cdots \ge a_m \ge 0$, $\eta \ge 0$ and $\sum_{i=1}^r a_i + \eta \ge \sum_{i=r+1}^m a_i$, then for all $\alpha \ge 1$,

$$\sum_{j=r+1}^{m} a_j^{\alpha} \le r \left(\sqrt[\alpha]{\frac{\sum_{i=1}^{r} a_i^{\alpha}}{r}} + \frac{\eta}{r} \right)^{\alpha}$$

With the above preparation, we may state and prove the critical results, which play key role in recovery estimation of regularized minimization.

Lemma 17. Let k be a positive integer and linear map \mathcal{A} obey the tk-order RIP with $\delta_{tk} \in (0,1)$ for certain integer t > 1. Then for any subset $T \subseteq [n]$ with $|T| \leq k$ and any matrix $H \in \mathbb{R}^{m \times n}$, we have

$$\|\boldsymbol{h}_{T}\|_{2} \leq \frac{\beta_{1}}{\sqrt{k}} \|\boldsymbol{h}_{T^{C}}\|_{1} + \gamma_{1} \|\mathcal{A}(\boldsymbol{H})\|_{2},$$
(26)

where $\boldsymbol{h} \in \mathbb{R}^d$ is the singular value vector of \boldsymbol{H} and

$$\beta_1 = \frac{\delta_{tk}}{\sqrt{(t-1)[1-\delta_{tk}^2]}}, \quad \gamma_1 = \frac{2}{(1-\delta_{tk})\sqrt{1+\delta_{tk}}}.$$
(27)

Proof. For given $H \in \mathbb{R}^{m \times n}$, let $U\mathcal{D}(h)V^{\top}$ be SVD of H, and for given t > 1, define

$$T_{1} = \left\{ i \in T^{c} \left| \left| \left(\boldsymbol{h}_{T^{c}} \right)_{i} \right| > \frac{\| \boldsymbol{h}_{T^{c}} \|_{1}}{(t-1)k} \right\},$$
(28)

$$T_{2} = \left\{ i \in T^{c} \big| \left(\boldsymbol{h}_{T^{c}} \right)_{i} \right| \leq \frac{\|\boldsymbol{h}_{T^{c}}\|_{1}}{(t-1)k} \right\}.$$
(29)

Note that $T^c = T_1 \cup T_2$ which yields

$$h = h_T + h_{T^c} = h_T + h_{T_1} + h_{T_2}.$$
 (30)

Besides, the fact $T_1 \cap T = \emptyset$ implies $\|h_T\|_2 \le \|h_{T \cup T_1}\|_2$. Thus, to show (26), it suffices to show that

$$\|\boldsymbol{h}_{T\cup T_{1}}\|_{2} \leq \frac{\beta_{1}}{\sqrt{k}} \|\boldsymbol{h}_{T^{c}}\|_{1} + \gamma_{1} \|\mathcal{A}(\boldsymbol{H})\|_{2},$$
(31)

where β_1 and γ_1 are defined as (27). We can apply Lemma 15 to show (31) to be true.

For this purpose, we first show that

$$|T_1| < (t-1)k. (32)$$

Obviously, the above inequality holds if $T_1 = \emptyset$. Otherwise, we can apply (28) to get that

$$\|\boldsymbol{h}_{T_1}\|_1 > |T_1| \frac{\|\boldsymbol{h}_{T^c}\|_1}{(t-1)k} = |T_1| \frac{\|\boldsymbol{h}_{T_1 \cup T_2}\|_1}{(t-1)k} \ge |T_1| \frac{\|\boldsymbol{h}_{T_1}\|_1}{(t-1)k},$$

which implies that (32) holds.

We now turn to show (31) holds. The relation $T^c = T_1 \cup T_2$ and $T_1 \cap T_2 = \emptyset$, along with the expression (28), indicates that

$$\begin{split} \|\boldsymbol{h}_{T_2}\|_1 &= \|\boldsymbol{h}_{T^c}\|_1 - \|\boldsymbol{h}_{T_1}\|_1 \\ &\leq \|\boldsymbol{h}_{T^c}\|_1 - |T_1| \frac{\|\boldsymbol{h}_{T^c}\|_1}{(t-1)k} \\ &= ((t-1)k - |T_1|) \frac{\|\boldsymbol{h}_{T^c}\|_1}{(t-1)k}. \end{split}$$

Since tk is an integer, it follows from (32) that $(t-1)k - |T_1|$ is a positive integer. Additionally, (29) implies that

$$\|m{h}_{T_2}\|_{\infty} \leq rac{\|m{h}_{T^c}\|_1}{(t-1)k}.$$

By setting

$$m{v} = m{h}_{T_2}, \,\, lpha = rac{\|m{h}_{T^c}\|_1}{(t-1)k}, \,\, s = (t-1)k - |T_1|$$

and applying Lemma 15, we have

$$\boldsymbol{h}_{T_2} = \sum_{i=1}^N \lambda_i \boldsymbol{u}_i \tag{33}$$

for some positive integer N, where $\sum_{i=1}^{N} \lambda_i = 1$ with $0 \le \lambda_i \le 1$ and

$$\boldsymbol{u}_{i} \in \mathcal{U}\left(\frac{\|\boldsymbol{h}_{T^{c}}\|_{1}}{(t-1)k}, (t-1)k - |T_{1}|, \boldsymbol{h}_{T_{2}}\right)$$
(34)

with \mathcal{U} being defined as (24). Moreover, the expression (24) implies that

$$\begin{aligned} \|\boldsymbol{u}_{i}\|_{2}^{2} &\leq \|\boldsymbol{u}_{i}\|_{0} \|\boldsymbol{u}_{i}\|_{\infty}^{2} \\ &\leq \left((t-1)k - |T_{1}|\right) \left(\frac{\|\boldsymbol{h}_{T^{c}}\|_{1}}{(t-1)k}\right)^{2} \\ &\leq \frac{\left(\|\boldsymbol{h}_{T^{c}}\|_{1}\right)^{2}}{(t-1)k}. \end{aligned}$$
(35)

For simplicity, denote

$$\boldsymbol{v}_i := (1 + \delta_{tk})\boldsymbol{h}_{T \cup T_1} + \delta_{tk}\boldsymbol{u}_i, \tag{36}$$

$$\bar{\boldsymbol{v}}_i := (1 - \delta_{tk})\boldsymbol{h}_{T \cup T_1} - \delta_{tk}\boldsymbol{u}_i. \tag{37}$$

The relation $\|\boldsymbol{u}_i\|_0 \leq (t-1)k - |T_1|$ combined with (32) and the fact $|T| \leq k$ yield that \boldsymbol{v}_i and $\bar{\boldsymbol{v}}_i$ are tk-sparse for $1 \leq i \leq N$.

Thus, we have

$$\sum_{i=1}^{N} \lambda_{i} \left[\left\| \mathcal{A} \left(U \mathcal{D}(\boldsymbol{v}_{i}) \boldsymbol{V}^{\top} \right) \right\|_{2}^{2} - \left\| \mathcal{A} \left(U \mathcal{D}(\bar{\boldsymbol{v}}_{i}) \boldsymbol{V}^{\top} \right) \right\|_{2}^{2} \right]$$

$$= \sum_{i=1}^{N} \lambda_{i} \left[\left\| \mathcal{A} \left(U \mathcal{D}(\boldsymbol{h}_{T \cup T_{1}}) \boldsymbol{V}^{\top} \right) + \delta_{tk} \mathcal{A} \left(U \mathcal{D}(\boldsymbol{u}_{i} + \boldsymbol{h}_{T \cup T_{1}}) \boldsymbol{V}^{\top} \right) \right\|_{2}^{2} - \left\| \mathcal{A} \left(U \mathcal{D}(\boldsymbol{h}_{T \cup T_{1}}) \boldsymbol{V}^{\top} \right) \right. \right. \right.$$

$$\left. - \delta_{tk} \mathcal{A} \left(U \mathcal{D}(\boldsymbol{u}_{i} + \boldsymbol{h}_{T \cup T_{1}}) \boldsymbol{V}^{\top} \right) \right\|_{2}^{2} \right]$$

$$= \sum_{i=1}^{N} \lambda_{i} \left[4 \delta_{tk} \left(\mathcal{A} (U \mathcal{D}(\boldsymbol{h}_{T \cup T_{1}}) \boldsymbol{V}^{\top} \right) \right)^{\top} \mathcal{A} \left(U \mathcal{D}(\boldsymbol{u}_{i} + \boldsymbol{h}_{T \cup T_{1}}) \boldsymbol{V}^{\top} \right) \right]$$

$$\left. \stackrel{(a)}{=} 4 \delta_{tk} \left(\mathcal{A} (U \mathcal{D}(\boldsymbol{h}_{T \cup T_{1}}) \boldsymbol{V}^{\top}) \right)^{\top} \mathcal{A} \left(U \mathcal{D}(\boldsymbol{h}) \boldsymbol{V}^{\top} \right)$$

$$= 4 \delta_{tk} \left(\mathcal{A} (U \mathcal{D}(\boldsymbol{h}_{T \cup T_{1}}) \boldsymbol{V}^{\top}) \right)^{\top} \mathcal{A} (H)$$

$$\left. \stackrel{(c)}{\leq} 4 \delta_{tk} \sqrt{1 + \delta_{tk}} \left\| \boldsymbol{h}_{T \cup T_{1}} \right\|_{2} \left\| \mathcal{A} (H) \right\|_{2},$$

$$(38)$$

where (a) follows from (25), (b) follows from (30) and (33), (c) follows from Cauchy-Schwartz inequality and (8). Next we will give a lower bound of the first formula in (38). That is

$$\sum_{i=1}^{N} \lambda_{i} \left[\| \mathcal{A} \left(\mathcal{U} \mathcal{D}(\boldsymbol{v}_{i}) \boldsymbol{V}^{\top} \right) \|_{2}^{2} - \| \mathcal{A} \left(\mathcal{U} \mathcal{D}(\bar{\boldsymbol{v}}_{i}) \boldsymbol{V}^{\top} \right) \|_{2}^{2} \right]$$

$$\stackrel{(a)}{\geq} \sum_{i=1}^{N} \lambda_{i} (1 - \delta_{tk}) \| \boldsymbol{v}_{i} \|_{2}^{2} - \sum_{i=1}^{N} \lambda_{i} (1 + \delta_{tk}) \| \bar{\boldsymbol{v}}_{i} \|_{2}^{2}$$

$$\stackrel{(b)}{=} \sum_{i=1}^{N} \lambda_{i} \left(2(\delta_{tk} - \delta_{tk}^{3}) \| \boldsymbol{h}_{T \cup T_{1}} \|_{2}^{2} - 2\delta_{tk}^{3} \| \boldsymbol{u}_{i} \|_{2}^{2} \right)$$

$$\stackrel{(c)}{\geq} 2(\delta_{tk} - \delta_{tk}^{3}) \| \boldsymbol{h}_{T \cup T_{1}} \|_{2}^{2} - 2\delta_{tk}^{3} \frac{(\| \boldsymbol{h}_{T^{c}} \|_{1})^{2}}{(t - 1)k},$$

where (a) follows from (8), (b) follows from (36) and (37), (c) follows from (25) and (35). The inclusion together with (38) indicates that

$$2(\delta_{tk} - \delta_{tk}^3) \|\boldsymbol{h}_{T \cup T_1}\|_2^2 - 2\delta_{tk}^3 \frac{(\|\boldsymbol{h}_{T^c}\|_1)^2}{(t-1)k} \le 4\delta_{tk} \sqrt{1+\delta_{tk}} \|\boldsymbol{h}_{T \cup T_1}\|_2 \|\mathcal{A}(\boldsymbol{H})\|_2,$$

which is exactly

$$(1-\delta_{tk}^2)\|\boldsymbol{h}_{T\cup T_1}\|_2^2 - 2\sqrt{1+\delta_{tk}}\|\boldsymbol{h}_{T\cup T_1}\|_2\|\mathcal{A}(\boldsymbol{H})\|_2 - \delta_{tk}^2\frac{(\|\boldsymbol{h}_{T^c}\|_1)^2}{(t-1)k} \leq 0.$$

Hence we have

$$\begin{split} \|\boldsymbol{h}_{T\cup T_{1}}\|_{2} \leq & \left[\left((2\sqrt{1+\delta_{tk}} \|\mathcal{A}(\boldsymbol{H})\|_{2})^{2} + 4\delta_{tk}^{2}(1-\delta_{tk}^{2})\frac{(\|\boldsymbol{h}_{T^{c}}\|_{1})^{2}}{(t-1)k} \right)^{\frac{1}{2}} + 2\sqrt{1+\delta_{tk}} \|\mathcal{A}(\boldsymbol{H})\|_{2} \right] \middle/ 2(1-\delta_{tk}^{2}) \\ & \leq & \frac{\delta_{tk}}{\sqrt{(1-\delta_{tk}^{2})(t-1)k}} \|\boldsymbol{h}_{T^{c}}\|_{1} + \frac{2}{(1-\delta_{tk})\sqrt{1+\delta_{tk}}} \|\mathcal{A}(\boldsymbol{H})\|_{2}, \end{split}$$

where the last inequality follows from $\sqrt{x^2 + y^2} \le x + y$ for any $x \ge 0$ and $y \ge 0$.

By virtue of (27) and (31), we obtain the desired results. This completes the proof.

Lemma 18. Let k be a positive integer and the desired matrix X^o satisfy $\mathcal{A}(X^o) + s = b$ with perturbation $||s||_2 \le \epsilon$. Set $H := X^* - X^o$ with $X^* \in \arg \min \mathcal{J}(X)$. Denote $U_o \mathcal{D}(x^0) V_o^{\top}$ and $U_H \mathcal{D}(h) V_H^{\top}$ as the SVD of X^o and H, respectively. Define $\Omega := \operatorname{supp}(x_{\max(k)}^o)$ and $\Gamma := \operatorname{supp}(h_{\max(k)})$, then we have

$$\|\mathcal{A}(\mathbf{H})\|_{2}^{2} - 2\epsilon \|\mathcal{A}(\mathbf{H})\|_{2} \leq 2\lambda (2\|\mathbf{X}^{o}\|_{*} + \|\mathbf{h}_{\Gamma}\|_{1} - \|\mathbf{h}_{\Gamma^{c}}\|_{1} + \|\mathbf{h}\|_{2})$$
(39)

and

$$\|\boldsymbol{h}_{\Gamma^{c}}\|_{1} \leq 2\|\boldsymbol{X}^{o}\|_{*} + \|\boldsymbol{h}_{\Gamma}\|_{1} + \|\boldsymbol{h}\|_{2} + \frac{\epsilon}{\lambda}\|\mathcal{A}(\boldsymbol{H})\|_{2}.$$
(40)

Proof. According to the definition of X^* , we obtain that $\mathcal{J}(X^*) \leq \mathcal{J}(X^o)$, i.e.,

$$\| \boldsymbol{X}^* \|_* - \| \boldsymbol{X}^* \|_F + rac{1}{2\lambda} \| \mathcal{A}(\boldsymbol{X}^*) - \boldsymbol{b} \|_2^2 \le \| \boldsymbol{X}^o \|_* - \| \boldsymbol{X}^o \|_F + rac{1}{2\lambda} \| \mathcal{A}(\boldsymbol{X}^o) - \boldsymbol{b} \|_2^2$$

which after simplification gives

$$\frac{1}{2\lambda} \left(\|\mathcal{A}(\boldsymbol{H})\|_{2}^{2} - 2\langle \boldsymbol{s}, \mathcal{A}(\boldsymbol{H}) \rangle \right) \leq \|\boldsymbol{X}^{o}\|_{*} - \|\boldsymbol{X}^{o}\|_{F} - \|\boldsymbol{X}^{*}\|_{*} + \|\boldsymbol{X}^{*}\|_{F}.$$

$$\tag{41}$$

Denote the left and right sides of (41) as ρ_l and ρ_r respectively. It is known from the Cauchy-Schwartz inequality that

$$\rho_{l} \geq \frac{1}{2\lambda} \left(\|\mathcal{A}(\boldsymbol{H})\|_{2}^{2} - 2\|\boldsymbol{s}\|_{2} \|\mathcal{A}(\boldsymbol{H})\|_{2} \right)$$

$$\geq \frac{1}{2\lambda} \left(\|\mathcal{A}(\boldsymbol{H})\|_{2}^{2} - 2\epsilon \|\mathcal{A}(\boldsymbol{H})\|_{2} \right).$$
(42)

Besides, direct calculations lead to the expression of $H + X^o = A + B$ with

$$oldsymbol{A} = oldsymbol{U}_o \mathcal{D}(oldsymbol{x}_\Omega^o)oldsymbol{V}_o^ op + oldsymbol{U}_H \mathcal{D}(oldsymbol{h}_\Omega)oldsymbol{V}_H^ op$$

and

$$oldsymbol{B} = oldsymbol{U}_o \mathcal{D}(oldsymbol{x}_{\Omega^c}^o)oldsymbol{V}_o^+ + oldsymbol{U}_H \mathcal{D}(oldsymbol{h}_{\Omega^c})oldsymbol{V}_H^+$$
 ,

Then, we have

$$\rho_{r} = \|\boldsymbol{X}^{o}\|_{*} - \|\boldsymbol{X}^{o}\|_{F} - \|\boldsymbol{H} + \boldsymbol{X}^{o}\|_{*} + \|\boldsymbol{H} + \boldsymbol{X}^{o}\|_{F} \\
\leq \|\boldsymbol{X}^{o}\|_{*} - \|\boldsymbol{X}^{o}\|_{F} - \|\boldsymbol{B}\|_{*} + \|\boldsymbol{A}\|_{*} + \|\boldsymbol{H}\|_{F} + \|\boldsymbol{X}^{o}\|_{F} \\
\leq \|\boldsymbol{X}^{o}\|_{*} - \|\boldsymbol{h}_{\Omega^{c}}\|_{1} + \|\boldsymbol{x}_{\Omega^{c}}^{o}\|_{1} + \|\boldsymbol{h}_{\Omega}\|_{1} + \|\boldsymbol{h}\|_{2} \\
\leq 2\|\boldsymbol{X}^{o}\|_{*} + \|\boldsymbol{h}_{\Gamma}\|_{1} - \|\boldsymbol{h}_{\Gamma^{c}}\|_{1} + \|\boldsymbol{h}\|_{2}.$$
(43)

Hence combining (42) with (43), we can get the desired inequalities (39) and (40) hold trivially from (39). \Box

We shall focus on investigating the recovery performance of problem (7) and characterizing the recovery errors of this method.

Theorem 19. Assume that k is a positive integer with $k \ge 6$, and b is given by (5) with $||s||_2 \le \epsilon$. If linear map A obeys

$$\delta_{tk} < \sqrt{\frac{t-1}{t+\theta_k^2 - 1}} \tag{44}$$

for a certain integer t > 1, where θ_k is given by

$$\theta_k = \frac{\sqrt{k} + \sqrt{2} - 1}{\sqrt{k} - \sqrt{2} - 1}.$$
(45)

Then we have

$$\|\boldsymbol{X}^* - \boldsymbol{X}^o\|_F \le C_1 \|\boldsymbol{X}^o\|_* + C_2 \lambda \tag{46}$$

where C_1 and C_2 are determined by

$$C_1 = \frac{2(\beta_1 \hat{\gamma}_1 + \xi_1)}{\sqrt{k}(1 - \beta_1)\hat{\gamma}_1 \kappa_1}, \quad C_2 = \frac{2\xi_1 \hat{\gamma}_1}{\sqrt{k}(1 - \beta_1)\kappa_1},$$

with $\hat{\beta}_1$ and $\hat{\gamma}_1$ being given by (49) and (51), and κ_1 and ξ_1 being given in (52) and (53), respectively.

Proof. Denote $H = X^* - X^o$. Let $X^o = U_o \mathcal{D}(x^0) V_o^{\top}$ and $H = U_H \mathcal{D}(h) V_H^{\top}$ be the SVD of X^o and H, respectively. Define $\Omega := \operatorname{supp}(x_{\max(k)}^o)$, $\Gamma := \operatorname{supp}(h_{\max(k)})$ and $\omega := 2 ||X^o||_* + ||h||_2$. On the one hand, noting the fact $||h_{\Gamma}||_1 \le \sqrt{k} ||h_{\Gamma}||_2$, together with Lemma 17 and Lemma 18, we obtain

$$\|\boldsymbol{h}_{\Gamma}\|_{2} \leq \frac{\beta_{1}}{\sqrt{k}} \left(\omega + \sqrt{k} \|\boldsymbol{h}_{\Gamma}\|_{2} + \frac{\epsilon}{\lambda} \|\mathcal{A}(\boldsymbol{H})\|_{2} \right) + \gamma_{1} \|\mathcal{A}(\boldsymbol{H})\|_{2},$$
(47)

which is exactly

$$\|\boldsymbol{h}_{\Gamma}\|_{2} \leq \frac{\beta_{1}\epsilon + \gamma_{1}\sqrt{k}\lambda}{(1-\beta_{1})\sqrt{k}\lambda} \|\mathcal{A}(\boldsymbol{H})\|_{2} + \frac{\beta_{1}}{(1-\beta_{1})\sqrt{k}}\omega$$
(48)

(49)

(52)

due to $\beta_1 < 1$ by (44) and (45). On the other hand, invoking Lemma 16 and (40), it yields

$$egin{aligned} \|m{h}_{\Gamma^c}\|_2 &\leq \sqrt{k} \left(\sqrt{rac{\|m{h}_{\Gamma}\|_2^2}{k}} + rac{\omega + rac{\epsilon}{\lambda} \|\mathcal{A}(m{H})\|_2}{k}
ight), \ \|m{h}_{\Gamma^c}\|_2 &\leq \|m{h}_{\Gamma}\|_2 + rac{1}{\sqrt{k}} \left(\omega + rac{\epsilon}{\lambda} \|\mathcal{A}(m{H})\|_2
ight). \ \hat{eta}_1 &:= (\sqrt{2} - 1)eta_1 + 1. \end{aligned}$$

that is,

Define

Therefore, we obtain that

$$\|\boldsymbol{h}\|_{2} = \sqrt{\|\boldsymbol{h}_{\Gamma}\|_{2}^{2} + \|\boldsymbol{h}_{\Gamma^{c}}\|_{2}^{2}}$$

$$\leq \sqrt{\|\boldsymbol{h}_{\Gamma}\|_{2}^{2} + \left(\|\boldsymbol{h}_{\Gamma}\|_{2} + \frac{1}{\sqrt{k}}(\omega + \frac{\epsilon}{\lambda}\|\mathcal{A}(\boldsymbol{H})\|_{2})\right)^{2}}$$

$$\leq \sqrt{2}\|\boldsymbol{h}_{\Gamma}\|_{2} + \frac{1}{\sqrt{k}}(\omega + \frac{\epsilon}{\lambda}\|\mathcal{A}(\boldsymbol{H})\|_{2})$$

$$\leq \frac{\sqrt{2k}\gamma_{1}\lambda + \epsilon\left((\sqrt{2}-1)\beta_{1}+1\right)}{\lambda\sqrt{k}(1-\beta_{1})}\|\mathcal{A}(\boldsymbol{H})\|_{2} + \frac{(\sqrt{2}-1)\beta_{1}+1}{\sqrt{k}(1-\beta_{1})}\omega$$

$$= \frac{\sqrt{2k}\gamma_{1}\lambda + \hat{\beta}_{1}\epsilon}{\lambda\sqrt{k}(1-\beta_{1})}\|\mathcal{A}(\boldsymbol{H})\|_{2} + \frac{\hat{\beta}_{1}}{\sqrt{k}(1-\beta_{1})}\omega.$$
(50)

By employing (39) and Lemma 17, we have

$$\|\mathcal{A}(\boldsymbol{H})\|_{2}^{2} - 2\epsilon \|\mathcal{A}(\boldsymbol{H})\|_{2} \leq 2\sqrt{k}\lambda(\frac{\beta_{1}}{\sqrt{k}}\|\boldsymbol{h}_{\Gamma^{c}}\|_{1} + \gamma_{1}\|\mathcal{A}(\boldsymbol{H})\|_{2}) - 2\lambda\|\boldsymbol{h}_{\Gamma^{c}}\|_{1} + 2\lambda\omega,$$

that is,

$$\|\mathcal{A}(\boldsymbol{H})\|_{2}^{2} - (2\sqrt{k}\lambda\gamma_{1} + 2\epsilon)\|\mathcal{A}(\boldsymbol{H})\|_{2} - 2\lambda\omega \leq 2\lambda(\beta_{1} - 1)\|\boldsymbol{h}_{\Gamma^{c}}\|_{1}.$$

 $\|\mathcal{A}(\boldsymbol{H})\|_{2}^{2} - 2\hat{\gamma}_{1}\lambda\|\mathcal{A}(\boldsymbol{H})\|_{2} - 2\lambda\omega \leq 0.$

Define

$$\eta := \frac{\epsilon}{\lambda}, \ \hat{\gamma}_1 := \sqrt{k}\gamma_1 + \eta.$$
(51)

It yields

Thus,

$$\begin{split} \|\mathcal{A}(\boldsymbol{H})\|_{2} &\leq \frac{2\hat{\gamma}_{1}\lambda + \sqrt{4\hat{\gamma}_{1}^{2}\lambda^{2} + 8\lambda\omega}}{2} \\ &\leq \hat{\gamma}_{1}\lambda + \sqrt{(\hat{\gamma}_{1}\lambda + \frac{\omega}{\hat{\gamma}_{1}})^{2}} \\ &= 2\hat{\gamma}_{1}\lambda + \frac{\omega}{\hat{\gamma}_{1}}, \end{split}$$

which indicates that

Define

Then, we have

$$\left(1 - \frac{\hat{\beta}_1 \hat{\gamma}_1 + \xi_1}{\sqrt{k}(1 - \beta_1)\hat{\gamma}_1}\right) \|\boldsymbol{h}\|_2 \le \frac{2(\hat{\beta}_1 \hat{\gamma}_1 + \xi_1)}{\sqrt{k}(1 - \beta_1)\hat{\gamma}_1} \|\boldsymbol{X}^o\|_* + \frac{2\hat{\gamma}_1 \xi_1}{\sqrt{k}(1 - \beta_1)} \lambda.$$

Denote

$$\kappa_1 := 1 - \frac{\hat{\beta}_1 \hat{\gamma}_1 + \xi_1}{\sqrt{k}(1 - \beta_1)\hat{\gamma}_1},\tag{53}$$

it is easy to verify that $\kappa_1 > 0$ by applying the relation (44) and (45) and the fact $\xi_1 \leq \sqrt{2}\hat{\gamma}_1$. Therefore,

$$\|\boldsymbol{H}\|_{F} \leq \frac{2(\beta_{1}\hat{\gamma}_{1} + \xi_{1})}{\sqrt{k}(1 - \beta_{1})\hat{\gamma}_{1}\kappa_{1}} \|\boldsymbol{X}^{o}\|_{*} + \frac{2\hat{\gamma}_{1}\xi_{1}}{\sqrt{k}(1 - \beta_{1})\kappa_{1}}\lambda$$

and we complete the proof.

IV. CONCLUSION

In this paper, we focused on analysis of theoretical guarantees for reconstruction of low rank matrix from noisy measurements via L_{*-F} minimization. Firstly, we briefly presented the relationship of the optimal solution among several L_{*-F} minimization problems. Secondly, we gave the sufficient conditions of stable recovery and the recovery error estimation for the general constrained L_{*-F} minimization problems. Besides, we considered the robust low rank matrix recovery by regularized L_{*-F} minimization model, approximate error estimation of which was obtained under the framework of powerful RIP tools. To our knowledge, this theoretical result has not been studied before. A further issue worth considering is developing a tighter recovery error estimation for this regularized model. It is also interesting to see if the techniques in this paper can be applied in other settings.

REFERENCES

- [1] X. Chang, Y. Zhong, Y. Wang, and S. Lin, "Unified low-rank matrix estimate via penalized matrix least squares approximation," IEEE Trans. Neural Netw. Learn. Syst., vol. 30, no. 2, pp. 474-485, 2018.
- [2] E. J. Candès and T. Tao, "The power of convex relaxation: Near-optimal matrix completion," IEEE Trans. Inf. Theory, vol. 56, no. 5, pp. 2053–2080, 2010.
- [3] E. J. Candès and B. Recht, "Exact matrix completion via convex optimization," Commun. ACM, vol. 55, no. 6, pp. 111–119, 2012.
- [4] E. J. Candès, X. Li, Y. Ma, and J. Wright, "Robust principal component analysis?" J. ACM, vol. 58, no. 3, pp. 1–37, 2011.
- [5] C. Tomasi and T. Kanade, "Shape and motion from image streams: a factorization method," Proceed. National Academy of Sciences, vol. 90, no. 21, pp. 9795-9802, 1993.
- [6] R. Meka, P. Jain, C. Caramanis, and I. S. Dhillon, "Rank minimization via online learning," in Proceedings of the 25th International Conference on Machine Learning, 2008, pp. 656-663.
- [7] M. Fazel, H. Hindi, and S. P. Boyd, "A rank minimization heuristic with application to minimum order system approximation," in Proceedings of the 2001 American Control Conference.(Cat. No. 01CH37148), vol. 6. IEEE, 2001, pp. 4734-4739.
- [8] D. L. Donoho, "Compressed sensing," *IEEE Trans. Inf. Theory*, vol. 52, no. 4, pp. 1289–1306, 2006.
 [9] J.-F. Cai, E. J. Candes, and Z. Shen, "A singular value thresholding algorithm for matrix completion," *SIAM J. Optim.*, vol. 20, no. 4, pp. 1956–1982, 2010
- [10] K.-C. Toh and S. Yun, "An accelerated proximal gradient algorithm for nuclear norm regularized linear least squares problems," Pacific J. Optim., vol. 6, no. 615-640, p. 15, 2010.
- [11] D. Zhang, Y. Hu, J. Ye, X. Li, and X. He, "Matrix completion by truncated nuclear norm regularization," in 2012 IEEE Conference on Computer Vision and Pattern Recognition. IEEE, 2012, pp. 2192-2199.
- [12] S. Gu, L. Zhang, W. Zuo, and X. Feng, "Weighted nuclear norm minimization with application to image denoising," in Proceedings of the IEEE conference on computer vision and pattern recognition, 2014, pp. 2862-2869.
- [13] B. Recht, M. Fazel, and P. A. Parrilo, "Guaranteed minimum-rank solutions of linear matrix equations via nuclear norm minimization," SIAM Rev., vol. 52, no. 3, pp. 471-501, 2010.
- [14] P. D. Tao and L. H. An, "Convex analysis approach to dc programming: theory, algorithms and applications," Acta Math. Vietn., vol. 22, no. 1, pp. 289-355, 1997.
- [15] P. Gong, C. Zhang, Z. Lu, J. Huang, and J. Ye, "A general iterative shrinkage and thresholding algorithm for non-convex regularized optimization problems," in International Conference on Machine Learning. PMLR, 2013, pp. 37-45.
- [16] E. J. Candès and T. Tao, "Decoding by linear programming," IEEE Trans. Inf. Theory, vol. 51, no. 12, pp. 4203-4215, 2005.
- [17] H. Ge, W. Chen, and M. K. Ng, "New rip bounds for recovery of sparse signals with partial support information via weighted ℓ_p -minimization," *IEEE* Trans. Inf. Theory, vol. 66, no. 6, pp. 3914-3928, 2020.
- [18] M. A. Davenport and M. B. Wakin, "Analysis of orthogonal matching pursuit using the restricted isometry property," IEEE Trans. Inf. Theory, vol. 56, no. 9, pp. 4395-4401, 2010.
- [19] L.-H. Chang and J.-Y. Wu, "An improved rip-based performance guarantee for sparse signal recovery via orthogonal matching pursuit," IEEE Trans. Inf. Theory, vol. 60, no. 9, pp. 5702-5715, 2014.
- [20] E. J. Candès and Y. Plan, "Tight oracle inequalities for low-rank matrix recovery from a minimal number of random measurements," IEEE Trans. Inf. Theory, vol. 57, no. 4, pp. 2342-2359, 2011.
- [21] T. T. Cai and A. Zhang, "Sparse representation of a polytope and recovery of sparse signals and low-rank matrices," IEEE Trans. Inf. Theory, vol. 60, no. 1, pp. 122-132, 2013.
- [22] S. Tu, R. Boczar, M. Simchowitz, M. Soltanolkotabi, and B. Recht, "Low-rank solutions of linear matrix equations via procrustes flow," in International Conference on Machine Learning. PMLR, 2016, pp. 964-973.
- [23] S. Bhojanapalli, B. Neyshabur, and N. Srebro, "Global optimality of local search for low rank matrix recovery," Adv. Neural Inf. Process. Syst., vol. 29, 2016.
- [24] X. Liu, J. Hou, and J. Wang, "Robust low-rank matrix recovery fusing local-smoothness," IEEE Signal Process. Letters, vol. 29, pp. 2552–2556, 2022.

- [25] F. Krahmer and R. Ward, "New and improved johnson-lindenstrauss embeddings via the restricted isometry property," SIAM J. Math. Anal., vol. 43, no. 3, pp. 1269–1281, 2011.
- [26] T. T. Do, L. Gan, N. H. Nguyen, and T. D. Tran, "Fast and efficient compressive sensing using structurally random matrices," *IEEE Trans. Signal Process.*, vol. 60, no. 1, pp. 139–154, 2011.
- [27] Y. Cai, "Minimization of the difference of nuclear and frobenius norms for noisy low rank matrix recovery," International Journal of Wavelets, Multiresolution and Information Processing, vol. 18, no. 02, p. 1950056, 2020.
- [28] T.-H. Ma, Y. Lou, and T.-Z. Huang, "Truncated ℓ_{1−2} models for sparse recovery and rank minimization," SIAM J. Imag. Sci., vol. 10, no. 3, pp. 1346–1380, 2017.
- [29] M. Zhang, Z.-H. Huang, and Y. Zhang, "Restricted p-isometry properties of nonconvex matrix recovery," IEEE Trans. Inf. Theory, vol. 59, no. 7, pp. 4316–4323, 2013.
- [30] H. Guo, Z.-H. Huang, and X. Zhang, "Low rank matrix recovery with impulsive noise," Appl. Math. Lett., vol. 134, p. 108364, 2022.
- [31] S. Boyd, S. P. Boyd, and L. Vandenberghe, Convex Optimization. Cambridge University Press, 2004.
- [32] Q. Yao, J. T. Kwok, and X. Guo, "Fast learning with nonconvex ℓ_{1-2} regularization," arXiv preprint arXiv:1610.09461, 2016.
- [33] E. J. Candès and Y. Plan, "Tight oracle inequalities for low-rank matrix recovery from a minimal number of noisy random measurements," *IEEE Trans. Inf. Theory*, vol. 57, no. 4, pp. 2342–2359, 2011.
- [34] T. T. Cai and A. Zhang, "Sharp rip bound for sparse signal and low rank matrix recovery," Appl. Comput. Harmon. Anal., vol. 35, no. 1, pp. 74–93, 2013.