

LAPLACIAN EIGEN VALUES OF CHARACTER DEGREE GRAPHS OF SOLVABLE GROUPS

G. SIVANESAN AND C. SELVARAJ

ABSTRACT. Let G be a finite solvable group, let $Irr(G)$ be the set of all complex irreducible characters of G and let $cd(G)$ be the set of all degrees of characters in $Irr(G)$. Let $\rho(G)$ be the set of primes that divide degrees in $cd(G)$. The character degree graph $\Delta(G)$ of G is the simple undirected graph with vertex set $\rho(G)$ and in which two distinct vertices p and q are adjacent if there exists a character degree $r \in cd(G)$ such that r is divisible by the product pq . In this paper, we obtain Laplacian eigen values and distance Laplacian eigen values of regular character degree graph, super graphs of regular character degree graph and character degree graph with diameter 2 has two blocks.

1. INTRODUCTION

In this paper, G is a finite solvable group with identity 1. The set of complex irreducible characters of G is denoted by $Irr(G)$. $cd(G) = \{\chi(1) \mid \chi \in Irr(G)\}$ is the set of all distinct degrees of irreducible characters in $Irr(G)$. Let $\rho(G)$ be the set of all primes that divide degrees in $cd(G)$. An extensive literature has been devoted to studying how graphs can be associated with groups, and that literature can be used to investigate the algebraic structure of groups using graph theoretical properties. It is one of these graphs that is called the character degree graph $\Delta(G)$ of G . In fact, $\Delta(G)$ is an undirected simple graph with vertex set $\rho(G)$ in which $p, q \in \rho(G)$ are joined by an edge if there exists a character degree $\chi(1) \in cd(G)$ which is divisible by pq . It was first defined in [12] and studied by many authors (see [16], [18], [9]). There have been some interesting results obtained on the character graph of G when G is a solvable group. Actually, Manz [11] proved that $\Delta(G)$ has at most two connected components. Also, Manz et al. [13] have proved that diameter of $\Delta(G)$ is at most 3. We obtained a necessary condition for the character degree graph $\Delta(G)$ of a finite solvable group G to be Eulerian [17]. Mahdi Ebrahimi et al. [5] studied all regular character-graphs whose eigenvalues are in the interval $[-2, \infty)$. Furthermore, Hafezieh et al. studied the effects of cut vertices and eigenvalues of $\Delta(G)$ on the group structure of G [6]. This study on character degree graphs of solvable groups motivated us to study Laplacian eigenvalues and distance Laplacian eigenvalues of regular character degree graphs, super graphs of regular character degree graphs, and character degree graphs with diameter 2 has two blocks.

2020 *Mathematics Subject Classification.* 20C15, 05C50, 05C25.

Key words and phrases. character degree graph, eigenvalue, Laplacian spectrum, regular graph.

2. PRELIMINARIES

This section presents some preliminary results that are used in the paper. The graphs are assumed to be simple, undirected, and finite. Let Γ be a graph with vertex set $V(\Gamma)$ and edge set $E(\Gamma)$. If Γ is connected, then the distance $d(u, v)$ between two distinct vertices $u, v \in V(\Gamma)$ is the length of the shortest path between them. A graph's diameter is the maximum distance between all pairs of distinct vertices. A graph with n vertices in which any two distinct vertices are adjacent is denoted by K_n . C_n denotes a cycle with n vertices. The vertex connectivity $k(\Gamma)$ of Γ is defined to be the minimum number of vertices whose removal from Γ results in a disconnected subgraph of Γ . A cut vertex v of a graph Γ is a vertex such that the number of connected components of $\Gamma - v$ is more than the number of connected components of Γ . A block is a maximally connected subgraph that does not have a cut vertex. By their maximality, different blocks of Γ overlap in at most one vertex, which is then a cut vertex. A vertex's degree is defined as the number of edges incident with it, and it is denoted by $d(v)$ or $degv$. A graph Γ is called k -regular, if the degree of each vertex is k . Cliques of graphs are complete subgraphs and the number of vertices on a clique of maximum size is called its clique number and it is denoted by $\omega(\Gamma)$. The spanning tree T of an undirected graph G is a subgraph that includes all of its vertices.

For a finite simple undirected graph Γ with vertex set $V(\Gamma) = \{v_1, v_2, \dots, v_n\}$, the adjacency matrix $A(\Gamma)$ is the $n \times n$ matrix with (i, j) th entry is 1 if v_i and v_j are adjacent and 0 otherwise. We denote the diagonal matrix $D(\Gamma) = \text{diag}(d_1, d_2, \dots, d_n)$ where d_i is the degree of the vertex v_i of Γ , $i = 1, 2, \dots, n$. The Laplacian matrix $L(\Gamma)$ of Γ is the matrix $D(\Gamma) - A(\Gamma)$. The matrix $L(\Gamma)$ is symmetric and positive semidefinite, so that its eigenvalues are real and nonnegative. Furthermore, the sum of each row (column) of $L(\Gamma)$ is zero. The eigenvalues of $L(\Gamma)$ are called the Laplacian eigenvalues of Γ and it is denoted by $\lambda_1(\Gamma) \geq \lambda_2(\Gamma) \geq \dots \geq \lambda_n(\Gamma) = 0$. Now let $\lambda_{n_1}(\Gamma) \geq \lambda_{n_2}(\Gamma) \geq \dots \geq \lambda_{n_r}(\Gamma) = 0$ be the distinct eigenvalues of Γ with multiplicities m_1, m_2, \dots, m_r , respectively. The Laplacian spectrum of Γ , that is, the spectrum of $L(\Gamma)$ is represented as

$$\begin{pmatrix} \lambda_{n_1}(\Gamma) & \lambda_{n_2}(\Gamma) & \cdots & \lambda_{n_r}(\Gamma) \\ m_1 & m_2 & \cdots & m_r \end{pmatrix}.$$

The distance between two vertices $u, v \in V$ is denoted by d_{uv} and is defined as the length of a shortest path between u and v in G . The distance matrix of G is denoted by $D(G)$ and defined by $D(G) = (d_{uv})_{u, v \in V}$. The transmission $Tr(v)$ of a vertex v is defined to be the sum of the distances from v to all other vertices in G , i.e.

$$Tr(v) = \sum_{u \in V} d_{uv}.$$

The distance Laplacian of a connected graph G is $D^L(G) = Tr(G) - D(G)$, where $Tr(G)$ denotes the diagonal matrix with $Tr(v_i)$ as the i -th diagonal entry.

We will take into account the following well known facts concerning character degree graphs and they are needed in the next section.

Remark 2.1. For results regarding $\Delta(G)$, we start with Pálffy's three prime theorem on the character degree graph of solvable groups. Pálffy theorem [16,

Theorem, p. 62] states that given a solvable group G and any three distinct vertices of $\Delta(G)$, there exists an edge incident with the other two vertices. On applying Pálffy's theorem, $\Delta(G)$ has at most two connected components.

Remark 2.2. Huppert [7, p. 25] listed all possible graphs $\Delta(G)$ for solvable groups G with at most 4 vertices. In fact, every graph with 3 or few vertices that satisfies Pálffy's condition occurs as $\Delta(G)$ for some solvable group G [9, p. 184].

Theorem 2.3. [5, Lemma 2.7] *Let G be a group with $|\rho(G)| \geq 3$. If $\Delta(G)$ is not a block and the diameter of $\Delta(G)$ is at most 2, then each block of $\Delta(G)$ is a complete graph.*

Theorem 2.4. [18, Theorem 5] *The graph with four vertices in Figure 1 is not the character degree graph of a solvable group.*




FIGURE 1. Graph with four vertices

Theorem 2.5. [10, Theorem 1.1] *Let G be a solvable group. Then $\Delta(G)$ has at most one cut vertex.*

Theorem 2.6. [15, Theorem A] *If $\Delta(G)$ is a non-complete and regular character degree graph of a finite solvable group G with n vertices, then $\Gamma(G)$ is $n - 2$ is a regular graph.*

Theorem 2.7. [8, Lemma 2.1] *Let p, q_1, \dots, q_n be distinct primes so that p is odd. Then there is a solvable group G of fitting height 2 so that $\Delta(G)$ has two connected components: $\{p\}$ and $\{q_1, \dots, q_n\}$.*

Theorem 2.8. [8, Theorem A] *Let Γ be a graph with n vertices. There exists a solvable group G of fitting height 2 with $\Delta(G) = \Gamma$ if and only if the vertices of degree less than $n - 1$ can be partitioned into two subsets, each of which induces a complete subgraph of Γ and one of which contains only vertices of degree $n - 2$.*

Theorem 2.9. [14, Corollary 4.2] *The number of spanning trees of the graph G of order $n = \frac{1}{n} \lambda_2(G) \lambda_3(G) \dots \lambda_n(G)$. Here $\lambda_2(G), \lambda_3(G), \dots, \lambda_n(G)$ are Laplacian eigen values of G .*

Theorem 2.10. [1, Theorem 3.1] *Let G be a connected graph on n vertices with diameter $D \leq 2$. Let $\lambda_1^L \geq \lambda_2^L \geq \dots \geq \lambda_{n-1}^L > \lambda_n^L = 0$ be the Laplacian spectrum of G . Then the distance Laplacian spectrum of G is $2n - \lambda_{n-1}^L \geq 2n - \lambda_{n-2}^L \geq \dots \geq 2n - \lambda_1^L > \delta_n^L = 0$. Moreover, for every $i \in \{1, 2, \dots, n - 1\}$ the eigenspaces corresponding to λ_i^L and to $2n - \lambda_{n-i}^L$ are the same.*

Remark 2.11. Operation D: Consider a character degree graph $\Delta(G)$. Select two distinct primes p and q such that p is odd prime and $p, q \notin \rho(G)$. By Theorem 2.7, there exists a solvable group H of fitting height 2 and $\Delta(H)$ is the graph containing two isolated vertices. Now one can make a obtain the direct product $\Delta(G \times H)$. Given $\Delta(G)$, the construction of such a direct product $\Delta(G \times H)$ is used in the proof of Theorem 3.1 repeatedly. We denote this process as Operation D.


Definition 2.12. [2, p. 502] *Using direct product, one can find bigger groups from smaller groups. The same may be used to construct higher order character degree graphs. For two groups A and B where $\rho(A)$ and $\rho(B)$ are disjoint, we have that $\rho(A \times B) = \rho(A) \cup \rho(B)$. Define an edge between vertices p and q in $\rho(A \times B)$ if any of the following is satisfied:*

- (i) $p, q \in \rho(A)$ and there is an edge between p and q in $\Delta(A)$;
- (ii) $p, q \in \rho(B)$ and there is an edge between p and q in $\Delta(B)$;
- (iii) $p \in \rho(A)$ and $q \in \rho(B)$;
- (iv) $p \in \rho(B)$ and $q \in \rho(A)$.

Now we get a higher order character degree graph and it is called direct product.

3. LAPLACIAN EIGEN VALUES OF CHARACTER DEGREE GRAPHS

If $\Delta(G)$ is a character degree graph and u, v are two non adjacent vertices of $\Delta(G)$, then $\Delta(G) + uv$ may or may not be character degree graph. Consider the following example:



(A) Character degree graph with 6 vertices

(B) Unknown six-vertex graph

FIGURE 2. Character degree graph with addition of line

Figure 2(A) shows a character degree graph as shown in [2, p. 502-503]. Figure 2(B) cannot be concluded as a character degree graph at this time, as shown in [2, p. 509].

Interestingly, all graphs which are obtained by adding two non-adjacent vertices in a $n - 2$ regular character degree graph are character degree graphs.

Theorem 3.1. *Let $n \geq 4$ be an even integer. Let $\Delta(G)$ be the $n - 2$ regular character degree graph, then all the super graphs of $\Delta(G)$ by addition of two non adjacent vertices are character degree graphs for some solvable group G .*

Proof. Using the induction method, let us prove the result based on the number of vertices in $\Delta(G)$.

Case 1. Let $n = 4$.

Figure 3 shows character degree graphs as shown in [9, p. 184]. Figures 3(B) and 3(C) are supergraphs of Figure 3(A). By adding two non-adjacent vertices to a 2-regular graph with four vertices, super graphs are formed.

Case 2. Let $n = 6$.

The graphs in Figure 4 are character degree graphs as shown in [2, p. 502-503]. It is important to note that Figures 4(B) through Figure 4(D) are super




FIGURE 3. Character degree graphs with four vertices




FIGURE 4. Character degree graphs with six vertices

graphs of Figure 4(A). Supergraphs are created by adding two non-adjacent vertices to four regular graphs with six vertices.

Case 3. Let $n = 8$. Through direct product construction, we can construct a six-regular graph with eight vertices and all of its supergraphs by adding two adjacent vertices.

As in Case 3, there are four character degree graphs with six vertices. Let us consider each of these graphs as a $\Delta(G)$. After applying Operation D 2.11, we

get four direct products $\Delta(G \times H)$ with 8 vertices. First, we have a 6-regular graph with 8 vertices. In the second graph, two vertices have a degree of 7 and six vertices have a degree of 6. On the third graph, four of the vertices are of degree 7 while the remaining four are of degree 6. The fourth graph contains six vertices of degree 7 and two vertices of degree 6. The complete graph occurs as $\Delta(G)$ for some solvable group G [3, p. 39]. As a result, a complete graph with 8 vertices is also a character degree graph. Thus, all the supergraphs of the 6-regular graph with 8 vertices are character degree graphs.

Case 4. Let $n = 10$. Four supergraphs of 6-regular graphs with 8 vertices and one 6-regular graph are constructed in case 3. We can take each of these graphs as a $\Delta(G)$. After applying Operation D 2.11, we obtain five direct products $\Delta(G \times H)$ with 10 vertices. An 8-regular graph with 10 vertices is the first graph. There are two vertices of degree 9 and eight vertices of degree 8 in the second graph. The third graph has four vertices of degree 9 and six vertices of degree 8. In the fourth graph, there are six vertices of degree 9 and four of degree 8. The fifth graph has eight vertices of degree 9 and two vertices of degree 8. The complete graph with 10 vertices is also known as the character degree graph. All supergraphs of the 8-regular graph with 10 vertices are character degree graphs.

Assume that as induction hypothesis, there exists $\frac{n-2}{2}$ super graphs of $n-4$ regular graph with $n-2$ vertices and one $n-4$ regular graph with $n-2$ vertices. Let us take each of these graphs as $\Delta(G)$. By applying Operation D 2.11, we obtain $\frac{n}{2}$ direct products $\Delta(G \times H)$ with n vertices. These $\frac{n}{2}$ direct product graphs are super graphs of $n-2$ regular graph with n vertices. As stated in [3, p. 39], the complete graph with n vertices is also a character degree graph. Therefore, all super graphs of $n-2$ regular graph with n vertices are character degree graphs. \square

Remark 3.2. *Each vertex in an $n-2$ regular graph is not adjacent to another vertex. Let us say that the vertex v_i ($1 \leq i \leq \frac{n}{2}$) is not adjacent to the vertex $v_{i+\frac{n}{2}}$. Let $\Delta(G)$ be the supergraph of a $n-2$ regular graph with n vertices. Initially, we label the vertices of $\Delta(G)$ as $\{v_1, v_2, v_3, \dots, v_{n-1}, v_n\}$. Assume that m is the number of vertices having a degree of $n-1$. Remaining $n-m$ vertices have degree $n-2$ can be partitioned into two subsets. Let $r = \frac{n-m}{2}$. There is no difficulty in finding r vertices that are adjacent to each other. As a result, it forms a complete subgraph of $\Delta(G)$. Let us refer to this vertex set as $U = \{u_1, \dots, u_r\}$. Each vertex in U has a degree $n-2$. Thus, for each vertex in U , there is correspondingly one vertex which is not adjacent to that vertex. We shall label this set of vertex values as $W = \{w_1, \dots, w_r\}$. In W , each vertex has a degree of $n-2$. Therefore, all the vertices in W form a complete subgraph of $\Delta(G)$. Based on Theorem 2.8, G is a solvable group of fitting height 2 for supergraphs of $n-2$ regular character degree graph. \square*

The next step is to find the Laplacian eigenvalues and multiplicities of an $n-2$ regular graph.

Theorem 3.3. *Let $n \geq 4$ be an even integer and G be a solvable group such that $\Delta(G)$ is $n-2$ regular graph. The characteristic polynomial of the Laplacian matrix of $\Delta(G)$ is given by $x(x - (n-2))^{\frac{n}{2}}(x-n)^{\frac{n}{2}-1}$.*

Proof. The Laplacian matrix of $\Delta(G)$ is $n \times n$ matrix. Each row and column is indexed based on its vertices $v_1, v_2, v_3, \dots, v_{\frac{n}{2}}, v_{\frac{n}{2}+1}, \dots, v_n$. In $n-2$ regular graph, each vertex is not adjacent with one vertex. Let's say the vertex v_i ($1 \leq i \leq \frac{n}{2}$) is not adjacent with the vertex $v_{i+\frac{n}{2}}$. Then

$$L(\Delta(G)) = \begin{bmatrix} n-2 & -1 & -1 \cdots -1 & 0 & -1 & -1 \cdots & -1 \\ -1 & n-2 & -1 \cdots -1 & -1 & 0 & -1 \cdots & -1 \\ -1 & -1 & n-2 \cdots -1 & -1 & -1 & 0 \cdots & -1 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ 0 & -1 & -1 \cdots -1 & n-2 & -1 & -1 \cdots & -1 \\ -1 & 0 & -1 \cdots -1 & -1 & n-2 & -1 \cdots & -1 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ -1 & -1 & -1 \cdots 0 & -1 & -1 & -1 \cdots & n-2 \end{bmatrix}$$

The characteristic polynomial of $L(\Delta(G))$ is $|L(\Delta(G) - xI)|$

$$= \begin{vmatrix} (n-2) - x & -1 & -1 \cdots -1 & 0 & -1 & -1 \cdots & -1 \\ -1 & (n-2) - x & -1 \cdots -1 & -1 & 0 & -1 \cdots & -1 \\ -1 & -1 & (n-2) - x \cdots -1 & -1 & -1 & 0 \cdots & -1 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ 0 & -1 & -1 \cdots -1 & (n-2) - x & -1 & -1 \cdots & -1 \\ -1 & 0 & -1 \cdots -1 & -1 & (n-2) - x & -1 \cdots & -1 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ -1 & -1 & -1 \cdots 0 & -1 & -1 & -1 \cdots & (n-2) - x \end{vmatrix}$$

$$= \begin{vmatrix} -x & -x & -x \cdots -x & -x & -x & -x \cdots & -x \\ -1 & (n-2) - x & -1 \cdots -1 & -1 & 0 & -1 \cdots & -1 \\ -1 & -1 & (n-2) - x \cdots -1 & -1 & -1 & 0 \cdots & -1 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ 0 & -1 & -1 \cdots -1 & (n-2) - x & -1 & -1 \cdots & -1 \\ -1 & 0 & -1 \cdots -1 & -1 & (n-2) - x & -1 \cdots & -1 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ -1 & -1 & -1 \cdots 0 & -1 & -1 & -1 \cdots & (n-2) - x \end{vmatrix}$$

(Apply the row operation $R_1 \rightarrow R_1 + R_2 + R_3 + \dots + R_n$)

$$= (-x) \begin{vmatrix} 1 & 1 & 1 \cdots 1 & 1 & 1 & 1 \cdots & 1 \\ -1 & (n-2) - x & -1 \cdots -1 & -1 & 0 & -1 \cdots & -1 \\ -1 & -1 & (n-2) - x \cdots -1 & -1 & -1 & 0 \cdots & -1 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ 0 & -1 & -1 \cdots -1 & (n-2) - x & -1 & -1 \cdots & -1 \\ -1 & 0 & -1 \cdots -1 & -1 & (n-2) - x & -1 \cdots & -1 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ -1 & -1 & -1 \cdots 0 & -1 & -1 & -1 \cdots & (n-2) - x \end{vmatrix}$$

$$= (-x) \begin{vmatrix} 1 & 1 & 1 \cdots 1 & 1 & 1 & 1 \cdots & 1 \\ 0 & (n-1) - x & 0 \cdots 0 & 0 & 1 & 0 \cdots & 0 \\ 0 & 0 & (n-1) - x \cdots 0 & 0 & 0 & 1 \cdots & 0 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ 0 & -1 & -1 \cdots -1 & (n-2) - x & -1 & -1 \cdots & -1 \\ 0 & 1 & 0 \cdots 0 & 0 & (n-1) - x & 0 \cdots & 0 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ 0 & 0 & 0 \cdots 1 & 0 & 0 & 0 \cdots & (n-1) - x \end{vmatrix}$$

(Apply row operations:

For $2 \leq i \leq \frac{n}{2}$, $R_i \rightarrow R_1 + R_i$

For $\frac{n}{2} + 2 \leq i \leq n$, $R_i \rightarrow R_1 + R_i$)

$$= (-x) \begin{vmatrix} 1 & 1 & 1 \cdots 1 & 1 & 1 & 1 \cdots & 1 \\ 0 & (n-1) - x & 0 \cdots 0 & 0 & 1 & 0 \cdots & 0 \\ 0 & 0 & (n-1) - x \cdots 0 & 0 & 0 & 1 \cdots & 0 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ 0 & 0 & 0 \cdots 0 & (n-2) - x & (n-2) - x & (n-2) - x \cdots & (n-2) - x \\ 0 & 1 & 0 \cdots 0 & 0 & (n-1) - x & 0 \cdots & 0 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ 0 & 0 & 0 \cdots 1 & 0 & 0 & 0 \cdots & (n-1) - x \end{vmatrix}$$

(Apply the row operation : $R_{\frac{n}{2}+1} \rightarrow R_{\frac{n}{2}+1} + \dots + R_n$)

$$= (-x)((n-2)-x) \begin{vmatrix} 1 & 1 & 1 \cdots 1 & 1 & 1 & 1 \cdots & 1 \\ 0 & (n-1) - x & 0 \cdots 0 & 0 & 1 & 0 \cdots & 0 \\ 0 & 0 & (n-1) - x \cdots 0 & 0 & 0 & 1 \cdots & 0 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ 0 & 0 & 0 \cdots 0 & 1 & 1 & 1 \cdots & 1 \\ 0 & 1 & 0 \cdots 0 & 0 & (n-1) - x & 0 \cdots & 0 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ 0 & 0 & 0 \cdots 1 & 0 & 0 & 0 \cdots & (n-1) - x \end{vmatrix}$$

(Apply the row operation : $R_{\frac{n}{2}+1} \rightarrow R_{\frac{n}{2}+1}/((n-2)-x)$)

$$= (-x)((n-2)-x) \begin{vmatrix} 1 & 1 & 1 \cdots 1 & 1 & 1 & 1 \cdots & 1 \\ 0 & n-x & 0 \cdots 0 & 0 & n-x & 0 \cdots & 0 \\ 0 & 0 & n-x \cdots 0 & 0 & 0 & n-x \cdots & 0 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ 0 & 0 & 0 \cdots 0 & 1 & 1 & 1 \cdots & 1 \\ 0 & 1 & 0 \cdots 0 & 0 & (n-1)-x & 0 \cdots & 0 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ 0 & 0 & 0 \cdots 1 & 0 & 0 & 0 \cdots & (n-1)-x \end{vmatrix}$$

(Apply Row operation: For $2 \leq i \leq \frac{n}{2}$, $R_i \rightarrow R_i + R_{\frac{n}{2}+i}$)

$$= (-x)((n-2)-x)(n-x)^{\frac{n}{2}-1} \begin{vmatrix} 1 & 1 & 1 \cdots 1 & 1 & 1 & 1 \cdots & 1 \\ 0 & 1 & 0 \cdots 0 & 0 & 1 & 0 \cdots & 0 \\ 0 & 0 & 1 \cdots 0 & 0 & 0 & 1 \cdots & 0 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ 0 & 0 & 0 \cdots 0 & 1 & 1 & 1 \cdots & 1 \\ 0 & 1 & 0 \cdots 0 & 0 & (n-1)-x & 0 \cdots & 0 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ 0 & 0 & 0 \cdots 1 & 0 & 0 & 0 \cdots & (n-1)-x \end{vmatrix}$$

(Apply Row operation: For $2 \leq i \leq \frac{n}{2}$, $R_i \rightarrow R_i/(n-x)$)

$$= (-x)((n-2)-x)(n-x)^{\frac{n}{2}-1} \begin{vmatrix} 1 & 1 & 1 \cdots 1 & 1 & 1 & 1 \cdots & 1 \\ 0 & 1 & 0 \cdots 0 & 0 & 1 & 0 \cdots & 0 \\ 0 & 0 & 1 \cdots 0 & 0 & 0 & 1 \cdots & 0 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ 0 & 0 & 0 \cdots 0 & 1 & 1 & 1 \cdots & 1 \\ 0 & 0 & 0 \cdots 0 & 0 & (n-2)-x & 0 \cdots & 0 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ 0 & 0 & 0 \cdots 0 & 0 & 0 & 0 \cdots & (n-2)-x \end{vmatrix}$$

(Apply the row operation: For $2 \leq i \leq \frac{n}{2}$, $R_{\frac{n}{2}+i} \rightarrow R_{\frac{n}{2}+i} - R_i$)

According to Schur's decomposition theorem [4], we have

$$|L(\Delta(G) - xI)| = (x)((n-2) - x)^{\frac{n}{2}}(n-x)^{\frac{n}{2}-1}.$$

So the characteristic polynomial is $(x)(x - (n-2))^{\frac{n}{2}}(x - (n))^{\frac{n}{2}-1}$. This completes the proof.

Corollary 3.4. *Let $n \geq 4$ be an even integer, the Laplacian spectrum of $\Delta(G)$ is given by $\begin{pmatrix} 0 & n-2 & n \\ 1 & \frac{n}{2} & \frac{n}{2}-1 \end{pmatrix}$.*

Based on [1, Theorem 3.1] we have the following corollary

Corollary 3.5. *Let $n \geq 4$ be an even integer, the distance Laplacian spectrum of $\Delta(G)$ is given by $\begin{pmatrix} 0 & n+2 & n \\ 1 & \frac{n}{2} & \frac{n}{2}-1 \end{pmatrix}$.*

According to [14, Theorem 2.8], the following corollary follows:

Corollary 3.6. *Let $n \geq 4$ be an even integer, the number of spanning trees of $\Delta(G)$ is $(n-2)^{\frac{n}{2}}(n)^{\frac{n}{2}-1}$.*

Theorem 3.7. *Let $n \geq 4$ be an even integer and G be a solvable group such that $\Delta(G)$ is $n-2$ regular graph. Let $\Delta(G_1)$ be the super character degree graph of $\Delta(G)$ obtained by the addition of n_1 edges ($1 \leq n_1 \leq \frac{n}{2}$). The characteristic polynomial of the Laplacian matrix of $\Delta(G_1)$ is given by $x(x - (n-2))^{\frac{n}{2}-n_1}(x-n)^{(\frac{n}{2}+n_1)-1}$.*

Proof. The Laplacian matrix of $\Delta(G)$ is an $n \times n$ matrix. The rows and columns are indexed according to the vertices $v_1, v_2, \dots, v_{n_1+1}, v_{n_1+2}, \dots, v_{\frac{n}{2}+1}, \dots, v_{\frac{n}{2}+n_1+1}, \dots, v_n$. In $n-2$ regular graph, each vertex is not adjacent with one vertex. Let us say that the vertex v_i ($1 \leq i \leq \frac{n}{2}$) is not adjacent with the vertex $v_{i+\frac{n}{2}}$. Let n_1 ($1 \leq n_1 \leq \frac{n}{2}$) be the number of edges added to $\Delta(G)$. That is, For $1 \leq i \leq n_1$, each v_i is adjacent with $v_{\frac{n}{2}+i}$. Then

$$L(\Delta(G_1)) = \begin{bmatrix} n-1 & -1 \cdots -1 & -1 \cdots & -1 \cdots & -1 \cdots & -1 \\ -1 & n-1 \cdots -1 & -1 \cdots & -1 \cdots & -1 \cdots & -1 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ -1 & -1 \cdots n-2 & -1 \cdots & -1 \cdots & 0 \cdots & -1 \\ -1 & -1 \cdots -1 & n-2 \cdots & -1 \cdots & -1 \cdots & -1 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ -1 & -1 \cdots -1 & -1 \cdots & n-1 \cdots & -1 \cdots & -1 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ -1 & -1 \cdots 0 & -1 \cdots & -1 \cdots & n-2 \cdots & -1 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ -1 & -1 \cdots -1 & -1 \cdots & -1 \cdots & -1 \cdots & n-2 \end{bmatrix}$$

So that the characteristic polynomial of $L(\Delta(G_1))$ is $|L(\Delta(G_1) - xI)|$

$$= \begin{vmatrix} (n-1) - x & -1 \cdots -1 & -1 \cdots & -1 \cdots & -1 \cdots & -1 \\ -1 & (n-1) - x \cdots -1 & -1 \cdots & -1 \cdots & -1 \cdots & -1 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ -1 & -1 \cdots (n-2) - x & -1 \cdots & -1 \cdots & 0 \cdots & -1 \\ -1 & -1 \cdots -1 & (n-2) - x \cdots & -1 \cdots & -1 \cdots & -1 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ -1 & -1 \cdots -1 & -1 \cdots & (n-1) - x \cdots & -1 \cdots & -1 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ -1 & -1 \cdots 0 & -1 \cdots & -1 \cdots & (n-2) - x \cdots & -1 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ -1 & -1 \cdots -1 & -1 \cdots & -1 \cdots & -1 \cdots & (n-2) - x \end{vmatrix}$$

$$= \begin{vmatrix} -x & -x \cdots -x & -x \cdots & -x \cdots & -x \cdots & -x \\ -1 & (n-1) - x \cdots - 1 & -1 \cdots & -1 \cdots & -1 \cdots & -1 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ -1 & -1 \cdots (n-2) - x & -1 \cdots & -1 \cdots & 0 \cdots & -1 \\ -1 & -1 \cdots - 1 & (n-2) - x \cdots & -1 \cdots & -1 \cdots & -1 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ -1 & -1 \cdots - 1 & -1 \cdots & (n-1) - x \cdots & -1 \cdots & -1 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ -1 & -1 \cdots 0 & -1 \cdots & -1 \cdots & (n-2) - x \cdots & -1 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ -1 & -1 \cdots - 1 & -1 \cdots & -1 \cdots & -1 \cdots & (n-2) - x \end{vmatrix}$$

(Apply the row operation: $R_1 \rightarrow R_1 + R_2 + R_3 + \dots + R_n$)

$$= (-x) \begin{vmatrix} 1 & 1 \cdots 1 & 1 \cdots & 1 \cdots & 1 \cdots & 1 \\ -1 & (n-1) - x \cdots - 1 & -1 \cdots & -1 \cdots & -1 \cdots & -1 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ -1 & -1 \cdots (n-2) - x & -1 \cdots & -1 \cdots & 0 \cdots & -1 \\ -1 & -1 \cdots - 1 & (n-2) - x \cdots & -1 \cdots & -1 \cdots & -1 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ -1 & -1 \cdots - 1 & -1 \cdots & (n-1) - x \cdots & -1 \cdots & -1 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ -1 & -1 \cdots 0 & -1 \cdots & -1 \cdots & (n-2) - x \cdots & -1 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ -1 & -1 \cdots - 1 & -1 \cdots & -1 \cdots & -1 \cdots & (n-2) - x \end{vmatrix}$$

$$= (-x) \begin{vmatrix} 1 & 1 \cdots 1 & 1 \cdots & 1 \cdots & 1 \cdots & 1 \\ 0 & n - x \cdots 0 & 0 \cdots & 0 \cdots & 0 \cdots & 0 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ 0 & 0 \cdots (n-1) - x & 0 \cdots & 0 \cdots & 1 \cdots & 0 \\ 0 & 0 \cdots 0 & (n-1) - x \cdots & 0 \cdots & 0 \cdots & 0 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ 0 & 0 \cdots 0 & 0 \cdots & n - x \cdots & 0 \cdots & 0 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ 0 & 0 \cdots 1 & 0 \cdots & 0 \cdots & (n-1) - x \cdots & 0 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ 0 & 0 \cdots 0 & 0 \cdots & 0 \cdots & 0 \cdots & (n-1) - x \end{vmatrix}$$

(Apply the row operations: For $2 \leq i \leq n$, $R_i \rightarrow R_1 + R_i$)

$$= (-x) (n-x)^{(2n_1-1)} \begin{vmatrix} 1 & 1 \dots 1 & 1 \dots & 1 \dots & 1 \dots & 1 \\ 0 & 1 \dots 0 & 0 \dots & 0 \dots & 0 \dots & 0 \\ \vdots & \dots & \dots & \dots & \dots & \vdots \\ 0 & 0 \dots (n-1) - x & 0 \dots & 0 \dots & 1 \dots & 0 \\ 0 & 0 \dots 0 & (n-1) - x \dots & 0 \dots & 0 \dots & 0 \\ \vdots & \dots & \dots & \dots & \dots & \vdots \\ 0 & 0 \dots 0 & 0 \dots & n-x \dots & 0 \dots & 0 \\ \vdots & \dots & \dots & \dots & \dots & \vdots \\ 0 & 0 \dots 1 & 0 \dots & 0 \dots & (n-1) - x \dots & 0 \\ \vdots & \dots & \dots & \dots & \dots & \vdots \\ 0 & 0 \dots 0 & 0 \dots & 0 \dots & 0 \dots & (n-1) - x \end{vmatrix}$$

(Apply the row operation:

For $2 \leq i \leq n_1$, $R_i \rightarrow R_i/(n-x)$)

$$= (-x) (n-x)^{(2n_1-1)} \begin{vmatrix} 1 & 1 \dots 1 & 1 \dots & 1 \dots & 1 \dots & 1 \\ 0 & 1 \dots 0 & 0 \dots & 0 \dots & 0 \dots & 0 \\ \vdots & \dots & \dots & \dots & \dots & \vdots \\ 0 & 0 \dots (n-1) - x & 0 \dots & 0 \dots & 1 \dots & 0 \\ 0 & 0 \dots 0 & (n-1) - x \dots & 0 \dots & 0 \dots & 0 \\ \vdots & \dots & \dots & \dots & \dots & \vdots \\ 0 & 0 \dots 0 & 0 \dots & 1 \dots & 0 \dots & 0 \\ \vdots & \dots & \dots & \dots & \dots & \vdots \\ 0 & 0 \dots 1 & 0 \dots & 0 \dots & (n-1) - x \dots & 0 \\ \vdots & \dots & \dots & \dots & \dots & \vdots \\ 0 & 0 \dots 0 & 0 \dots & 0 \dots & 0 \dots & (n-1) - x \end{vmatrix}$$

(Apply the row operation:

For $\frac{n}{2} + 1 \leq i \leq \frac{n}{2} + n_1$, $R_i \rightarrow R_i/(n-x)$)

$$= (-x) (n-x)^{(2n_1-1)} (n-x)^{\left(\frac{n}{2}-n_1\right)} \begin{vmatrix} 1 & 1 \dots 1 & 1 \dots & 1 \dots & 1 \dots & 1 \\ 0 & 1 \dots 0 & 0 \dots & 0 \dots & 0 \dots & 0 \\ \vdots & \dots & \dots & \dots & \dots & \vdots \\ 0 & 0 \dots 1 & 0 \dots & 0 \dots & 1 \dots & 0 \\ 0 & 0 \dots 0 & 1 \dots & 0 \dots & 0 \dots & 0 \\ \vdots & \dots & \dots & \dots & \dots & \vdots \\ 0 & 0 \dots 0 & 0 \dots & 1 \dots & 0 \dots & 0 \\ \vdots & \dots & \dots & \dots & \dots & \vdots \\ 0 & 0 \dots 1 & 0 \dots & 0 \dots & (n-1) - x \dots & 0 \\ \vdots & \dots & \dots & \dots & \dots & \vdots \\ 0 & 0 \dots 0 & 0 \dots & 0 \dots & 0 \dots & (n-1) - x \end{vmatrix}$$

(Apply the row operations consecutively,
 For $n_1 + 1 \leq i \leq \frac{n}{2}$, $R_i \rightarrow R_i + R_{\frac{n}{2}+i}$
 For $n_1 + 1 \leq i \leq \frac{n}{2}$, $R_i \rightarrow R_i/(n-x)$)

$$= (-x)(n-x)^{\left(\frac{n}{2}+n_1\right)-1} \begin{vmatrix} 1 & 1 \cdots 1 & 1 \cdots & 1 \cdots & 1 \cdots & 1 \\ 0 & 1 \cdots 0 & 0 \cdots & 0 \cdots & 0 \cdots & 0 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ 0 & 0 \cdots 1 & 0 \cdots & 0 \cdots & 1 \cdots & 0 \\ 0 & 0 \cdots 0 & 1 \cdots & 0 \cdots & 0 \cdots & 0 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ 0 & 0 \cdots 0 & 0 \cdots & 1 \cdots & 0 \cdots & 0 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ 0 & 0 \cdots 1 & 0 \cdots & 0 \cdots & (n-1) - x \cdots & 0 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ 0 & 0 \cdots 0 & 0 \cdots & 0 \cdots & 0 \cdots & (n-1) - x \end{vmatrix}$$

$$= (-x)(n-x)^{\left(\frac{n}{2}+n_1\right)-1} \begin{vmatrix} 1 & 1 \cdots 1 & 1 \cdots & 1 \cdots & 1 \cdots & 1 \\ 0 & 1 \cdots 0 & 0 \cdots & 0 \cdots & 0 \cdots & 0 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ 0 & 0 \cdots 1 & 0 \cdots & 0 \cdots & 1 \cdots & 0 \\ 0 & 0 \cdots 0 & 1 \cdots & 0 \cdots & 0 \cdots & 0 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ 0 & 0 \cdots 0 & 0 \cdots & 1 \cdots & 0 \cdots & 0 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ 0 & 0 \cdots 0 & 0 \cdots & 0 \cdots & (n-2) - x \cdots & 0 \\ \vdots & \cdots & \cdots & \cdots & \cdots & \vdots \\ 0 & 0 \cdots 0 & 0 \cdots & 0 \cdots & 0 \cdots & (n-2) - x \end{vmatrix}$$

(Apply the row operation: For $n_1 + 1 \leq i \leq \frac{n}{2}$, $R_{\frac{n}{2}+i} \rightarrow R_{\frac{n}{2}+i} - R_i$)

By using Schur's decomposition theorem [4], we have

$$|L(\Delta(G_1) - xI)| = (-x)((n-2) - x)^{\frac{n}{2}-n_1}(n-x)^{\left(\frac{n}{2}+n_1\right)-1}.$$

So the characteristic polynomial is $x(x - (n-2))^{\frac{n}{2}-n_1}(x - (n))^{\left(\frac{n}{2}+n_1\right)-1}$ and hence the result holds.

Corollary 3.8. *Let $n \geq 4$ be an even integer, the Laplacian spectrum of $\Delta(G_1)$ is given by $\begin{pmatrix} 0 & n-2 & n \\ 1 & \frac{n}{2} - n_1 & (\frac{n}{2} + n_1) - 1 \end{pmatrix}$.*

[1, Theorem 3.1] demonstrates the following corollary:

Corollary 3.9. *Let $n \geq 4$ be an even integer, the distance Laplacian spectrum of $\Delta(G_1)$ is given by $\begin{pmatrix} 0 & n+2 & n \\ 1 & \frac{n}{2} - n_1 & (\frac{n}{2} + n_1) - 1 \end{pmatrix}$.*

[14, Theorem 2.8] implies following corollary:

Corollary 3.10. *Let $n \geq 4$ be an even integer, the number of spanning trees of $\Delta(G_1)$ is $(n-2)^{\binom{n}{2}-n_1} (n)^{\binom{n}{2}+n_1-1}$.*

Theorem 3.11. *Let $n \geq 3$ be an integer and G be a solvable group. Let $\Delta(G)$ be a character degree graph with diameter 2 and it is not a block. $\Delta(G)$ has the structure: $\Delta(G) = K_{n-(n_1+1)} - v - K_{n_1}$, for some integer $1 \leq n_1 \leq n - (n_1 + 1)$, where $V(K_{n-(n_1+1)}) \cup \{v\}$ and $V(K_{n_1}) \cup \{v\}$ generates a clique. The characteristic polynomial of $L(\Delta(G))$ is given by $(x)(x-1)(x-n)(x-(n_1+1))^{n_1-1}(x-(n-n_1))^{n-(n_1+2)}$.*

Proof. The Laplacian matrix of $\Delta(G)$ is $n \times n$ matrix. The rows and columns are indexed in order by the vertices $v_1, v_2, \dots, v_{n_1}, v_{n_1+1}, v_{n_1+2}, \dots, v_n$. Complete graph K_{n_1} is not adjacent with any vertex of $K_{n-(n_1+1)}$ and vice versa.

$$L(\Delta(G)) = \begin{bmatrix} n_1 & -1 \cdots -1 & -1 & 0 \cdots & 0 \\ -1 & n_1 \cdots -1 & -1 & 0 \cdots & 0 \\ \vdots & \cdots & \cdots & \vdots & \\ -1 & -1 \cdots n_1 & -1 & 0 \cdots & 0 \\ -1 & -1 \cdots -1 & n-1 & -1 \cdots & -1 \\ 0 & 0 \cdots 0 & -1 & n-(n_1+1) \cdots & -1 \\ \vdots & \cdots & \cdots & \cdots & \vdots \\ 0 & 0 \cdots 0 & -1 & -1 \cdots & n-(n_1+1) \end{bmatrix}$$

The characteristic polynomial of $L(\Delta(G))$ is $|L(\Delta(G) - xI)|$

$$= \begin{vmatrix} n_1 - x & -1 \cdots -1 & -1 & 0 \cdots & 0 \\ -1 & n_1 - x \cdots -1 & -1 & 0 \cdots & 0 \\ \vdots & \cdots & \cdots & \vdots & \\ -1 & -1 \cdots n_1 - x & -1 & 0 \cdots & 0 \\ -1 & -1 \cdots -1 & (n-1) - x & -1 \cdots & -1 \\ 0 & 0 \cdots 0 & -1 & (n-(n_1+1)) - x \cdots & -1 \\ \vdots & \cdots & \cdots & \cdots & \vdots \\ 0 & 0 \cdots 0 & -1 & -1 \cdots & (n-(n_1+1)) - x \end{vmatrix}$$

$$= \begin{vmatrix} -x & -x \cdots -x & -x & -x \cdots & -x \\ -1 & n_1 - x \cdots -1 & -1 & 0 \cdots & 0 \\ \vdots & \cdots & \cdots & \vdots & \\ -1 & -1 \cdots n_1 - x & -1 & 0 \cdots & 0 \\ -1 & -1 \cdots -1 & (n-1) - x & -1 \cdots & -1 \\ 0 & 0 \cdots 0 & -1 & (n-(n_1+1)) - x \cdots & -1 \\ \vdots & \cdots & \cdots & \cdots & \vdots \\ 0 & 0 \cdots 0 & -1 & -1 \cdots & (n-(n_1+1)) - x \end{vmatrix}$$

(Apply the row operation: $R_1 \rightarrow R_1 + R_2 + R_3 + \dots + R_n$)

$$= (-x) \left| \begin{array}{cccccc} 1 & 1 \cdots 1 & 1 & 1 \cdots & 1 \\ -1 & n_1 - x \cdots - 1 & -1 & 0 \cdots & 0 \\ \vdots & \cdots & \cdots & \vdots & \\ -1 & -1 \cdots n_1 - x & -1 & 0 \cdots & 0 \\ -1 & -1 \cdots - 1 & (n-1) - x & -1 \cdots & -1 \\ 0 & 0 \cdots 0 & -1 & (n - (n_1 + 1)) - x \cdots & -1 \\ \vdots & \cdots & \cdots & \cdots & \vdots \\ 0 & 0 \cdots 0 & -1 & -1 \cdots & (n - (n_1 + 1)) - x \end{array} \right|$$

$$= (-x)(n-x) \left| \begin{array}{cccccc} 1 & 1 \cdots 1 & 1 & 1 \cdots & 1 \\ 0 & (n_1 + 1) - x \cdots 0 & 0 & 1 \cdots & 1 \\ \vdots & \cdots & \cdots & \vdots & \\ 0 & 0 \cdots (n_1 + 1) - x & 0 & 1 \cdots & 1 \\ 0 & 0 \cdots 0 & 1 & 0 \cdots & 0 \\ 0 & 0 \cdots 0 & -1 & (n - (n_1 + 1)) - x \cdots & -1 \\ \vdots & \cdots & \cdots & \vdots & \\ 0 & 0 \cdots 0 & -1 & -1 \cdots & (n - (n_1 + 1)) - x \end{array} \right|$$

(Apply the row operations consecutively,

For $2 \leq i \leq n_1 + 1$, $R_i \rightarrow R_1 + R_i$

$R_{n_1+1} \rightarrow R_{n_1+1}/(n-x)$

$$= (-x)(n-x) \left| \begin{array}{cccccc} 1 & 1 \cdots 1 & 1 & 1 \cdots & 1 \\ 0 & (n_1 + 1) - x \cdots 0 & 0 & 1 \cdots & 1 \\ \vdots & \cdots & \cdots & \vdots & \\ 0 & 0 \cdots (n_1 + 1) - x & 0 & 1 \cdots & 1 \\ 0 & 0 \cdots 0 & 1 & 0 \cdots & 0 \\ 0 & 0 \cdots 0 & 0 & (n - (n_1 + 1)) - x \cdots & -1 \\ \vdots & \cdots & \cdots & \vdots & \\ 0 & 0 \cdots 0 & 0 & -1 \cdots & (n - (n_1 + 1)) - x \end{array} \right|$$

(Apply the row operation: For $n_1 + 2 \leq i \leq n$, $R_i \rightarrow R_{n_1+1} + R_i$)

$$= (-x)(n-x)(1-x) \left| \begin{array}{cccccc} 1 & 1 \cdots 1 & 1 & 1 \cdots & 1 \\ 0 & (n_1 + 1) - x \cdots 0 & 0 & 1 \cdots & 1 \\ \vdots & \cdots & \cdots & \vdots & \\ 0 & 0 \cdots (n_1 + 1) - x & 0 & 1 \cdots & 1 \\ 0 & 0 \cdots 0 & 1 & 0 \cdots & 0 \\ 0 & 0 \cdots 0 & 0 & 1 \cdots & 1 \\ \vdots & \cdots & \cdots & \vdots & \\ 0 & 0 \cdots 0 & 0 & -1 \cdots & (n - (n_1 + 1)) - x \end{array} \right|$$

(Apply the row operations consecutively,
 $R_{n_1+2} \rightarrow R_{n_1+2} + R_{n_1+3} + \dots + R_n$
 $R_{n_1+2} \rightarrow R_{n_1+2}/(1-x)$)

$$= (-x)(n-x)(1-x) \begin{vmatrix} 1 & 1 \cdots 1 & 1 & 1 \cdots & 1 \\ 0 & (n_1+1) - x \cdots 0 & 0 & 1 \cdots & 1 \\ \vdots & \cdots & \cdots & \vdots & \\ 0 & 0 \cdots (n_1+1) - x & 0 & 1 \cdots & 1 \\ 0 & 0 \cdots 0 & 1 & 0 \cdots & 0 \\ 0 & 0 \cdots 0 & 0 & 1 \cdots & 1 \\ \vdots & \cdots & \cdots & \vdots & \\ 0 & 0 \cdots 0 & 0 & 0 \cdots & (n-n_1) - x \end{vmatrix}$$

(Apply the row operation: For $n_1+2 \leq i \leq n$, $R_i \rightarrow R_i + R_{n_1+2}$)

By using Schur's decomposition theorem [4], we have

$|L(\Delta(G) - xI)| = (-x)(n-x)(1-x)((n_1+1) - x)^{n_1-1}((n-n_1) - x)^{n-(n_1+2)}$.
 So the characteristic polynomial is
 $(x)(x-1)(x-n)(x-(n_1+1))^{n_1-1}(x-(n-n_1))^{n-(n_1+2)}$ and hence the result holds.

Corollary 3.12. *Let $n \geq 3$ be an integer, the Laplacian spectrum of $\Delta(G)$ is given by $\begin{pmatrix} 0 & 1 & n & n_1+1 & n-n_1 \\ 1 & 1 & 1 & n_1-1 & n-(n_1+2) \end{pmatrix}$.*

According to [1, Theorem 3.1] the following corollary follows:

Corollary 3.13. *Let $n \geq 3$ be an integer, the distance Laplacian spectrum of $\Delta(G)$ is given by $\begin{pmatrix} 0 & 2n-1 & n & 2n-(n_1+1) & n+n_1 \\ 1 & 1 & 1 & n_1-1 & n-(n_1+2) \end{pmatrix}$.*

From [14, Theorem 2.8] follows the following corollary:

Corollary 3.14. *Let $n \geq 3$ be an integer, the number of spanning trees of $\Delta(G)$ is $n(n_1+1)^{(n_1-1)}(n-n_1)^{(n-(n_1+2))}$.*

REFERENCES

- [1] M. Aouchiche, P. Hansen, Two Laplacians for the distance matrix of a graph, *Linear Algebra Appl.* **439** (2013), 21–33.
- [2] M. Bissler, J. Laubacher and M.L. Lewis, Classifying character degree graphs with six vertices, *Beitr. Algebra Geom.* **60** (2019), 499 – 511.
- [3] M. Bissler, J. Laubacher, Classifying families of character degree graphs of solvable groups, *Int. J. Group Theory* **8** (4)(2019), 37– 46.
- [4] D. Cvetkovic, S. Simic and P. Rowlinson, *An Introduction to the Theory of Graph Spectra*. Cambridge University Press, Cambridge (2009).
- [5] M. Ebrahimi, M. Khatami and Z. Mirzaei, Regular character-graphs whose eigenvalues are greater than or equal to -2 , *Discrete Math.* **346** (1) (2023), 113137.
- [6] R. Hafezieh, M.A. Hosseinzadeh, S. Hossein-zadeh and A. Iranmanesh, *On cut vertices and eigenvalues of character graphs of solvable groups*, *Discrete Appl. Math.* **303** (2021), 86–93.
- [7] B. Huppert, *Research in representation theory at Mainz (1984-1990)*, *Progr. Math.* **95**(1991), 17–36.

- [8] M.L. Lewis, *Character degree graphs of solvable groups of fitting height 2*, *Canad. Math. Bull.* **49**(1) (2006), 127–133.
- [9] M.L. Lewis, An overview of graphs associated with character degrees and conjugacy class sizes in finite groups, *Rocky Mountain J. Math.* **38** (1) (2008), 175–211.
- [10] M.L. Lewis and Q. Meng, Solvable groups whose prime divisor character degree graphs are 1-connected, *Monatsh. Math.* **190** (2019), 541–548.
- [11] O. Manz, Degree problems II: π -separable character degrees, *Comm. Algebra* **13** (1985), 2421–2431.
- [12] O. Manz, R. Staszewski and W. Willems, On the number of components of a graph related to character degrees, *Proc. AMS* **103** (1) (1988), 31–37.
- [13] O. Manz, W. Willems and T.R. Wolf, The diameter of the character degree graph, *J. Reine Angew. Math.* **402** (1989), 181–198.
- [14] B. Mohar, The Laplacian spectrum of graphs. *Graph Theory Combin. Appl.* **vol. 2** (Kalamazoo, MI, 1988), Wiley-Intersci. Publ., Wiley, New York, (1991), 871–898.
- [15] C.P. Morresi Zuccari, Regular character degree graphs, *J. Algebra* **411**(2014), 215–224.
- [16] P.P. Pálffy, On the character degree graph of solvable groups I: three primes, *Period. Math. Hungar.* **36** (1) (1998), 61–65.
- [17] G. Sivanesan, C. Selvaraj and T. Tamizh chelvam, Eulerian Character degree graphs of solvable groups (2023), arXiv: 2304.13472.
- [18] J. Zhang, On a problem by Huppert, *Beijing Daxue Xuebao Ziran Kexue Ban*, **34 (2-3)** (1998) , 143–150.

DEPARTMENT OF MATHEMATICS, GOVERNMENT COLLEGE OF ENGINEERING, SALEM 636011, TAMIL NADU, INDIA, ORCID: 0000-0001-7153-960X.

Email address: sivanesan@gcesalem.edu.in

DEPARTMENT OF MATHEMATICS, PERIYAR UNIVERSITY, SALEM 636011, TAMIL NADU, INDIA, ORCID: 0000-0002-4050-3177.

Email address: selvavlr@yahoo.com