Fine-tuning Games: Bargaining and Adaptation for General-Purpose Models

Benjamin Laufer*

Jon Kleinberg^{†,§}

Hoda Heidari^{‡,§}

Abstract

Major advances in Machine Learning (ML) and Artificial Intelligence (AI) increasingly take the form of developing and releasing general-purpose models. These models are designed to be adapted by other businesses and agencies to perform a particular, domain-specific function. This process has become known as adaptation or fine-tuning. This paper offers a model of the fine-tuning process where a Generalist brings the technological product (here an ML model) to a certain level of performance, and one or more Domain-specialist(s) adapts it for use in a particular domain. Both entities are profit-seeking and incur costs when they invest in the technology, and they must reach a bargaining agreement on how to share the revenue for the technology to reach the market. For a relatively general class of cost and revenue functions, we characterize the conditions under which the fine-tuning game yields a profit-sharing solution. We observe that any potential domain-specialization will either contribute, free-ride, or abstain in their uptake of the technology, and we provide conditions yielding these different strategies. We show how methods based on bargaining solutions and sub-game perfect equilibria provide insights into the strategic behavior of firms in these types of interactions, and we find that profit-sharing can still arise even when one firm has significantly higher costs than another. We also provide methods for identifying Pareto-optimal bargaining arrangements for a general set of utility functions.

1 Introduction

Generative machine-learning (ML) models have garnered a great deal of excitement because they are considered to be *general purpose* (Goldfarb et al., 2023; Eloundou et al., 2023; Crafts, 2021; Pan et al., 2020; Wei et al., 2022; Ouyang et al., 2022). Some have referred to these technologies as *foundation models* (Bommasani et al., 2021, 2023; Fei et al., 2022) because they are designed as massive, centralized models that support potentially many downstream uses. For example, Bommasani et al. (2021) write, "a foundation model is itself incomplete but serves as the common basis from which many task-specific models are built via adaptation."

There is palpable excitement about these technologies. But to turn their potential into actual use and impact, one needs to specialize and tweak the technology to particular application domains. This process takes various names, including *adaptation* (Peters et al., 2019) and, in some contexts, *fine-tuning* (Tajbakhsh et al., 2016; Zhang et al., 2020; Kumar et al., 2022).

Notably, the process of fine-tuning a technology involves multiple parties. Technology teams developing ML and Artificial Intelligence (AI) technologies rely on outside entities to adapt, tweak, transfer, and integrate the general-purpose model. This dynamic suggests a latent strategic interaction between producers of a foundational, general-purpose technology and specialists considering whether and how to adopt the technology in a particular context. Understanding this interaction is necessary to study the social and economic consequences of introducing the technology.

This paper brings methods from economic theory to model and analyze the fine-tuning process. We put forward a model of fine-tuning where the interaction between two agents, a generalist and a specialist, determines how they'll bring a general-purpose technology to market (Figure 1). The result of this interaction

^{*}Cornell Tech, New York, NY, USA

[†]Cornell University, Ithaca, NY, USA

[‡]Carnegie Mellon University, Pittsburgh, PA, USA

[§] Joint senior author.




Figure 1: An illustration of the fine-tuning game. In the first step, players bargain over the revenue-sharing agreement δ . In this example, they agree that G will receive 80% of the revenue and D will receive 20%. In the second step, G develops the technology to performance level $\alpha_0 = 21$. In the third step, D 'fine-tunes' the technology to $\alpha_1 = 25$. If the players collectively receive revenue of 25, they'd share so that G receives 20 and D receives 5.

is a domain-adapted product that offers a certain level of *performance* to consumers, in exchange for a certain level of surplus revenue for the producers. Crucially, the producers must decide how to distribute the surplus, and engage in a bargaining process to do so. An immediate intuition might be to divide this surplus based on contribution to the technology — however, this is one of many potential bargaining solutions, each with different implications for the technology's performance and the distribution of utility. For example, splitting based on contribution can yield a worse-performing technology compared to other bargaining arrangements.

Through this analysis, we discover several general principles that apply not just to today's generative machine learning technologies, but to a potentially wide swath of models that exhibit a similar structure — i.e., developed for general use and adapted to one or more domains. Thus, even as these technologies improve and develop, our proposed model of fine-tuning may continue to describe how they may be adapted for real-world use(s).

Further, as we'll discuss in Section 6.1, some of our findings apply to other general-purpose technologies outside machine learning context. For example, cloud computing infrastructure enables a number of consumer-facing services that use web hosting, database services, and other on-demand computing resources. Additive manufacturing (e.g., 3D printing) requires the production of a general-purpose technology that other people and entities use to create valuable products in particular domains. Finally, digital marketplaces are similarly general, market-making technologies that enable specialist vendors and producers with whom they bargain over surplus.

Our main conceptual contribution is modeling the fine-tuning process as a combination of 1) a **multi-stage game** and 2) a **bargaining process** between a general-purpose technology producer and one or more domain specialist(s). Both players bargain over how to share revenue, and each takes a turn contributing to the technology's performance before it reaches the market. Within the set of Pareto-optimal bargaining agreements, we introduce a number of *bargaining solutions* that represent potential arrangements for how entities involved in AI's development should distribute profit and effort. These bargaining solutions represent different normative assumptions about the appropriate way to distribute welfare.

A significant, high-level take-away from our analysis is a characterization of the specialist fine-tuning strategy for any particular domain. We find that any potential adaptor of a technology falls into one of three groups: **Contributors**, who invest effort before selling the technology; **Free-riders**, who sell the technology without investing any additional effort; and **Abstainers**, who do not enter any fine-tuning agreement and opt not to bring the technology to their particular domain. It turns out, using only marginal information about a domain (0th- and 1st-order approximations of cost and revenue), it is possible to reliably determine which strategy the adaptor will take for a notably broad set of scenarios and cost and revenue functions (Section 5.3).

Our analysis consists in deriving the subgame perfect equilibrium strategies, identifying the set of Pareto-

optimal bargaining agreements, and then solving for various bargaining solutions proposed by economists. Even in the presence of significant cost differentials, we find bargaining leads to profit-sharing agreements because specialists can leverage their power to exit the deal, reducing the reach of the technology — or, in the case of one specialist, preventing the technology from being produced altogether. For fine-tuning games with a somewhat general set of cost and revenue functions, we develop a method for identifying Pareto-optimal bargains. We find that the Pareto-optimal set is a single interval when there is only one specialist, but may be multiple disjoint intervals in the multi-specialist generalization. The potentially disjoint set arises in cases with multiple specialists because a bargaining deal might be reached with some subset of domain specialists while others abstain.

Some have suggested that scholarship on AI and data-driven technologies focuses predominantly on the technical developments without situating these developments in political economy (Abebe et al., 2020; Trajtenberg, 2018; Brevini and Pasquale, 2020). We propose a model that accounts for the different interests and interactions involved in the development of new, general-purpose AI technology. Our model enables analysis on how these interactions affect market outcomes like performance in practice. Understanding these interactions may also inform future regulation of harms when they arise from generative machine-learning technologies.

1.1 Related Work

Existing approaches to fine-tuning. Fine-tuning a base model (e.g., a language model (Howard and Ruder, 2018)) often consists of several steps: (1) gathering and processing domain-specific data, (2) choosing and adjusting the base model's architecture (including number of layers (Wang et al., 2017) and parameters (Sanh et al., 2020)) and the appropriate objective function (Gunel et al., 2020), (3) Updating the model parameters using techniques like gradient descent or transfer learning, and (4) evaluating the resulting model and refining if necessary. Fine-tuning is an instance of the broader concept of transfer learning (Zhuang et al., 2020).

Existing economic models of general-purpose technology production. Several lines of work in growth economics address the development and diffusion of general-purpose technologies (or GPTs). See (Bresnahan, 2010) for a survey and (Jovanovic and Rousseau, 2005) for a historic account of electricity and IT as GPTs with major impacts on the US economy. Scholars have examined the effects of factors such as knowledge accumulation, entrepreneurial activity, network effects, and sectoral interactions on the creation of GPTs (Helpman, 1998). The model presented here abstracts away the forces giving rise to the creation of general-purpose technologies, such as LLMs, in the first place, and instead focuses on the later-stage decision of when (or at what performance level) to release the GPT to market for domain-specialization.

Some have suggested that general-purpose technologies create the need for new business models that describe their impact on individual sectors (Lipsey et al., 2005). Gambardella and McGahan (2010) proposed a similar model of domain adaptation for general-purpose technology that is based on revenue sharing — however, they do not use bargaining or multi-stage strategy to describe how the technology is developed and brought to market. Our notion of *performance* as it relates to model technologies is inspired by economic models of product innovation (Viscusi and Moore, 1993; Cooper, 1984).

Bargaining solutions. Bargaining solutions in economics are used to analyze the outcome of the negotiation/bargaining process through which agents reach mutually beneficial agreements to share gains of collective ventures. The bargaining solutions commonly studied in economics include the Nash Bargaining Solution (NBS) (Nash, 1950) and the Kalai-Smorodinsky Bargaining Solution (KSBS) (Kalai and Smorodinsky, 1975). NBS maximizes the product of the bargainers' individual gains. It proposes that the final outcome of a negotiation should result in an equal division of the surplus between the parties, assuming they have equal bargaining power. The Kalai-Smorodinsky Bargaining Solution proposes an outcome that lies on the Pareto frontier and equalizes the proportion of maximal gains. The Egalitarian (or 'Kalai') Bargaining Solution is another solution where the agreement maximizes the minimum gain for each party (Kalai and Samet, 1985). More general multiplayer bargaining games have been proposed and analyzed with analogous solutions (Moulin, 1991; Chatterjee et al., 1993). A related—but distinct—body of work is referred to as the hold-up problem (Rogerson, 1992). In this literature, two (or more) agents negotiate over an incomplete contract and distribute surplus (Hart and Moore, 1988). In these models, after an initial agreement, players are able to re-negotiate and alter parts of the contract, yielding shifts in strategy.

2 A Model of Fine-Tuning

In this section, we put forward a model of fine-tuning a data-driven technology for use in a domain-specific context. The technology is developed in two steps: First, a general-purpose producer develops a technology up to a certain level of performance. Then, a domain-specific producer decides whether to adopt the technology, and how much to invest in the technology to further improve its performance beyond the general-purpose baseline. After these steps, the two entities share a payout.

Generalist. Player G (for General-purpose producer) is the first to invest in the technology's performance, and brings the performance level to $\alpha_0 \in \mathbb{R}$. G is motivated to invest in the technology because, ultimately, the technology's performance level determines the revenue G earns.

Domain Specialist. After investing in the technology, G can offer the technology to a domain-specialist, denoted D, who fine-tunes the model to their specific use case. If D and G enter an agreement, D will invest in improving the technology's performance from α_0 to $\alpha_1 \in \mathbb{R}$.

Revenue and costs. The technology's performance, α_1 , determines the total revenue that can be gained from fine-tuning the technology in that domain. In particular, we assume there is a monotonic function $r: \mathbb{R}^+ \to \mathbb{R}^+$ such that $r(\alpha_1)$ is the total revenue generated by performance level α_1 . Unless otherwise specified, we assume $r(\cdot)$ is the identity function, that is, the total revenue brought by technology is α_1 . The cost associated with producing α_1 requires considering the two steps involved with developing the technology: general production and fine-tuning. We say that G faces cost function $\phi_0(\alpha_0): \mathbb{R}^+ \to \mathbb{R}^+$ to produce a general technology at performance-level α_0 . D faces cost function $\phi_1(\alpha_1; \alpha_0): \mathbb{R}^+ \to \mathbb{R}^+$ to bring the technology from performance α_0 to performance α_1 . We assume these cost functions are publicly known. Unless otherwise specified, we also assume r(0) = 0, $\phi_0(0) = 0$, and $\phi_1(\alpha_1 = \alpha_0; \alpha_0) = 0$, meaning that not investing in the technology is free and brings in zero revenue.

The fine-tuning game. The players are G and D. In deciding whether to purchase the technology, D negotiates revenue sharing with G. G and D share revenue $r(\alpha_1)$ according to a bargaining parameter $\delta \in [0,1]$. At the end of the game, G receives $\delta r(\alpha_1)$ in revenue, and D receives $(1-\delta)r(\alpha_1)$. The model fine-tuning game consists in each player deciding their level of investment and collectively bargaining to decide δ . The game proceeds as follows:

- 1. G and D negotiate bargaining coefficient $\delta \in [0, 1]$.
- 2. G invests in a general-purpose technology, subject to cost $\phi_0(\alpha_0)$, yielding performance-level α_0 .
- 3. D fine-tunes the technology, subject to cost $\phi_1(\alpha_1;\alpha_0)$, yielding performance-level α_1 .

The steps of the game are illustrated in Figure 1. Players earn the following utilities, defined as revenue share minus cost:

$$U_G(\delta) := \delta r(\alpha_1) - \phi_0(\alpha_0), \tag{1}$$

$$U_D(\delta) := (1 - \delta)r(\alpha_1) - \phi_1(\alpha_1; \alpha_0). \tag{2}$$

If the players do not agree to a feasible bargain $\delta \in [0, 1]$, then the bargaining outcome is referred to as disagreement. In this scenario, the generalist receives d_0 and the specialist receives d_1 . We assume, unless otherwise specified, that the disagreement scenario is described by $d_0 = d_1 = 0$.

2.1 Primer on Bargaining Games

Bargaining games are a potentially useful method for computer science (CS) research. In this section we include a primer on these methods before demonstrating their use in our model.

Often, in game theory and mechanism design, a model consists of a strategic interaction between two or more agents with individual utility functions. When the players must reach an agreement over a set of alternatives, game theoretic tools provide methods to identify candidate agreements. For example, the concept of Pareto efficiency helps guide players to only agreement(s) for which no alternative exists that

¹It could be the case that a general-purpose technology producer receives positive payout even if the specialist abstains from a bargain. This case is, essentially, a second 'path' for bringing a general technology to market. We discuss this possibility further in the multi-specialist generalization of our model (Section 5).

makes one or more players strictly better off. As is the case in our current setting, Pareto efficiency often identifies a *set of multiple solutions*, within which further tools and methods are needed to identify potential strategies and agreements.

A bargain is a process for identifying joint agreements between two or more agents on how to share payoff. The *Bargaining Problem*, formalized by Nash (1950), consists of two players that must jointly decide how to share surplus profit. The problem consists of a set of feasible agreements and a 'disagreement' alternative, which specifies the utilities players receive if they do not come to an agreement.

Bargaining solutions are established ways to select among candidate agreements on how to share surplus. Different bargaining solutions, proposed over the years by mathematicians and economists, aim to satisfy certain desiderata like fairness, Pareto optimality, and utility-maximization. Typically, solving for bargaining solutions consists in defining some measure of *joint utility* between players (e.g. take the sum, product, or minimum of the players' utilities). The feasible, Pareto-optimal solution that maximizes this joint utility is known as a *bargaining solution*.

Bargaining solutions are normative: they provide guidelines for how surplus payoffs should be distributed. Solutions are inspired by moral theories like utilitarianism (which aims to maximize the sum of utilities) and egalitarianism (which aims to maximize the worst-off agent). We demonstrate the use of bargaining solutions in the subsequent sections.

2.2 Pareto-Optimal Bargains

Our model of the fine-tuning process unfolds in two stages: the first stage is a bargain where the players must jointly agree on δ , and the second stage is a sequential game where the players make decisions individually in order (i.e., G moves first and D moves second). In order to derive solutions, it is important to define Pareto domination and Pareto efficiency. Once we've defined a few preliminary qualities, we'll state our first finding deriving the set of Pareto-optimal solutions for a general set of cost and revenue functions.

Definition 2.1 (Pareto-dominant agreements). A bargaining agreement δ_i Pareto-dominates an alternative agreement $\delta_i \neq \delta_i$ iff at least one player gains utility by switching from δ_i to δ_i , and no players lose utility.

Definition 2.2 (Pareto-optimal agreements). A **Pareto-optimal** agreement is one where no alternative agreement would improve the utility of one player without decreasing the utility of the other player. In other words, it is an agreement that is not Pareto-dominated by any other agreement.

Definition 2.3 (Strictly Unimodal Function). A function $f : \mathbb{R} \to \mathbb{R}$ is called a **strictly unimodal function** over a real domain $x \in \mathbf{D}$ if there exists some value $m \in \mathbf{D}$ such that f is strictly increasing $\forall x \leq m$ and f is strictly decreasing $\forall x \geq m$.

When reasoning about how two agents can jointly reach an agreement, it is useful to start by considering the scenario where one player is *all-powerful*, meaning the bargain is determined solely to maximize one player's utility. The formal definition of this sort of bargaining arrangement is provided below.


Definition 2.4 (Powerful-P solution). For a given fine-tuning game player $P \in \{G, D\}$, the powerful-P solution is the revenue-sharing agreement $\delta^{Powerful\ P} \in [0,1]$ that maximizes the utility of P:

$$\delta^{Powerful\ P} = argmax_{\delta \in [0,1]} U_P(\delta).$$

We are now in a position to state our first theorem, which characterizes the Pareto-optimal solutions to any fine-tuning game with strictly unimodal utility functions.

Theorem 2.1. Consider a fine-tuning game where players bargain over a parameter δ . If the players' utilities are strictly unimodal functions of δ , the set of Pareto-optimal agreements is the interval between their optima $\{\delta^{Powerful\ D}, \delta^{Powerful\ G}\}$, where both players' utilities are greater than the disagreement scenario. If no such interval exists, then disagreement is Pareto-optimal.

The proof is provided in Appendix 7. To provide some intuition for the proof, consider the range of agreements δ between the point which maximizes one player's utility (say, $\delta^{Powerful\ D}$) and the point which maximizes the other ($\delta^{Powerful\ G}$). Agreements within this range exhibit a trade-off between the two utilities.






Figure 2: Example to illustrate Theorem 2.1. Left: For two strictly unimodal, positive utility functions over a bargaining parameter δ , the set of Pareto-optimal bargaining agreements is the interval between their optima. Within this interval, players' utilities are characterized by a trade-off, whereas outside this interval, any agreement is Pareto-dominated. Right: The non-numerical bargaining outcome ('disagreement') consists in players receiving 0 utility. Thus, if a potential bargaining agreement yields negative utility for one or both players, they prefer to exit the agreement. The relevant set of Pareto-optimal agreements is constrained to only the interval where both players receive payout that exceeds the disagreement scenario. If no such interval exists, disagreement is Pareto-optimal.

Agreements outside this range, however, leave both players worse-off than, e.g., the nearest powerful-P solution, so they are Pareto-dominated. This intuition is illustrated in Figure 2.

Theorem 2.1 applies to a notably broad set of utility functions. To illustrate some of these forms, and for ease of reference, we provide the following immediate corollary:

Corollary 2.1. If U_D is either strictly increasing, strictly decreasing, or strictly concave in δ , and U_G is either strictly increasing, strictly decreasing, or strictly concave in δ , then the set of Pareto-optimal agreements is the interval between their optima $\{\delta^{Powerful\ D}, \delta^{Powerful\ G}\}$, where both players' utilities are greater than the disagreement scenario. If no such interval exists, then disagreement is Pareto-optimal.

Notice this follows immediately from Theorem 2.1 because any strictly increasing, strictly decreasing or strictly concave function on the interval $\delta \in [0, 1]$ is strictly unimodal on the same interval.

Equipped with the theorem above, solving the fine-tuning game consists of the following steps: (1) Use backward induction to solve for D and G's strategies, represented by α_1^* and α_0^* , in terms of δ . (2) Find the set of Pareto-optimal bargaining agreements δ between the powerful-D and powerful-G solutions. (3) Within the Pareto set, solve for bargaining agreements that maximize some joint function of the players' utilities.

3 Analysis for Polynomial Costs

Our model applies to general cost and revenue functions, and in Section 4 we will provide results at this general level. But to understand how the central parameters of the model interact in closed form, it is also useful to study instantiations of the model with specific functional forms. Accordingly, we show in this section how to solve the model with a set of polynomial cost functions as a paradigmatic instance of convex cost functions, where the marginal costs increase as the technology is improved. Following this, we show how to draw conclusions about the model with general costs.

Thus, in this section, cost functions take the following polynomial function forms:

$$\phi_0(\alpha_0) := c_0 \alpha_0^{k_0},\tag{3}$$

$$\phi_1(\alpha_1; \alpha_0) := c_1(\alpha_1 - \alpha_0)^{k_1}. \tag{4}$$

Here, $c_0, c_1 > 0$ since costs should increase with investment, and $k_0, k_1 > 1$, meaning that an incremental improvement grows costlier at higher levels of performance. We will continue to assume that $r(\alpha_1) = \alpha_1$ throughout this section's analysis.

First (3.1), we derive the subgame perfect equilibrium strategies α_0^* , α_1^* for fixed δ . Second (3.2), we find the set of Pareto-optimal revenue-sharing schemes δ^{Pareto} . Reaching a revenue-sharing agreement $\delta^* \in \delta^{Pareto}$ is modeled as a bargaining problem because the players must decide how to share surplus utility. So, third (3.3), we define five potential bargaining solutions: Best-performing-model, Vertical Monopoly, Egalitarian, Nash Bargaining Solution, and Kalai-Smorodinsky. Where possible, we derive closed-form expressions for these solutions. We end by discussing the implications of these different revenue-sharing schemes.

3.1 Derivation of the Subgame Perfect Equilibrium for a Given δ

We use backward induction to determine the fine-tuning game's subgame perfect equilibrium (which we will refer to as a 'solution' or 'equilibrium'). Fixing the outcome of the initial negotiation, δ , it is possible establish the following closed-form solution:

Theorem 3.1. For a fixed δ , the sub-game perfect equilibrium of the fine-tuning game with polynomial costs yields the following best-response strategies:

$$\alpha_0^* = \left(\frac{\delta}{k_0 c_0}\right)^{\frac{1}{k_0 - 1}}, \quad \alpha_1^* = \left(\frac{\delta}{k_0 c_0}\right)^{\frac{1}{k_0 - 1}} + \left(\frac{1 - \delta}{k_1 c_1}\right)^{\frac{1}{k_1 - 1}}.$$

Notice that the domain-specific performance, α_1^* , is equal to the general-purpose performance, α_0^* , plus a term, $(\frac{1-\delta}{k_1c_1})^{\frac{1}{k_1-1}}$, independent of the G's choice over α_0^* . This is because the cost of marginal improvements for D only depends on the difference $(\alpha_1 - \alpha_0)$, and is not affected by a large or small initial investment by G. Though we assume, in this section, that D's cost is not defined solely in terms of marginal improvement, Section 4 contains findings that generalize beyond this assumption.

As an immediate corollary of Theorem 3.1, we derive players' utilities as a function of δ alone.

Corollary 3.1. For a fixed bargaining parameter δ , the players' utilities are as follows:

$$U_G(\delta) = \left(\frac{1}{k_0 c_0}\right)^{\frac{1}{k_0 - 1}} \left(1 - \frac{1}{k_0}\right) \delta^{\frac{k_0}{k_0 - 1}} + \left(\frac{1}{k_1 c_1}\right)^{\frac{1}{k_1 - 1}} \delta(1 - \delta)^{\frac{1}{k_1 - 1}},\tag{5}$$

$$U_D(\delta) = \left(\frac{1}{k_1 c_1}\right)^{\frac{1}{k_1 - 1}} \left(1 - \frac{1}{k_1}\right) (1 - \delta)^{\frac{k_1}{k_1 - 1}} + \left(\frac{1}{k_0 c_0}\right)^{\frac{1}{k_0 - 1}} (1 - \delta) \delta^{\frac{1}{k_0 - 1}}.$$
 (6)

In order to determine the set of Pareto-optimal agreements, we first find that the utility functions derived above are strictly unimodal for all c_0, c_1 and $k_0, k_1 \ge 2$.

Proposition 3.1. In the fine-tuning game with polynomial costs, if $k_0, k_1 \ge 2$, then U_G and U_D are strictly unimodal functions of $\delta \in [0, 1]$.

The above findings are proven in Appendix 8.3. They suggest that a general set of cost functions yield strictly unimodal utility curves. The set of Pareto-optimal solutions to these games can therefore be identified using Theorem 2.1. It is easy to show that the strict unimodality finding further generalizes to linear combinations of polynomial terms of the form provided in Equations 3 and 4, so long as all exponents are greater than or equal to 2. However, when the condition is not met and $k_0, k_1 < 2$, numerical simulations suggest that there are counter-examples to the strict unimodality property. When the strict unimodality property does not hold, it is still possible to analyze players' strategies—for example, our analysis in Sections 4 and 5.3 stands even in cases where utility functions that are not unimodal in δ .

Solving the powerful-G, powerful-D, vertical monopoly or other bargaining solutions consists in maximizing players' utilities either separately or combined into a joint utility. This is possible once parameters are specified; however, we cannot produce a closed-form expression for the general polynomial case because doing so would require solving for the zeroes of a polynomial of high degree. Therefore, for the remainder of this section, we will demonstrate the solution steps using parameter values $k_0, k_1 = 2$. We call this the case of quadratic costs. We choose the quadratic case for clarity and exposition, though we note that other solutions with other parameter values can be calculated using analogous steps.




Figure 3: Utilities of the Generalist (a) and Domain Specialist (b) for different bargaining parameters and costs. The general-purpose producer prefers to share revenue when $\frac{c_1}{c_0} < 1$ and the domain-specific producer prefers to share revenue when $\frac{c_1}{c_0} > 1$. (c) depicts the technology performance α_1^* . Color bar scales are defined assuming $c_0 = 1$.

3.2 Deriving the Set of Pareto-optimal Bargaining Agreements

We've derived both players' optimal strategies for fixed δ . Now, we consider the process where players agree on a particular value of δ . Since both players must enter an agreement in order for the technology to be viable, the determination of δ is a two-player bargaining game. We start by solving for the set of Pareto-optimal bargaining agreements, which is the interval between the 'powerful-player' solutions, defined below.

3.2.1 Powerful-Player Solutions

As we showed in Theorem 2.1, identifying the 'powerful-player' agreements is important for characterizing the set of Pareto-optimal bargaining solutions. Thus, we begin this section of analysis by solving for the powerful-G and powerful-D solutions (as defined in Definition 2.4).

Proposition 3.2 (Powerful-G Solution). The Powerful-G solution to the model fine-tuning game with quadratic costs is as follows:

$$\delta^{Powerful \ G} = \begin{cases} \frac{c_0}{2c_0 - c_1} & for \ c_1 < c_0, \\ 1 & for \ c_1 \ge c_0. \end{cases}$$

Proposition 3.3 (Powerful-D Solution). The Powerful-D solution to the model fine-tuning game with quadratic costs is as follows:

$$\delta^{Powerful \ D} = \begin{cases} 0 & for \ c_1 < c_0, \\ \frac{c_1 - c_0}{2c_1 - c_0} & for \ c_1 \ge c_0. \end{cases}$$

Now, using Theorem 2.1 and Proposition 3.1, we can define the set of Pareto-optimal solutions as: $\delta^{Pareto} \in \{\delta : \delta \leq \delta^{Powerful\ G} \cap \delta \geq \delta^{Powerful\ D}\}$. A visual representation of these solutions for the fine-tuning game with quadratic costs is given in Figure 5.

3.3 Bargaining Solutions

If neither player dominates in a bargain, how do they decide how to share surplus profit? Solutions to bargaining problems tend to find an agreement that maximizes some joint utility function or satisfies certain desirable properties. In this section, we define the various bargaining solutions that the two players could plausibly arrive at within the set of Pareto-optimal solutions. These solutions mostly use a joint utility function to guide the bargaining agreement, as depicted in Figure 4. A visual representation of the bargaining solutions is provided in Figure 5. Definitions and closed-form solutions are provided below, and the proofs and steps yielding the solutions are included in Appendix 8.

Solution that maximizes the technology's performance. The first solution we propose presumes the joint goal of the two players is to collectively produce a technology with maximum performance α_1^* . There




Figure 4: Various joint-utility functions for finding bargaining solutions. Gray regions are δ values that are not Pareto-optimal and therefore not candidate bargaining solutions. Color bar scales are defined assuming $c_0 = 1$.

are a few ways to think of this quantity: It is the performance of the technology, and, equivalently, it is also the amount of revenue the two players collect. Though we do not formally specify a social welfare function, the technological performance can be thought of as the total utility offered to society by firms G and D.

Definition 3.1 (Maximum-performance solution). For the fine-tuning game, the maximum-performance bargaining solution is the feasible revenue-sharing agreement $\delta^{max-\alpha_1^*} \in [0,1]$ that maximizes the technology's performance α_1^* : $\delta^{max-\alpha_1^*} = argmax_{\delta \in [0,1]}\alpha_1^*$.

Proposition 3.4 (Maximum- α_1^* Solution). A bargaining solution that maximizes the technology's performance is given by:

$$\delta^{Max-\alpha_1^*} = \begin{cases} 0 & for \ c_1 < c_0, \\ 1 & for \ c_1 \ge c_0. \end{cases}$$

Vertical Monopoly Solution. A perhaps intuitive approach to bargaining is to choose a revenue-sharing agreement that maximizes the sum of utilities $U_G + U_D$. This solution imagines that the two players are jointly controlled by a single entity who simply wishes to maximize the sum of utility. This solution is known as either the 'vertical monopoly' solution or the 'utilitarian' solution.

Definition 3.2 (Vertical Monopoly Solution). For the fine-tuning game, the Vertical Monopoly (or 'Utilitarian') Solution is the feasible revenue-sharing agreement $\delta^{VM} \in [0,1]$ that maximizes the sum of the players' utilities: $\delta^{VM} = argmax_{\delta \in [0,1]} (U_G(\delta) + U_D(\delta))$.

Proposition 3.5 (Vertical Monopoly Solution). The Vertical Monopoly Bargaining Solution to the fine-tuning game with quadratic costs is as follows:

$$\delta^{\textit{Vertical Monopoly}} = \frac{c_1}{c_1 + c_0}.$$

Egalitarian Bargaining Solution. An alternative bargaining approach tries to help the worst-off player. This bargaining solutions is known as the 'egalitarian' solution.

Definition 3.3 (Egalitarian Bargaining Solution). For the fine-tuning game, the Egalitarian Bargaining Solution is the feasible revenue-sharing agreement $\delta^{Egal.} \in [0,1]$ that maximizes the minimum of the players' utilities: $\delta^{Egal.} = argmax_{\delta \in [0,1]} \left(min_{P \in \{G,D\}} \left(U_P(\delta) \right) \right)$.

Proposition 3.6 (Egalitarian Bargaining Solution to the fine-tuning game with quadratic costs). The Egalitarian Bargaining Solution to the fine-tuning game with quadratic costs is given by:

$$\delta^{Egal.} = \frac{-\sqrt{c_0^2 - c_0 c_1 + c_1^2} - c_1 + 2c_0}{3(c_0 - c_1)}.$$

Nash Bargaining Solution. The Nash Bargaining solution maximizes the product between the two players' utilities. This arrangement satisfies a number of desiderata, originally laid out by Nash (1950).




Figure 5: Bargaining agreements for the fine-tuning game with quadratic costs (i.e., the polynomial cost game where $k_0 = k_1 = 2$). The set of Pareto-optimal agreements is the interval between the Powerful G (blue) and Powerful D (orange) solutions. Most bargaining solutions suggest revenue sharing, even at the limit where one player faces much higher costs. For example, the Nash bargaining solution (red) is limited at 0.75 for $c_1 >> c_0$. One exception is the vertical monopoly solution (green), which is limited at 1.

Definition 3.4 (Nash Bargaining Solution). For the fine-tuning game, the Nash Bargaining Solution is the feasible revenue-sharing agreement $\delta^{NBS} \in [0,1]$ that maximizes the product of the players' utilities: $\delta^{NBS} = argmax_{\delta \in [0,1]} (U_G(\delta) * U_D(\delta)).$

Though a closed-form solution for quadratic functions is possible, it involves solving the roots of a cubic function and yields a solution that is clunky and uninterpretable. Therefore, we refer the reader to our numerical findings on this solution, which are depicted in Figures 4 and 5.

Kalai-Smorodinsky Bargaining Solution.

Another solution suggested in economic literature is known as the 'Kalai-Smorodinsky' bargaining solution. This solution equalizes the ratio of maximal gains. More formally:

Definition 3.5 (Kalai-Smorodinsky Bargaining Solution). (Kalai and Smorodinsky (1975)) For the fine-tuning game, the Kalai-Smorodinsky Bargaining Solution is the feasible revenue-sharing agreement $\delta^{KSBS} \in [0,1]$ that satisfies the following relation:

$$\frac{U_G(\delta^{KSBS})}{\max_{\delta \in \delta^{Pareto}} U_G(\delta)} = \frac{U_D(\delta^{KSBS})}{\max_{\delta \in \delta^{Pareto}} U_D(\delta)}.$$

Notice the denominators in the above equation are simply the utilities associated with the powerful-G and powerful-D solutions. Despite this simplifying step, the closed form Kalai-Smorodinsky solution is clunky and uninterpretable, so we omit it from this paper. We refer the reader to our numerical findings on this solution, which are depicted in Figure 5.

3.4 Discussion on Bargaining Solutions

Above we solve for a number of bargaining solutions revealing different possible configurations of fine-tuning arrangements. The general technology-producer and the domain specialist each have different optimal arrangements, between which any agreement is Pareto-optimal in the case of polynomial costs.

The first notable take-away is that players do not necessarily opt to maximize their own share of the profit. Even if one player has full control over the bargaining solution, depending on the relative cost of production, they may benefit from a profit-sharing agreement in order to encourage investment by the other player. If bargaining is conceptualized as splitting a pie, one player prefers to cede some portion of the pie if it means the entire pie grows to a size that justifies profit-sharing. This phenomenon arises in real-world settings. For instance, Apple allows third party developers to build software on iPhones. Opening up the tasks of application development to third parties improves consumer experience so much that consumers are willing to purchase apps or other capabilities within apps. This additional revenue is then shared between Apple and the developer, leaving Apple with higher profits and a better product. Revenue sharing arises, often, because doing so is lucrative.

Profit-sharing is present even when both players have exceedingly different costs of production (i.e., when c_i approaches 0 or ∞). In these limiting instances, we find that the Nash bargaining solution, Kalai-Smorodinski, and Egalitarian solutions all suggest profit-sharing. Only the Utilitarian solution—which models the two players as a vertical monopoly that is centrally controlled—yields the intuitively performance-optimal bargain, where the player with lower costs receives the entire profit. However, the vertical monopoly solution is not always performance-optimal. It underperforms the Kalai-Smorodinsky solution when the players' face similar costs ($\sim 0.5 < \frac{c_1}{c_0} < 2.5$).

The bargaining solutions are neither binding rules nor descriptive observations; instead, they are normative. Identifying joint utility functions can help guide agents towards decisions that serve collective interests. For example, utilitarian and egalitarian solutions offer different visions for the appropriate distribution of welfare. In the same vein, specifying and committing to a social welfare function would allow us to identify a bargaining solution that might be referred to as 'socially optimal.' Unsurprisingly, however, specifying social interests in a single function is a difficult undertaking. In any given context, all the relevant interests, requirements, and aims must be accounted for. In our present case, a social welfare function would need to balance the interests of (at least) 1) the technology's producers 2) consumers who value performance and 3) other external stakeholders. The procedure demonstrated in this section (and generalized in the coming sections) provides a road map for a social welfare analysis of the deployment of general-purpose models. Such an analysis might uncover how fine-tuning processes can be configured to serve collective, societal interests.

4 Findings for General Revenue and Costs

The fine-tuning game may be defined for any cost and revenue functions ϕ_0, ϕ_1, r . We demonstrate in Section 3 that closed-form solutions are attainable for certain polynomial function forms. In this section, we provide general results that suggest the existence of solutions for a broad set of cost and revenue functions. We put forward an existence finding that suggests meaningful cooperation (profit sharing agreement without free-riding) is viable in a swath of fine-tuning games. Our result is that for any non-decreasing cost and revenue functions, as long as $r'(0) > \lambda_0 \phi'_0(0)$ and $r'(\alpha_0) > \lambda_1 \phi'_1(\alpha_0)$ where $\lambda_0, \lambda_1 \geq 2$, there exists a profit-sharing solution that (a) Pareto-dominates disagreement and (b) does not lead to free-riding.

Before we state the theorem formally, we have to define free riding for the fine-tuning game:

Definition 4.1 (Free riding). A solution to the fine-tuning game $\delta_i \in [0,1]$ exhibits **free riding** if at least one player receives profit without investing any effort in improving the technology. That is, either $\alpha_0 = 0$ or $\alpha_1 = \alpha_0$.

The formal theorem statement is below:

Theorem 4.1 (Revenue sharing solution for the fine-tuning game). Consider any fine-tuning game where $r(\alpha_1)$, $\phi_0(\alpha_0)$, and $\phi_1(\alpha_1)$ are non-decreasing. If the following two marginal conditions are met:

- Condition 1: $r'(0) > \lambda_0 \phi'_0(0)$ where $\lambda_0 \geq 2$,
- Condition 2: $r'(\alpha_0) > \lambda_1 \phi_1'(\alpha_0)$ where $\lambda_1 \geq 2$,

Then there exists a solution δ^* to the fine-tuning game with the following properties: (A) Players share revenue $0 < \delta^* < 1$. (B) Players do not free ride. (C) δ^* Pareto-dominates disagreement.

The full proof is provided in Appendix 9.1. The intuition is that, as long as it is marginally profitable for both players to invest some finite positive amount of effort, then both players have positive utility (Pareto-dominating the disagreement alternative), neither player free rides, and both receive some profit. To provide further exposition of the logical argument, we include the component Lemmas below.

Lemma 4.1. If $r'(0) > \lambda_0 \phi_0'(0)$ for a constant $\lambda_0 \ge 2$, then there exists a set $A^* \subseteq (0,1)$ such that $\frac{1}{2} \in A^*$ and for all $\delta \in A^*$, $\alpha_0^*(\delta) > 0$, $U_G(\delta) > 0$.

Lemma 4.2. If $r'(\alpha_0) > \lambda_1 \phi_1'(\alpha_0)$ for a constant $\lambda_1 \geq 2$, then there exists a set $B^* \subseteq (0,1)$ such that $\frac{1}{2} \in B^*$ and for all $\delta \in B^*$, $\alpha_1^*(\delta) > \alpha_0^*$, $U_D(\delta) > 0$.

Corollary 4.1. $A^* \cap B^*$ is a non-empty set where any solution $\delta^* \in A^* \cap B^*$ satisfies the three properties: $0 < \delta_i < 1$; $0 < \alpha_0^* < \alpha_1^*$; and $U_D, U_G > 0$.

This theorem perhaps helps explain why real-world situations arise where two different entities collaboratively invest in technologies. Cooperation is advantageous in a wide variety of fine-tuning scenarios. In the next section, we will further generalize our findings to a new set of games, for which the above theorems may be adapted.

5 Extension to Multiple Domain Specialists

So far, we've modeled the fine-tuning process as a two-player game between a generalist and a specialist. However, an important feature of general-purpose AI models is that they can be developed without fully anticipating the set of possible downstream use-cases. To capture the possibly many use-cases for general-purpose models, in this section, we generalize our model to the case where $n \ge 1$ domain specialists adapt the technology.

The multi-specialist fine-tuning game. Consider a game with $n \geq 1$ specialists. The players are G, D_1 , D_2 ,..., D_n and we'll use i to index the specialists. G develops a technology to general performance α_0 , after which domain specialist D_i specializes the technology to reach performance α_i in their domain. G and D_i share revenue $r_i(\alpha_i)$ according to bargaining parameter $\delta \in [0, 1]$. The bargaining parameter is fixed, which captures common scenarios where the generalist simply has to set a certain pricing agreement for model access. In other words, G cannot price discriminate depending on domain i. The game proceeds as follows:

- Players collectively bargain to decide $\delta \in [0, 1]$.²
- G invests in a general-purpose technology yielding performance-level α_0 and subject to cost $\phi_0(\alpha_0)$.
- Each specialist D_i may fine-tune the technology by choosing a performance level α_i subject to cost $\phi_i(\alpha_i; \alpha_0)$.

Players earn the following utilities, defined as revenue share minus cost:

$$U_G(\delta) := \sum_i \delta r_i(\alpha_i) - \phi_0(\alpha_0), \tag{7}$$

$$U_{D_i}(\delta) := (1 - \delta)r_i(\alpha_i) - \phi_i(\alpha_i; \alpha_0). \tag{8}$$

If the general-purpose producer does not agree to a feasible bargain $\delta \in [0,1]$, then all players receive 0 utility. If any particular specialist does not agree to a feasible bargain, this does not preclude other specialists from reaching a deal. Note that some general-purpose technologies might be marketed directly to consumers by the generalist, meaning that reaching a deal with an individual specialist is not, necessarily, needed for G to receive revenue. We believe this scenario can be captured by an additional specialist engaged in a vertical monopoly agreement with G.

In the remainder of this section, we generalize the Pareto-characterization, subgame perfect equilibria, and bargaining solutions that were demonstrated for the one-specialist case. Although some steps require

²In cases with multiple domain specialists, the bargain over δ is a multi-player bargaining game. We arrive at bargaining solutions for games with more than two players through a similar process to the two-player case, where a joint utility function is specified and then optimized. For more on collective welfare-driven decisions, see Sen (2018).




Figure 6: Illustration of Theorem 5.1. When there are multiple specialists, the set of Pareto-optimal bargaining agreements is not, necessarily, a single interval, even when utitilities are strictly unimodal. In this illustrated example, the region in red represents values of δ for which all relevant players' utilities are increasing. In the red region, D_1 would abstain from the agreement. Any value of δ in this region is Pareto-dominated by $\delta^{\text{Powerful } D_2}$

additional nuance and complexity (for example, the Pareto-region is no longer, necessarily, a single interval), the high-level take-away is that the solution steps demonstrated in this paper generalize to cases with multiple specialists. Solving the bargaining agreements proceeds similarly to Section 3.

5.1 Pareto-Optimal Bargains

First we show that even in cases where all players' utility functions are strictly unimodal, the set of Pareto-optimal bargains is no longer necessarily a single interval—as was the case with one specialist. Generalizing our findings about the set of Pareto-optimal bargaining solutions to multiple-specialists requires additional steps to exclude solutions that are Pareto-dominated within the region between players' optima.

Below we state the theorem that generalizes 2.1 for the multi-player fine-tuning game.

Theorem 5.1. Consider the multi-player fine-tuning game where players' utility functions are strictly unimodal over the bargaining parameter δ . The Pareto set of bargaining agreements δ^{Pareto} is the feasible region consisting of all δ values in [0,1] excluding:

- all values of δ for which $U_G(\delta) < 0$.
- all values of δ for which $U_G(\delta)$ is positive and increasing and no values of $U_{D_i}(\delta)$ are positive and decreasing $\forall i$.
- all values of δ for which U_G(δ) is positive and decreasing and no values of U_{Di}(δ) are positive and increasing ∀i.

Disagreement is Pareto-optimal, if and only if all values of $\delta \in [0,1]$ are excluded through the above criteria.

A proof of the above theorem is provided in Appendix 10.1. Figure 6 provides an illustration of a 2-specialist fine-tuning game where the Pareto-optimal region is not a single interval but instead consists of two disconnected intervals. The finding highlights the added complexity of the multi-specialist fine-tuning game, where individual specialists can abstain but a deal can still be reached among other players.

5.2 Demonstration of Equilibria and Bargaining Solutions for Polynomial Costs

For the multi-specialist game, we demonstrate our results on a set of polynomial cost functions, defined below for $c_0, c_i > 0$ and $k_0, k_i \ge 2$:

$$\phi_0(\alpha_0) := c_0 \alpha_0^{k_0},\tag{9}$$

$$\phi_i(\alpha_i; \alpha_0) := c_i(\alpha_i - \alpha_0)^{k_i}. \tag{10}$$

The equilibria and bargaining solutions for this set of cost functions follow a similar set of steps to those demonstrated in the one-specialist fine-tuning game (Section 3). When the players' utility functions are positive unimodal functions of $\delta \in [0,1]$ (e.g., in the case shown in Observation 10.1), we can use Theorem 5.1 to find Pareto-optimal solutions and report bargaining solutions by maximizing various expressions for joint utility as a function of δ . We include the full set of results in Table 1. We refer the reader to Appendix 10 for the full description of steps and results.

Description	Solution	Result that generalizes
Generalist Strategy	$lpha_0^* = \sqrt[k_0-1]{rac{n\delta}{c_0k_0}}$	Theorem 3.1
Specialists' Strategies	$\alpha_i^* = \sqrt[k_0 - 1]{\frac{n\delta}{c_0 k_0}} + \sqrt[k_i - 1]{\frac{1 - \delta}{c_i k_i}}$	
Generalist Utility	$U_G(\delta) = n \left(\frac{n}{c_0 k_0}\right)^{\frac{1}{k_0 - 1}} \left(1 - \frac{1}{k_0}\right) \delta^{\frac{k_0}{k_0 - 1}} + \delta \sum_i (c_i k_i)^{\frac{-1}{k_i - 1}} (1 - \delta)^{\frac{1}{k_i - 1}}$	Corollary 3.1
Specialists' Utilities	$U_{D_{i}}(\delta) = \left(\frac{n}{c_{0}k_{0}}\right)^{\frac{1}{k_{0}-1}} \left(\delta^{\frac{1}{k_{0}-1}} - \delta^{\frac{k_{0}}{k_{0}-1}}\right) + \left(c_{i}k_{i}\right)^{\frac{-1}{k_{i}-1}} \left(1 - \frac{1}{k_{i}}\right) \left(1 - \delta\right)^{\frac{k_{i}}{k_{i}-1}}$	Corollary 3.1
Powerful- G Solution	$\delta^{\text{P.G}} = \begin{cases} \frac{\sum_{i} \frac{1}{c_{i}}}{2\sum_{i} \frac{1}{c_{i}} - \frac{n^{2}}{c_{0}}} & for \frac{1}{n} \sum_{i} \frac{c_{0}}{c_{i}} > n, \\ 1 & else. \end{cases}$ $\delta^{\text{P.D}_{i}} = \begin{cases} \frac{2c_{0} - 2nc_{i}}{c_{0} - 2nc_{i}} & for \frac{c_{0}}{c_{i}} < n, \\ 0 & else. \end{cases}$	Proposition 3.2
Powerful- D_i Solutions	$\delta^{\text{P.}D_i} = \begin{cases} \frac{2c_0 - 2nc_i}{c_0 - 2nc_i} & for \frac{c_0}{c_i} < n, \\ 0 & else. \end{cases}$	Proposition 3.3
Max-performance Solution	$\delta^{ ext{Max-}\sum_i lpha_i^*} = egin{cases} 0 & for rac{1}{n}\sum_i rac{c_0}{c_i} > n, \ 1 & else. \end{cases}$	Proposition 3.4
Vertical Monopoly Solution	$\delta^{ ext{V.M.}} = rac{n^2}{n^2 + \sum_i rac{c_0}{c_i}}$	Proposition 3.5
Egalitarian Solution	$\delta^{Egal.} = \frac{\sqrt{c_0^2(\sum_i \frac{1}{c_i})^2 - 2c_0n(\sum_i \frac{1}{c_i}) + \frac{c_0}{c_{\max}}n^2 + n^2}}{n^2 + 2n - \frac{c_0}{c_{\max}} - 2c_0(\sum_i \frac{1}{c_i})}{n^2 + 2n - \frac{c_0}{c_{\max}} - 2c_0(\sum_i \frac{1}{c_i})} + \frac{n - \frac{c_0}{c_{\max}}n - c_0(\sum_i \frac{1}{c_i})}{n^2 + 2n - \frac{c_0}{c_{\max}} - 2c_0(\sum_i \frac{1}{c_i})}$	Proposition 3.6

Table 1: Solutions to the multi-specialist fine-tuning game with polynomial costs. Though the equilibria strategies and utility functions are solved for general polynomial costs, the bargaining solutions represent the quadratic case where $k_0 = k_i = 2$. The max-performance solution for multiple specialists maximizes the sum of all downstream performances (see Definition 10.1). For the Egalitarian solution, $c_{\text{max}} := \max_i c_i$.

The results suggest that meaningful bargaining solutions are possible for a wide set of potential domain specialties, so long as costs start out reasonably low. In particular, Theorem 4.1 would generalize to the multi-specialist fine-tuning, where if the generalist and at least one specialist meet the marginal cost conditions stated in the Theorem, the players would have at least one profit-sharing solution that does not consist in free-riding and Pareto-dominates disagreement.

5.3 Characterizing Domain Specialists' Strategies

When there are potentially many domains where a technology may prove useful or marketable, different strategies around investment levels and fine-tuning can arise. In some domains, a technology may be adopted 'as-is' without significant additional investment or specialization. In other domains, it might be in everyone's interest for a technology to receive significant investment and specialization. Finally, in some domains, a technology might not be viable for any use at all. In this section, we explore the different sorts of cooperation (or non-cooperation) that might arise in domains with different characteristics. Our next general finding is a theorem on the different regimes of domain-specialist strategies, depending on particular attributes of revenue and cost functions.

First, we'll offer a set of relevant definitions to help characterize the different possible regimes of strategies for the specialist. Then, we'll state the formal theorem.

Definition 5.1 (Contributor). A domain specialist D_i is a **contributor** at the profit-sharing agreement δ if, given the generalist optimal investment α_0 at δ , D_i 's optimal strategy is to bring the technology to performance $\alpha_i^* > \alpha_0$.

Definition 5.2 (Free-rider). A domain specialist D_i is a **free-rider** at the profit-sharing agreement δ if, given the generalist optimal investment α_0 at δ , D_i 's optimal strategy is to enter the deal without improving its performance, so $\alpha_i^* = \alpha_0$.

Definition 5.3 (Abstainer). A domain specialist D_i is an **abstainer** at the profit-sharing agreement δ if, given the generalist optimal investment α_0 at δ , D_i 's optimal strategy is to exit the deal and opt for disagreement.

Notice that any specialist with any cost and revenue function is inevitably either a contributor, free-rider, or abstainer. These three regimes span the possible strategies for D_i in the fine-tuning game. In the Theorem below, we outline conditions that characterize a specialist's strategy depending on their domain's cost and revenue functions $\{r, \phi_i\}$.

Theorem 5.2. Say a generalist has produced a general-purpose technology operating at performance level α_0 and available at profit-sharing parameter δ . For any specialist with utility unimodal in α_i , the following two conditions characterize their strategy, as shown in Table 2.

- "Fixed Costs Under Control": At zero investment $(\alpha_i = \alpha_0)$, the domain specialist i's cost is less than its share of the revenue. Formally, $r_i(\alpha_0) > \frac{1}{1-\delta}\phi_i(\alpha_0)$.
- "Marginally Profitable Investment": At zero investment ($\alpha_i = \alpha_0$), a marginal investment from the domain specialist i increases its revenue share more than its costs. Formally, $r'_i(\alpha_0) > \frac{1}{1-\delta}\phi'_i(\alpha_0)$.

"Fixed Costs Under Control"	"Marginally Profitable Investment"	Type of
$r_i(\alpha_0) > \frac{1}{1-\delta}\phi_i(\alpha_0)$	$r_i'(\alpha_0) > \frac{1}{1-\delta}\phi_i'(\alpha_0)$	Specialist
T	T	Contributor
${ m T}$	F	Free-rider
\mathbf{F}	${ m T}$	Contributor or Abstainer*
F	F	Abstainer

Table 2: Types of specialists. In the third case (*), more information would be needed to conclude whether the specialist is a contributor or abstainer.

A proof of the above theorem is provided in Appendix 10.4. The requirement that specialist utility is unimodal in α_i is, in our view, quite natural and broad. It covers three possible scenarios: 1) specialist utility is increasing with investment, 2) specialist utility is decreasing with investment, or 3) specialist utility increases with investment up to a certain point, beyond which any further investment is not cost-justified.

It is important to note that the three 'regimes' defined in this section can describe a specialist's strategy in either the 1-specialist or multi-specialist fine-tuning game. In the 1-specialist case, the potential strategies describe counter-factuals that depend on the particular cost and revenue functions of the specialist. In the multi-specialist game, all of these regimes are ways of grouping domains and all can exist simultaneously.




Figure 7: Examples illustrating Theorem 5.2. When fixed costs are under control and investment is marginally profitable (upper left), the domain specialist will contirbute to the technology. When fixed costs are under control but investment is not marginally profitable (upper right), the domain specialist will free-ride. When fixed costs are too high and investment is marginally costly (lower right), the domain specialist will not bring the technology to market. Finally, when the fixed costs are too high but investment yields marginal returns (lower left quadrant), the domain specialist might abstain or contribute, depending on whether revenue sufficiently exceeds cost at any level of investment. In this final case, the specialist prefers abstaining to free-riding.

One scenario portrayed in Table 2 does not determine cleanly which regime the specialist falls into. In the scenario labeled with an asterisk (*), fixed costs are not under control but it is marginally profitable to invest in the technology. At zero investment, the technology is not ready to bring to market profitably, and it is unclear only from the marginal return on an initial investment whether it is worthwhile for the specialist to invest. In this case, the technology is potentially viable with some non-zero effort spend or, alternatively, not viable for the domain at any level of investment. More information would be needed to conclude whether the specialist would contribute. In particular, if $(1 - \delta)r_i(\alpha_i) - \phi_i(\alpha_i)$ has positive real roots (above α_0), then we could conclude that the technology is viable.

An illustration of Theorem 5.2 is provided in Figure 7. The indeterminate case contains two possible scenarios, one where the specialist would abstain and one where the specialist would contribute. A neat feature of this result is that these behaviors about particular domain adaptations depend only on attributes about the domain around $\alpha_i = \alpha_0$. It uses only 0th- and 1st-order approximations of $U_{D_i}|_{\alpha_i = \alpha_0}$, when the domain has invested no investment into the technology.

This theorem perhaps explains why technologies see significant uptake in some domains and not others. It could, potentially, help identify domains that are particularly likely or unlikely to adopt a general purpose technology. It also may explain why some technologies are re-sold without additional investment while others require significant fine-tuning.

6 Discussion

Our theoretical model of fine-tuning provides a way of understanding the process whereby firms adapt a general-purpose technology to a particular domain. Notably, the model highlights some of the *normative* considerations in these arrangements. There are many potential mechanisms for distributing surplus between technological developers. Different bargaining solutions can highlight the ways that developers share upsides, with implications for a technology's performance (i.e., the value provided to the consumer).

Our model provides a starting point for considering the different interests and choices involved in the development of general-purpose models. By putting forward this model, we attempt to invoke the political economy of the development of general AI technologies. These technologies are produced by a number of entities with different interests, and may potentially affect many individuals. This paper models agents' different interests explicitly, and proposes different methods for weighing between them in light of societal values.

Further, our model may prove useful in suggesting economic and incentive-based levers that might be available to the regulatory process. Interventions and incentives could help align the production of general-purpose models with social welfare.

6.1 Examples of Fine-Tuning and Adaptation Processes

A number of real-world scenarios may be described as a negotiation between a general-purpose technology producer and one or more specialist(s) seeking to adapt the technology to a particular domain. Our model and findings may apply in any case where a computational technology relies on adaptation in order to reach the market. We discuss a few examples in this section.

Generative AI Models: The development of machine learning models is the main motivation for our paper, as discussed above. Recent developments in ML have yielded new tools and capabilities including text and image generation. Large language models and generative models trained on image data have been developed by a handful of technology companies including OpenAI and Google. Their development is, for the most part, untethered to particular domain contexts or commodities, though a number of potential uses are imaginable. These uses include, for example, automating customer service intake, writing tasks, creating design mock-ups, and more. Task-specific model tweaks (i.e., fine-tuning) may be needed to use these models in a certain domain. Other adaptation steps include software and hardware integration. These tweaks and steps are performed by entities other than the general technology provider.

Digital Marketplaces: Digital marketplaces can be thought of as technologies that support and enable specialist producers to bring valuable products to consumers. For example, Apple's App Store is an intermediary between producers and a massive consumer audience. An adaptation model of the app store

might consider the iPhone hardware as a general-purpose technology that can be tweaked and improved by new software apps with particular capabilities. Instead of building all this software in-house, Apple opens the app store to developers who can tweak the technology (i.e., build applications). Apple takes a fixed 30% profit-sharing cut on all app purchases (Roma and Ragaglia, 2016), an arrangement leaving all parties (Apple, developers, and consumers) with some value payoff. Opening up a technology to independent developers is extremely common in digital marketplaces: web hosting, retail, video streaming – all these marketplaces might be described as adapting a general, market-making technology.

Additive Manufacturing: Printing three-dimensional objects has become a mainstay technology developed over the past 50 years. 3D printers have become available as commodities for hobbyists, companies, and research labs to use. The process of 3D printing consists of creating a model and printing in one or more materials (plastics, metals, ceramics) according to the specified model (Gibson et al., 2021). The technology can be thought of as a general-purpose technology, where adaptors may buy the technology in order to make downstream, specialized goods like toys, car parts or other goods. Consumer 3D printers are bought with fixed prices while large-scale, industrial manufacturers might conceivably bargain over hardware (e.g., Roos and Fusco (2014)).

Cloud Computing Infrastructure: On-demand computing resources have revolutionized internet-mediated software services. This technology uses shared computer systems distributed over potentially many locations. A cloud service provider like Amazon Web Services sells computing resources on-demand to software companies who wish to perform operations, like manipulating a database, potentially subject to fluctuations in demand for servers. As software companies have evolved, different sorts of workloads are needed, and cloud services have had to invest in optimizing their technology in order to support clients' needs. For example, cloud computing has moved from fixed computing resources to kubernetes, allowing a more distributed and fluid computing access (Bernstein, 2014). The cloud computing market can be thought of as an adaptation process where a provider like AWS produces a general technology that enables specific software companies to function. The specific companies, then, can be thought of as adaptors of cloud computing technology.

6.2 Conclusion

In this work, we introduced the notion of a *fine-tuning game* as a way of modeling the economic interaction between a firm creating a general-purpose technology and one or more additional firms that specialize the technology to specific domains. We have shown how this formalism leads to an interesting class of bargaining games with rich structure, with insights into how the relative capabilities of the different parties translate into the effort invested in development and the eventual performance of the technology.

The work suggests a number of interesting directions for further research. One direction is to identify further general existence results for bargaining solutions with general functions in this model. More broadly, we also believe that formalizing the societal interests involved in AI regulation is an important direction; such a formalism would need to build on an underlying model that contains the economic interests of the firms producing the AI technology. Our model may therefore help form the foundation for such work.

References

Rediet Abebe, Solon Barocas, Jon Kleinberg, Karen Levy, Manish Raghavan, and David G Robinson. 2020. Roles for computing in social change. In *Proceedings of the 2020 conference on fairness, accountability, and transparency*. 252–260.

David Bernstein. 2014. Containers and cloud: From lxc to docker to kubernetes. *IEEE cloud computing* 1, 3 (2014), 81–84.

Rishi Bommasani, Drew A Hudson, Ehsan Adeli, Russ Altman, Simran Arora, Sydney von Arx, Michael S Bernstein, Jeannette Bohg, Antoine Bosselut, Emma Brunskill, et al. 2021. On the opportunities and risks of foundation models. arXiv preprint arXiv:2108.07258 (2021).

Rishi Bommasani, Dilara Soylu, Thomas I Liao, Kathleen A Creel, and Percy Liang. 2023. Ecosystem Graphs: The Social Footprint of Foundation Models. arXiv preprint arXiv:2303.15772 (2023).

- Timothy Bresnahan. 2010. General purpose technologies. *Handbook of the Economics of Innovation* 2 (2010), 761–791.
- Benedetta Brevini and Frank Pasquale. 2020. Revisiting the Black Box Society by rethinking the political economy of big data., 2053951720935146 pages.
- Kalyan Chatterjee, Bhaskar Dutta, Debraj Ray, and Kunal Sengupta. 1993. A noncooperative theory of coalitional bargaining. The Review of Economic Studies 60, 2 (1993), 463–477.
- Robert G Cooper. 1984. The strategy-performance link in product innovation. *R&D Management* 14, 4 (1984), 247–259.
- Nicholas Crafts. 2021. Artificial intelligence as a general-purpose technology: an historical perspective. Oxford Review of Economic Policy 37, 3 (2021).
- Tyna Eloundou, Sam Manning, Pamela Mishkin, and Daniel Rock. 2023. Gpts are gpts: An early look at the labor market impact potential of large language models. arXiv preprint arXiv:2303.10130 (2023).
- Nanyi Fei, Zhiwu Lu, Yizhao Gao, Guoxing Yang, Yuqi Huo, Jingyuan Wen, Haoyu Lu, Ruihua Song, Xin Gao, Tao Xiang, et al. 2022. Towards artificial general intelligence via a multimodal foundation model. *Nature Communications* 13, 1 (2022), 3094.
- Alfonso Gambardella and Anita M McGahan. 2010. Business-model innovation: General purpose technologies and their implications for industry structure. Long range planning 43, 2-3 (2010), 262–271.
- Ian Gibson, David W Rosen, Brent Stucker, Mahyar Khorasani, David Rosen, Brent Stucker, and Mahyar Khorasani. 2021. Additive manufacturing technologies. Vol. 17. Springer.
- Avi Goldfarb, Bledi Taska, and Florenta Teodoridis. 2023. Could machine learning be a general purpose technology? a comparison of emerging technologies using data from online job postings. *Research Policy* 52, 1 (2023), 104653.
- Beliz Gunel, Jingfei Du, Alexis Conneau, and Ves Stoyanov. 2020. Supervised contrastive learning for pre-trained language model fine-tuning. arXiv preprint arXiv:2011.01403 (2020).
- Oliver Hart and John Moore. 1988. Incomplete contracts and renegotiation. *Econometrica: Journal of the Econometric Society* (1988), 755–785.
- Elhanan Helpman. 1998. General purpose technologies and economic growth. MIT press.
- Jeremy Howard and Sebastian Ruder. 2018. Universal language model fine-tuning for text classification. arXiv preprint arXiv:1801.06146 (2018).
- Boyan Jovanovic and Peter L Rousseau. 2005. General purpose technologies. In *Handbook of economic growth*. Vol. 1. Elsevier, 1181–1224.
- Ehud Kalai and Dov Samet. 1985. Monotonic solutions to general cooperative games. *Econometrica: Journal of the Econometric Society* (1985), 307–327.
- Ehud Kalai and Meir Smorodinsky. 1975. Other solutions to Nash's bargaining problem. *Econometrica: Journal of the Econometric Society* (1975), 513–518.
- Ananya Kumar, Aditi Raghunathan, Robbie Jones, Tengyu Ma, and Percy Liang. 2022. Fine-tuning can distort pretrained features and underperform out-of-distribution. arXiv preprint arXiv:2202.10054 (2022).
- Richard G Lipsey, Kenneth I Carlaw, and Clifford T Bekar. 2005. Economic transformations: general purpose technologies and long-term economic growth. Oup Oxford.
- Hervi Moulin. 1991. Axioms of cooperative decision making. Number 15. Cambridge university press.
- John F Nash. 1950. The bargaining problem. Econometrica: Journal of the econometric society (1950), 155–162.

- Long Ouyang, Jeffrey Wu, Xu Jiang, Diogo Almeida, Carroll Wainwright, Pamela Mishkin, Chong Zhang, Sandhini Agarwal, Katarina Slama, Alex Ray, et al. 2022. Training language models to follow instructions with human feedback. *Advances in Neural Information Processing Systems* 35 (2022), 27730–27744.
- Xudong Pan, Mi Zhang, Shouling Ji, and Min Yang. 2020. Privacy risks of general-purpose language models. In 2020 IEEE Symposium on Security and Privacy (SP). IEEE, 1314–1331.
- Matthew E Peters, Sebastian Ruder, and Noah A Smith. 2019. To tune or not to tune? adapting pretrained representations to diverse tasks. arXiv preprint arXiv:1903.05987 (2019).
- William P Rogerson. 1992. Contractual solutions to the hold-up problem. The Review of Economic Studies 59, 4 (1992), 777–793.
- Paolo Roma and Daniele Ragaglia. 2016. Revenue models, in-app purchase, and the app performance: Evidence from Apple's App Store and Google Play. *Electronic commerce research and applications* 17 (2016), 173–190.
- Göran Roos and Mark Fusco. 2014. Strategic implications of additive manufacturing (AM) on traditional industry business models. In *Proceedings of Additive Manufacturing with Powder Metallurgy Conference*, Orlando.
- Victor Sanh, Thomas Wolf, and Alexander Rush. 2020. Movement pruning: Adaptive sparsity by fine-tuning. Advances in Neural Information Processing Systems 33 (2020), 20378–20389.
- Amartya Sen. 2018. Collective choice and social welfare. Harvard University Press.
- Nima Tajbakhsh, Jae Y Shin, Suryakanth R Gurudu, R Todd Hurst, Christopher B Kendall, Michael B Gotway, and Jianming Liang. 2016. Convolutional neural networks for medical image analysis: Full training or fine tuning? *IEEE transactions on medical imaging* 35, 5 (2016), 1299–1312.
- Manuel Trajtenberg. 2018. AI as the next GPT: a Political-Economy Perspective. Technical Report. National Bureau of Economic Research.
- W Kip Viscusi and Michael J Moore. 1993. Product liability, research and development, and innovation. *Journal of Political Economy* 101, 1 (1993), 161–184.
- Yu-Xiong Wang, Deva Ramanan, and Martial Hebert. 2017. Growing a brain: Fine-tuning by increasing model capacity. In *Proceedings of the IEEE Conference on Computer Vision and Pattern Recognition*. 2471–2480.
- Jason Wei, Yi Tay, Rishi Bommasani, Colin Raffel, Barret Zoph, Sebastian Borgeaud, Dani Yogatama, Maarten Bosma, Denny Zhou, Donald Metzler, et al. 2022. Emergent abilities of large language models. arXiv preprint arXiv:2206.07682 (2022).
- Tianyi Zhang, Felix Wu, Arzoo Katiyar, Kilian Q Weinberger, and Yoav Artzi. 2020. Revisiting few-sample BERT fine-tuning. arXiv preprint arXiv:2006.05987 (2020).
- Fuzhen Zhuang, Zhiyuan Qi, Keyu Duan, Dongbo Xi, Yongchun Zhu, Hengshu Zhu, Hui Xiong, and Qing He. 2020. A comprehensive survey on transfer learning. *Proc. IEEE* 109, 1 (2020), 43–76.

7 Section 2 Materials

7.1 Pareto set characterization and Theorem 2.1

Proof of Theorem 2.1. Consider three non-overlapping intervals that collectively span the feasible set $\delta \in [0, 1]$. These intervals are:

- 1. $0 \le \delta < \min(\delta^{\text{Powerful } D}, \delta^{\text{Powerful } G})$
- 2. $\min(\delta^{\text{Powerful } D}, \delta^{\text{Powerful } G}) \leq \delta \leq \max(\delta^{\text{Powerful } D}, \delta^{\text{Powerful } G})$
- 3. $\max(\delta^{\text{Powerful }D}, \delta^{\text{Powerful }G}) < \delta \leq 1$

We will characterize each of these intervals in turn, finding that intervals (1) and (3) are always Pareto dominated, and interval (2) is characterized by a trade-off in utilities.

- 1. Within interval (1), the domain is characterized by $\delta < \min(\delta^{\text{Powerful }D}, \delta^{\text{Powerful }G}) \Rightarrow \delta < \delta^{\text{Powerful }D}$ and $\delta < \delta^{\text{Powerful }G}$. By the definition of a strictly unimodal function (2.3), this means that both utility functions $\{U_D, U_G\}$ are strictly increasing over interval 1. Thus, there exists some quantity $\epsilon > 0$ such that, for any value δ in interval (1), $U_D(\delta + \epsilon) > U_D(\delta)$ and $U_G(\delta + \epsilon) > U_G(\delta)$. Thus, every potential agreement in interval (1) is Pareto-dominated.
- 2. Within interval (2), the domain is characterized by $\min(\delta^{\operatorname{Powerful}\ D}, \delta^{\operatorname{Powerful}\ G}) \leq \delta$, and also $\delta \leq \max(\delta^{\operatorname{Powerful}\ D}, \delta^{\operatorname{Powerful}\ G})$. If $\delta^{\operatorname{Powerful}\ D} = \delta^{\operatorname{Powerful}\ G}$, then the value $\delta = \delta^{\operatorname{Powerful}\ D} = \delta^{\operatorname{Powerful}\ G}$ is the unique Pareto-optimal agreement because it is optimal for both players. Otherwise if $\delta^{\operatorname{Powerful}\ D} \neq \delta^{\operatorname{Powerful}\ G}$, then interval (2) can be characterized as follows: For one player $P \in \{G, D\}$, the utility U_P one utility function is strictly decreasing because $\delta \geq \delta^{\operatorname{Powerful}\ P}$ and $U_P(\delta)$ is a strictly unimodal function. For the other player $\{G, D\} \setminus P$, the utility $U_{\{G, D\} \setminus P}$ is strictly increasing because $\delta \leq \delta^{\operatorname{Powerful}\ \{G, D\} \setminus P}$ and $U_{\{G, D\} \setminus P}(\delta)$ is a strictly unimodal function. Since one player's utility is strictly increasing and the other's is strictly decreasing, any perturbation of δ within interval (2) constitutes a utility gain for one player and a utility loss for the other. For any value of δ within this interval, if both players' utilities exceed the disagreement payoff (i.e., positive utility), then δ is Pareto-optimal.
- 3. Within interval (3), the domain is characterized by $\delta > \max(\delta^{\text{Powerful }D}, \delta^{\text{Powerful }G}) \Rightarrow \delta > \delta^{\text{Powerful }D}$ and $\delta > \delta^{\text{Powerful }G}$. By the definition of a strictly unimodal function (2.3), this means that both utility functions $\{U_D, U_G\}$ are strictly decreasing over interval (3). Thus, there exists some quantity $\epsilon > 0$ such that, for any value δ in interval (3), $U_D(\delta \epsilon) > U_D(\delta)$ and $U_G(\delta \epsilon) > U_G(\delta)$. Thus, every potential agreement in interval (3) is Pareto-dominated.

Thus interval (2) is Pareto-efficient among the set of feasible bargaining agreements.

8 Section 3 Materials (Polynomial fine-tuning game)

8.1 Subgame perfect equilibrium findings

Proof of Theorem 3.1. We solve the game using backward induction as follows: First, starting with the last stage (3), we solve for α_1^* given α_0, δ, c_1 :

$$\alpha_{1}^{*} = \operatorname{argmax}_{\alpha_{1}} U_{D}(\alpha_{1}, \alpha, \delta)$$

$$\Rightarrow \frac{\partial U_{D}}{\partial \alpha_{1}} \Big|_{\alpha_{1} = \alpha_{1}^{*}} = 0$$

$$\Rightarrow \frac{\partial}{\partial \alpha_{1}} \left((1 - \delta)\alpha_{1} - c_{1}(\alpha_{1} - \alpha_{0})^{k_{1}} \right) \Big|_{\alpha_{1} = \alpha_{1}^{*}} = 0$$

$$\Rightarrow (1 - \delta) - k_{1}c_{1}(\alpha_{1}^{*} - \alpha_{0})^{k_{1} - 1} = 0$$

$$\Rightarrow \alpha_{1}^{*} = \alpha_{0} + \left(\frac{1 - \delta}{k_{1}c_{1}} \right)^{\frac{1}{k_{1} - 1}}.$$
(11)

Note that $\frac{\partial^2 U_D}{\partial \alpha_1^2} = -k_1(k_1 - 1)c_1(\alpha_1 - \alpha_0)^{k_1 - 2}$. This quantity is negative as long as k > 1, which is assumed. Thus, the α_1^* derived above yields a global maximum of U_D .

Second, knowing D's choice of α_1^* above, we solve for α_0^* as follows:

$$\alpha_0^* = \operatorname{argmax}_{\alpha_0} U_G(\alpha_0, \delta)$$

$$\Rightarrow \frac{\partial U_G}{\partial \alpha_0} \Big|_{\alpha_0 = \alpha_0^*} = 0$$

$$\Rightarrow \frac{\partial}{\partial \alpha_0} \left(\delta \alpha_1^* - c_0 \alpha_0^{k_0} \right) \Big|_{\alpha_0 = \alpha_0^*} = 0$$

$$\Rightarrow \frac{\partial}{\partial \alpha_0} \left(\delta \left(\alpha_0 + \frac{1 - \delta}{2c_1} \right) - c_0 \alpha_0^{k_0} \right) \Big|_{\alpha_0 = \alpha_0^*} = 0$$

$$\Rightarrow \frac{\partial}{\partial \alpha_0} \left(\delta \alpha_0 + [\operatorname{const}] - c_0 \alpha_0^{k_0} \right) \Big|_{\alpha_0 = \alpha_0^*} = 0$$

$$\Rightarrow \delta - k_0 c_0 (\alpha_0^*)^{k_0 - 1} = 0$$

$$\Rightarrow \alpha_0^* = \left(\frac{\delta}{k_0 c_0} \right)^{\frac{1}{k_0 - 1}}.$$

The second derivative $\frac{\partial^2 U_G}{\partial \alpha^2} = -k_0(k_0 - 1)c_0(\alpha_0)^{k_0 - 2}$. This quantity is negative as long as k > 1, which is assumed. Thus, the value of α_0^* derived above yields a global maximum of U_G .

Finally, plugging in $\alpha_0^* = \left(\frac{\delta}{k_0 c_0}\right)^{\frac{1}{k_0 - 1}}$ into Equation 11, we obtain the following expression for α_1^* as a function of δ only:

$$\alpha_1^* = \left(\frac{\delta}{k_0 c_0}\right)^{\frac{1}{k_0 - 1}} + \left(\frac{1 - \delta}{k_1 c_1}\right)^{\frac{1}{k_1 - 1}}.$$

This finishes the proof.

8.2 Utilities as a function of δ

Proof of Corollary 3.1. Plugging the formulas from Theorem 3.1 into Equation 1, we obtain:

$$\begin{split} U_G &= \delta\alpha_1 - \phi_0(\alpha_0) \\ &= \delta\left(\left(\frac{\delta}{k_0c_0}\right)^{\frac{1}{k_0-1}} + \left(\frac{1-\delta}{k_1c_1}\right)^{\frac{1}{k_1-1}}\right) - c_0\left(\left(\frac{\delta}{k_0c_0}\right)^{\frac{1}{k_0-1}}\right)^{k_0} \\ &= \delta\left(\frac{\delta}{k_0c_0}\right)^{\frac{1}{k_0-1}} + \delta\left(\frac{1-\delta}{k_1c_1}\right)^{\frac{1}{k_1-1}} - c_0\left(\frac{\delta}{k_0c_0}\right)^{\frac{k_0}{k_0-1}} \\ &= \left[\left(\frac{1}{k_0c_0}\right)^{\frac{1}{k_0-1}} - c_0\left(\frac{1}{k_0c_0}\right)^{\frac{k_0}{k_0-1}}\right] \delta^{\frac{k_0}{k_0-1}} + \left(\frac{1}{k_1c_1}\right)^{\frac{1}{k_1-1}} \delta(1-\delta)^{\frac{1}{k_1-1}} \\ &= \left(\frac{1}{k_0c_0}\right)^{\frac{1}{k_0-1}} \left(1 - \frac{1}{k_0}\right) \delta^{\frac{k_0}{k_0-1}} + \left(\frac{1}{k_1c_1}\right)^{\frac{1}{k_1-1}} \delta(1-\delta)^{\frac{1}{k_1-1}}. \end{split}$$

Plugging the formulas from Theorem 3.1 into Equation 2, we obtain:

$$\begin{split} U_D &= \qquad (1-\delta)\alpha_1 - \phi_i(\alpha_1;\alpha_0) \\ &= \qquad (1-\delta)\left[\left(\frac{\delta}{k_0c_0}\right)^{\frac{1}{k_0-1}} + \left(\frac{1-\delta}{k_1c_1}\right)^{\frac{1}{k_1-1}}\right] - c_1\left[\left(\frac{\delta}{k_0c_0}\right)^{\frac{1}{k_0-1}} + \left(\frac{1-\delta}{k_1c_1}\right)^{\frac{1}{k_1-1}} - \left(\frac{\delta}{k_0c_0}\right)^{\frac{1}{k_0-1}}\right]^{k_1} \\ &= \qquad (1-\delta)\left(\frac{\delta}{k_0c_0}\right)^{\frac{1}{k_0-1}} + (1-\delta)\left(\frac{1-\delta}{k_1c_1}\right)^{\frac{1}{k_1-1}} - c_1\left(\frac{1-\delta}{k_1c_1}\right)^{\frac{k_1}{k_1-1}} \\ &= \qquad \left(\frac{1}{k_0c_0}\right)^{\frac{1}{k_0-1}} (1-\delta)\delta^{\frac{1}{k_0-1}} + \left(\frac{1}{k_1c_1}\right)^{\frac{1}{k_1-1}} (1-\delta)^{\frac{k_1}{k_1-1}} - \left(\frac{1}{k_1c_1}\right)^{\frac{1}{k_1-1}} \left(\frac{1}{k_1}\right) (1-\delta)^{\frac{k_1}{k_1-1}} \\ &= \qquad \left(\frac{1}{k_1c_1}\right)^{\frac{1}{k_1-1}} \left(1-\frac{1}{k_1}\right) (1-\delta)^{\frac{k_1}{k_1-1}} + \left(\frac{1}{k_0c_0}\right)^{\frac{1}{k_0-1}} (1-\delta)\delta^{\frac{1}{k_0-1}}. \end{split}$$

8.3 Utilities are strictly unimodal functions of δ

Proof of Proposition 3.1. We'll start by proving U_G is strictly unimodal, and then extend the results to U_D . Beginning with U_G : The proof relies on the following Lemma:

Lemma 8.1. A differentiable continuous function $f(\delta)$ is strictly unimodal over $\delta \in [a, b]$ if the following conditions are met: i) f'(a) > 0, ii) f' is concave over the domain.

Proof of Lemma 8.1. By the definition of strict unimodality, we can conclude a function $f(\delta)$ is strictly unimodal over an interval [a,b] if one of the following properties hold: 1) $f'(\delta) > 0 \,\,\forall\,\,\delta \in [0,1]$, meaning the function is strictly increasing over the interval, or 2) For some value $c \in (a,b)$, the function is strictly increasing for values [a,c) and strictly decreasing for values (c,b]. Notice that, so long as condition (i) holds (i.e., the function starts out strictly increasing at a), the function is strictly unimodal as long as its derivative crosses the f'=0 axis at no more than one point in [a,b]. So, the remainder of the proof finds a contradiction when we assume conditions (i) and (ii) and that there are two values in (a,b) for which f'=0.




Figure 8: Illustration of the proof for Lemma 8.1. If the derivative of a function f is positive at a and concave, it cannot cross the axis more than 1 time, meaning f is strictly unimodal.

Consider the curve $f' = \frac{df}{d\delta}$. We specify a point on this curve using $\{(x,y)|x=\delta,y=\frac{df}{d\delta}\}$. Given condition (i), There is some point $x_0=(a,p)$ where p>0 and $x_0\in\frac{df}{d\delta}$. Assume for the sake of contradiction that there are two points $x_1=(q,0), x_2=(r,0)$ where $x_1,x_2\in\frac{df}{d\delta}$ and a< q< r< b (without loss of generality). We can plug these three points into the definition of concavity and find our contradiction: First, notice

 $l = \frac{q-a}{r-a} \in (0,1)$ because a < q < r. Next, plugging in the definition of concavity:

$$\frac{df}{d\delta}((1-l)a+lr) \geq (1-l)\frac{df}{d\delta}(a) + l\frac{df}{d\delta}(r)$$

$$\frac{df}{d\delta}\left(\left(1 - \frac{q-a}{r-a}\right)a + \frac{q-a}{r-a}r\right) \geq \left(1 - \frac{q-a}{r-a}\right)[p] + \frac{q-a}{r-a}[0]$$

$$\frac{df}{d\delta}\left(a - \frac{q-a}{r-a}a + \frac{q-a}{r-a}r\right) \geq p - \frac{q-a}{r-a}p$$

$$\frac{df}{d\delta}\left(a + (r-a)\frac{q-a}{r-a}\right) \geq p - \frac{q-a}{r-a}p$$

$$\frac{df}{d\delta}\left(a + (r-a)\frac{q-a}{r-a}\right) \geq p - \frac{q-a}{r-a}p$$

$$\frac{df}{d\delta}\left(a + (q-a)\right) \geq p - \frac{q-a}{r-a}p$$

$$\frac{df}{d\delta}\left(a\right) \geq p - \frac{q-a}{r-a}p$$

$$0 \geq p - \frac{q-a}{r-a}p > 0.$$

Hence the contradiction: We know $p - \frac{q-a}{r-a}p > 0$ is strictly greater than 0 because p > 0 and $l \in (0,1)$. Thus a function characterized by conditions (i) and (ii) cannot contain these three points. This concludes Lemma 8.1's proof: If f' is concave, and starts positive at a, it must cross the axis at most once meaning f is unimodal.

Now, we prove the utilities are unimodal by showing that $U_G(\delta)$ satisfies both conditions in Lemma 8.1 for the domain $\delta \in [0, 1]$.

We first differentiate U_G with respect to δ .

$$\frac{\partial U_G}{\partial \delta} = \frac{\partial}{\partial \delta} \left[\left(\frac{1}{k_0 c_0} \right)^{\frac{1}{k_0 - 1}} \left(1 - \frac{1}{k_0} \right) \delta^{\frac{k_0}{k_0 - 1}} + \left(\frac{1}{k_1 c_1} \right)^{\frac{1}{k_1 - 1}} \delta(1 - \delta)^{\frac{1}{k_1 - 1}} \right]
= \frac{\partial}{\partial \delta} \left[A \delta^{\frac{k_0}{k_0 - 1}} + B \delta(1 - \delta)^{\frac{1}{k_1 - 1}} \right],$$

where $A := \left(\frac{1}{k_0c_0}\right)^{\frac{1}{k_0-1}} \left(1 - \frac{1}{k_0}\right) > 0$ and $B := \left(\frac{1}{k_1c_1}\right)^{\frac{1}{k_1-1}} > 0$. A is positive as long as $k_0 > 1$ and $c_0 > 0$, which is given. B is positive as long as $k_1 > 1$ and $c_1 > 0$, which is given. Continuing:

$$= A\left(\frac{k_0}{k_0-1}\right)\delta^{\frac{1}{k_0-1}} + B\left[-\delta\left(\frac{1}{k_1-1}\right)(1-\delta)^{\frac{1}{k_1-1}-1} + (1-\delta)^{\frac{1}{k_1-1}}\right]$$

$$= A\left(\frac{k_0}{k_0-1}\right)\delta^{\frac{1}{k_0-1}} + \frac{(1-\delta)^{\frac{1}{k_1-1}-1}}{k_1-1}\left[-\delta + (k_1-1)(1-\delta)\right]$$

$$= A\left(\frac{k_0}{k_0-1}\right)\delta^{\frac{1}{k_0-1}} + \frac{(1-\delta)^{\frac{1}{k_1-1}-1}}{k_1-1}\left[k_1-k_1\delta-1+\delta-\delta\right]$$

$$= A\left(\frac{k_0}{k_0-1}\right)\delta^{\frac{1}{k_0-1}} + \frac{(1-\delta)^{\frac{1}{k_1-1}-1}}{k_1-1}\left[k_1(1-\delta)-1\right]$$

$$\frac{\partial U_G}{\partial \delta} = A\left(\frac{k_0}{k_0-1}\right)\delta^{\frac{1}{k_0-1}} + B\left(\frac{k_1}{k_1-1}\right)(1-\delta)^{\frac{1}{k_1-1}} - B\left(\frac{1}{k_1-1}\right)(1-\delta)^{\frac{1}{k_1-1}-1}.$$

Now we can show the first condition (a) in Lemma 8.1 holds:

$$\left. \frac{\partial U_G}{\partial \delta} \right|_{\delta=0} = [0] + B\left(\frac{k_1}{k_1 - 1}\right) - B\left(\frac{1}{k_1 - 1}\right) = B\left(\frac{k_1 - 1}{k_1 - 1}\right) = B > 0.$$

To show the second condition (b) in Lemma 8.1 holds, we perform the second-derivative test, which requires differentiating the function two more times:

$$\frac{\partial^{3} U_{G}}{\partial \delta^{3}} = A \left(\frac{k_{0}}{k_{0}-1} \right) \left(\frac{1}{k_{0}-1} \right) \left(\frac{1}{k_{0}-1} - 1 \right) \delta^{\frac{1}{k_{0}-1}-2}$$

$$+ B \left(\frac{k_{1}}{k_{1}-1} \right) \left(\frac{1}{k_{1}-1} \right) \left(\frac{1}{k_{1}-1} - 1 \right) (1-\delta)^{\frac{1}{k_{1}-1}-2}$$

$$- B \left(\frac{1}{k_{1}-1} \right) \left(\frac{1}{k_{1}-1} - 1 \right) \left(\frac{1}{k_{1}-1} - 2 \right) (1-\delta)^{\frac{1}{k_{1}-1}-3}.$$

The above expression is never positive. First, notice all three coefficients are less than or equal to zero:

- $A(\frac{k_0}{k_0-1})(\frac{1}{k_0-1})(\frac{1}{k_0-1}-1)$ is the product of one negative and otherwise non-negative numbers: Given $k_0 \ge 2$, observe A > 0, $(\frac{k_0}{k_0-1}) > 0$, $(\frac{1}{k_0-1}) > 0$, $(\frac{1}{k_0-1}-1) \le 0$.
- $B(\frac{k_1}{k_1-1})(\frac{1}{k_1-1})(\frac{1}{k_1-1}-1)$ is the product of one negative and otherwise non-negative numbers: Given $k_1 \ge 2$, observe B > 0, $(\frac{k_1}{k_1-1}) > 0$, $(\frac{1}{k_1-1}) > 0$, $(\frac{1}{k_1-1}-1) \le 0$.
- $-B(\frac{1}{k_1-1})(\frac{1}{k_1-1}-1)(\frac{1}{k_1-1}-2)$ is the product of three negative and otherwise non-negative numbers: Given $k_1 \ge 2$, observe -B < 0, $(\frac{1}{k_1-1}) > 0$, $(\frac{1}{k_1-1}-1) \le 0$, $(\frac{1}{k_1-1}-1) < 0$.

Second, notice all three expressions of δ are defined and positive on the interval (0,1):

- $\delta^{\frac{1}{k_0-1}-2}$ is positive and defined $\forall \delta > 0$.
- $(1-\delta)^{\frac{1}{k_1-1}-2}$ is positive and defined $\forall \delta < 1$.
- $(1-\delta)^{\frac{1}{k_1-1}-3}$ is positive and defined $\forall \delta < 1$.

Every term in our derived expression for $\frac{\partial^3 U_G}{\partial U_G^3}$ is non-positive. Thus the function $\frac{\partial U_G}{\partial U_G}$ is concave satisfying condition (b) in Lemma 8.1. This completes the proof that U_G is unimodal.

Moving on to U_D : Notice the formulation of U_G in Equation 5 is almost exactly the same functional form as U_D in Equation 6. If we define a variable $\gamma = (1 - \delta)$, we can use the identical proof completed above to show U_D is unimodal in γ . Since we prove unimodality on the interval [0,1], a function defined over $\gamma \in [0,1]$ is simply a function of $\delta \in [0,1]$ reflected over the vertical line $\delta = 0.5$. A transform that reflects a univariate function over the vertical line passing through the midpoint of its domain preserves strict unimodality.

8.4 Powerful-G Bargaining Solution

Proof of Proposition 3.2. The powerful-G solution is the solution $\delta^{Powerful G}$ that maximizes U_G over the feasible set of $\delta \in [0,1]$:

$$\delta^{Powerful} G = \operatorname{argmax}_{\delta} U_{G}(\delta)$$

$$\Rightarrow \frac{\partial U_{G}}{\partial \delta} = 0$$

$$\Rightarrow \frac{\partial}{\partial \delta} \left[\frac{\delta^{2}}{4c_{0}} + \frac{\delta}{2c_{1}} - \frac{\delta^{2}}{2c_{1}} \right] = 0 \quad Corr. \ 3.1$$

$$\Rightarrow \frac{\delta}{2c_{0}} + \frac{1}{2c_{1}} - \frac{\delta}{c_{1}} = 0$$

$$\Rightarrow \delta \left(\frac{1}{2c_{0}} - \frac{1}{c_{1}} \right) = -\frac{1}{2c_{1}}$$

$$\Rightarrow \delta = -\frac{1}{2c_{1}} \left(\frac{c_{1} - 2c_{0}}{2c_{1}c_{0}} \right)^{-1}$$

$$\Rightarrow \delta = \frac{c_{0}}{2c_{0} - c_{1}}.$$

The second partial derivative $\frac{\partial^2 U_G}{\partial \delta^2} = \frac{1}{2c_0} - \frac{1}{c_1}$, which is negative as long as $0 < \frac{c_1}{c_0} < 2$. Since there is only one root, the derived equation is a global maximum for $0 < c_1 < 2c_0$. However, notice that the derived

expression is only feasible for the values $c_1 \leq c_0$, since the value δ must be in the range [0,1] (Specialist would not take a negative share of the profit). Thus, $\delta^{Powerful\ G} = \frac{1}{2-c_1}$ for $0 < c_1 < c_0$.

The remainder of the proof will show that, for $c_1 \geq c_0$, within the feasible set $0 \leq \delta \leq 1$, $\delta = 1$ maximizes U_G . We'll do so by showing that the partial derivative $\frac{\partial U_G}{\partial \delta}$ is non-negative for all $c_1 \geq c_0$ and $0 \leq \delta \leq 1$. Assume for sake of contradiction:

$$\frac{\partial U_G}{\partial \delta} < 0$$

$$\Rightarrow \frac{\delta}{2c_0} + \frac{1}{2c_1} - \frac{\delta}{c_1} < 0$$

$$\Rightarrow \delta \left(\frac{1}{2c_0} - \frac{1}{c_1}\right) + \frac{1}{2c_1} < 0$$

$$\Rightarrow \delta \left(\frac{c_1 - 2c_0}{2c_1c_0}\right) + \frac{c_0}{2c_1c_0} < 0$$

$$\Rightarrow \delta(c_1 - 2c_0) + c_0 < 0 \quad because \ c_1, c_0 > 0.$$

$$\Rightarrow \delta(c_1 - 2c_0) < -c_0.$$

Notice the resulting expression is only met when $\delta < 0$ or $c_1 \le c_0$. However, we're given $\delta \in [0,1] \cap c_1 \ge c_0$.

Powerful-D Bargaining Solution 8.5

Proof of Proposition 3.3. The powerful-D solution is the solution $\delta^{Powerful D}$ that maximizes U_D over the feasible set:

$$\delta^{Powerful\ D} = \operatorname{argmax}_{\delta} U_D(\delta)$$

$$\Rightarrow \frac{\partial U_D}{\partial \delta} = 0$$

$$\Rightarrow \frac{\partial}{\partial \delta} \left[\left(\frac{1}{4c_1} \right) + \left(\frac{1}{2c_0} - \frac{1}{2c_1} \right) \delta + \left(\frac{1}{4c_1} - \frac{1}{2c_0} \right) \delta^2 \right] = 0 \quad Corr.\ 3.1$$

$$\Rightarrow \left(\frac{1}{2c_0} - \frac{1}{2c_1} \right) + 2 \left(\frac{1}{4c_1} - \frac{1}{2c_0} \right) \delta = 0$$

$$\Rightarrow \left(\frac{1}{2c_1} - \frac{1}{c_0} \right) \delta = \frac{1}{2c_1} - \frac{1}{2c_0}$$

$$\Rightarrow \delta = \frac{\left(\frac{c_0 - c_1}{2c_0 c_1} \right)}{\left(\frac{c_0 - 2c_1}{2c_0 c_1} \right)}$$

$$\Rightarrow \delta = \frac{c_1 - c_0}{2c_0 c_0}.$$

The second partial derivative $\frac{\partial^2 U_D}{\partial \delta^2} = \frac{1}{2c_1} - \frac{1}{c_0}$, which is negative as long as $2c_1 > c_0$. Since there is only one root, the derived equation is a global maximum for $2c_1 > c_0$. However, notice that the derived expression

is only feasible for the values $c_1 \geq c_0$, since the value δ must be in the range [0,1] (Generalist would not take a negative share of the profit). Thus, $\delta^{Powerful\ D} = \frac{c_1 - c_0}{2c_1 - c_0}$ for $c_1 \geq c_0$.

The remainder of the proof will show that, for $c_1 < c_0$, within the feasible set $0 \leq \delta \leq 1$, $\delta = 0$ maximizes U_D . We'll do so by showing that the partial derivative $\frac{\partial U_D}{\partial \delta} \leq 0$ for all $0 < c_1 < 1$ and $0 \leq \delta \leq 1$. Assume for sake of contradiction:

$$\frac{\partial U_D}{\partial \delta} > 0$$

$$\Rightarrow \left(\frac{1}{2c_0} - \frac{1}{2c_1}\right) + \left(\frac{1}{2c_1} - \frac{1}{c_0}\right) \delta > 0$$

$$\Rightarrow \left(\frac{1}{2c_1} - \frac{1}{c_0}\right) \delta > \frac{1}{2c_1} - \frac{1}{2c_0}$$

$$\Rightarrow \left(\frac{c_0 - 2c_1}{2c_0c_1}\right) \delta > \frac{c_0 - c_1}{2c_0c_1}$$

$$\Rightarrow (c_0 - 2c_1) \delta > c_0 - c_1 \qquad because \ c_0, c_1 > 0.$$

We show the contradiction $\forall \frac{c_1}{c_0} \in (0,1)$ (equivalently, every scenario where $0 < c_1 \le c_0$):

- For $\frac{1}{2} < \frac{c_1}{c_0} < 1$, the final step implies $\delta > \frac{c_1 c_0}{2c_1 c_0}$, which contradicts the global optimum finding above.
- For $0 < \frac{c_1}{c_0} < \frac{1}{2}$, the final step implies $\delta \le \frac{c_1 c_0}{2c_1 c_0}$. But notice the right-hand-side must be negative, contradicting the given range $\delta \in [0, 1]$.

• Finally, for $\frac{c_1}{c_0} = \frac{1}{2}$, the final step implies $0 > \frac{1}{2}$.

Thus we've established the contradiction.

8.6 Maximum-performance bargaining solution

Proof of Proposition 3.4. We will show that within the feasible region $\delta \in [0, 1], c_1 > 0$, the following three properties hold:

- 1. $\frac{\partial \alpha_1}{\partial \delta}(c_1) < 0 \quad \forall \quad c_1 < c_0$
- 2. $\frac{\partial \alpha_1}{\partial \delta}(c_1) > 0 \quad \forall \quad c_1 > c_0$
- 3. $\frac{\partial \alpha_1}{\partial \delta}(c_1) = 0$ for $c_1 = c_0$

First, we differentiate our expression for $\alpha_1(\delta; c_0, c_1)$ with respect to δ , using the expression attained in Theorem 3.1:

$$\frac{\frac{\partial \alpha_1}{\partial \delta}}{=\frac{\partial}{\partial \delta} \left[\frac{\delta}{2c_0} + \frac{1-\delta}{2c_1} \right] }$$

$$= \frac{1}{2c_0} - \frac{1}{2c_1}.$$

Now notice each of the three properties are satisfied in turn:

- 1. For $c_1 < c_0$, $\frac{\partial \alpha_1}{\partial \delta} = \frac{1}{2c_0} \frac{1}{2c_1} < 0$.
- 2. For $c_1 > c_0$, $\frac{\partial \alpha_1}{\partial \delta} = \frac{1}{2c_0} \frac{1}{2c_1} > 0$.
- 3. For $c_1 = c_0$, $\frac{\partial \alpha_1}{\partial \delta} = \frac{1}{2c_0} \frac{1}{2c_1} = 0$.

8.7 Vertical monopoly bargaining solution

Proof of Proposition 3.5. The vertical monopoly or "utilitarian" solution is the solution that maximizes the sum of utilities:

$$\delta^{VerticalMonopoly} = \operatorname{argmax}_{\delta} U_{G}(\delta) + U_{D}(\delta)$$

$$\Rightarrow \frac{\partial}{\partial \delta} \left(U_{G}(\delta) + U_{D}(\delta) \right) = 0$$

$$\Rightarrow \frac{\delta}{2c_{0}} + \frac{1}{2c_{1}} - \frac{\delta}{c_{1}} + \left(\frac{1}{2c_{0}} - \frac{1}{2c_{1}} \right) + 2 \left(\frac{1}{4c_{1}} - \frac{1}{2c_{0}} \right) \delta = 0 \quad Corr. \ 3.1$$

$$\Rightarrow \delta \left(\frac{1}{2c_{0}} - \frac{1}{c_{1}} + \frac{1}{2c_{1}} - \frac{1}{c_{0}} \right) = -\frac{1}{2c_{1}} - \frac{1}{2c_{0}} + \frac{1}{2c_{1}}$$

$$\Rightarrow \delta \left(c_{1} - 2c_{0} + c_{0} - 2c_{1} \right) = -c_{1}$$

$$\Rightarrow \delta = \frac{c_{1}}{c_{1} + c_{0}}.$$

The second partial derivative is $\frac{\partial^2 U_G}{\partial \delta^2} = -\frac{1}{2c_1} - \frac{1}{2c_0}$ which is negative for any $c_0, c_1 > 0$, meaning $\delta^{VerticalMonopoly} = \frac{c_1}{c_1 + c_0}$ maximizes the sum of utilities.

8.8 Egalitarian bargaining solution

Proof of Proposition 3.6. The Kalai (egalitarian) solution $\delta^{Egal.}$ is the solution that maximizes the minimum utility among players.

First, observe that if there exists a point in the Pareto solution set where the two utilities are equal, this point must be the egalitarian solution. Pareto means that an increase in any player's utility must correspond to a decrease in another player's utility. If a solution within the Pareto set equalizes utilities, then any other solution must inevitably trade off one player's utility for the other's, meaning any alternative solution would yield a lower utility for at least one player.

So, setting the utilities equal we get:

$$U_G(\delta) = U_D(\delta)$$

$$\frac{\delta^2}{4c_0} + \frac{\delta}{2c_1} - \frac{\delta^2}{2c_1} = \frac{1}{4c_1} + \frac{\delta}{2c_0} - \frac{\delta}{2c_1} + \frac{\delta^2}{4c_1} - \frac{\delta^2}{2c_0} \quad Corr. \quad 3.1$$

$$\delta^2(3c_1 - 3c_0) + \delta(4c_0 - 2c_1) - c_0 = 0$$

Plugging into the quadratic formula, we get two candidate solutions:

$$\delta^{Egal.} \stackrel{?}{=} \left\{ \frac{\sqrt{c_0^2 - c_0 c_1 + c_1^2} - c_1 + 2c_0}{3(c_0 - c_1)}, \frac{-\sqrt{c_0^2 - c_0 c_1 + c_1^2} - c_1 + 2c_0}{3(c_0 - c_1)} \right\}$$

Notice that the first of these solutions, for $c_0, c_1 > 0$, is not in the feasible set $0 \le \delta \le 1$. Thus, the Egalitarian solution is given by:

$$\delta^{Egal.} = \frac{-\sqrt{c_0^2 - c_0 c_1 + c_1^2} - c_1 + 2c_0}{3(c_0 - c_1)}.$$

9 Section 4 Materials

9.1 Existence theorem

Proof of Theorem 4.1. We prove this theorem via a sequence of Lemmas.

Lemma 9.1. If $r'(0) > \lambda_0 \phi_0'(0)$ for a constant $\lambda_0 \geq 2$, then there exists a set $A^* \subseteq (0,1)$ such that $\frac{1}{2} \in A^*$ and for all $\delta \in A^*$, $\alpha_0^*(\delta) > 0$, $U_G(\delta) > 0$.

Let's presume $\delta^* = \frac{1}{2}$. If we can show that this solution yields $\alpha_0^* > 0$, then this means the generalist's utility is greater than 0 for investing some non-zero effort spend. α_0^* is the value of α_0 that maximizes U_G , so if U_G has positive slope at $\alpha_0 = 0$, then $\alpha_0^* > 0$.

Notice that this positive-utility outcome is met as long as $\frac{\partial U_G}{\partial \alpha_0}|_{\alpha_0=0} > 0$. This condition would necessarily mean that there exists some positive $\alpha_0 > 0$ which maximizes U_G . So, formally:

$$\begin{aligned} r'(0) &> 2\phi_0'(0) \\ \frac{1}{2}r'(0) - \phi_0'(0) &> 0 \\ \frac{\partial}{\partial \alpha_0} \left(\delta r - \phi_0\right) \bigg|_{\alpha_0 = 0} &> 0 \\ \frac{\partial U_G}{\partial \alpha_0} \bigg|_{\alpha_0 = 0} &> 0. \end{aligned}$$

Notice that this inequality is met as long as $r'(0) > \lambda_0 \phi'_0(0)$ where $\lambda_0 > 2$. Thus, there is a non-empty set A^* of solutions with $\frac{1}{2} \in A^*$ that yield positive α_0 and positive U_G .

Lemma 9.2. If $r'(\alpha_0) > \lambda_1 \phi_1'(\alpha_0)$ for a constant $\lambda_1 \geq 2$, then there exists a set $B^* \subseteq (0,1)$ such that $\frac{1}{2} \in B^*$ and for all $\delta \in B^*$, $\alpha_1^*(\delta) > \alpha_0^*$, $U_D(\delta) > 0$.

Let's presume $\delta^* = \frac{1}{2}$. If we can show that this solution yields $\alpha_1^* > \alpha_0^*$, then this means the domain specialist's utility is greater than 0 for investing some non-zero effort spend. α_1^* is the value of α_1 that maximizes U_D , so if U_D has positive slope at $\alpha_1 = \alpha_0^*$, then $\alpha_1^* > \alpha_0^*$.

maximizes U_D , so if U_D has positive slope at $\alpha_1 = \alpha_0^*$, then $\alpha_1^* > \alpha_0^*$. Notice that this positive-utility outcome is met as long as $\frac{\partial U_D}{\partial \alpha_1}\Big|_{\alpha_1 = \alpha_0^*} > 0$ – this condition would necessarily mean that there exists some $\alpha_1^* > \alpha_0^*$ which maximizes U_D . So, formally:

$$r'(\alpha_0) > 2\phi_1'(\alpha_0)$$

$$\frac{1}{2}r'(\alpha_0) - \phi_1'(\alpha_0) > 0$$

$$\frac{\partial}{\partial \alpha_1} \left((1 - \delta)r - \phi_1 \right) \Big|_{\alpha_1 = \alpha_0} > 0$$

$$\frac{\partial U_D}{\partial \alpha_1} \Big|_{\alpha_1 = \alpha_0} > 0.$$

Notice that this inequality is met as long as $r'(\alpha_0) > \lambda_1 \phi_1'(\alpha_0)$ where $\lambda_1 \geq 2$. Thus, there is a non-empty set B^* of solutions with $\frac{1}{2} \in B^*$ that yield $\alpha_1^* > \alpha_0^*$ and positive U_D .

Corollary 9.1. $A^* \cap B^*$ is a non-empty set where any solution $\delta^* \in A^* \cap B^*$ satisfies the three properties: $0 < \delta_i < 1$; $0 < \alpha_0^* < \alpha_1^*$; and $U_D, U_G > 0$.

This Corollary follows from the findings that have already been shown in the former Lemmas.

First, $A^* \cap B^*$ is non-empty because $\frac{1}{2} \in A^*$ (as shown in Lemma 9.1) and $\frac{1}{2} \in B^*$ (as shown in Lemma 9.2) so it follows that $\frac{1}{2} \in A^* \cap B^*$.

The three properties are each met for the following reasons:

- 1. Property 1 $(0 < \delta^* < 1)$ is met because $\frac{1}{2} \in A^* \cap B^*$ and $0 < \frac{1}{2} < 1$ so the existence finding is satisfied for a non-extreme value of δ . This means, for $\delta^* \in A^* \cap B^*$, players share revenue.
- 2. Property 2 ($0 < \alpha_0^* < \alpha_1^*$) is met because any solution in A^* yields $\alpha_0^* > 0$ (as shown in Lemma 9.1) and any solution in B^* yields $\alpha_1^* > \alpha_0^*$ (as shown in Lemma 9.2). This means, for $\delta^* \in A^* \cap B^*$, players do not free-ride (they both act to improve the technology).
- 3. Property 3 $(U_G, U_D > 0)$ is met because any solution in A^* yields $U_G > 0$ (as shown in Lemma 9.1) and any solution in B^* yields $U_D > 0$ (as shown in Lemma 9.2). This means any solution $\delta^* \in A^* \cap B^*$ yields positive utility for both players, which Pareto-dominates the disagreement scenario in which both players have zero utility.

10 Section 5 Materials

10.1 Proof of Pareto Set Characterization in the Multi-specialist Game

Proof of Theorem 5.1. Say there are $n \ge 2$ specialists. For each player in the game, at a given value of δ , the utility curve can exhibit one of three regimes: 1) negative utility 2) increasing utility 3) decreasing utility. Thus, at a particular value of δ , the utilities as a function of delta can exhibit 3^{n+1} different combinations of regimes. For example, if there are n = 2 domain specialists, a given value of δ might exhibit one of $3^3 = 27$ regimes, including $\{D_1 \text{ negative}, D_2 \text{ decreasing}, G \text{ increasing}\}$ as one example.

We systematically analyze all possible regimes:

1. All regimes where the generalist's utility are negative $U_G < 0$ would lead the generalist to opt for the no-deal solution – this captures a total of 3^n combinations. In the n=2 example, these scenarios are described by $\{\cdot,\cdot,G \text{ negative}\}$, so a total of $3^2=9$ out of 27 possible regimes are accounted for.

- 2. All regimes where the U_G is positive-increasing and no specialist is positive-decreasing are Pareto-dominated. This is because, if no specialist is decreasing, then for some small increment of δ , all players are better off. This captures a total of 2^n combinations. In the n=2 example, these scenarios account for a total of $2^2=4$ combinations. The four combinations would be: $\{D_1 \text{ increasing}, D_2 \text{ increasing}, G \text{ increasing}\}, \{D_1 \text{ negative}, D_2 \text{ increasing}\}, \{D_1 \text{ increasing}\}, \{D_1 \text{ negative}, G \text{ increasing}\}.$
- 3. All regimes where the U_G is positive-decreasing and no specialist is positive-increasing are Pareto-dominated. This is because, if no specialist is decreasing, then for some small decrease of δ , all players are better off. This captures a total of 2^n combinations. In the n=2 example, these scenarios account for a total of $2^2=4$ combinations.
- 4. All regimes where U_G is positive-increasing and at least one specialist is positive-decreasing are Paretooptimal. This is because, for unimodal curves, it is impossible to improve one players' utility without a
 corresponding decrease in the other's. This secnario is exactly the set outside scenario (2) above. Thus,
 this accounts for $3^n 2^n$ cases. In the case where n = 2, this would be 5 cases.
- 5. All regimes where U_G is positive-decreasing and at least one specialist is positive-increasing are Paretooptimal. This is because, for unimodal curves, it is impossible to improve one players' utility without a
 corresponding decrease in the other's. This secnario is exactly the set outside scenario (3) above. Thus,
 this accounts for $3^n 2^n$ cases. In the case where n = 2, this would be 5 cases.

In total, we've defined 5 non-overlapping cases, collectively accounting for $3^n + 2^n + 2^n$ cases where the value of δ is not Pareto-optimal, and $(3^n - 2^n) + (3^n - 2^n)$ cases where the value of δ is Pareto-optimal. Overall these cases sum to $3^n + 2^n + 2^n + (3^n - 2^n) + (3^n - 2^n) = 3 * 3^n = 3^{n+1}$, so we've accounted for every possible regime.

10.2 Derivation of the Subgame Perfect Equilibrium for a Given δ

We use backward induction to determine the multi-specialist fine-tuning game's subgame perfect equilibrium, as we did in Section 3.1.

Theorem 10.1. For a fixed δ , the sub-game perfect equilibrium of the fine-tuning game with polynomial costs yields the following best-response strategies: $\alpha_0^* = {}^{k_0-1}\sqrt{\frac{n\delta}{c_0k_0}}$; $\alpha_i^* = {}^{k_0-1}\sqrt{\frac{n\delta}{c_0k_0}} + {}^{k_i-1}\sqrt{\frac{1-\delta}{c_ik_i}}$.

Proof of Theorem 10.1. We solve the game using backward induction as follows: First, starting with the last stage (3), we solve for α_i^* given α and δ :

$$\alpha_{i}^{*} = \operatorname{argmax}_{\alpha_{i}} U_{D_{i}}(\alpha_{i}, \alpha_{0}, \delta)$$

$$\Rightarrow \frac{\partial U_{D_{i}}}{\partial \alpha_{i}} \Big|_{\alpha_{i} = \alpha_{i}^{*}} = 0$$

$$\Rightarrow (1 - \delta) - \frac{\partial \phi_{i}}{\partial \alpha_{i}} = 0$$

$$\Rightarrow (1 - \delta) - c_{i}k_{i}(\alpha_{i} - \alpha_{0})^{k_{i} - 1}$$

$$\Rightarrow \alpha_{i}^{*} = \alpha_{0} + {}^{k_{i}} \sqrt{\frac{1 - \delta}{c_{i}k_{i}}}.$$
(12)

Note that $\frac{\partial^2 U_{D_i}}{\partial \alpha_i^2} = -c_i k_i (k_i - 1) (\alpha_i - \alpha_0)^{k_i - 2}$ is negative for all $c_i > 0$ and $k_i > 1$ meaning that α_i^* derived above yields a global maximum of U_{D_i} .

Second, knowing D_i 's choice of α_i^* above, we solve for α_0^* as follows:

$$\alpha_0^* = \operatorname{argmax}_{\alpha_0} U_G(\alpha_0, \delta)$$

$$\Rightarrow \frac{\partial U_G}{\partial \alpha_0} \Big|_{\alpha_0 = \alpha_0^*} = 0$$

$$\Rightarrow \frac{\partial}{\partial \alpha_0} \left(\sum_{i=1}^n \delta \alpha_i - c_0(\alpha_0)^{k_0} \right) \Big|_{\alpha_0 = \alpha_0^*} = 0$$

$$\Rightarrow n\delta - c_0 k_0 \alpha_0^{k_0 - 1} = 0$$

$$\Rightarrow \alpha_0^* = {}^{k_0 - 1} \sqrt{\frac{n\delta}{c_0 k_0}}.$$

The second derivative $\frac{\partial^2 U_G}{\partial \alpha_i^2} = -c_0 k_0 (k_0 - 1) \alpha_0^{k_0 - 2}$ is negative for all $c_i > 0$ and $k_i > 1$ meaning that α_0^* derived above yields a global maximum of U_G .

Finally, plugging in $\alpha_0^* = \sqrt[k_0-1]{\frac{n\delta}{c_0k_0}}$ into Equation 12, we obtain the following expression for α_i^* as a function of $\delta, c_0, c_1, k_0, k_1$ only:

$$\alpha_i^* = \sqrt[k_0 - 1]{\frac{n\delta}{c_0 k_0}} + \sqrt[k_i - 1]{\frac{1 - \delta}{c_i k_i}}.$$

This finishes the proof.

As a corollary to Theorem 10.1, we can specify the utilities of each player as a function of δ alone:

Corollary 10.1. For a fixed bargaining parameter δ , the players' utilities are as follows:

$$U_G(\delta) = n \left(\frac{n}{c_0 k_0}\right)^{\frac{1}{k_0 - 1}} \left(1 - \frac{1}{k_0}\right) \delta^{\frac{k_0}{k_0 - 1}} + \delta \sum_i \left(c_i k_i\right)^{\frac{-1}{k_i - 1}} \left(1 - \delta\right)^{\frac{1}{k_i - 1}},\tag{13}$$

$$U_{D_i}(\delta) = \left(\frac{n}{c_0 k_0}\right)^{\frac{1}{k_0 - 1}} \delta^{\frac{1}{k_0 - 1}} \left(1 - \delta\right) + \left(c_i k_i\right)^{\frac{-1}{k_i - 1}} \left(1 - \frac{1}{k_i}\right) \left(1 - \delta\right)^{\frac{k_i}{k_i - 1}}.$$
 (14)

Proof of Corollary 10.1. Plugging in the formulas from Theorem 10.1 into equation 9, we obtain:

$$\begin{split} U_G &= \sum_{i} \delta r_i(\alpha_i) - \phi_0(\alpha_0) \\ &= \delta \sum_{i} \left(\sqrt[k_0 - 1]{\frac{n\delta}{c_0 k_0}} + \sqrt[k_1 - 1]{\frac{1-\delta}{c_i k_i}} \right) - c_0 \left(\sqrt[k_0 - 1]{\frac{n\delta}{c_0 k_0}} \right)^{k_0} \\ &= \delta \sum_{i} \left(\frac{n\delta}{c_0 k_0} \right)^{\frac{1}{k_0 - 1}} + \delta \sum_{i} \left(\frac{1-\delta}{c_i k_i} \right)^{\frac{1}{k_i - 1}} - c_0 \left(\frac{n\delta}{c_0 k_0} \right)^{\frac{k_0}{k_0 - 1}} \\ &= n \left(\frac{n}{c_0 k_0} \right)^{\frac{1}{k_0 - 1}} \delta^{\frac{k_0}{k_0 - 1}} - c_0 \left(\frac{n}{c_0 k_0} \right)^{\frac{k_0}{k_0 - 1}} \delta^{\frac{k_0}{k_0 - 1}} + \delta \sum_{i} \left(\frac{1}{c_i k_i} \right)^{\frac{1}{k_i - 1}} (1 - \delta)^{\frac{1}{k_i - 1}} \\ &= n \left(\frac{n}{c_0 k_0} \right)^{\frac{1}{k_0 - 1}} \left(1 - \frac{1}{k_0} \right) \delta^{\frac{k_0}{k_0 - 1}} + \delta \sum_{i} \left(c_i k_i \right)^{\frac{-1}{k_i - 1}} (1 - \delta)^{\frac{1}{k_i - 1}}. \end{split}$$

Plugging in the formulas from Theorem 10.1 into equation 10, we obtain:

$$\begin{split} U_{D_i} &= & (1-\delta)r_i(\alpha_i) - \phi_i(\alpha_i;\alpha_0) \\ &= & (1-\delta)\left(\sqrt[k_0-1]{\frac{n\delta}{c_0k_0}} + \sqrt[k_i-1]{\frac{1-\delta}{c_ik_i}} \right) - c_i\left(\sqrt[k_0-1]{\frac{n\delta}{c_0k_0}} + \sqrt[k_i-1]{\frac{1-\delta}{c_ik_i}} - \sqrt[k_0-1]{\frac{n\delta}{c_0k_0}} \right)^{k_i} \\ &= & (1-\delta)\left(\frac{n\delta}{c_0k_0} \right)^{\frac{1}{k_0-1}} + (1-\delta)\left(\frac{1-\delta}{c_ik_i} \right)^{\frac{1}{k_i-1}} - c_i\left(\frac{1-\delta}{c_ik_i} \right)^{\frac{k_i}{k_i-1}} \\ &= & \left(\frac{n}{c_0k_0} \right)^{\frac{1}{k_0-1}} \delta^{\frac{1}{k_0-1}} (1-\delta) + (c_ik_i)^{\frac{-1}{k_i-1}} \left(1 - \frac{1}{k_i} \right) (1-\delta)^{\frac{k_i}{k_i-1}}. \end{split}$$

10.3Deriving the Set of Pareto-Optimal Solutions

In order to determine the set of Pareto-optimal bargaining agreements, first notice that when $k_0 = 2$ and $k_i = 2 \,\forall i$ (i.e., the case of quadratic costs), the utilities are strictly unimodal.

Observation 10.1. In the multi-specialist fine-tuning game with quadratic costs, U_G and U_{D_i} are strictly unimodal functions of δ , for all domain-specialists i.

Proof of Observation 10.1. Notice that the utility functions take the same form as the one-specialist case, whose analogous observation is proven in Appendix 8.3. There, we proved that a utility of the form $A\delta^{\frac{k_0}{k_0-1}} + B\delta(1-\delta)^{\frac{1}{k_1-1}}$ is unimodal for $A, B > 0, k_0, k_1 \ge 2, \delta \in [0,1].$

For simplicity, in the multi-specialist case, we prove unimodality for the case of quadratic costs (this is all we need to arrive at the bargaining solutions reported in the paper). We show that the same proof holds for both the generalist and specialists' utilities:

- For the specialist, the utility function given in Equation 14 is already of the form proven unimodal in Proposition 3.1.
- For the generalist, using Equation 13, $U_G(\delta) = n\left(\frac{n}{c_0k_0}\right)\left(1-\frac{1}{2}\right)\delta^2 + \delta\sum_i\left(c_ik_i\right)^{-1}\left(1-\delta\right) = A'\delta^2 + C_{ij}(\delta)$ $B'\delta(1-\delta)$. This matches the form already proven unimodal in Proposition 3.1.

This suggests that this multi-specialist fine-tuning game has a set of Pareto-optimal solutions that are characterized by Theorem 5.1.

10.3.1Powerful-Player Solutions

Proposition 10.1 (Powerful-G Solution). The Powerful-G solution to the multi-specialist fine-tuning game with quadratic costs is as follows:

$$\delta^{Powerful\ G} = \begin{cases} \frac{\sum_{i} \frac{1}{c_{i}}}{2\sum_{i} \frac{1}{c_{i}} - \frac{n^{2}}{c_{0}}} & for \ \frac{1}{n} \sum_{i} \frac{c_{0}}{c_{i}} > n, \\ 1 & else. \end{cases}$$

Proof of Proposition 10.1. The powerful-G solution is the solution that maximizes G's utility:

$$\begin{split} \delta^{\text{Powerful }G} &= \arg\max_{\delta} U_G(\delta) \\ \Rightarrow & \frac{\partial U_G}{\partial \delta} = 0 \\ \Rightarrow & \frac{\partial}{\partial \delta} \left[n \left(\frac{n}{2c_0} \right) \left(1 - \frac{1}{2} \right) \delta^2 + \delta \sum_i \left(2c_i \right)^{-1} \left(1 - \delta \right) \right] = 0 \\ \Rightarrow & \frac{n^2}{4c_0} (2\delta) + \sum_i \frac{1}{2c_i} \left(1 - 2\delta \right) = 0 \\ \Rightarrow & \frac{n^2}{2c_0} (\delta) + \sum_i \frac{1}{2c_i} - \sum_i \frac{1}{c_i} (\delta) = 0 \\ \Rightarrow & \delta \left(\frac{n^2}{2c_0} - \sum_i \frac{1}{c_i} \right) = -\sum_i \frac{1}{2c_i} \\ \Rightarrow & \delta^* = \frac{\sum_i \frac{1}{2c_i}}{\sum_i \frac{1}{c_i} - \frac{n^2}{2c_0}}. \end{split}$$

The second partial derivative $\frac{\partial^2 U_G}{\partial \delta^2} = \frac{n^2}{2c_0} - \sum_i \frac{1}{c_i}$ suggesting the critical point identified represents a maximum as long as $\frac{1}{n}\sum_i \frac{c_0}{c_i} > \frac{n}{2}$. We therefore have the following three cases:

• For $\frac{1}{n}\sum_{i}\frac{c_0}{c_i} \geq n$, the expression maximizes U_G and $\delta^* \in [0,1]$ so $\delta^{\text{Powerful } G} = \delta^*$ for this case.

- For $n > \frac{1}{n} \sum_{i} \frac{c_0}{c_i} > \frac{n}{2}$, the expression maximizes U_G but $\delta^* > 1$. So, notice the expression is strictly increasing in $\delta \in [0, 1]$, so $\delta^{\text{Powerful } G} = 1$ for this case.
- For $\frac{1}{n}\sum_{i}\frac{c_0}{c_i} \leq \frac{n}{2}$, the expression minimizes U_G and the expression is strictly decreasing over $\delta \in [0,1]$ so $\delta^{\text{Powerful }G} = 1$ for this case.

Thus:

$$\delta^{\text{Powerful }G} = \begin{cases} \frac{\sum_{i} \frac{1}{c_{i}}}{2\sum_{i} \frac{1}{c_{i}} - \frac{n^{2}}{c_{0}}} & \textit{for } \frac{1}{n} \sum_{i} \frac{c_{0}}{c_{i}} > n, \\ 1 & \textit{else}. \end{cases}$$

Proposition 10.2 (Powerful- D_i Solution). The Powerful- D_i solution to the multi-specialist fine-tuning game with quadratic costs is as follows:

$$\delta^{Powerful\ D_i} = \begin{cases} \frac{2c_0 - 2nc_i}{c_0 - 2nc_i} & for \ \frac{c_0}{c_i} < n, \\ 0 & else. \end{cases}$$

Proof of Proposition 10.2. The powerful- D_i solution maximizes D_i 's utility:

$$\delta^{\text{Powerful }D_{i}} = \arg\max_{\delta} U_{D_{i}}(\delta)$$

$$\Rightarrow \frac{\partial U_{D_{i}}}{\partial \delta} = 0$$

$$\Rightarrow \frac{\partial}{\partial \delta} \left[\left(\frac{n}{2c_{0}} \right) \delta(1 - \delta) + \frac{1}{2c_{i}} \left(1 - \frac{1}{2} \right) (1 - \delta)^{2} \right] = 0$$

$$\Rightarrow \left(\frac{n}{2c_{0}} \right) (1 - 2\delta) - 2\frac{1}{2c_{i}} \left(\frac{1}{2} \right) (1 - \delta) = 0$$

$$\Rightarrow \frac{n}{2c_{0}} - \frac{n}{c_{0}} \delta - \frac{1}{2c_{i}} + \frac{1}{2c_{i}} \delta = 0$$

$$\Rightarrow \left(\frac{1}{2c_{i}} - \frac{n}{c_{0}} \right) \delta = \frac{1}{2c_{i}} - \frac{n}{2c_{0}}$$

$$\Rightarrow \delta^{*} = \frac{\frac{1}{2c_{i}} - \frac{n}{2c_{0}}}{\frac{1}{2c_{i}} - \frac{n}{c_{0}}}$$

$$\Rightarrow \delta^{*} = \frac{2c_{0} - 2nc_{i}}{c_{0} - 2nc_{i}}.$$

The second partial derivative $\frac{\partial^2 U_{D_i}}{\partial \delta^2} = \frac{1}{2c_i} - \frac{n}{c_0}$, meaning the critical point identified represents a maximum so long as $\frac{c_0}{c_i} < 2n$.

We therefore have the following three cases:

- For $c_i \geq \frac{c_0}{n}$, the expression maximizes U_{D_i} and $\delta^* \in [0,1]$, so $\delta^{\text{Powerful } D_i} = \delta^*$ for this case.
- For $\frac{c_0}{n} > c_i > \frac{c_0}{2n}$, the expression maximizes U_{D_i} but $\delta^* < 0$. So, notice the expression is strictly decreasing in $\delta \in [0, 1]$, so $\delta^{\text{Powerful } D_i} = 0$ for this case.
- For $\frac{c_0}{2n} \ge c_i$, the expression minimizes U_{D_i} and the expression is strictly decreasing in $\delta \in [0,1]$ so $\delta^{\text{Powerful } D_i} = 0$ for this case.

Thus:

$$\delta^{\text{Powerful } D_i} = \begin{cases} \frac{2c_0 - 2nc_i}{c_0 - 2nc_i} & \text{ for } \frac{c_0}{c_i} < n, \\ 0 & \text{ else.} \end{cases}$$

10.3.2 Solution that maximizes the technology's performance

Definition 10.1 (Maximum-performance solution). For the multi-specialist fine-tuning game, the maximum-performance bargaining solution is the feasible revenue-sharing agreement $\delta^{max-\sum_i \alpha_i^*} \in [0,1]$ that maximizes the sum of the technology's domain-specific performances: $\delta^{max-\sum_i \alpha_i^*} = \arg \max_{\delta \in [0,1]} \sum_i \alpha_i^*$.

Proposition 10.3 (Maximum- $\sum_i \alpha_i^*$ Solution). A bargaining solution that maximizes the technology's domain-specific performances is given by:

$$\delta^{\text{Max-}\sum_{i}\alpha_{i}^{*}} = \begin{cases} 0 & \text{for } \frac{1}{n}\sum_{i}\frac{c_{0}}{c_{i}} > n, \\ 1 & \text{else.} \end{cases}$$

Proof of Proposition 10.3. The maximum-performance solution to the multi-specialist fine-tuning game maximizes the sum of performances:

$$\begin{split} \delta^{\text{Max-}\sum_i \alpha_i^*} &= \arg\max_{\delta} \sum_i \alpha_i^* \\ \Rightarrow & \frac{\partial}{\partial \delta} \sum_i \alpha_i \\ &= & \frac{\partial}{\partial \delta} \sum_i \left[\frac{n}{2c_0} \delta + (1-\delta) \frac{1}{2} \sum_j \frac{1}{c_j} \right] \\ &= & \frac{n^2}{2c_0} + (-1) \frac{1}{2} \sum_i \frac{1}{c_i} \\ &= & \frac{n^2}{2c_0} - \sum_i \frac{1}{2c_i}. \end{split}$$

Notice this quantity is constant over the domain $\delta \in [0, 1]$ and is either non-positive or non-negative. We therefore have the following cases:

- If $\sum_{i} \frac{1}{c_i} > \frac{n^2}{c_0}$, then the sum of downstream performances $\sum_{i} \alpha_i$ is decreasing in δ so $\delta^{\text{Max-}\sum_{i} \alpha_i^*} = 0$.
- Otherwise, the sum of downstream performances $\sum_i \alpha_i$ is increasing in δ so $\delta^{\text{Max-}\sum_i \alpha_i^*} = 1$.

10.3.3 Vertical Monopoly Solution

Proposition 10.4 (Vertical Monopoly Solution). The Vertical Monopoly Bargaining Solution to the multi-specialist fine-tuning game with quadratic costs is as follows:

$$\delta^{\textit{Vertical Monopoly}} = \frac{n^2}{n^2 + \sum_i \frac{c_0}{c_i}}.$$

Proof of Proposition 10.4. The vertical monopoly bargaining solution maximizes the sum of utilities:

$$\delta^{VM} = \arg\max_{\delta} \left(U_G + \sum_i U_{D_i}\right)$$

$$\Rightarrow \frac{\partial}{\partial \delta} \left(U_G + \sum_i U_{D_i}\right) = 0$$

$$\Rightarrow \frac{\partial}{\partial \delta} \left[\frac{n^2}{4c_0} \delta^2 + \delta(1-\delta) \sum_i \frac{1}{2c_i} + \sum_i \left(\frac{1}{4c_i} + \left(\frac{n}{2c_0} - \frac{1}{2c_i}\right) \delta + \left(\frac{1}{4c_i} - \frac{n}{2c_0}\right) \delta^2\right)\right] = 0$$

$$\Rightarrow \frac{n^2}{2c_0} \delta + (1-2\delta) \left(\sum_i \frac{1}{2c_i}\right) + \sum_i \left(\frac{n}{2c_0} - \frac{1}{2c_i} + 2\left(\frac{1}{4c_i} - \frac{n}{2c_0}\right) \delta\right) = 0$$

$$\Rightarrow \frac{n^2}{2c_0} \delta + (1-2\delta) \left(\sum_i \frac{1}{2c_i}\right) + \frac{n^2}{2c_0} - \frac{2n^2\delta}{2c_0} + \sum_i \left(\frac{1}{2c_i} \delta - \frac{1}{2c_i}\right) = 0$$

$$\Rightarrow \left(\frac{1}{2} - \delta\right) \left(\sum_i \frac{1}{c_i}\right) + (1-\delta) \frac{n^2}{2c_0} + \frac{1}{2}(\delta - 1) \left(\sum_i \frac{1}{c_i}\right) = 0$$

$$\Rightarrow \frac{1}{2} \delta \left(\sum_i \frac{1}{c_i}\right) + \delta \frac{n^2}{2c_0} = 0$$

$$\Rightarrow \frac{1}{2} \delta \left(\sum_i \frac{1}{c_i}\right) + \delta \frac{n^2}{2c_0} = \frac{n^2}{2c_0}$$

$$\Rightarrow \delta^* = \frac{n^2}{n^2 + \sum_i \frac{c_0}{c_i}}.$$

The second partial derivative $\frac{\partial^2}{\partial \delta^2} \left(U_G + \sum_i U_{D_i} \right) = -\frac{1}{2} \left(\sum_i \frac{1}{c_i} \right) - \frac{n^2}{2c_0}$ which is negative $\forall c_0, c_i > 0$. Thus, $\delta^{VM} = \delta^*$ maximizes the sum of utilities.

10.3.4 Egalitarian Bargaining Solution

Proposition 10.5 (Egalitarian Solution). The Egalitarian Bargaining Solution to the multi-specialist fine-tuning game with quadratic costs is given by:

$$\delta^{Egal.} = \frac{\sqrt{c_0^2 K^2 - 2c_0 nK + \frac{c_0}{c_{max}} n^2 + n^2} + n - \frac{c_0}{c_{max}} n - c_0 K}{n^2 + 2n - \frac{c_0}{c_0} - 2c_0 K}.$$

Where $K = \sum_{i} \frac{1}{c_i}$ and $c_{max} = \max_{i} c_i$.

Proof of Proposition 10.5. The egalitarian solution is given by:

$$\delta^{\text{Egal.}} = \arg \max_{\delta} \left[\min \left(U_G, \min_i \left(U_{D_i} \right) \right) \right].$$

Finding a closed-form solution for the egalitarian bargaining agreement proves more challenging in the multi-specialist case because many players in the fine-tuning game, could, potentially, be worst-off. Instead of simply equalizing utilities, the multi-specialist case might have an egalitarian solution where only the two worst-off players have equal utility.

The proof for quadratic costs starts off by observing that the specialist with the lowest utility must be the specialist j with the highest cost coefficient c_j . This is because revenue function $r(\cdot) = I(\cdot)$ is constant across all domains, as is the bargaining parameter δ and coefficient $k_i = 2$. Thus, the highest-cost-coefficient domain specialist is given by $j = \arg \max_i c_i$. This observation leaves us with two possibilities for the lowest-utility player: 1) Player D_j 2) Player G. Thus, the egalitarian solution is the bargaining agreement that equalizes the utilities between the generalist and the costliest player:

$$U_G(\delta^{\text{Egal.}}) = U_{D_i}(\delta^{\text{Egal.}}).$$

Plugging in, we get:

$$\begin{split} &\frac{n^2}{4c_0}\delta^2 + \delta \sum_i \frac{1}{2c_i}(1-\delta) = \frac{1}{4c_{\max}} + \left(\frac{n}{2c_0} - \frac{1}{2c_{\max}}\right)\delta + \left(\frac{1}{4c_{\max}} - \frac{n}{2c_0}\right)\delta^2 \\ \Rightarrow & \left[\frac{n^2}{4c_0} - \sum_i \frac{1}{2c_i} + \frac{n}{2c_0} - \frac{1}{4c_{\max}}\right]\delta^2 + \left[\sum_i \frac{1}{2c_i} + \frac{1}{2c_{\max}} - \frac{n}{2c_0}\right]\delta - \frac{1}{4c_{\max}} = 0. \end{split}$$

Plugging in to the quadratic formula, we get two candidate solutions:

$$\delta^{Egal.} \stackrel{?}{=} \left\{ \frac{\pm \sqrt{c_0^2(\sum_i \frac{1}{c_i})^2 - 2c_0 n(\sum_i \frac{1}{c_i}) + \frac{c_0}{c_{\max}} n^2 + n^2} + n - \frac{c_0}{c_{\max}} n - c_0(\sum_i \frac{1}{c_i})}{n^2 + 2n - \frac{c_0}{c_{\max}} - 2c_0(\sum_i \frac{1}{c_i})} \right\}.$$

As a last step, we select the candidate solution that yields a feasible $\delta \in [0, 1]$:

$$\delta^{Egal.} = \frac{\sqrt{c_0^2(\sum_i \frac{1}{c_i})^2 - 2c_0 n(\sum_i \frac{1}{c_i}) + \frac{c_0}{c_{\max}} n^2 + n^2} + n - \frac{c_0}{c_{\max}} n - c_0(\sum_i \frac{1}{c_i})}{n^2 + 2n - \frac{c_0}{c_{\max}} - 2c_0(\sum_i \frac{1}{c_i})}.$$

10.4 Theorem on the Three Specialist Regimes

Proof of Theorem 5.2. We prove this theorem in a sequence of Lemmas. The proof follows for any given specialist D_i and revenue-sharing parameter δ .

Lemma 10.1. If fixed costs are under control, meaning $r_i(\alpha_0) > \frac{1}{1-\delta}\phi_i(\alpha_0)$, then D_i will not abstain – instead, D_i would always prefer to free-ride.

If $r_i(\alpha_0) > \frac{1}{1-\delta}\phi_i(\alpha_0)$, then $U_{D_i}|_{\alpha_i=\alpha_0} = r_i(\alpha_0) - \frac{1}{1-\delta}\phi_i(\alpha_0)$ is simply the RHS minus the LHS of the inequality. This means U_{D_i} must be positive at $\alpha_i = \alpha_0$. Thus, as long as fixed costs are under control, the specialist prefers free-riding to abstaining.

Lemma 10.2. If fixed costs are not under control, meaning $r_i(\alpha_0) < \frac{1}{1-\delta}\phi_i(\alpha_0)$, then D_i will not free-ride instead, D_i would always prefer to abstain.

If $r_i(\alpha_0) < \frac{1}{1-\delta}\phi_i(\alpha_0)$, then $U_{D_i}\big|_{\alpha_i=\alpha_0} = r_i(\alpha_0) - \frac{1}{1-\delta}\phi_i(\alpha_0)$ is simply the RHS minus the LHS of the inequality. This means U_{D_i} must be negative at $\alpha_i = \alpha_0$. Thus, as long as fixed costs are not under control, the specialist prefers abstaining to free-riding.

Lemma 10.3. If it is marginally profitable to invest in the technology, meaning $r'_i(\alpha_0) > \frac{1}{1-\delta}\phi'_i(\alpha_0)$, then D_i will not free-ride – instead, D_i would always prefer to contribute.

If $r_i'(\alpha_0) > \frac{1}{1-\delta}\phi_i'(\alpha_0)$, then $\frac{\partial U_{D_i}}{\partial \alpha_i}\big|_{\alpha_i=\alpha_0} = r_i'(\alpha_0) - \frac{1}{1-\delta}\phi_i'(\alpha_0)$ is simply the RHS minus the LHS of the inequality. This means U_{D_i} is increasing at $\alpha_i=\alpha_0$. Thus, as long as it is marginally profitable to improve the technology, the specialist prefers contributing to free-riding.

Lemma 10.4. If it is marginally costly to invest in the technology, meaning $r'_i(\alpha_0) < \frac{1}{1-\delta}\phi'_i(\alpha_0)$, then D_i will not contribute – instead, D_i would always prefer to free-ride.

If $r_i'(\alpha_0) < \frac{1}{1-\delta}\phi_i'(\alpha_0)$, then $\frac{\partial U_{D_i}}{\partial \alpha_i}\big|_{\alpha_i=\alpha_0} = r_i'(\alpha_0) - \frac{1}{1-\delta}\phi_i'(\alpha_0)$ is simply the RHS minus the LHS of the inequality. This means U_{D_i} is decreasing at $\alpha_i=\alpha_0$. Thus, as long as it is marginally costly to improve the technology, the specialist prefers free-riding to contributing.

Taken together, we can conclude the following about combinations of conditions:

- Fixed costs under control, marginally profitable investment: A<F, F<C (Lemmas 10.1 and 10.3). Thus the specialist would contribute.
- Fixed costs under control, marginally costly: A<F, C<F (Lemmas 10.1 and 10.4). Thus the specialist would free-ride.
- Fixed costs not under control, marginally profitable: F<A, F<C (Lemmas 10.2 and 10.3). Thus the specialist would either abstain or contribute.
- Fixed costs not under control, marginally costly: F<A, C<F (Lemmas 10.2 and 10.4). Thus the specialist would abstain.

Above, the short-hand notation 'A,' 'F,' and 'C' refer to the strategies of abstaining, free-riding, and contributing, respectively. The optimal strategies follow from the two marginal conditions. This completes the proof.