

Vacuum radiation versus shortcuts to adiabaticity

Ricardo R. Ancheyta ^{1,*}

¹ *Centro de Física Aplicada y Tecnología Avanzada, Universidad Nacional Autónoma de México, Boulevard Juriquilla 3001, Querétaro 76230, Mexico*

(Dated: August 11, 2023)

The nonadiabatic dynamic of the electromagnetic field triggers photons generation from the quantum vacuum. Shortcuts to adiabaticity, instead, are protocols that mimic the field's adiabatic dynamic in a finite time. Here, we show how the counterdiabatic term of the transitionless tracking algorithm cancels out, exactly, the term responsible for the photon production in the dynamical Casimir effect. This result suggests that the energy of producing photons out of the vacuum is related to the energetic cost of the shortcut. Furthermore, if the system operates under a quantum thermodynamic cycle, we confirm the equivalence between the adiabatic and nonadiabatic work outputs. Finally, our study reveals that identifying these unreported observations can only be possible using the so-called effective Hamiltonian approach.

I. INTRODUCTION

From where there is no light (vacuum), photons can be created if the electromagnetic field is subject to nonadiabatic changes. This impressive phenomenon is known as the dynamical Casimir effect (DCE) [1, 2]. It was first predicted using a variable-length cavity [3] and experimentally confirmed within the architecture of superconducting quantum circuits [4, 5]. Besides being fundamental to quantum field theory [6], the DCE may have applications in nanophotonics, nanomechanics, and chemistry since it modifies the static Casimir force [1, 7, 8]. Moreover, the DCE is a finite-time quantum electrodynamics process and is expected to be relevant in finite-time quantum thermodynamics [9–11]. For example, in a quantum refrigerator at low temperatures, the DCE dominates and imposes the ultimate limit for cooling [12]. The reason is a heating process produced by the nonadiabatic excitations of the DCE enforcing the third law [13].

On the other hand, shortcuts to adiabaticity (STA) are protocols that imitate adiabatic systems' dynamics (quantum or classical) at finite times [14–16]. STA are receiving much attention in quantum thermodynamics since they increase the power output of quantum heat engines by shortening their adiabatic (isentropic) strokes [17–21]. STA act as quantum lubricants [22], producing frictionless evolution by counteracting nonadiabatic excitations. Theoretically, we may study the STA and the DCE with the harmonic oscillator. For instance, the work by Muga et al. [23] introduced one of the first STA based on transitionless quantum driving [24]; it changed an oscillator's frequency without transitions in finite time. For the DCE, one of its simplest versions occurs in a single-mode nonstationary cavity modeled by an oscillator with time-dependent frequency.

In this work, using an effective Hamiltonian approach [25], we describe in detail how the nonlocal potential arising from the STA combats the generation of

photons in the DCE. We show that the squeezing terms appearing in the DCE and the STA are the same but with opposite signs, implying that the energy of producing photons from the vacuum relates to the energetic cost of the shortcut. Furthermore, with the effective Hamiltonian approach, we make transparent results of the STA in quantum thermodynamics, for instance, the equivalence between the adiabatic and nonadiabatic work outputs.

To make our article as self-consistent as possible, we structure it as follows. First, Sec. II A reviews the STA based on the transitionless tracking algorithm, and Sec. II B discusses the STA of the quantum harmonic oscillator with time-dependent frequency. Next, Sec. III A describes the effective Hamiltonian approach of the DCE and shows that the squeezing term responsible for generating photons is the same (with the opposite sign) as the counterdiabatic term obtained in Sec. II B. This observation simplifies the corresponding Hamiltonian structure and makes it easier to compute the work output when the system operates under a quantum thermodynamic cycle, Sec. III B. Finally, we show our conclusions in Sec. IV.

II. SHORTCUTS TO ADIABATICITY

A. Transitionless tracking algorithm

We review Berry's formulation of the shortcuts to adiabaticity based on the transitionless tracking algorithm since it is more straightforward and enlightening [14, 24]. For the time-dependent Hamiltonian $\hat{H}_0(t)$ satisfying $\hat{H}_0(t)|n(t)\rangle = E_n(t)|n(t)\rangle$, the adiabatic approximation states that $|\psi_n(t)\rangle = e^{i\gamma_n(t)} e^{i\theta_n(t)} |n(t)\rangle$ is an *approximate* solution of the Schrödinger equation when $\hat{H}_0(t)$ varies slowly, see Appendix A. Here, $\gamma_n(t) = i \int_0^t dt' \langle n(t') | [\partial_{t'} |n(t')\rangle]$ and $\theta_n(t) = - \int_0^t dt' E_n(t') \hbar^{-1}$ are the geometrical and dynamical phases. The aim of the transitionless tracking algorithm is to find a quantum Hamiltonian $\hat{H}(t)$ such that $|\psi_n(t)\rangle$ is an *exact* solution of the corresponding Schrödinger equation $i\hbar \partial_t |\psi_n(t)\rangle = \hat{H}(t) |\psi_n(t)\rangle$. The time-evolution operator $\hat{U}(t)$ also satis-

* ancheyta@fata.unam.mx

fies the Schrödinger equation $i\hbar \partial_t \hat{U}(t) = \hat{H}(t) \hat{U}(t)$, then $\hat{H}(t)$ can be extracted from $\hat{H}(t) = i\hbar \partial_t \hat{U}(t) \hat{U}^\dagger(t)$. The next step is to find an explicit form of the evolution operator. Fortunately, by choosing $\hat{U}(t) = \sum_n |\psi_n(t)\rangle \langle n(0)|$, we obtain $\hat{U}(t) |m(0)\rangle = |\psi_m(t)\rangle$. Therefore, $\hat{U}(t)$ is indeed an appropriate choice, where its time-derivative is

$$\begin{aligned} \frac{\partial \hat{U}(t)}{\partial t} &= \sum_n \left\{ i \dot{\gamma}_n(t) |\psi_n(t)\rangle \langle n(0)| \right. \\ &\quad \left. + i \dot{\theta}_n(t) |\psi_n(t)\rangle \langle n(0)| \right. \\ &\quad \left. + e^{i\gamma_n(t)} e^{i\theta_n(t)} \left[\frac{\partial}{\partial t} |n(t)\rangle \right] \langle n(0)| \right\}. \end{aligned} \quad (1)$$

Multiplying the above equation with $\hat{U}^\dagger(t)$ we get

$$\begin{aligned} \frac{\partial \hat{U}(t)}{\partial t} \hat{U}^\dagger(t) &= \sum_n \left\{ i \dot{\gamma}_n(t) |n(t)\rangle \langle n(t)| \right. \\ &\quad \left. + i \dot{\theta}_n(t) |n(t)\rangle \langle n(t)| + \left[\frac{\partial}{\partial t} |n(t)\rangle \right] \langle n(t)| \right\}. \end{aligned} \quad (2)$$

Therefore, the searched Hamiltonian is

$$\hat{H}(t) = \sum_n \left\{ E_n |n\rangle \langle n| + i\hbar (|\partial_t n\rangle \langle n| - \langle n| \partial_t n) |n\rangle \langle n| \right\}. \quad (3)$$

Note we have adopted the simplified notation $|n\rangle = |n(t)\rangle$, $E_n = E_n(t)$ and $|\partial_t n\rangle = \partial |n(t)\rangle / \partial t$. One can split $\hat{H}(t)$ in two parts, $\hat{H}(t) \equiv \hat{H}_0(t) + \hat{H}_1(t)$, where $\hat{H}_0(t) = \sum_n E_n |n\rangle \langle n|$ and $\hat{H}_1(t) = \sum_n i\hbar (|\partial_t n\rangle \langle n| - \langle n| \partial_t n) |n\rangle \langle n|$ are known as the reference and counterdiabatic Hamiltonian, respectively [24].

B. Time-dependent quantum harmonic oscillator

We choose the reference Hamiltonian $\hat{H}_0(t)$ as the quantum harmonic oscillator Hamiltonian with time-dependent frequency, $\omega(t)$ (with $m = 1$)

$$\hat{H}_0(t) = \frac{\hat{p}^2}{2} + \frac{1}{2} \omega^2(t) \hat{x}^2, \quad (4)$$

where \hat{x} and \hat{p} are, respectively, the position and momentum quantum mechanical operators satisfying the canonical commutation relation $[\hat{x}, \hat{p}] = i\hbar$. The corresponding instantaneous eigenvalues are $E_n(t) = \omega(t)(n + 1/2)$. Note we did the mass of the oscillator to the unit to connect with the quantum Hamiltonian of the non-stationary electromagnetic field in Sec. III A. To get $\hat{H}_1(t)$ one needs to know the instantaneous eigenstates $|n(t)\rangle$ of $\hat{H}_0(t)$. In the coordinate representation, these are

$$\begin{aligned} \langle x | n(t) \rangle &= \frac{1}{\sqrt{2^n n!}} \left[\frac{\omega(t)}{\pi \hbar} \right]^{\frac{1}{4}} H_n \left(x \sqrt{\omega(t)/\hbar} \right) \\ &\quad \times \exp \left(-\frac{\omega(t)}{2\hbar} x^2 \right), \end{aligned} \quad (5)$$

where H_n are the Hermite polynomials. Using the recurrence relation of H_n , Muga et al. [23] proved that

$\langle n | \partial_t n \rangle = 0$, simplifying $\hat{H}_1(t) = i\hbar \sum_n |\partial_t n\rangle \langle n|$. By computing the time-derivative of (5) one can show that the counterdiabatic term for the time-dependent quantum harmonic oscillator is [23]

$$\hat{H}_1(t) = -\frac{\dot{\omega}(t)}{4\omega(t)} (\hat{x}\hat{p} + \hat{p}\hat{x}), \quad (6)$$

where the dot in $\omega(t)$ represents its time-derivative. The above result implies that $|\psi_n(t)\rangle$, see Sec. II A, is an exact solution of the Schrödinger equation when using

$$\hat{H}(t) = \frac{\hat{p}^2}{2} + \frac{1}{2} \omega^2(t) \hat{x}^2 - \frac{\dot{\omega}(t)}{4\omega(t)} (\hat{x}\hat{p} + \hat{p}\hat{x}). \quad (7)$$

Hamiltonian (6) is a non-local operator, representing a non-local potential. Since it has a quadratic structure in terms of \hat{x} and \hat{p} , Hamiltonian (7) can be viewed as a generalized harmonic oscillator. Although the finite time dynamics generated by $\hat{H}(t)$ follows, exactly, the adiabatic solution of the reference Hamiltonian $\hat{H}_0(t)$, we want to emphasize that by only looking at the algebraic structure of Eq. (7), it is not obvious understanding why this happens. In other words, at this point, we do not know in detail how the new counterdiabatic term, proportional to $\hat{x}\hat{p} + \hat{p}\hat{x}$, combats the nonadiabatic evolution generated by the Hamiltonian of the time-dependent harmonic oscillator (4). Clarifying such a question is, precisely, the aim of this work.

First, note that $\hat{H}_1(t)$ in (6) is associated with the squeezing operator of quantum optics [27–29], as demonstrated in [23, 30, 31]. Second, it is well known that the time-dependent quantum harmonic oscillator generates, during its evolution, a squeezing effect in the system's quadratures. Therefore, it is reasonable to think that in (7) the counterdiabatic term $\hat{H}_1(t)$ should somehow compensate for the squeezing evolution generated by $\hat{H}_0(t)$.

Due to the above discussion, we want to write (7) in terms of ladder operators. However, writing the proper creation and annihilation operators of the time-dependent quantum harmonic oscillator, equivalent to a time-dependent spacetime metric, is by no means a trivial task [31–36]. We, for instance, may work with the “instantaneous” ladder operators given by [23] ($\hbar = 1$)

$$\hat{a}_t = \frac{1}{\sqrt{2\omega(t)}} [\omega(t) \hat{x} + i\hat{p}], \quad \hat{a}_t^\dagger = \frac{1}{\sqrt{2\omega(t)}} [\omega(t) \hat{x} - i\hat{p}]. \quad (8)$$

Evidently, for any given protocol that starts at $t = t_0$ and ends at $t = t_f$, the oscillator's frequency $\omega(t)$ has fixed values ω_0 and ω_f for $t \leq t_0$ and $t \geq t_f$, respectively. For those cases, Eq. (8) reduces to the well-known definition of the ladder operators of the harmonic oscillator with constant frequency [26]. Equation (8) implies that $\hat{x} = (\hat{a}_t^\dagger + \hat{a}_t) / \sqrt{2\omega(t)}$ and $\hat{p} = i(\hat{a}_t^\dagger - \hat{a}_t) \sqrt{\omega(t)}/2$, which upon substitution in (7) one gets [23]

$$\hat{H}(t) = \omega(t) \left(\hat{a}_t^\dagger \hat{a}_t + \frac{1}{2} \right) - i \frac{\dot{\omega}(t)}{4\omega(t)} (\hat{a}_t^{\dagger 2} - \hat{a}_t^2). \quad (9)$$

Again, the above equation does not provide us with a clear view of how it generates the adiabatic dynamics of reference Hamiltonian $\hat{H}_0(t)$. Instead, it brings us an important result, the second (squeezing) term of (9) can be written without the temporal subscript t [23]. This can be done due to the fact that for any frequency modulation $\omega(t)$, the combination $i\hat{a}_t^{\dagger 2} - i\hat{a}_t^2$ is always equal to the time-independent term $\hat{x}\hat{p} + \hat{p}\hat{x}$, thus, the former can be safely computed at any given time. In particular, and for convenience of the discussion of Sec. III A, we evaluate it at the beginning of a given time-dependent protocol, i.e., throughout the article we shall always write $\hat{H}_1(t) = -i\dot{\omega}(t)(\hat{a}_0^{\dagger 2} - \hat{a}_0^2)/4\omega(t)$. On the contrary, the first term of (9), $\hat{a}_t^\dagger \hat{a}_t + 1/2$, does not equal any time-independent combination of \hat{x} and \hat{p} . Actually, the reference Hamiltonian (4) written in terms of the instantaneous ladder operators (8) at $t = t_0$ looks like

$$\begin{aligned} \hat{H}_0(t) = \omega_0 \left(\hat{a}_0^\dagger \hat{a}_0 + \frac{1}{2} \right) & \left[\frac{\omega^2(t)}{2\omega_0^2} + \frac{1}{2} \right] \\ & + \omega_0 (\hat{a}_0^{\dagger 2} + \hat{a}_0^2) \left[\frac{\omega^2(t)}{4\omega_0^2} - \frac{1}{4} \right]. \end{aligned} \quad (10)$$

To avoid any confusion, keep in mind that $\omega(t)$ in above comes from the bare Hamiltonian (4), while ω_0 is the frequency at which the instantaneous ladder operators (8) are defined. As expected, if $\omega(t) \rightarrow \omega_0$, $\hat{H}_1(t) = 0$ and $\hat{H}(t) \rightarrow \hat{H}_0(t) = \omega_0(\hat{a}_0^\dagger \hat{a}_0 + 1/2)$. When the frequency modulation is of the form $\omega(t) = \omega_0 + f(t)$, with $f(t)$ an arbitrary (usually periodic) time-dependent function, $\hat{H}_0(t) = \omega_0(\hat{a}_0^\dagger \hat{a}_0 + 1/2) + f^2(t)(\hat{a}_0^{\dagger 2} + \hat{a}_0^2 + 2\hat{a}_0^\dagger \hat{a}_0 + 1)/4\omega_0$ has an algebraic structure that has been used to investigate the evolution of coherent states in a Kerr medium [37]. It is worth mentioning that (10) possesses a time-independent squeezing term $(\hat{a}_0^{\dagger 2} + \hat{a}_0^2)$, which, unfortunately, cannot be eliminated with the one contained in $\hat{H}_1(t)$. However, in the next section, we explicitly show how it is possible to obtain an effective reference Hamiltonian $\hat{H}_0(t)$ containing the squeezing term that cancels out $\hat{H}_1(t)$. This idea naturally comes from the studies of non-stationary cavity fields in which the dynamical Casimir effect occurs.

III. DYNAMICAL CASIMIR EFFECT

A. Effective Hamiltonian approach

The DCE, a name introduced by Schwinger [38], was predicted by Moore in 1970 [3]. However, the original theory did not have a Hamiltonian, so for many years the DCE was studied in the Heisenberg representation. It was not until 1994 that C. K. Law derived an effective Hamiltonian of the DCE, allowing an equivalent Schrödinger description [39]. Essential features of the DCE, as the generation of photons from the vacuum,

are captured by the single-mode version of such effective Hamiltonian, which is the Hamiltonian of the time-dependent harmonic oscillator [1, 2]. In what follows, we show the basic steps to obtain the single-mode effective Hamiltonian of the time-dependent oscillator.

First, we need the Heisenberg representation for the instantaneous ladder operators. We replace in (8) the Heisenberg representation of the position and momentum operators, i.e., $\hat{a}_H \equiv [\omega(t)\hat{x}_H + i\hat{p}_H]/\sqrt{2\omega(t)}$, where the subscript H in \hat{x}_H and \hat{p}_H indicates these are in the Heisenberg picture. Second, we take the time-derivative

$$\begin{aligned} \frac{d\hat{a}_H}{dt} &= \frac{1}{\sqrt{2\omega(t)}} \left[\omega(t) \frac{d\hat{x}_H}{dt} + i \frac{d\hat{p}_H}{dt} \right] \\ &+ \frac{1}{\sqrt{2\omega(t)}} \frac{d\omega(t)}{dt} \hat{x}_H \\ &- \frac{1}{[2\omega(t)]^{3/2}} \frac{d\omega(t)}{dt} [\omega(t)\hat{x}_H + i\hat{p}_H], \\ &= \frac{1}{\sqrt{2\omega(t)}} [\omega(t)\hat{p}_H - i\omega^2(t)\hat{x}_H] \\ &+ \frac{1}{[2\omega(t)]^{3/2}} \frac{d\omega(t)}{dt} [\omega(t)\hat{x}_H - i\hat{p}_H], \\ &= -i\omega(t)\hat{a}_H + \frac{1}{2\omega(t)} \frac{d\omega(t)}{dt} \hat{a}_H^\dagger. \end{aligned} \quad (11)$$

In the same manner,

$$\frac{d\hat{a}_H^\dagger}{dt} = +i\omega(t)\hat{a}_H^\dagger + \frac{1}{2\omega(t)} \frac{d\omega(t)}{dt} \hat{a}_H. \quad (12)$$

To get (11) and (12) we have substituted the corresponding equations of motion for the position and momentum operators $d\hat{x}_H/dt = \hat{p}_H$ and $d\hat{p}_H/dt = -\omega^2(t)\hat{x}_H$. At this point, we stress that Eqs. (11) and (12) are not proper quantum mechanical equations of motion [26]. Instead, we obtained them by deriving with respect to time the definition of the instantaneous ladder operator. However, we did use the proper equations of motion of the position and momentum operators generated by the Hamiltonian of the time-dependent harmonic oscillator, i.e., $\hat{H}_0(t)$ of (4) in the Heisenberg equation $d\hat{O}_H/dt = i[\hat{H}_0(t), \hat{O}_H]$, where \hat{O}_H is an arbitrary operator in the Heisenberg representation. Note that contrary to the constant frequency case, Eqs. (11) and (12) display an extra term due to the explicit time-dependency of the ladder operators (8).

Second, to understand the idea of how the effective Hamiltonian approach works, we illustrate it with the constant frequency case. If $\omega(t) \rightarrow \omega_0$, $d\hat{a}_H/dt = -i\omega_0\hat{a}_H$ and $d\hat{a}_H^\dagger/dt = i\omega_0\hat{a}_H^\dagger$. Assuming we are, for some reason, only provided with the information on these equations rather than with the Hamiltonian itself. The crucial question is, which Hamiltonian in the Schrödinger picture can generate the equations above? If such a Hamiltonian exists, the provided equations may be interpreted as proper quantum mechanical equations of motion. For this simple case, it is easy to see that $\hat{H}_{\text{eff}} = \omega_0\hat{a}_0^\dagger\hat{a}_0$ does the desired task since $[\hat{H}_{\text{eff}}, \hat{a}_0] = -\omega_0\hat{a}_0$.

Now, for an arbitrary frequency drive $\omega(t)$ and noting that $[\hat{a}_0^{\dagger 2}, \hat{a}_0] = -2\hat{a}_0^{\dagger}$, it is not difficult to show that the effective Hamiltonian,

$$\hat{H}_{\text{eff}}(t) = \omega(t)\hat{a}_0^{\dagger}\hat{a}_0 + i\frac{1}{4\omega(t)}\frac{d\omega(t)}{dt}(\hat{a}_0^{\dagger 2} - \hat{a}_0^2), \quad (13)$$

indeed generates Eqs. (11) and (12) [1]. This effective Hamiltonian first derived almost three decades ago in the context of the DCE, describes an electromagnetic cavity with a moving mirror, where its frequency is given by $\omega(t) = \pi/q(t)$ and $q(t)$ is the prescribed mirror's trajectory [39]. The Hamiltonian of [39] displays additional terms accounting for an intermode interaction induced by the non-stationary field's boundary conditions. Nevertheless, subsequent studies [40–45] show that when the non-stationary cavity field supports one-single mode, (13) can be safely considered as the simplest version where the DCE can be manifested. For example, under the resonant conditions $\omega(t) = \omega_0[1 + \varepsilon \sin(2\omega_0 t)]$, (13) predicts an exponential photon growth [42–44]

$$\langle 0|\hat{U}^{\dagger}\hat{a}_0^{\dagger}\hat{a}_0\hat{U}|0\rangle = \sinh^2(\varepsilon\omega_0 t/2), \quad (14)$$

where \hat{U} is the corresponding time-evolution operator of $\hat{H}_{\text{eff}}(t)$, $\hat{a}_0|0\rangle = 0$ defines the vacuum state, and $\varepsilon \ll 1$ is a small amplitude modulation depth. Equation (14) is known as the Casimir or vacuum radiation and is the consequence of the nonadiabatic boundary conditions of the field represented by the second term of (13).

Remarkably, when replacing $\hat{H}_0(t)$ by its physically equivalent effective Hamiltonian $\hat{H}_{\text{eff}}(t)$ in the definition of $\hat{H}(t)$, i.e., $\hat{H}(t) = \hat{H}_{\text{eff}}(t) + \hat{H}_1(t)$, we obtain

$$\hat{H}(t) = \omega(t)\hat{a}_0^{\dagger}\hat{a}_0. \quad (15)$$

This equation shows, crystal clear, that the finite-time dynamics generated by the shortcut to adiabaticity is, indeed, the one given by the quantum harmonic oscillator with an instantaneous frequency $\omega(t)$. Since the counterdiabatic term $\hat{H}_1(t) = -i\dot{\omega}(t)(\hat{a}_0^{\dagger 2} - \hat{a}_0^2)/4\omega(t)$ cancels out the induced nonadiabatic squeezing term in (13), the time evolution operator is $\hat{U} = \exp[-i\int dt'\omega(t')\hat{a}_0^{\dagger}\hat{a}_0]$, implying $\langle 0|\hat{U}^{\dagger}\hat{a}_0^{\dagger}\hat{a}_0\hat{U}|0\rangle = 0$. Therefore, no matter what frequency drive is used, photons generation from the vacuum is impossible when performing a shortcut to adiabaticity. Notice that from a different approach, a similar conclusion was recently obtained by exploiting the conformal symmetry of the system [46].

Using (15) it is easy to write the final energy, $\langle \hat{H}(t_f) \rangle = \omega_f \langle \hat{a}_0^{\dagger}\hat{a}_0 \rangle$, in terms of the initial energy $\langle \hat{H}(t_0) \rangle$ as

$$\langle \hat{H}(t_f) \rangle = (\omega_f/\omega_0)\langle \hat{H}(t_0) \rangle. \quad (16)$$

The above expression is a common result encountered in the adiabatic limit of systems with scale-invariance dynamical symmetry, where $(\omega_0/\omega_f)^{1/2}$ is known as the

adiabatic scaling factor [47, 48]. However, since (15) contains the use of an STA, i.e., $\hat{H}(t) = \hat{H}_{\text{eff}}(t) + \hat{H}_1(t)$, the relation between initial and final energies is valid for any physically unitary finite-time protocol driving $\hat{H}_{\text{eff}}(t)$ through $\omega(t)$ and not just for the adiabatic approximation. Certainly, $\hat{H}(t)$ strictly follows the adiabatic solution of the reference Hamiltonian $\hat{H}_0(t)$ but at a finite time. An immediate consequence of this result is in finite-time quantum thermodynamics [9]. For instance, we may choose $\hat{H}(t)$ as the time-dependent Hamiltonian associated with the working substance of a reciprocating quantum heat engine [49, 50]. When the nonstationary cavity suffers an expansion (compression), the cavity length increases (decreases), and the frequency decreases (increases), ending with a lower (higher) internal energy (16). This behavior resembles the one encounter with a thermodynamic piston. Note that the previously discussed situation is not evident when using the standard expressions of $\hat{H}(t)$ encounter in (7) or (9).

Even though the counterdiabatic term $\hat{H}_1(t)$ typically vanishes at the start (t_0) and end (t_f) of an STA, getting a relationship between initial and final energies using (9) is not straightforward. This is because the initial (final) $\hat{H}(t)$ depends on the instantaneous operators at $t = t_0$ (t_f). In contrast, $\hat{H}(t)$ in (15) is always written in terms of operators at some specific time, we used t_0 but the result holds for other times.

B. Quantum thermodynamic implications

The quantum Otto cycle operates between a hot and a cold reservoir and is a paradigmatic thermodynamic cycle extensively used by the quantum thermodynamics community [51, 52]. The cycle consists of four branches: two adiabatic (isentropic) strokes where the working substance (the nonstationary cavity in our case) suffers a compression and expansion while isolated from the heat reservoirs; two quantum isochoric strokes where the system is put in contact with one reservoir, here the cavity frequency has fixed values and heat transfer occurs (thermalization) but no work is performed. During the compression stroke, the cavity frequency, $\omega(t)$, increases from the initial value ω_1 to ω_2 , while in the expansion, it goes from ω_2 to ω_1 . At the hot (cold) isochoric stroke, the cavity relaxes to a thermal state with temperature T_h (T_c).

For the thermal state $\hat{\rho}_{\text{th}} = \exp(-\hbar\omega_j\hat{a}_0^{\dagger}\hat{a}_0/k_B T)Z^{-1}$, where Z is the partition function, the average number of photons is $\langle \hat{a}_0^{\dagger}\hat{a}_0 \rangle = \text{tr}\{\hat{\rho}_{\text{th}}\hat{a}_0^{\dagger}\hat{a}_0\} = \frac{1}{2} \coth(\hbar\omega_j/2k_B T) - 1/2$. With this and with the help of (16), we compute the average energies $\langle \hat{H}(t) \rangle$ of the nonstationary cavity at each of the four strokes of the Otto cycle. Since we use an STA, these values coincide with the ones obtained during the slow adiabatic process [49]. The mean work $\langle W \rangle$, calculated as $\text{tr}\{\hat{\rho}\Delta\hat{H}\}$ with $\Delta\hat{H}$ the change in the Hamiltonian [19], during the compression and expansion strokes

is $\langle W \rangle_{\text{comp}} = \frac{\hbar}{2}(\omega_2 - \omega_1) \coth(\hbar\omega_1/2k_B T_c)$ and $\langle W \rangle_{\text{exp}} = \frac{\hbar}{2}(\omega_1 - \omega_2) \coth(\hbar\omega_2/2k_B T_h)$, respectively. Then, the total work per (finite-time) cycle, $\langle W \rangle_{\text{comp}} + \langle W \rangle_{\text{exp}}$, equals the work output in the slow adiabatic cycle. This expected result confirms that we have used an STA in the thermodynamic cycle. When computing the corresponding engine's efficiency, without the cost of the STA, one gets the maximum possible value $\eta = 1 - \omega_1/\omega_2$ at finite power [17]. Of course, to be fair with the slow adiabatic cycle, we still need to consider the thermodynamic cost of implementing the shortcut [17–21, 53]. However, there is still an ongoing debate on how to compute this cost correctly and, just as important, how to incorporate it into the engine's efficiency, see [14]. Interestingly, any proposal willing to use the counterdiabatic term $\hat{H}_1(t)$ to compute the cost of the STA will tell us that the energetic cost of producing photons out of the vacuum is related to the cost of the shortcut since $\hat{H}_1(t)$ is the same as the squeezing term of the dynamical Casimir effect.

IV. CONCLUSIONS

We show that with the effective Hamiltonian approach of quantum optics, one can quickly identify that the counterdiabatic term of a shortcut to adiabaticity associated with the time-dependent quantum harmonic oscillator equals (with the opposite sign) the squeezing term inherent to the dynamical Casimir effect. We exhibit the combat between the nonadiabatic evolution generated by the DCE and the STA, the latter trying to enforce an adiabatic dynamic at all times. We confirm the equivalence between the adiabatic and nonadiabatic work outputs when using an STA in a quantum thermodynamic cycle. We also show that the energy cost of producing photons from the vacuum can be related to the shortcut cost. As we mainly deal with ladder operators, these results may be difficult to obtain only with the position and momentum operators, predominately used in the STA literature. Finally, it would be interesting to see other counterdiabatic Hamiltonians, believed to be challenging to implement, and look for their effective Hamiltonian counterpart and vice versa.

ACKNOWLEDGMENTS

The author acknowledges Barış Çakmak for indicating that the counterdiabatic Hamiltonian of the time-dependent harmonic oscillator is a squeezing term.

Appendix A: Adiabatic approximation

For pedagogical purposes, here we follow the procedure carried out in [26] describing the adiabatic approximation in quantum mechanics. One starts with the Schrödinger equation $i\hbar \partial_t |\psi(t)\rangle = \hat{H}_0(t) |\psi(t)\rangle$. A general solution is $|\psi(t)\rangle = \sum_n c_n(t) e^{i\theta_n(t)} |n(t)\rangle$, where $\theta_n(t) = -\frac{1}{\hbar} \int_0^t dt' E_n(t')$ is known as the dynamical phase,

and $|n(t)\rangle$ satisfies $\hat{H}_0(t) |n(t)\rangle = E_n(t) |n(t)\rangle$. $c_n(t)$ are time-dependent expansion coefficients to be determined. Taking the time derivative of $|\psi(t)\rangle$ we get

$$\begin{aligned} \frac{\partial}{\partial t} |\psi(t)\rangle &= \sum_n \dot{c}_n(t) e^{i\theta_n(t)} |n(t)\rangle \\ &\quad + \sum_n i\dot{\theta}_n(t) c_n(t) e^{i\theta_n(t)} |n(t)\rangle \\ &\quad + \sum_n c_n(t) e^{i\theta_n(t)} \frac{\partial}{\partial t} |n(t)\rangle \\ &= \frac{1}{i\hbar} \hat{H}_0(t) |\psi(t)\rangle. \end{aligned} \quad (\text{A1})$$

Using the eigenvalue equation and noting that $\dot{\theta}_n(t) = -E_n(t)/\hbar$, the above equation simplifies to $\sum_n e^{i\theta_n(t)} [\dot{c}_n(t) |n(t)\rangle + c_n(t) \partial_t |n(t)\rangle] = 0$. Taking the inner product with $\langle m(t) |$ and using the orthonormality associated with the eigenstates at equal times, one obtains the following differential equation

$$\dot{c}_m(t) + \sum_n e^{i[\theta_n(t) - \theta_m(t)]} \langle m(t) | \left[\frac{\partial}{\partial t} |n(t)\rangle \right] = 0. \quad (\text{A2})$$

On the other hand, by taking the time derivative for the eigenvalue equation we get $\dot{\hat{H}}_0(t) |n(t)\rangle + \hat{H}_0(t) \partial_t |n(t)\rangle = \dot{E}_n(t) |n(t)\rangle + E_n(t) \partial_t |n(t)\rangle$. It implies that

$$\langle m(t) | \dot{\hat{H}}_0(t) |n(t)\rangle = [E_n(t) - E_m(t)] \langle m(t) | \left[\frac{\partial}{\partial t} |n(t)\rangle \right]. \quad (\text{A3})$$

This means that

$$\langle m(t) | \left[\frac{\partial}{\partial t} |n(t)\rangle \right] = \frac{\langle m(t) | \dot{\hat{H}}_0(t) |n(t)\rangle}{E_n(t) - E_m(t)}. \quad (\text{A4})$$

Substituting this in the equation for $\dot{c}_m(t)$ we get

$$\begin{aligned} \dot{c}_m(t) &= -c_m(t) \langle m(t) | \left[\frac{\partial}{\partial t} |m(t)\rangle \right] \\ &\quad - \sum_{n \neq m} c_n(t) e^{i[\theta_n(t) - \theta_m(t)]} \frac{\langle m(t) | \dot{\hat{H}}_0(t) |n(t)\rangle}{E_n(t) - E_m(t)}. \end{aligned} \quad (\text{A5})$$

The adiabatic approximation consists in neglecting the second term in the right-hand side of the above equation [26], $\dot{c}_m(t) \approx -c_m(t) \langle m(t) | \left[\frac{\partial}{\partial t} |m(t)\rangle \right]$. This has the following solution

$$c_m(t) = \exp \left(- \int_0^t dt' \langle m(t') | \left[\frac{\partial}{\partial t'} |m(t')\rangle \right] \right) c_m(0), \quad (\text{A6})$$

which we can rewrite as $c_m(t) = e^{i\gamma_m(t)} c_m(0)$, where $\gamma_m(t) \equiv i \int_0^t dt' \langle m(t') | \left[\frac{\partial}{\partial t'} |m(t')\rangle \right]$ is known as the geometrical phase or Berry's phase when the evolution is cyclic. Therefore, $|\psi(t)\rangle = \sum_n c_n(0) e^{i\gamma_n(t)} e^{i\theta_n(t)} |n(t)\rangle$. If the system starts in an eigenstate $|n\rangle$ of $\hat{H}_0(t=0)$, then it continues in the eigenstate $|n(t)\rangle$ of $\hat{H}_0(t)$, since $c_i(0) = 0$ unless $i = n$, in that case $c_n(0) = 1$. The solution in the adiabatic approximation is $|\psi_n(t)\rangle = e^{i\gamma_n(t)} e^{i\theta_n(t)} |n(t)\rangle$.

- [1] Viktor Dodonov, “Fifty years of the dynamical Casimir effect,” *Physics* **2**, 67–104 (2020).
- [2] V V Dodonov, “Current status of the dynamical Casimir effect,” *Physica Scripta* **82**, 038105 (2010).
- [3] Gerald T. Moore, “Quantum theory of the electromagnetic field in a variable-length one-dimensional cavity,” *Journal of Mathematical Physics* **11**, 2679–2691 (1970).
- [4] C. M. Wilson, G. Johansson, A. Pourkabirian, M. Simoen, J. R. Johansson, T. Duty, F. Nori, and P. Delsing, “Observation of the dynamical Casimir effect in a superconducting circuit,” *Nature* **479**, 376–379 (2011).
- [5] Pasi Lähteenmäki, G. S. Paraoanu, Juha Hassel, and Pertti J. Hakonen, “Dynamical Casimir effect in a Josephson metamaterial,” *Proceedings of the National Academy of Sciences* **110**, 4234–4238 (2013).
- [6] P. D. Nation, J. R. Johansson, M. P. Blencowe, and Franco Nori, “Colloquium: Stimulating uncertainty: Amplifying the quantum vacuum with superconducting circuits,” *Rev. Mod. Phys.* **84**, 1–24 (2012).
- [7] George Palasantzas, Mehdi Sedighi, and Vitaly B. Svetovoy, “Applications of Casimir forces: Nanoscale actuation and adhesion,” *Applied Physics Letters* **117** (2020).
- [8] Tao Gong, Matthew R. Corrado, Ahmed R. Mahbub, Calum Shelden, and Jeremy N. Munday, “Recent progress in engineering the Casimir effect—applications to nanophotonics, nanomechanics, and chemistry,” *Nanophotonics* **10**, 523–536 (2021).
- [9] Bjarne Andresen and Peter Salamon, “Future perspectives of finite-time thermodynamics,” *Entropy* **24** (2022).
- [10] A V Dodonov, D Valente, and T Werlang, “Quantum power boost in a nonstationary cavity-qed quantum heat engine,” *Journal of Physics A: Mathematical and Theoretical* **51**, 365302 (2018).
- [11] Zhaoyu Fei, Jingning Zhang, Rui Pan, Tian Qiu, and H. T. Quan, “Quantum work distributions associated with the dynamical casimir effect,” *Phys. Rev. A* **99**, 052508 (2019).
- [12] Nahuel Freitas and Juan Pablo Paz, “Fundamental limits for cooling of linear quantum refrigerators,” *Phys. Rev. E* **95**, 012146 (2017).
- [13] Giuliano Benenti and Giuliano Strini, “Dynamical Casimir effect and minimal temperature in quantum thermodynamics,” *Phys. Rev. A* **91**, 020502 (2015).
- [14] D. Guéry-Odelin, A. Ruschhaupt, A. Kiely, E. Torrontegui, S. Martínez-Garaot, and J. G. Muga, “Shortcuts to adiabaticity: Concepts, methods, and applications,” *Rev. Mod. Phys.* **91**, 045001 (2019).
- [15] Erik Torrontegui, Sara Ibáñez, Sofia Martínez-Garaot, Michele Modugno, Adolfo del Campo, David Guéry-Odelin, Andreas Ruschhaupt, Xi Chen, and Juan Gonzalo Muga, “Chapter 2 - shortcuts to adiabaticity,” in *Advances in Atomic, Molecular, and Optical Physics*, Advances In Atomic, Molecular, and Optical Physics, Vol. 62, edited by Ennio Arimondo, Paul R. Berman, and Chun C. Lin (Academic Press, 2013) pp. 117–169.
- [16] Adolfo del Campo and Kihwan Kim, “Focus on shortcuts to adiabaticity,” *New Journal of Physics* **21**, 050201 (2019).
- [17] A. del Campo, J. Goold, and M. Paternostro, “More bang for your buck: Super-adiabatic quantum engines,” *Scientific Reports* **4**, 6208 (2014).
- [18] Obinna Abah and Eric Lutz, “Performance of shortcut-to-adiabaticity quantum engines,” *Phys. Rev. E* **98**, 032121 (2018).
- [19] Barış Çakmak and Özgür E. Müstecaplıoğlu, “Spin quantum heat engines with shortcuts to adiabaticity,” *Phys. Rev. E* **99**, 032108 (2019).
- [20] Obinna Abah and Mauro Paternostro, “Shortcut-to-adiabaticity Otto engine: A twist to finite-time thermodynamics,” *Phys. Rev. E* **99**, 022110 (2019).
- [21] Obinna Abah, Mauro Paternostro, and Eric Lutz, “Shortcut-to-adiabaticity quantum Otto refrigerator,” *Phys. Rev. Res.* **2**, 023120 (2020).
- [22] Tom Feldmann and Ronnie Kosloff, “Quantum lubrication: Suppression of friction in a first-principles four-stroke heat engine,” *Phys. Rev. E* **73**, 025107 (2006).
- [23] J G Muga, X Chen, S Ibáñez, I Lizuain, and A Ruschhaupt, “Transitionless quantum drivings for the harmonic oscillator,” *Journal of Physics B: Atomic, Molecular and Optical Physics* **43**, 085501 (2010).
- [24] M V Berry, “Transitionless quantum driving,” *J. Phys. A Math. Theor.* **42**, 365303 (2009).
- [25] Ralf Schützhold, Günter Plunien, and Gerhard Soff, “Trembling cavities in the canonical approach,” *Phys. Rev. A* **57**, 2311–2318 (1998).
- [26] J. J. Sakurai and Jim Napolitano, *Modern Quantum Mechanics*, 3rd ed. (Cambridge University Press, 2020).
- [27] Christopher Gerry and Peter Knight, *Introductory Quantum Optics* (Cambridge University Press, 2004).
- [28] Girish S Agarwal, *Quantum Optics* (Cambridge University Press, 2012).
- [29] Andrei B Klimov and Sergei M Chumakov, *A Group-theoretical Approach to Quantum Optics: Models of Atoms* (John Wiley & Sons, 2009).
- [30] Héctor Moya-Cessa and Manuel Fernández Guasti, “Coherent states for the time dependent harmonic oscillator: the step function,” *Phys. Lett. A* **311**, 1–5 (2003).
- [31] M. Fernández Guasti and H Moya-Cessa, “Solution of the Schrödinger equation for time-dependent 1D harmonic oscillators using the orthogonal functions invariant,” *Journal of Physics A: Mathematical and General* **36**, 2069–2076 (2003).
- [32] H R Lewis, “Classical and quantum systems with time-dependent harmonic-oscillator-type hamiltonians,” *Phys. Rev. Lett.* **18**, 510–512 (1967).
- [33] I.A. Malkin and V.I. Man’ko, “Coherent states and excitation of N-dimensional non-stationary forced oscillator,” *Physics Letters A* **32**, 243–244 (1970).
- [34] S. Ibáñez, Xi Chen, E. Torrontegui, J. G. Muga, and A. Ruschhaupt, “Multiple Schrödinger pictures and dynamics in shortcuts to adiabaticity,” *Phys. Rev. Lett.* **109**, 100403 (2012).
- [35] Natalia Sánchez-Kuntz, Álvaro Parra-López, Mireia Tolosa-Simeón, Tobias Haas, and Stefan Floerchinger, “Scalar quantum fields in cosmologies with 2 + 1 space-time dimensions,” *Phys. Rev. D* **105**, 105020 (2022).

- [36] Mireia Tolosa-Simeón, Álvaro Parra-López, Natalia Sánchez-Kuntz, Tobias Haas, Celia Viermann, Marius Sparn, Nikolas Liebster, Maurus Hans, Elinor Kath, Helmut Strobel, Markus K. Oberthaler, and Stefan Floerchinger, “Curved and expanding spacetime geometries in bose-einstein condensates,” *Phys. Rev. A* **106**, 033313 (2022).
- [37] R. Román-Ancheyta, M. Berrondo, and J. Récamier, “Parametric oscillator in a Kerr medium: evolution of coherent states,” *J. Opt. Soc. Am. B* **32**, 1651–1655 (2015).
- [38] J Schwinger, “Casimir energy for dielectrics.” *Proceedings of the National Academy of Sciences* **89**, 4091–4095 (1992).
- [39] C. K. Law, “Effective Hamiltonian for the radiation in a cavity with a moving mirror and a time-varying dielectric medium,” *Phys. Rev. A* **49**, 433–437 (1994).
- [40] A. V. Dodonov and V. V. Dodonov, “Dynamical Casimir effect via modulated Kerr or higher-order nonlinearities,” *Phys. Rev. A* **105**, 013709 (2022).
- [41] R. Román-Ancheyta, O. de los Santos-Sánchez, and C. González-Gutiérrez, “Damped Casimir radiation and photon correlation measurements,” *J. Opt. Soc. Am. B* **35**, 523–527 (2018).
- [42] Ricardo Román-Ancheyta, Irán Ramos-Prieto, Armando Perez-Leija, Kurt Busch, and Roberto J. de León-Montiel, “Dynamical Casimir effect in stochastic systems: Photon harvesting through noise,” *Phys. Rev. A* **96**, 032501 (2017).
- [43] R. Román-Ancheyta, C. González-Gutiérrez, and J. Récamier, “Influence of the Kerr nonlinearity in a single nonstationary cavity mode,” *J. Opt. Soc. Am. B* **34**, 1170–1176 (2017).
- [44] A V Dodonov and V V Dodonov, “Photon statistics in the dynamical Casimir effect modified by a harmonic oscillator detector,” *Physica Scripta* **2013**, 014017 (2013).
- [45] A. V. Dodonov and V. V. Dodonov, “Approximate analytical results on the cavity dynamical Casimir effect in the presence of a two-level atom,” *Phys. Rev. A* **85**, 015805 (2012).
- [46] Nicolás F. Del Grosso, Fernando C. Lombardo, Francisco D. Mazzitelli, and Paula I. Villar, “Shortcut to adiabaticity in a cavity with a moving mirror,” *Phys. Rev. A* **105**, 052217 (2022).
- [47] J Jaramillo, M Beau, and A del Campo, “Quantum supremacy of many-particle thermal machines,” *New Journal of Physics* **18**, 75019 (2016).
- [48] A del Campo, Aurélia Chenu, Shujin Deng, and Haibin Wu, “Friction-free quantum machines,” in *Thermodynamics in the Quantum Regime: Fundamental Aspects and Applications*, edited by Felix Binder, Luis A. Correa, Christian Gogolin, Janet Anders, and Gerardo Adesso (Springer International Publishing, Cham, 2018) pp. 127–148.
- [49] O. Abah, J. Roßnagel, G. Jacob, S. Deffner, F. Schmidt-Kaler, K. Singer, and E. Lutz, “Single-ion heat engine at maximum power,” *Phys. Rev. Lett.* **109**, 203006 (2012).
- [50] Johannes Roßnagel, Samuel T Dawkins, Karl N Tolazzi, Obinna Abah, Eric Lutz, Ferdinand Schmidt-Kaler, and Kilian Singer, “A single-atom heat engine,” *Science* **352**, 325–329 (2016).
- [51] H. T. Quan, Yu-xi Liu, C. P. Sun, and Franco Nori, “Quantum thermodynamic cycles and quantum heat engines,” *Phys. Rev. E* **76**, 031105 (2007).
- [52] Ronnie Kosloff and Yair Rezek, “The quantum harmonic Otto cycle,” *Entropy* **19** (2017), 10.3390/e19040136.
- [53] Qiongyuan Wu, Mario A. Ciampini, Mauro Paternostro, and Matteo Carlesso, “Quantifying protocol efficiency: A thermodynamic figure of merit for classical and quantum state-transfer protocols,” *Phys. Rev. Res.* **5**, 023117 (2023).