

DESCRIPTION OF ALL FROBENIUS ALGEBRA STRUCTURES ON TWO-DIMENSIONAL VECTOR SPACE OVER ANY BASIC FIELD

D.ASROROV¹, U.BEKBAEV², I.RAKHIMOV³

¹Samarqand State University, Samarqand, Uzbekistan

²Turin Polytechnic University in Tashkent, Tashkent, Uzbekistan

³Universiti Teknologi MARA (UiTM), Shah Alam, Malaysia

ABSTRACT. In the paper we classify all possible Frobenius algebra structures on two-dimensional vector space over any basic field. We give the lists of canonical representatives of isomorphism classes, explicitly, over basic field with the characteristics not 2,3, characteristic 2 and characteristic 3, respectively.

1. INTRODUCTION

In mathematics, especially in the fields of representation theory and module theory, a Frobenius algebra is a finite-dimensional unital associative algebra with a special kind of bilinear form which gives the algebras particularly nice duality theories. Frobenius algebras began to be studied in the 1930s by Richard Brauer and Cecil Nesbitt, and were named after Georg Frobenius [3]. T. Nakayama discovered the beginnings of a rich duality theory [5, 6]. J. Dieudonné used this to characterize Frobenius algebras [4]. Frobenius algebras were generalized to quasi-Frobenius rings, those Noetherian rings whose right regular representation is injective. In recent times, interest has been renewed in Frobenius algebras due to connections to topological quantum field theory. It has been found that they play an important role in the algebraic treatment and axiomatic foundation of topological quantum field theory [1, 10].

2. PRELIMINARIES

Let \mathbb{A} be a PI-algebra, with a given set of polynomial identities $\{P_j[x_1, x_2, \dots, x_n] = 0 : j \in J\}$, over a field \mathbb{F} . In some applications this kind of algebras appear with a non-degenerate bilinear form $\sigma : \mathbb{A} \times \mathbb{A} \rightarrow \mathbb{F}$ for which $\sum_{i=1}^{k_j} \sigma(Q_j^i[a_1, a_2, \dots, a_n], R_j^i[a_1, a_2, \dots, a_n]) = 0$ at all $a_1, a_2, \dots, a_n \in \mathbb{A}$, where $P_j[x_1, x_2, \dots, x_n] = \sum_{i=1}^{k_j} Q_j^i[x_1, x_2, \dots, x_n]R_j^i[x_1, x_2, \dots, x_n]$, $j \in J$. Such pair (\mathbb{A}, σ) is said to be a Frobenius PI-algebra. The classification of a given class of Frobenius PI-algebras is of great interest. In this paper we consider as a class PI-algebras the class of associative algebras and provide a complete description of the corresponding class of Frobenius algebras on two-dimensional space over arbitrary field.

Definition 1. *Let (\mathbb{A}, σ) and (\mathbb{B}, τ) be finite-dimensional algebras over a field \mathbb{F} with non-degenerate bilinear forms $\sigma : \mathbb{A} \times \mathbb{A} \rightarrow \mathbb{F}$ and $\tau : \mathbb{B} \times \mathbb{B} \rightarrow \mathbb{F}$, respectively. Pairs (\mathbb{A}, σ) and (\mathbb{B}, τ) are said to be isomorphic if there exists isomorphism of algebras $f : \mathbb{A} \rightarrow \mathbb{B}$ such that $\sigma(xy) = \tau(f(x)f(y))$ for all $x, y \in \mathbb{A}$.*

emails: ¹96asrorovdiorjon@mail.ru; ²uralbekbaev@gmail.com; ³isamidin@uitm.edu.my.

Definition 2. A pair (\mathbb{A}, σ) , where \mathbb{A} is a finite-dimensional, associative algebra, $\sigma : \mathbb{A} \times \mathbb{A} \rightarrow \mathbb{F}$ is a nondegenerate bilinear form, is said to be a Frobenius algebra if $\sigma(x \cdot y, z) = \sigma(x, y \cdot z)$ for all $x, y, z \in \mathbb{A}$.

The bilinear form σ is called the Frobenius form of the algebra \mathbb{A} .

Examples.

- (1) Any matrix algebra defined over a field \mathbb{F} is a Frobenius algebra with Frobenius form $\sigma(A, B) = \text{tr}(A \times B)$, where $\text{tr}(X)$ is the trace of matrix X .
- (2) Any finite-dimensional unital associative algebra \mathbb{A} has a natural homomorphism to its own endomorphism ring $\text{End}(\mathbb{A})$. A bilinear form can be defined on \mathbb{A} in the sense of the previous example. If this bilinear form is nondegenerate, then it equips \mathbb{A} with the structure of a Frobenius algebra.
- (3) Every group ring $\mathbb{F}[G]$ of a finite group G over a field \mathbb{F} is a symmetric Frobenius algebra, with Frobenius form $\sigma(f, g)$ given by the coefficient of the identity element in fg .

Properties.

- The direct product and tensor product of Frobenius algebras are Frobenius algebras.
- A finite-dimensional commutative local algebra over a field is Frobenius if and only if the right regular module is injective, if and only if the algebra has a unique minimal ideal.
- Commutative, local Frobenius algebras are precisely the zero-dimensional local Gorenstein rings containing their residue field and finite-dimensional over it.
- Frobenius algebras are quasi-Frobenius rings, and in particular, they are left and right Artinian and left and right self-injective.
- For a field \mathbb{F} , a finite-dimensional, unital, associative algebra is Frobenius if and only if the injective right \mathbb{A} -module $\text{Hom}_{\mathbb{F}}(\mathbb{A}, \mathbb{F})$ is isomorphic to the right regular representation of \mathbb{A} .
- For an infinite field \mathbb{F} , a finite-dimensional, unital, associative \mathbb{F} -algebra is a Frobenius algebra if it has only finitely many minimal right ideals.

In this paper we focus on two-dimensional algebras.

Let \mathbb{A} be a two-dimensional algebra over a field \mathbb{F} and $\mathbf{e} = (e_1, e_2)$ be a basis and

$$A = \begin{pmatrix} \alpha_1 & \alpha_2 & \alpha_3 & \alpha_4 \\ \beta_1 & \beta_2 & \beta_3 & \beta_4 \end{pmatrix}$$

be MSC of \mathbb{A} on the basis \mathbf{e} , that is $e_1 \mathbf{e} = (e_1^2, e_1 e_2) = \mathbf{e} \begin{pmatrix} \alpha_1 & \alpha_2 \\ \beta_1 & \beta_2 \end{pmatrix}$, $e_2 \mathbf{e} = (e_2 e_1, e_2^2) = \mathbf{e} \begin{pmatrix} \alpha_3 & \alpha_4 \\ \beta_3 & \beta_4 \end{pmatrix}$.

Any non-degenerate bilinear form $\sigma : \mathbb{A} \times \mathbb{A} \rightarrow \mathbb{F}$ in $\mathbf{e} = (e_1, e_2)$ is defined as $\sigma(x, y) = x^t \sigma y$, where $\sigma = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in GL(2, \mathbb{F})$, $x = \mathbf{e}x, y = \mathbf{e}y$ and $x = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}, y = \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} \in \mathbb{F}^2$ are column coordinate vectors of x, y , respectively. Note that further we do not make difference between an algebra \mathbb{A} and its MSC A , between bilinear form σ and its matrix σ on the basis \mathbf{e} , I stands for the second order identity matrix.

In [9] the classification, up to isomorphism, of two-dimensional associative algebras, over any basic field \mathbb{F} and their automorphism groups were given. The result was based on technique applied in [2]. Therefore, we give the result here keeping the notations of [2] as the following theorems for the characteristic of \mathbb{F} to be not two and three, to be two and to be three, respectively.

Theorem 1. Any non-trivial 2-dimensional associative algebra over a field \mathbb{F} ($\text{Char}(\mathbb{F}) \neq 2, 3$) is isomorphic to only one of the following listed, by their matrices of structure constants, algebras:

- $A_3(1,0,0) = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$,
- $A_3(1,0,1) = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}$,
- $A_3(\frac{1}{2},0,0) = \begin{pmatrix} \frac{1}{2} & 0 & 0 & 0 \\ 0 & 0 & \frac{1}{2} & 0 \end{pmatrix}$,
- $A_3(\frac{1}{2},\alpha_4,\frac{1}{2}) = \begin{pmatrix} \frac{1}{2} & 0 & 0 & \alpha_4 \\ 0 & \frac{1}{2} & \frac{1}{2} & 0 \end{pmatrix} \simeq \begin{pmatrix} \frac{1}{2} & 0 & 0 & a^2\alpha_4 \\ 0 & \frac{1}{2} & \frac{1}{2} & 0 \end{pmatrix}$, where $\alpha_4, a \in \mathbb{F}$, $a \neq 0$,
- $A_{13} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}$.

Among the listed above algebras only $A_3(\frac{1}{2},\alpha_4,\frac{1}{2})$ is unital and its unite element is $\mathbf{1} = \begin{pmatrix} 2 \\ 0 \end{pmatrix}$.

Theorem 2. Any non-trivial 2-dimensional associative algebra over a field \mathbb{F} , ($\text{Char}(\mathbb{F}) = 2$) is isomorphic to only one of the following listed by their matrices of structure constants, such algebras:

- $A_{3,2}(1,0,0) = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$,
- $A_{3,2}(1,0,1) = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}$,
- $A_{4,2}(1,\beta_1,0) = \begin{pmatrix} 1 & 1 & 1 & 0 \\ \beta_1 & 0 & 0 & 1 \end{pmatrix} \cong \begin{pmatrix} 1 & 1 & 1 & 0 \\ \beta_1 + a + a^2 & 0 & 0 & 1 \end{pmatrix}$, where $a, \beta_1 \in \mathbb{F}$,
- $A_{6,2}(1,0) = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}$,
- $A_{11,2}(\beta_1) = \begin{pmatrix} 0 & 1 & 1 & 0 \\ \beta_1 & 0 & 0 & 1 \end{pmatrix} \cong \begin{pmatrix} 0 & 1 & 1 & 0 \\ b^2\beta_1 + a^2 & 0 & 0 & 1 \end{pmatrix}$, where $\beta_1, a, b \in \mathbb{F}$, $b \neq 0$,
- $A_{12,2} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}$.

Among the algebras listed above only $A_{4,2}(1,\beta_1,0)$ and $A_{11,2}(\beta_1)$ are unital and theirs unite element is $\mathbf{1} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$.

Theorem 3. Any non-trivial 2-dimensional associative algebra over a field \mathbb{F} , ($\text{Char}(\mathbb{F}) = 3$) is isomorphic to only one of the following listed by their matrices of structure constants, such algebras:

- $A_{3,3}(1,0,0) = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$,
- $A_{3,3}(1,0,1) = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}$,
- $A_{3,3}(2,0,0) = \begin{pmatrix} 2 & 0 & 0 & 0 \\ 0 & 0 & 2 & 0 \end{pmatrix}$,
- $A_{3,3}(2,\alpha_4,2) := \begin{pmatrix} 2 & 0 & 0 & \alpha_4 \\ 0 & 2 & 2 & 0 \end{pmatrix} \simeq \begin{pmatrix} 2 & 0 & 0 & a^2\alpha_4 \\ 0 & 2 & 2 & 0 \end{pmatrix}$, where $\alpha_4, a \in \mathbb{F}$, $a \neq 0$,
- $A_{13,3} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}$.

Here only $A_{3,3}(2,\alpha_4,2)$ is unital and its unite element is $\mathbf{1} = \begin{pmatrix} 2 \\ 0 \end{pmatrix}$.

Theorem 4. *The automorphism groups of the algebras listed in Theorem 1 are given as follows*

$$\begin{aligned} \text{Aut}(A_3(1, 0, 0)) &= \left\{ \begin{pmatrix} 1 & 0 \\ 0 & d \end{pmatrix} : 0 \neq d \in \mathbb{F} \right\}, \\ \text{Aut}(A_3(1, 0, 1)) &= \left\{ \begin{pmatrix} 1 & 0 \\ c & d \end{pmatrix} : c, d \in \mathbb{F}, d \neq 0 \right\}, \\ \text{Aut}(A_3(\frac{1}{2}, 0, 0)) &= \left\{ \begin{pmatrix} 1 & 0 \\ c & d \end{pmatrix} : c, d \in \mathbb{F}, d \neq 0 \right\}, \\ \text{Aut}(A_3(\frac{1}{2}, 0, \frac{1}{2})) &= \left\{ \begin{pmatrix} 1 & 0 \\ 0 & d \end{pmatrix} : d \neq 0 \right\}, \\ \text{Aut}(A_3(\frac{1}{2}, \alpha_4, \frac{1}{2})) &= \left\{ \begin{pmatrix} 1 & 0 \\ 0 & \pm 1 \end{pmatrix} \right\}, \text{ whenever } \alpha_4 \neq 0, \\ \text{Aut}(A_{13}) &= \left\{ \begin{pmatrix} a & 0 \\ c & a^2 \end{pmatrix} : a, c \in \mathbb{F}, a \neq 0 \right\}. \end{aligned}$$

Theorem 5. *The automorphism groups of the algebras listed in Theorem 2 are given as follows*

$$\begin{aligned} \text{Aut}(A_{3,2}(1, 0, 0)) &= \left\{ \begin{pmatrix} 1 & 0 \\ 0 & d \end{pmatrix} : 0 \neq d \in \mathbb{F} \right\}, \\ \text{Aut}(A_{3,2}(1, 0, 1)) &= \left\{ \begin{pmatrix} 1 & 0 \\ c & d \end{pmatrix} : c, d \in \mathbb{F}, d \neq 0 \right\}, \\ \text{Aut}(A_{4,2}(1, 0, 0)) &= \left\{ \begin{pmatrix} a & 0 \\ c & 1 \end{pmatrix}, a \neq 0, c \in \mathbb{F} \right\}, \\ \text{Aut}(A_{4,2}(1, \beta_1, 0)) &= \left\{ \begin{pmatrix} 1 & 0 \\ c & 1 \end{pmatrix} : c \in \mathbb{F} \right\}, \\ \text{Aut}(A_{6,2}(1, 0)) &= \left\{ I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \right\}, \\ \text{Aut}(A_{11,2}(\beta_1)) &= \left\{ \begin{pmatrix} a & 0 \\ \beta_1(a-1) & 1 \end{pmatrix} : a \neq 0 \in \mathbb{F} \right\}, \\ \text{Aut}(A_{12,2}) &= \left\{ \begin{pmatrix} a & 0 \\ c & a^2 \end{pmatrix} : a, c \in \mathbb{F}, a \neq 0 \right\}. \end{aligned}$$

Theorem 6. *The automorphism groups of the algebras listed in Theorem 3 are given as follows*

$$\begin{aligned} \text{Aut}(A_{3,3}(1, 0, 1)) &= \left\{ \begin{pmatrix} 1 & 0 \\ c & d \end{pmatrix} : c, d \in \mathbb{F}, d \neq 0 \right\}, \\ \text{Aut}(A_{3,3}(1, 0, 0)) &= \left\{ \begin{pmatrix} 1 & 0 \\ 0 & d \end{pmatrix} : 0 \neq d \in \mathbb{F} \right\}, \\ \text{Aut}(A_{3,3}(2, 0, 0)) &= \left\{ \begin{pmatrix} 1 & 0 \\ 1+2d & d \end{pmatrix} : 0 \neq d \in \mathbb{F} \right\}, \\ \text{Aut}(A_{3,3}(2, 0, 2)) &= \left\{ I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \right\}, \\ \text{Aut}(A_{3,3}(2, \alpha_4, 2)) &= \left\{ \begin{pmatrix} 1 & 0 \\ c & 1 \end{pmatrix} : c \in \mathbb{F} \right\}, \text{ provided that } \alpha_4 \neq 0, \\ \text{Aut}(A_{13,3}) &= \left\{ \begin{pmatrix} a & 0 \\ c & 2a^2 \end{pmatrix} : a, c \in \mathbb{F}, a \neq 0 \right\}. \end{aligned}$$

Note that if

$$x = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}, \quad y = \begin{pmatrix} y_1 \\ y_2 \end{pmatrix}, \quad z = \begin{pmatrix} z_1 \\ z_2 \end{pmatrix}, \quad A = \begin{pmatrix} \alpha_1 & \alpha_2 & \alpha_3 & \alpha_4 \\ \beta_1 & \beta_2 & \beta_3 & \beta_4 \end{pmatrix}, \quad \sigma = \begin{pmatrix} a & b \\ c & d \end{pmatrix}, \quad \sigma(x, y) = x^t \sigma y$$

then the condition $\sigma(x \cdot y, z) = \sigma(x, y \cdot z)$ is nothing but $(x^t \otimes y^t)A^t gz = x^t gA(y \otimes z)$ and therefore, due to

$$(2.1) \quad \begin{aligned} (x^t \otimes y^t)A^t gz &= (\alpha_1 a + \beta_1 c)x_1 y_1 z_1 + (\alpha_2 a + \beta_2 c)x_1 y_2 z_1 + (\alpha_3 a + \beta_3 c)x_2 y_1 z_1 \\ &\quad + (\alpha_4 a + \beta_4 c)x_2 y_2 z_1 + (\alpha_1 b + \beta_1 d)x_1 y_1 z_2 + (\alpha_2 b + \beta_2 d)x_1 y_2 z_2 \\ &\quad + (\alpha_3 b + \beta_3 d)x_2 y_1 z_2 + (\alpha_4 b + \beta_4 d)x_2 y_2 z_2 \\ x^t gA(y \otimes z) &= (\alpha_1 a + \beta_1 b)x_1 y_1 z_1 + (\alpha_3 a + \beta_3 b)x_1 y_2 z_1 + (\alpha_1 c + \beta_1 d)x_2 y_1 z_1 \\ &\quad + (\alpha_3 c + \beta_3 d)x_2 y_2 z_1 + (\alpha_2 a + \beta_2 b)x_1 y_1 z_2 \\ &\quad + (\alpha_4 a + \beta_4 b)x_1 y_2 z_2 + (\alpha_2 c + \beta_2 d)x_2 y_1 z_2 + (\alpha_4 c + \beta_4 d)x_2 y_2 z_2 \end{aligned}$$

the condition $(x^t \otimes y^t)A^t gz = x^t gA(y \otimes z)$ is equivalent to the following system of equalities

$$(2.2) \quad \begin{cases} \alpha_1 a + \beta_1 c - \alpha_1 a - \beta_1 b = 0 \\ \alpha_2 a + \beta_2 c - \alpha_3 a - \beta_3 b = 0 \\ \alpha_3 a + \beta_3 c - \alpha_1 c - \beta_1 d = 0 \\ \alpha_4 a + \beta_4 c - \alpha_3 c - \beta_3 d = 0 \\ \alpha_1 b + \beta_1 d - \alpha_2 a - \beta_2 b = 0 \\ \alpha_2 b + \beta_2 d - \alpha_4 a - \beta_4 b = 0 \\ \alpha_3 b + \beta_3 d - \alpha_2 c - \beta_2 d = 0 \\ \alpha_4 b + \beta_4 d - \alpha_4 c - \beta_4 d = 0 \end{cases}$$

as far as the system of monomial functions $\{x_1 y_1 z_1, x_1 y_2 z_1, x_2 y_1 z_1, x_1 y_1 z_2, x_2 y_2 z_1, x_1 y_2 z_2, x_2 y_1 z_2, x_2 y_2 z_2\}$ is linear independent over \mathbb{F} .

3. CLASSIFICATION

In this section first we classify non-degenerate bilinear forms given as $\sigma = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix}$ with respect to the transformations $g^t \sigma g$, $g \in G$, where G is a fixed nontrivial automorphism's group from Theorems 4 - 6. For the further usage we enumerate as G_i , $i = 1, 2, \dots, 8$ the list of all nontrivial autmorphisms' groups in Theorems 4 - 6 as follows

$$\begin{aligned} G_1 &= \left\{ \begin{pmatrix} 1 & 0 \\ 0 & d \end{pmatrix} : 0 \neq d \in \mathbb{F} \right\}, \quad G_2 = \left\{ \begin{pmatrix} 1 & 0 \\ c & d \end{pmatrix} : c, d \in \mathbb{F}, d \neq 0 \right\}, \quad G_3 = \left\{ \begin{pmatrix} 1 & 0 \\ c & 1 \end{pmatrix} : c \in \mathbb{F} \right\}, \\ G_4 &= \left\{ \begin{pmatrix} a & 0 \\ c & a^2 \end{pmatrix} : a, c \in \mathbb{F}, a \neq 0 \right\}, \quad G_5 = \left\{ \begin{pmatrix} a & 0 \\ c & 2a^2 \end{pmatrix} : a, c \in \mathbb{F}, a \neq 0 \right\} (\text{Char}(\mathbb{F}) = 3), \\ G_{6, \beta_1} &= \left\{ \begin{pmatrix} a & 0 \\ \beta_1(a-1) & 1 \end{pmatrix} : a, c \in \mathbb{F}, a \neq 0 \right\}, \quad G_7 = \left\{ \begin{pmatrix} 1 & 0 \\ 1+2d & d \end{pmatrix} : d \in \mathbb{F}, d \neq 0 \right\} (\text{Char}(\mathbb{F}) = 3), \\ G_8 &= \left\{ \begin{pmatrix} 1 & 0 \\ 0 & \pm 1 \end{pmatrix} \right\} (\text{Char}(\mathbb{F}) \neq 2). \end{aligned}$$

Now we treat the action $g^t \sigma g$ for each G_i , $i = 1, 2, \dots, 8$ ($g \in G_i$) one by one.

Let $g = \begin{pmatrix} 1 & 0 \\ 0 & d \end{pmatrix} \in G_1$. Then $g^t \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} g = \begin{pmatrix} \alpha & d\beta \\ d\gamma & d^2\delta \end{pmatrix}$ and the following canonical forms occur:

- $\begin{pmatrix} \alpha & 1 \\ \gamma & \delta \end{pmatrix}$, where $\alpha\delta - \gamma \neq 0$,

- $\begin{pmatrix} \alpha & 0 \\ 1 & \delta \end{pmatrix}$, where $\alpha\delta \neq 0$,
- $\begin{pmatrix} \alpha & 0 \\ 0 & \delta \end{pmatrix} \simeq \begin{pmatrix} \alpha & 0 \\ 0 & d^2\delta \end{pmatrix}$, where $0 \neq d \in \mathbb{F}, \alpha\delta \neq 0$.

If $g = \begin{pmatrix} 1 & 0 \\ c & d \end{pmatrix} \in G_2$ then $g^t \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} g = \begin{pmatrix} \alpha + c\gamma + c\beta + c^2\delta & d\beta + cd\delta \\ d\gamma + cd\delta & d^2\delta \end{pmatrix}$ and one comes to the following canonical forms:

- $\begin{pmatrix} \alpha & 0 \\ 1 & \delta \end{pmatrix}$, where $\alpha\delta \neq 0$:
- $\begin{pmatrix} \alpha & 0 \\ 0 & \delta \end{pmatrix} \simeq \begin{pmatrix} \alpha & 0 \\ 0 & d^2\delta \end{pmatrix}$, where $0 \neq d \in \mathbb{F}, \alpha\delta \neq 0$,
- $\begin{pmatrix} 0 & 1 \\ \gamma & 0 \end{pmatrix}$, where $\gamma \neq 0, \gamma + 1 \neq 0$,
- $\begin{pmatrix} \alpha & 1 \\ -1 & 0 \end{pmatrix}$, where $\alpha \in \mathbb{F}$.

Let $g = \begin{pmatrix} 1 & 0 \\ c & 1 \end{pmatrix} \in G_3$. Then $g^t \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} g = \begin{pmatrix} \alpha + c\gamma + c\beta + c^2\delta & \beta + c\delta \\ \gamma + c\delta & \delta \end{pmatrix}$ and we get the following canonical forms:

- $\begin{pmatrix} \alpha & 0 \\ \gamma & \delta \end{pmatrix}$, where $\alpha\delta \neq 0$,
- $\begin{pmatrix} 0 & \beta \\ \gamma & 0 \end{pmatrix}$, where $\beta\gamma \neq 0, \beta + \gamma \neq 0$,
- $\begin{pmatrix} \alpha & \beta \\ -\beta & 0 \end{pmatrix}$, where $\beta \neq 0$.

If $g = \begin{pmatrix} a & 0 \\ c & a^2 \end{pmatrix} \in G_4$, where $a, c \in \mathbb{F}, a \neq 0$, then $g^t \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} g = \begin{pmatrix} a^2\alpha + ac\gamma + ac\beta + c^2\delta & a^3\beta + a^2c\delta \\ a^3\gamma + a^2c\delta & a^4\delta \end{pmatrix}$

and the following canonical forms occur:

- $\begin{pmatrix} \alpha & 0 \\ \gamma & \delta \end{pmatrix} \simeq \begin{pmatrix} a^2\alpha & 0 \\ a^3\gamma & a^4\delta \end{pmatrix}$, where $a \neq 0$,
- $\begin{pmatrix} 0 & \beta \\ \gamma & 0 \end{pmatrix} \simeq \begin{pmatrix} 0 & a^3\beta \\ a^3\gamma & 0 \end{pmatrix}$, where $a \neq 0$,
- $\begin{pmatrix} \alpha & \beta \\ -\beta & 0 \end{pmatrix} \simeq \begin{pmatrix} a^2\alpha & a^3\beta \\ -a^3\beta & 0 \end{pmatrix}$, where $a \neq 0$.

Let now $g = \begin{pmatrix} a & 0 \\ c & 2a^2 \end{pmatrix} \in G_5$, where $a, c \in \mathbb{F}, a \neq 0$ and $Char(\mathbb{F}) = 3$. Then we have

$$g^t \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} g = \begin{pmatrix} a^2\alpha + ac\gamma + ac\beta + c^2\delta & 2a^3\beta + 2a^2c\delta \\ 2a^3\gamma + 2a^2c\delta & a^4\delta \end{pmatrix}$$

and the canonical forms are given as follows:

- $\begin{pmatrix} \alpha & 0 \\ \gamma & \delta \end{pmatrix} \simeq \begin{pmatrix} a^2\alpha & 0 \\ a^3\gamma & a^4\delta \end{pmatrix}$, where $\alpha\delta \neq 0$,
- $\begin{pmatrix} 0 & \beta \\ \gamma & 0 \end{pmatrix} \simeq \begin{pmatrix} 0 & a^3\beta \\ a^3\gamma & 0 \end{pmatrix}$, where $\beta\gamma \neq 0$,

- $\begin{pmatrix} \alpha & \beta \\ -\beta & 0 \end{pmatrix} \simeq \begin{pmatrix} a^2\alpha & a^3\beta \\ -a^3\beta & 0 \end{pmatrix}$, where $\beta \neq 0$.

If $g = \left\{ \begin{pmatrix} a & 0 \\ \beta_1(a-1) & 1 \end{pmatrix} : a \neq 0 \in \mathbb{F} \right\} \in G_{6,\beta_1}$ then

$$g^t \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} g = \begin{pmatrix} a^2\alpha + \beta_1(a+1)(a\gamma + a\beta + \beta_1\delta(a+1)) & a\beta + \beta_1\delta(a+1) \\ a\gamma + \beta_1\delta(a+1) & \delta \end{pmatrix}$$

and one gets the following canonical forms:

- $\begin{pmatrix} \alpha & 0 \\ 1 & \delta \end{pmatrix}$,
- $\begin{pmatrix} \alpha & 0 \\ 0 & \delta \end{pmatrix} \simeq \begin{pmatrix} a^2\alpha & 0 \\ 0 & \delta \end{pmatrix}$, where $0 \neq a \in \mathbb{F}$,
- $\begin{pmatrix} 0 & 1 \\ \gamma & 0 \end{pmatrix}$, where $\gamma \neq 0, \gamma + 1 \neq 0$,
- $\begin{pmatrix} \alpha & 1 \\ -1 & 0 \end{pmatrix}$, where $\alpha \in F$.

Let $G_7 = \left\{ \begin{pmatrix} 1 & 0 \\ 1+2d & d \end{pmatrix} : d \in \mathbb{F}, d \neq 0 \right\} (Char(\mathbb{F}) = 3)$. Then one has

$$g^t \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} g = \begin{pmatrix} \alpha + \gamma(1+2d) + (1+2d)(\beta + \delta(1+2d)) & d(\beta + \delta(1+2d)) \\ d\gamma + d\delta(1+2d) & d^2\delta \end{pmatrix}$$

and the following canonical forms occur:

- $\begin{pmatrix} \alpha & 0 \\ \gamma & \delta \end{pmatrix}$, where $\alpha\delta \neq 0$,
- $\begin{pmatrix} \alpha & \beta \\ 0 & \delta \end{pmatrix}$, where $\alpha\delta \neq 0$,
- $\begin{pmatrix} \alpha & -\delta \\ -\delta & \delta \end{pmatrix} \simeq \begin{pmatrix} \alpha - \delta + d^2\delta & -d^2\delta \\ -d^2\delta & d^2\delta \end{pmatrix}$, where $\delta(\alpha - \delta) \neq 0$,
- $\begin{pmatrix} \alpha & 1 \\ \gamma & 0 \end{pmatrix}$, where $\gamma \neq 0$.

Finally, taking $g = \begin{pmatrix} 1 & 0 \\ 0 & \pm 1 \end{pmatrix} \in G_8$ we get $g^t \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} g = \begin{pmatrix} \alpha & \pm\beta \\ \pm\gamma & \delta \end{pmatrix}$ and only the canonical form appears:

- $\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \simeq \begin{pmatrix} \alpha & -\beta \\ -\gamma & \delta \end{pmatrix}$, where $\alpha\delta - \beta\gamma \neq 0$.

Now we turn to pairs (\mathbb{A}, σ) , where \mathbb{A} is a two-dimensional associative algebra, $\sigma : \mathbb{A} \times \mathbb{A} \rightarrow \mathbb{F}$ is a non-degenerate bilinear form, up to isomorphism.

Therefore, Theorems 1-6 and the classification canonical forms of bilinear forms above enable us to state the following results.

Lemma 3.1. *Every pair (\mathbb{A}, σ) , where \mathbb{A} is a two dimensional associative algebra over a field \mathbb{F} ($Char(\mathbb{F}) \neq 2, 3$), $\sigma : \mathbb{A} \times \mathbb{A} \rightarrow \mathbb{F}$ is a nondegerate form, is isomorphic to only one of the following pairs*

$$(1) \left(A_3(1, 0, 0), \begin{pmatrix} \alpha & 1 \\ \gamma & \delta \end{pmatrix} \right), \text{ where } \alpha, \gamma, \delta \in \mathbb{F}, \alpha\delta - \gamma \neq 0,$$

$$(2) \left(A_3(1, 0, 0), \begin{pmatrix} \alpha & 0 \\ 1 & \delta \end{pmatrix} \right), \text{ where } \alpha, \delta \in \mathbb{F}, \alpha\delta \neq 0,$$

$$(3) \left(A_3(1, 0, 0), \begin{pmatrix} \alpha & 0 \\ 0 & \delta \end{pmatrix} \right) \simeq \left(A_3(1, 0, 0), \begin{pmatrix} \alpha & 0 \\ 0 & d^2\delta \end{pmatrix} \right), \text{ where } \alpha, \delta, d \in \mathbb{F}, \alpha\delta \neq 0, d \neq 0,$$

$$(4) \left(A_3(1, 0, 1), \begin{pmatrix} \alpha & 0 \\ 1 & \delta \end{pmatrix} \right), \text{ where } \alpha, \delta \in \mathbb{F}, \alpha\delta \neq 0,$$

$$(5) \left(A_3(1, 0, 1), \begin{pmatrix} \alpha & 0 \\ 0 & \delta \end{pmatrix} \right) \simeq \left(A_3(1, 0, 1), \begin{pmatrix} \alpha & 0 \\ 0 & d^2\delta \end{pmatrix} \right), \text{ where } \alpha, \delta, d \in \mathbb{F}, \alpha\delta \neq 0, d \neq 0,$$

$$(6) \left(A_3(1, 0, 1), \begin{pmatrix} 0 & 1 \\ \gamma & 0 \end{pmatrix} \right), \text{ where } \gamma \in \mathbb{F}, \gamma \neq 0, \gamma + 1 \neq 0,$$

$$(7) \left(A_3(1, 0, 1), \begin{pmatrix} \alpha & 1 \\ -1 & 0 \end{pmatrix} \right), \text{ where } \alpha \in \mathbb{F},$$

$$(8) \left(A_3\left(\frac{1}{2}, 0, 0\right), \begin{pmatrix} \alpha & 0 \\ 1 & \delta \end{pmatrix} \right), \text{ where } \alpha, \delta \in \mathbb{F}, \alpha\delta \neq 0,$$

$$(9) \left(A_3\left(\frac{1}{2}, 0, 0\right), \begin{pmatrix} \alpha & 0 \\ 0 & \delta \end{pmatrix} \right) \simeq \left(A_3\left(\frac{1}{2}, 0, 0\right), \begin{pmatrix} \alpha & 0 \\ 0 & d^2\delta \end{pmatrix} \right), \text{ where } \alpha, \delta, d \in \mathbb{F}, \alpha\delta \neq 0, d \neq 0,$$

$$(10) \left(A_3\left(\frac{1}{2}, 0, 0\right), \begin{pmatrix} 0 & 1 \\ \gamma & 0 \end{pmatrix} \right), \text{ where } \gamma \in \mathbb{F}, \gamma \neq 0, \gamma + 1 \neq 0,$$

$$(11) \left(A_3\left(\frac{1}{2}, 0, 0\right), \begin{pmatrix} \alpha & 1 \\ -1 & 0 \end{pmatrix} \right), \text{ where } \alpha \in \mathbb{F},$$

$$(12) \left(A_3\left(\frac{1}{2}, 0, \frac{1}{2}\right), \begin{pmatrix} \alpha & 1 \\ \gamma & \delta \end{pmatrix} \right), \text{ where } \alpha, \gamma, \delta \in \mathbb{F}, \alpha\delta - \gamma \neq 0,$$

$$(13) \left(A_3\left(\frac{1}{2}, 0, \frac{1}{2}\right), \begin{pmatrix} \alpha & 0 \\ 1 & \delta \end{pmatrix} \right), \text{ where } \alpha, \delta \in \mathbb{F}, \alpha\delta \neq 0,$$

$$(14) \left(A_3\left(\frac{1}{2}, 0, \frac{1}{2}\right), \begin{pmatrix} \alpha & 0 \\ 0 & \delta \end{pmatrix} \right) \simeq \left(A_3\left(\frac{1}{2}, 0, \frac{1}{2}\right), \begin{pmatrix} \alpha & 0 \\ 0 & d^2\delta \end{pmatrix} \right), \text{ where } \alpha, \delta, d \in \mathbb{F}, \alpha\delta \neq 0, d \neq 0,$$

$$(15) \left(A_3\left(\frac{1}{2}, \alpha_4, \frac{1}{2}\right), \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \right) \simeq \left(A_3\left(\frac{1}{2}, \alpha_4, \frac{1}{2}\right), \begin{pmatrix} \alpha & -\beta \\ -\gamma & \delta \end{pmatrix} \right), \text{ where } \alpha_4, \alpha, \beta, \gamma, \delta \in \mathbb{F}, \alpha_4 \neq 0, \alpha\delta - \beta\gamma \neq 0,$$

$$(16) \left(A_{13}, \begin{pmatrix} \alpha & 0 \\ \gamma & \delta \end{pmatrix} \right) \simeq \left(A_{13}, \begin{pmatrix} a^2\alpha & 0 \\ a^3\gamma & a^4\delta \end{pmatrix} \right), \text{ where } \alpha, \gamma, \delta, a \in \mathbb{F}, \alpha\delta \neq 0, a \neq 0,$$

$$(17) \left(A_{13}, \begin{pmatrix} 0 & \beta \\ \gamma & 0 \end{pmatrix} \right) \simeq \left(A_{13}, \begin{pmatrix} 0 & a^3\beta \\ a^3\gamma & 0 \end{pmatrix} \right), \text{ where } \beta, \gamma, \delta, a \in \mathbb{F}, \beta\gamma \neq 0, a \neq 0,$$

$$(18) \left(A_{13}, \begin{pmatrix} \alpha & \beta \\ -\beta & 0 \end{pmatrix} \right) \simeq \left(A_{13}, \begin{pmatrix} a^2\alpha & a^3\beta \\ -a^3\beta & 0 \end{pmatrix} \right), \text{ where } \alpha, \beta, a \in \mathbb{F}, \alpha \neq 0, \beta \neq 0, a \neq 0.$$

Lemma 3.2. *Every pair (\mathbb{A}, σ) , where \mathbb{A} is a two dimensional associative algebra over a field \mathbb{F} ($\text{Char}(\mathbb{F}) = 2$), $\sigma : \mathbb{A} \times \mathbb{A} \rightarrow \mathbb{F}$ is a nondegerate form, is isomorphic to only one of the following pairs*

- (1) $\left(A_{12,2}, \begin{pmatrix} \alpha & 0 \\ \gamma & \delta \end{pmatrix} \right) \simeq \left(A_{12,2}, \begin{pmatrix} a^2\alpha & 0 \\ a^3\gamma & a^4\delta \end{pmatrix} \right)$, where $\alpha, \gamma, \delta, a \in \mathbb{F}$, $\alpha\delta \neq 0, a \neq 0$,
- (2) $\left(A_{12,2}, \begin{pmatrix} 0 & \beta \\ \gamma & 0 \end{pmatrix} \right) \simeq \left(A_{12,2}, \begin{pmatrix} 0 & a^3\beta \\ a^3\gamma & 0 \end{pmatrix} \right)$, where $\beta, \gamma, \delta, a \in \mathbb{F}$, $\beta\gamma \neq 0, a \neq 0$,
- (3) $\left(A_{12,2}, \begin{pmatrix} \alpha & \beta \\ -\beta & 0 \end{pmatrix} \right) \simeq \left(A_{12,2}, \begin{pmatrix} a^2\alpha & a^3\beta \\ -a^3\beta & 0 \end{pmatrix} \right)$, where $\alpha, \beta, a \in \mathbb{F}$, $\beta^2 \neq 0, a \neq 0$,
- (4) $\left(A_{11,2}(\beta_1), \begin{pmatrix} \alpha & 0 \\ 1 & \delta \end{pmatrix} \right)$, where $\alpha, \delta \in \mathbb{F}$, $\alpha\delta \neq 0$,
- (5) $\left(A_{11,2}(\beta_1), \begin{pmatrix} \alpha & 0 \\ 0 & \delta \end{pmatrix} \right) \simeq \left(A_{11,2}(\beta_1), \begin{pmatrix} a^2\alpha & 0 \\ 0 & \delta \end{pmatrix} \right)$, where $\alpha, \delta, a \in \mathbb{F}$, $\alpha\delta \neq 0, a \neq 0$.
- (6) $\left(A_{11,2}(\beta_1), \begin{pmatrix} 0 & 1 \\ \gamma & 0 \end{pmatrix} \right)$, where $\gamma \in \mathbb{F}$, $\gamma \neq 0, \gamma + 1 \neq 0$,
- (7) $\left(A_{11,2}(\beta_1), \begin{pmatrix} \alpha & 1 \\ -1 & 0 \end{pmatrix} \right)$, where $\alpha \in \mathbb{F}$,
- (8) $\left(A_{6,2}(1, 0), \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \right)$, where $\alpha, \beta, \gamma, \delta \in \mathbb{F}$, $\alpha\delta - \beta\gamma \neq 0$,
- (9) $\left(A_{4,2}(1, 0, 0), \begin{pmatrix} \alpha & 0 \\ 1 & \delta \end{pmatrix} \right)$, where $\alpha, \delta \in \mathbb{F}$, $\alpha\delta \neq 0$,
- (10) $\left(A_{4,2}(1, 0, 0), \begin{pmatrix} \alpha & 0 \\ 0 & \delta \end{pmatrix} \right) \simeq \left(A_{4,2}(1, 0, 0), \begin{pmatrix} a^2\alpha & 0 \\ 0 & \delta \end{pmatrix} \right)$, where $\alpha, \delta, a \in \mathbb{F}$, $\alpha\delta \neq 0, a \neq 0$,
- (11) $\left(A_{4,2}(1, 0, 0), \begin{pmatrix} 0 & 1 \\ \gamma & 0 \end{pmatrix} \right)$, where $\gamma \in \mathbb{F}$, $\gamma \neq 0, \gamma + 1 \neq 0$,
- (12) $\left(A_{4,2}(1, 0, 0), \begin{pmatrix} \alpha & 1 \\ -1 & 0 \end{pmatrix} \right)$, where $\alpha \in \mathbb{F}$,
- (13) $\left(A_{4,2}(1, \beta_1, 0), \begin{pmatrix} \alpha & 0 \\ \gamma & \delta \end{pmatrix} \right)$, where $\alpha, \gamma, \delta \in \mathbb{F}$, $\alpha\delta \neq 0$,
- (14) $\left(A_{4,2}(1, \beta_1, 0), \begin{pmatrix} 0 & \beta \\ \gamma & 0 \end{pmatrix} \right)$, where $\beta, \gamma \in \mathbb{F}$, $\beta\gamma \neq 0, \beta + \gamma \neq 0$,
- (15) $\left(A_{4,2}(1, \beta_1, 0), \begin{pmatrix} \alpha & \beta \\ -\beta & 0 \end{pmatrix} \right)$, where $\alpha, \beta \in \mathbb{F}, \beta \neq 0$,
- (16) $\left(A_{3,2}(1, 0, 0), \begin{pmatrix} \alpha & 1 \\ \gamma & \delta \end{pmatrix} \right)$, where $\alpha, \gamma, \delta \in \mathbb{F}$, $\alpha\delta - \gamma \neq 0$,
- (17) $\left(A_{3,2}(1, 0, 0), \begin{pmatrix} \alpha & 0 \\ 1 & \delta \end{pmatrix} \right)$, where $\alpha, \delta \in \mathbb{F}$, $\alpha\delta \neq 0$,
- (18) $\left(A_{3,2}(1, 0, 0), \begin{pmatrix} \alpha & 0 \\ 0 & \delta \end{pmatrix} \right) \simeq \left(A_{3,2}(1, 0, 0), \begin{pmatrix} \alpha & 0 \\ 0 & d^2\delta \end{pmatrix} \right)$, where $\alpha, \delta, d \in \mathbb{F}$, $\alpha\delta \neq 0, d \neq 0$,
- (19) $\left(A_{3,2}(1, 0, 1), \begin{pmatrix} \alpha & 0 \\ 1 & \delta \end{pmatrix} \right)$, where $\alpha, \delta \in \mathbb{F}$, $\alpha\delta \neq 0$,

$$(20) \left(A_{3,2}(1, 0, 1), \begin{pmatrix} \alpha & 0 \\ 0 & \delta \end{pmatrix} \right) \simeq \left(A_{3,2}(1, 0, 1), \begin{pmatrix} \alpha & 0 \\ 0 & d^2\delta \end{pmatrix} \right), \text{ where } \alpha, \delta \in \mathbb{F}, \alpha\delta \neq 0, d \neq 0,$$

$$(21) \left(A_{3,2}(1, 0, 1), \begin{pmatrix} 0 & 1 \\ \gamma & 0 \end{pmatrix} \right), \text{ where } \gamma \in \mathbb{F}, \gamma \neq 0, \gamma + 1 \neq 0,$$

$$(22) \left(A_{3,2}(1, 0, 1), \begin{pmatrix} \alpha & 1 \\ -1 & 0 \end{pmatrix} \right), \text{ where } \alpha \in \mathbb{F}.$$

Lemma 3.3. *Every pair (\mathbb{A}, σ) , where \mathbb{A} is a two dimensional associative algebra over a field \mathbb{F} ($\text{Char}(\mathbb{F}) = 3$), $\sigma : \mathbb{A} \times \mathbb{A} \rightarrow \mathbb{F}$ is a nondegerate form, is isomorphic to only one of the following pairs*

$$(1) \left(A_{13,3}, \begin{pmatrix} \alpha & 0 \\ \gamma & \delta \end{pmatrix} \right) \simeq \left(A_{13,3}, \begin{pmatrix} a^2\alpha & 0 \\ a^3\gamma & a^4\delta \end{pmatrix} \right), \text{ where } \alpha, \gamma, \delta, a \in \mathbb{F}, \alpha\delta \neq 0, a \neq 0,$$

$$(2) \left(A_{13,3}, \begin{pmatrix} 0 & \beta \\ \gamma & 0 \end{pmatrix} \right) \simeq \left(A_{13,3}, \begin{pmatrix} 0 & a^3\beta \\ a^3\gamma & 0 \end{pmatrix} \right), \text{ where } \beta, \gamma, a \in \mathbb{F}, \beta + \gamma \neq 0, \beta\gamma \neq 0, a \neq 0,$$

$$(3) \left(A_{13,3}, \begin{pmatrix} \alpha & \beta \\ -\beta & 0 \end{pmatrix} \right) \simeq \left(A_{13,3}, \begin{pmatrix} a^2\alpha & a^3\beta \\ -a^3\beta & 0 \end{pmatrix} \right), \text{ where } \alpha, \beta, a \in \mathbb{F}, \beta^2 \neq 0, a \neq 0,$$

$$(4) \left(A_{3,3}(1, 0, 1), \begin{pmatrix} \alpha & 0 \\ 1 & \delta \end{pmatrix} \right), \text{ where } \alpha, \delta \in \mathbb{F}, \alpha\delta \neq 0,$$

$$(5) \left(A_{3,3}(1, 0, 1), \begin{pmatrix} \alpha & 0 \\ 0 & \delta \end{pmatrix} \right) \simeq \left(A_{3,3}(1, 0, 1), \begin{pmatrix} \alpha & 0 \\ 0 & d^2\delta \end{pmatrix} \right), \text{ where } \alpha, \delta, d \in \mathbb{F}, \alpha\delta \neq 0, d \neq 0,$$

$$(6) \left(A_{3,3}(1, 0, 1), \begin{pmatrix} 0 & 1 \\ \gamma & 0 \end{pmatrix} \right), \text{ where } \gamma \in \mathbb{F}, \gamma \neq 0, \gamma + 1 \neq 0,$$

$$(7) \left(A_{3,3}(1, 0, 1), \begin{pmatrix} \alpha & 1 \\ -1 & 0 \end{pmatrix} \right), \text{ where } \alpha \in \mathbb{F},$$

$$(8) \left(A_{3,3}(1, 0, 0), \begin{pmatrix} \alpha & 1 \\ \gamma & \delta \end{pmatrix} \right), \text{ where } \alpha, \gamma, \delta \in \mathbb{F}, \alpha\delta - \gamma \neq 0,$$

$$(9) \left(A_{3,3}(1, 0, 0), \begin{pmatrix} \alpha & 0 \\ 1 & \delta \end{pmatrix} \right), \text{ where } \alpha, \delta \in \mathbb{F}, \alpha\delta \neq 0,$$

$$(10) \left(A_{3,3}(1, 0, 0), \begin{pmatrix} \alpha & 0 \\ 0 & \delta \end{pmatrix} \right) \simeq \left(A_{3,3}(1, 0, 0), \begin{pmatrix} \alpha & 0 \\ 0 & d^2\delta \end{pmatrix} \right), \text{ where } \alpha, \delta, d \in \mathbb{F}, \alpha\delta \neq 0, d \neq 0,$$

$$(11) \left(A_{3,3}(2, 0, 0), \begin{pmatrix} \alpha & 0 \\ \gamma & \delta \end{pmatrix} \right), \text{ where } \alpha, \delta, \gamma \in \mathbb{F}, \alpha\delta \neq 0,$$

$$(12) \left(A_{3,3}(2, 0, 0), \begin{pmatrix} \alpha & \beta \\ 0 & \delta \end{pmatrix} \right), \text{ where } \alpha, \beta, \delta \in \mathbb{F}, \alpha\delta \neq 0,$$

$$(13) \left(A_{3,3}(2, 0, 0), \begin{pmatrix} \alpha & -\delta \\ -\delta & \delta \end{pmatrix} \right) \simeq \left(A_{3,3}(2, 0, 0), \begin{pmatrix} \alpha - \delta + d^2\delta & -d^2\delta \\ -d^2\delta & d^2\delta \end{pmatrix} \right), \text{ where } \alpha, \delta, d \in \mathbb{F},$$

$$\delta(\alpha) + \delta \neq 0,$$

$$(14) \left(A_{3,3}(2, 0, 0), \begin{pmatrix} \alpha & 1 \\ \gamma & 0 \end{pmatrix} \right), \text{ where } \alpha, \gamma \in \mathbb{F}, \gamma \neq 0,$$

$$(15) \left(A_{3,3}(2, 0, 2), \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \right), \text{ where } \alpha, \beta, \gamma, \delta \in \mathbb{F}, \alpha\delta - \beta\gamma \neq 0,$$

$$(16) \left(A_{3,3}(2, \alpha_4, 2), \begin{pmatrix} \alpha & 0 \\ \gamma & \delta \end{pmatrix} \right), \text{ where } \alpha, \gamma, \delta \in \mathbb{F}, \alpha\delta \neq 0,$$

$$(17) \left(A_{3,3}(2, \alpha_4, 2), \begin{pmatrix} 0 & \beta \\ \gamma & 0 \end{pmatrix} \right), \text{ where } \beta, \gamma \in \mathbb{F}, \beta\gamma \neq 0, \beta + \gamma \neq 0,$$

$$(18) \left(A_{3,3}(2, \alpha_4, 2), \begin{pmatrix} \alpha & \beta \\ -\beta & 0 \end{pmatrix} \right), \text{ where } \alpha, \beta \in \mathbb{F}, \beta \neq 0.$$

Now we specify those pairs from the lmmas above which are Frobenius algebras.

Theorem 7.

- If $\text{Char}(\mathbb{F}) \neq 2, 3$ then any two-dimensional Frobenius algebra over \mathbb{F} is isomorphic to only one of the following such algebras,
 - * $\left(A_3(1, 0, 0), \begin{pmatrix} \alpha & 0 \\ 0 & \delta \end{pmatrix} \right) \simeq \left(A_3(1, 0, 0), \begin{pmatrix} \alpha & 0 \\ 0 & d^2\delta \end{pmatrix} \right)$, where $\alpha, \delta, d \in \mathbb{F}, d \neq 0, \alpha\delta \neq 0$,
 - * $\left(A_3\left(\frac{1}{2}, 0, \frac{1}{2}\right), \begin{pmatrix} \alpha & 1 \\ 1 & 0 \end{pmatrix} \right)$, where $\alpha \in \mathbb{F}$,
 - * $\left(A_3\left(\frac{1}{2}, \alpha_4, \frac{1}{2}\right), \begin{pmatrix} \alpha & \beta \\ \beta & 2\alpha_4\alpha \end{pmatrix} \right) \simeq \left(A_3\left(\frac{1}{2}, \alpha_4, \frac{1}{2}\right), \begin{pmatrix} \alpha & -\beta \\ -\beta & 2\alpha_4\alpha \end{pmatrix} \right)$, where $\alpha_4, \alpha, \beta \in \mathbb{F}, 2\alpha_4\alpha^2 - \beta^2 \neq 0, \alpha_4 \neq 0$.
 - * $\left(A_{13}, \begin{pmatrix} 0 & \beta \\ \beta & 0 \end{pmatrix} \right) \simeq \left(A_{13}, \begin{pmatrix} 0 & a^3\beta \\ a^3\beta & 0 \end{pmatrix} \right)$, where $\beta, a \in \mathbb{F}, a \neq 0, \beta \neq 0$.
- If $\text{Char}(\mathbb{F}) = 2$ then any two-dimensional Frobenius algebra over \mathbb{F} is isomorphic to only one of the following such algebras,
 - * $\left(A_{3,2}(1, 0, 0), \begin{pmatrix} \alpha & 0 \\ 0 & \delta \end{pmatrix} \right) \simeq \left(A_{3,2}(1, 0, 0), \begin{pmatrix} \alpha & 0 \\ 0 & d^2\delta \end{pmatrix} \right)$, where $\alpha, \delta, d \in \mathbb{F}, d \neq 0, \alpha\delta \neq 0$,
 - * $\left(A_{4,2}(1, 0, 0), \begin{pmatrix} 1 & 1 \\ -1 & 0 \end{pmatrix} \right)$,
 - * $\left(A_{4,2}(1, \beta_1, 0), \begin{pmatrix} \beta_1\delta & 0 \\ 0 & \delta \end{pmatrix} \right)$, where $\beta_1, \delta \in \mathbb{F}, \beta_1\delta \neq 0$,
 - * $\left(A_{4,2}(1, \beta_1, 0), \begin{pmatrix} \beta & \beta \\ -\beta & 0 \end{pmatrix} \right)$, where $\beta_1, \beta \in \mathbb{F}, \beta \neq 0$,
 - * $\left(A_{11,2}(\beta_1), \begin{pmatrix} \beta_1\delta & 0 \\ 0 & \delta \end{pmatrix} \right) \simeq \left(A_{11,2}(\beta_1), \begin{pmatrix} a^2\beta_1\delta & 0 \\ 0 & \delta \end{pmatrix} \right)$, where $\beta_1, \delta, a \in \mathbb{F}, a \neq 0, \beta_1\delta \neq 0$,
 - * $\left(A_{11,2}(\beta_1), \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \right)$, where $\beta_1 \in \mathbb{F}$,
 - * $\left(A_{12,2}, \begin{pmatrix} 0 & \beta \\ \beta & 0 \end{pmatrix} \right) \simeq \left(A_{12,2}, \begin{pmatrix} 0 & a^3\beta \\ a^3\beta & 0 \end{pmatrix} \right)$, where $\beta, a \in \mathbb{F}, a \neq 0, \beta \neq 0$.
- If $\text{Char}(\mathbb{F}) = 3$ then any two-dimensional Frobenius algebra over \mathbb{F} is isomorphic to only one of the following such algebras,
 - * $\left(A_{3,3}(1, 0, 0), \begin{pmatrix} \alpha & 0 \\ 0 & \delta \end{pmatrix} \right) \simeq \left(A_{3,3}(1, 0, 0), \begin{pmatrix} \alpha & 0 \\ 0 & d^2\delta \end{pmatrix} \right)$, where $\alpha, \delta, d \in \mathbb{F}, d \neq 0, \alpha\delta \neq 0$,
 - * $\left(A_{3,3}(2, 0, 2), \begin{pmatrix} \alpha & \beta \\ \beta & 0 \end{pmatrix} \right)$, where $\alpha, \beta \in \mathbb{F}, \beta \neq 0$,

- * $\left(A_{3,3}(2, \alpha_4, 2), \begin{pmatrix} \alpha & 0 \\ 0 & 2\alpha_4\alpha \end{pmatrix} \right)$, where $\alpha_4, \alpha \in \mathbb{F}, \alpha_4\alpha \neq 0$,
- * $\left(A_{3,3}(2, \alpha_4, 2), \begin{pmatrix} 0 & \beta \\ \beta & 0 \end{pmatrix} \right)$, where $\alpha_4, \beta \in \mathbb{F}, \alpha_4 \neq 0, \beta \neq 0$,
- * $\left(A_{13,3}, \begin{pmatrix} 0 & \beta \\ \beta & 0 \end{pmatrix} \right) \simeq \left(A_{13,3}, \begin{pmatrix} 0 & a^3\beta \\ a^3\beta & 0 \end{pmatrix} \right)$, where $\beta, a \in \mathbb{F}, a \neq 0, \beta \neq 0$.

Proof. Let now check the condition $\sigma(x \cdot y, z) = \sigma(x, y \cdot z)$ for the pairs appeared in Lemma 3.1 ($Char\mathbb{F} \neq 2, 3$), Lemma 3.2 ($Char\mathbb{F} = 2$) and Lemma 3.3 ($Char\mathbb{F} = 3$), i.e., find the solutions to the system of equations (2.2).

Let $Char\mathbb{F} \neq 2, 3$.

In this case the system of equations (2.2) is consistent only for pairs (3), (12), (15) and (17). The solutions to the system are as follows:

For (3) $\left(A_3(1, 0, 0), \begin{pmatrix} \alpha & 0 \\ 0 & \delta \end{pmatrix} \right) \simeq \left(A_3(1, 0, 0), \begin{pmatrix} \alpha & 0 \\ 0 & d^2\delta \end{pmatrix} \right)$, where $\alpha, \delta, d \in \mathbb{F}, \alpha\delta \neq 0, d \neq 0$ the system becomes an identity, therefore, the pair (3) is a Frobenius algebra.

Consider (12) $\left(A_3\left(\frac{1}{2}, 0, \frac{1}{2}\right), \begin{pmatrix} \alpha & 1 \\ \gamma & \delta \end{pmatrix} \right)$, where $\alpha, \gamma, \delta \in \mathbb{F}, \alpha\delta - \gamma \neq 0$. As a solution the system (2.2) we get $\gamma = 1$ and $\delta = 0$ and the corresponding Frobenius algebras are

$$\left(A_3\left(\frac{1}{2}, 0, \frac{1}{2}\right), \begin{pmatrix} \alpha & 1 \\ 1 & 0 \end{pmatrix} \right), \text{ where } \alpha \in \mathbb{F}.$$

Among the pairs (15) $\left(A_3\left(\frac{1}{2}, \alpha_4, \frac{1}{2}\right), \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \right) \simeq \left(A_3\left(\frac{1}{2}, \alpha_4, \frac{1}{2}\right), \begin{pmatrix} \alpha & -\beta \\ -\gamma & \delta \end{pmatrix} \right)$, where $\alpha, \beta, \gamma, \delta, \alpha_4 \in \mathbb{F}, \alpha\delta - \beta\gamma \neq 0, \alpha_4 \neq 0$ the algebras satisfying $\gamma = \beta$, and $\delta = 2\alpha_4\alpha$ are Frobenius. Thus, we get

$$\left(A_3\left(\frac{1}{2}, \alpha_4, \frac{1}{2}\right), \begin{pmatrix} \alpha & \beta \\ \beta & 2\alpha_4\alpha \end{pmatrix} \right) \simeq \left(A_3\left(\frac{1}{2}, \alpha_4, \frac{1}{2}\right), \begin{pmatrix} \alpha & -\beta \\ -\beta & 2\alpha_4\alpha \end{pmatrix} \right),$$

where $\alpha, \beta, \alpha_4 \in \mathbb{F}, 2\alpha_4\alpha^2 - \beta^2 \neq 0, \alpha_4 \neq 0$.

Considering the system of equations (2.2) for (17) $\left(A_{13}, \begin{pmatrix} 0 & \beta \\ \gamma & 0 \end{pmatrix} \right) \simeq \left(A_{13}, \begin{pmatrix} 0 & a^3\beta \\ a^3\gamma & 0 \end{pmatrix} \right)$, where $a \neq 0$ we obtain $\gamma = \beta$ and hence,

$$\left(A_{13}, \begin{pmatrix} 0 & \beta \\ \beta & 0 \end{pmatrix} \right) \simeq \left(A_{13}, \begin{pmatrix} 0 & a^3\beta \\ a^3\beta & 0 \end{pmatrix} \right) \text{ where } \beta, a \in \mathbb{F} \ a \neq 0, \text{ and } \beta^2 \neq 0$$

is a Frobenius algebra.

Let now treat the case of $Char\mathbb{F} = 2$.

From the pairs (2) $\left(A_{12,2}, \begin{pmatrix} 0 & \beta \\ \gamma & 0 \end{pmatrix} \right) \simeq \left(A_{12,2}, \begin{pmatrix} 0 & a^3\beta \\ a^3\gamma & 0 \end{pmatrix} \right)$, where $\beta, \gamma, \delta, a \in \mathbb{F}, \beta\gamma \neq 0, a \neq 0$ here we generate the following Frobenius algebras $\left(A_{12,2}, \begin{pmatrix} 0 & \beta \\ \beta & 0 \end{pmatrix} \right) \simeq \left(A_{12,2}, \begin{pmatrix} 0 & a^3\beta \\ a^3\beta & 0 \end{pmatrix} \right)$, where $\beta, a \in \mathbb{F}, a \neq 0, \beta^2 \neq 0$ since the solution to the system of equations (2.2) in this case is $\gamma = \beta = 0$.

From (5) $\left(A_{11,2}(\beta_1), \begin{pmatrix} \alpha & 0 \\ 0 & \delta \end{pmatrix} \right) \simeq \left(A_{11,2}(\beta_1), \begin{pmatrix} a^2\alpha & 0 \\ 0 & \delta \end{pmatrix} \right)$, where $\alpha, \delta, a \in \mathbb{F}, \alpha\delta \neq 0, a \neq 0$ subjecting to the system (2.2) we get $\alpha = \beta_1\delta$ and obtain the following Frobenius algebra

$$\left(A_{11,2}(\beta_1), \begin{pmatrix} \beta_1\delta & 0 \\ 0 & \delta \end{pmatrix} \right) \simeq \left(A_{11,2}(\beta_1), \begin{pmatrix} a^2\beta_1\delta & 0 \\ 0 & \delta \end{pmatrix} \right), \text{ where } \delta, a \in \mathbb{F}, \beta_1\delta^2 \neq 0, a \neq 0.$$

For (7) $\left(A_{11,2}(\beta_1), \begin{pmatrix} \alpha & 1 \\ -1 & 0 \end{pmatrix}\right)$, where $\alpha \in \mathbb{F}$,

the solution to the system of equations (2.2) is $\alpha = 0$. Therefore, $\left(A_{11,2}(\beta_1), \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}\right)$ is a Frobenius algebra.

Considering (12) $\left(A_{4,2}(1,0,0), \begin{pmatrix} \alpha & 1 \\ -1 & 0 \end{pmatrix}\right)$, where $\alpha \in \mathbb{F}$, we obtain $\alpha = 1$, i.e.,

$$\left(A_{4,2}(1,0,0), \begin{pmatrix} 1 & 1 \\ -1 & 0 \end{pmatrix}\right)$$

is a Frobenius algebra here.

Let us consider (13) $\left(A_{4,2}(1,\beta_1,0), \begin{pmatrix} \alpha & 0 \\ \gamma & \delta \end{pmatrix}\right)$, where $\alpha\delta \neq 0$. Then the system of equations (2.2) is equivalent to

$$\begin{cases} \gamma = 0 \\ \alpha - \beta_1\delta = 0 \end{cases}$$

Therefore, $\left(A_{4,2}(1,\beta_1,0), \begin{pmatrix} \beta_1\delta & 0 \\ 0 & \delta \end{pmatrix}\right)$, where $\beta_1, \delta \in \mathbb{F}, \beta_1\delta^2 \neq 0$ are Frobenius algebras.

From (15) $\left(A_{4,2}(1,\beta_1,0), \begin{pmatrix} \alpha & \beta \\ -\beta & 0 \end{pmatrix}\right)$, where $\alpha, \beta \in \mathbb{F}$ and $\beta \neq 0$ as a solution to the system (2.2) we obtain $\beta = \alpha$. Hence,

$$\left(A_{4,2}(1,\beta_1,0), \begin{pmatrix} \beta & \beta \\ -\beta & 0 \end{pmatrix}\right), \text{ where } \beta \in \mathbb{F} \text{ and } \beta \neq 0.$$

are Frobenius algebras.

For (18) $\left(A_{3,2}(1,0,0), \begin{pmatrix} \alpha & 0 \\ 0 & \delta \end{pmatrix}\right) \simeq \left(A_{3,2}(1,0,0), \begin{pmatrix} \alpha & 0 \\ 0 & d^2\delta \end{pmatrix}\right)$, where $\alpha, \delta, d \in \mathbb{F}, d \neq 0, \alpha\delta \neq 0$ the system of equations (2.2) is an identity and they are Frobenius algebras.

Note that for all the other classes from Lemma 3.2 the system of equations (2.2) is inconsistent.

Let $\text{Char}\mathbb{F} = 3$.

We verify the list of pairs from Lemma 3.3 to be Frobenius algebras. From (2) $\left(A_{13,3}, \begin{pmatrix} 0 & \beta \\ \gamma & 0 \end{pmatrix}\right) \simeq \left(\begin{pmatrix} 0 & a^3\beta \\ a^3\gamma & 0 \end{pmatrix}\right)$, where $\beta\gamma \neq 0$ as a solution to the system (2.2) we get $\gamma = \beta$ that means $\left(A_{13,3}, \begin{pmatrix} 0 & \beta \\ \beta & 0 \end{pmatrix}\right) \simeq \left(\begin{pmatrix} 0 & a^3\beta \\ a^3\beta & 0 \end{pmatrix}\right)$, where $\beta, a \in \mathbb{F}, a \neq 0, \beta^2 \neq 0$ are Frobenius algebras.

All the pairs (10) $\left(A_{3,3}(1,0,0), \begin{pmatrix} \alpha & 0 \\ 0 & \delta \end{pmatrix}\right) \simeq \left(A_{3,3}(1,0,0), \begin{pmatrix} \alpha & 0 \\ 0 & d^2\delta \end{pmatrix}\right)$, where $0 \neq d \in \mathbb{F}, \alpha\delta \neq 0$ are Frobenius algebras.

It is easy to see that among the pairs (15) $\left(A_{3,3}(2,0,2), \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix}\right)$, where $\alpha\delta - \beta\gamma \neq 0$ Frobenius algebras are

$$\left(A_{3,3}(2,0,2), \begin{pmatrix} \alpha & \beta \\ \beta & 0 \end{pmatrix}\right), \text{ where } \alpha, \beta \in \mathbb{F}, \beta^2 \neq 0$$

From the pairs (16) $\left(A_{3,3}(2, \alpha_4, 2), \begin{pmatrix} \alpha & 0 \\ \gamma & \delta \end{pmatrix} \right)$, where $\alpha\delta \neq 0$, we generate the following Frobenius algebras

$$\left(A_{3,3}(2, \alpha_4, 2), \begin{pmatrix} \alpha & 0 \\ 0 & 2\alpha_4\alpha \end{pmatrix} \right), \text{ where } \alpha_4, \alpha \in \mathbb{F}, 2\alpha_4\alpha^2 \neq 0.$$

Finally, from (17) $\left(A_{3,3}(2, \alpha_4, 2), \begin{pmatrix} 0 & \beta \\ \gamma & 0 \end{pmatrix} \right)$, where $\beta\gamma \neq 0$, $\beta + \gamma \neq 0$, we obtain the following Frobenius algebras

$$\left(A_{3,3}(2, \alpha_4, 2), \begin{pmatrix} 0 & \beta \\ \beta & 0 \end{pmatrix} \right), \text{ where } \beta \in \mathbb{F}, \beta \neq 0.$$

For all the other pairs the system (2.2) is inconsistent. □

REFERENCES

- [1] D. BAR-NATAN, Khovanov's homology for tangles and cobordisms, *Geom. Topol.*, 9 (3) (2005), 1443–1499, arXiv:math/0410495, doi:10.2140/gt.2005.9.1443.
- [2] U. BEKBAEV, Classification of two-dimensional algebras over any basic field, 2023, *J. Phys.: Conf. Ser.*
- [3] R. BRAUER, C. NESBITT, On the regular representations of algebras, *Proc. Natl. Acad. Sci. USA*, 23 (4)(1937), 236–240, doi:10.1073/pnas.23.4.236.
- [4] J. DIEUDONNÉ, Remarks on quasi-Frobenius rings, *Illinois Journal of Mathematics*, 2(3)(1958), 346–354, doi:10.1215/ijm/1255454538.
- [5] T. NAKAYAMA, On Frobeniusean algebras I, *Ann. of Math.*, 40 (1939), 611–633,
- [6] T. NAKAYAMA, On Frobeniusean algebras II, *Ann. of Math.*, 42 (1941), 1–21.
- [7] T. NAKAYAMA, On the complete cohomology theory of Frobenius algebras, *Osaka Math. J.*, 9 (1957), 165–187.
- [8] T. NAKAYAMA, C. NESBITT, Note on symmetric algebras, *Ann. of Math.*, 39 (1938), 659–668.
- [9] I.S.RAKHIMOV, *Complete classification of two-dimensional associative and diassociative algebras over any basic field*, arXiv:2307.09927 v1 [math.RA], 19 July, 2023
- [10] P. TURNER, *Five Lectures on Khovanov Homology*, arXiv:math/0606464, (2006).