THE CENTRE OF A FINITELY GENERATED STRONGLY VERBALLY CLOSED GROUP IS ALMOST ALWAYS PURE

Filipp D. Denissov Anton A. Klyachko Faculty of Mechanics and Mathematics of Moscow State University Moscow 119991, Leninskie gory, MSU. Moscow Center for Fundamental and Applied Mathematics klyachko@mech.math.msu.su denissov.filipp@gmail.com

The assertion in the title implies that many interesting groups (e.g., all non-abelian braid groups or $\mathbf{SL}_{100}(\mathbb{Z}))$ are not strongly verbally closed, i.e., they embed into some finitely generated groups as verbally closed subgroups, which are not retracts.

1. Introduction

A subgroup H of a group G is called *verbally closed* [MR14] if any equation of the form

 $w(x, y, ...) = h$, where w is an element of a free group $F(x, y, ...)$ and $h \in H$,

admits a solution in G if and only if it admits a solution in H. Similarly, H is called *algebraically closed* in G if each finite system of equations with coefficients from H

 $\{w_1(x, y, \ldots) = 1, \ldots, w_m(x, y, \ldots) = 1\},\$ where $w_i \in H * F(x, y, \ldots)$ (and $*$ means the free product)

admits a solution in G if and only if it admits a solution in H .

The algebraic closedness is a stronger property than the verbal closedness, but, often, these properties turn out to be equivalent (see [Rom12], [RKh13], [MR14], [Mazh17], [RKhK17], [KM18], [KMM18], [Mazh18], [Bog22], [Mazh19], $[RT20]$, $[Tim21]$, $[KMO23]$, $[KK24]$). A group H is called *strongly verbally closed* [Mazh18] if it is algebraically closed in any group containing H as a verbally closed subgroup. Thus, the verbal closedness (as well as algebraic closedness) is a property of a subgroup, while the strong verbal closedness is a property of an abstract group. The class of strongly verbally closed groups is fairly wide. For example, the following groups are strongly verbally closed:

– all abelian groups [Mazh18],

- all free groups [KM18],
	- and even all virtually free groups containing no nonidentity finite normal subgroups [KM18], [KMM18],
- all free products of nontrivial groups [Mazh19],
- the fundamental group of any connected surface, except for the Klein bottle [Mazh18], [K21],
	- and even all acylindrically hyperbolic groups without nonidentity finite normal subgroups [Bog23] (this generalises several results above),
- all finite groups with nonabelian monoliths (in particular, all finite simple groups) [KMO23],
- all finite symmetric groups [KK24],
- all dihedral groups of order not divisible by eight [KMO23].

There is also a general embedding theorem [KMO23]:

any group H embeds into a strongly verbally closed group of cardinality $|H| + \aleph_0$ that satisfies all identities of H.

Actually, neither proving nor disproving the strong verbal closedness of a given group is an easy task.

Until quite lately, only few examples of non-strongly-verbally-closed groups were known. However, the following theorem has recently been proven.

Theorem KMO [KMO23] . *The centre of a finite strongly verbally closed group is its direct factor.*

This theorem implies immediately that, e.g., all finite nilpotent non-abelian groups are not strongly verbally closed. The same is true for finitely generated nilpotent groups with non-abelian torsion subgroups [D23]. However, the following question from [KMO23] remains open:

does there exist a finitely generated non-abelian nilpotent strongly verbally closed group?

(The embedding theorem [KMO23], see above, shows that the finitely-generated condition cannot be dropped.)

The simplest infinite finitely generated non-abelian nilpotent group (with abelian torsion subgroup) is the discrete Heisenberg group $UT_3(\mathbb{Z})$, which turned out to be non-strongly-verbally-closed [KMO23], but the proof of this fact is not easy.

Let us ask a more daring question.

The work of the second author was supported by the Russian Science Foundation, project no. 22-11-00075.

Question. *Is it true that the centre of any finitely generated strongly verbally closed group is its direct factor?*

For now, we are able to prove only a weaker property — the purity of the centre, and under some additional conditions (not very onerous though). Recall that a subgroup B of a group A is called *pure* (or *servant*) if, for every integer n and for every $b \in B$, the element b is an nth power in A if and only if it is an nth power in B.

Main theorem (a simplified form). *The centre of a finitely generated strongly verbally closed group is pure if at least one of the following conditions holds*:

- *all finite-order elements lie in the centre* (in this case, the purity of the centre is equivalent to its isolatedness)*,*
- *or the group is not infinitely decomposable.*

(The full statement can be found in the next section.) Here, we call a group *infinitely decomposable* if, for any $n \in \mathbb{N}$, it decomposes into a product of n pairwise commuting noncentral subgroups (which are normal in this case of course).

Finitely generated infinitely decomposable groups are exotic creatures, but they do exist; there exists even a nontrivial finitely generated group isomorphic to its Cartesian square [TJ74].

The central quotient of an infinitely decomposable group decomposes into a direct product of arbitrarily many nontrivial subgroups; this means that it does not satisfy the maximality condition for direct factors by virtue of the following theorem ([O95], Corollary $3 +$ Proposition 1):

a group with finitely generated commutator quotient group decomposes into a finite direct product of indecomposable groups if and only if it satisfies the maximality condition for direct factors; and the number of such decompositions (*up to isomorphisms of the factors*) *is always finite.*

Thus, we obtain the following fact.

Corollary 1. *The centre of a finitely generated strongly verbally closed group is pure if its central quotient satisfies the maximality condition for direct factors* (*or, equivalently, decomposes into a finite direct product of indecomposable groups*)*. In particular, the centre of any finitely generated solvable strongly verbally closed group is pure.*

This corollary of course contains the mentioned above fact that the Klein-bottle group is not strongly verbally closed [K21] (because the Klein-bottle group $K = \langle a, b | a^2 = b^2 \rangle$ is solvable, and its centre $\langle a^2 \rangle$ is not pure: a^2 is a square in K but not in $\langle a^2 \rangle$). Moreover, our theorem gives many new interesting examples of non-strongly-verballyclosed groups, e.g.,

the following groups are not strongly verbally closed:

– all non-abelian braid groups (because these groups are torsion-free, and the centre is generated by the square of the "fundamental" braid [G69])*, and, in particular, it is not 2-pure, i.e., not pure with respect to squares,* $-$ **SL**_n(\mathbb{Z}) for even n; indeed, for even n the group $SL_n(\mathbb{Z})$ is not pure (precisely not 2-pure), because its centre is generated by the square of the block-diagonal matrix with blocks $\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$, and the group $PSL_n(\mathbb{Z})$ is

indecomposable into a direct product, as it is easy to show*.* ∗)

The following corollary (which we prove in the last section) is a direct generalisation of Theorem KMO.

Corollary 2. *If the commutator subgroup of a finitely generated strongly verbally closed group is finite* (*or, equivalently, the index of the centre is finite***))*, then the centre is a direct factor of the group.*

Our approach to the proof of the main theorem may be called similar to the proof from [K21], but our argument is significantly trickier: in particular, we use Chevalley's theorem on zeros of polynomials [Che35] (to be more precise, we use its generalisation, see Section 3); in this aspect, our argument is similar to the proof of Theorem KMO [KMO23]. The notation being used in the paper is mainly standard. Note only that, if $k \in \mathbb{Z}$, and x and y are elements of a group, then x^y , x^{ky} , and x^{-y} denote $y^{-1}xy$, $y^{-1}x^ky$, and $y^{-1}x^{-1}y$, respectively. The commutator $[x, y]$ is $x^{-1}y^{-1}xy$. The centre of a group G is denoted by $Z(G)$. The cardinality of a set X is denoted by |X|. If X is a subset of a group, then $\langle X \rangle$ denotes the subgroup generated by X. The symbol $\langle g \rangle_n$ denotes the cyclic subgroup of order n generated by an element g. The letters $\mathbb Z$ and $\mathbb N$ stand for the sets of integers and positive integers, respectively. $\mathbb Z_n$ is the residues ring modulo n (or the additive group of this ring). The free product of groups A and B is denoted by $A * B$.

^{*)} The braid groups are linear [Big01], [Kra02] (as well as $SL_n(\mathbb{Z})$); for linear (and even for all equationally Noetherian) subgroups H of finitely generated groups G , the algebraic closedness is equivalent to the existence of a retraction (i.e., a homomorphism $G \to H$ that is identical on H) [MR14]; hence, these examples can be stated in a "more categorical" language (see the abstract).

^{**}) *any group whose centre is of finite index has finite commutator subgroup* by Schur's theorem [Sch04]. The converse for finitely generated groups was proven by B. Neumann [Neu51] (but the history of this question is somewhat odd [Ya16]).

The authors thank an anonymous referee and A. A. Skutin for useful comments allowing us to improve the readability of the paper. The second author thanks the Theoretical Physics and Mathematics Advancement Foundation "BASIS".

2. Main theorem

First note that, to prove the main result as it is stated in the introduction, we may restrict ourselves to prime powers.

An abelian subgroup B of a group A is pure (=servant) if and only if it is p-pure (= p-servant) for each prime p, i.e.,
$$
B \cap \{a^{p^k} \mid a \in A\} = \{b^{p^k} \mid b \in B\}
$$
 for all positive integers k.

Indeed, if $a^{mn} = b = b_1^m = b_2^n$, where $m, n \in \mathbb{Z}$ are coprime, $a \in A$, and $b, b_1, b_2 \in B$, then $(b_1^k b_2^l)^{mn} = b^{kn+lm} = b$, where integers k and l are chosen such that $kn + lm = 1$.

 M ain theorem. If a strongly verbally closed group H is finitely generated modulo its centre and a central element b^{p^k} *is not the* p^k th power of any central element (where $b \in H$, the integer p is prime, and $k \in \mathbb{N}$), then, for any positive *integer* n*, the group* H *decomposes into a product of* n *pairwise commuting subgroups not contained in the centraliser of the set* $\{h \in H \mid h^{p^k} = 1, [h, b] = 1\}$. (In particular, this set is not contained in the centre of *H*.)

Proof. This theorem follows immediately from the following three lemmata, which we prove in the next section.

Verbal-closedness lemma. Suppose that H is a group, $b \in H$, and $b^{p^k} \in Z(H)$, where p is prime and $k \in \mathbb{N}$. Then, *in the semidirect product* $G = F \times \bigtimes_{s \in S} H_s$, where

- *–* S *is a finite set,*
- *– F* is a subgroup in the group of functions $S \to \mathbb{Z}_{p^k}$ (but *F* is written multiplicatively),
- H_s are isomorphic copies of H (and the corresponding isomorphisms are denoted by the subscripts: $h \mapsto h_s$),
- *–* and the action is the following: $h_s^f = \left(h^{b^{f(s)}}\right)$ s ,

the "diagonal" subgroup $\left\{ \Pi \right\}$ $\prod_{s\in S}h_s\ \Big|\ h\in H$ \mathcal{L} \simeq *H* is verbally closed, if each function $f \in F$ vanishes at some point $s \in S$.

Algebraic-non-closedness lemma. *Suppose that a group* H *is finitely generated modulo its centre,* $b \in H$ *, and an element* b^{p^k} , where p is prime and $k \in \mathbb{N}$, lies in the centre of H but is not the p^k th power of any central element. Let *n* be the maximal number of pairwise commuting subgroups whose product is H and none of which centralises the set $\{h \in H \mid h^{p^k} = 1, [h, b] = 1\}$. Then the diagonal subgroup H of the semidirect product G from the verbal*closedness lemma is not algebraically closed in* G *if, for any subset* $T \subseteq S$ *of cardinality at most* max $(n, 1)$ *, there exists a* function $f_T \in F$ *such that* $f_T(s) = 1$ for all $s \in T$ *.*

To complete the proof of the theorem (modulo these lemmata), it remains to show that the conditions of two lemmata above can be satisfied simultaneously (for suitable S and F); the following assertion shows that they can.

Function lemma. *For any positive integers* k *and* n *and any prime* p*, there exists a finite set* S *and a subgroup* F *in the group of functions* $S \to \mathbb{Z}_{p^k}$ *such that*

- 1) *each function from* F *vanishes at some point of* S*;*
- 2) *but, for any* $s_1, \ldots, s_n \in S$, there exists a function $f \in F$ such that $f(s_1) = \ldots = f(s_n) = 1$.

In the next section, we prove these three lemmata, and thereby complete the proof of the theorem.

3. Proofs of the three lemmata

Proof of the verbal-closedness lemma. To show that H is verbally closed in G , we have to prove that any equation of the form

 $w(x, y, \ldots) = h$, where $h \in H$ and $w(x, y, \ldots)$ is an element of a free group $F(x, y, \ldots)$,

which is solvable in G , is solvable in H . A change of variables reduces such an equation to the form

 $x^m u(x, y, \ldots) = h$, where $h \in H$ and $u(x, y, \ldots)$ lies in the commutator subgroup of a free group $F(x, y, \ldots)$.

(To obtain this change of variables, note that the free group modulo its commutator subgroup is the free abelian group $\mathbb{Z} \oplus \ldots \oplus \mathbb{Z}$; and, for any element \widehat{w} of a free abelian group, there is a basis B such that $\widehat{w} = mb$, where $m \in \mathbb{Z}$ and $b \in B$.)

Suppose that this equation has a solution $(\tilde{x}, \tilde{y}, \ldots)$ in G; in particular, $\tilde{x} = f^{\varepsilon}d$, where $\varepsilon \in \mathbb{Z}$, $f \in F$, and $d \in \bigtimes_{s \in S} H_s$. By the condition, the function f vanishes at a point $s \in S$. Consider the homomorphism (taking the sth coordinate):

$$
\varphi: G \to \widetilde{H} = \langle \beta \rangle_{p^k} \angle H \text{ (with the action } h^\beta = h^b \text{), where } \varphi(h_s) = h, \, \varphi(h_{s'}) = 1 \text{ for } s' \neq s, \, \varphi(f') = \beta^{f'(s)} \text{ for } f' \in F.
$$

Applying this homomorphism, we obtain that the equation $w(\varphi(\tilde{x}), y, z, \ldots) = h$ over H with unknowns y, z, \ldots (whose exponent sums are zero in the word w) has a solution in the group H. (This is an equation over H, because $f(s) = 0$ and, hence, $\varphi(\widetilde{x}) \in H$.) It remains to apply the following almost obvious fact:

> *if the exponent sum of each variable in a word* $v \in H * F(y, z, \ldots)$ *is zero, and the equation* $v = 1$ *is solvable in the group* \hat{H} *, then it is solvable in* H *too.*

Indeed, replacing β with b in a solution, we obtain a solution from H. This completes the proof.

Proof of the algebraic-non-closedness lemma. Suppose that $H = \langle h_1, \ldots, h_m \rangle \cdot Z(H)$. The system of equations

$$
\left\{ x_{f}^{p^{k}} \stackrel{\text{(1)}}{=} 1, y_{i\,s}^{x_{f}} \stackrel{\text{(2)}}{=} y_{i\,s}^{b^{f(s)}}, \prod_{q \in S} y_{i\,q} \stackrel{\text{(3)}}{=} h_{i}, [y_{i\,s}, y_{j\,s'}] \stackrel{\text{(4)}}{=} 1, [x_{f}, b] \stackrel{\text{(5)}}{=} 1 \middle| f \in F, s \in S \ni s' \neq s, i, j = 1, \ldots, m \right\}
$$

over H has an obvious solution in G $(x_f = f, y_{is} = (h_i)_s)$ but no solutions in H. Indeed,

- equations (4) mean that the subgroups $K_s = \langle y_{1s}, \ldots, y_{m s} \rangle \cdot \mathcal{Z}(H)$ pairwise commute;
- equations (3) say that $H = \prod K_s$;
- the elements x_f commute with b by virtue of equation (5);
- all the aforesaid observations and equation (1) imply (by the condition of the lemma) that the elements x_f commute with all subgroups K_s , except for some n (or less) subgroups K_t , where $t \in T \subseteq S$, $|T| \leq n$ (this set T is independent of f);
- thus, (by the condition of the lemma) there exists a function $f_T \in F$ such that $f_T(t) = 1$ for all $t \in T$;
- − hence, equations (2) show that the element $x_{f_T}b^{-1}$ commutes with $y_{i,s}$ for $s \in T$;
	- − but, for $s \notin T$, the elements $x_{fr}b^{-1}$ and y_{is} commute too, because, for such s, the element y_{is} commutes with all x_f (as mentioned above), therefore, choosing a function $f \in F$ such that $f(s) = 1$ (which exists by the condition), we obtain $y_{i s} = y_{i s}^{x_f} \stackrel{\text{(2)}}{=} y_{i s}^{b^{f(s)}} = y_{i s}^{b};$
- − thus, equations (3) show that the element $x_{f_T}b^{-1}$ is central (because h_i generate H modulo $\mathcal{Z}(H)$);
- raising this central element to the power p^k , we obtain b^{-p^k} (because $(x_{f_T}b^{-1})^{p^k} \stackrel{(5)}{=} x_{f_T}^{p^k}$ $p_{f_T}^{\kappa}b^{-p^k} \stackrel{(1)}{=} b^{-p^k}$) that contradicts the condition $b^{p^k} \notin Z(H)^{p^k}$. This completes the proof of the lemma.

Proof of the function lemma. Put $S = \mathbb{Z}_{p^k}^m \setminus p\mathbb{Z}_{p^k}^m$, where m is sufficiently large (with respect to k and p, see below). Take as F the set of (polynomial) mappings $\mathbb{Z}_{p^k}^m \to \mathbb{Z}_{p^k}$ defined by polynomials (in m variables) without free terms and of degree at most $n(p-1)p^{k-1}$.

Condition 2) holds. Indeed,

- we can choose a basis in the free \mathbb{Z}_{p^k} -module $\mathbb{Z}_{p^k}^m$ and assume that all coordinates of vectors s_i vanish, except for the first n coordinates (because the definition of F does not depend on the choice of a basis);
- consider the reduction modulo $p \varphi : \mathbb{Z}_{p^k}^m \to \mathbb{Z}_{p^k}^m / p\mathbb{Z}_{p^k}^m \simeq \mathbb{Z}_p^m$;
- each function from an *n*-dimensional vector space (containing all $\varphi(s_i)$) to \mathbb{Z}_p is represented by a polynomial whose degree with respect to each variable is at most $p-1$ (because the polynomials x^p and x define the same function $\mathbb{Z}_p \to \mathbb{Z}_p$);
- i.e., we have a polynomial $g \in \mathbb{Z}[x_1,\ldots,x_n]$ of degree at most $n(p-1)$ such that $g(0) \in p\mathbb{Z}_{p^k}$ and $g(s_i) \in 1+p\mathbb{Z}_{p^k}$ for all i ;
- hence, the polynomial $g^{p^{k-1}}$ takes (in \mathbb{Z}_{p^k})
	- value 0 at zero, because $p^{k-1} > k$ (so this polynomial has no free term);
	- value 1 at all points s_i (because $g(s_i)$ lie in $1+p\mathbb{Z}_{p^k}$, and this set is a (multiplicative) group of order p^{k-1} , so we raise an element of a group to the power equal to the order of the group);

- thus, deg $g^{p^{k-1}} = p^{k-1} \cdot \deg g \leqslant n(p-1)p^{k-1}$, and $g^{p^{k-1}} \in F$ by the definition of F.

To verify Condition 1), we apply a theorem of Schanuel [Sch74], which says (in particular) that:

a polynomial $f \in \mathbb{Z}_{p^k}[x_1,\ldots,x_m]$ without free term has a nonzero root in $\{0,1\}^m$ if $m > (p^k-1) \cdot \deg f$.

Thus, if we choose m larger than $n(p-1)p^{k-1}(p^k-1)$, then the both conditions are fulfilled. This completes the proof of the lemma (and of the main theorem).

4. Proof of Corollary 2

This result follows easily from the main theorem and some known facts. Indeed,

- the group H is finitely generated, and the index of the centre is finite, therefore, the centre is a finitely generated abelian group and, hence, decomposes into a direct product of a finite abelian group and a free abelian group F (and the index $|H : F|$ is finite);
- as was shown by Schur [Sch04] (see also [Rob96], Theorem 10.1.3),

for any central finite-index subgroup F *of* (*any*) *group* H*, the mapping* $\tau: x \mapsto x^{|H:F|}$ (called the transfer) is a homomorphism from H to F.

- $-\tau(H) = \tau(F)$, because the centre is pure in H (by the main theorem), and F is pure in the centre (obviously);
- therefore, the composition of τ and a (suitable) isomorphism $\tau(H) = \tau(F) \to F$ is a retraction $H \to F$;
- i.e., F (being a central subgroup) is a direct factor: $H = F \times H_1$;
- the second factor H_1 must be finite (because F is of finite index in H) and strongly verbally closed:
	- *any direct factor of a strongly verbally closed group is strongly verbally closed**) (since both verbal and algebraic closedness of a subgroup A in a group \widetilde{A} are equivalent to the corresponding properties of $A \times B$ in $A \times B$);
- it remains to note that $Z(H) = F \times Z(H_1)$, and the centre of H_1 is its direct factor by Theorem KMO.

REFERENCES

- [Big01] S. Bigelow, Braid groups are linear, Journal of the American Mathematical Society, 14:2 (2001), 471-486.
- [Bog22] O. Bogopolski, Equations in acylindrically hyperbolic groups and verbal closedness, Groups, Geometry, and Dynamics, 16:2 (2022), 613-682. See also arXiv:1805.08071.
- [Che35] C. Chevalley, Démonstration d'une hypothèse de M. Artin, Abh. Math. Semin. Univ. Hambg., 11 (1935), 73-75.
- [D23] F. D. Denissov, Finite normal subgroups of strongly verbally closed groups, Journal of Group Theory (to appear). https://doi.org/10.1515/jgth-2023-0015 . See also arXiv:2301.02752.
- [G69] F. A. Garside, The braid group and other groups, The Quarterly Journal of Mathematics, 20:1 (1969), 235-254.
- [KK24] O. K. Karimova, A. A. Klyachko, Finite symmetric groups are strongly verbally closed, arXiv:2405.01179.
- [K21] A. A. Klyachko, The Klein bottle group is not strongly verbally closed, though awfully close to being so, Canadian Mathematical Bulletin, 64:2 (2021), 491-497. See also arXiv:2006.15523.
- [KM18] A. A. Klyachko, A. M. Mazhuga, Verbally closed virtually free subgroups, Sbornik: Mathematics, 209:6 (2018), 850-856. See also arXiv:1702.07761.
- [KMM18] A. A. Klyachko, A. M. Mazhuga, V. Yu. Miroshnichenko, Virtually free finite-normal-subgroup-free groups are strongly verbally closed, J. Algebra, 510 (2018), 319-330. See also arXiv:1712.03406.
- [KMO23] A. A. Klyachko, V. Yu. Miroshnichenko, A. Yu. Olshanskii, Finite and nilpotent strongly verbally closed groups, Journal of Algebra and Its Applications 22:09 (2023), 2350188. See also arXiv:2109.12397.
- [Kra02] D. Krammer, Braid groups are linear, Annals of Mathematics, 155:1 (2002), 131-156. See also arXiv:math/0405198.
- [Mazh17] A. M. Mazhuga, On free decompositions of verbally closed subgroups of free products of finite groups, J. Group Theory, 20:5 (2017), 971-986. See also arXiv:1605.01766.
- [Mazh18] A. M. Mazhuga, Strongly verbally closed groups, J. Algebra, 493 (2018), 171-184. See also arXiv:1707.02464.
- [Mazh18d] A. M. Mazhuga, Verbally closed subgroups, κ andudamcka duccepmayun (\approx Ph.D. thesis), MSU, Moscow, 2018 (in Russian), https://istina.msu.ru/dissertations/150563554/.
- [Mazh19] A. M. Mazhuga, Free products of groups are strongly verbally closed, Sbornik: Mathematics, 210:10 (2019), 1456-1492. See also arXiv:1803.10634.
- [MR14] A. Myasnikov, V. Roman'kov, Verbally closed subgroups of free groups, J. Group Theory, 17:1 (2014), 29-40. See also arXiv:1201.0497.
- [Neu51] B. H. Neumann, Groups with finite classes of conjugate elements, Proc. London Math. Soc., s3-1:1 (1951), 178-187.
- [O95] F. Oger, The direct decompositions of a group G with G/G' finitely generated, Trans. Amer. Math. Soc., 347:6 (1995), 1997-2010.
- [Rob96] D. J. S. Robinson, A Course in the theory of groups. Graduate texts in mathematics 80. Springer, 1996.
- [Rom12] V. A. Roman'kov, Equations over groups, Groups Complexity Cryptology, 4:2 (2012), 191-239.

^{*}) The "opposite" question of Mazhuga is still open: *is the class of strongly verbally closed groups closed with respect to direct products?* See [Mazh18d] for some results on this topic.

- [RKh13] V. A. Roman'kov, N. G. Khisamiev. Verbally and existentially closed subgroups of free nilpotent groups. Algebra and Logic, 52:4 (2013), 336-351.
- [RKhK17] V. A. Roman'kov, N. G. Khisamiev, A. A. Konyrkhanova, Algebraically and verbally closed subgroups and retracts of finitely generated nilpotent groups, Siberian Math. J., 58:3 (2017), 536-545.
- [RT20] V. A. Roman'kov, E. I. Timoshenko, Verbally Closed Subgroups of Free Solvable Groups, Algebra and Logic, 59:3 (2020), 253-265. See also arXiv:1906.11689.
- [Sch74] S. H. Schanuel, An extension of Chevalley's theorem to congruences modulo prime powers, Journal of Number Theory, 6:4 (1974), 284-290.
- [Sch04] J. Schur, Uber die Darstellung der endlichen Gruppen durch gebrochene lineare Substitutionen, Journal für die reine und angewandte Mathematik, 127 (1904), 20-50.
- [Tim21] E. I. Timoshenko, Retracts and verbally closed subgroups with respect to relatively free soluble groups, Sib. Math. J., 62:3, 537-544 (2021).
- [TJ74] J. M. Tyrer-Jones, Direct products and the Hopf property, J. Austral. Math. Soc., 17:2 (1974), 174-196.
- [Ya16] M. K. Yadav, Central Quotient Versus Commutator Subgroup of Groups, In: S. Rizvi, A. Ali, V. Filippis, (eds) Algebra and its Applications. Springer Proceedings in Mathematics & Statistics, 174 (2016). Springer, Singapore. See also arXiv:1011.2083.