# arXiv:2308.07714v1 [cs.AI] 15 Aug 2023

# Flashpoints Signal Hidden Inherent Instabilities in Land-Use Planning

Hazhir Aliahmadi,<sup>1</sup> Maeve Beckett,<sup>1</sup> Sam Connolly,<sup>1</sup> Dongmei Chen,<sup>2</sup> and Greg van Anders<sup>1</sup>

<sup>1</sup>Department of Physics, Engineering Physics, and Astronomy, Queen's University, Kingston ON, K7L 3N6, Canada

<sup>2</sup>Department of Geography and Planning, Queen's University, Kingston ON, K7L 3N6, Canada

Land-use decision-making processes have a long history of producing globally pervasive systemic equity and sustainability concerns. Quantitative, optimization-based planning approaches, e.g. Multi-Objective Land Allocation (MOLA), seemingly open the possibility to improve objectivity and transparency by explicitly evaluating planning priorities by the type, amount, and location of land uses. Here, we show that optimization-based planning approaches with generic planning criteria generate a series of unstable "flashpoints" whereby tiny changes in planning priorities produce large-scale changes in the amount of land use by type. We give quantitative arguments that the flashpoints we uncover in MOLA models are examples of a more general family of instabilities that occur whenever planning accounts for factors that coordinate use on- and between-sites, regardless of whether these planning factors are formulated explicitly or implicitly. We show that instabilities lead to regions of ambiguity in land-use type that we term "gray areas". By directly mapping gray areas between flashpoints, we show that quantitative methods retain utility by reducing combinatorially large spaces of possible land-use patterns to a small, characteristic set that can engage stakeholders to arrive at more efficient and just outcomes.

# I. INTRODUCTION

Sustainable, equitable land use will be a major challenge in this century due to the effects of urbanization [1, 2], climate change [3?, 4], and renewable energy generation [5]. Historically, allocating land-use qualitatively has produced outcomes that have raised serious, continuing issues regarding sustainability [6] and equity[7]. Inequitable and unsustainable land use can arise through planning processes that are topdown, subjective, and opaque. These shortcomings suggest a step forward through decision-making processes that explicitly and transparently articulate planning criteria for land use by type, amount, and location. These land-use factors can be modelled quantitatively, and optimized via techniques such as Multi-Objective Land Allocation (MOLA) [8]. However, although MOLA models for land-use planning problems make criteria clear and precise, solving MOLA and similar models typically relies on the use of optimization techniques [9-15] in which the relationship between the priorities of planning criteria and the resulting land-use allocations is obscured. Obscuring priority-outcome relationships undermines the promise of process-transparency that explicit models potentially provide for stakeholders. Moreover, this obscurity of optimization methods could mask dangerous instabilities in the models themselves. If that were the case, a standard, optimizationsupported land-use planning approach would not just be unclear, it would also be unreliable.

Here, we show that common land-use planning criteria produce outcomes that are extremely sensitive to the relative priority (weighting) of the criteria. We analyze a MOLA model [16] based on the two most commonly used planning criteria: compactness and suitability [8], and show that it exhibits a series of discrete, large-scale changes in land-use allocation outcomes in response to minute changes in planning priorities. This creates a series of instabilities we term "flashpoints".

For each flashpoint, we compare the land-use patterns on either side and map the discrepancies between the patterns. These discrepancies create a set of locations we term "gray areas" that are regions of land-use that are sensitive to infinitesimal changes in planning priorities.

By comparing with mathematically identical models of magnetic materials, we find that land-use flashpoints are equivalent to the most abrupt state changes known in nature. This mathematical equivalence, coupled with the general understanding developed over several decades of the study of phenomena in magnetism, indicates that an extremely broad class of MOLA models will exhibit flashpoints of the type we observe. This generality is driven by a conflict between the coordination of use between multiple land parcels and the coordination of a parcel's use with preexisting human or natural factors. These results indicate that conventional approaches to optimization-supported land-use planning are unreliable.

Though the inherent instability of MOLA means that there is no globally optimal land-use pattern for all suitability priority levels, our analysis yields a distribution of patterns that provides insight into the underlying trade-offs between planning priorities. We use techniques from statistical physics to construct explicit relationships between sets of planning priorities and the resulting land-use allocations they produce. By formulating priority-outcome relationships explicitly, we reduce the large space of possible land-use outcomes to a small set of representative patterns that are reliable over identifiable ranges of planning priorities. Explicit priority-pattern relationships of the form we construct here restore transparency to explicit models, and provide concrete information about planning trade-offs that can empower stakeholders to engage in conversation that could lead to more sustainable and equitable land-use decision making.

### II. RESULTS

**Compactness-suitability trade-off produces flashpoints.** To determine the dependence of globally optimal land-use distributions on the choice of priorities, we analyzed land-use patterns from 82,500 statistically independent simulations of the multi-objective land allocation model from Ref. [16] for various levels of suitability priority relative to compactness. Results are shown in Fig. 1.

In principle, land-use patterns could depend continuously or discontinuously on the priority of the suitability criterion,  $P_S$ . We determine this dependence in Fig. 1a, which shows the globally optimal land-use pattern in terms of the landuse fraction of land-use types. We aggregated an average of  $2 \times 10^5$  land-use patterns for each suitability priority. We determined global optima from inferred optimization landscapes, which we computed via Landau free energy minima (see Methods), using standard accumulation techniques, e.g. [17]. Apart from minor continuous changes in the land-use distribution for  $2.0 < P_S < 3.9$  and  $5.4 < P_S < 6.8$ , the response to changes of suitability priority was characterized by a series of abrupt, discontinuous changes we refer to as flashpoints (see Methods for cutoff).

Flashpoint instabilities produce gray areas. According to the criteria outlined in Methods, we identified six flashpoints between  $0.1 < P_S < 8$ . The existence of a flashpoint signals a significant redistribution of the land-use pattern in response to a small change in suitability priority. The corresponding redistribution at a flashpoint will produce regions of uncertainty in land use. Fig. 1b-g, middle images, depict these regions, called "gray areas".

We determined gray areas by aggregating land-use patterns at Landau free energy minima. We represent each parcel's land-use type by a point on a circle where the three land-use types are separated by 120 degrees (see Methods). Taking the average of those points across several thousands of land-use patterns gives an average for each parcel within the circle, as shown in the bottom right of Fig. 1. Suitability priority just below a flashpoint (Fig. 1b-g, left images) or above that (right images) results in a consistent land-use pattern, i.e., little to no gray, across samples. However, comparing land-use patterns on either side of the flashpoints shows significant gray areas associated with different land-use patterns. This finding indicates that there are large regions with extreme parametric sensitivity.

**Spontaneous Symmetry Breaking Creates the Conditions** for Flashpoints. To understand how flashpoints arise, we consider a simplified model with a vanishing suitability priority,  $P_S = 0$ , so that land-use patterns optimize spatial compactness alone. In the analysis above, we worked at fixed annealing threshold, T. Now, instead we vary the threshold and track emergent land-use patterns as a function of the threshold. Results for the compactness-only model are depicted in Fig. 2. Fig. 2a shows randomly selected land-use snapshots at a range of annealing thresholds. This compactnessonly model shows non-compact solutions at high thresholds, while land-use patterns become compact at low thresholds, with one land-use type dominating the entire map. Prevailing single-use patterns at low annealing thresholds, in a model that makes no explicit preferential distinction between landuse types, occur because multiple local minima emerge corresponding to each possible land use. The emergence of multiple local minima signals a phenomenon termed spontaneous symmetry breaking [18] in physics.

Fig. 2b plots distributions of all sampled land-use patterns at thresholds corresponding to T = 1.0, indicating broken

symmetry with minimal fluctuations, validating our choice of T = 1.0 in the flashpoint analysis shown in Fig. 1. A similar analysis of land-use distributions is shown in Fig. 2c-d suggests symmetry breaking occurs between  $2.2 \le T \le 2.3$ . Results for all thresholds are shown in Supplementary Movie 2.

To confirm that spontaneous symmetry braking is occurring, we computed inferred optimization landscapes, via Landau free energies (see Methods), as a function of land-use fraction for each studied threshold. Fig. 3a shows locations of free energy minima as a function of threshold, and panels b and c show results at T = 2.2 and 2.3 confirming spontaneous symmetry breaking occurs between those thresholds. The increase in the statistical uncertainty of the land-use fraction near the symmetry-breaking threshold is a known physical effect that comes from diverging fluctuation magnitudes near a so-called critical point [18].

Comparing the compactness-only model with the compactness-suitability trade-off model suggests that increasing suitability priority at low annealing-thresholds disrupts "local" balances between compactness and suitability. When local balance is disrupted, the result is a discrete, large-scale use reallocation of many land parcels in a region between single-use and multi-use patterns.

Flashpoints from Generic Disrupted Multi-site/On-site Coordination Balance. It is possible to substantiate this disrupted compactness–suitability balance argument quantitatively, and more generally. Generic planning objectives can arise from human factors (e.g. social, economic, or demographic) and natural factors (e.g. geographic, ecological, environmental). Both human and natural factors can drive the land use of a parcel to either coordinate with multiple adjacent parcels or to coordinate with the features or properties of the parcel itself [8]. Examples of drivers of multi-site coordination include the provision of social or community services and infrastructure, [?]; or contiguity of habitat for wildlife. Examples of drivers of on-site coordination include prior land use, flood risk, terrain, soil quality, etc.

To formulate the balance between multi-site and on-site coordination quantitatively, note that, generically, multi-site coordination will drive spontaneous symmetry breaking that will induce regions with homogeneous, coordinated use. For parcels in a region R of the map, the objective "cost" of real-location of the land use is given by

$$\Delta F = \Delta E_M L(\partial R) - \Delta E_O A(R) , \qquad (1)$$

where  $\Delta E_M$  is the average objective increase for breaking multi-site coordination per unit length of the boundary,  $\partial R$ , of region R, L is the length of the boundary,  $\Delta E_O$  is the average objective reduction for on-site coordination per unit area, and A is the area of the region. A region will switch its land-use if  $\Delta F < 0$ , i.e. if it reduces overall objective cost. Reducing objective cost by breaking homogeneity in a region will occur whenever the average on-site cost reduction is sufficient that

$$\Delta E_O > \frac{\Delta E_M L(\partial R)}{A(R)} . \tag{2}$$



FIG. 1. Changing priorities in quantitative land-use allocation generates flashpoints: priority sets in which small changes generate largescale reorganization in land use patterns. **a** The optimal land-use fraction computed via simulated annealing-like methods with low annealing threshold (T = 1) as a function of suitability priority level  $P_S$ . With changing priority, the land-use fraction shows relatively stable regions punctuated by a series of discrete jumps. **b-g** Comparing land-use patterns on either side of each flashpoint shows that areas of change are generally clusters of parcels rather than peripherals or borders between land-use types. Maps are computed by labelling each land-use type with an integer, transforming it to a point on a circle in the complex plane, and then averaging over hundreds or thousands of statistically independent realizations of land-use patterns, according to the key below panels **b-g**.

In this relationship, which is an application of general principles of nucleation theory [19], the left-hand side scales linearly with the priority for on-site coordination, whereas the right-hand side scales linearly with the priority for multi-site coordination. For a region of any geometry defined by a given L/A, there will be some set of weights that cross the condition defined in Eq. (2). Thus flashpoints are inherent in any land allocation problem with allocation objectives that involve multi-site and on-site coordination, regardless of the details of the optimization method or the model.

# III. DISCUSSION

We analyzed several tens of thousands of statistically independent multi-objective land allocation patterns spread over a range of planning priorities. We found optimal land-use amounts by type exhibited extreme sensitivity to the priorities of planning criteria, with use type undergoing a series of discrete, large-scale changes we termed flashpoints. Flashpoint instabilities signal that optimization-based modelling like MOLA, conventionally deployed, is not a reliable support for land-use planning. We mapped corresponding spatial locations of land-use instability, called gray areas.

Our results were produced using simulated annealing-like methods applied to a specific model that is known in the literature, however, the effects we found are more general. We compared allocation models with mathematically analogous systems that model the physics of magnetism. The mathematical equivalence between quantitative land-use models and a broad class of magnetic systems for which a consistent understanding developed through decades of theoretical, computational, and experimental investigation with multiple approaches [20] means that the instabilities we observed are not artifacts of our specific model or methods. Instead, our results signal that flashpoints are a generic feature of any land allocation problem that combines objectives for compactness and suitability.

To relate our findings to other methods for similar problems, it is important to note that the approach we took to MOLA here maps the parameters of the model to a statistical distribution of candidate solutions. This distribution is determined by the structure of the underlying solution space, so it is a property of the land-use model, not the optimization method. Other methods such as genetic algorithms [10, 11, 13, 15], particle swarm optimization [9, 12], or ant colony optimization [14] might explore the solution space differently, but that space would still be marked by the flashpoint instabilities we reported here. To relate our findings to other models, note that although the model we used here had particular forms for multi-site and on-site coordination, mathematically similar forms of coordination result from disparate human or natural factors. For example, factors driving multi-site coordination such as the importance of contiguity in wildlife habitats or the benefits of contiguous communities for providing infrastructure or public services suggest spatial compactness in

4



FIG. 2. Compactness-only land-use allocation model shows spontaneous symmetry breaking with decreasing annealing threshold, indicating the underlying optimization landscape is rough. **a** Selected annealing replicates at fixed, decreasing annealing threshold (corresponding to T; decreasing from right to left) show compactness-only models at low annealing thresholds are characterized by a predominance of uniform, single land-use patterns. Data at each threshold are shown for 10 statistically independent annealing replicates. Data from all replicates are aggregated in ternary plots (see Methods) in SI Movie 2 and panels **b-d**. Panels **b-d** show land-use where the vertices of the triangle correspond to single-use patterns. Each simulation sample is plotted as an ordered triple, representing land-use fractions of land-use types (Agriculture, Construction, Conservation). Coloured lines indicate lines of constant land-use fractions of the corresponding land-use type, and the centroid of the triangle represents an equal distribution of land-uses among the three types. Panel **b** shows that at a low threshold (T = 1) data points localize the ternary vertices, indicating near single-use maps. In contrast, at a higher threshold (T = 2.3; panel **d**), data localize with triangular symmetry at the centroid but are instead becoming peripheral, signalling that so-called spontaneous symmetry breaking underlies the rough optimization landscape.

a generic form similar to the present model. Indeed, a recent systematic review found that compactness is the most common MOLA-modelling criterion [8]. The same systematic review has also shown that suitability is a frequent model criterion [8], but any other factor that produces on-site coordination, e.g., distributions of prior use, will lead to similar effects. Some form of spatial inhomogeneity is also inevitable since regions of land frequently have differences in, e.g., flood risk, soil quality, proximity to water, or terrain. The interaction of multi-site and on-site coordination will generally produce flashpoints (see Fig. 4).

Our investigation indicates that, generically, land-use al-

location models reduce the large space of possible land-use patterns to a small set of basic arrangements that are punctuated by extreme sensitivity to the relative priority of competing planning criteria. Simply making a choice about the priority of various objectives and then determining an optimal solution will produce results that can exhibit spectacular dependence on arbitrarily small changes in planning priorities. In quantitative models, the abruptness of these changes in optimization outcomes is mathematically equivalent to so-called first-order phase transitions, the most abrupt changes of state that are known in nature [21]. Abruptly triggered instabilities run directly counter to the resilience that has long been sought

5



FIG. 3. Inferred optimization landscapes (Landau free energy) at vanishing suitability priority confirms spatial compactness drives spontaneous symmetry breaking. **a** Landau free energy minima at a range of thresholds indicates low thresholds spontaneously fix single land use patterns. Data points indicate positions of free energy minima extracted from sampling more than 2,000 statistically independent simulations spread across the indicated range of annealing thresholds. **b,c** Landau free energy computed as a function of land use fraction confirms symmetry breaking occurs at a threshold  $2.2 \le T \le 2.3$ . Statistical error is on the order of the "jitter" in the data traces.

in managing the link between social and ecological systems [22, 23].

Our analysis of land-use change was quantitative. However, generic features of the broad class of models that are similar to ours indicate that land-use decision-making that confronts multi-site and on-site coordination is subject to a perilous balance regardless of whether these factors are explicitly quantified or not. Rather than using quantitative approaches to land-use allocation for top-down optimization, our analysis suggests quantitative approaches could more profitably confront an unstable balance among planning priorities by charting relationships between the priorities and land-use outcomes (see Fig. 5).

Finally, top-down, allocation modelling alone cannot drive

sustainable land-use planning. In land-use planning, as in other domains, sustainability requires iterative, reciprocal interaction that drives coproduction between science and policy [24]. Our approach expands allocation techniques because it moves beyond singular, putative optimal outcomes of a given model, and renders explicit priority–outcome relationships among families of underlying land-use models. Making these modelling relationships explicit is a critical step forward in nurturing conversations between scientists and policymakers that inform refined or revised models to better support sustainable land-use planning.



FIG. 4. Land-use flashpoints arise generically when common human and natural considerations drive conflict between multi-site land use coordination and on-site coordination. **a** Both human factors and natural factors induce planning considerations about the coordination between adjacent land parcels and coordination of a land parcel's use with the, possibly pre-existing, properties. Multi-site and on-site forms of coordination combine to generate the potential for instability that produces flashpoints. **b** Multi-site coordination arises because a given land use for a particular parcel generates an influence area on adjacent parcels. **c** On-site coordination drives the use of a parcel according to local factors.



FIG. 5. Substantive science–policy dialogue requires charting explicit connections between planning goals (pie charts, bottom), planning priorities (axis, middle), and planning outcomes (maps, top). We match pie charts showing possible targets for fractions of land allocated for given uses to corresponding families of land-allocation models that differ in planning priorities but yield similar allocations. We further match fractional allocations with spatial distributions of land-use types. Colour shading indicates regions of commonality across a family of models, and gray shading indicates variation.

# IV. METHODS

# A. Multi-Objective Land Allocation Model

We model land-use patterns as a set of discrete parcels each with a potentially different use subject to overall planning criteria following established conventions for multi-objective land allocation (MOLA) models [8]. Systematic review [8] has shown that spatial compactness and suitability are generic MOLA planning criteria. For concreteness, we work with a model described in Ref. [16] with compactness and suitability criteria that are sufficiently representative to facilitate extracting generalizable lessons. The data used for this study are based on a semi-hypothetical dataset provided in [10], which pertains to a square area of 9 km<sup>2</sup> located in the Xin'andu township of Dongxihu District, Wuhan, China.

Following Ref. [16], we take the criterion for compactness as

$$O_1 = -\sum_{s=0}^{S} \sum_{i=1}^{N} \sum_{j=1}^{M} b_{ijs} x_{ijs} , \qquad (3)$$

where M and N are the number of columns and rows of the map, S is the number of land-use types, and  $x_{ijs}$  is 1 if the parcel at (i, j) is allocated for land-use s, and 0 otherwise. The coefficient  $b_{ijs}$  counts the number matching neighbours for the (i, j) parcel

$$b_{ijs} = x_{i+1j+1s} + x_{ij+1s} + x_{i+1js} + x_{i-1js} + x_{ij-1s} + x_{i-1j-1s} + x_{i-1j-1s}$$

We take suitability criterion as

$$O_2 = -\sum_{s=0}^{S} \sum_{i=1}^{N} \sum_{j=1}^{M} c_{ijs} x_{ijs} , \qquad (5)$$

where  $c_{ijs}$  parametrizes the suitability of the (i, j) parcel for land-use type s. Here, we follow Ref. [16] using M = N =30 to give a 30×30 map, and three land-use types; agriculture, construction, and conservation labelled with the integers s =0, 1, 2, respectively. Suitability data  $c_{ijs}$  are given in SI.

We form a weighted combination of the planning criteria Eqs. (3) and (5) as a Hamiltonian

$$H = P_C O_1 + P_S O_2 , (6)$$

where  $P_C$  and  $P_S$  are weights that correspond to priorities for the compactness and suitability objectives, and are physical analogues of pressure [25].  $P_C \gg P_S$  or  $P_C \ll P_S$  generate allocations dominated by one criterion or another. Trade-offs occur when  $P_C$  and  $P_S$  are of a similar order. We fixed units by setting  $P_C = 1$  and taking  $0.1 < P_S < 8$ , which covers the essential region of compactness/suitability trade-offs. Similar to techniques from simulated annealing [26], we sample patterns using a Markov Chain Monte Carlo algorithm.

### B. Markov Chain Monte Carlo Sampling

We study the multi-objective land allocation model using Markov Chain Monte Carlo (MCMC) techniques, using the Wolff algorithm.[27] The presence of suitability objective creates a spatial inhomogeneity that violates a symmetry assumption of the standard Wolff algorithm, so we implemented a modified version described in Ref. [28]. Land-use patterns are sampled according to an annealing threshold  $k_BT$ , where T is equivalent to the temperature in simulated annealing approaches to MOLA (e.g. Ref. [16]), and  $k_B$  is a constant expressing the conversion between energy and temperature units (we set  $k_B = 1$  without loss of generality). C++/Python source code is available at Ref. [29].

To ensure accurate results, for our flashpoint analysis we thermalized systems of  $30 \times 30$  land units at a temperature of T = 15 for  $10^3$  Monte Carlo (MC) sweeps. We cooled systems to a target temperature of T = 1 over  $3.5 \times 10^4$  MC

sweeps. We equilibrated systems at the target temperature for  $10^4$  MC sweeps. We measured land-use distributions at intervals of 50 MC sweeps over  $10^4$  sweeps. We computed an average of more than  $10^3$  statistically-independent simulations over the entire suitability priority range using a minimum of 300 random seeds at priorities far from flashpoints, and many more near flashpoints. Data are available at Ref. [30].

## C. Ternary Plots

Our MC sampling approach produced more than  $1.6 \times 10^7$ <sup>i</sup>Tahd-thse patterns!: A complete display in the form of Fig. 2a would require more than  $1.4 \times 10^{10}$  pixels. We, therefore, analyzed land-use patterns by aggregating land use of parcels across each pattern for each land-use type ( $N_X$  where X represents either agriculture, construction, or conservation). We represented each pattern as an ordered triple. Considering ordered triples as defining a three-dimensional space, ordered triples for fixed map size all lie on a plane that is normal to the vector (1,1,1). We projected our ordered triples to this plane, where all possible land-use patterns fall inside an equilateral triangle, with single-use patterns at the vertices.

We plotted land-use totals by placing a marker for observed patterns in Fig. 2b-d, and in SI Movies 1 and 2. In order to not violate the limits of our plotting software, we plotted 5000 randomly selected patterns at each choice of the priority set for a fixed annealing threshold.

# D. Landau Free Energy

In generic multi-objective, non-convex optimization problems, possible outcomes depend non-trivially on system configurations and on the relative priority of objectives. To determine the form of the outcome landscape in physical systems it is conventional to infer its form from statistical sampling where the landscape is known as the Landau free energy [18].

We determine optimal land use patterns for a given set of planning priorities by generating patterns via MCMC sampling as described above. We aggregated sampled patterns by the count of uses of each type. We accumulated these counts over thousands of samples to compute a histogram by landuse type. This histogram produces a Monte Carlo estimate of the probability of sampling patterns of fixed-use counts. The negative logarithm of this distribution gives the inferred optimization landscape (Landau free energy [18]), and the global minimum of this landscape corresponds to the optimal land use pattern.

Additionally, the structure of a Landau free energy surface as a function of system parameters encodes distinct regimes of system behaviour and the abruptness of transitions between them [18]. Continuous, so-called symmetry-breaking, transitions occur when changes in system parameters induce a single, global minimum to split into multiple minima [18], e.g., as seen in Fig. 3. Abrupt, so-called first-order, transitions occur when changes in system parameters induce an exchange between local and global minima [18], e.g., which drives the behaviour in Fig. 1.

# E. Flashpoint Cutoff

A flashpoint is an abrupt change in land use outcomes in response to slight changes in planning priorities. For this investigation, we define a flashpoint as a 10% change in any land-use type across a single incremental change in suitability priority according to

$$\frac{|N_X(P_S^{n+1}) - N_X(P_S^n)|}{\frac{1}{2}[N_X(P_S^{n+1}) + N_X(P_S^n)]} > \alpha \tag{7}$$

where  $N_X(P_S^n)$  is the number of land use X at the n<sup>th</sup>  $P_S$ , and  $\alpha = 0.1$  is the flashpoint cutoff.

To ensure accurate  $N_X$  we computed Landau Free Energy minima [18] at T = 1.0, sufficiently low that thermal fluctuations are small. Lower temperatures would only sharpen the behaviour we observed. We averaged results from 500 statistically independent replicates for each  $0.1 < P_S < 8.0$  in increments of 0.1. For priorities near the flashpoints, we used 2000 replications to ensure accuracy.

# F. Gray Areas

We mapped gray areas via a quantity that expresses local land-use patterns. We did that by labelling each parcel of a pattern by an integer s = 0, 1, 2, corresponding to the landuse types; agriculture, construction, and conservation, respectively. Following the Landau free energy computation for the

- K. C. Seto, B. Güneralp, and L. R. Hutyra, Proceedings of the National Academy of Sciences 109, 16083 (2012).
- [2] United Nations Department of Economic and Social Affairs, Population Division, *World Urbanization Prospects, the 2018 Revision* (United Nations, New York, 2019).
- [3] A. F. Prein, R. M. Rasmussen, K. Ikeda, C. Liu, M. P. Clark, and G. J. Holland, Nature Climate Change 7, 48 (2017).
- [4] J. Huang, H. Yu, X. Guan, G. Wang, and R. Guo, Nature Climate Change 6, 166 (2016).
- [5] D.-J. van de Ven, I. Capellan-Peréz, I. Arto, I. Cazcarro, C. de Castro, P. Patel, and M. Gonzalez-Eguino, Sci Rep 11, 2907 (2021).
- [6] S. E. Bibri, J. Krogstie, and M. Kärrholm, Developments in the Built Environment 4, 100021 (2020).
- [7] I. Anguelovski, L. Shi, E. Chu, D. Gallagher, K. Goh, Z. Lamb, K. Reeve, and H. Teicher, Journal of Planning Education and Research 36, 333 (2016).
- [8] M. M. Rahman and G. Szabó, Sustainable Cities and Society 74, 103214 (2021).
- [9] Y. Liu, J. Peng, L. Jiao, and Y. Liu, PLOS ONE 11, e0157728 (2016).
- [10] M. Song and D. Chen, Geo-spatial Information Science 21, 273 (2018).

flashpoint threshold, described above, we aggregate patterns at the free energy global minimum and map each parcel's land-use type to a point on the unit circle in the complex plane via

$$z = \exp\left(\frac{2\pi i(s-1)}{3}\right) \,. \tag{8}$$

We then average z of each parcel at the free energy minimum for the suitability priority immediately to either side of the flashpoint, and the combination of both sides. This average  $\langle z \rangle$  over the set of mapped points gives a point on the unit circle for each parcel, which we shade according to the colour map shown in Fig. 1.

# V. ACKNOWLEDGEMENTS

We acknowledge the support of the Natural Sciences and Engineering Research Council of Canada (NSERC) grants RGPIN-2019-05655, DGECR-2019-00469, and RGPIN-2019-05773. Computations were performed on resources and with support provided by the Centre for Advanced Computing (CAC) at Queen's University in Kingston, Ontario. The CAC is funded by: the Canada Foundation for Innovation, the Government of Ontario, and Queen's University. GvA acknowledges the hospitality of the Kavli Institute for Theoretical Physics (KITP) where this work was completed. This research was supported in part by the National Science Foundation under Grant No. NSF PHY-1748958. We also thank Dr. Mingjie Song for providing the land use and suitability data.

- [11] T. J. Stewart, R. Janssen, and M. van Herwijnen, Computers & Operations Research **31**, 2293 (2004).
- [12] H. Zhang, Y. Zeng, X. Jin, B. Shu, Y. Zhou, and X. Yang, Ecological Modelling 320, 334 (2016).
- [13] M. Mohammadi, M. Nastaran, and A. Sahebgharani, INDJST 8, 1 (2015).
- [14] X. Liu, X. Li, X. Shi, K. Huang, and Y. Liu, International Journal of Geographical Information Science 26, 1325 (2012).
- [15] P. Gao, H. Wang, S. A. Cushman, C. Cheng, C. Song, and S. Ye, Landscape Ecol 36, 1877 (2021).
- [16] M. Song and D. Chen, Annals of GIS 24, 19 (2018).
- [17] G. van Anders, D. Klotsa, N. K. Ahmed, M. Engel, and S. C. Glotzer, Proc. Natl. Acad. Sci. U.S.A. **111**, E4812 (2014).
- [18] N. Goldenfeld, Lectures on Phase Transitions and the Renormalization Group (Addison-Wesley, Reading MA, 1992).
- [19] P. G. Debenedetti, *Metastable Liquids: Concepts and Principles*, Physical Chemistry (Princeton University Press, Princeton, N.J, 1996).
- [20] S. Chikazumi, *Physics of Ferromagnetism*, second edition, second edition ed., International Series of Monographs on Physics (Oxford University Press, Oxford, New York, 2009).
- [21] J. Sethna, *Statistical Mechanics: Entropy, Order Parameters,* and Complexity (Oxford University Press, USA, 2021).

- [22] F. Berkes, C. Folke, and J. Colding, *Linking Social and Ecological Systems: Management Practices and Social Mechanisms* for Building Resilience (Cambridge University Press, 2000).
- [23] R. S. DeFries, J. A. Foley, and G. P. Asner, Frontiers in Ecology and the Environment **2**, 249 (2004).
- [24] W. C. Clark, L. van Kerkhoff, L. Lebel, and G. C. Gallopin, Proceedings of the National Academy of Sciences 113, 4570 (2016).
- [25] A. A. Klishin, C. P. Shields, D. J. Singer, and G. van Anders, New J. Phys. 20, 103038 (2018), arxiv:1709.03388

[physics.soc-ph].

- [26] S. Kirkpatrick, C. D. Gelatt, and M. P. Vecchi, Science 220, 671 (1983).
- [27] U. Wolff, Phys. Rev. Lett. 62, 361 (1989).
- [28] J. Kent-Dobias and J. P. Sethna, Phys. Rev. E 98, 063306 (2018).
- [29] S. Connolly, M. Beckett, H. Aliahmadi, and G. van Anders, (2023), 10.5281/ZENODO.7846107.
- [30] H. Aliahmadi, M. Beckett, S. Connolly, D. Chen, and G. van Anders, (2023), 10.5683/SP3/KVXIMS.