THE ALGEBRAIC DEGREE OF SPARSE POLYNOMIAL OPTIMIZATION

JULIA LINDBERG, LEONID MONIN, AND KEMAL ROSE

ABSTRACT. We study a broad class of polynomial optimization problems whose constraints and objective functions exhibit sparsity patterns. We give two characterizations of the number of critical points to these problems, one as a mixed volume and one as an intersection product on a toric variety. As a corollary, we obtain a convex geometric interpretation of polar degrees, a classical invariant of algebraic varieties, as well as Euclidean distance degrees. Furthermore, we prove the BKK generality of Lagrange systems in many instances.

Keywords: sparse polynomial optimization, toric varieties, algebraic degree, ED degree, polar degree MSC2020: 90C26, 14M25, 14Q20, 52B20

1. INTRODUCTION

We consider polynomial optimization problems of the form:

(POP)
$$
\min_{y \in \mathbb{R}^n} f_0(y) \quad \text{subject to} \quad f_1(y) = 0, \dots, f_m(y) = 0
$$

where $f_i \in \mathbb{R}[y_1, \ldots, y_n]$. Polynomial programs have broad modeling power and are found in applications such as signal processing, combinatorial optimization, power systems engineering, and more [\[TR01,](#page-28-0) [PRW95,](#page-28-1) [MH19\]](#page-27-0). In general, these problems are NP-hard to solve [\[Vav90\]](#page-28-2) but there has been much work in recent decades studying various solution techniques for [\(POP\)](#page-0-0). Inspired by recent improvements in numerical methods for polynomial system solving, in this work we focus on solving the critical point equations for [\(POP\)](#page-0-0).

When the first-order optimality conditions hold, there are finitely many complex critical points to [\(POP\)](#page-0-0). For a specified objective function f_0 and fixed constraints f_1, \ldots, f_m we denote $\mathbf{F} = (f_0, \ldots, f_m)$. The number of complex critical points of **F** is called the *algebraic degree* of **F**. While [\(POP\)](#page-0-0) is a real optimization problem, we consider complex critical points since for polynomials \bf{F} with fixed monomial support, the algebraic degree is generically^{[1](#page-0-1)} constant. When the algebraic degree of [\(POP\)](#page-0-0) is finite, the coordinates of an optimal solution of [\(POP\)](#page-0-0) can be expressed as the roots of polynomial functions of the coefficients of **. Under sufficient genericity** conditions these polynomial functions are all of the same degree and the algebraic degree of F has an additional interpretation as the degree of these polynomials.

The algebraic degree of \bf{F} also has practical implications for optimization algorithms as it gives an upper bound on the number of critical points of [\(POP\)](#page-0-0). This gives a bound on the number of local extrema at which local optimization algorithms can terminate.

When **F** consists of generic polynomials with full monomial support, a formula for the algebraic degree of \bf{F} was given in [\[NR09\]](#page-27-1). This was then specialized

¹By generic, we mean generic with respect to the Zariski topology. See Remark [2.1](#page-4-0) for a detailed explanation.

for many classes of convex polynomial optimization problems in [\[GvBR09\]](#page-27-2) and [\[NRS10\]](#page-28-3). When the objective function of [\(POP\)](#page-0-0) is the Euclidean distance function, i.e. $f_0 = ||y - u||_2^2$ for a generic point $u \in \mathbb{R}^n$, the algebraic degree of **F** is called the ED degree of (f_1, \ldots, f_m) . The study of ED degrees began with $[DHO+16]$ $[DHO+16]$. Follow up work then studied the ED degree of real algebraic groups [\[BD15\]](#page-26-1), Fermat hypersurfaces [\[Lee17\]](#page-27-3), orthogonally invariant matrices [\[DLOT17\]](#page-26-2), smooth complex projective varieties [\[AH18\]](#page-26-3), the multiview variety [\[MRW20a\]](#page-27-4), hypersurfaces [\[BSW22\]](#page-26-4) and when the data u and polynomials f_1, \ldots, f_m are not generic [\[MRW20b\]](#page-27-5).

A related problem coming from statistics considers the likelihood objective function, $f_0 = y_1^{u_1} \cdots y_n^{u_n}$. This problem is called *maximum likelihood estimation* and the number of complex critical points to [\(POP\)](#page-0-0) when $f_0 = y_1^{u_1} \cdots y_n^{u_n}$ for generic $u \in \mathbb{R}^n$ is called the *ML degree* of (f_1, \ldots, f_m) . Relationships between ML degrees and Euler characteristics as well as the ML degree of various statistical models have been studied in [\[CHKS06,](#page-26-5) [HKS05,](#page-27-6) [Huh13,](#page-27-7) [ABB](#page-26-6)⁺19, [DM21,](#page-27-8) [MMW21,](#page-27-9) [MMM](#page-27-10)⁺23].

Inspired by recent results on the ED and ML degrees of sparse polynomial systems [\[BSW22,](#page-26-4) [LNRW23\]](#page-27-11), we study the algebraic degree of [\(POP\)](#page-0-0) when each polynomial $f_i \in \mathbb{R}[y_1,\ldots,y_n]$ is assumed to be a *sparse polynomial* with generic coefficients. Given an optimization problem of the form (POP) where \bf{F} is a list of sparse polynomial equations with generic coefficients, the Lagrangian of \bf{F} is

$$
\Phi_F(\lambda, y) := f_0 - \sum_{i=1}^m \lambda_i f_i.
$$

We consider the *Lagrange system* of **F**, namely $\mathbf{L}_{\mathbf{F}} = (f_1, \ldots, f_m, \ell_1, \ldots, \ell_n)$, where

$$
\ell_j = \frac{\partial}{\partial y_j} \left(f_0 - \sum_{i=1}^m \lambda_i f_i \right).
$$

Analogous to the *algebraic degree of polynomial optimization* from [\[NR09\]](#page-27-1), we generalize the common term to the algebraic degree of sparse polynomial optimization. It is the number of complex critical points of f_0 restricted to $\mathcal{V}(f_1, \ldots, f_m)$, where each $f_i \in \mathbb{R}[y_1, \ldots, y_n]$ is a sparse polynomial. When all critical points are smooth, it is:

$$
(1.1) \quad \# \mathcal{V} (\mathbf{L}_{\mathrm{F}}) = \# \left\{ (y, \lambda) \in \mathbb{C}^n \times \mathbb{C}^m : \quad 0 = f_1 = \cdots = f_m = \ell_1 = \cdots = \ell_n \right\}.
$$

There exist classical results in algebraic geometry bounding the number of isolated solutions to a square polynomial system. A result of Bézout says that $\#\mathcal{V}(L_F)$ is bounded above by the product of the degrees of the polynomials in L_F . If $deg(f_i) = d_i$ and $deg(\ell_j) = h_j$, $0 \leq i \leq m, 1 \leq j \leq n$, Bézout's bound reduces to $d_1 \cdots d_m \cdot h_1 \cdots h_n$ where $h_j \leq \max_{i \in [m]} \{d_0 - 1, d_i\}$. Work of Nie and Ranestad refined this bound and showed that

$$
\# \mathcal{V}(\mathbf{L}_{\mathrm{F}}) \leq d_1 \cdots d_m \cdot D_{n-m}(d_0-1,\ldots,d_m-1)
$$

where $D_r(n_1,\ldots,n_k) = \sum_{i_1+\cdots+i_k=r} n_1^{i_1}\cdots n_k^{i_k}$ [\[NR09\]](#page-27-1). While this bound is generically tight when F consists of polynomials with full monomial support, the following example shows that it can be a strict upper bound when \bf{F} is sparse.

Example 1.1. Consider the following optimization problem:

(1.2)
$$
\min_{y \in \mathbb{R}^n} c^T y \text{ subject to } f = \alpha_1 y_1^3 + \sum_{j=2}^{n-1} \alpha_j y_j^2 + \alpha_n y_n = 1,
$$

where $c, \alpha \in \mathbb{R}^n$ are generic parameters. The corresponding Lagrange system is given by $\mathbf{L}_{\text{F}} = (\ell_1, \ldots, \ell_n, f)$ where

$$
\ell_1 = c_1 - 3\lambda \alpha_1 y_1^2, \quad \ell_n = c_n - \alpha_n \lambda, \quad \ell_j = c_i - 2\lambda \alpha_j y_j, \quad 2 \le j \le n - 1.
$$

Bézout's bound gives that $\#\mathcal{V}(L_F) \leq 9 \cdot 2^{n-2}$, which is worse than the Nie-Ranestad bound $\#\mathcal{V}(\mathbf{L}_{\mathrm{F}}) \leq 3 \cdot D_{n-1}(0,2) = 3 \cdot 2^{n-1}$. In this case, the sparsity of the Lagrange equations allows one to solve the Lagrange system explicitly. One can see that for generic values of c and α , $\#\mathcal{V}(\mathbf{L}_{\mathrm{F}}) = 2$.

Motivated by the previous example, we are interested in obtaining tighter bounds for the algebraic degree of sparse polynomial optimization problems. We focus on the following questions.

Question 1. How many smooth critical points does [\(POP\)](#page-0-0) have for sparse F?

The motivation for Question [1](#page-2-0) is that if we know how many critical points [\(POP\)](#page-0-0) has, and we find them all, then we can globally solve [\(POP\)](#page-0-0). Currently, the only way to *provably* find all smooth critical points is to find all complex zeros of L_F .

The field of computational algebraic geometry has traditionally been associated with symbolic computations based on Gröbner bases. Recent developments in numerical frameworks, such as homotopy continuation [\[BHSW13\]](#page-26-7), provide algorithms that are able to solve problems intractable with symbolic methods. Moreover, numerical algorithms can not only provide the floating point approximation of a solution, but also certify that a given approximation represents a unique solution, and provide guarantees for correctness [\[BRT23,](#page-26-8) [Rum99,](#page-28-4) [Lee19\]](#page-27-12). Therefore, numerical computation can be used to prove lower bounds on $\#\mathcal{V}(L_F)$. However, ensuring the absence of additional solutions requires establishing an upper bound, which can be achieved through intersection theory.

Such an intersection theoretic bound for sparse polynomial systems was given by the celebrated Bernstein-Kouchnirenko-Khovanskii (BKK) theorem. The BKK the-orem [\[Ber75\]](#page-26-9) relates the number of $\mathbb{C}^* := \mathbb{C} \setminus \{0\}$ zeros of a system of polynomial equations to the mixed volume of their corresponding Newton polytopes (see Section [2.1\)](#page-3-0). While the BKK bound is generically tight, we note that the coefficients of the system L_F are linearly dependent, so a priori it is not clear that the system L_F has the expected number of solutions. This motivates the second question guiding this work.

Question 2. Is the Lagrange system of [\(POP\)](#page-0-0) BKK general?

An affirmative answer to Question [2](#page-2-1) would show that polyhedral homotopy algorithms are optimal for finding all complex critical points for generic instances of [\(POP\)](#page-0-0) in the sense that for every solution to $L_F = 0$, exactly one *homotopy path* is tracked. For more details on polyhedral homotopy continuation see [\[HS95\]](#page-27-13).

A second consequence of Question [2](#page-2-1) has to do with the complexity of algorithms in real algebraic geometry. A fundamental question in real algebraic geometry is to find a point on each connected component of a real algebraic variety, Y . In [\[Hau13\]](#page-27-14), Hauenstein showed that when f_0 is the Euclidean distance function, the zeros of the Lagrange system of f_0 restricted to Y cover all connected components. He then used this result to suggest a homotopy algorithm that would find a point on each connected component of Y . In Hauenstein's homotopy algorithm, he partitions the homotopy paths into two sets, E and E_1 , where E denotes the set of homotopy paths that converge and E_1 denotes the set of homotopy paths that diverge. By showing that polyhedral homotopy algorithms are optimal for a large class of Euclidean distance optimization problems, we would further the analysis of Hauenstein's algorithm by showing that the set E_1 is empty.

Finally, we note that understanding BKK exactness of non generic polynomial systems is of increasing interest in the applied algebraic geometry community. In this field, researchers study systems of polynomial equations coming from a diverse set of fields including biology, optimization, data science and power systems engineering. Recent results also study BKK exactness in these settings [\[CCG23,](#page-26-10) [BSW22,](#page-26-4) [LNRW23,](#page-27-11) [LAR23,](#page-27-15) [CKL22\]](#page-26-11).

Contribution. Our contribution consists of multiple results determining the algebraic degree of generic sparse polynomial programs. First, we show in Theorem [B,](#page-9-0) that the answer to Question [2](#page-2-1) is positive for a wide class of sparse polynomial programs having strongly admissible monomial support (see Definition [3.4\)](#page-8-0). In particular, our results show that the BKK bound is tight for Example [1.1.](#page-1-0) Further, in Corollary [3.6](#page-9-1) we show that algebraic degrees of generic sparse polynomial programs are determined by the Newton polytopes of F. Corollary [3.6](#page-9-1) also has algorithmic implications which were studied in [\[LMR23\]](#page-27-16).

As a corollary, we prove an analogous result to that in [\[BSW22\]](#page-26-4), and show that the ED degree of a variety defined by polynomials with strongly admissible support is equal to the BKK bound of its corresponding Lagrange system (Corollary [4.1\)](#page-12-0). We also prove similar results for (the sum of) polar degrees (Corollary [4.3\)](#page-13-0), giving the first convex geometric interpretation of this algebraic invariant.

In Theorem [C](#page-10-0) we loosen the assumption on \bf{F} of strongly admissible monomial support and provide a different formula for the algebraic degree of \bf{F} when \bf{F} is in a larger family of sparse polynomial programs. Our main tool here is Porteous' formula which computes the fundamental class of the degeneracy locus of a morphism between two vector bundles as a polynomial of their Chern classes. Using Porteous' formula, Theorem [C](#page-10-0) expresses the algebraic degree in terms of the intersection theory of a certain toric compactification of $(\mathbb{C}^*)^n$. The formula for the algebraic degree in Theorem [C](#page-10-0) can be expressed as a (not necessarily positive) linear combination of mixed volumes. However, the explicit connection to the mixed volume of the Lagrange system is still mysterious.

2. Preliminaries and notation

2.1. Sparse polynomials and polyhedral geometry. We specify a sparse poly*nomial* $f \in \mathbb{C}[y_1,\ldots,y_n]$ by its monomial support, $\mathcal{A} \subset \mathbb{N}^n := \mathbb{Z}_{\geq 0}^n$. Specifically, for a finite subset $\mathcal{A} \subset \mathbb{N}^n$, we write

$$
f = \sum_{\alpha \in \mathcal{A}} c_{\alpha} y^{\alpha}
$$

where $y^{\alpha} := y_1^{\alpha_1} \cdots y_n^{\alpha_n}$ and $c_{\alpha} \in \mathbb{C}$. Given a sparse polynomial $f \in \mathbb{C}[y_1, \ldots, y_n]$ with monomial support \mathcal{A} , the *Newton polytope of f* is defined as the convex hull of A. Explicitly, the Newton polytope of f is

$$
Newt(f) = conv\{\alpha : \alpha \in \mathcal{A}\}.
$$

A sparse polynomial system $\mathbf{F} = (f_0, \ldots, f_m)$ is defined by a tuple $\mathcal{A} = (\mathcal{A}_0, \ldots, \mathcal{A}_m)$ where $f_i = \sum_{\alpha \in A_i} c_{\alpha,i} y^{\alpha} \in \mathbb{C}[y_1, \ldots, y_n].$

Remark 2.1. In this paper, we consider *generic* sparse polynomial systems. A statement holds for a generic sparse polynomial system if it holds for all systems $\mathbf{F}=(f_0,\ldots,f_m)$ whose coefficients $\{c_{\alpha,i}\;:\alpha\in\mathcal{A}_i,\;0\leq i\leq m\}$ lie in some nonempty Zariski open subset of the space $\mathbb{C}^{\mathcal{A}_0} \times \cdots \times \mathbb{C}^{\mathcal{A}_m}$ of coefficients. This means that the non-generic behavior occurs on a set of measure zero in the space $\mathbb{C}^{\mathcal{A}_0} \times \cdots \times \mathbb{C}^{\mathcal{A}_m}$.

Given polytopes $P_1, \ldots, P_n \subset \mathbb{R}^n$, the mixed volume of P_1, \ldots, P_n is the coefficient of the monomial $\lambda_1 \cdots \lambda_n$ in the volume polynomial

$$
Vol_n(\lambda_1 P_1 + \ldots + \lambda_n P_n),
$$

where $P + Q = \{p + q : p \in P, q \in Q\}$ is the Minkowski sum and Vol_n is the standard n-dimensional Euclidean volume.

In a series of celebrated results [\[Ber75,](#page-26-9) [Kou76,](#page-27-17) [Kho78\]](#page-27-18) the connection between the number of solutions over \mathbb{C}^* to a system of sparse polynomial equations and the underlying convex geometry of the polynomials was made.

Theorem 2.2 (BKK Bound [\[Ber75,](#page-26-9) [Kou76,](#page-27-17) [Kho78\]](#page-27-18)). Let $\mathbf{F} = (f_1, \ldots, f_n)$ $\mathbb{C}[y_1,\ldots,y_n]$ be a sparse polynomial system with η isolated solutions in $(\mathbb{C}^*)^n$, counted with multiplicity and let $P_i = \text{Newt}(f_i)$. Then

$$
\eta \leq \mathrm{MV}(P_1,\ldots,P_n).
$$

Moreover, if the coefficients of **F** are generic then $\eta = MV(P_1, \ldots, P_n)$.

If for a sparse polynomial system \bf{F} the BKK bound holds with equality, we say F is BKK general. Bernstein gave explicit degeneracy conditions under which the above inequality is tight by considering the initial systems of \bf{F} [\[Ber75\]](#page-26-9).

Given a polytope $P \subset \mathbb{R}^n$ and a vector $w \in \mathbb{Z}^n \setminus \{0\}$, let P_w denote the face exposed by w . Specifically,

$$
P_w = \{ v \in P \; : \; \langle v, w \rangle \le \langle z, w \rangle \; \forall z \in P \}.
$$

For a sparse polynomial f we call

$$
\text{init}_{w}(f) = \sum_{\alpha \in (\text{Newt}(f))_w} c_{\alpha} y^{\alpha}
$$

the *initial polynomial* of f with respect to w. For a sparse polynomial system \mathbf{F} , we denote $\text{init}_w(\mathbf{F}) = (\text{init}_w(f_1), \dots, \text{init}_w(f_n)).$

Theorem 2.3. [\[Ber75,](#page-26-9) Theorem 2] Let $\mathbf{F} = (f_1, \ldots, f_n) \subset \mathbb{C}[y_1, \ldots, y_n]$ be a sparse polynomial system with η isolated \mathbb{C}^* solutions counted with multiplicity and let $P_i = \text{Newt}(f_i)$. All \mathbb{C}^* solutions of $F(y) = 0$ are isolated and $\eta = \text{MV}(P_1, \ldots, P_n)$ if and only if for every $w \in \mathbb{Z}^n \setminus \{0\}$, $\text{init}_w(\mathbf{F})$ has no \mathbb{C}^* solutions.

Theorem [2.2](#page-4-1) and Theorem [2.3](#page-4-2) demonstrate the intimate connection between solutions to systems of polynomial equations and polyhedral geometry. In the remainder of this section, we define a few more objects that are helpful when using this connection. Given $\mathcal{A} = (\mathcal{A}_0, \ldots, \mathcal{A}_m) \in \mathbb{N}^n$ we define the Cayley polytope of \mathcal{A} as

$$
Cay(\mathcal{A}) = conv(\{(z, e_i) : z \in \mathcal{A}_i, i = 0, \dots, m\}) \subset \mathbb{R}^{n+m}
$$

where e_i is the *i*-th standard basis vector of \mathbb{R}^m and e_0 is the vector of all zeroes. Similarly, for a sparse polynomial system **F** with support $\mathcal{A} = (\mathcal{A}_0, \dots, \mathcal{A}_m)$, we define $\text{Cav}(\mathbf{F}) = \text{Cav}(\mathcal{A}).$

For a face of a convex polytope, $F \subset P \subset \mathbb{R}^n$, the normal cone of F is the set of linear functionals which achieve their minimum on F:

$$
\sigma(F) = \{c \in \mathbb{R}^n : \langle c, x \rangle \le \langle c, z \rangle, \ \forall x \in F, \ z \in P\}.
$$

The normal cones of each face of P form a fan, denoted $\Sigma(P) \subset \mathbb{R}^n$.

Finally, in this paper we consider the operation of taking the "partial derivative" of a polytope which we define as follows. Let $P \subset \mathbb{R}_{\geq 0}^n$ be a polytope contained in the positive orthant. Then

$$
\partial_j P = (P - e_j) \cap \mathbb{R}_{\geq 0}^n = \{ \alpha - e_j \ : \ \alpha \in P, \ \alpha_j \geq 1 \},
$$

where e_j is the j-th standard basis vector of \mathbb{R}^n .

The definition of $\partial_i P$ is motivated by the partial differentiation operation of a polynomial $f \in \mathbb{R}[y_1,\ldots,y_n]$ with $\text{Newt}(f) = P$. Indeed, one always has $\text{Newt}(\frac{\partial}{\partial y_j}f) \subset$ ∂_j Newt(f). However, Newt($\frac{\partial}{\partial y_j} f$) and ∂_j Newt(f) can be quite different. For instance, let $f(y) = 1 + y^3$, then $Newt(f) = [0, 3]$ and $\partial Newt(f) = [0, 2]$. However, $\frac{\partial}{\partial y}f = 3y^2$ and Newt $(\frac{\partial}{\partial y}f) = \{2\}$. In general, even if P is integral, the polytope $\partial_i P$ does not have to be integral.

For a polynomial $f = \sum_{\alpha \in A} c_{\alpha} y^{\alpha}$ having full monomial support with respect to Newt(f) (i.e. $c_{\alpha} \neq 0$ for any $\alpha \in \text{Newt}(f) \cap \mathbb{N}^n$) the two constructions are connected in the following way:

$$
\text{Newt}\left(\frac{\partial}{\partial y_j}f\right) = \text{conv}\{\partial_j \text{Newt}(f) \cap \mathbb{N}^n\}.
$$

In particular, if f is a degree d polynomial with all monomials of degree $\leq d$ having non-zero coefficients, then

$$
\text{Newt}\left(\frac{\partial}{\partial y_j}f\right) = \partial_j \text{Newt}(f).
$$

2.2. Toric varieties. Theorem [2.2](#page-4-1) is a statement about intersection theory on toric varieties. A *toric variety* X is an irreducible variety such that $(\mathbb{C}^*)^n$ is a Zariski open subset of X and the action of $(\mathbb{C}^*)^n$ on itself extends to an action of $(\mathbb{C}^*)^n$ on X. We can also associate normal toric varieties with polyhedral fans.

Let $\sigma \in \mathbb{R}^n$ be a rational polyhedral cone which does not contain any vector subspace and denote

$$
S_{\sigma} = \sigma^{\vee} \cap \mathbb{Z}^n
$$

where $\sigma^{\vee} = \{y \in \mathbb{R}^n : \langle y, x \rangle \geq 0 \,\forall x \in \sigma\}$ is the dual cone of σ . Then the affine toric variety associated with σ is

$$
V_{\sigma} = \text{Spec}(\mathbb{C}[S_{\sigma}])
$$

where $\mathbb{C}[S_{\sigma}]$ is the semigroup algebra associated with S_{σ} .

Given a polyhedral fan Σ we have a collection of affine toric varieties indexed by cones in Σ , denoted $\{V_{\sigma} : \sigma \in \Sigma\}$. Specifically, for $\sigma, \tau \in \Sigma$ with $\sigma \subseteq \tau$ we have the inclusion $V_{\sigma} \subseteq V_{\tau}$. We denote X_{Σ} as the toric variety that is the colimit of all such inclusion maps. This means that X_{Σ} is obtained by gluing all affine varieties V_{τ} , $V_{\tau'}$ along V_{σ} , where $\sigma = \tau \cap \tau'$. For a more complete treatment of toric varieties, see [\[CLS11\]](#page-26-12).

In this paper, we will work with the *Cox ring* of a toric variety which is a generalization of the homogeneous coordinate ring of projective space and was introduced in [\[Cox95\]](#page-26-13). First, let us denote by $\Sigma(1)$ the set of all rays of Σ . By abuse of notation we often do not distinguish between rays ρ and their primitive ray generators. The Cox ring of X_{Σ} is

$$
S = \mathbb{C}[x_{\rho} : \rho \in \Sigma(1)].
$$

To every ray ρ we associate the corresponding torus invariant Weyl divisor D_{ρ} . Note that every torus invariant Weyl divisor D on X has a unique presentation of the form $D = \sum_{\rho \in \Sigma(1)} a_{\rho} D_{\rho}$. The global sections of the associated sheaf $\mathcal{O}_X(D)$ are spanned by monomials:

$$
H^{0}(\mathcal{O}_{X}(D), y) = \langle y^{m} \mid m \in \mathbb{Z}^{n}, \langle m, \rho \rangle \geq -a_{\rho} \rangle \subseteq \mathbb{C}[y^{\pm}_{1}, \dots, y^{\pm}_{n}].
$$

Given a global section $f = \sum_{m \in \mathbb{Z}^n} c_m y^m$ of $\mathcal{O}_X(D)$ we define the homogenization f of f to be the following element of S :

(2.1)
$$
\widetilde{f} = \prod_{\rho \in \Sigma(1)} x_{\rho}^{a_{\rho}} f(z_1, \dots, z_n) = \sum_{m \in \mathbb{Z}^n} c_m \prod_{\rho \in \Sigma(1)} x_{\rho}^{\langle m, \rho \rangle + a_{\rho}}.
$$

Here the variables z_i are defined by $z_i = \prod x_i^{\rho_i}$. Note that a Laurent polynomial $f \in \mathbb{C}[y_1^{\pm},\ldots,y_n^{\pm}]$ can be a section of the sheaf $\mathcal{O}_X(D)$ for a certain choices of D, and the homogenization f depends on the choice of D .

Example 2.4. Consider the toric variety $\mathbb{P}^1 \times \mathbb{P}^1$, associated with the dual fan $\Sigma(P)$ of the square P. The four generators $x_{-e_1}, x_{e_1}, x_{-e_2}, x_{e_2}$ of the Cox ring S are in bijection to the four rays.

Homogenizing the bivariate polynomial $f = 1 + y_1 + y_2 + y_1y_2$ yields the bihomogeneous polynomial $f = x_{-e_1}x_{-e_2} + x_{e_1}x_{-e_2} + x_{-e_1}x_{e_2} + x_{e_1}x_{e_2}$.

2.3. Chern and Segre classes of vector bundles. The main ingredient of the intersection theoretic formulas for the algebraic degree of polynomial optimization problems given in [\[NR09\]](#page-27-1) is Porteous' formula. Porteous' formula computes the expected cohomology class of the degeneracy locus of maps of vector bundles. Loosely speaking, vector bundles are families of vector spaces that are parameterized by another space, and cohomology classes are algebraic invariants of topological spaces. In this paper, all vector spaces will be parameterized by algebraic varieties and the vector spaces will all have the same dimension, called the rank of the vector bundle.

To formulate Porteous' formula one needs to use the Chern and Segre classes, which are well-studied characteristic classes of vector bundles. Here we list some main properties of these classes. For a more detailed introduction we refer to [\[EH16\]](#page-27-19).

For a vector bundle $\mathcal E$ of rank r on a variety X of dimension d and any $i = 0, \ldots, d$, one can define its i-th Chern class as an element of the 2i-th cohomology class of $X, c_i(\mathcal{E}) \in H^{2i}(X)$. For any vector bundle \mathcal{E} , one has $c_0(\mathcal{E}) = 1$ and $c_i(\mathcal{E}) = 0$ for any $i > r$. We will denote by $c(\mathcal{E})$ the *total Chern class* of \mathcal{E} , that is

$$
c(\mathcal{E})=c_0(\mathcal{E})+\ldots+c_{\max(d,r)}(\mathcal{E}).
$$

The crucial property of total Chern classes, known as Whitney's formula, is that they are multiplicative with respect to taking direct sums of line bundles:

$$
c(\mathcal{E} \oplus F) = c(\mathcal{E}) \cdot c(\mathcal{F}).
$$

In what follows we will mostly work with vector bundles coming as a direct sum of line bundles $E = \mathcal{L}_1 \oplus \ldots \oplus \mathcal{L}_n$. By applying the Whitney formula to such vector bundles we obtain a convenient formula for their total Chern class:

$$
c(\mathcal{E}) = \prod_{i=1}^n (1 + c_1(\mathcal{L}_i)),
$$
 These are graded pieces: $c_k(\mathcal{E}) = \sum_{I \in \binom{[n]}{k}} \prod_{i \in I} c_1(\mathcal{L}_i).$

Finally, let us recall the definition of Segre classes. Note that, the total Chern class $c(\mathcal{E})$ is an invertible element in the cohomology ring of X as its 0-th degree part is equal to 1. Using this one defines a total Segre class $s(\mathcal{E})$ of a vector bundle $\mathcal E$ on X to be the inverse of the total Chern class of $\mathcal E$:

$$
s(\mathcal{E}) = c(\mathcal{E})^{-1}
$$

.

Individual Segre classes $s_i(\mathcal{E})$ are defined as homogeneous components of the total Segre class. Note that, unlike Chern classes, one could have non-trivial Segre class $s_i(\mathcal{E})$ even for $i > r$.

3. Statement of the main result

In this section, we give an overview of the main results of this paper and defer the proofs of Theorem [A](#page-9-2) and Theorem [C](#page-10-0) to Section [6.](#page-20-0) The results in this paper require certain assumptions on the monomial support of \bf{F} , which we define in this section. While computational experiments suggest that both Theorem [A](#page-9-2) and Theorem [B](#page-9-0) are true under milder assumptions, we show in Example [3.9](#page-10-1) that they are needed for Theorem [C.](#page-10-0)

Definition 3.1. We call a point configuration $A = (A_0, \ldots, A_m) \in \mathbb{N}^n$ admissible if for every $i = 0, \ldots, m$:

- (1) $\mathbb{R}^n_{\geq 0}$ is a cone in the normal fan of the polytope conv (\mathcal{A}_i) , and
- (2) conv (\mathcal{A}_i) meets every coordinate hyperplane of \mathbb{R}^n .

When passing to a toric compactification, for various technical reasons we need to ensure that the constraints f_1, \ldots, f_m define a variety with a smooth closure. This is guaranteed by the following notion of an appropriate toric variety.

Definition 3.2. Let X be a proper, normal toric variety with underlying polyhedral fan Σ in \mathbb{R}^n and $\mathcal{A} = (\mathcal{A}_0, \ldots, \mathcal{A}_m)$ an admissible point configuration. We call X *appropriate* for $\mathcal A$ if the following three properties hold:

(1) the normal fan $\Sigma(\text{conv}(\mathcal{A}_i))$ is refined by Σ for each $i = 0, \ldots, m$,

- (2) the fan Σ contains the non-negative orthant $\mathbb{R}^n_{\geq 0}$, and
- (3) for generic functions f_i with monomial support \mathcal{A}_i the closure of $\mathcal{V}(f_1, \ldots, f_m)$ in X is disjoint from the singular locus of X .

Remark 3.3. Note that for every admissible point configuration A , there exists a smooth appropriate toric variety X. To construct X consider the normal fan Σ' of the Minkowski sum $conv(\mathcal{A}_0) + \cdots + conv(\mathcal{A}_m)$. A resolution of singularities can be performed by subdividing each singular cone of Σ' , resulting in a smooth, complete polyhedral fan Σ which contains the positive orthant. For more details on toric resolution of singularities consider Chapter 11 of [\[CLS11\]](#page-26-12).

In the proof of Theorem [A](#page-9-2) and Theorem [C](#page-10-0) we consider a natural choice for the toric compactification X , given by the coarsest refinement of all normal fans of the Newton polytopes $\text{Newt}(f_0), \ldots, \text{Newt}(f_m)$. To state our main results, we need one final definition.

Definition 3.4. We call a point configuration $A = (A_0, \ldots, A_m) \in \mathbb{N}^n$ strongly admissible if it is admissible and if the toric variety

$$
X = X(\Sigma(\text{conv}(\mathcal{A}_0) + \cdots + \text{conv}(\mathcal{A}_m)))
$$

is appropriate for A . Here X is the toric variety associated with the common refinement of the normal fans $\Sigma(\text{conv}(\mathcal{A}_i)).$

We give examples for the above definitions.

Example 3.5. Consider the tetrahedron $S = \text{conv}(0, e_1, e_2, e_3) \subset \mathbb{R}^3$, the cube $C = \text{conv}(0, e_1, e_2, e_3, e_1 + e_2, e_1 + e_3, e_2 + e_3, e_1 + e_2 + e_3) \subset \mathbb{R}^3$ and the bipyramid $\mathcal{B} = \text{conv}(0, e_1, e_2, e_3, e_1 + e_2 + e_3) \subset \mathbb{R}^3.$

- The point configuration $(S, -S + (1, 1, 1))$ is not an admissible point configuration since the normal fan of $-\mathcal{S}$ does not contain $\mathbb{R}^3_{\geq 0}$ as a cone.
- The point configuration (S, \mathcal{C}) is a strongly admissible point configuration since it is admissible and the singular locus of the toric variety defined by $S + C$ is zero-dimensional.
- The point configuration (S, \mathcal{B}) is an admissible, but not strongly admissible, point configuration. In particular, the toric variety defined by $S + \mathcal{B}$ is not appropriate for (S, \mathcal{B}) . To see this consider the three-dimensional toric variety X defined by $S + \mathcal{B}$. The torus orbit X_{σ} defined by the cone $\sigma =$ $\mathbb{R}_+(-1,-1,1)+\mathbb{R}_+(-1,-1,-1)$, dual to the face conv $((2,1,1),(1,2,1))$ of $S + B$, is contained in the singular locus. Let now f_1 be a generic polynomial with Newton polytope S. Then the vanishing locus $\mathcal{V}(f)$ in X intersects X_{σ} .

The polytopes $S + \mathcal{B}$ and $S + \mathcal{C}$ are displayed below, with the faces that define singular torus orbits colored in green.

The next two results express the number of solutions of L_F as mixed volumes.

Theorem A. Let $F = (f_0, \ldots, f_m)$ be a generic sparse system of polynomials in $\mathbb{C}[x_1,\ldots,x_n]$ with strongly admissible support. Then the algebraic degree of sparse polynomial optimization of [\(POP\)](#page-0-0) is equal to

(3.1) MV (Newt(f_1), ..., Newt(f_m), ∂_1 Newt(Φ_F), ..., ∂_n Newt(Φ_F)).

Here Φ_F denotes the Lagrangian $\Phi_F(\lambda, y) := f_0 - \sum_{i=1}^m \lambda_i f_i$, as above. Note that the polytopes ∂_j Newt (Φ_F) may be strictly larger than the Newton polytopes Newt (ℓ_j) of the partial derivatives $\ell_j = \frac{\delta}{\partial i}$ $\frac{\partial}{\partial y_j} (f_0 - \sum_{i=1}^m \lambda_i f_i)$. The equality in Theo-rem [A](#page-9-2) shows that L_F is BKK general, giving the next theorem.

Theorem B. Under the assumptions of Theorem [A](#page-9-2) the Lagrange system L_F is BKK general. Further, all critical points are smooth and lie in the algebraic torus $(\mathbb{C}^*)^n$. The number of critical points is equal to the mixed volume

(3.2) MV (Newt
$$
(f_1),...,
$$
Newt $(f_m),$ Newt $(\ell_1),...,$ Newt $(\ell_n))$.

If \bf{F} is not generic then (3.2) is an upper bound to the number of isolated, smooth critical points of f_0 restricted to $\mathcal{V}(f_1, \ldots, f_m)$.

Proof. For every $j = 1, ..., n$ we have the inclusion $\text{Newt}(\ell_i) \subseteq \partial_j \text{Newt}(\Phi_F)$ of polytopes, showing the inequality $(3.2) \leq (3.1)$ $(3.2) \leq (3.1)$ $(3.2) \leq (3.1)$. On the other hand, by Theorem [A,](#page-9-2) the BKK bound [\(3.2\)](#page-9-3) of L_F constitutes an upper bound to [\(1.1\)](#page-1-1) and we obtain

$$
(1.1) \le (3.2) \le (3.1) = (1.1).
$$

By Lemma [6.4,](#page-22-0) all critical points are smooth and all critical points are in the torus. □

By Theorem [A](#page-9-2) the monomial supports A_0, \ldots, A_m contribute to the mixed volume [\(3.1\)](#page-9-4) only by means of their convex hulls. This gives the following corollary.

Corollary 3.6. Under the assumptions of Theorem [A](#page-9-2) the algebraic degree of the sparse polynomial optimization problem [\(POP\)](#page-0-0) and the mixed volume [\(3.2\)](#page-9-3) depend only on the Newton polytopes Newt(f_i) for $i = 0, \ldots, m$.

Remark 3.7. Corollary [3.6](#page-9-1) has algorithmic consequences if one wishes to numerically find all critical points to [\(POP\)](#page-0-0) (as opposed to counting them). We leverage this in [\[LMR23\]](#page-27-16) and efficiently compute polyhedral start systems for L_F by imposing maximal sparsity on the Newton polytopes of F.

Example 3.8. Recall the optimization problem [\(1.2\)](#page-2-2) in Example [1.1.](#page-1-0) Theorem [B](#page-9-0) shows that the algebraic degree of this problem is equal to the mixed volume of its Lagrange system. A property of mixed volumes is that if $P_1, \ldots, P_n \subset \mathbb{R}^n$ and $Q_1, \ldots, Q_m \subset \mathbb{R}^{n+m}$, then

$$
\mathrm{MV}(P_1,\ldots,P_n,Q_1,\ldots,Q_m)=\mathrm{MV}(P_1,\ldots,P_n)\cdot\mathrm{MV}(\pi(Q_1),\ldots,\pi(Q_m)),
$$

where $\pi : \mathbb{R}^{n+m} \to \mathbb{R}^m$ is the projection onto the last m coordinates. Observe that Newt $(\ell_1), \ldots, \text{Newt}(\ell_n) \subset \mathbb{R}^{n+1}$ have *n*th coordinate zero. Therefore,

 $MV(Newt(\ell_1), \ldots, Newt(\ell_n), Newt(f)) = MV(Newt(\ell_1), \ldots, Newt(\ell_n)) \cdot MV(\pi_n(Newt(f)))$

where $\pi_n : \mathbb{R}^{n+1} \to \mathbb{R}$ is the projection onto the *n*th coordinate.

Since for $j = 1, ..., n$, Newt $(\ell_j) = \text{conv}(0, \alpha_j)$ for some $\alpha_j \in \mathbb{R}^n$, we can compute $MV(Newt(\ell_1), \ldots, Newt(\ell_n)) = det(M)$ where M is the matrix with jth column equal to α_j . In our case, this amounts to computing

$$
\det \left(\begin{bmatrix} 2 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ 0 & 0 & \ddots & 0 \\ 1 & 1 & \dots & 1 \end{bmatrix} \right) = 2.
$$

Finally, observe that $\pi_n(\text{Newt}(f)) = [0, 1]$ so it has (mixed) volume one. This gives a geometric proof that the optimization degree of (1.2) is 2, agreeing with the result we computed in Example [1.1.](#page-1-0)

Our final result, Theorem [C,](#page-10-0) weakens the assumption of strongly admissible support needed in Theorem [A](#page-9-2) and Theorem [B](#page-9-0) and characterizes the algebraic degree of F as a more general product in the Chow ring of a toric variety.

To formulate Theorem [C](#page-10-0) we first need some notation. We refer to Section [2](#page-3-1) and references therein for a brief introduction to the objects we use. Let $\mathcal{A} =$ $(\mathcal{A}_0, \ldots, \mathcal{A}_m)$ be an admissible point configuration (Definition [3.1\)](#page-7-0) and let X be a smooth toric variety given by the fan Σ which is appropriate for $\mathcal A$ (Definition [3.2\)](#page-7-1). As usual, the convex hull of each point configuration A_i defines a line bundle \mathcal{L}_{A_i} .

Further, since we assume that the fan Σ contains the positive orthant as one of its cones, we know that Σ contains the rays generated by the standard basis vectors e_1, \ldots, e_n of \mathbb{R}^n . We denote by D_{e_1}, \ldots, D_{e_n} the corresponding torus-invariant divisors on X and by $\mathcal{O}_X(D_{e_1}), \ldots, \mathcal{O}_X(D_{e_n})$ the corresponding line bundles. For further reading on toric line bundles, see [\[CLS11,](#page-26-12) Chapter 6].

Theorem C. Let $F = (f_0, \ldots, f_m)$ be a generic sparse system of polynomials in $\mathbb{C}[x_1,\ldots,x_n]$ with admissible support A and let X be as above. Then the algebraic degree of sparse polynomial optimization [\(POP\)](#page-0-0) is finite and equal to the degree of the following cycle class:

(3.3)
$$
c_1(\mathcal{L}_{\mathcal{A}_1}) \cdots c_1(\mathcal{L}_{\mathcal{A}_n}) \left(s(\mathcal{E}) c(\mathcal{F}) \right)_{n-m},
$$

where $\mathcal{E} = \mathcal{L}_{\mathcal{A}_0}^{-1} \oplus \cdots \oplus \mathcal{L}_{\mathcal{A}_m}^{-1}$ and $\mathcal{F} = \mathcal{O}_X(-D_{e_1}) \oplus \cdots \oplus \mathcal{O}_X(-D_{e_n}).$

Moreover, if \bf{F} is not generic then (3.3) is an upper bound to the number of isolated, smooth critical points of f_0 restricted to $\mathcal{V}(f_1, \ldots, f_m)$.

The purpose of the following example is to demonstrate that the assumptions for Theorem [C](#page-10-0) are necessary.

Example 3.9. Consider [\(POP\)](#page-0-0) where

$$
f_0 = 7 + 11y_1 - 13y_2 - 19y_1y_2 - 2y_1^2 - 5y_2^2
$$

$$
f_1 = -5y_1y_2 + 29y_1y_2^2 - 17y_1^2y_2 + 61y_1^2y_2^2 + y_1^2 - 3y_2^2.
$$

Note that the normal fan of Newt (f_1) does not contain $\mathbb{R}^2_{\geq 0}$, so the monomial support of (f_0, f_1) does not form an admissible point configuration, meaning the assumptions of Theorem [C](#page-10-0) do not hold in this case.

To evaluate [\(3.3\)](#page-10-2) we denote the Chern classes

$$
c_1(\mathcal{L}_{\mathcal{A}_0}) = [2D_2 + 4D_3 + 2D_4 - 2D_6] \text{ and } c_1(\mathcal{L}_{\mathcal{A}_1}) = [2D_2 + 2D_3 + 2D_4],
$$

and the vector bundles

$$
\mathcal{E} = \mathcal{L}_{\mathcal{A}_0}^{-1} \oplus \mathcal{L}_{\mathcal{A}_1}^{-1} \text{ and } \mathcal{F} = \mathcal{O}_X(D_1)^{-1} \oplus \mathcal{O}_X(D_5)^{-1}.
$$

Here, X is the smooth toric variety defined by the complete fan Σ with ray generators

$$
\rho_1 = (0, 1), \ \rho_2 = (1, 1), \ \rho_3 = (1, 0), \ \rho_4 = (0, -1), \ \rho_5 = (-1, -1), \ \rho_6 = (-1, 0).
$$

We have $c_1(\mathcal{F}) = [-D_1 - D_5]$, and

$$
s_1(\mathcal{E}) = \frac{-1}{c_1(\mathcal{E})} = c_1(\mathcal{L}_{\mathcal{A}_0}) + c_1(\mathcal{L}_{\mathcal{A}_1}) = [4D_2 + 6D_3 + 4D_4 - 2D_6].
$$

Finally, direct computation shows

$$
\deg (c_1(\mathcal{L}_{\mathcal{A}_1}) \cdot (s(\mathcal{E}) c(\mathcal{F}))_1) = \deg (c_1(\mathcal{L}_{\mathcal{A}_1}) \cdot (c_1(\mathcal{F}) + s_1(\mathcal{E}))) = 12.
$$

In this case, [\(3.3\)](#page-10-2) gives 12, while the algebraic degree of $\mathbf{F} = (f_0, f_1)$ is 10. In particular, the this shows the assumptions of Theorem [C](#page-10-0) are necessary. On the other hand, the [B](#page-9-0)KK bound of L_F is 10, so the bound in Theorem [A](#page-9-2) and Theorem B is tight. Although the assumptions in Theorem [A](#page-9-2) and Theorem [B](#page-9-0) are stronger than the ones in Theorem [C,](#page-10-0) we believe that Theorem [A](#page-9-2) and Theorem [B](#page-9-0) hold in greater generality than Theorem [C.](#page-10-0)

4. Sparse ED, polar and sectional degrees

In this section, we discuss important corollaries of Theorem [A](#page-9-2) and Theorem [B](#page-9-0) which relate Euclidean distance optimization, polar degrees, and sectional degrees to mixed volumes.

We first consider polynomial optimization problems where the objective function is $f_0 = ||y - u||_2^2$ for a generic point $u \in \mathbb{R}^n$. Let (f_1, \ldots, f_m) be a sparse polynomial system with variety $Y = \mathcal{V}(f_1, \ldots, f_m) \subset \mathbb{C}^n$. The *ED degree* of Y is the number of complex critical points of the optimization problem:

(ED)
$$
\min_{y \in \mathbb{R}^n} \|y - u\|_2^2 \text{ subject to } f_1(y) = \ldots = f_m(y) = 0.
$$

Equivalently, it is the algebraic degree of $\mathbf{F} = (f_0, \ldots, f_m)$. This brings us to the main result of this section, which relates ED degrees and mixed volumes. To simplify notation, for polynomials $\mathbf{F} = (f_1, \ldots, f_m)$ we write

$$
\mathrm{MV}(F) := \mathrm{MV}(\mathrm{Newt}(f_1), \ldots, \mathrm{Newt}(f_m)).
$$

Corollary 4.1 (Euclidean distance objective function). Let (f_1, \ldots, f_m) be a generic sparse polynomial system with variety $V(f_1, \ldots, f_m) = Y \subset \mathbb{C}^n$ and $f_0 = ||y - u||_2^2$ where u is a generic point in \mathbb{R}^n . If the monomial support of $\mathbf{F} = (f_0, \ldots, f_m)$ is strongly admissible, then

$$
MV(\mathbf{L}_F) = ED \ degree(Y).
$$

Proof. Consider the *weighted Euclidean distance function*, $f_C = ||Cy - u||_2^2$, where C is an $n \times n$ diagonal matrix with generic entries and define $\mathbf{F}_C = (f_C, f_1, \ldots, f_m)$. Theorem [A](#page-9-2) implies that the degree and mixed volume of L_{F_C} are equal. Call this value η .

Observe that the variety of L_{F_c} is in bijection with the critical points of

(4.1)
$$
\min_{y \in \mathbb{R}^n} \|y - u\|_2^2 \quad \text{subject to} \quad f_1(C^{-1}y) = \dots = f_m(C^{-1}y) = 0.
$$

This gives that there are η critical points to [\(4.1\)](#page-12-1). Notice that the monomial support of $(f_1(C^{-1}(y)), \ldots, f_m(C^{-1}(y))$ is the same as that of $(f_1(y), \ldots, f_m(y)),$ since C is diagonal. Therefore, the degree of L_F is equal to its mixed volume. \Box

Now that we have established a relationship between ED degrees and mixed volumes, we next recall that in $[DHO+16]$ $[DHO+16]$ a relationship between ED degrees and polar degrees was established. Let $\overline{Y} \subset \mathbb{P}^n$ be a projective variety. For a point $y \in \overline{Y}$, denote $T_x Y$ as the tangent space of Y at y. Denote the conormal variety of Y as

(4.2)
$$
\mathcal{N}_{\overline{Y}} = \overline{\{(y,z) \in \mathbb{P}^n \times (\mathbb{P}^n)^* \; : \; y \in \overline{Y}_{sm}, \; z \perp T_y \overline{Y}\},\}
$$

where $(\mathbb{P}^n)^*$ is the *dual projective space*, which is the projective space consisting of all hyperplanes in \mathbb{P}^n . For a proper subvariety, $\overline{Y} \subset \mathbb{P}^n$, the dimension of $\mathcal{N}_{\overline{Y}}$ is independent of the dimension of Y and $\dim(\mathcal{N}_{\overline{Y}}) = n - 1$.

For an irreducible variety $\overline{Y} \subset \mathbb{P}^n$, the *i*-th polar degree of \overline{Y} is

$$
\delta_i(\overline{Y}) = |\mathcal{N}_{\overline{Y}} \cap (L_1 \times L_2)|,
$$

where $L_1 \subseteq \mathbb{P}^n$ is a generic linear space of dimension $n + 1 - i$ and $L_2 \subseteq (\mathbb{P}^n)^*$ is a generic linear space of dimension i. The intersection $\mathcal{N}_X \cap (L_1 \times L_2)$ is finite, since

$$
\dim(\mathcal{N}_{\overline{Y}}) + \dim(L_1 \times L_2) = 2n = \dim(\mathbb{P}^n \times (\mathbb{P}^n)^*).
$$

While in our situation, the requirement that \overline{Y} is an irreducible projective variety does not typically hold, we remark that by considering an affine variety $Y \subset \mathbb{C}^n$ defined by polynomial equations with strongly admissible support, we can simply consider the *projective closure* of Y, which we denote \overline{Y} . Under mild genericity conditions, the projective closure of Y is defined by homogenizing the defining equations of Y with respect to a new variable, y_0 .

In general the ED degree of Y need not be equal to that of \overline{Y} [\[DHO](#page-26-0)⁺16, Example 6.6], but under certain transversality conditions they are. Given two varieties $V, W \subset$ \mathbb{P}^n , we say that the the intersection $V \cap W$ is *transversal* if the scheme theoretic

intersection $V \cap W$ is smooth. Let $H_{\infty} = \mathbb{P}^n \setminus \mathbb{C}^n = \mathcal{V}(y_0)$ denote the hyperplane at infinity and denote $Y_{\infty} = \overline{Y} \cap H_{\infty}$ and $Q_{\infty} = \{y_0^2 + \ldots + y_n^2 = 0\}.$

Theorem 4.2 (Theorem 6.11 [\[DHO](#page-26-0)⁺16]). Let $Y \subset \mathbb{C}^n$ be an irreducible, affine variety and $\overline{Y} \subset \mathbb{P}^n$ its projective closure. Assume that the intersections $\overline{Y} \cap H_{\infty}$ and $Y_\infty \cap Q_\infty$ are both transversal. Then,

$$
ED \text{ Degree}(Y) = \delta_0(\overline{Y}) + \delta_1(\overline{Y}) + \cdots + \delta_{n-1}(\overline{Y})
$$

where $\delta_i(\overline{Y})$ is the *i*-th polar degree of \overline{Y} .

As a consequence of Corollary [4.1](#page-12-0) and Theorem [4.2](#page-13-1) we establish a relationship between polar degrees and mixed volumes. To our knowledge, this is the first time a connection between convex geometry and polar degrees has been made.

Corollary 4.3. Let $Y \subset \mathbb{C}^n$ be an irreducible affine variety defined by polynomials (f_1, \ldots, f_m) and let $\overline{Y} \subset \mathbb{P}^n$ be its projective closure. Assume that the intersections $Y_{\infty} = \overline{Y} \cap H_{\infty}$ and $Y_{\infty} \cap Q_{\infty}$ are both transversal. Let L_F be the Lagrange system of $\mathbf{F} = (f_0, \ldots, f_m)$ where $f_0 = ||y - u||_2^2$ for generic $u \in \mathbb{R}^n$ and \mathbf{F} has strongly admissible support. Then

$$
MV(\mathbf{L}_F) = \delta_0(\overline{Y}) + \cdots + \delta_{n-1}(\overline{Y}),
$$

where $\delta_i(\overline{Y})$ is the *i*-th polar degree of the projective closure \overline{Y} .

Example 4.4. Consider the Euclidean distance optimization problem

$$
\min_{y \in \mathbb{R}^2} \left\| \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} - \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\|_2^2 \quad \text{subject to} \quad 4y_1^2 + 2y_2^2 - y_1y_2 - 1 = 0.
$$

The Lagrange system of this optimization problem is

$$
\mathbf{L}_{\mathrm{F}} = (2(y_1 - 1) - \lambda(8y_1 - y_2), 2(y_2 - 1) - \lambda(-y_1 + 4y_2), 4y_1^2 + 2y_2^2 - y_1y_2 - 1).
$$

The mixed volume of \mathbf{L}_{F} is four and there are indeed four complex solutions (y_1, y_2, λ) , two of which are real:

$$
[0.3864, 0.5557, -0.4840],
$$

\n $[-0.3050, -0.6418, 1.4516],$
\n $[-0.1407 - 1.6318i, 2.3431 - 0.5950i, 0.2904 - 0.1023i],$
\n $[-0.1407 + 1.6318i, 2.3431 + 0.5950i, 0.2904 + 0.1023i]$

Now we consider the projective closure of the variety

$$
Y = \mathcal{V}(4y_1^2 + 2y_2^2 - y_1y_2 - 1) \subset \mathbb{C}^2
$$

which is defined as

$$
\overline{Y} = \mathcal{V}(4y_1^2 + 2y_2^2 - y_1y_2 - y_0^2) \subset \mathbb{P}^2.
$$

The conormal variety of \overline{Y} , $\mathcal{N}_{\overline{Y}} \subset \mathbb{P}^2 \times (\mathbb{P}^2)^*$, is defined as

$$
\mathcal{N}_{\overline{Y}} = \{([y_0:y_1:y_2],[z_0:z_1:z_2]) \in \mathbb{P}^2 \times (\mathbb{P}^2)^* : f_1 = f_2 = f_3 = f_4 = f_5 = f_6 = 0\}
$$

FIGURE 1. The ellipse $4y_1^2 + 2y_2^2 - y_1y_2 - 1 = 0$ along with the critical points (green) of the Euclidean distance problem from the point $(1, 1)$ (blue).

where the polynomials $f_1, f_2, f_3, f_4, f_5, f_6$ are

$$
f_1 = -y_0^2 + 4y_1^2 - y_1y_2 + 2y_2^2
$$

\n
$$
f_2 = y_1z_1 - 4y_2z_1 + 8y_1z_2 - y_2z_2
$$

\n
$$
f_3 = 31z_0^2 - 8z_1^2 - 4z_1z_2 - 16z_2^2
$$

\n
$$
f_4 = 31y_2z_0 + 2y_0z_1 + 16y_0z_2
$$

\n
$$
f_5 = 31y_1z_0 + 8y_0z_1 + 2y_0z_2
$$

\n
$$
f_6 = y_0z_0 + 4y_2z_1 - 8y_1z_2 + 2y_2z_2.
$$

Direct computation shows that $\delta_0(\overline{Y}) = 2$ and $\delta_1(\overline{Y}) = 2$, and that the ED degree of Y equals the sum of the polar degrees of its projective closure as expected.

We conclude this section by making a final connection to sectional degrees. The notion of *sectional degrees* was recently studied in [\[MRWW24\]](#page-27-20). Given an affine variety $Y \subset \mathbb{C}^n$, the *i-th sectional degree* of Y, denoted $s_i(Y)$, is defined as the algebraic degree of the optimization problem

(SO_i)
$$
\min_{y \in \mathbb{R}^n} \langle u, y \rangle \quad \text{subject to} \quad y \in Y \cap H_1 \cap H_2 \cap \dots \cap H_i
$$

where $\langle u, \cdot \rangle$ is a generic linear function and H_1, \ldots, H_i are generic affine linear hyperplanes. As an immediate consequence of Theorem [B](#page-9-0) we have a convex algebraic interpretation of $s_i(Y)$.

Corollary 4.5. Let $Y \subset \mathbb{C}^n$ be an affine variety defined by generic polynomials (f_1,\ldots,f_m) where for generic $u \in \mathbb{R}^n$, $\mathbf{F} = (\langle u, y \rangle, f_1, \ldots, f_m)$ has strongly admissible support. Let L_F be the Lagrange system of F corresponding to the sectional optimization problem (SO_i) (SO_i) . Then

$$
\mathrm{MV}(\mathbf{L}_\mathrm{F}) = s_i(Y)
$$

where $s_i(Y)$ is the *i*-th sectional degree of Y.

Furthermore, [\[MRWW24\]](#page-27-20) gives a condition where sectional and polar degrees agree. This condition relies on the *dual variety* of $\overline{Y} \subseteq \mathbb{P}^n$ which is defined as

$$
\overline{Y}^{\vee} = \pi_2(\mathcal{N}_{\overline{Y}})
$$

where $\pi_2 : \mathbb{P}^n \times (\mathbb{P}^n)^* \to (\mathbb{P}^n)^*$ is the projection onto the second coordinate and $\mathcal{N}_{\overline{Y}}$ is the conormal variety of \overline{Y} as defined in [\(4.2\)](#page-12-2).

Specifically, [\[MRWW24,](#page-27-20) Corollary 6.8] states that if $Y \subset \mathbb{C}^n$ is an affine variety with projective closure $\overline{Y} \subset \mathbb{P}^n$ such that H_{∞} is not contained in \overline{Y}^{\vee} , then $s_i(Y) =$ $\delta_i(\overline{Y})$ for all $0 \leq i \leq \dim(Y)$. Therefore, as a corollary of this, Corollary [4.3](#page-13-0) and Corollary [4.5](#page-14-1) we get the following equality of mixed volumes.

Corollary 4.6. Let $Y \subset \mathbb{C}^n$ be an affine variety defined by polynomials (f_1, \ldots, f_m) such that for generic $u \in \mathbb{R}^n$ the polynomial systems

$$
\mathbf{E} = (\|y - u\|_2^2, f_1 \dots, f_m), \quad and
$$

$$
\mathbf{F} = (\langle u, y \rangle, f_1, \dots, f_m)
$$

have strongly admissible support. Let L_E be the Lagrange system of **E** corresponding to the Euclidean distance optimization problem [\(ED\)](#page-11-0) and \mathbf{L}_{F_i} the Lagrange system of \mathbf{F}_i corresponding to the *i*-th sectional optimization problem [\(SO](#page-14-0)_i). Assume that H_{∞} is not contained in the dual variety of \overline{Y} . Then

$$
\mathrm{MV}(\mathbf{L}_{\mathrm{E}}) = \sum_{i=0}^{n-1} \mathrm{MV}(\mathbf{L}_{\mathrm{F}_i}).
$$

Observe that under certain genericity conditions, the results in [\[MRWW24\]](#page-27-20) show that sectional degrees are the affine analog of polar degrees. With this in mind and the aforementioned results, we have the following conjecture.

Conjecture 7. Let $Y \subseteq \mathbb{C}^n$ be an irreducible, affine variety and $\overline{Y} \subset \mathbb{P}^n$ its projective closure. If \overline{Y} intersects Q_{∞} transversely then the ED degree of Y is equal to $s_0(Y) + \ldots + s_{n-1}(Y)$.

To provide one piece of evidence for Conjecture [7](#page-15-0) and highlight its distinction from Theorem [4.2,](#page-13-1) we give an example of a variety Y where Conjecture [7](#page-15-0) is true but Theorem [4.2](#page-13-1) gives a strict upper bound on the ED degree of Y .

Example 4.8. Consider the affine variety $Y = \mathcal{V}(y_1^2 - y_2) \subset \mathbb{R}^2$. We can directly compute the ED degree of Y to be three. The sectional degrees of Y are $s_0(Y) = 1$ and $s_1(Y) = 2$. In this case Conjecture [7](#page-15-0) holds.

Conversely, we can consider the polar degrees of $\overline{Y} = \mathcal{V}(y_1^2 - y_2y_0)$ and compute that $\delta_0(\overline{Y}) = 2$ and $\delta_1(\overline{Y}) = 2$. This provides an example where the sum of the polar degrees of \overline{Y} is a strict upper bound on the ED degree of Y but the sum of the sectional degrees is exact.

5. Homogeneous equations for critical points

In this section, we define homogeneous critical point equations for the optimization problem [\(POP\)](#page-0-0). We give two different sets of critical point equations for [\(POP\)](#page-0-0). On the one hand, in [\[NR09\]](#page-27-1) critical points are characterized as an intersection of the vanishing locus of homogeneous equations $\{\widetilde{f}_1 = \cdots = \widetilde{f}_m = 0\}$ with a projective determinantal variety W . We generalize this approach by replacing projective

space with an appropriate toric variety X . On the other hand, we homogenize the Lagrange equations $\mathbf{L}_{\mathbf{F}} = (f_1, \ldots, f_m, \ell_1, \ldots, \ell_n)$ in the Cox ring of a toric variety, $\mathbb{P}(\mathcal{E})$, which we introduce now. We show that both approaches define the desired critical point equations in Lemma [5.8](#page-19-0) with [\(5.2\)](#page-19-1) concerning the former approach and [\(5.3\)](#page-19-2) the latter.

5.1. Toric projective bundles. We now describe the toric structure on the projectivization of a direct sum of line bundles on toric varieties. Let X be a complete toric variety given by a fan Σ and let $\mathcal{E} = \mathcal{L}_0 \oplus \ldots \oplus \mathcal{L}_m$ be a direct sum of line bundles on X . In this subsection, we will describe the fan of the total space of the projectivization $\mathbb{P}(\mathcal{E})$. For a more general study of line bundles with toric variety fiber see [\[HKM23\]](#page-27-21).

Lemma 5.1. Let X be a toric variety and let $\mathcal{E} = \mathcal{L}_0 \oplus \ldots \oplus \mathcal{L}_m$ be a direct sum of line bundles. The total spaces of $\mathcal E$ and $\mathbb P(\mathcal E)$ can be given the structure of a toric variety.

Proof. For every line bundle \mathcal{L}_i , there exists a torus invariant divisor D_i such that \mathcal{L}_i is isomorphic to $\mathcal{O}(D_i)$. Therefore, each line bundle \mathcal{L}_i on X can be equipped with an equivariant structure, i.e. the action of T on the total space of \mathcal{L}_i where T is the torus acting on X . This makes the projection map equivariant.

By fixing an equivariant structure on each of the line bundles \mathcal{L}_i , we obtain a T-action on the total space of $\mathcal E$. Finally, we extend the T-action on $\mathcal E$ to the action of $T \times (\mathbb{C}^*)^{m+1}$ by making the second component act fiberwise in a natural way. This action is faithful and has an open-dense orbit in the total space of \mathcal{E} .

Moreover, the action of $T \times (\mathbb{C}^*)^{m+1}$ on $\mathcal E$ descends to an action on $\mathbb P(\mathcal E)$. The latter action has a one-dimensional kernel given by the diagonal subtorus in $(\mathbb{C}^*)^{m+1}$. Hence $\mathbb{P}(\mathcal{E})$ has the structure of a toric variety with respect to the factor torus

$$
T \times ((\mathbb{C}^*)^{m+1}/\mathbb{C}^* \cdot (1,\ldots,1)). \square
$$

Remark 5.2. Note that the divisor D_i is defined up to addition of the principal divisor $div(u)$ of character $u \in \mathbb{Z}^n$ or, equivalently, any two equivariant structures on \mathcal{L}_i differ by the action of a character of T. Therefore, the toric varieties defined by different choices of D_i are isomorphic.

We conclude by describing the defining fan of the projectivized total space $\mathbb{P}(\mathcal{E})$, when the defining line bundles of $\mathcal E$ are torus equivariant. More precisely, for $0 \leq$ $j \leq m$, we denote D_j as a torus invariant divisor such that $\mathcal{L}_i = \mathcal{O}(D_i)$. Each divisor D_i defines a cone-wise linear function

$$
\psi_i \colon \mathbb{R}^n = |\Sigma| \to \mathbb{R}.
$$

Let $\Psi = (\psi_0, \dots, \psi_m) : |\Sigma| \to \mathbb{R}^{m+1}$ be the corresponding piece-wise linear map.

Denote $\widetilde{\Sigma} \subset \mathbb{R}^n \times \mathbb{R}^{m+1}$ as the fan that is obtained as the graph of the function Ψ. That is, $\widetilde{\Sigma}$ consists of cones $\widetilde{\sigma}$ where

$$
\widetilde{\sigma} = \{(x, \Psi(x)) \, | \, x \in \sigma\}, \text{ for } \sigma \in \Sigma.
$$

We now abuse notation and denote $\mathbb{R}^n_{\geq 0}$ as the fan supported on the positive orthant, whose cones are all of the form $\sigma_j = \{x \in \mathbb{R}^n_{\geq 0}, \mid \forall j \in J : x_j = 0\}, \ J \subseteq [n].$ The fan defining the total space of $\mathcal E$ consists of cones

$$
\widetilde{\sigma} + \tau \text{ for } \sigma \in \Sigma, \tau \in \mathbb{R}_{\geq 0}^n \times \{0\}.
$$

Similarly, the fan F defining the total space of $\mathcal E$ with the zero section removed is given by

$$
\widetilde{\sigma} + \tau \text{ for } \sigma \in \Sigma, \tau \in \partial \mathbb{R}^n_{\geq 0},
$$

where $\partial \mathbb{R}^n_{\geq 0} = \mathbb{R}^n_{\geq 0} \setminus {\mathbb{R}^n_{\geq 0}}$ denotes the fan consisting of all cones in $\mathbb{R}^n_{\geq 0}$, except for the one of dimension n. Finally, let $S_0 \subset \mathcal{E}$ be the image of the zero section of \mathcal{E} . Now \mathcal{S}_0 is a torus invariant subset of \mathcal{E} and thus the natural projection of $\mathcal{E} \setminus \mathcal{S}_0 \to \mathbb{P}(\mathcal{E})$ is a toric morphism. On the level of fans, consider the projection

$$
\tau \colon \mathbb{R}^n \times \mathbb{R}^{m+1} \longrightarrow \mathbb{R}^n \times (\mathbb{R}^{m+1}/\mathbb{R} \cdot (1,\ldots,1)) \cong \mathbb{R}^n \times \mathbb{R}^m.
$$

Now let S be the Cox ring of X, and let $S_{\mathcal{E}}$ be the Cox ring of $\mathbb{P}(\mathcal{E})$. By the above discussion, each ray of Σ is either of the form $\tilde{\rho}$, where ρ is a ray of Σ , or of the form $\{0\} \times e_i$, $i = 1, \ldots, m + 1$. This splits the generators of $S_{\mathcal{E}}$ over $\mathbb C$ into two groups. The first group of generators $x_{\tilde{\rho}}$ is bijective to the generators x_{ρ} of S. We denote members of the second group by $\lambda_i = x_{\{0\} \times e_{i+1}}$ and obtain the following proposition.

Proposition 5.3. The Cox ring $S_{\mathcal{E}}$ is isomorphic to the free S-algebra $S[\lambda_0, \ldots, \lambda_m]$.

Remark 5.4. In the following, we are often in the situation where f is a global section of a torus invariant line bundle $\mathcal{O}_X(D)$ on X. Now f denotes an element of $S \subseteq S_{\mathcal{E}}$. At the same time, f can be identified with a section of the bundle $\pi^* \mathcal{O}_X(D)$ on $\mathbb{P}(\mathcal{E})$, where $\pi : \mathbb{P}(\mathcal{E}) \longrightarrow X$ is the natural projection. When homogenizing, this gives rise to another element $\tilde{f} \in S_{\mathcal{E}}$. Direct computation shows that there is no need for disambiguation since both expressions are equal.

5.2. Constructing critical point equations in Cox rings. We start by fixing some notation and definitions. For the rest of this section, let $\mathbf{F} = (f_0, \ldots, f_m)$ be a generic sparse system of polynomials in $\mathbb{C}[y_1, \ldots, y_n]$ with admissible support $\mathcal{A} = (\mathcal{A}_0, \ldots, \mathcal{A}_m)$. Furthermore, X denotes a toric variety that is appropriate for $\mathcal A$, with fan Σ .

Remark 5.5. Note that, since Σ contains the positive orthant $\mathbb{R}^n_{\geq 0}$ as a cone, there is a distinct copy of the affine space \mathbb{C}^n contained in X. For clarity, we denote the affine variables in the coordinate ring $\mathbb{C}[y_1,\ldots,y_n]$ of \mathbb{C}^n as y_1,\ldots,y_n , while the generators of the Cox ring S of X as x_{ρ} . By slight abuse of notation, we will denote the element x_{e_j} in S by x_j for each $j = 1 \ldots, n$.

For every $i = 0, \ldots, m$ let $\mathcal{L}_i = \mathcal{O}(-D_{f_i})$ denote the dual line bundle associated with D_{f_i} . Here D_{f_i} is the torus invariant Weyl divisor on X, corresponding to the Newton polytope $\text{Newt}(f_i)$:

(5.1)
$$
D_{f_i} = \sum_{\rho \in \Sigma(1)} a_{\rho,i} D_{\rho}, \text{ where } a_{\rho,i} = -\min\{\langle m, \rho \rangle : m \in \text{Newt}(f_i)\}.
$$

We denote by $\mathcal E$ the vector bundle $\mathcal E = \mathcal L_0 \oplus \cdots \oplus \mathcal L_m$ with projectivized total space

$$
\mathbb{P}(\mathcal{E}) = \{ (x, [\lambda]) \mid x \in X, [\lambda] \in \mathbb{P}(\mathcal{E}(x)) \}.
$$

The rest of this section is devoted to giving two different, but related, systems of homogeneous critical point equations for (POP) , one in the Cox ring S of X, and one in the Cox ring $S_{\mathcal{E}}$ of $\mathbb{P}(\mathcal{E})$. On the one hand, critical points are characterized

by the vanishing of the Lagrange system L_F . It describes the intersection of the incidence variety

$$
Z^{\circ} := \{ (x, [\lambda]) \in \mathbb{C}^n \times \mathbb{P}^n : (\nabla f_0 \mid \dots \mid \nabla f_m) \lambda = 0 \}
$$

with the vanishing locus of f_1, \ldots, f_m . On the other hand, critical points are characterized by the Jacobian $(\nabla f_0 | \cdots | \nabla f_m)$ dropping rank. They form the intersection $V^{\circ} \cap W^{\circ}$, where $V^{\circ} := \mathcal{V}(f_1, \ldots, f_m) \subseteq \mathbb{C}^n$, and W° is the determinantal variety

$$
W^{\circ} := \{ x \in \mathbb{C}^n : \text{ rank} (\nabla f_0 \mid \dots \mid \nabla f_m) \le m \}.
$$

We proceed by giving homogeneous equations for V°, W° and Z° in S and $S_{\mathcal{E}}$ respectively. Every polynomial f_i is a global section of the line bundle $\mathcal{O}_X(D_{f_i}),$ and its homogeneous form can be written as

$$
\widetilde{f}_i = \sum_{m \in \text{Newt}(f_i) \cap \mathbb{Z}^n} c_{m,i} \prod_{\rho \in \Sigma(1)} x_{\rho}^{\langle m, \rho \rangle + a_{\rho, i}}.
$$

Here we homogenize f_i as in [\(2.1\)](#page-6-0) in Section [2.2.](#page-5-0) In particular, f_i is defined by our choice of line bundle $\mathcal{O}_X(D_{f_i})$.

We denote V as the closure of $V^{\circ} = \mathcal{V}(f_1, \ldots, f_m)$ in X. Observe that by genericity of \mathbf{F}, V is equal to the vanishing locus of the homogeneous equations $V = \mathcal{V}\left(\widetilde{f}_1, \ldots \widetilde{f}_m\right).$

We denote M as a homogeneous version of the Jacobian matrix:

$$
M = \left(\widetilde{\nabla} \widetilde{f}_0 \mid \cdots \mid \widetilde{\nabla} \widetilde{f}_m \right).
$$

Here $\widetilde{\nabla}$ denotes the vector $\left(\frac{\partial}{\partial x_1}, \ldots, \frac{\partial}{\partial x_n}\right)$ $\frac{\partial}{\partial x_n}$ ^T. We use the notation $\tilde{\nabla}$ instead of ∇ to indicate that we differentiate with respect to coordinates in the Cox ring. So M has columns $\left(\frac{\partial}{\partial x}\right)$ $\frac{\partial}{\partial x_1} \widetilde{f}_i, \ldots, \frac{\partial}{\partial x_n}$ $\left(\frac{\partial}{\partial x_n}\widetilde{f}_i\right)^T$. We define

$$
W \coloneqq \{ x \in X \; : \; \text{rank}(M) \le m \}
$$

to be the vanishing locus of the maximal minors of M . Furthermore, we let

$$
Z := \{(x, [\lambda]) \in \mathbb{P}(\mathcal{E}) \ : \ M(x)\lambda = 0\}
$$

be the associated incidence variety, contained in the projectivized total space $\mathbb{P}(\mathcal{E})$.

The rest of this Section is devoted to proving Lemma [5.8.](#page-19-0) It shows that the homogeneous critical point equations agree with the affine ones when restricted to affine space \mathbb{C}^n .

We need an observation about differentiating homogeneous polynomials. Let D be a torus invariant Weyl divisor on X (or on $\mathbb{P}(\mathcal{E})$), and f a global section of $\mathcal{O}_X(D)$. Observe that for $j = 1, \ldots, n$ the Newton polytope of the differential $y_j\frac{\partial}{\partial n}$ $\frac{\partial}{\partial y_j} f$ is contained in the rational polytope $e_j + \partial_j \text{Newt}(f)$, and in particular $\frac{\partial}{\partial y_j} f$ is a global section of the sheaf $\frac{1}{y_j}\mathcal{O}_X(D - D_{e_j})$ (or of the sheaf $\frac{1}{y_j}\mathcal{O}_{\mathbb{P}(\mathcal{E})}(D - D_{\tilde{e}_j})$). Direct computation shows the following proposition.

Proposition 5.6. Homogenization and differentiation commute: $\frac{\partial}{\partial y_j} f = \frac{\partial}{\partial x_j}$ $\frac{\partial}{\partial x_j} f$.

We denote $\widetilde{\Phi}_F$ as the homogenization of the Lagrangian, Φ , in the Cox ring $S_{\mathcal{E}}$. This makes sense since $\mathbb{P}(\tilde{\mathcal{E}})$ defines a global section of the sheaf $\mathcal{O}_{\mathbb{P}(\mathcal{E})}(D_{\Phi_F}),$

associated with the Cayley polytope $\text{Newt}(\Phi_F) = \text{Cay}(\text{Newt}(f_0), \ldots, \text{Newt}(f_m)).$ By Proposition [5.6,](#page-18-0) each of the defining equations

$$
0 = (M(x)\lambda)_j = \lambda_0 \frac{\partial}{\partial x_j} \widetilde{f}_0 + \dots + \lambda_m \frac{\partial}{\partial x_j} \widetilde{f}_m = \frac{\partial}{\partial x_j} \widetilde{\Phi}_F(\lambda, x)
$$

of Z is equal to the homogenization $\widetilde{\ell}_j$ of $\ell_j = \widetilde{\frac{\partial}{\partial y_j}} \widetilde{\Phi}_F(\lambda, x)$. Here $\widetilde{\ell}_j$ is considered as a global section of the sheaf $\mathcal{O}_{\mathbb{P}(\mathcal{E})}(D_{\Phi_F} - D_{\tilde{e}_j}).$

We denote $\widetilde{\mathbf{L}}_F = (\widetilde{f}_1, \ldots, \widetilde{f}_m, \widetilde{\ell}_1, \ldots, \widetilde{\ell}_n)$ as the homogenized Lagrange system. On one hand, we observed above that Z is equal to the vanishing locus of $\tilde{\ell}_1, \ldots, \tilde{\ell}_n$. On the other hand, the vanishing locus of $\widetilde{f}_1, \ldots, \widetilde{f}_m$ in $\mathbb{P}(\mathcal{E})$ is the preimage $\pi^{-1}(V)$ of the vanishing locus V of $\tilde{f}_1, \ldots, \tilde{f}_m$ in X. We obtain the following Proposition.

Proposition 5.7. The vanishing locus of $\widetilde{\mathbf{L}}_F$ in $\mathbb{P}(\mathcal{E})$ is the intersection $Z \cap \pi^{-1}(V)$.

The following lemma shows that the homogeneous critical point equations introduced in this chapter restrict on \mathbb{C}^n to the expected affine critical point equations.

Lemma 5.8. The following three equalities hold:

- $V \cap \mathbb{C}^n = V^\circ$, $W \cap \mathbb{C}^n = W^\circ$ (5.2)
- (5.3) $Z \cap \pi^{-1}(\mathbb{C}^n) = Z^\circ,$

where the intersection in [\(5.2\)](#page-19-1) is in X and the intersection in [\(5.3\)](#page-19-2) is in $\mathbb{P}(\mathcal{E})$.

Proof. The first of the equalities is clear, by the definition of V as the closure of V° . To see the second equality, we prove that the entries of M are homogenizations of the entries of the Jacobian $(\nabla f_0, \ldots, \nabla f_m)$. This is a direct consequence of Proposition [5.6,](#page-18-0) since for every $i = 0, \ldots, m$ and $j = 1, \ldots, n$, $\widetilde{\frac{\partial}{\partial y_j}} f_i = \frac{\delta}{\partial x_j}$ $\frac{\partial}{\partial x_j} f_i$. The third equality is analogous, since homogenizing the defining equations $\ell_1, \ldots \ell_n$ of Z° yields the defining equations $\widetilde{\ell}_1, \ldots \widetilde{\ell}_n$ of Z .

We close this section with the following generalization of Euler's equation.

Proposition 5.9. Let $D = \sum_{\rho \in \Sigma(1)} a_{\rho} D_{\rho}$ be a torus invariant Weyl divisor, f a global section of $\mathcal{O}_X(D)$ and $\tau \in \Sigma(1)$ a ray. Then the generalized Euler equation

(5.4)
$$
-x_{\tau} \frac{\partial}{\partial x_{\tau}} \widetilde{f} + \tau_1 x_1 \frac{\partial}{\partial x_1} \widetilde{f} + \cdots + \tau_n x_n \frac{\partial}{\partial x_n} \widetilde{f} = -a_{\tau} \widetilde{f}
$$

holds for the homogenization $\widetilde{f} \in S$ of f.

Proof. Equation (2.1) reads
$$
\tilde{f} = \sum_{m \in \mathbb{Z}^n} c_m \prod_{\rho \in \Sigma(1)} x_{\rho}^{\langle m, \rho \rangle + a_{\rho}}
$$
 and we have
\n
$$
- x_{\tau} \frac{\partial}{\partial x_{\tau}} \tilde{f} + \tau_1 x_1 \frac{\partial}{\partial x_1} \tilde{f} + \dots + \tau_n x_n \frac{\partial}{\partial x_n} \tilde{f}
$$
\n
$$
= \sum_{m \in \mathbb{Z}^n} c_m \left(-x_{\tau} \frac{\partial}{\partial x_{\tau}} + \tau_1 x_1 \frac{\partial}{\partial x_1} + \dots + x_n \frac{\partial}{\partial x_n} \right) \prod_{\rho \in \Sigma(1)} x_{\rho}^{\langle m, \rho \rangle + a_{\rho}}
$$
\n
$$
= \sum_{m \in \mathbb{Z}^n} c_m(-\langle m, \tau \rangle - a_{\tau} + m_1 + a_{e_1} + \dots + m_n + a_{e_n}) \prod_{\rho \in \Sigma(1)} x_{\rho}^{\langle m, \rho \rangle + a_{\rho}}
$$
\n
$$
= \sum_{m \in \mathbb{Z}^n} c_m(-a_{\tau}) \prod_{\rho \in \Sigma(1)} x_{\rho}^{\langle m, \rho \rangle + a_{\rho}} = -a_{\tau} \tilde{f}.
$$

6. Computing the number of critical points

In this section we prove Theorem [A](#page-9-2) and Theorem [C,](#page-10-0) relying on the results from Section [5.](#page-15-1) In Lemma [5.8](#page-19-0) we characterized critical points of [\(POP\)](#page-0-0) in two ways. First, in [\(5.2\)](#page-19-1) we characterized the critical points of [\(POP\)](#page-0-0) as an intersection $V \cap W$ in X. Second, we characterized the critical points of (POP) in (5.3) by means of homogenized Lagrange equations $\widetilde{\mathbf{L}}_F$ in the Cox ring of $\mathbb{P}(\mathcal{E})$. In this section, we show that all intersections are transversal and happen in \mathbb{C}^n . This characterizes the number of critical points as products of cohomology classes. In the case of Theorem [A](#page-9-2) this product is a mixed volume. The proof of Theorem [C](#page-10-0) relies on a characterization of [W] as a Porteous class.

Before diving into the proofs, we wish to highlight where the assumptions of our main theorem are used. We use the assumption that A be admissible in the proof of Proposition [6.2.](#page-21-0) For the proof of Proposition [6.1](#page-20-1) we need that the closure V of the constraint locus $V(f_1, \ldots, f_m)$ is smooth. This is guaranteed by the stronger assumption that A is strongly admissible in the proof of Theorem A and Theorem [B.](#page-9-0) For Theorem [C](#page-10-0) we assume X is smooth in order to employ Porteous' formula.

6.1. Preliminary results. We start by proving some technical statements that are needed for the desired transversality results. For the rest of the section we again fix the assumptions from Section 5.2 , and assume that V does not intersect the singular locus of X . It follows from Bertini's Theorem, that V is smooth, which is the motivation for Definition [3.4](#page-8-0) and Definition [3.2.](#page-7-1)

For the next proposition, we use the following notion. Similar to the projection $\mathbb{C}^{n+1}\setminus\{0\} \to \mathbb{P}^n$, there exists the open subset $U_{\Sigma} \subseteq \mathbb{C}^{\Sigma(1)}$ with a projection τ : $U_{\Sigma} \to X$. For a subvariety Z of X, we define the cone $C(Z)$ over Z to be the closure of the preimage $\tau^{-1}(Z)$ in $\mathbb{C}^{\Sigma(1)}$. The intersection $C(Z) \cap U_{\Sigma}$ forms a torus principal bundle over Z. In particular, $C(Z) \cap U_{\Sigma}$ is smooth if Z is smooth.

Proposition 6.1. The matrices $(\widetilde{\nabla}\widetilde{f}_0,\ldots,\widetilde{\nabla}\widetilde{f}_m)$ and $(\widetilde{\nabla}\widetilde{f}_1,\ldots,\widetilde{\nabla}\widetilde{f}_m)$ have full ranks $m + 1$ and m everywhere on $\mathcal{V}(\tilde{f}_0, \ldots, \tilde{f}_m)$ and $\mathcal{V}(\tilde{f}_1, \ldots, \tilde{f}_m)$ respectively.

Proof. The proof for the second matrix is analogous, so we only present the proof for

$$
M=\left(\widetilde{\nabla}\widetilde{f}_0,\ldots,\widetilde{\nabla}\widetilde{f}_m\right).
$$

Let $x \in \mathcal{V}(\tilde{f}_0, \ldots, \tilde{f}_m)$ be arbitrary and $\sigma \in \Sigma$ the unique cone such that x is contained in the torus orbit $O(\sigma)$. Let \widetilde{M} denote the matrix with rows

(6.1)
$$
\left(\frac{\partial}{\partial x_{\rho}}\tilde{f}_0,\ldots,\frac{\partial}{\partial x_{\rho}}\tilde{f}_m\right)
$$

for each ray ρ in $\Sigma(1)$. The left kernel of \widetilde{M} is the tangent space of the cone $C\left(\mathcal{V}\left(\widetilde{f}_0,\ldots,\widetilde{f}_m\right)\right)$ in $\mathbb{C}^{\Sigma(1)}$. The Jacobian \widetilde{M}_{σ} , of the cone over the variety

$$
O(\sigma) \cap \mathcal{V}\left(\widetilde{f}_0,\ldots,\widetilde{f}_m\right)
$$

is a submatrix of M. Its rows correspond to those rays ρ that are not contained in σ . By our assumption at the beginning of this section, V is disjoint from the singular

locus of X, and we can apply Bertini's Theorem to show that $\mathcal{V}(\tilde{f}_0,\ldots,\tilde{f}_m)$ is a smooth variety. Furthermore, the intersection $O(\sigma) \cap V(\tilde{f}_0, \ldots, \tilde{f}_m)$ is transversal by [\[Kho78\]](#page-27-18), so \widetilde{M}_{σ} is of full rank $m + 1$ at x. We now finish the proof by showing that the row span of M_{σ} is contained in the row span of M. Let ρ be any ray that is not contained in σ . To show that the corresponding row [\(6.1\)](#page-20-2) of M_{σ} is contained in the row span of M, we apply Proposition [5.9](#page-19-3) to all functions $\widetilde{f}_0, \ldots, \widetilde{f}_m$. The right side of equation (5.4) vanishes, and we obtain

$$
\begin{pmatrix} \rho_1 x_1 \\ \vdots \\ \rho_n x_n \end{pmatrix}^T M = x_{\rho} \begin{pmatrix} \frac{\partial}{\partial x_{\rho}} \widetilde{f}_0 \\ \vdots \\ \frac{\partial}{\partial x_{\rho}} \widetilde{f}_m \end{pmatrix}^T.
$$

Proposition 6.2. The gradient $\widetilde{\nabla}\widetilde{f}_0 = \left(\frac{\partial}{\partial x_0}\right)^{\frac{1}{2}}$ $\frac{\partial}{\partial x_j} \widetilde{f}_0\bigg)$ $does not vanish on any orbit.$

Proof. Towards a contradiction, we assume that there exists a cone $\sigma \in \Sigma$ such that for every $j = 1, ..., n$ the polynomial $\frac{\partial}{\partial x_j} \tilde{f}_0$ vanishes on the associated torus orbit $O(\sigma)$ of X. We denote $\frac{\partial}{\partial x_j} \tilde{f}_0 \Big|_{O(\sigma)}$ as the restriction of $\frac{\partial}{\partial x_j} \tilde{f}_0$ to the cone over $O(\sigma)$. It is obtained by substituting all variables x_{ρ} with zero, where ρ is contained in σ .

Now consider the face $\text{Newt}(f_0)^\sigma$ of $\text{Newt}(f_0)$ exposed by σ . For every lattice point m of $\text{Newt}(f_0)^\sigma$ the monomial

$$
\frac{\partial}{\partial x_j} \prod_{\rho \in \Sigma(1)} x_\rho^{\langle m, \rho \rangle + a_{\rho, 0}}, \quad a_{\rho, 0} = -\min \{ \langle m, \rho \rangle : m \in \text{Newt}(f_0) \}
$$

of $\frac{\partial}{\partial x_j} f_0$ only vanishes on $O(\sigma)$ if $m_j = 0$. In particular, the face Newt $(f_0)^\sigma$ can only contain the single element 0. By assumption [3.2](#page-7-1) on X , 0 is a smooth vertex of Newt (f_0) , and dual to the cone $\mathbb{R}^n_{\geq 0}$. Since σ reveals the vertex $\underline{0}$, it has to intersect the interior of the positive orthant $\mathbb{R}^n_{\geq 0}$ and in fact both cones are equal. This leaves us with the case where the torus orbit is {0}. But the gradient $\tilde{\nabla} \tilde{f}_0$ does not vanish
uniformly at 0 uniformly at 0 .

Let V and W denote the varieties from Section [5.2:](#page-17-0)

$$
V = \mathcal{V}\left(\widetilde{f}_1, \ldots, \widetilde{f}_m\right) \quad \text{and} \quad W = \{x \in X \; : \; \mathrm{rank}\left(\widetilde{\nabla}\widetilde{f}_0 \mid \cdots \mid \widetilde{\nabla}\widetilde{f}_m\right) \leq m\}.
$$

Proposition 6.3. The variety $V \cap W$ is of dimension zero.

Proof. Towards a contradiction we assume that there exists a torus orbit $O(\sigma)$, and a curve C such that C is contained in the intersection $W \cap V \cap O(\sigma)$. We denote by $\widetilde{f}_0\Big|_{O(\sigma)}$ the restriction of \widetilde{f}_0 to $O(\sigma)$. It is obtained by substituting all variables x_ρ with zero, where ρ is contained in σ . We now distinguish two cases: either $\widetilde{f}_0\Big|_{O(\sigma)}$ vanishes somewhere on $O(\sigma)$, or it is a scalar multiple of a monomial. In the first case $\widetilde{f}_0\Big|_{O(\sigma)}$ vanishes on C by genericity. In particular, the matrix M drops rank somewhere on the vanishing locus $\mathcal{V}(\tilde{f}_0,\ldots,\tilde{f}_m)$, contradicting Proposition [6.1.](#page-20-1)

In the second case, we now derive a contradiction from Proposition [6.1](#page-20-1) by showing that the matrix $(\tilde{\nabla} \tilde{f}_1, \ldots, \tilde{\nabla}, \tilde{f}_m)$ drops rank somewhere on C. Suppose $\tilde{f}_0\Big|_{O(\sigma)}$ is a monomial. Then each restriction $\frac{\partial}{\partial x_j} \widetilde{f}_0 \Big|_{O(\sigma)}$ is either a monomial or zero, and by Proposition [6.2](#page-21-0) there is an index $\ell = 1, \ldots, n$ such that $\frac{\partial}{\partial x_{\ell}} \widetilde{f}_0 \Big|_{O(\sigma)}$ is not zero. Without loss of generality, we assume $\ell = 1$. Consider the following matrix, M^* , obtained by subtracting from the j−th row of M the multiple

$$
\frac{\frac{\partial}{\partial x_j}\widetilde{f}_0}{\frac{\partial}{\partial x_1}\widetilde{f}_0}\left(\frac{\partial}{\partial x_1}\widetilde{f}_0,\ldots,\frac{\partial}{\partial x_1}\widetilde{f}_m\right)
$$

of the first row, for each $j = 2, \ldots, n$. This eliminates the first entry in all but the first row:

$$
M^* = \begin{bmatrix} \frac{\partial}{\partial x_1} \widetilde{f}_0 & \frac{\partial}{\partial x_1} \widetilde{f}_1 \cdots \frac{\partial}{\partial x_1} \widetilde{f}_m \\ 0 & \vdots & A \\ 0 & \end{bmatrix}.
$$

Since M drops rank everywhere on C and $\frac{\partial}{\partial x_1} \tilde{f}_0$ is not identically zero, A also drops rank on C. Let $\mu = (\mu_1, \ldots, \mu_n)^T$ be a vector of rational functions on $O(\sigma)$ satisfying $Au = 0$

everywhere on C. Since the expression $\mu_1 \frac{\delta}{\partial x}$ $\frac{\partial}{\partial x_1} f_1 + \cdots + \mu_n \frac{\partial}{\partial x_n}$ $\frac{\partial}{\partial x_n} f_1$, is not a monomial on $O(\sigma)$, it vanishes at some point x in C by genericity. This shows that $\mu(x)$ is in the right kernel of $(\tilde{\nabla} \tilde{f}_1(x), \ldots, \tilde{\nabla} \tilde{f}_m(x))$, finishing the proof. \Box

Lemma 6.4. The intersection $Z \cap V(\tilde{f}_1, \ldots, \tilde{f}_m)$ is transversal and contained in the big torus $(\mathbb{C}^*)^{n+m}$ in the toric variety $\mathbb{P}(\mathcal{E})$.

Proof. The image of $Z \cap \mathcal{V}(\widetilde{f}_1,\ldots,\widetilde{f}_m)$ under the natural projection $\pi : \mathbb{P}(\mathcal{E}) \longrightarrow X$ is $W \cap \mathcal{V}(\tilde{f}_1,\ldots,\tilde{f}_m)$, which by Proposition [6.3](#page-21-1) is finite. We prove below that π bijectively identifies both sets. In particular, the n defining equations of Z , given by $M(x)\lambda = 0$, form a complete intersection when restricted to $\mathcal{V}(\widetilde{f}_1, \ldots, \widetilde{f}_m)$.

To inductively apply Bertini to the equations $(M(x)\lambda)_j = 0$ we now show that, for varying coefficients of $f_0, Z \cap V(\tilde{f}_1, \ldots, \tilde{f}_m)$ defines a basepoint-free family of varieties on the vanishing locus $\mathcal{V}(\widetilde{f}_1,\ldots,\widetilde{f}_m)$ in $\mathbb{P}(\mathcal{E})$. To do this, we fix any element x of V and show that Z does not have a fixed point in the fiber $\pi^{-1}(x)$. By Proposition [6.1](#page-20-1) the last m columns $(\tilde{\nabla}\tilde{f}_1,\ldots,\tilde{\nabla}\tilde{f}_m)$ of M are linearly independent. In particular, varying the first column ∇f_0 changes the unique solution $[\lambda]$ to $M(x)\lambda = 0$. It now suffices to see that the gradient $\tilde{\nabla} \tilde{f}_0$ does not vanish uniformly at x, which by Proposition [6.2](#page-21-0) is true for generic coefficients of f_0 . To see that $Z \cap V(\tilde{f}_1, \ldots, \tilde{f}_m)$ is contained in the big torus, $(\mathbb{C}^*)^{n+m}$, orbit, we apply the same Bertini type argument

to show transversality of the intersection $O \cap \mathcal{V}(\tilde{f}_1,\ldots,\tilde{f}_m) \cap Z$. Here O denotes any torus orbit on $\mathbb{P}(\mathcal{E})$. For dimensional reasons, this intersection can only be nonempty for the big torus $(\mathbb{C}^*)^{n+m}$ orbit.

6.2. The proof of Theorem [A.](#page-9-2) The idea behind the proof of Theorem A is to study the system of homogenized Lagrange equations $\widetilde{\mathbf{L}}_F = (\widetilde{f}_1, \ldots, \widetilde{f}_m, \widetilde{\ell}_1, \ldots, \widetilde{\ell}_n)$. We show that it comprises global sections of \mathbb{Q} -Cartier divisors, that intersect transversely and away from infinity. This expresses the number of solutions as a product of Chern classes, which is a mixed volume.

For the rest of this subsection, we impose the assumptions of Theorem [A.](#page-9-2) Let Σ be the normal fan of the Minkowski sum of the polytopes $\text{Newt}(f_0), \ldots, \text{Newt}(f_n)$, and let X be the associated normal toric variety. Then the assumptions from the beginning of Section [5.2](#page-17-0) are fulfilled, since X is appropriate for A , and V does not intersect the singular locus of X.

Again, let Φ_F denote the Lagrangian $\Phi_F(\lambda, y) = f_0 - \sum_{i=1}^m \lambda_i f_i$, and ℓ_j the partial differentials $\frac{\partial}{\partial y_j}(f_0 - \sum_{i=1}^m \lambda_i f_i)$ of Φ_F . For each $j = 1, \ldots, n$, we define the homogenization $\tilde{\ell}_j$ of ℓ_j as a section of the divisor $D_{\Phi_F} - D_{\tilde{e}_j}$ on $\mathbb{P}(\mathcal{E})$. We want to prove that this divisor is the torus invariant divisor associated with a translate of the rational polytope ∂_i Newt (Φ_F) but in order to do that we need a few propositions.

Proposition 6.5. Let X be a toric variety and $\mathcal{E} = \bigoplus \mathcal{L}_{P_i}$ be a direct sum of line bundles on X. Then the the relative $\mathcal{O}(1)$ bundle of $\mathbb{P}(\mathcal{E})$ is represented by the Cayley polytope $\text{Cay}(P_1, \ldots, P_m)$.

Proof. The space of sections $H^0(\mathbb{P}(\mathcal{E}), \mathcal{O}(1))$ is canonically isomorphic to

$$
H^0(X, \mathcal{E}) = \bigoplus_i H^0(X, \mathcal{L}_{P_i}).
$$

Moreover, $H^0(X, \mathcal{L}_{P_i})$ is the weight space of $H^0(X, \mathcal{E})$ with respect to $(\mathbb{C}^*)^m$ acting fiber-wise on $\mathbb{P}(\mathcal{E})$ corresponding to the *i*-th basis vector of \mathbb{Z}^m . Since the weights of the base torus acting on $H^0(X, \mathcal{L}_{P_i})$ is given by the lattice points of P_i , we obtain the result. \Box

Proposition 6.6. Let σ be a cone in the normal fan $\Sigma(P_1 + \cdots + P_n)$. Then the face of $Cay(P_1, \ldots, P_n)$ exposed by $\tilde{\sigma}$ is equal to the Cayley polytope of the faces $P_1^{\sigma}, \ldots, P_n^{\sigma}$:

(6.2)
$$
\text{Cay}(P_1,\ldots,P_n)^{\widetilde{\sigma}} = \text{Cay}(P_1^{\sigma},\ldots,P_n^{\sigma}).
$$

Proof. This can be done by direct computation. A different argument relies on Proposition [6.5.](#page-23-0) For this, denote by X_{σ} the closure of the torus orbit of X_{Σ} corresponding to $\sigma \in \Sigma$. By Proposition [6.5,](#page-23-0) the equation [\(6.2\)](#page-23-1) is equivalent to

$$
\mathcal{O}_{\mathbb{P}(\mathcal{E})}(1)|_{X_{\Sigma}} = \mathcal{O}_{\mathbb{P}(\mathcal{E}|_{X_{\Sigma}})}(1).
$$

Lemma 6.7. For all $j = 1, ..., n$, the divisor $D_{\Phi_F} - D_{\tilde{e}_j}$ on $\mathbb{P}(\mathcal{E})$ is rationally equivalent to the divisor associated with the rational polytope ∂_j Newt (Φ_F) .

Proof. We now prove that the divisor $D_{\Phi_F} - D_{\tilde{e}_j}$ is associated with the polytope $e_j + \partial_j \text{Newt}(\Phi_F)$. Note that $e_j + \partial_j \text{Newt}(\Phi_F)$ is the intersection of Newt (Φ_F) with the affine halfspace $\{x_i \geq 1\}$. We have to prove that the support function of Newt $(\Phi_F) \cap \{x_j \geq 1\}$ takes the same value on all rays of $\Sigma_{\mathcal{E}}$, except for \tilde{e}_j , where it differs by one I of *x* be any element of \mathbb{R}^{n+m} . The value differs by one. Let v be any element of \mathbb{R}^{n+m} . The value

$$
-\min\{\langle w,v\rangle\ :\ w\in\mathrm{Newt}(\Phi_F)\}\
$$

of the support function of $\text{Newt}(\Phi_F)$ on v can only differ if the face $\text{Newt}(\Phi_F)^v$ is contained in the facet

$$
Newt(\Phi_F)^{\widetilde{e}_j} = Newt(\Phi_F) \cap \{x_j = 0\}.
$$

Note that a face of the form $\text{Newt}(\Phi_F)^{\{0\}\times e_i}$ is equal to the Cayley polytope

 $Cay(\mathcal{A}_0, \ldots, \mathcal{A}_{i-1}, \mathcal{A}_{i+1}, \ldots, \mathcal{A}_m),$

where we omit one of the constraints. In particular, it is always a facet, so we may restrict to rays of the form $\tilde{\rho}$. Let $\tilde{\rho}$ be a ray such that Newt $(\Phi_F)^{\tilde{\rho}}$ is contained in
Newt $(\Phi_F)^{\tilde{e}_i}$. By Proposition 6.6 we have Newt $(\Phi_F)^{\widetilde{e}_j}$. By Proposition [6.6](#page-23-2) we have

Cay (Newt
$$
(f_0)^{\rho}, \ldots
$$
, Newt $(f_m)^{\rho}$)
= Newt $(\Phi_F)^{\widetilde{\rho}}$
 \subseteq Newt $(\Phi_F)^{\widetilde{e}_j}$
= Cay (Newt $(f_0)^{e_j}, \ldots$, Newt $(f_m)^{e_j})$,

implying $\text{Newt}(f_i)^{\rho} \subseteq \text{Newt}(f_i)^{e_j}$ for all $i = 0, \ldots, m$. We obtain

$$
\left(\sum_{i=0}^m \text{Newt}(f_i)\right)^{\rho} = \sum_{i=0}^m \text{Newt}(f_i)^{\rho} \subseteq \sum_{i=0}^m \text{Newt}(f_i)^{e_j} = \left(\sum_{i=0}^m \text{Newt}(f_i)\right)^{e_j}.
$$

This is an inclusion of facets of the Minkowski sum $\sum_{i=0}^{m} \text{Newt}(f_i)$, so $\rho = e_j$ \Box

Before proving Theorem [A](#page-9-2) we need to prove a statement about the intersection of Q-Cartier divisors.

Lemma 6.8 (Generic intersection of \mathbb{Q} -Cartier divisors). Let X be a normal, proper variety of dimension n with Weyl divisors D_1, \ldots, D_n , and let k be an integer such that $\mathcal{O}(kD_i)$ is a line-bundle for each $i = 1, \ldots, n$. Let f_i be a global section of $\mathcal{O}(D_i)$ for $i = 1, \ldots, n$ such that $\mathcal{V}(\widetilde{f}_1, \ldots, \widetilde{f}_n)$ is a zero-dimensional smooth scheme contained in the smooth locus of X. Then

$$
k^{n} \# \mathcal{V}(\widetilde{f}_1,\ldots,\widetilde{f}_n) = c_1(\mathcal{O}(kD_1))\cdots c_1(\mathcal{O}(kD_n)).
$$

Proof. The length of the zero-dimensional scheme $\mathcal{V}((f_1)^k, \ldots, (f_n)^k)$ is equal to the product $c_1(\mathcal{O}(kD_1))\cdots c_1(\mathcal{O}(kD_n))$ of Chern classes. On the other hand, since f_1, \ldots, f_n intersect transversely, each isolated point of $\mathcal{V}((f_1)^k, \ldots, (f_n)^k)$ is isomorphic to the scheme $Spec \mathbb{C}[X_1, \ldots, X_n]/\langle X_1^k, \ldots, X_n^k \rangle$. In particular, we have

$$
k^n \cdot \text{length}(\mathcal{V}(\widetilde{f}_1,\ldots,\widetilde{f}_m)) = \text{length}(\mathcal{V}((\widetilde{f}_1)^k,\ldots,(\widetilde{f}_n)^k)),
$$

finishing the proof. \Box

Proof of Theorem [A.](#page-9-2) The vanishing locus of the homogenized system of Lagrange equations $\widetilde{\mathbf{L}}_F = (\widetilde{f}_1, \ldots, \widetilde{f}_m, \widetilde{\ell}_1, \ldots, \widetilde{\ell}_n)$ is the intersection of Z with the vanishing locus of $\tilde{f}_1, \ldots, \tilde{f}_m$ and by Lemma [6.4](#page-22-0) this is a smooth zero-dimensional variety, contained in $(\mathbb{C}^*)^{n+m}$. By Lemma [5.8,](#page-19-0) the algebraic degree of sparse polynomial optimization is equal to its cardinality. According to Lemma [6.7,](#page-23-3) the system L_F

comprises global sections of Q-Cartier divisors, associated with the respective, rational, polytopes $\text{Newt}(f_1), \ldots, \text{Newt}(f_m), \partial_1 \text{Newt}(\Phi_F), \ldots, \partial_n \text{Newt}(\Phi_F)$. As a consequence of Lemma [6.8,](#page-24-0) and using multilinearity of the mixed volume, we can express the number of solutions to L_F as the mixed volume [\(3.1\)](#page-9-4) of these polytopes. \Box

6.3. The proof of Theorem C . In this section, we study the intersection of the determinantal variety W with the vanishing locus V of f_1, \ldots, f_m in X. The proof of Theorem [C](#page-10-0) rests on a proof of transversality, and a characterisation of the cohomology class $[W]$ as a Porteous class.

We start by recalling Porteous' formula, also called the Giambelli–Thom–Porteous formula. For more details refer to chapter 14 in [\[Ful98\]](#page-27-22) and chapter 12 in [\[EH16\]](#page-27-19). The following statement is a special case of Theorem 12.4 in [\[EH16\]](#page-27-19).

Theorem 6.9. (Porteous' formula) Let $\varphi : \mathcal{E} \longrightarrow \mathcal{F}$ be a morphism of vector bundles of ranks $m + 1 \leq n$ on a smooth proper variety X of dimension n. We denote by W the (possibly non reduced) degeneracy locus of φ , supported on the set

$$
|W| = \{x \in X : \varphi_x : \mathcal{E}(x) \longrightarrow \mathcal{F}(x) \text{ is not injective}\}.
$$

If W is pure of codimension $n - m$ then the cohomology class of W is the $n - m$ qraded part of the product of the total Seqre class $s(\mathcal{E})$ and the total Chern class $c(\mathcal{F})$:

$$
[W] = (\mathbf{s}(\mathcal{E}) \mathbf{c}(\mathcal{F}))_{n-m}.
$$

We need the following modified version of Porteous' formula which only requires W to be pure dimensional after restricting to a subvariety \hat{X} of X.

Corollary 6.10. Under the assumptions of Theorem [6.9,](#page-25-0) let \hat{X} be an irreducible closed subvariety of X of codimension k , which intersects W transversely and let the intersection $\hat{X} \cap W$ be pure of codimension $n - m + k$. Then the cohomology class $\left[\hat{X} \cap W\right]$ is given by

$$
\left[\hat{X} \cap W\right] = \left[\hat{X}\right] \cdot \left(\textbf{s}(\mathcal{E}) \, \textbf{c}(\mathcal{F})\right)_{n-m}.
$$

Proof. Note that $\hat{X} \cap W$ is the degeneracy locus of the restriction $\varphi |_{\hat{X}}$. By applying Porteous' formula to $\phi|_{\hat{X}}: \mathcal{E}|_{\hat{X}} \to \mathcal{F}|_{\hat{X}}$ we get

$$
[\hat{X} \cap W] = \left(\mathbf{s}(\mathcal{E}|_{\hat{X}}) \,\mathbf{c}(\mathcal{F}|_{\hat{X}})\right)_{n-m+k}.
$$

Lastly we notice that $(s(\mathcal{E}|_{\hat{X}}) c(\mathcal{F}|_{\hat{X}}))_{n-m+k} = \left[\hat{X}\right] \cdot (s(\mathcal{E}) c(\mathcal{F}))_{n-m}$ by naturality of characteristic classes. □

Lemma 6.11. Under the assumptions of Theorem [C](#page-10-0) the intersection $V \cap W$ is transversal and contained in $(\mathbb{C}^*)^{\hat{n}}$.

Proof. Under the assumptions of Theorem [C,](#page-10-0) the assumptions from Section [5.2](#page-17-0) are satisfied. The inclusion in $(\mathbb{C}^*)^n$ follows from Lemma [6.4.](#page-22-0) We now show that transversality of the intersection $V \cap W$ follows from transversality of the intersection of Z with $\mathcal{V}(\tilde{f}_1,\ldots,\tilde{f}_m)$. Let $\pi: \mathbb{P}(\mathcal{E}) \longrightarrow X$ denote the natural projection and let $z = (x, [\lambda])$ be any element of $\mathbb{P}(\mathcal{E})$. If Z intersects $\pi^{-1}(V)$ transversely at z, then for the tangent spaces $T_{Z,z}$ and $T_{\pi^{-1}(V),z}$ at z it holds

(6.3)
$$
T_{Z,z} + T_{\pi^{-1}(V),z} = T_{\mathbb{P}(\mathcal{E}),z}.
$$

To see that W and V intersect transversely at x we show $T_{W,x} + T_{V,x} = T_{X,x}$. We apply the differential $d\pi$ to both sides of [\(6.3\)](#page-25-1) and note that we have the inclusions

$$
d\pi(T_{Z,z}) \subseteq T_{W,x}, \ d\pi(T_{\pi^{-1}(V),z}) \subseteq T_{V,x}, \ d\pi(T_{\mathbb{P}(\mathcal{E}),z}) = T_{X,x}.
$$

Proof of Theorem [C.](#page-10-0) By Lemma [5.8](#page-19-0) the algebraic degree of sparse polynomial optimization is the cardinality of $V \cap W \cap \mathbb{C}^n$. By Lemma [6.11,](#page-25-2) the scheme theoretic intersection $V \cap W$ is a smooth variety of dimension zero, contained in $(\mathbb{C}^*)^n$. We now finish the proof by verifying the assumptions of Corollary [6.10.](#page-25-3)

Let D_{f_i} be the Weyl divisors introduced in equation [\(5.1\)](#page-17-1). By the assumptions of Theorem [C,](#page-10-0) X is smooth. In particular, all divisors considered in this proof are Cartier. The variety W is defined to be the degeneracy locus of the matrix M , whose entries are global sections of the bundle $\mathcal{O}_X(D_{f_i}-D_{e_j})$. The transpose of M defines a morphism $\varphi : \mathcal{E} \to \mathcal{F}$ of vector bundles, where

$$
\mathcal{E} = \mathcal{O}_X(-D_{f_0}) \oplus \cdots \oplus \mathcal{O}_X(-D_{f_m}), \text{ and } \mathcal{F} = \mathcal{O}_X(-D_{e_1}) \oplus \cdots \oplus \mathcal{O}_X(-D_{e_n}).
$$

Now W is the degeneracy locus of φ , further $V \cap W$ is pure of dimension zero. Finally, $\mathcal{L}_{A_i} = \mathcal{O}_X(D_{f_i})$ which finishes the proof. \Box

REFERENCES

- [ABB+19] Carlos Am´endola, Nathan Bliss, Isaac Burke, Courtney R. Gibbons, Martin Helmer, Serkan Hosten, Evan D. Nash, Jose Israel Rodriguez, and Daniel Smolkin. The maximum likelihood degree of toric varieties. J. Symbolic Comput., 92:222–242, 2019.
- [AH18] Paolo Aluffi and Corey Harris. The Euclidean distance degree of smooth complex projective varieties. Algebra Number Theory, 12(8):2005–2032, 2018.
- [BD15] Jasmijn A. Baaijens and Jan Draisma. Euclidean distance degrees of real algebraic groups. Linear Algebra Appl., 467:174–187, 2015.
- [Ber75] David N. Bernstein. The number of roots of a system of equations. Funkcional. Anal. $i \ Priložen., 9(3):1-4, 1975.$
- [BHSW13] Daniel J. Bates, Jonathan D. Hauenstein, Andrew J. Sommese, and Charles W. Wampler. Numerically solving polynomial systems with Bertini, volume 25 of Software, Environments, and Tools. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2013.
- [BRT23] Paul Breiding, Kemal Rose, and Sascha Timme. Certifying zeros of polynomial systems using interval arithmetic. ACM Trans. Math. Software, 49(1):Art. 11, 14, 2023.
- [BSW22] Paul Breiding, Frank Sottile, and James Woodcock. Euclidean distance degree and mixed volume. Found. Comput. Math., 22(6):1743–1765, 2022.
- [CCG23] Jane Ivy Coons, Mark Curiel, and Elizabeth Gross. Mixed volumes of networks with binomial steady-states, 2023.
- [CHKS06] Fabrizio Catanese, Serkan Hosten, Amit Khetan, and Bernd Sturmfels. The maximum likelihood degree. Amer. J. Math., 128(3):671–697, 2006.
- [CKL22] Tianran Chen, Evgeniia Korchevskaia, and Julia Lindberg. On the typical and atypical solutions to the kuramoto equations, 2022.
- [CLS11] David A. Cox, John B. Little, and Henry K. Schenck. Toric varieties, volume 124 of Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, 2011.
- [Cox95] David A. Cox. The homogeneous coordinate ring of a toric variety. J. Algebraic Geom., 4(1):17–50, 1995.
- [DHO⁺16] Jan Draisma, Emil Horobet, Giorgio Ottaviani, Bernd Sturmfels, and Rekha R. Thomas. The Euclidean distance degree of an algebraic variety. Found. Comput. *Math.*, $16(1):99-149$, 2016 .
- [DLOT17] Dmitriy Drusvyatskiy, Hon-Leung Lee, Giorgio Ottaviani, and Rekha R. Thomas. The Euclidean distance degree of orthogonally invariant matrix varieties. Israel J. Math., 221(1):291–316, 2017.

- [NRS10] Jiawang Nie, Kristian Ranestad, and Bernd Sturmfels. The algebraic degree of semidefinite programming. Math. Program., 122(2, Ser. A):379–405, 2010.
- [PRW95] S. Poljak, F. Rendl, and H. Wolkowicz. A recipe for semidefinite relaxation for (0, 1) quadratic programming. J. Global Optim., 7(1):51–73, 1995.
- [Rum99] Siegfried M. Rump. INTLAB — INTerval LABoratory, pages 77–104. Springer Netherlands, Dordrecht, 1999.
- [TR01] Peng Hui Tan and L.K. Rasmussen. The application of semidefinite programming for detection in CDMA. IEEE Journal on Selected Areas in Communications, 19(8):1442– 1449, 2001.
- [Vav90] Stephen A. Vavasis. Quadratic programming is in NP. Inform. Process. Lett., 36(2):73–77, 1990.

Julia Lindberg, University of Texas-Austin Email address: julia.lindberg@math.utexas.edu URL: https://sites.google.com/view/julialindberg/home

LEONID MONIN, ÉCOLE POLYTECHNIQUE FÉDÉRALE DE LAUSANNE Email address: leonid.monin@epfl.ch URL: http://www.math.toronto.edu/lmonin/

Kemal Rose, KTH Royal Institute of Technology Email address: kemalr@kth.se URL: https://kemalrose.github.io/