Interpretable Distribution-Invariant Fairness Measures for Continuous Scores

Ann-Kristin Becker SCHUFA Holding AG ann-kristin.becker@schufa.de Oana Dumitrasc SCHUFA Holding AG oana.dumitrasc@schufa.de

Klaus Broelemann SCHUFA Holding AG klaus.broelemann@schufa.de

Abstract

Measures of algorithmic fairness are usually discussed in the context of binary decisions. We extend the approach to continuous scores. So far, ROC-based measures have mainly been suggested for this purpose. Other existing methods depend heavily on the distribution of scores, are unsuitable for ranking tasks, or their effect sizes are not interpretable. Here, we propose a distributionally invariant version of fairness measures for continuous scores with a reasonable interpretation based on the Wasserstein distance. Our measures are easily computable and well suited for quantifying and interpreting the strength of group disparities as well as for comparing biases across different models, datasets, or time points. We derive a link between the different families of existing fairness measures for scores and show that the proposed distributionally invariant fairness measures outperform ROC-based fairness measures because they are more explicit and can quantify significant biases that ROC-based fairness measures miss. Finally, we demonstrate their effectiveness through experiments on the most commonly used fairness benchmark datasets.

1 Introduction

In recent years, many decision-making processes in areas such as finance, education, social media or medicine have been automated, often at least in part with the goal of making those decisions more comparable, objective, and non-discriminatory [12, 16, 1, 5, 33]. For high-risk business transactions between individuals and companies (e.g. in the lending industry), often predictions of machine learning algorithms are incorporated into those decisions. Such algorithms aim to differentiate individuals as optimally as possible based on historical data and in terms of future behavior. They assign risk scores or risk categories to individuals. Even with good intentions, the approach runs the risk of directly or indirectly discriminating against individuals on the basis of protected characteristics, such as gender, ethnicity, political background or sexual orientation [24, 9, 20]. That may be the case, for example, if the data reflect biased social circumstances or include prejudicial historical decisions. Such discriminatory predictions manifest as disparities among protected groups and may occur in different forms and for different reasons. For example, individuals belonging to different protected groups may be assigned different scores even if they have the same outcome, or predictions may turn out to have different levels of consistency with the ground-truth risk. Unfortunately, in most cases different notions of algorithmic fairness are incompatible [2, 19, 34, 6, 32]. Various measures for algorithmic fairness have been developed that aim to quantify different kinds of group disparities [38, 18, 26]. So far, most of the available literature discusses the problem in the context of binary decision tasks [30, 2, 21]. However, in many applications, neither a final decision is known, nor is the explicit cost of false predictions. It may also be that a score is never applied as a pure decision

but only as a quantitative prediction that affects, e.g. the cost of a product (risk-based pricing). In these cases, fairness can only be fully assessed if the disparities between groups are summarized across the entire score model. Since regularly measuring group disparities of score models can help increase the visibility of group effects that might otherwise go unnoticed - especially in the case of underrepresented or historically disadvantaged groups - we need flexible and transparent measures that correctly quantify group disparities for continuous scores and allow us to interpret their implications. This paper presents a new approach to quantifying group disparities for continuous score models in a well interpretable and mathematically sound manner. We present a distribution-invariant framework, that allows for monitoring bias over time or between models and populations, even if there is a shift in the score distribution. Moreover, the setting allows to bridge the gap between common fairness-metrics stemming directly from three parity concepts [19, 15, 2, 26] and ROC-based approaches [36, 17, 37, 3]. As not all group disparities arise from discriminatory circumstances even large disparities between groups may be explainable or justifiable otherwise - assessing whether disparities are unfair should entail a more detailed analysis of their underlying causes and drivers. Thus, to be explicit, we use the term *disparity measure* instead of *fairness measure* throughout the rest of the paper to underline that all discussed measures are purely observational.

The paper is structured as follows: Most of the available quantitative disparity metrics for classifiers reduce down to three main parity concepts that are based on conditional independence: Independence, separation and sufficiency [2, 26, 21]. In Section 2, we discuss these concepts and existing related measures in terms of binary classifiers first, and generalize them to continuous scores in Section 3. We show that our proposed measures are more flexible than many existing metrics and we discuss their interpretability. In Section 4, we compare the presented measures to ROC-based disparity measures, and we prove that our proposed measures impose a stronger objective and are better suited to detect bias. We outline published related work throughout each section. Section 5 contains results of experiments using benchmark data and Section 6 includes final discussion and outlook. All proofs of technical results are deferred to the appendix.

2 Parity concepts and fairness measures for classifiers

Let Y denote a binary target variable with favorable outcome class Y = 0 and unfavorable class Y = 1, and X a set of predictors. Let $S \in S \subset \mathbb{R}$ denote an estimate of the posterior probability of the favorable outcome of Y, $\mathbb{P}(Y = 0 | X)$ or some increasing function of this quantity, in the following called *score*, with cumulative distribution function F_S and density function f_S . We assume S to be bounded with $|S| = \sup S - \inf S$ denoting the length of the score range. Let A be a (protected) attribute of interest defining two (protected) groups $(A \in \{a, b\}$ binary w.l.o.g.). We choose A = b as the group of interest, e.g. the expected discriminated group. All discussed measures are purely observational and based on the joint distribution of (S, A, Y). They can be easily calculated if a random sample of the joint distribution is available.

Note that each continuous score S induces an infinite set of binary classifiers by choosing a threshold $s \in S$ and accepting every sample with S > s. We define disparity measures for binary classifiers in dependence of such a threshold value s. For a group A, the *positive rate* at a threshold s is given by $\operatorname{PR}_A(s) = \mathbb{P}(S > s|A) = 1 - F_{S|A}(s)$, the *true positive and false positive rates* by $\operatorname{TPR}_A(s) = 1 - F_{S|A,Y=0}(s)$ and $\operatorname{FPR}_A(s) = 1 - F_{S|A,Y=1}(s)$, respectively. We will write in short $F := F_S$ and $f := f_S$, as well as $S_{ay} := S|A = a, Y = y$, $F_{ay} := F_{S|A=a,Y=y}$ and $S_{by}, F_{by}, f_{ay}, f_{by}$ for the conditional random variables, distribution functions and density functions. For a cumulative distribution function G, we denote by G^{-1} the related quantile function (generalized inverse) with $G^{-1}(p) = \inf\{x \in \mathbb{R} : p \leq G(x)\}$ which fulfills $G^{-1}(G(X)) = X$ almost surely. If G is continuous and strictly monotonically increasing, then the quantile function is the inverse.

Independence (selection rate parity) The random variables S and A satisfy independence if $S \perp A$, which implies $F_{S|A=a} = F_{S|A=b} = F_S$. Group disparity of classifiers can be quantified by the difference between the positive rates [26, 38, 11]

$$c-\text{bias}_{\text{IND}}(S_a, S_b; s) = \text{PR}_b(s) - \text{PR}_a(s) = F_a(s) - F_b(s).$$
(1)

The concept of independence contradicts optimality S = Y, if $Y \not\perp A$ and is, thus, not an intuitive fairness measure in most cases. On the other hand, the following two measures, separation and sufficiency, are both compatible with optimality and allow $A \not\perp Y$, as they include the target variable

Y in the independence statements and allow for disparities that can be explained by group differences in the ground-truth.

Separation (error rate parity) The random variables S, A and Y satisfy separation if $S \perp A \mid Y$. For a binary outcome Y, the separation condition splits into *true positive rate parity* $F_{S|A=a,Y=0} = F_{S|A=b,Y=0} = F_{S|Y=0}$ (equal opportunity, EO) [40, 15] and *false positive rate parity* $F_{S|A=a,Y=1} = F_{S|A=b,Y=1} = F_{S|Y=1}$ (predictive equality, PE) [8, 26]. If both hold, the condition is also known as equalized odds [26, 15]. Group disparity of classifiers can be quantified by the difference between the true and false positive rates

$$c-bias_{EO}(S_a, S_b; s) = TPR_b(s) - TPR_a(s) = F_{a0}(s) - F_{b0}(s),$$
(2)

$$c-\text{bias}_{\text{PE}}(S_a, S_b; s) = \text{FPR}_b(s) - \text{FPR}_a(s) = F_{a1}(s) - F_{b1}(s).$$
(3)

If separation is violated, the model output includes more information about the group A as is justified by the ground truth Y alone. So, different groups carry different costs of misclassification. It is therefore a reasonable concept for surfacing potential inequities. Conversely, a violation of sufficiency results in a different calibration and a different meaning of identical score values per group. That is the case, if the relation of A and Y is not properly modeled by the score.

Sufficiency (predictive value parity) The random variables S, A and Y satisfy sufficiency if $Y \perp A \mid S$ (in words, S is sufficient to optimally predict Y). Sufficiency implies group parity of positive and negative predictive values. However, especially in case of continuous scores, usually, *calibration* within each group [19] (resp. *test fairness* [6]), as an equivalent concept, is used instead [2]. The calibration bias examines if the model's predicted probability deviates similarly strongly from the true outcome rates within each group:

c-bias_{CALI}
$$(S_a, S_b; s) = \mathbb{P}(Y = 0 | A = b, S = s) - \mathbb{P}(Y = 0 | A = a, S = s).$$
 (4)

Well-calibration [19, 32] additionally requires the prediction of both groups to accurately reflect the ground truth $\mathbb{P}(Y = 0|A, S = s) = s$. For determining the calibration difference, the score range is usually binned into a fixed number of intervals. A high calibration bias reflects the fact that (for a given score s) the lower-risk group carries the costs of the higher-risk group. The concept of sufficiency is especially important if the model is applied in a context, where both, the score and the group membership are available to the decision maker. Then, a high calibration bias will evoke a group-specific interpretation and handling of identical score values. On the other hand, sufficiency does not prevent discrimination: high- and low-risk individuals of a group can be mixed and assigned an intermediate risk score without violating sufficiency. Moreover, sufficiency is often naturally fulfilled as a consequence of unconstrained supervised learning, especially if the group membership is (at least to some extent) encoded in the input data. Thus, it is usually not a constraint and not a trade-off with predictive performance [25].

In general, independence, separation and sufficiency are opposing concepts. It can be shown that for a given dataset, except for special cases (like perfect prediction or equal base rates), every pair of the three parity concepts is mathematically incompatible [2, 19, 34, 6, 32].

3 Generalization to continuous scores

We propose to use the expected absolute classifier bias as a disparity measure for scores. Note, that an expected value of zero implies that every classifier derived from the score by choosing a groupunspecific threshold will be bias-free. By evaluating and aggregating the bias across all possible decision thresholds, this generalization serves as a useful diagnostic tool in fairness analyses and follows a similar idea as used in ROC analyses. The two proposed versions can be seen as generalized rate differences. They differ only in the way, in which possible thresholds are weighted. We show, that for the concepts independence and separation, the proposed disparity measures are identical to Wasserstein distances between the groupwise score-distributions.

3.1 Expected classifier bias with uniformly weighted thresholds

Definition 3.1. By assuming each threshold $s \in S$ is equally important, we define

$$\operatorname{bias}_{x}^{\mathcal{U}}(S_{a}, S_{b}) := \mathbb{E}_{S \sim \mathcal{U}}[|\operatorname{c-bias}_{x}(S_{a}, S_{b}; S)|] = \frac{1}{|\mathcal{S}|} \int_{\mathcal{S}} |\operatorname{c-bias}_{x}(S_{a}, S_{b}; s)| \, ds.$$
(5)

Theorem 3.2. For the concepts independence and separation, i.e for $x \in \{IND, PE, EO\}$, it holds:

(i) $\operatorname{bias}_{x}^{\mathcal{U}}(S|A=a, S|A=b)$ is equal to the normalized Wasserstein-1-distance between the conditional score distributions in the groups over the (finite) score region S i.e.

$$\operatorname{bias}_{x}^{\mathcal{U}}(S_{a}, S_{b}) = \frac{1}{|\mathcal{S}|} \cdot W_{1}(S_{ay}, S_{by}), \tag{6}$$

where y = 0 for x = EO, y = 1 for x = PE, and $y = \cdot$ for x = IND.

(ii) As a consequence, we can derive the disparity between average scores per group (known as *balance for the positive / negative class* [19]) as a lower bound, i.e.

$$\operatorname{bias}_{x}^{\mathcal{U}}(S_{a}, S_{b}) \geq \frac{1}{|\mathcal{S}|} \left| \mathbb{E}[S_{by}] - \mathbb{E}[S_{ay}] \right|.$$
(7)

The Wasserstein distance was proposed as a fairness measure recently by Miroshnikov et al. [29] or Kwegyir-Aggrey et al. [23], and it was especially used for debiasing purposes earlier [28, 14, 7]. A formal definition and properties of W_1 can be found in the appendix. For calibration, the bias $\text{bias}_{CALI}^{\mathcal{U}}$ is equal to the two-sample version of the the l_1 -calibration error [22].

3.2 Distribution-invariant setting

It can be difficult to compare the expected classifier bias of datasets with distinct score distributions. Especially for imbalanced datasets score distributions are often highly skewed. In this case, disparities in dense score areas may be more critical as they effect more samples. That is why we propose an alternative generalization that weights the thresholds by their frequency observed in the population. By this, the resulting fairness measures become distribution-free and evaluate the fairness of a bipartite ranking task, similar to ROC measures. Each sample is equally important in this scenario.

Definition 3.3.

$$\operatorname{bias}_{x}^{S}(S_{a}, S_{b}) := \mathbb{E}_{S \sim F}[|\operatorname{c-bias}_{x}(S_{a}, S_{b}; S)|]$$
(8)

$$= \int_{0}^{1} |\operatorname{c-bias}_{x}(S_{a}, S_{b}; F^{-1}(r))| \, dr = \int_{\mathcal{S}} |\operatorname{c-bias}_{x}(S_{a}, S_{b}; s)| \cdot f(s) \, ds \quad (9)$$

Note that $bias_x^S(S|A = a, S|A = b)$ is invariant under monotonic score transformations as it is a purely ranking-based metric, $bias_x^{\mathcal{U}}(S|A = a, S|A = b)$ is not. If $S \sim \mathcal{U}$ it holds $bias^S = bias^{\mathcal{U}}$. We show, that the distribution-free bias is equal to the Wasserstein-1-distance between quantile-transformed distributions. To our knowledge, this is the first introduction of a distribution-free Wasserstein-distance as a fairness measure.

Theorem 3.4. For the concepts independence and separation, i.e. for $x \in \{IND, PE, EO\}$, it holds:

(i) bias_x^S is equal to the Wasserstein-1-distance using the push-forward by the quantile function $F^{-1} \# \mathcal{L}_1$ as ground metric (with y = 0 for $x = \operatorname{EO}$, y = 1 for $x = \operatorname{PE}$, and $y = \cdot$ for $x = \operatorname{IND}$)

$$\operatorname{bias}_{x}^{S}(S_{a}, S_{b}) = W_{1}(F(S_{ay}), F(S_{by})) = \int_{0}^{1} |F_{ay} \circ F^{-1}(t) - F_{by} \circ F^{-1}(t)| \, dt.$$
(10)

(ii) We can derive the disparity between the average relative rank per group as a lower bound.

For reasons of simplicity, we will use the notation $W_Z(X, Y) := W_1(F_Z(X), F_Z(Y))$.

3.3 Interpretation of the score bias

In general, $bias^{\mathcal{U}}$ and $bias^{S}$ take values in the interval [0, 1] as they are expected values over rate differences. The optimal value, a bias of zero, indicates group parity for all decision thresholds with respect to the analyzed type of classifier error. When comparing multiple score models or one model over multiple populations, a smaller bias is preferable. The distribution-free method allows

direct comparison of models with different score distributions with respect to group parity in bipartite ranking tasks. $bias^{\mathcal{U}}$ and $bias^{S}$ can be interpreted as the classifier bias to be expected at a randomly chosen threshold - either randomly selected from all available score values ($bias^{\mathcal{U}}$) or by randomly selecting one sample and assigning the favorable label to all samples that are ranked higher ($bias^{S}$).

In addition, the separation and independence biases can be interpreted in terms of the Wasserstein distance (or *Earth Mover distance*): The bias is measured as the minimum cost of aligning the two groups with respect to the analyzed type of classifier error. Here, the baseline distance is measured in normalized scores for bias^U or in ranks for bias^S. It indicates what proportion of a group must be scored (how) differently in order to equalize the groups.

3.4 Positive and negative components of the score bias

Unlike a classifier bias, a score bias does not have to be overall positive or negative for a particular group. Instead, there may be thresholds at which one group is disadvantaged and others at which the opposing group is disadvantaged. To further analyze the bias, we can decompose the total bias into a positive and a negative component (positive and negative from the point of view of the chosen disadvantaged group, here b). For this purpose, the classifier bias is divided into a positive and a negative part for each threshold

$$c-bias^+(s) = max(c-bias(s), 0)$$
 and $c-bias^-(s) = -min(c-bias(s), 0)$.

This allows to derive a decomposition of both score bias types into two components:

$$\text{pos-bias}_x(S_a, S_b) = \mathbb{E}[\text{c-bias}_x^+(S_a, S_b; S)], \tag{11}$$

neg-bias_x(S_a, S_b) =
$$\mathbb{E}[\text{c-bias}_x^-(S_a, S_b; S)],$$
 (12)

where $bias_x(S_a, S_b) = pos bias_x(S_a, S_b) + neg bias_x(S_a, S_b)$. By dividing each component by the total bias, a percentage can be calculated. The decomposition helps to interpret, which of the two compared groups is affected predominantly negatively by the observed bias. A similar decomposition of a Wasserstein bias was proposed by Miroshnikov et al. [29].

4 ROC-based fairness measures and relations

Furthermore, there exists a wide variety of (separation) fairness metrics which are calculated based on ROC curves or the area under the curves. We show, that the proposed distribution-free bias measures outperform these ROC-based measures as they are more explicit, easier to interpret, and can measure biases, that ROC-based fairness measures cannot catch. We define the ROC curve between two arbitrary random variables G, H, similar to Vogel et al. [36]. In a bipartite ranking or scoring task, the ROC curve is usually used to evaluate the separability between positive and negative outcome class. In this case, $G = S_0, H = S_1$.

Definition 4.1 (ROC). Let G and H be two random variables with cumulative distribution functions F_G, F_H on \mathbb{R} with quantile functions F_G^{-1}, F_H^{-1} . Then the ROC curve of G and H is the mapping

$$\operatorname{ROC}_{G,H} : p \in [0,1] \mapsto 1 - F_G(F_H^{-1}(1-p))$$
 (13)

with the area under the curve (AUROC) and the Gini coefficient defined as

$$\operatorname{AUROC}(G,H) = \int_0^1 \operatorname{ROC}_{G,H}(p) \, dp \quad \text{and} \quad \operatorname{Gini}(G,H) = 2 \cdot \operatorname{AUROC}(G,H) - 1.$$
(14)

Definition 4.2. Similar to the above introduced biases, a ROC-based disparity-measure for score models can be defined as the expected absolute difference between two ROC curves

$$\operatorname{bias}_{\operatorname{ROC}}(S_a, S_b) = \mathbb{E}[|\operatorname{ROC}_{S_{b0}, S_{b1}} - \operatorname{ROC}_{S_{a0}, S_{a1}}|] = \int_0^1 |\operatorname{ROC}_{S_{b0}, S_{b1}}(s) - \operatorname{ROC}_{S_{a0}, S_{a1}}(s)| \, ds$$

 $bias_{ROC}(S|A = a, S|A = b)$ is equal to the absolute between ROC area (*ABROCA*) [13]. In general, $bias_{ROC}(S|A = a, S|A = b) \ge |AUROC(S_{b0}, S_{b1}) - AUROC(S_{a0}, S_{a1})|$, which is known as *intra-group fairness* and often used as a fairness measure for scores [36, 3, 4, 37]. If the ROC curves

of two groups do not cross (i.e. one group gets uniformly better scores than the other), equality holds. As the thresholds that lead to certain ROC values (pair of FPR and TPR at a certain score threshold) are group-specific, it is not sufficient to compare intra-group ROC curves [36]. Thus, we define a second ROC-based measure that compares the discriminatory power across groups and is based on the cross-ROC curve [17].

Definition 4.3. We define the cross-ROC bias as the expected difference of the ROC curves across groups

$$\operatorname{bias}_{\operatorname{xROC}}(S_a, S_b) = \mathbb{E}[|\operatorname{ROC}_{S_{b0}, S_{a1}} - \operatorname{ROC}_{S_{a0}, S_{b1}}|] = \int_0^1 |\operatorname{ROC}_{S_{b0}, S_{a1}}(s) - \operatorname{ROC}_{S_{a0}, S_{b1}}(s)| \, ds$$

The cross-ROC bias evaluates the difference in separability of negatives samples in one group versus positive samples of the other group. $bias_{xROC}$ is always greater or equal to the related AUROC-based fairness-measure $|AUROC(S_{a0}, S_{b1}) - AUROC(S_{b0}, S_{a1})|$, that is known as subgroup positive background negative (BPSN) or inter-group fairness [4, 36, 3, 37].

4.1 Relating Wasserstein and ROC biases

We now reveal some connections of the distribution-free Wasserstein disparity measures with the ROC-based disparity measures. We first consider the general case of the Wasserstein distance between two random variables X, Y quantile-transformed by Z.

For the following section, we require $ROC_{X,X}(r) = r$. This is fulfilled, whenever F_X is continuous and strictly monontonic increasing, so it permits a well-defined inverse, or if the ROC-curve is interpolated linearly from finite data.

Theorem 4.4. The quantile-transformed Wasserstein distance can be rewritten in terms of ROC

$$W_Z(X,Y) = \int_0^1 |F_X(F_Z^{-1}(t)) - F_Y(F_Z^{-1}(t))| dt = \int_0^1 |\operatorname{ROC}_{X,Z}(t) - \operatorname{ROC}_{Y,Z}(t)| dt.$$
(15)

Moreover, we easily get the following result.

Proposition 4.5. Let Z_i , i = 1, ..., n be random variables with values in S and with densities f_i . Let Z_K be their mixture, where K is a random variable with values in $\{1, ..., n\}$. Then their joint density is given by $f_{Z_K}(x) = \sum_{i=1}^n \mathbb{P}(K=i)f_i(x)$ and it holds

$$W_{Z_K}(X,Y) = \sum_{i=1}^{n} \mathbb{P}(K=i) W_{Z_i}(X,Y).$$
(16)

Formulating S as a mixture of the two groups and two outcome classes $S_{a0}, S_{a1}, S_{b0}, S_{b1}$, we get

$$W_{S}(S_{ay}, S_{by}) = w_{a0} \cdot W_{S_{a0}}(S_{ay}, S_{by}) + w_{b0} \cdot W_{S_{b0}}(S_{ay}, S_{by})$$

$$+ w_{a1} \cdot W_{S_{a1}}(S_{ay}, S_{by}) + w_{b1} \cdot W_{S_{b1}}(S_{ay}, S_{by}).$$
(17)

By looking at the different mixture components, we can reveal a connection to the ROC-based disparity measures.

Lemma 4.6. $W_{S_{ay}}(S_{ay}, S_{by})$ and $W_{S_{a\tilde{y}}}(S_{ay}, S_{by})$ for $\tilde{y} \neq y$ can be rewritten in terms of ROC

$$W_{S_{ay}}(S_{ay}, S_{by}) = \int_{0}^{1} \left| \text{ROC}_{S_{by}, S_{ay}}(r) - r \right| dr,$$
(18)

$$W_{S_{a\bar{y}}}(S_{ay}, S_{by}) = \int_0^1 |\operatorname{ROC}_{S_{by}, S_{a\bar{y}}}(r) - \operatorname{ROC}_{S_{ay}, S_{a\bar{y}}}(r)| dr.$$
(19)

Lemma 4.7. From Jensen inequality, it follows

$$W_{S_{ay}}(S_{ay}, S_{by}) \ge |\operatorname{AUROC}(S_{by}, S_{ay}) - \frac{1}{2}| = \frac{1}{2} \cdot |\operatorname{Gini}(S_{by}, S_{ay})|.$$
 (20)

If the ROC curve does not cross the diagonal, then equality holds.

Theorem 4.8. We can now decompose each separation bias into a sum of four ROC statements. Let $w_{ay} = \mathbb{P}(Y = y, A = a)$ and $w_{by} = \mathbb{P}(Y = y, A = b)$, as well as $w_y = \mathbb{P}(Y = y)$, then it holds:

$$bias_{EO}^{S}(S_{a}, S_{b}) = w_{a0} \int_{0}^{1} |ROC_{S_{b0}, S_{a0}}(r) - r| dr + w_{b0} \int_{0}^{1} |ROC_{S_{a0}, S_{b0}}(r) - r| dr + w_{a1} \int_{0}^{1} |ROC_{S_{a0}, S_{a1}}(r) - ROC_{S_{b0}, S_{a1}}(r)| dr + w_{b1} \int_{0}^{1} |ROC_{S_{a0}, S_{b1}}(r) - ROC_{S_{b0}, S_{b1}}(r)| dr,$$
(21)

and analogously for $\text{bias}_{\text{PE}}^{S}(S_a, S_b)$ by exchanging w_{a0} with w_{a1} , w_{b0} with w_{b1} , S_{a0} with S_{a1} and S_{b0} with S_{b1} .

Corollary 4.9. From Theorem 4.8 we can infer upper bounds of the separation biases and their sum

$$bias_{EO}^{S}(S_a, S_b) \le 1 - \frac{w_0}{2} \text{ and } bias_{PE}^{S}(S_a, S_b) \le 1 - \frac{w_1}{2}$$
 (22)

$$\Rightarrow \operatorname{bias}_{\mathrm{EO}}^{S}(S_a, S_b) + \operatorname{bias}_{\mathrm{PE}}^{S}(S_a, S_b) \le \frac{3}{2}.$$
(23)

Moreover, we show that the sum of the separation biases is an upper bound (up to population-specific constants) to both ROC biases and the separability of the groups within each outcome class.

Theorem 4.10. The following inequality holds ¹

$$\operatorname{bias}_{\mathrm{EO}}^{S}(S_{a}, S_{b}) + \operatorname{bias}_{\mathrm{PE}}^{S}(S_{a}, S_{b}) = W_{S}(S_{a0}, S_{b0}) + W_{S}(S_{a1}, S_{b1})$$

$$\geq \frac{\min(w_{a0}, w_{a1}, w_{b0}, w_{b1})}{2} \cdot (\operatorname{bias}_{\mathrm{ROC}} + \operatorname{bias}_{\mathrm{xROC}} + \operatorname{Gini}(S_{a0}, S_{b0}) + \operatorname{Gini}(S_{a1}, S_{b1})). \quad (24)$$

Note, that the constant $\min(w_{a0}, w_{a1}, w_{b0}, w_{b1})/2$ is fixed for each dataset. Thus, decreasing both separation biases leads to a decrease of the sum of both ROC biases as well as the separability of the groups within each outcome class. Especially, separation biases of zero also diminish both ROC biases.

Corollary 4.11. Zero separation biases imply zero ROC biases

$$\operatorname{bias}_{\mathrm{EO}}^{S}(S_a, S_b) = \operatorname{bias}_{\mathrm{PE}}^{S}(S_a, S_b) = 0 \Rightarrow \operatorname{bias}_{\mathrm{ROC}}(S_a, S_b) = \operatorname{bias}_{\mathrm{xROC}}(S_a, S_b) = 0.$$
(25)

The inverse does not hold.

Theorem 4.12. Moreover, if only one separation bias is zero, ROC and cross-ROC bias become equal

$$\operatorname{bias}_{\mathrm{EO}}^{S}(S_{a}, S_{b}) = 0 \text{ or } \operatorname{bias}_{\mathrm{PE}}^{S}(S_{a}, S_{b}) = 0 \Rightarrow \operatorname{bias}_{\mathrm{ROC}}(S_{a}, S_{b}) = \operatorname{bias}_{\mathrm{xROC}}(S_{a}, S_{b}).$$
(26)

5 Experiments

We use the COMPAS dataset, the Adult dataset and the German Credit dataset to demonstrate the application of the fairness measures for continuous scores. For each bias, we perform permutation tests to determine statistical significance under the null hypothesis of group parity [10, 35]. Experimental details and complete results including all presented bias types can be found in appendix.

5.1 COMPAS

We calculate the different types of biases for the famous COMPAS decile score (n = 7214), which predicts the risk of violent recidisvism within two years following release. We choose race as protected attribute and set African-America as the expected discriminated group versus Caucasian race. To be consistent with the notation in this paper, we calculate the counter-score, so that a high score stands for the favorable outcome. In constrast to the original analysis [24] we calculate

¹Note, that if F_{ay} and F_{by} have identical supports and permit an inverse, then $Gini(S_{ay}, S_{by}) = Gini(S_{by}, S_{ay})$. If this symmetry is not fulfilled, the minimum of both must be used on the right side.

type of bias	total	pos.	neg.	p-value
$bias^{S}_{EO}$	0.1613	0%	100%	<0.01
$bias_{PE}^{S}$	0.1543	0%	100%	< 0.01
$bias_{CALI}^S$	0.0340	79%	21%	0.19
bias _{ROC}	0.0160	46%	54%	0.38
$bias_{xROC}$	0.3723	0%	100%	< 0.01

Table 1: Bias of COMPAS score of African-American vs. Caucasian.

Table 2: Gender bias of logistic regression (trained with and without sex) scores on German Credit Risk dataset; positive and negative component from the point of view of female persons.

	LogR			LogR (debiased)				
type of bias	total bias	pos.	neg.	p-value	total bias	pos.	neg.	p-value
$bias_{EO}^S$	0.083	0%	100%	0.05	0.048	93%	7%	0.29
$\mathrm{bias}_{\mathrm{PE}}^{S}$	0.092	0%	100%	0.12	0.010	62%	38%	1
$bias_{CALI}^S$	0.291	46%	54%	0.28	0.299	58%	42%	0.29

the bias over the entire score area. Results (Table 1) show a significant separation bias against the African-American and in favor of the Caucasian race. The disadvantaged group experiences a much lower true-positive rate (rate difference in average $bias_{EO}^S = 0.16$) as well as false positive rate (rate difference in average $bias_{PE}^S = 0.15$). The calibration bias is lower and not statistically significant but predominantly in favor of the African-American race. While the ROC bias is also low (implying that the separability is equally good in both groups considered independently), the cross-ROC bias is again high. In this case, there is not much difference between $bias^{\mathcal{U}}$ and $bias^{\mathcal{S}}$ (complete results can be found in appendix).

5.2 German Credit Data

Moreover, we trained two logistic regression scores on the German Credit Risk dataset (n = 1000) to predict if a borrower belongs to the good risk class. The first model *LogR* uses all available nine predictors including the feature *sex*, which we choose as protected attribute. For the second score *LogR* (*debiased*), the protected attribute was removed from the model input. We set *female* as the expected discriminated group. The scores achieve an AUROC of 0.772 and 0.771.

Compared to COMPAS, the separation biases of both models are lower (all below 0.1) whereas the calibration biases are higher (close to 0.3). Removing the attribute decreases the separation bias (Table 2), while it slightly increases the calibration bias. Note that while *LogR* contains bias to the detriment of female, the debiased model predominantly favors female over male.

5.3 UCI Adult

Moreover, we used the UCI Adult dataset (n = 32561) to train three different scores that predict the probability of the income being above 50k\$. Again, we choose *sex* as the protected attribute and *female* as the expected discriminated group. As before, a logistic regression was trained including (*logR*) and excluding (*logR* (*debiased*)) the protected attribute *sex*. Moreover, an XGBoost model (*XGB*) was trained with the complete feature set. The logistic regression achieved an AUROC of 0.895 with and of 0.892 without the protected attribute, the XGBoost model achieved an AUROC of 0.923 on the testset. Resulting biases are shown in Table 3, with the lowest bias in bold.

Removing the protected attribute from the model input improves all biases of LogR except $bias_{ROC}$ but separation biases are still against female while the calibration bias of the debiased model is predominantly in favor of female. XGB outperforms the logistic regression model that was trained on the same data in terms of fairness. In most cases, the bias of the XGB model is even smaller than the bias of logR (debiased). Here, due to the high sample size, all biases are statistically significant. We see a difference between $bias^{\mathcal{U}}$ and $bias^{S}$ that is due to the skewed score distributions on the

	LogR		LogR (debiased)			XGB			
type of bias	total bias	pos.	neg.	total bias	pos.	neg.	total bias	pos.	neg.
$bias_{EO}^S$	0.107	0%	100%	0.069	0%	100%	0.057	0%	100%
$bias_{PE}^{S}$	0.164	0%	100%	0.121	0%	100%	0.143	0%	100%
$bias_{CALI}^{\overline{S}}$	0.053	21%	79%	0.046	56%	44%	0.041	50%	50%
bias _{ROC}	0.049	98%	2%	0.051	98%	2%	0.032	99%	1%
bias _{xROC}	0.206	0%	100%	0.151	0%	100%	0.129	0%	100%
$\mathrm{bias}_{\mathrm{EO}}^{\mathcal{U}}$	0.162	0%	100%	0.104	0%	100%	0.089	0%	100%
$\mathrm{bias}_{\mathrm{PE}}^{\mathcal{U}}$	0.118	0%	100%	0.098	0%	100%	0.101	0%	100%
$\mathrm{bias}_{\mathrm{CALI}}^{\overline{\mathcal{U}}}$	0.057	25%	75%	0.054	58%	42%	0.053	51%	49%

Table 3: Gender bias of logistic regression (trained with and without sex) and XGBoost on Adult dataset; positive and negative component from the point of view of female persons. Each permutation tests gives p < 0.01.

imbalanced dataset (appendix Fig. C1-C3): in general rate differences in the range of low scores are weighted higher for bias^S as they effect more people (appendix Fig. C4-C5). Note that bias_{ROC} is in favor of female persons: Looking only at groupwise ROC curves (bias_{ROC}) suggests an advantage for females. However, female persons experience lower true- and false positive rates at every possible threshold that is chosen independently of the group, as bias^S_{EO} and bias^S_{PE} clearly show.

6 Discussion and Outlook

In this paper, we introduced a family of distribution-invariant group disparity measures for continuous scores that have an intuitive interpretation and theoretical grounding based on the Wasserstein distance. We derived their relation to well-established parity concepts and to ROC-based measures and we proved, that reducing the proposed separation biases is a stronger objective than reducing ROC-based measures and, hence, is better suited to cover different sorts of bias. Moreover, we demonstrated the practical application on fairness benchmark datasets. Our results show that removing information about the attribute influences the fairness of a model and also which group is affected by it. They also show that debiasing often leads to a shift between different bias types and should be monitored carefully. XGBoost results may indicate that flexible models can produce fairer results than simpler models. The proposed measures generalize rate differences from classification tasks to entire score models. As a future extension, a generalization of rate ratios is another option that is to be explored. Moreover, the discussed decision model errors (TPR/FPR/Calibration) could be summed or related to each other (i.e., TPR/FPR) to create further disparity measures. Note also, that the given definitions of the classifier biases are based on the l_1 -norm. Especially when used for bias mitigation, that we did not cover here, it may also be useful to replace the l_1 -norm by l_p with p > 1, especially l_2 or l_{∞} , to penalize large disparities more than small ones. However, the score bias is then no longer a Wasserstein-distance. Another option is to use the Wasserstein-p-distance with p > 1. Typically, the outcome of fairness analyses is to assess whether certain groups are discriminated against by a score model. All the proposed disparity measures can be used to assess the group disparity of the errors made by the model. While parity, i.e. a small bias, can be taken as a sign that there is no algorithmic unfairness in a sample with respect to a particular type of error, not all disparities are discriminatory. For practical applications we propose not to use hard thresholds to decide whether a model is fair or unfair. If needed, such thresholds can be chosen similarly to the thresholds for classification biases and should be task-specific. Once a high bias is detected, the causes of the disparities should be analyzed in detail to decide for follow-up actions. The relation to the field of causal fairness criteria (i.e. [31, 39, 27]) is out of scope of this manuscript. Further studies should investigate the relation and how they can be used to perform follow-up analyses in case of significant group disparities.

References

- Oscar Alvarado and Annika Waern. Towards Algorithmic Experience: Initial Efforts for Social Media Contexts. In Proceedings of the 2018 CHI Conference on Human Factors in Computing Systems, pages 1–12, Montreal QC Canada, April 2018. ACM. ISBN 978-1-4503-5620-6. doi: 10.1145/3173574.3173860. URL https://dl.acm.org/doi/10.1145/3173574.3173860.
- [2] Solon Barocas, Moritz Hardt, and Arvind Narayanan. Fairness and Machine Learning: Limitations and Opportunities. fairmlbook.org, 2019. URL http://www.fairmlbook.org.
- [3] Alex Beutel, Jilin Chen, Tulsee Doshi, Hai Qian, Li Wei, Yi Wu, Lukasz Heldt, Zhe Zhao, Lichan Hong, Ed H. Chi, and Cristos Goodrow. Fairness in Recommendation Ranking through Pairwise Comparisons. In Proceedings of the 25th ACM SIGKDD International Conference on Knowledge Discovery & Data Mining, pages 2212–2220, Anchorage AK USA, July 2019. ACM. ISBN 978-1-4503-6201-6. doi: 10.1145/3292500.3330745. URL https://dl.acm. org/doi/10.1145/3292500.3330745.
- [4] Daniel Borkan, Lucas Dixon, Jeffrey Sorensen, Nithum Thain, and Lucy Vasserman. Nuanced Metrics for Measuring Unintended Bias with Real Data for Text Classification. In *Companion Proceedings of The 2019 World Wide Web Conference*, pages 491–500, San Francisco USA, May 2019. ACM. ISBN 978-1-4503-6675-5. doi: 10.1145/3308560.3317593. URL https: //dl.acm.org/doi/10.1145/3308560.3317593.
- [5] Taina Bucher. The algorithmic imaginary: Exploring the ordinary affects of Facebook algorithms. Information, Communication & Society, 20(1):30–44, January 2017. ISSN 1369-118X, 1468-4462. doi: 10.1080/1369118X.2016.1154086. URL https://www.tandfonline.com/doi/ full/10.1080/1369118X.2016.1154086.
- [6] Alexandra Chouldechova. Fair Prediction with Disparate Impact: A Study of Bias in Recidivism Prediction Instruments. *Big Data*, 5(2):153-163, June 2017. ISSN 2167-6461, 2167-647X. doi: 10.1089/big.2016.0047. URL http://www.liebertpub.com/doi/10. 1089/big.2016.0047.
- [7] Evgenii Chzhen, Christophe Denis, Mohamed Hebiri, Luca Oneto, and Massimiliano Pontil. Fair Regression with Wasserstein Barycenters. In *Proceedings of the 34th International Conference* on Neural Information Processing Systems, {NIPS'20}. {Curran Associates Inc.}, June 2020. doi: 10.5555/3495724.3496338. URL http://arxiv.org/abs/2006.07286.
- [8] Sam Corbett-Davies, Emma Pierson, Avi Feller, Sharad Goel, and Aziz Huq. Algorithmic decision making and the cost of fairness. In *Proceedings of the 23rd Acm Sigkdd International Conference on Knowledge Discovery and Data Mining*, pages 797–806, June 2017. doi: 10.1145/3097983.309809. URL http://arxiv.org/abs/1701.08230.
- [9] Amit Datta, Michael Carl Tschantz, and Anupam Datta. Automated experiments on ad privacy settings: A tale of opacity, choice, and discrimination. *arXiv preprint arXiv:1408.6491*, 2014.
- [10] Cyrus DiCiccio, Sriram Vasudevan, Kinjal Basu, Krishnaram Kenthapadi, and Deepak Agarwal. Evaluating Fairness Using Permutation Tests. Proceedings of the 26th ACM SIGKDD International Conference on Knowledge Discovery & Data Mining, pages 1467–1477, August 2020. URL https://doi.org/10.1145/3394486.3403199.
- [11] Cynthia Dwork, Moritz Hardt, Toniann Pitassi, Omer Reingold, and Rich Zemel. Fairness Through Awareness. ITCS '12: Proceedings of the 3rd Innovations in Theoretical Computer Science Conference, pages 214–226, 2012. doi: 10.1145/2090236.2090255. URL http: //arxiv.org/abs/1104.3913.
- [12] Andre Esteva, Brett Kuprel, Roberto A. Novoa, Justin Ko, Susan M. Swetter, Helen M. Blau, and Sebastian Thrun. Dermatologist-level classification of skin cancer with deep neural networks. *Nature*, 542(7639):115–118, February 2017. ISSN 0028-0836, 1476-4687. doi: 10.1038/ nature21056. URL http://www.nature.com/articles/nature21056.

- [13] Josh Gardner, Christopher Brooks, and Ryan Baker. Evaluating the Fairness of Predictive Student Models Through Slicing Analysis. In *Proceedings of the 9th International Conference* on Learning Analytics & Knowledge, pages 225–234, Tempe AZ USA, March 2019. ACM. ISBN 978-1-4503-6256-6. doi: 10.1145/3303772.3303791. URL https://dl.acm.org/ doi/10.1145/3303772.3303791.
- [14] Xiaotian Han, Zhimeng Jiang, Hongye Jin, Zirui Liu, Na Zou, Qifan Wang, and Xia Hu. Retiring ΔDP: New distribution-level metrics for demographic parity. In arXiv:2301.13443. arXiv, January 2023. doi: 10.48550/arXiv.2301.13443. URL http://arxiv.org/abs/2301. 13443.
- [15] Moritz Hardt, Eric Price, and Nati Srebro. Equality of Opportunity in Supervised Learning. In Proceedings of the 30th International Conference on Neural Information Processing Systems, 29, page 9, 2016. doi: 10.5555/3157382.3157469.
- [16] Kenneth Holstein, Bruce M. McLaren, and Vincent Aleven. Student Learning Benefits of a Mixed-Reality Teacher Awareness Tool in AI-Enhanced Classrooms. In Carolyn Penstein Rosé, Roberto Martínez-Maldonado, H. Ulrich Hoppe, Rose Luckin, Manolis Mavrikis, Kaska Porayska-Pomsta, Bruce McLaren, and Benedict Du Boulay, editors, *Artificial Intelligence in Education*, volume 10947, pages 154–168. Springer International Publishing, Cham, 2018. ISBN 978-3-319-93842-4 978-3-319-93843-1. doi: 10.1007/978-3-319-93843-1_12. URL http://link.springer.com/10.1007/978-3-319-93843-1_12.
- [17] Nathan Kallus and Angela Zhou. The Fairness of Risk Scores Beyond Classification: Bipartite Ranking and the XAUC Metric. In H. Wallach, H. Larochelle, A. Beygelzimer, F. d'Alché-Buc, E. Fox, and R. Garnett, editors, *Proceedings of the 33rd International Conference on Neural Information Processing Systems*, volume 32. Curran Associates, Inc., 2019. doi: Proceedingsofthe33rdInternationalConferenceonNeuralInformationProcessingSystems.
- [18] Toshihiro Kamishima, Shotaro Akaho, Hideki Asoh, and Jun Sakuma. Fairness-Aware Classifier with Prejudice Remover Regularizer. In David Hutchison, Takeo Kanade, Josef Kittler, Jon M. Kleinberg, Friedemann Mattern, John C. Mitchell, Moni Naor, Oscar Nierstrasz, C. Pandu Rangan, Bernhard Steffen, Madhu Sudan, Demetri Terzopoulos, Doug Tygar, Moshe Y. Vardi, Gerhard Weikum, Peter A. Flach, Tijl De Bie, and Nello Cristianini, editors, *Machine Learning and Knowledge Discovery in Databases*, volume 7524, pages 35–50. Springer Berlin Heidelberg, Berlin, Heidelberg, 2012. ISBN 978-3-642-33485-6 978-3-642-33486-3_3. URL http://link.springer.com/10.1007/978-3-642-33486-3_3.
- [19] Jon Kleinberg, Sendhil Mullainathan, and Manish Raghavan. Inherent Trade-Offs in the Fair Determination of Risk Scores. In arXiv:1609.05807, 2018. URL http://arxiv.org/abs/ 1609.05807.
- [20] Alina Köchling and Marius Claus Wehner. Discriminated by an algorithm: a systematic review of discrimination and fairness by algorithmic decision-making in the context of hr recruitment and hr development. *Business Research*, 2020.
- [21] Nikita Kozodoi, Johannes Jacob, and Stefan Lessmann. Fairness in credit scoring: Assessment, implementation and profit implications. *European Journal of Operational Research*, 297(3): 1083–1094, 2022.
- [22] Ananya Kumar, Percy Liang, and Tengyu Ma. Verified Uncertainty Calibration. In Advances in Neural Information Processing SystemsNeurIPS, volume 32. arXiv, 2019. URL http: //arxiv.org/abs/1909.10155.
- [23] Kweku Kwegyir-Aggrey, Rebecca Santorella, and Sarah M. Brown. Everything is Relative: Understanding Fairness with Optimal Transport. *arXiv:2102.10349 [cs]*, February 2021. URL http://arxiv.org/abs/2102.10349.
- [24] Jeff Larson, Surya Mattu, Lauren Kirchner, and Julia Angwin. How we analyzed the compas recidivism algorithm. *ProPublica* (5 2016), 9(1):3–3, 2016.

- [25] Lydia T. Liu, Max Simchowitz, and Moritz Hardt. The implicit fairness criterion of unconstrained learning. In *International Conference on Machine Learning*, pages 4051–4060. arXiv, January 2019. URL http://arxiv.org/abs/1808.10013.
- [26] Karima Makhlouf and Sami Zhioua. On the Applicability of Machine Learning Fairness Notions. ACM SIGKDD Explorations Newsletter, 1(23):14–23, 2021. URL https://doi. org/10.1145/3468507.3468511.
- [27] Karima Makhlouf, Sami Zhioua, and Catuscia Palamidessi. Survey on causal-based machine learning fairness notions. *arXiv preprint arXiv:2010.09553*, 2020.
- [28] Alexey Miroshnikov, Konstandinos Kotsiopoulos, Ryan Franks, and Arjun Ravi Kannan. Modelagnostic bias mitigation methods with regressor distribution control for Wasserstein-based fairness metrics. arXiv:2111.11259 [cs, math], November 2021. URL http://arxiv.org/ abs/2111.11259.
- [29] Alexey Miroshnikov, Konstandinos Kotsiopoulos, Ryan Franks, and Arjun Ravi Kannan. Wasserstein-based fairness interpretability framework for machine learning models. *Machine Learning*, 111(9):3307–3357, September 2022. ISSN 0885-6125, 1573-0565. doi: 10.1007/s10994-022-06213-9. URL https://link.springer.com/10.1007/ s10994-022-06213-9.
- [30] Shira Mitchell, Eric Potash, Solon Barocas, Alexander D'Amour, and Kristian Lum. Algorithmic Fairness: Choices, Assumptions, and Definitions. Annual Review of Statistics and Its Application, 8(1):141–163, March 2021. ISSN 2326-8298, 2326-831X. doi: 10.1146/annurev-statistics-042720-125902. URL https://www.annualreviews.org/doi/ 10.1146/annurev-statistics-042720-125902.
- [31] Hamed Nilforoshan, Johann D Gaebler, Ravi Shroff, and Sharad Goel. Causal conceptions of fairness and their consequences. In *International Conference on Machine Learning*, pages 16848–16887. PMLR, 2022.
- [32] Geoff Pleiss, Manish Raghavan, Felix Wu, Jon Kleinberg, and Kilian Q Weinberger. On fairness and calibration. *Advances in neural information processing systems*, 30, 2017.
- [33] Emilee Rader and Rebecca Gray. Understanding User Beliefs About Algorithmic Curation in the Facebook News Feed. In *Proceedings of the 33rd Annual ACM Conference on Human Factors in Computing Systems*, pages 173–182, Seoul Republic of Korea, April 2015. ACM. ISBN 978-1-4503-3145-6. doi: 10.1145/2702123.2702174. URL https://dl.acm.org/ doi/10.1145/2702123.2702174.
- [34] Kailash Karthik Saravanakumar. The Impossibility Theorem of Machine Fairness A Causal Perspective, January 2021. URL http://arxiv.org/abs/2007.06024.
- [35] Roman Schefzik, Julian Flesch, and Angela Goncalves. Fast identification of differential distributions in single-cell RNA-sequencing data with waddR. *Bioinformatics*, 37(19):3204– 3211, October 2021. ISSN 1367-4803, 1460-2059. doi: 10.1093/bioinformatics/btab226. URL https://academic.oup.com/bioinformatics/article/37/19/3204/6207964.
- [36] Robin Vogel, Aurélien Bellet, and Stephan Clémençon. Learning Fair Scoring Functions: Bipartite Ranking under ROC-based Fairness Constraints. In arXiv:2002.08159, pages 784–792, February 2021. URL http://arxiv.org/abs/2002.08159.
- [37] Zhenhuan Yang, Yan Lok Ko, Kush R Varshney, and Yiming Ying. Minimax auc fairness: Efficient algorithm with provable convergence. *arXiv preprint arXiv:2208.10451*, 2022.
- [38] Muhammad Bilal Zafar, Isabel Valera, Manuel Gomez Rodriguez, and Krishna P. Gummadi. Fairness Beyond Disparate Treatment & Disparate Impact: Learning Classification without Disparate Mistreatment. In *Proceedings of the 26th International Conference on World Wide Web*, pages 1171–1180, Perth Australia, April 2017. International World Wide Web Conferences Steering Committee. ISBN 978-1-4503-4913-0. doi: 10.1145/3038912.3052660. URL https: //dl.acm.org/doi/10.1145/3038912.3052660.

- [39] Junzhe Zhang and Elias Bareinboim. Fairness in decision-making—the causal explanation formula. In Proceedings of the AAAI Conference on Artificial Intelligence, volume 32, 2018.
- [40] Junzhe Zhang and Elias Bareinboim. Equality of Opportunity in Classification: A Causal Approach. In S. Bengio, H. Wallach, H. Larochelle, K. Grauman, N. Cesa-Bianchi, and R. Garnett, editors, *Advances in Neural Information Processing Systems*, volume 31. Curran Associates, Inc., 2018. URL https://proceedings.neurips.cc/paper/2018/file/ ff1418e8cc993fe8abcfe3ce2003e5c5-Paper.pdf.

A Background definitions and results

A.1 Wasserstein-p-Distance

Definition A.1 (Wasserstein-p-Distance). The p^{th} Wasserstein distance between two probability measures μ and ν in $\mathcal{P}_p(\mathbb{R}^d)$ is defined as

$$W_p(\mu,\nu) := \left(\inf_{\gamma \in \Gamma(\mu,\nu)} \int_{\mathbb{R}^d \times \mathbb{R}^d} d(x,y)^p \,\mathrm{d}\gamma(x,y)\right)^{1/p},\tag{27}$$

where $\Gamma(\mu, \nu)$ denotes the collection of all measures on $\mathbb{R}^d \times \mathbb{R}^d$ with marginals μ and ν on the first and second factors respectively.

Corollary A.2. The Wasserstein metric may be equivalently defined by

$$W_p(\mu,\nu) = \left(\inf \mathbb{E}\left[d(X,Y)^p\right]\right)^{1/p},\tag{28}$$

where $\mathbb{E}[Z]$ denotes the expected value of a random variable Z and the infimum is taken over all joint distributions of the random variables X and Y with marginals μ and ν respectively.

If d = 1, the Wasserstein distance has a closed form. For this special case, we define W as a measure between two random variables.

Corollary A.3. Let X and Y be two random variables on \mathbb{R} and let F_X and F_Y denote their cumulative distribution functions. Then

$$W_p(X,Y) = \left(\int_0^1 |F_X^{-1}(s) - F_Y^{-1}(s)|^p \, ds\right)^{\frac{1}{p}}$$
(29)

Proposition A.4. Properties of the Wasserstein-Distance for d = 1:

- 1. For any real number a, $W_p(aX, aY) = |a|W_p(X, Y)$.
- 2. For any fixed vector x, $W_p(X + x, Y + x) = W_p(X, Y)$.
- 3. For independent X_1, \ldots, X_n and independent Y_1, \ldots, Y_n ,

$$W_p\left(\sum_{i=1}^n X_i, \sum_{i=1}^n Y_i\right) \le \sum_{i=1}^n W_p(X_i, Y_i).$$

A.2 Special case: One-dimensional Wasserstein-1-Distance

Corollary A.5. If p = 1 and X, Y are random variables on \mathbb{R} with cumulative distribution functions F_X and F_Y , then

$$W_1(X,Y) = \int_0^1 |F_X^{-1}(p) - F_Y^{-1}(p)|dp$$
(30)

$$= \int_{\mathbb{R}} |F_X(t) - F_Y(t)| dt.$$
(31)

Remark. The Wasserstein-1-distance is not invariant under monotone transformations (for instance, under scale transformations).

Remark. The Wasserstein distance is insensitive to small wiggles. For example if P is uniform on [0, 1] and Q has density $1 + \sin(2\pi kx)$ on [0, 1] then their Wasserstein distance is $\mathcal{O}(1/k)$.

Theorem A.6 (lower bound of W_1). The Wasserstein-distance is always greater or equal to the distance of the means:

$$W_1(X,Y) \ge |\mathbb{E}[X] - \mathbb{E}[Y]| \tag{32}$$

Proof. By Jensen inequality, as norm is convex.

Theorem A.7 (upper bound of W_1). For integers $p \leq q$,

$$W_p(X,Y) \le W_q(X,Y),\tag{33}$$

especially

$$W_1(X,Y) \le W_q(X,Y) \quad \forall q \ge 1.$$
(34)

Proof. By Jensen inequality, as $z \to z^{q/p}$ is convex.

A.3 Wasserstein-Distance of Quantile-Transformed Variables

Definition A.8 (Quantile-Transformed Wasserstein Distance). Let X, Y, Z be random variables on \mathbb{R} and let $F_X, F_Y, F_Z : \mathbb{R} \to [0, 1]$ denote their distribution functions and f_Z denote the density of Z. The (by Z) quantile-transformed Wasserstein Distance is then given by:

$$W_Z(X,Y) := W_1(F_Z(X), F_Z(Y))$$
(35)

$$= \int_{0}^{1} \left| F_{F_{Z}(X)}(t) - F_{F_{Z}(Y)}(t) \right| dt$$
(36)

$$= \int_{0}^{1} \left| F_X(F_Z^{-1}(t)) - F_Y(F_Z^{-1}(t)) \right| dt$$
(37)

$$= \int_{\mathbb{R}} |F_X(s) - F_Y(s)| f_Z(s) \, ds \tag{38}$$

Proposition A.9. Properties of the quantile-transformed Wasserstein-distance

- 1. For any real number $a \neq 0$, $W_Z(aX, aY) = W_{Z/|a|}(X, Y)$.
- 2. For any fixed vector x, $W_Z(X + x, Y + x) = W_{Z-x}(X, Y)$.

Remark. The quantile-transformed Wasserstein-1-distance is invariant under monotone transformations, for instance, under scale transformations: For a > 0:

$$W_Z(X,Y) = W_{aZ}(aX,aY).$$
(39)

A.4 Pushforward

The pushforward of a measure along a measurable function assigns to a subset the original measure of the preimage under the function of that subset.

Definition A.10. Let (X_1, Σ_1) and (X_2, Σ_2) be two measurable spaces, $f : X_1 \to X_2$ a measurable function and $\mu : \Sigma_1 \to [0, \infty]$ a measure on (X_1, Σ_1) . The pushforward of μ is defined as

$$f \# \mu : \Sigma_2 \to [0, \infty], f \# \mu(A) = \mu(f^{-1}(A)) \, \forall A \in \Sigma_2$$
 (40)

Corollary A.11. Let again (X_1, Σ_1) and (X_2, Σ_2) be two measurable spaces, $f : X_1 \to X_2$ a measurable function and $\mu : \Sigma_1 \to [0, \infty]$ a measure on (X_1, Σ_1) . If g is another measurable function on X_2 , then

$$\int_{X_2} g \circ f \, d\mu = \int_{X_1} g \, d(f \# \mu) \tag{41}$$

B Complete proofs

Lemma B.1. If we quantile-transform a continuous random variable $X \in \mathbb{R}$ by its own distribution F_X , the result will follow a uniform distribution in [0, 1]:

$$F_X(X) \sim \mathcal{U}[0,1], \text{ so } F_{F_X}(x) = x.$$

$$\tag{42}$$

Lemma B.2. Let X, Y be two random variables in \mathbb{R} with cumulative distribution functions F_X, F_Y . The cumulative distribution function of a random variable $Z = F_X(Y)$ is given by $F_Y(F_X^{-1}(z))$:

$$F_{F_X(Y)}(z) = F_Z(z) = \mathbb{P}(Z \le z) = \mathbb{P}(F_X(Y) \le z)$$

$$= \mathbb{P}(Y \le F_X^{-1}(z)) = F_Y(F_X^{-1}(z))$$

$$(43)$$

If F_X and F_Y are bijective and have the same support, then

$$F_{F_X(Y)} = F_{F_Y(X)}^{-1}.$$
(44)

Proof of Theorem 3.2. For x = EO:

$$\operatorname{bias}_{x}^{\mathcal{U}}(S|A=a, S|A=b) = \frac{1}{|\mathcal{S}|} \int_{\mathcal{S}} |\operatorname{c-bias}_{x}(S|A=a, S|A=b; s)| ds \tag{45}$$

$$= \frac{1}{|\mathcal{S}|} \int_{\mathcal{S}} |\mathbb{P}(S > s | A = b, Y = 0) - \mathbb{P}(S > s | A = a, Y = 0)| ds$$

$$\tag{46}$$

$$= \frac{1}{|\mathcal{S}|} \int_{\mathcal{S}} |(1 - F_{b0}(s)) - (1 - F_{a0}(s))| ds$$
(47)

$$= \frac{1}{|S|} \int_{S} |F_{a0}(s) - F_{b0}(s)| ds$$
(48)

$$\stackrel{A.5}{=} \frac{1}{|\mathcal{S}|} \cdot W_1(S|A = a, Y = 0, S|A = b, Y = 0).$$
(49)

For x = PE and x = IND the result follows similary. (ii) follows from Theorem A.6.

Proof of Theorem 3.4. For x = EO:

$$\operatorname{bias}_{x}^{S}(S|A=a, S|A=b) = \int_{S} |\operatorname{c-bias}_{x}(S|A=a, S|A=b; s)|f(s)ds \tag{50}$$

$$\stackrel{(41)}{=} \int_{\mathcal{S}} |\operatorname{c-bias}_{x}(S|A=a, S|A=b; s)| d(F^{-1} \# \mu)$$
(51)

$$= \int_{0}^{1} |\operatorname{c-bias}_{x}(S|A = a, S|A = b; F^{-1}(p))|dp$$
 (52)

$$\stackrel{3.2}{=} W_1(F_{F_S(S_{ay})}, F_{F_S(S_{by})}) \tag{53}$$

$$\stackrel{B.2}{=} W_1(F_{ay} \circ F_S^{-1}, F_{by} \circ F_S^{-1}) \tag{54}$$

For x = PE and x = IND the result follows similary. (ii) follows from Theorem A.6.

Proof of Theorem 4.4.

$$W_Z(X,Y) = \int_0^1 \left| F_{F_Z(X)}(s) - F_{F_Z(Y)}(s) \right| ds$$
(55)

$$\stackrel{B.2}{=} \int_{0}^{1} \left| F_X(F_Z^{-1}(s)) - F_Y(F_Z^{-1}(s)) \right| ds \tag{56}$$

$$= \int_{0}^{1} \left| (1 - F_X(F_Z^{-1}(1 - r)) - (1 - F_Y(F_Z^{-1}(1 - r))) \right| dr$$
(57)

$$= \int_0^1 |\operatorname{ROC}_{X,Z}(r) - \operatorname{ROC}_{Y,Z}(r)| \, dr \tag{58}$$

Proof of Theorem 4.5. Results directly from Def. 3.3 by using the additivity of the density in (9). \Box

Proof of Lemma 4.6. Using Theorem 4.4 and
$$\operatorname{ROC}_{X,X}(r) = r$$
.

Under additional assumptions, we can follow that a quantile-transformation by group a and b result in equal distances:

Lemma B.3. If S_{ay} and S_{by} have bijective cdfs and identical support, then

$$W_{S_{ay}}(S_{ay}, S_{by}) = W_{S_{by}}(S_{ay}, S_{by}) = W_{S_y}(S_{ay}, S_{by}).$$
(59)

Proof of Lemma B.3. We show more general: If X, Y are two random variables on an interval I in \mathbb{R} with cdfs F_X and F_Y that are bijective on I

$$W_X(X,Y) = W_Y(X,Y)$$

By Lemma B.1 and by Lemma B.2, it follows

$$W_X(X,Y) \stackrel{(38)}{=} \int_0^1 \left| F_{F_X(X)}(t) - F_{F_X(Y)}(t) \right| dt \tag{60}$$

$$\stackrel{B.1}{=} \int_0^1 \left| t - F_{F_X(Y)}(t) \right| dt \tag{61}$$

$$\stackrel{B.2}{=} \int_{0}^{1} \left| t - F_{F_{Y}(X)}^{-1}(t) \right| dt \tag{62}$$

$$\stackrel{B.1}{=} \int_0^1 \left| F_{F_Y(Y)}^{-1}(t) - F_{F_Y(X)}^{-1}(t) \right| dt \tag{63}$$

$$\stackrel{(31)}{=} W_Y(X,Y) \tag{64}$$

It follows for $Z = w_1 X + w_2 Y$:

$$W_Z(X,Y) = w_1 W_X(X,Y) + w_2 W_Y(X,Y)$$

= $W_X(X,Y) = W_Y(X,Y)$ (65)

and Lemma B.3 as a special case.

Lemma B.3 implies that quantile-transformation can under the above assumptions be performed on either of the two groups or the whole sample with the same result. Under the same assumptions, ROC, AUROC and Gini become symmetrical, i.e. $\text{ROC}_{S_{ay},S_{by}} = \text{ROC}_{S_{by},S_{ay}}$.

Proof of Theorem 4.8. Using Proposition 4.5 and Lemma 4.6:

$$bias_{EO}^{S}(S|A = a, S|A = b) = W_{S}(S_{a0}, S_{b0})$$
(66)

$$\stackrel{4.5}{=} w_{a0} W_{S_{a0}}(S_{a0}, S_{b0}) + w_{b0} W_{S_{b0}}(S_{a0}, S_{b0})$$

$$+ w_{a0} W_{G_{a0}}(S_{a0}, S_{b0}) + w_{b0} W_{G_{a0}}(S_{a0}, S_{b0})$$

$$(67)$$

$$+ w_{a1} W_{S_{a1}}(S_{a0}, S_{b0}) + w_{b1} W_{S_{b1}}(S_{a0}, S_{b0})$$

$$= w_{a0} \int |\operatorname{ROC}_{S_{a0},S_{b0}}(r) - r| dr + w_{b0} \int |\operatorname{ROC}_{S_{b0},S_{a0}}(r) - r| dr$$

$$+ w_{a1} \int |\operatorname{ROC}_{S_{a0},S_{a1}}(r) - \operatorname{ROC}_{S_{b0},S_{a1}}(r)| dr$$

$$+ w_{b1} \int |\operatorname{ROC}_{S_{a0},S_{b1}}(r) - \operatorname{ROC}_{S_{b0},S_{b1}}(r)| dr$$

$$(68)$$

For predictive equality, the result follows similarly.

Proof of Theorem 4.10. From theorem 4.8 follows by triangle-Inequality:

$$\operatorname{bias}_{\mathrm{EO}}^{S}(S|A=a, S|A=b) + \operatorname{bias}_{\mathrm{PE}}^{S}(S|A=a, S|A=b) = W_{S}(S_{a0}, S_{b0}) + W_{S}(S_{a1}, S_{b1})$$
(69)

$$\geq w_0 \int |\operatorname{ROC}_{S_{a0}, S_{b0}}(r) - r| \, dr + w_1 \int |\operatorname{ROC}_{S_{a1}, S_{b1}}(r) - r| \, dr \tag{70}$$

$$+\min(w_{a0}, w_{a1}) \cdot \operatorname{bias_{xROC}} + \min(w_{b0}, w_{b1}) \cdot \operatorname{bias_{xROC}} \\ \ge w_0 \int |\operatorname{ROC}_{S_{a0}, S_{b0}}(r) - r| \, dr + w_1 \int |\operatorname{ROC}_{S_{a1}, S_{b1}}(r) - r| \, dr$$
(71)

 $+2\min(w_{a0}, w_{a1}, w_{b0}, w_{b1}) \cdot \operatorname{bias}_{\mathrm{xROC}}$

and also

$$\operatorname{bias}_{\mathsf{EO}}^{S}(S|A=a, S|A=b) + \operatorname{bias}_{\mathsf{PE}}^{S}(S|A=a, S|A=b) = W_{S}(S_{a0}, S_{b0}) + W_{S}(S_{a1}, S_{b1})$$
(72)

$$\geq w_0 \int |\operatorname{ROC}_{S_{a0},S_{b0}}(r) - r| \, dr + w_1 \int |\operatorname{ROC}_{S_{a1},S_{b1}}(r) - r| \, dr \tag{73}$$

$$+ \min(w_{a1}, w_{b1}) \cdot \operatorname{blas}_{\mathrm{ROC}} + \min(w_{a0}, w_{b0}) \cdot \operatorname{blas}_{\mathrm{ROC}}$$

$$\geq w_0 \int |\operatorname{ROC}_{S_{a0}, S_{b0}}(r) - r| \, dr + w_1 \int |\operatorname{ROC}_{S_{a1}, S_{b1}}(r) - r| \, dr$$

$$+ 2\min(w_{a0}, w_{a1}, w_{b0}, w_{b1}) \cdot \operatorname{blas}_{\mathrm{ROC}}$$

$$(74)$$

By additionally using Corollary 4.7, we get

$$bias_{EO}^{S} + bias_{PE}^{S} = W_{S}(S_{a0}, S_{b0}) + W_{S}(S_{a1}, S_{b1})$$

$$\geq \min(w_{a0}, w_{a1}, w_{b0}, w_{b1}) \cdot (bias_{ROC} + bias_{xROC}) + \frac{w_{0}}{2} \operatorname{Gini}(S_{a0}, S_{b0}) + \frac{w_{1}}{2} \operatorname{Gini}(S_{a1}, S_{b1})$$
(75)

As $\frac{w_i}{2} \leq \frac{\min(w_{a0}, w_{a1}, w_{b0}, w_{b1})}{2}$ for i = 0, 1, we can combine all weights to get

$$bias_{EO}^{S} + bias_{PE}^{S} = W_{S}(S_{a0}, S_{b0}) + W_{S}(S_{a1}, S_{b1})$$

$$(76)$$

$$w_{0} = w_{1} + w_{2} + w_{3} + w_{4} + w_{5} + w_{5$$

$$\geq \min(w_{a0}, w_{a1}, w_{b0}, w_{b1}) \cdot (\operatorname{bias}_{\operatorname{ROC}} + \operatorname{bias}_{\operatorname{xROC}}) + \frac{w_0}{2} \operatorname{Gini}(S_{a0}, S_{b0}) + \frac{w_1}{2} \operatorname{Gini}(S_{a1}, S_{b1})$$
(77)
$$\min(w_{a0}, w_{a1}, w_{b0}, w_{b1}) \cdot (\operatorname{bias}_{\operatorname{ROC}} + \operatorname{bias}_{\operatorname{xROC}}) + \frac{w_0}{2} \operatorname{Gini}(S_{a0}, S_{b0}) + \frac{w_1}{2} \operatorname{Gini}(S_{a1}, S_{b1})$$
(77)

$$\geq \frac{\min(w_{a0}, w_{a1}, w_{b0}, w_{b1})}{2} (\operatorname{bias}_{\operatorname{ROC}} + \operatorname{bias}_{\operatorname{xROC}} + \operatorname{Gini}(S_{a0}, S_{b0}) + \operatorname{Gini}(S_{a1}, S_{b1}))$$
(78)

Note, that if F_{ay} and F_{by} have identical supports and permit an inverse, then $\text{Gini}(S_{ay}, S_{by}) = \text{Gini}(S_{by}, S_{ay})$. If this symmetry is not fulfilled, the minimum of both must be used on the right side.

Proof of Theorem 4.12. Let $bias_{EO}(S|A = a, S|A = b) = 0$, it follows $F_{b0} = F_{a0}$ almost everywhere. Then

$$\operatorname{bias}_{\operatorname{ROC}}(S|A=a, S|A=b) \tag{79}$$

$$= \int_{0}^{1} |F_{b0}(F_{b1}^{-1}(s)) - F_{a0}(F_{a1}^{-1}(s))| ds$$
(80)

$$= \int_{0}^{1} |F_{a0}(F_{b1}^{-1}(s)) - F_{b0}(F_{a1}^{-1}(s))| ds$$
(81)

$$\operatorname{bias}_{\mathsf{xROC}}(S|A=a, S|A=b) \tag{82}$$

For predictive equality, the statement follows similarly.

type of bias	total	pos.	neg.	p-value
$bias^{S}_{EO}$	0.1613	0%	100%	<0.01
$bias^{S}_{PE}$	0.1543	0%	100%	<0.01
$bias^{S}_{CALL}$	0.0340	79%	21%	0.19
bias _{ROC}	0.0160	46%	54%	0.38
bias _{xROC}	0.3723	0%	100%	<0.01
$\mathrm{bias}_{\mathrm{EO}}^{\mathcal{U}}$	0.1515	0%	100%	<0.01
$\mathrm{bias}_{\mathrm{PE}}^{\mathcal{U}}$	0.1633	0%	100%	<0.01
$\mathrm{bias}_{\mathrm{CALI}}^{\mathcal{U}}$	0.0370	78%	22%	0.17

Table C1: Bias of COMPAS score (complete table)

C Experiments

We perform experiments in python using the COMPAS dataset², the Adult dataset³ and the German Credit dataset⁴. Empirical implementations of Wasserstein-distance (*scipy.wasserstein_distance*), calibration curves (*sklearn.calibration_curve*) and ROC curves (*sklearn.metrics.roc_curve*) were used.

C.1 Statistical Testing

We perform permutation tests [10, 35] with 100 permutations and one pseudocount to determine the statistical significance of the calculated biases under the null hypothesis of group parity. The calibration biases were calculated using 50 bins.

C.2 Details on COMPAS experiments

Full results are shown in Table C1.

C.3 Details on German Credit data experiments

Both models have been trained on 70% of the dataset and evaluated on the remaining samples. We used min-max-scaling on continuous features and one-hot-encoding for categorical features. Full results are shown in Table C2. As the sample size is relatively small, it happens that even the large calibration biases are not statistically significant.

C.4 Details on Adult experiments

All three models have been trained on 70% of the dataset and evaluated on the remaining samples. We removed the feature *relationship*, which is highly entangled with *sex* through the categories *husband* and *wife* and we engineered the remaining features to merge rare categories. We used min-max-scaling on continuous features and one-hot-encoding for categorical features. Fig. C1-C3 show the score distributions of the three scores on the testset. Fig. C4 and C5 show the classifier bias over the score range and the quantile-transformed score range.

²https://raw.githubusercontent.com/propublica/compas-analysis/master/compas-scores-two-years.csv

³https://archive.ics.uci.edu/ml/machine-learning-databases/adult/adult.data

⁴https://www.kaggle.com/datasets/uciml/german-credit?resource=download

type of bias	Model	total bias	pos.	neg.	p-value
$bias_{EO}^S$	LogR	0.083	0%	100%	0.05
	LogR (debiased)	0.048	93%	7%	0.29
$bias_{PE}^{S}$	LogR	0.092	0%	100%	0.12
	LogR (debiased)	0.010	62%	38%	1
$bias_{CALI}^S$	LogR	0.291	46%	54%	0.28
	LogR (debiased)	0.299	58%	42%	0.29
bias _{ROC}	LogR	0.049	98%	2%	0.75
	LogR (debiased)	0.051	98%	2%	0.71
bias _{xROC}	LogR	0.133	0%	100%	0.01
	LogR (debiased)	0.056	93%	7%	0.60
$\mathrm{bias}_{\mathrm{EO}}^{\mathcal{U}}$	LogR	0.041	3%	97%	0.19
	LogR (debiased)	0.036	97%	3%	0.23
$\mathrm{bias}_{\mathrm{PE}}^{\mathcal{U}}$	LogR	0.078	1%	99%	0.15
	LogR (debiased)	0.024	74%	26%	0.99
$\mathrm{bias}^{\mathcal{U}}_{\mathrm{CALI}}$	LogR	0.212	36%	64%	0.68
	LogR (debiased)	0.193	67%	33%	0.85

Table C2: Bias of models for German Credit data (complete table)



Figure C1: Distribution of logistic regression scores, trained on Adult data.



Figure C2: Distribution of logistic regression scores, trained on Adult data without protected attribute.



Figure C3: Distribution of XGBoost scores trained on Adult data.



Figure C4: Equal opportunity bias $bias_{EO}^S$ of the logistic regression model trained on the Adult dataset. $bias_{EO}^S$ is equal to the area under the curve of the true positive rate difference. The area is colored according to the group for which the bias part is favorable.



Figure C5: Equal opportunity bias $\operatorname{bias}_{EO}^{\mathcal{U}}$ of the logistic regression model trained on the Adult dataset. $\operatorname{bias}_{EO}^{S}$ is equal to the area under the curve of the true positive rate difference. The area is colored according to the group for which the bias part is favorable.