Space-time hybridizable discontinuous Galerkin method for advection-diffusion on deforming domains: The advection-dominated regime

Y. Wang[∗] S. Rhebergen[†]

August 24, 2023

Abstract

We analyze a space-time hybridizable discontinuous Galerkin method to solve the timedependent advection-diffusion equation on deforming domains. We prove stability of the discretization in the advection-dominated regime by using weighted test functions and derive a priori space-time error estimates. A numerical example illustrates the theoretical results.

1 Introduction

In this paper we analyze a space-time hybridizable discontinuous Galerkin (HDG) method for the time-dependent advection-diffusion problem on a time-dependent polygonal $(d = 2)$ or polyhedral $(d=3)$ domain $\Omega(t) \subset \mathbb{R}^d$, that evolves continuously in the time interval $t \in [0, T]$. This problem is given by

$$
\partial_t u + \overline{\nabla} \cdot (\overline{\beta} u) - \varepsilon \overline{\nabla}^2 u = f \quad \text{in } \Omega(t), \ 0 < t \le T,\tag{1.1}
$$

in which $\overline{\nabla}=(\partial_{x_1},\partial_{x_2},\ldots,\partial_{x_d})$ denotes the spatial gradient, $\bar{\beta}$ is the given divergence-free advective field, $\varepsilon > 0$ is the constant diffusion coefficient, and f is a forcing term. The focus in this paper is the advection-dominated regime ($\varepsilon \ll 1$).

Many different finite element methods have been designed and analyzed for advection-diffusion problems. These include, for example, streamline upwind Petrov–Galerkin (SUPG) [\[7,](#page-29-0) [9,](#page-29-1) [29,](#page-30-0) [17\]](#page-29-2), continuous interior penalty (CIP) [\[11,](#page-29-3) [8,](#page-29-4) [10\]](#page-29-5), discontinuous Galerkin (DG) [\[14,](#page-29-6) [15,](#page-29-7) [19,](#page-29-8) [36\]](#page-31-0), and HDG [\[13,](#page-29-9) [18,](#page-29-10) [34,](#page-30-1) [44\]](#page-31-1) methods. Of particular interest to the current work is the analysis presented by Ayuso and Marini [\[2\]](#page-28-0). They introduce a weighted test function to analyze a DG method for the stationary advection-diffusion-reaction problem. For this they assume that $\bar{\beta}$ has no closed curves nor stationary points which implies the existence of a smooth function ψ such that $\bar{\beta} \cdot \overline{\nabla} \psi(x) > b_0$ for some constant $b_0 > 0$ depending on the inverse of the diameter of the domain. The function ψ is used to define a weighting function $\varphi := \exp(-\psi) + \chi$, with χ a free to choose positive constant, which is used to show stability of the DG method. They are able to show stability for the advection-diffusion-reaction problem independent of the diffusion parameter ε , and therefore stability of the DG method in the advection-dominated regime, in a norm that provides control of the streamline derivative. Furthermore, their analysis also holds for the advection-diffusion problem

^{*}Department of Applied Mathematics, University of Waterloo, ON, Canada (<yuan.wang@uwaterloo.ca>), [http:](http://orcid.org/0009-0006-8092-4378) [//orcid.org/0009-0006-8092-4378](http://orcid.org/0009-0006-8092-4378)

[†]Department of Applied Mathematics, University of Waterloo, ON, Canada (<srheberg@uwaterloo.ca>), [http:](http://orcid.org/0000-0001-6036-0356) [//orcid.org/0000-0001-6036-0356](http://orcid.org/0000-0001-6036-0356)

with solenoidal advective field, thereby relaxing the usual coercivity condition that $\mu(x) + \frac{1}{2}\overline{\nabla}\cdot\overline{\beta} > 0$, where $\mu(x)$ is the reaction coefficient. The same idea is also used to analyze the HDG method for the stationary advection-diffusion problem in the advection-dominated regime in Fu et al. [\[21\]](#page-30-2).

To discretize the advection-diffusion problem on a time-dependent domain we consider a fully discrete space-time formulation of eq. (1.1) using DG time stepping [\[28,](#page-30-3) [27\]](#page-30-4). DG time stepping can be combined with different spatial discretizations. On fixed domains, for example, DG time stepping combined with SUPG was analyzed for the advection-diffusion equation in Hughes et al. [\[26\]](#page-30-5), while space-time DG, in which DG is applied both in space and time, was analyzed for a nonlinear advection-diffusion problem in Feistauer et al. [\[20\]](#page-30-6). The space-time DG method for the (linear) advection-diffusion problem on a time-dependent domain was analyzed in Sudirham et al. [\[42\]](#page-31-2) by considering the space-time discretization on a space-time mesh consisting of anisotropic (in space and time) elements. This enabled them to obtain error estimates in terms of the spatial mesh size and the time step, extending the analysis of DG methods for the stationary advection-diffusion problem on anisotropic spatial meshes [\[22\]](#page-30-7) to space-time.

DG methods are known to be expensive due to the large number of degrees-of-freedom (dofs) compared to, for example, a continuous finite element method on the same mesh. This resulted in the introduction of the HDG method by Cockburn et al. [\[16\]](#page-29-11), a DG method that uses static condensation to reduce the number of globally coupled dofs. The extension of HDG to spacetime, in which HDG is used to discretize a PDE in both space and time, was presented in [\[37,](#page-31-3) [38\]](#page-31-4). Furthermore, it was demonstrated in Sivas et al. [\[40\]](#page-31-5) that nonsymmetric algebraic multigrid, based on approximate ideal restriction [\[32,](#page-30-8) [31\]](#page-30-9), is an effective preconditioner for space-time HDG discretizations of the advection-diffusion problem in the advection-dominated regime.

The space-time HDG method for eq. [\(1.1\)](#page-0-0) was analyzed in Kirk et al. [\[30\]](#page-30-10) following the spacetime anisotropic framework used in Sudirham et al. [\[42\]](#page-31-2). However, despite the space-time HDG method performing well in practice for $\varepsilon \ll 1$, the well-posedness result proven in Kirk et al. [\[30\]](#page-30-10) does not hold in the advection dominated regime. To address this discrepancy between practice and theory, we revisit the analysis in Kirk et al. [\[30\]](#page-30-10) and focus on its extension to the advectiondominated regime. Like the analysis of Ayuso and Marini [\[2\]](#page-28-0) discussed above, we will use a weighted test function. However, our weighting function is constructed explicitly for the time-dependent advection-diffusion equation and only depends on the time variable t and final time T , not on the space variable x. We prove stability of the space-time HDG method in a norm providing control on the streamline derivative and present error estimates for the advection-dominated regime that are also valid on moving meshes.

The paper is organized as follows. In section [2,](#page-1-0) we introduce the space-time formulation of the time-dependent advection-diffusion problem. In section [3](#page-2-0) we describe the finite element spaces, present inequalities that will be used in the analysis, and the space-time HDG discretization of the advection-diffusion problem. The main result of section [4](#page-7-0) is inf-sup stability of the discretization. The a priori error analysis is then presented in section [5.](#page-18-0) We provide pre-asymptotic and asymptotic convergence rates, with the transition from the former to the latter featuring a drop in the rate of convergence from $p + 1/2$ to p, with p the degree of the polynomial approximation. In section [6,](#page-21-0) we numerically confirm the error analysis by solving a time-dependent advection-diffusion problem on a deforming mesh that contains hanging nodes in both spatial and temporal directions.

2 The space-time formulation of the advection-diffusion problem

In this section we formulate eq. (1.1) as a problem in $(d+1)$ -dimensional space-time. For this, we define the $(d+1)$ -dimensional polyhedral space-time domain as $\mathcal{E} := \{(t,x) : x \in \Omega(t), 0 < t < T\} \subset$

 \mathbb{R}^{d+1} . Its boundary, $\partial \mathcal{E}$, consists of $\Omega(0) := \{(t,x) \in \partial \mathcal{E} : t = 0\}$, $\Omega(T) := \{(t,x) \in \partial \mathcal{E} : t = T\}$, and $\mathcal{Q}_{\mathcal{E}} := \{(t,x) \in \partial \mathcal{E} : 0 < t < T\}$. The outward space-time normal vector to $\partial \mathcal{E}$ is denoted by $n := (n_t, \bar{n})$, where n_t and \bar{n} are the temporal and spatial components of the space-time normal vector, respectively. Introducing the space-time advective field $\beta := (1, \beta)$ and space-time gradient operator $\nabla := (\partial_t, \overline{\nabla})$, the space-time formulation of eq. (1.1) is given by

$$
\nabla \cdot (\beta u) - \varepsilon \overline{\nabla}^2 u = f \text{ in } \mathcal{E}.
$$
 (2.1a)

We consider a nonoverlapping partition of the domain boundary, $\partial \mathcal{E} = \partial \mathcal{E}_D \cup \partial \mathcal{E}_N$, and impose the boundary conditions

$$
-\zeta^- u\beta \cdot n + \varepsilon \overline{\nabla} u \cdot \bar{n} = g \text{ on } \partial \mathcal{E}_N, \qquad u = 0 \text{ on } \partial \mathcal{E}_D.
$$
 (2.1b)

The Dirichlet $\partial \mathcal{E}_D$ and Neumann $\partial \mathcal{E}_N$ boundaries are defined here by:

$$
\partial \mathcal{E}_D := \left\{ (t, x) : x \in \Gamma_D(t), 0 < t \le T \right\}, \qquad \partial \mathcal{E}_N := \left\{ (t, x) : x \in \Gamma_N(t) \cup \Omega(0) \cup \Omega(T), 0 < t \le T \right\},\
$$

where we also prescribe a nonoverlapping partition of the boundary of $\Omega(t)$, i.e., $\partial \Omega(t) = \Gamma_D(t) \cup$ $\Gamma_N(t)$. Furthermore, ζ^- is an indicator function for the inflow (where $\beta \cdot n < 0$) part of the boundary of E. Therefore, the boundary condition on $\partial \mathcal{E}_N$ also imposes the initial condition $u(x, 0) = g(x)$ on $\Omega(0)$.

We assume that the forcing term f lies in $L^2(\mathcal{E})$ and that the Neumann boundary data g lies in $L^2(\partial \mathcal{E}_N)$. Furthermore, we assume that $\bar{\beta} \in [W^{1,\infty}(\mathcal{E})]^d$, $\|\bar{\beta}\|_{L^{\infty}(\mathcal{E})} \leq 1$ and, following Ayuso and Marini [\[2\]](#page-28-0), that $\|\bar{\beta}\|_{W^{1,\infty}(\mathcal{E})} \leq c \|\bar{\beta}\|_{L^{\infty}(\mathcal{E})} \leq c.$

3 The space-time hybridizable discontinuous Galerkin method

3.1 Description of space-time slabs, faces, and elements

An initial partition of the space-time domain $\mathcal E$ consists of dividing the time interval $[0, T]$ into time levels $0 = t_0 < t_1 < \cdots < t_N = T$ and defining the *n*th time interval as $I_n = (t_n, t_{n+1})$. The space-time domain is divided into space-time slabs $\mathcal{E}^n := \mathcal{E} \cap (I_n \times \mathbb{R}^d)$, which are then divided into space-time elements, $\mathcal{E}^n = \cup_j \mathcal{K}_j^n$. To construct the space-time element \mathcal{K}_j^n , we divide the domain $\Omega(t_n)$ into nonoverlapping spatial elements K_j^n so that $\Omega(t_n) = \cup_j K_j^{n'}$. Let Υ be the transformation describing the deformation of the domain. The spatial elements K_j^{n+1} at t_{n+1} are obtained by mapping the nodes of the elements K_j^n into their new position via the transformation T. Each space-time element \mathcal{K}_j^n is obtained by connecting the elements K_j^n and K_j^{n+1} via linear interpolation in time following van der Vegt and van der Ven [\[43\]](#page-31-6). We denote the set of all spacetime elements tessellating the space-time domain by \mathcal{T}_h .

The boundary of a space-time element K with $t \in (t_*, t^*)$ is partitioned as $\partial \mathcal{K} = \mathcal{Q}_{\mathcal{K}} \cup \mathcal{R}_{\mathcal{K}}$ where $\mathcal{R}_\mathcal{K} := K_* \cup K^*$, $\mathcal{Q}_\mathcal{K} \cap \mathcal{R}_\mathcal{K} = \emptyset$, and where K_* denotes the face of \mathcal{K} at time $t = t_*$ and K^* denotes the face of K at time t^* . On ∂K , the outward unit space-time normal vector is denoted by $n^{\mathcal{K}} = (n^{\mathcal{K}}_t, \bar{n}^{\mathcal{K}})$, where $n^{\mathcal{K}}_t$ and $\bar{n}^{\mathcal{K}}$ are the temporal and spatial components of the space-time normal vector, respectively. Note that \bar{n} is the zero vector on an R-face, i.e., that $\bar{n} = 0$ on K^* and K_* , and that $\bar{n} \neq 0$ on a $\mathcal{Q}\text{-face}$.

The set of all faces in the mesh shared between at most two elements is denoted by \mathcal{F}_h . Within this set, we denote by \mathcal{F}_h^i , \mathcal{F}_{h}^b , $\mathcal{F}_{Q,h}$, and $\mathcal{F}_{R,h}$ all the interior faces, boundary faces, Q -faces, and R-faces respectively. When the mesh is viewed from an element's perspective, we denote by $\partial \mathcal{T}_h$ the set of element boundaries. Within this set, we denote by \mathcal{Q}_h and \mathcal{R}_h the set of element boundary

(a) A spatial mesh $\Omega(t_n) = \bigcup_j K_j^n$ (b) A space-time slab \mathcal{E}_n . The faces dyed in red form $\mathcal{Q}_{\mathcal{K}}$ while $\mathcal{R}_{\mathcal{K}}$ is the union of K^* and K_* .

Figure 1: Illustration of a moving spatial domain $\Omega(t) \subset \mathbb{R}^d$ for $t \in I_n$ resulting in the space-time slab $\mathcal{E}^n \subset \mathbb{R}^{(d+1)}$ (with $d=2$). Local time-stepping within a space-time slab is featured in fig. [1b.](#page-3-0) Here $K^* = K^{n+1}$ and $K_* = K^n$.

Q-faces and element boundary R-faces, respectively. Finally, the union of all faces in \mathcal{F}_h is denoted by Γ. An illustration of a $(d+1)$ -dimensional space-time mesh in slab \mathcal{E}^n , with $d=2$, is shown in fig. [1.](#page-3-0)

To define the finite element spaces, we require the mapping $\Phi_{\mathcal{K}}$ between a fixed reference element $\hat{\mathcal{K}} = (-1, 1)^{d+1}$ and space-time element $\mathcal{K} \in \mathcal{T}_h$. Following Georgoulis [\[22\]](#page-30-7) and Sudirham et al. [\[42\]](#page-31-2), this mapping $\Phi_{\mathcal{K}}(\widehat{\mathcal{K}}) = \mathcal{K}$ is decomposed into two parts. First, $G_{\mathcal{K}}(\widehat{\mathcal{K}}) = \widetilde{\mathcal{K}}$ denotes the affine mapping defined by $G_K(\hat{x}) = A_K \hat{x} + b$, where $A_K = \text{diag}(\delta t_K/2, h_K/2, \ldots, h_K/2)$ and $b \in \mathbb{R}^{d+1}$ as constant translation vector such that the brick $\tilde{\mathcal{K}} := (0, h_K)^d \times (0, \delta t_{\mathcal{K}})$, see fig. [2.](#page-4-0) In the following, h_K is used to denote the spatial mesh size and δt_K the time-step. We then define $\Phi_K := \phi_K \circ G_K$, where ϕ_K is a diffeomorphism such that $\phi_K(\mathcal{K}) = \mathcal{K}$ (see fig. [2\)](#page-4-0). Note that G_K sets the size of the element K while ϕ_K sets its shape. Following Georgoulis [\[22\]](#page-30-7) and Sudirham et al. [\[42\]](#page-31-2), we assume that ϕ_K is close to the identity, i.e., we will assume that ϕ_K satisfies:

$$
c^{-1} \le |\det J_{\phi_K}| \le c, \quad \left\| (J_{\phi_K})_{ij} \right\|_{L^\infty(\widetilde{\mathcal{K}})} \le c \quad 0 \le i, j \le d, \quad \forall \mathcal{K} \in \mathcal{T}_h,
$$
\n(3.1)

where c is a generic constant independent of h_K , δt_K , ε , and T, where $J_{\phi_K} \in \mathbb{R}^{(n+1)\times(n+1)}$ is the Jacobian of the diffeomorphism ϕ_K , and where the index 0 denotes the time coordinate direction. Since t only depends on t ,

$$
\|\partial_{\widetilde{x}_k}t\|_{L^{\infty}(\widetilde{\mathcal{K}})}=\left\|\left(J_{\phi_{\mathcal{K}}}\right)_{0k}\right\|_{L^{\infty}(\widetilde{\mathcal{K}})}=0,\quad 1\leq k\leq d,\quad \forall \mathcal{K}\in\mathcal{T}_h.
$$

For the inverse of J_{ϕ_K} , let det $J_{\phi_K\setminus mn}$ denote the (m, n) minor of J_{ϕ_K} . We will assume that:

$$
c^{-1} \le |\det J_{\phi_{\mathcal{K}}}^{-1}| \le c, \quad \|\det J_{\phi_{\mathcal{K}}\backslash mn}\|_{L^{\infty}(\widetilde{\mathcal{K}})} \le c, \quad \forall \mathcal{K} \in \mathcal{T}_h.
$$
 (3.2)

Let F_Q^j be a Q-face where \tilde{x}_j is fixed in its affine domain. The parametrization of F_Q^j , obtained from the restriction of ϕ_K to the boundary of K where \tilde{x}_j is fixed, is denoted by ϕ_{F_Q} . Then, [see
22. Theorem 21.3 and Definition on page 180] [33,](#page-30-11) Theorem 21.3 and Definition on page 189],

$$
\int_{F_{\mathcal{Q}}^{j}} f(x) ds = \int_{\widetilde{F}_{\mathcal{Q}}^{j}} f\left(\phi_{F_{\mathcal{Q}}}(\widetilde{x})\right) \left(\det\left(\left(J_{\phi_{\mathcal{K}}}^{j}\right)^{\mathsf{T}} J_{\phi_{\mathcal{K}}}^{j}\right)\right)^{1/2} d\widetilde{s},\tag{3.3}
$$

where J^j_{ϕ} $\psi_{\phi_K}^j \in \mathbb{R}^{(n+1)\times n}$ is obtained by removing the jth column vector from J_{ϕ_K} . We will assume that

$$
c^{-1} \le (\det ((J^i_{\phi_K})^\mathsf{T} J^i_{\phi_K}))^{1/2} \le c, \quad 0 \le i \le d. \tag{3.4}
$$

Figure 2: Construction of the space-time element K through an affine mapping $G_{\mathcal{K}} : \widehat{\mathcal{K}} \to \widetilde{\mathcal{K}}$ and a diffeomorphism $\phi_{\mathcal{K}} : \mathcal{K} \to \mathcal{K}$ [\[42\]](#page-31-2). Note that the front and back faces of \mathcal{K} have constant t-coordinate and hence are parallel to each other.

We will assume that $\delta t_{\mathcal{K}} \leq h_K$. A general bound $\delta t_{\mathcal{K}} \leq \mathcal{O}(h_K)$ poses no difficulty to our analysis but is avoided for the ease of presentation. Our spatial mesh on $\Omega(t)$, for $0 \le t \le T$, is shape-regular and we allow 1-irregularly refined space-time elements, i.e., we allow at most one level of refinement difference between neighboring space-time elements. It will also be useful to introduce a second time-step, $\Delta t_{\mathcal{K}}$, which is the time-step size of the space-time slab to which K belongs. When performing uniform time-stepping, the ratio $\Delta t_K/\delta t_K \equiv 1$ for all $K \in \mathcal{T}_h$. When using local time-stepping, we will assume that the ratio between the time-step of a space-time slab and the minimum local time-step size in that slab is bounded: $\Delta t_K/\delta t_K \leq c$ for all $K \in \mathcal{T}_h$.

3.2 Spaces and useful inequalities

Let $\partial_x^{\alpha}v$, with α a multi-index, be the weak derivative of v and let $H^s(U) := \{v \in L^2(U) :$ $\partial_x^{\alpha} v \in L^2(U)$ for $|\alpha| \leq s$, where s is a nonnegative integer and $U \subset \mathbb{R}^r$ is an open domain with $x := (x_1, \ldots, x_r)$ denoting the coordinates of \mathbb{R}^r . The norm of $H^s(U)$ is defined by $||v||^2_{H^s(U)} :=$ $\sum_{|\alpha|\leq s}\|\partial_x^{\alpha}v\|_L^2$ \mathcal{U}^2 , where $\left\|\cdot\right\|_U$ is the usual L^2 -norm on U.

We also require anisotropic Sobolev spaces. Following Sudirham et al. [\[42\]](#page-31-2) we only consider anisotropy between spatial and temporal variables with no anisotropy between the spatial variables. As such, let s_s and s_t denote the spatial and temporal Sobolev indices, respectively. For $\alpha_t, \alpha_{s_i} \geq 0$, $1 \leq i \leq d$, the anisotropic Sobolev space of order (s_t, s_s) is defined on an open domain $U \subset \mathbb{R}^{d+1}$ by [see 22]:

$$
H^{(s_t,s_s)}(U) := \{ v \in L^2(U) : \partial_t^{\alpha_t} \partial_x^{\alpha_s} v \in L^2(U) \text{ for } \alpha_t \le s_t, |\alpha_s| \le s_s \},
$$

where $\alpha_s = (\alpha_{s_1}, \ldots, \alpha_{s_d})$ and $x := (x_1, \ldots, x_d)$ denotes the spatial coordinates. The anisotropic Sobolev norm reads $||v||^2_{H^{(s_t,s_s)}(U)} := \sum_{\alpha_t \leq s_t, |\alpha_s| \leq s_s} ||\partial_t^{\alpha_t} \partial_x^{\alpha_s} v||^2_{U}$ $_U^2.$

For the HDG method, we require the following finite element spaces

$$
V_h^{(p_t, p_s)} := \{v_h \in L^2(\mathcal{E}) : v_h | \kappa \circ \phi_K \circ G_K \in Q_{(p_t, p_s)}(\widehat{\mathcal{K}}) \quad \forall K \in \mathcal{T}_h\},
$$

$$
M_h^{(p_t, p_s)} := \{\mu_h \in L^2(\Gamma) : \mu_h |_{F} \circ \phi_K \circ G_K \in Q_{(p_t, p_s)}(\widehat{F}) \quad \forall F \in \mathcal{F}_h, \mu_h = 0 \text{ on } \partial \mathcal{E}_D\},
$$

where $Q^{(p_t,p_s)}(U)$ denotes the set of all tensor product polynomials of degree p_t in the temporal direction and p_s in each spatial direction on a domain U. Furthermore, we define $\boldsymbol{V}_h = V_h^{(p_t, p_s)} \times$

 $M_h^{(p_t,p_s)}$ $h_h^{(p_t, p_s)}$. We will denote the pairs $(v, \mu) \in V_h$ and $(u, \lambda) \in V_h$ as $v = (v, \mu)$ and $u = (u, \lambda)$. Furthermore, we denote by [v] the HDG jump $(v - \mu)$.

Adapting [\[22,](#page-30-7) Corollaries 3.49, 3.54] to the space-time context, specifically taking into account the spatial mesh size h_K and time step δt_K of a space-time element $K \in \mathcal{T}_h$, we have the following anisotropic inverse and trace inequalities, which hold for all $v_h \in V_h$:

$$
\left\|\partial_t v_h\right\|_{\mathcal{K}} \le c \left(\delta t_{\mathcal{K}}^{-1} + h_K^{-1}\right) \left\|v_h\right\|_{\mathcal{K}},\tag{3.5a}
$$

$$
\|\overline{\nabla}v_h\|_{\mathcal{K}} \le ch_K^{-1} \|v_h\|_{\mathcal{K}},\tag{3.5b}
$$

$$
||v_h||_{\mathcal{Q}_\mathcal{K}} \le c_\star h_K^{-1/2} ||v_h||_{\mathcal{K}},\tag{3.5c}
$$

$$
||v_h||_{\partial \mathcal{K}} \le c \left(\delta t_{\mathcal{K}}^{-1/2} + h_K^{-1/2}\right) ||v_h||_{\mathcal{K}},
$$
\n(3.5d)

where c_{\star} is a constant independent of h_K , δt_K , ε , and T. (We distinguish c_{\star} from c to prove lemma [4.2.](#page-7-1)) The following lemma introduces an additional inequality.

Lemma 3.1. Let $K \in \mathcal{T}_h$ be a space-time element. For all $\mu_h \in M_h$,

$$
\|\partial_t \mu_h\|_{F_{\mathcal{Q}}} \le c \left(\delta t_{\mathcal{K}}^{-1} + h_K^{-1}\right) \|\mu_h\|_{F_{\mathcal{Q}}}.
$$
\n(3.6)

Proof. The d-dimensional hypersurface F_Q is embedded in \mathbb{R}^{d+1} and in general it may be curved. Therefore, we cannot use eq. $(3.5a)$ directly to conclude eq. (3.6) . Instead, we first map $F_{\mathcal{Q}}$ to the affine domain. For this, let $\phi_{F_Q}(F_Q) = F_Q$, i.e., the transformation of a face from the affine domain to the physical domain. We then observe that one of the spatial coordinates, which is denoted by \widetilde{x}_j without loss of generality, of $\widetilde{F}_{\mathcal{Q}}$ is fixed. This means we can view $\widetilde{F}_{\mathcal{Q}}$ in the \mathbb{R}^d domain with coordinates $(\tilde{t}, \tilde{x}_1, \ldots, \tilde{x}_{j-1}, \tilde{x}_{j+1}, \ldots, \tilde{x}_d)$ and apply the d-dimensional versions of eq. [\(3.5a\)](#page-5-0) and eq. $(3.5b)$ to $F_{\mathcal{O}}$:

$$
\|\partial_{\tilde{t}}\tilde{\mu}_h\|_{\tilde{F}_{\mathcal{Q}}} \le c \left(\delta t_{\mathcal{K}}^{-1} + h_K^{-1}\right) \|\tilde{\mu}_h\|_{\tilde{F}_{\mathcal{Q}}}, \quad \|\overline{\nabla}\tilde{\mu}_h\|_{\tilde{F}_{\mathcal{Q}}} \le ch_K^{-1} \|\tilde{\mu}_h\|_{\tilde{F}_{\mathcal{Q}}}.
$$
\n(3.7)

Using the mapping ϕ_{F_Q} , eqs. [\(3.3\)](#page-3-1) and [\(3.4\)](#page-3-2),

$$
\|\partial_t \mu_h\|_{F_{\mathcal{Q}}}^2 = \int_{\widetilde{F}_{\mathcal{Q}}} \left[\left(\partial_t \left(\widetilde{\mu}_h \circ \phi_{F_{\mathcal{Q}}}^{-1} \right) \right) \circ \phi_{F_{\mathcal{Q}}} \right]^2 \left[\det \left(\left(J_{\phi_K}^j \right)^\mathsf{T} J_{\phi_K}^j \right) \right]^{1/2} d\widetilde{s}
$$

$$
\leq c \int_{\widetilde{F}_{\mathcal{Q}}} \left[\left(\partial_t \left(\widetilde{\mu}_h \circ \phi_{F_{\mathcal{Q}}}^{-1} \right) \right) \circ \phi_{F_{\mathcal{Q}}} \right]^2 d\widetilde{s}.
$$

By the chain rule,

$$
\partial_t \left(\widetilde{\mu}_h \circ \phi_{F_{\mathcal{Q}}}^{-1} \right) = \left(\left(\partial_{\widetilde{t}} \widetilde{\mu}_h \right) \circ \phi_{F_{\mathcal{Q}}}^{-1} \right) \frac{\partial \widetilde{t}}{\partial t} + \sum_{1 \leq i \leq d, i \neq j} \left(\left(\partial_{\widetilde{x}_i} \widetilde{\mu}_h \right) \circ \phi_{F_{\mathcal{Q}}}^{-1} \right) \frac{\partial \widetilde{x}_i}{\partial t}.
$$

We note that $\frac{\partial \tilde{x}_i}{\partial t}$ is the $(i,0)$ -element of $J_{\phi_{\mathcal{K}}}^{-1}$ $\int_{\phi_K}^{-1}$ which equals $(-1)^i \det J_{\phi_{K\setminus i0}} / \det J_{\phi_K}$. Similarly, $\frac{\partial \tilde{t}}{\partial t}$ corresponds to the $(0,0)$ -element of $J_{\phi\kappa}^{-1}$ $\psi_{\phi_K}^{-1}$ which equals det $J_{\phi_{\mathcal{K}\setminus 00}}/\det J_{\phi_{\mathcal{K}}}$. Now using eq. [\(3.1\)](#page-3-3), eq. (3.2) , eq. (3.7) , definition eq. (3.3) , and eq. (3.4) we find that:

$$
\|\partial_t \mu_h\|_{F_{\mathcal{Q}}}^2 \le c \left(\int_{\widetilde{F}_{\mathcal{Q}}} (\partial_{\widetilde{t}} \widetilde{\mu}_h)^2 d\widetilde{s} + \int_{\widetilde{F}_{\mathcal{Q}}} (\widetilde{\nabla} \widetilde{\mu}_h)^2 d\widetilde{s} \right) \n\le c \left(\delta t_{\mathcal{K}}^{-2} + h_K^{-2} \right) \int_{\widetilde{F}_{\mathcal{Q}}} \widetilde{\mu}_h^2 \left[\det \left(\left(J_{\phi_{\mathcal{K}}}^j \right)^{\mathsf{T}} J_{\phi_{\mathcal{K}}}^j \right) \right]^{1/2} d\widetilde{s} \le c \left(\delta t_{\mathcal{K}}^{-2} + h_K^{-2} \right) \|\mu_h\|_{F_{\mathcal{Q}}}^2,
$$

which is eq. (3.6) .

 \Box

6

To end this section we introduce $V := V \times M$, where $V := \{v \in H^1(\mathcal{E}) \mid v|_{\partial \mathcal{E}_D} = 0\} \cap H^2(\mathcal{E})$ and M its trace space, and define the extended function space $V(h) := V(h) \times M(h)$ where $V(h) := V_h^{(p_t, p_s)} + V$ and $M(h) := M_h^{(p_t, p_s)} + M$. We will require the following three norms on $V(h)$:

$$
\|\boldsymbol{v}\|_{v}^{2} := \sum_{\mathcal{K}\in\mathcal{T}_{h}} \|v\|_{\mathcal{K}}^{2} + \sum_{\mathcal{K}\in\mathcal{T}_{h}} \|\beta_{s} - \frac{1}{2}\beta \cdot n\|^{1/2} \left[v\right]\|_{\partial\mathcal{K}}^{2} + \sum_{F\in\partial\mathcal{E}_{N}} \|\|\frac{1}{2}\beta \cdot n\|^{1/2} \mu\|_{F}^{2} \qquad (3.8a)
$$

$$
+ \sum_{\mathcal{K}\in\mathcal{T}_{h}} \varepsilon \|\overline{\nabla}v\|_{\mathcal{K}}^{2} + \sum_{\mathcal{K}\in\mathcal{T}_{h}} \varepsilon h_{K}^{-1} \|\left[v\right]\|_{\mathcal{Q}_{\mathcal{K}}}^{2},
$$

$$
\|\boldsymbol{v}\|_{s}^{2} := \|\boldsymbol{v}\|_{v}^{2} + \sum_{\mathcal{K}\in\mathcal{T}_{h}} \tau_{\varepsilon} \|\partial_{t} v\|_{\mathcal{K}}^{2},\tag{3.8b}
$$

$$
\|\|\mathbf{v}\|_{ss}^2 := \|\|\mathbf{v}\|_{s}^2 + \|v\|_{sd}^2 := \|\|\mathbf{v}\|_{s}^2 + \sum_{\mathcal{K}\in\mathcal{T}_h} \frac{\delta t_{\mathcal{K}} h_{\mathcal{K}}^2}{\delta t_{\mathcal{K}} + h_{\mathcal{K}}} \|\Pi_h(\beta \cdot \nabla v)\|_{\mathcal{K}}^2, \tag{3.8c}
$$

where Π_h denotes the L^2 -projection onto $V_h^{(p_t, p_s)}$ $\mathcal{L}_h^{(p_t, p_s)}$ and the parameter τ_{ε} in the definition of $\Vert \vert \mathbf{v} \Vert \vert_s$ depends on the size of the space-time element compared to the diffusion parameter ε :

$$
\tau_{\varepsilon} := \Delta t_{\mathcal{K}} \tilde{\varepsilon} := \begin{cases} \Delta t_{\mathcal{K}} & \text{if } \mathcal{K} \in \mathcal{T}_h^d := \left\{ \mathcal{K} \in \mathcal{T}_h | \delta t_{\mathcal{K}} \le h_K \le \varepsilon \right\}, \\ \Delta t_{\mathcal{K}} \varepsilon^{1/2} & \text{if } \mathcal{K} \in \mathcal{T}_h^x := \left\{ \mathcal{K} \in \mathcal{T}_h | \delta t_{\mathcal{K}} \le \varepsilon < h_K \right\}, \\ \Delta t_{\mathcal{K}} \varepsilon & \text{if } \mathcal{K} \in \mathcal{T}_h^c := \left\{ \mathcal{K} \in \mathcal{T}_h | \varepsilon < \delta t_{\mathcal{K}} \le h_K \right\}. \end{cases}
$$

Finally, $\beta_s := \sup_{(x,t) \in F} |\beta \cdot n|$, for $F \subset \partial K$. It is useful to remark that

$$
\inf_{(x,t)\in F} (\beta_s - \frac{1}{2}\beta \cdot n) \ge \frac{1}{2} \max_{(x,t)\in F} |\beta \cdot n| \qquad \forall F \in \partial \mathcal{K}, \ \forall \mathcal{K} \in \mathcal{T}_h.
$$
\n(3.9)

3.3 The discretization

Let $u, v \in [L^2(U)]^r$ for $1 \le r \le d+1$. We will write $(u, v)_U = \int_U u \cdot v \, dx$ if $U \subset \mathbb{R}^{d+1}$ and $\langle u, v \rangle_U =$ Let $u, v \in [D]$ (v)] for $1 \leq t \leq u+1$. We will write $(u, v)_U = J_U u \cdot v$ dx if $U \subset \mathbb{R}^d$. Furthermore, we define $(u, v)_{\mathcal{T}_h} := \sum_{\mathcal{K} \in \mathcal{T}_h} \langle u, v \rangle_{\partial \mathcal{K}}$, $\langle u, v \rangle_{\partial \mathcal{T}_h} := \sum_{\mathcal{K} \in \mathcal{T}_h} \langle u, v \rangle_{\partial \mathcal{K}}$, $\langle u, v \rangle_{\mathcal{Q}_h} := \sum_{\mathcal{K} \in \mathcal{T}_h} \langle u, v \rangle_{\mathcal{Q}_\mathcal{K}}, \text{ and } \langle u, v \rangle_{\partial \mathcal{E}_N} := \sum_{F \in \mathcal{F}_h^b \cap \partial \mathcal{E}_N} \langle u, v \rangle_F.$

The space-time HDG method for eq. [\(2.1\)](#page-2-1) is given by: Find $u_h \in V_h$ such that

$$
a_h(\boldsymbol{u}_h, \boldsymbol{v}_h) = (f, v_h)_{\mathcal{T}_h} + \langle g, \mu_h \rangle_{\partial \mathcal{E}_N} \quad \forall \boldsymbol{v}_h \in \boldsymbol{V}_h, \tag{3.10}
$$

with $a_h(\boldsymbol{u}_h, \boldsymbol{v}_h) := a_{h,d}(\boldsymbol{u}_h, \boldsymbol{v}_h) + a_{h,c}(\boldsymbol{u}_h, \boldsymbol{v}_h)$ and where

$$
a_{h,d}(\boldsymbol{u},\boldsymbol{v}) := (\varepsilon \overline{\nabla} u, \overline{\nabla} v)_{\mathcal{T}_h} + \langle \varepsilon \alpha h_K^{-1}[\boldsymbol{u}], [\boldsymbol{v}] \rangle_{\mathcal{Q}_h} - \langle \varepsilon[\boldsymbol{u}], \overline{\nabla}_{\bar{n}} v \rangle_{\mathcal{Q}_h} - \langle \varepsilon \overline{\nabla}_{\bar{n}} u, [\boldsymbol{v}] \rangle_{\mathcal{Q}_h},
$$

$$
a_{h,c}(\boldsymbol{u},\boldsymbol{v}) := -(\beta u, \nabla v)_{\mathcal{T}_h} + \langle \zeta^{\dagger} \beta \cdot n \lambda, \mu \rangle_{\partial \mathcal{E}_N} + \langle (\beta \cdot n) \lambda + \beta_s[\boldsymbol{u}], [\boldsymbol{v}] \rangle_{\partial \mathcal{T}_h}.
$$

Here $\alpha > 0$ is a penalty parameter and ζ^+ denotes the outflow boundary indicator on a facet. The following boundedness result, which follows from the Cauchy–Schwarz inequality and eq. [\(3.5c\)](#page-5-4), will be useful in the proof of stability in section [4:](#page-7-0)

$$
|a_{h,d}(\boldsymbol{u}_h,\boldsymbol{v}_h)| \leq c \|\|\boldsymbol{u}_h\|_{\boldsymbol{v}} \|\|\boldsymbol{v}_h\|_{\boldsymbol{v}} \qquad \forall \boldsymbol{u}_h, \boldsymbol{v}_h \in \boldsymbol{V}_h. \tag{3.11}
$$

4 Stability

The main goal of this section is to prove theorem [4.1](#page-7-2) which states stability of the space-time HDG method for the advection-diffusion equation with respect to a norm that measures the streamline derivative, i.e., $\|\cdot\|_{ss}$. We will prove that this result is robust with respect to ε . A similar result for the stationary problem is shown in Ayuso and Marini [\[2,](#page-28-0) Theorem 4.6]. For this and following sections, c_T denotes a constant independent of h_K , δt_K , and ε , but linear in T.

Theorem 4.1. There exists δt_0 , independent of ε and T, such that when $\delta t_K \leq \min(h_K, \delta t_0)$ on all $\mathcal{K} \in \mathcal{T}_h$, and for all $\mathbf{w}_h \in \mathbf{V}_h$,

$$
c_T^{-1} \|\|\boldsymbol{w}_h\|\|_{ss} \leq \sup_{\boldsymbol{v}_h \in \boldsymbol{V}_h} \frac{a_h(\boldsymbol{w}_h, \boldsymbol{v}_h)}{\|\|\boldsymbol{v}_h\|\|_s}.
$$
\n(4.1)

The following two inf-sup conditions with respect to, respectively, $\|\cdot\|_v$ and $\|\cdot\|_s$, and which hold under the same conditions as theorem [4.1,](#page-7-2) are used to prove theorem [4.1:](#page-7-2)

$$
c_T^{-1} \|\|\boldsymbol{w}_h\|\|_v \leq \sup_{\boldsymbol{v}_h \in \boldsymbol{V}_h} \frac{a_h(\boldsymbol{w}_h, \boldsymbol{v}_h)}{\|\|\boldsymbol{v}_h\||_v} \qquad \forall \boldsymbol{w}_h \in \boldsymbol{V}_h,
$$
\n(4.2a)

$$
c_T^{-1} \|\|\boldsymbol{w}_h\|\|_{s} \leq \sup_{\boldsymbol{v}_h \in \boldsymbol{V}_h} \frac{a_h(\boldsymbol{w}_h, \boldsymbol{v}_h)}{\|\|\boldsymbol{v}_h\|\|_{s}} \qquad \forall \boldsymbol{w}_h \in \boldsymbol{V}_h. \tag{4.2b}
$$

We prove eqs. $(4.2a)$ and $(4.2b)$ in sections [4.1](#page-7-2) and [4.2,](#page-12-0) respectively. We then prove theorem 4.1 in section [4.3.](#page-16-0) To prove these results we introduce, for $T \geq 1$, the weighting function

$$
\varphi = e \exp(-t/T) + \chi,\tag{4.3}
$$

where the positive constant χ will be determined later. For $0 < T < 1$ we propose $\varphi(t) =$ $e \exp(-t) + \chi$. In our analysis, however, we will only consider $T \geq 1$; the analysis for $T < 1$ follows identical steps as the $T \geq 1$ case, resulting in inf-sup conditions theorem [4.1](#page-7-2) and eqs. [\(4.2a\)](#page-7-3) and [\(4.2b\)](#page-7-4) independent of T. Denoting the cell mean of $\bar{\beta}$ by $\bar{\beta}_0$, we will also use that [see [12\]](#page-29-12),

$$
\|\bar{\beta} - \bar{\beta}_0\|_{L^\infty(\mathcal{K})} \le ch_K|\bar{\beta}|_{W^{1,\infty}(\mathcal{K})} \quad \forall \mathcal{K} \in \mathcal{T}_h. \tag{4.4}
$$

4.1 The inf-sup condition with respect to $\|\cdot\|_v$

To prove eq. [\(4.2a\)](#page-7-3) we first require the following lemmas.

Lemma 4.2. Let φ be defined as in eq. [\(4.3\)](#page-7-6) with χ chosen such that $\chi > (e -$ √ $2) T /$ (√ $(2-1).$ Furthermore, choose the penalty parameter α in eq. [\(3.10\)](#page-6-0) such that $\alpha > 1+4c_\star^2$, with c_\star the constant in eq. [\(3.5c\)](#page-5-4). Then for all $w_h := (w_h, \varkappa_h) \in \mathbf{V}_h$:

$$
a_h(\boldsymbol{w}_h, \varphi \boldsymbol{w}_h) \geq \frac{1}{2} (T + \chi) \left(\sum_{\mathcal{K} \in \mathcal{T}_h} \varepsilon \left\| \overline{\nabla} w_h \right\|_{\mathcal{K}}^2 + \sum_{\mathcal{K} \in \mathcal{T}_h} \varepsilon h_K^{-1} \left\| \left[\boldsymbol{w}_h \right] \right\|_{\mathcal{Q}_\mathcal{K}}^2 \right) + \frac{1}{2} \sum_{\mathcal{K} \in \mathcal{T}_h} \left\| w_h \right\|_{\mathcal{K}}^2 + (T + \chi) \left(\left\| \left| \frac{1}{2} \beta \cdot n \right|^{1/2} \varkappa_h \right\|_{\partial \mathcal{E}_N}^2 + \sum_{\mathcal{K} \in \mathcal{T}_h} \left\| \left| \beta_s - \frac{1}{2} \beta \cdot n \right|^{1/2} \left[\boldsymbol{w}_h \right] \right\|_{\partial \mathcal{K}}^2 \right).
$$

Proof. On an element $K \in \mathcal{T}_h$ we have $-w_h \beta \cdot \nabla (\varphi w_h) = -\frac{1}{2} \nabla \cdot (\varphi \beta w_h^2) - \frac{1}{2} w_h^2 \beta \cdot \nabla \varphi$. Using Gauss's theorem, $[\varphi \mathbf{w}_h] = \varphi[\mathbf{w}_h]$, and that $\zeta^+ \beta \cdot n = (\beta \cdot n + |\beta \cdot n|)/2$, we note that

$$
a_{h,c}(\mathbf{w}_h, \varphi \mathbf{w}_h) = -(\frac{1}{2}w_h^2, \beta \cdot \nabla \varphi)_{\mathcal{T}_h} + \langle \frac{1}{2} \varphi \varkappa_h^2, \beta \cdot n + |\beta \cdot n| \rangle_{\partial \mathcal{E}_N} - \langle \frac{1}{2} \varphi w_h^2, \beta \cdot n \rangle_{\partial \mathcal{T}_h} + \langle \varphi [\mathbf{w}_h]^2, \sup |\beta \cdot n| \rangle_{\partial \mathcal{T}_h} + \langle \varphi \varkappa_h [\mathbf{w}_h], \beta \cdot n \rangle_{\partial \mathcal{T}_h}.
$$

Since $-\frac{1}{2}w_h^2 + \varkappa_h w_h] = -\frac{1}{2}$ $\frac{1}{2}\left[\boldsymbol{w}_h\right]^2-\frac{1}{2}$ $\frac{1}{2}\varkappa_h^2$, \varkappa_h is single-valued on element boundaries, $\varkappa_h = 0$ on $\partial \mathcal{E}_D$, we have by definition of φ and using that $-\beta \cdot \nabla \varphi \geq 1$:

$$
a_{h,c}(\boldsymbol{w}_h, \varphi \boldsymbol{w}_h) \ge (T + \chi) \left\| \left| \frac{1}{2} \beta \cdot n \right|^{1/2} \varkappa_h \right\|_{\partial \mathcal{E}_N}^2 + \frac{1}{2} \sum_{\mathcal{K} \in \mathcal{T}_h} \left\| w_h \right\|_{\mathcal{K}}^2 + (T + \chi) \sum_{\mathcal{K} \in \mathcal{T}_h} \left\| \left| \beta_s - \frac{1}{2} \beta \cdot n \right|^{1/2} \left[\boldsymbol{w}_h \right] \right\|_{\partial \mathcal{K}}^2. \tag{4.5}
$$

Next, noting that $\overline{\nabla}\varphi = 0$, and using the Cauchy–Schwarz inequality and eq. [\(3.5c\)](#page-5-4),

$$
a_{h,d}(\mathbf{w}_h, \varphi \mathbf{w}_h) \ge (T + \chi) \sum_{\mathcal{K} \in \mathcal{T}_h} \varepsilon \|\overline{\nabla} w_h\|_{\mathcal{K}}^2 + (T + \chi) \alpha \sum_{\mathcal{K} \in \mathcal{T}_h} \varepsilon h_K^{-1} \|[\mathbf{w}_h]\|_{\mathcal{Q}_\mathcal{K}}^2
$$

$$
- \sum_{\mathcal{K} \in \mathcal{T}_h} 2\varepsilon^{1/2} c_\star (eT + \chi) \|\overline{\nabla} w_h\|_{\mathcal{K}} \varepsilon^{1/2} h_K^{-1/2} \|[\mathbf{w}_h]\|_{\mathcal{Q}_\mathcal{K}}.
$$

Using Hölder's inequality for sums and the inequality $ax^2 - 2bxy + dy^2 \geq (ad - b^2)(x^2 + y^2)/(a + d)$, which holds for positive real numbers a, b, d and $ad > b^2$ [see [35\]](#page-31-7) allows us to obtain

$$
a_{h,d}(\boldsymbol{w}_h, \varphi \boldsymbol{w}_h) \geq (T + \chi) \frac{\alpha - \left(\frac{c_{\star}(eT + \chi)}{T + \chi}\right)^2}{1 + \alpha} \left(\sum_{\mathcal{K} \in \mathcal{T}_h} \varepsilon \left\|\overline{\nabla}w_h\right\|_{\mathcal{K}}^2 + \sum_{\mathcal{K} \in \mathcal{T}_h} \varepsilon h_K^{-1} \left\|\left[\boldsymbol{w}_h\right]\right\|_{\mathcal{Q}_{\mathcal{K}}}^2\right).
$$

Since χ and α are chosen such that $\chi > (e -$ √ $2)T/$ (√ $(2-1)$, so that $T + \chi > (eT + \chi)$ √ 2, and $\alpha > 1 + 4c_\star^2$, it follows that

$$
a_{h,d}(\boldsymbol{w}_h, \varphi \boldsymbol{w}_h) \geq \frac{1}{2}(T + \chi) \left(\sum_{\mathcal{K} \in \mathcal{T}_h} \varepsilon \left\| \overline{\nabla} w_h \right\|_{\mathcal{K}}^2 + \sum_{\mathcal{K} \in \mathcal{T}_h} \varepsilon h_K^{-1} \left\| \left[\boldsymbol{w}_h \right] \right\|_{\mathcal{Q}_{\mathcal{K}}}^2 \right). \tag{4.6}
$$

The result follows after combining eqs. [\(4.5\)](#page-8-0) and [\(4.6\)](#page-8-1).

Let $\Pi_h^{\mathcal{F}}$ be the L^2 -projection onto $M_h^{(p_t, p_s)}$ $h_h^{(p_t, p_s)}$. In appendix [A](#page-22-0) we show that for $u \in H^1(\mathcal{K}),$

$$
\left\|\overline{\nabla}\left(u-\Pi_h u\right)\right\|_{\mathcal{K}} \le c \left\|\overline{\nabla}u\right\|_{\mathcal{K}},\tag{4.7a}
$$

 \Box

$$
\|\partial_t (u - \Pi_h u)\|_{\mathcal{K}} \le c \left(\|\partial_t u\|_{\mathcal{K}} + \|\overline{\nabla} u\|_{\mathcal{K}} \right),\tag{4.7b}
$$

$$
\|\Pi_h u - \Pi_h^{\mathcal{F}} u\|_{\mathcal{Q}_{\mathcal{K}}} \le ch_K^{1/2} \|\overline{\nabla} u\|_{\mathcal{K}}.
$$
\n(4.7c)

The following lemma extends the L^2 -projection estimates of [\[2,](#page-28-0) Lemma 4.2] to space-time elements, taking into account the spatial mesh size h_K and time step $\delta t_{\mathcal{K}}.$

Lemma 4.3. Let φ be the function defined in eq. [\(4.3\)](#page-7-6). For any $w_h \in V_h^{(p_t, p_s)}$ $h^{(pt, p_s)}$ the following estimates hold:

$$
\left\| \left(I - \Pi_h\right) \left(\varphi w_h\right) \right\|_{\mathcal{K}} \leq c \delta t_{\mathcal{K}} \left\| w_h \right\|_{\mathcal{K}},\tag{4.8a}
$$

$$
\|\overline{\nabla}\left(\left(I - \Pi_h\right)(\varphi w_h)\right)\|_{\mathcal{K}} \le c\delta t_{\mathcal{K}} h_K^{-1} \|w_h\|_{\mathcal{K}},\tag{4.8b}
$$

$$
\|\overline{\nabla}\left(\left(I-\Pi_h\right)(\varphi w_h)\right)\|_{\mathcal{Q}_{\mathcal{K}}} \leq c\delta t_{\mathcal{K}} h_K^{-3/2} \|w_h\|_{\mathcal{K}},\tag{4.8c}
$$

$$
\left\| \left(I - \Pi_h\right) \left(\varphi w_h\right) \right\|_{\mathcal{Q}_{\mathcal{K}}} \leq c \delta t_{\mathcal{K}} h_K^{-1/2} \left\| w_h \right\|_{\mathcal{K}},\tag{4.8d}
$$

$$
||(I - \Pi_h)(\varphi w_h)||_{\mathcal{R}_{\mathcal{K}}} \le c\delta t_{\mathcal{K}}^{1/2} ||w_h||_{\mathcal{K}}.
$$
\n(4.8e)

Proof. See appendix **B**.

Lemma 4.4. Let $\Pi_h(\varphi w_h) := (\Pi_h(\varphi w_h), \Pi_h^{\mathcal{F}}(\varphi \varkappa_h))$ for all $w_h := (w_h, \varkappa_h) \in V_h$. The following holds:

$$
\left\|\left[\mathbf{\Pi}_{h}\left(\varphi\boldsymbol{w}_{h}\right)\right]\right\|_{v}\leq c_{T}\|\left|\hspace{-1.5pt}\left|\boldsymbol{w}_{h}\right|\right|_{v}.
$$

Proof. See appendix [C.](#page-26-0)

Lemma 4.5. For any $w := (w, \varkappa) \in L^2(\mathcal{E}) \times L^2(\Sigma)$, let $\boldsymbol{\delta w} := (w - \Pi_h w, \varkappa - \Pi_h^{\mathcal{F}} \varkappa)$. The following holds for all $\mathbf{w}_h \in \mathbf{V}_h$:

$$
a_h(\boldsymbol{w}_h, \boldsymbol{\delta}(\varphi \boldsymbol{w}_h)) \leq c_T \left(\sum_{\mathcal{K} \in \mathcal{T}_h} \varepsilon \left\| \overline{\nabla} w_h \right\|_{\mathcal{K}}^2 + \sum_{\mathcal{K} \in \mathcal{T}_h} \varepsilon h_K^{-1} \left\| \left[\boldsymbol{w}_h \right] \right\|_{\mathcal{Q}_\mathcal{K}}^2 \right) + \sum_{\mathcal{K} \in \mathcal{T}_h} \left(1/8 + \delta t_\mathcal{K} \right) \left\| w_h \right\|_{\mathcal{K}}^2
$$

+
$$
c_T \left(\sum_{F \in \partial \mathcal{E}_N} \left\| \left| \frac{1}{2} \beta \cdot n \right|^{1/2} \varkappa_h \right\|_F^2 + \sum_{\mathcal{K} \in \mathcal{T}_h} \left\| \left| \beta_s - \frac{1}{2} \beta \cdot n \right|^{1/2} \left[\boldsymbol{w}_h \right] \right\|_{\partial \mathcal{K}}^2 \right).
$$

Proof. Let $z \in H^1(\mathcal{T}_h)$ and $\varpi \in L^2(\mathcal{F}_h)$ such that $\varpi|_{\partial \mathcal{E}_D} = 0$. Let $z := (z, \varpi)$. Integrating $(\beta w_h, \nabla z)_{\mathcal{T}_h}$ by parts and using that $\langle (\beta \cdot n) \varkappa_h, \varpi \rangle_{\partial \mathcal{T}_h} = \langle (\beta \cdot n) \varkappa_h, \varpi \rangle_{\partial \mathcal{E}_N}$, because \varkappa_h and ϖ are single-valued on Γ and zero on $\partial \mathcal{E}_D$, we have:

$$
a_{h,c}(\boldsymbol{w}_h, \boldsymbol{z}) = (\beta \cdot \nabla w_h, z)_{\mathcal{T}_h} - \langle \frac{1}{2} (\beta \cdot n) [\boldsymbol{w}_h], z \rangle_{\partial \mathcal{T}_h} + \langle \left(\beta_s - \frac{1}{2} (\beta \cdot n) \right) [\boldsymbol{w}_h], z \rangle_{\partial \mathcal{T}_h} - \langle \beta_s [\boldsymbol{w}_h], \varpi \rangle_{\partial \mathcal{T}_h} + \langle \frac{1}{2} (|\beta \cdot n| - \beta \cdot n) \times_h, \varpi \rangle_{\partial \mathcal{E}_N}.
$$
\n(4.9)

At this point, note that $\delta(\varphi w_h) = \delta(eT \exp(-t/T) w_h)$ because $\delta(\chi w_h) = 0$. Furthermore, let $\beta_0 = (1, \bar{\beta}_0)$. By definition of Π_h , the following vanishes: $(\beta_0 \cdot \nabla w_h, (I - \Pi_h) (eT \exp(-t/T)) w_h)_{\mathcal{T}_h} =$ 0. From eq. [\(4.9\)](#page-9-0), with $\mathbf{z} = \delta(eT \exp(-t/T)\mathbf{w}_h)$, we now find that:

$$
a_{h,c}(\mathbf{w}_h, \delta (eT \exp(-t/T)) \mathbf{w}_h) = ((\beta - \beta_0) \cdot \nabla w_h, (I - \Pi_h) (eT \exp(-t/T) w_h))_{\mathcal{T}_h}
$$

\n
$$
- \langle \frac{1}{2} (\beta \cdot n) [\mathbf{w}_h], (I - \Pi_h) (eT \exp(-t/T) w_h) \rangle_{\partial \mathcal{T}_h}
$$

\n
$$
+ \langle (\beta_s - \frac{1}{2} (\beta \cdot n)) [\mathbf{w}_h], (I - \Pi_h) (eT \exp(-t/T) w_h) \rangle_{\partial \mathcal{T}_h}
$$

\n
$$
+ \langle \frac{1}{2} (|\beta \cdot n| - \beta \cdot n) \varkappa_h, (I - \Pi_h^{\mathcal{F}}) (eT \exp(-t/T) \varkappa_h) \rangle_{\partial \mathcal{E}_N}
$$

\n
$$
=: M_1 + M_2 + M_3 + M_4,
$$

where, by definitions of $\Pi_h^{\mathcal{F}}$ and β_s , $\langle \beta_s [\boldsymbol{w}_h] , (I - \Pi_h^{\mathcal{F}}) (eT \exp(-t/T) \varkappa_h) \rangle_{\partial \mathcal{T}_h} = 0$. We will now bound each of the terms M_i , $i = 1, ..., 4$ separately.

We observe that $(\beta - \beta_0) \cdot \nabla w_h = (\bar{\beta} - \bar{\beta}_0) \cdot \nabla w_h$ because the first components of β and β_0 are 1. We then bound M_1 using the Cauchy–Schwarz inequality, eq. (4.4) , eq. $(4.8a)$, and eq. $(3.5b)$:

$$
M_1 \leq \sum_{\mathcal{K}\in\mathcal{T}_h} ch_K h_K^{-1} \|w_h\|_{\mathcal{K}} \delta t_{\mathcal{K}} \|w_h\|_{\mathcal{K}} = c \sum_{\mathcal{K}\in\mathcal{T}_h} \delta t_{\mathcal{K}} \|w_h\|_{\mathcal{K}}^2.
$$
 (4.10)

We proceed with bounding M_2 and M_3 . Using the Cauchy–Schwarz inequality, eq. [\(3.9\)](#page-6-1), eq. [\(4.8d\)](#page-8-3), and eq. [\(4.8e\)](#page-8-4), we find that

$$
M_2 + M_3 \leq c \sum_{\mathcal{K} \in \mathcal{T}_h} \delta t_{\mathcal{K}} h_K^{-1/2} \left\| \left| \beta_s - \frac{1}{2} \beta \cdot n \right|^{1/2} \left[\boldsymbol{w}_h \right] \right\|_{\mathcal{Q}_\mathcal{K}} \left\| w_h \right\|_{\mathcal{K}} + c \sum_{\mathcal{K} \in \mathcal{T}_h} \delta t_{\mathcal{K}}^{1/2} \left\| \left| \beta_s - \frac{1}{2} \beta \cdot n \right|^{1/2} \left[\boldsymbol{w}_h \right] \right\|_{\mathcal{R}_\mathcal{K}} \left\| w_h \right\|_{\mathcal{K}}.
$$

 \Box

 \Box

Since $\delta t_{\mathcal{K}} \leq h_K$, $\delta t_{\mathcal{K}} h_K^{-1/2}$ can be bounded by 1. Therefore, applying Young's inequality,

$$
M_2 + M_3 \leq c \sum_{\mathcal{K} \in \mathcal{T}_h} |||\beta_s - \frac{1}{2} (\beta \cdot n)|^{1/2} \left[\mathbf{w}_h\right]||_{\partial \mathcal{K}} ||\mathbf{w}_h||_{\mathcal{K}}
$$

$$
\leq \frac{1}{2} \delta \sum_{\mathcal{K} \in \mathcal{T}_h} ||\mathbf{w}_h||_{\mathcal{K}}^2 + c\delta^{-1} \sum_{\mathcal{K} \in \mathcal{T}_h} |||\beta_s - \frac{1}{2} (\beta \cdot n)|^{1/2} \left[\mathbf{w}_h\right]||_{\partial \mathcal{K}}^2.
$$
 (4.11)

For M_4 , we first apply the Cauchy–Schwarz inequality and the triangle inequality:

$$
M_4 \leq c \sum_{F \in \partial \mathcal{E}_N} |||_{2}^{\frac{1}{2}} \beta \cdot n||_{F}^{1/2} \varkappa_h||_{F} (|||_{2}^{\frac{1}{2}} \beta \cdot n|_{F}^{1/2} \varkappa_h||_{F} + |||_{2}^{\frac{1}{2}} \beta \cdot n|_{F}^{1/2} \Pi_h^{\mathcal{F}} (eT \exp(-t/T) \varkappa_h)||_{F}) \qquad (4.12)
$$

The second term in parentheses on the right-hand side of eq. [\(4.12\)](#page-10-0) is bounded following identical steps in showing eq. [\(C.4\)](#page-27-0). Applying also Young's inequality, and denoting by \mathcal{K}_F the space-time element of which F is a facet,

$$
M_{4} \leq c \sum_{F \in \partial \mathcal{E}_{N}} |||_{\frac{1}{2}}^{\frac{1}{2}} \beta \cdot n|^{1/2} \times_{h} ||_{F}^{2}
$$

+ $c \sum_{F \in \partial \mathcal{E}_{N}} |||_{\frac{1}{2}}^{\frac{1}{2}} \beta \cdot n|^{1/2} \times_{h} ||_{F} (c_{T} |||_{\frac{1}{2}}^{\frac{1}{2}} \beta \cdot n|^{1/2} \times_{h} ||_{F} + c_{T} ||(\beta_{s} - \frac{1}{2}\beta \cdot n)^{1/2} [\boldsymbol{w}_{h}]||_{\partial K} + ||\boldsymbol{w}_{h}||_{\mathcal{K}_{F}})= (c + c_{T}) \sum_{F \in \partial \mathcal{E}_{N}} |||_{\frac{1}{2}}^{\frac{1}{2}} \beta \cdot n|^{1/2} \times_{h} ||_{F}^{2}+ c_{T} \sum_{F \in \partial \mathcal{E}_{N}} |||_{\frac{1}{2}}^{\frac{1}{2}} \beta \cdot n|^{1/2} \times_{h} ||_{F} ||(\beta_{s} - \frac{1}{2}\beta \cdot n)^{1/2} [\boldsymbol{w}_{h}]||_{\partial K} + c \sum_{F \in \partial \mathcal{E}_{N}} |||_{\frac{1}{2}}^{\frac{1}{2}} \beta \cdot n|^{1/2} \times_{h} ||_{F} ||\boldsymbol{w}_{h}||_{\mathcal{K}_{F}}\leq \frac{\delta}{2} \sum_{K \in \mathcal{T}_{h}} ||\boldsymbol{w}_{h}||_{K}^{2} + (c_{T} + c\delta^{-1}) \sum_{F \in \partial \mathcal{E}_{N}} |||_{\frac{1}{2}}^{\frac{1}{2}} \beta \cdot n|^{1/2} \times_{h} ||_{F}^{2} + c_{T} \sum_{F \in \partial \mathcal{E}_{N}} ||(\beta_{s} - \frac{1}{2}\beta \cdot n)^{1/2} [\boldsymbol{w}_{h}]||_{\partial K}^{2}.$ \n(4.13)

We proceed with the diffusive term $a_{h,d}$. With test function $\boldsymbol{z} = \boldsymbol{\delta} (eT \exp(-t/T)\boldsymbol{w}_h)$,

$$
a_{h,d}(\mathbf{w}_h, \delta(\epsilon T \exp(-t/T)) \mathbf{w}_h)
$$

= $(\epsilon \overline{\nabla} w_h, \overline{\nabla} ((I - \Pi_h) (\epsilon T \exp(-t/T)) w_h))_{\mathcal{T}_h} - \langle \epsilon \alpha h_K^{-1} [\mathbf{w}_h], (\Pi_h - \Pi_h^{\mathcal{F}}) (\epsilon T \exp(-t/T) w_h) \rangle_{\mathcal{Q}_h}$
+ $\langle \epsilon \alpha h_K^{-1} [\mathbf{w}_h], (I - \Pi_h^{\mathcal{F}}) (\epsilon T \exp(-t/T) [\mathbf{w}_h]) \rangle_{\mathcal{Q}_h} - \langle \epsilon [\mathbf{w}_h], \overline{\nabla}_{\bar{n}} ((I - \Pi_h) (\epsilon T \exp(-t/T) w_h)) \rangle_{\mathcal{Q}_h}$
+ $\langle \epsilon \overline{\nabla}_{\bar{n}} w_h, (\Pi_h - \Pi_h^{\mathcal{F}}) (\epsilon T \exp(-t/T) w_h) \rangle_{\mathcal{Q}_h} - \langle \epsilon \overline{\nabla}_{\bar{n}} w_h, (I - \Pi_h^{\mathcal{F}}) (\epsilon T \exp(-t/T) [\mathbf{w}_h]) \rangle_{\mathcal{Q}_h}$
=: $M_5 + M_6 + M_7 + M_8 + M_9 + M_{10}$.

To bound M_5 we use the Cauchy–Schwarz inequality, eq. [\(4.8b\)](#page-8-5), the assumption that $\delta t_K \leq h_K$, and Young's inequality:

$$
M_5 \leq c \sum_{\mathcal{K}\in\mathcal{T}_h} \varepsilon \left\| \overline{\nabla} w_h \right\|_{\mathcal{K}} \delta t_{\mathcal{K}} h_K^{-1} \left\| w_h \right\|_{\mathcal{K}} \leq \frac{1}{2} \delta \sum_{\mathcal{K}\in\mathcal{T}_h} \left\| w_h \right\|_{\mathcal{K}}^2 + c\varepsilon \delta^{-1} \sum_{\mathcal{K}\in\mathcal{T}_h} \varepsilon \left\| \overline{\nabla} w_h \right\|_{\mathcal{K}}^2. \tag{4.14}
$$

To bound M_6 we apply the Cauchy–Schwarz inequality, eq. $(4.7c)$, and Young's inequality:

$$
M_6 \leq c \sum_{\mathcal{K} \in \mathcal{T}_h} \varepsilon h_K^{-1} \left\| [\boldsymbol{w}_h] \right\|_{\mathcal{Q}_\mathcal{K}} h_K^{1/2} \left\| \overline{\nabla} \left(e^T \exp(-t/T) w_h \right) \right\|_{\mathcal{K}} \leq c_T \sum_{\mathcal{K} \in \mathcal{T}_h} \varepsilon h_K^{-1} \left\| [\boldsymbol{w}_h] \right\|_{\mathcal{Q}_\mathcal{K}}^2 + c_T \sum_{\mathcal{K} \in \mathcal{T}_h} \varepsilon \left\| \overline{\nabla} w_h \right\|_{\mathcal{K}}^2.
$$
\n(4.15)

 M_7 can be bounded using the boundedness of $\Pi_b^{\mathcal{F}}$:

$$
M_7 \leq c \sum_{\mathcal{K}\in\mathcal{T}_h} \varepsilon h_K^{-1} \left\| \left[\mathbf{w}_h\right]\right\|_{\mathcal{Q}_\mathcal{K}} \left\| \left(I - \Pi_h^{\mathcal{F}}\right) \left(eT\exp(-t/T)\left[\mathbf{w}_h\right]\right) \right\|_{\mathcal{Q}_\mathcal{K}} \leq c_T \sum_{\mathcal{K}\in\mathcal{T}_h} \varepsilon h_K^{-1} \left\| \left[\mathbf{w}_h\right]\right\|_{\mathcal{Q}_\mathcal{K}}^2. \tag{4.16}
$$

Terms M_9 and M_{10} are bounded in a similar way as M_6 and M_7 , and using eq. [\(3.5c\)](#page-5-4):

$$
M_9 + M_{10} \leq c_T \sum_{\mathcal{K} \in \mathcal{T}_h} \varepsilon h_K^{-1} ||[\boldsymbol{w}_h]||_{\mathcal{Q}_\mathcal{K}}^2 + c_T \sum_{\mathcal{K} \in \mathcal{T}_h} \varepsilon ||\overline{\nabla} w_h||_{\mathcal{K}}^2.
$$
 (4.17)

Finally, we bound M_8 using the Cauchy–Schwarz inequality, eq. [\(4.8c\)](#page-8-7), the assumption that $\delta t_{\mathcal{K}} \leq$ h_K , and Young's inequality:

$$
M_8 \leq \sum_{\mathcal{K}\in\mathcal{T}_h} \varepsilon \left\| [\boldsymbol{w}_h] \right\|_{\mathcal{Q}_\mathcal{K}} \left\| \overline{\nabla} \left((I - \Pi_h) \left(eT \exp(-t/T) w_h \right) \right) \right\|_{\mathcal{Q}_\mathcal{K}} \n\leq c \sum_{\mathcal{K}\in\mathcal{T}_h} \varepsilon \left\| [\boldsymbol{w}_h] \right\|_{\mathcal{Q}_\mathcal{K}} \delta t \kappa h_K^{-3/2} \left\| w_h \right\|_{\mathcal{K}} \n\leq \frac{1}{2} \delta \sum_{\mathcal{K}\in\mathcal{T}_h} \left\| w_h \right\|_{\mathcal{K}}^2 + c\varepsilon \delta^{-1} \sum_{\mathcal{K}\in\mathcal{T}_h} \varepsilon h_K^{-1} \left\| [\boldsymbol{w}_h] \right\|_{\mathcal{Q}_\mathcal{K}}^2.
$$
\n(4.18)

Collecting eqs. (4.10) , (4.11) and (4.13) to (4.18) we find that

$$
a_h(\boldsymbol{w}_h, \boldsymbol{\delta}(\varphi \boldsymbol{w}_h)) \leq c \sum_{K \in \mathcal{T}_h} \delta t_K \left\| w_h \right\|_K^2 + 2 \delta \sum_{K \in \mathcal{T}_h} \left\| w_h \right\|_K^2
$$

+ $(c_T + c\delta^{-1}) \left(\sum_{F \in \partial \mathcal{E}_N} \left\| |\frac{1}{2} \beta \cdot n|^{1/2} \right\|_F^2 + \sum_{K \in \mathcal{T}_h} \left\| |\beta_s - \frac{1}{2} \beta \cdot n|^{1/2} \left[\boldsymbol{w}_h \right] \right\|_{\partial K}^2 \right)$
+ $(c_T + c\epsilon \delta^{-1}) \left(\sum_{K \in \mathcal{T}_h} \epsilon \left\| \nabla w_h \right\|_K^2 + \sum_{K \in \mathcal{T}_h} \epsilon h_K^{-1} \left\| [\boldsymbol{w}_h] \right\|_{\mathcal{Q}_K}^2 \right).$

The result follows by choosing $\delta = 1/16$.

We are now ready to prove eq. $(4.2a)$.

Proof of eq. [\(4.2a\)](#page-7-3). Choose $\delta t_0 = 1/8$. When $\delta t_{\mathcal{K}} \leq \delta t_0$ for all $\mathcal{K} \in \mathcal{T}_h$ we find, by combining lemmas 4.2 and 4.5 ,

$$
a_h(\boldsymbol{w}_h, \boldsymbol{\Pi}_h (\varphi \boldsymbol{w}_h)) \geq (\frac{1}{4}(T + \chi) - c_T) \left(\sum_{\mathcal{K} \in \mathcal{T}_h} \varepsilon \left\| \overline{\nabla} w_h \right\|_{\mathcal{K}}^2 + \sum_{\mathcal{K} \in \mathcal{T}_h} \varepsilon h_K^{-1} \left\| [\boldsymbol{w}_h] \right\|_{\mathcal{Q}_\mathcal{K}}^2 \right) + \frac{1}{4} \sum_{\mathcal{K} \in \mathcal{T}_h} \left\| w_h \right\|_{\mathcal{K}}^2
$$

+ $(T + \chi - c_T) \left(\sum_{F \in \partial \mathcal{E}_N} \left\| |\frac{1}{2} \beta \cdot n|^{1/2} \right\|_{F}^2 + \sum_{\mathcal{K} \in \mathcal{T}_h} \left\| |\beta_s - \frac{1}{2} \beta \cdot n|^{1/2} \left[\boldsymbol{w}_h \right] \right\|_{\partial \mathcal{K}}^2 \right).$

Choosing χ to satisfy $\chi \geq 4c$ in addition to the conditions of lemma [4.2,](#page-7-1) we obtain

$$
a_h(\boldsymbol{w}_h, \boldsymbol{\Pi}_h(\varphi \boldsymbol{w}_h)) \geq \frac{1}{4} {\|\|\boldsymbol{w}_h\|_v^2} \geq c_T^{-1} {\|\|\boldsymbol{w}_h\|_v} {\|\|\boldsymbol{\Pi}_h(\varphi \boldsymbol{w}_h)\|_v},
$$
\n(4.19)

 \Box

where the second inequality is due to lemma [4.4.](#page-9-3) We therefore conclude eq. $(4.2a)$. \Box

4.2 The inf-sup condition with respect to $\|\cdot\|_{s}$

To prove eq. [\(4.2b\)](#page-7-4), we first construct the test function $y_h := (y_h, \vartheta_h)$ as a function of $w_h =$ $(w_h, \varkappa_h) \in \mathbf{V}_h$. The elemental test function y_h is defined as:

$$
y_h := \tau_\varepsilon \partial_t w_h. \tag{4.20a}
$$

To define the facet test function ϑ_h we consider four different sets of facets. First we consider facets F in $\partial\mathcal{K}_1\cap\partial\mathcal{K}_2\cap\mathcal{Q}_h^i$ and such that there is no difference in the refinement level in the time direction between \mathcal{K}_1 and \mathcal{K}_2 . This means that $\delta t_{\mathcal{K}_1} = \delta t_{\mathcal{K}_2} := \delta t_{\mathcal{K}}$ and, since \mathcal{K}_1 and \mathcal{K}_2 must come from the same space-time slab, $\Delta t_{\mathcal{K}_1} = \Delta t_{\mathcal{K}_2} := \Delta t_{\mathcal{K}}$. We then define:

$$
\vartheta_h := \begin{cases}\n\Delta t_{\mathcal{K}} \partial_t \varkappa_h, & \delta t_{\mathcal{K}} \le h_{K_1} \le \varepsilon, \delta t_{\mathcal{K}} \le h_{K_2} \le \varepsilon, \\
\Delta t_{\mathcal{K}} \varepsilon^{1/2} \partial_t \varkappa_h, & \delta t_{\mathcal{K}} \le \varepsilon < h_{K_1}, \delta t_{\mathcal{K}} \le \varepsilon < h_{K_2}, \\
0, & \text{otherwise.}\n\end{cases} \tag{4.20b}
$$

We next consider facets F in $\partial \mathcal{K}_1 \cap \partial \mathcal{K}_2 \cap \mathcal{Q}_h^i$ and such that there is one level of refinement difference between \mathcal{K}_1 and \mathcal{K}_2 in the time direction. Without loss of generality, we assume that $2\delta t_{\mathcal{K}_1} = \delta t_{\mathcal{K}_2}$. Furthermore, since \mathcal{K}_1 and \mathcal{K}_2 must come from the same space-time slab, $\Delta t_{\mathcal{K}_1} = \Delta t_{\mathcal{K}_2} := \Delta t_{\mathcal{K}}$. We then define:

$$
\vartheta_h := \begin{cases} \Delta t_{\mathcal{K}} \partial_t \varkappa_h, & \delta t_{\mathcal{K}_1} \le h_{K_1} \le \varepsilon, \delta t_{\mathcal{K}_2} \le h_{K_2} \le \varepsilon, \\ \Delta t_{\mathcal{K}} \varepsilon^{1/2} \partial_t \varkappa_h, & \delta t_{\mathcal{K}_1} \le \varepsilon < h_{K_1}, \delta t_{\mathcal{K}_2} \le \varepsilon < h_{K_2}, \\ 0, & \text{otherwise.} \end{cases} \tag{4.20c}
$$

For facets F in $\partial \mathcal{K} \cap \mathcal{Q}_h^b$, we define:

$$
\vartheta_h := \begin{cases} \Delta t_{\mathcal{K}} \partial_t \varkappa_h, & \delta t_{\mathcal{K}} \le h_K \le \varepsilon, \\ \Delta t_{\mathcal{K}} \varepsilon^{1/2} \partial_t \varkappa_h, & \delta t_{\mathcal{K}} \le \varepsilon < h_K, \\ 0, & \text{otherwise.} \end{cases} \tag{4.20d}
$$

Finally, for facets F in \mathcal{R}_h , we define:

$$
\vartheta_h := 0. \tag{4.20e}
$$

We observe from definition eq. [\(4.20\)](#page-12-1) that $\vartheta_h \equiv 0$ on $\partial \mathcal{T}_h^c$, which denotes the set of element boundaries of space-time elements in \mathcal{T}_h^c . Furthermore, for any space-time element $\mathcal{K} \in \mathcal{T}_h^{dx} :=$ $\mathcal{T}_h^d \cup \mathcal{T}_h^x$, we introduce $\mathcal{Q}_{\mathcal{K}}^0$ to denote those \mathcal{Q} -faces on which ϑ_h is prescribed in eqs. [\(4.20b\)](#page-12-2) and [\(4.20c\)](#page-12-3) to be zero. We will define $\mathcal{Q}_h^0 := \cup_{\mathcal{K} \in \mathcal{T}_h} \mathcal{Q}_{\mathcal{K}}^0$. Consider now $\mathcal{K} \in \mathcal{T}_h^{dx,0}$, which denotes the set of space-time elements in \mathcal{T}_h^{dx} for which $\mathcal{Q}_{\mathcal{K}}^0 \neq \emptyset$. Then, there exists a K' such that $\partial \mathcal{K}' \cap \partial \mathcal{K} \neq \emptyset$ and that either $h_K \leq \varepsilon \leq h_{K'}$ (or $h_{K'} \leq \varepsilon \leq h_K$), or $\delta t_K \leq \varepsilon \leq \delta t_{K'}$ (or $\delta t_{\mathcal{K}'} \leq \varepsilon \leq \delta t_{\mathcal{K}}$. For the former case, since spatial elements are shape-regular and the difference of refinement levels in the spatial direction between two adjacent space-time elements is at most one, we have $c^{-1}h_{K'} \leq h_K \leq ch_{K'}$. If the latter case holds, since $\delta t_{\mathcal{K}} = \frac{1}{2}$ $\frac{1}{2}\delta t_{\mathcal{K}'},$ it holds that $\delta t_{\mathcal{K}} \sim \varepsilon$. Therefore,

$$
c^{-1}h_K \le \varepsilon \le ch_K \quad \text{or} \quad c^{-1}\delta t_K \le \varepsilon \le c\delta t_K \qquad \forall K \in \mathcal{T}_h^{dx,0}.\tag{4.21}
$$

Lemmas [4.6](#page-13-0) and [4.7](#page-14-0) will be used to prove eq. [\(4.2b\)](#page-7-4). The proofs of these lemmas will repeatedly use the following set of inequalities: For all $\mathcal{K} \in \mathcal{T}_h$,

$$
h_K^{-1} \le \delta t_K^{-1}, \quad \Delta t_K \le c\delta t_K, \quad \tau_\varepsilon \le \Delta t_K, \quad \varepsilon \le 1. \tag{4.22}
$$

Lemma 4.6. Assume that $\delta t_K \leq h_K$ for all space-time elements $K \in \mathcal{T}_h$. Let $\mathbf{w}_h = (w_h, \varkappa_h) \in \mathbf{V}_h$ and let y_h be defined by eq. [\(4.20\)](#page-12-1). The following holds:

$$
\|\boldsymbol{y}_h\|_{s} \leq c \|\boldsymbol{w}_h\|_{s}.\tag{4.23}
$$

Proof. We start with the volume terms of $\|\cdot\|_s$. Using eq. [\(3.5a\)](#page-5-0) and eq. [\(4.22\)](#page-13-1), we have:

$$
\sum_{\mathcal{K}\in\mathcal{T}_h} \|y_h\|_{\mathcal{K}}^2 \le \sum_{\mathcal{K}\in\mathcal{T}_h} \tau_{\varepsilon} \|\partial_t w_h\|_{\mathcal{K}}^2.
$$
\n(4.24)

For the diffusive volume term, using commutativity of $\overline{\nabla}$ and ∂_t , eq. [\(3.5a\)](#page-5-0) and eq. [\(4.22\)](#page-13-1):

$$
\sum_{\mathcal{K}\in\mathcal{T}_h} \varepsilon \left\| \overline{\nabla} y_h \right\|_{\mathcal{K}}^2 = \sum_{\mathcal{K}\in\mathcal{T}_h} \tau_{\varepsilon}^2 \varepsilon \left\| \partial_t \left(\overline{\nabla} w_h \right) \right\|_{\mathcal{K}}^2 \leq c \sum_{\mathcal{K}\in\mathcal{T}_h} \varepsilon \left\| \overline{\nabla} w_h \right\|_{\mathcal{K}}^2. \tag{4.25}
$$

The time-derivative volume term is treated similarly, using eq. [\(4.22\)](#page-13-1):

$$
\sum_{\mathcal{K}\in\mathcal{T}_h} \tau_{\varepsilon} \left\| \partial_t y_h \right\|_{\mathcal{K}}^2 \leq c \sum_{\mathcal{K}\in\mathcal{T}_h} \tau_{\varepsilon} \left\| \partial_t w_h \right\|_{\mathcal{K}}^2.
$$
\n(4.26)

For the diffusive facet term in the definition of $\|\|\cdot\|_s$, we use lemma [3.1,](#page-5-5) eq. [\(3.5c\)](#page-5-4), that ϑ_h vanishes on $\partial \mathcal{T}_h^c$, that ϑ_h vanishes on $\mathcal{Q}_{\mathcal{K}}^0$ when $\mathcal{K} \in \mathcal{T}_h^{dx}$, that $\varepsilon \leq \delta t_{\mathcal{K}}$ and $\varepsilon h_K^{-2} \tau_{\varepsilon} \leq 1$ on \mathcal{T}_h^c , and that $\varepsilon h_K^{-2} \tau_\varepsilon \leq c$ on $\mathcal{T}_h^{dx,0}$ $\eta_h^{\text{cat},0}$ due to eq. [\(4.21\)](#page-12-4):

$$
\sum_{\mathcal{K}\in\mathcal{T}_{h}}\varepsilon h_{K}^{-1} \left\|[\mathbf{y}_{h}]\right\|_{\mathcal{Q}_{\mathcal{K}}}^{2}
$$
\n
$$
= \sum_{\mathcal{K}\in\mathcal{T}_{h}^{dx}}\varepsilon h_{K}^{-1}\tau_{\varepsilon}^{2} \left\|[\partial_{t}\mathbf{w}_{h}]\right\|_{\mathcal{Q}_{\mathcal{K}}\backslash\mathcal{Q}_{\mathcal{K}}^{0}}^{2} + \sum_{\mathcal{K}\in\mathcal{T}_{h}^{dx}}\varepsilon h_{K}^{-1}\tau_{\varepsilon}^{2} \left\|\partial_{t}\mathbf{w}_{h}\right\|_{\mathcal{Q}_{\mathcal{K}}}^{2} + \sum_{\mathcal{K}\in\mathcal{T}_{h}^{dx}}\varepsilon h_{K}^{-1}\tau_{\varepsilon}^{2} \left\|\partial_{t}\mathbf{w}_{h}\right\|_{\mathcal{Q}_{\mathcal{K}}}^{2} \right\}
$$
\n
$$
\leq c \sum_{\mathcal{K}\in\mathcal{T}_{h}}\varepsilon h_{K}^{-1} \left\|[\mathbf{w}_{h}]\right\|_{\mathcal{Q}_{\mathcal{K}}}^{2} + c \sum_{\mathcal{K}\in\mathcal{T}_{h}}\tau_{\varepsilon} \left\|\partial_{t}\mathbf{w}_{h}\right\|_{\mathcal{K}}^{2}.
$$
\n
$$
(4.27)
$$

For the advective facet term, using lemma [3.1,](#page-5-5) eq. [\(3.5d\)](#page-5-6), eq. [\(4.22\)](#page-13-1), that $\tilde{\varepsilon}^2 \leq \varepsilon h_K^{-1}$ on $K \in \mathcal{T}_h^{dx}$
since $h_K \leq \varepsilon$ on \mathcal{T}_h^d and $\tilde{\varepsilon} = \varepsilon^{1/2}$ on \mathcal{T}_h^x :

$$
\sum_{\mathcal{K}\in\mathcal{T}_h} ||\beta_s - \frac{1}{2}\beta \cdot n|^{1/2} [\mathbf{y}_h]||_{\partial \mathcal{K}}^2
$$
\n
$$
\leq c \Big(\sum_{\mathcal{K}\in\mathcal{T}_h^{dx}} \tau_{\varepsilon}^2 ||[\partial_t \mathbf{w}_h]||_{\mathcal{Q}_{\mathcal{K}}\setminus\mathcal{Q}_{\mathcal{K}}^0}^2 + \sum_{\mathcal{K}\in\mathcal{T}_h^{dx}} \tau_{\varepsilon}^2 ||\partial_t w_h||_{\mathcal{R}_{\mathcal{K}}\cup\mathcal{Q}_{\mathcal{K}}^0}^2 + \sum_{\mathcal{K}\in\mathcal{T}_h^{c}} \tau_{\varepsilon}^2 ||\partial_t w_h||_{\partial \mathcal{K}}^2 \Big) \Big) \tag{4.28}
$$
\n
$$
\leq c \Big(\sum_{\mathcal{K}\in\mathcal{T}_h^{dx}} \varepsilon h_K^{-1} ||[\mathbf{w}_h]||_{\mathcal{Q}_{\mathcal{K}}\setminus\mathcal{Q}_{\mathcal{K}}^0}^2 + c \sum_{\mathcal{K}\in\mathcal{T}_h^{c}} \tau_{\varepsilon} ||\partial_t w_h||_{\mathcal{K}}^2.
$$

Finally, the Neumann boundary term is bounded using the triangle inequality, Young's inequality, lemma [3.1,](#page-5-5) eq. [\(3.5c\)](#page-5-4), eq. [\(4.22\)](#page-13-1), and that $h_K \leq \varepsilon$ for $K \in \mathcal{T}_h^d$:

$$
\sum_{\mathcal{K}\in\mathcal{T}_{h}^{dx}}\left\|\frac{1}{2}\beta\cdot n\right\|^{1/2}\vartheta_{h}\right\|_{\mathcal{Q}_{\mathcal{K}}\cap\partial\mathcal{E}_{N}}^{2}\leq c\sum_{\mathcal{K}\in\mathcal{T}_{h}^{dx}}\tau_{\varepsilon}^{2}\left\|\left[\partial_{t}\mathbf{w}_{h}\right]\right\|_{\mathcal{Q}_{\mathcal{K}}\cap\partial\mathcal{E}_{N}}^{2}+c\sum_{\mathcal{K}\in\mathcal{T}_{h}^{dx}}\tau_{\varepsilon}^{2}\left\|\partial_{t}w_{h}\right\|_{\mathcal{Q}_{\mathcal{K}}\cap\partial\mathcal{E}_{N}}^{2}
$$
\n
$$
\leq c\sum_{\mathcal{K}\in\mathcal{T}_{h}^{dx}}\varepsilon h_{K}^{-1}\left\|\left[\mathbf{w}_{h}\right]\right\|_{\mathcal{Q}_{\mathcal{K}}\cap\partial\mathcal{E}_{N}}^{2}+c\sum_{\mathcal{K}\in\mathcal{T}_{h}^{dx}}\tau_{\varepsilon}\left\|\partial_{t}w_{h}\right\|_{\mathcal{K}}^{2}.
$$
\n(4.29)

Combining eqs. (4.24) to (4.29) yields eq. (4.23) .

Lemma 4.7. Assume that $\delta t_K \leq h_K$ for all space-time elements $K \in \mathcal{T}_h$. Let $\mathbf{w}_h = (w_h, \mathbf{z}_h) \in \mathbf{V}_h$, let y_h be defined as in eq. [\(4.20\)](#page-12-1), and let $\Pi_h(\varphi\bm{w}_h)$ be defined as in lemma [4.4.](#page-9-3) There exists a positive constant c such that

$$
\left\|\left|\boldsymbol{w}_h\right\|\right\|_s^2 \leq a_h(\boldsymbol{w}_h, 2(\boldsymbol{y}_h + c\boldsymbol{\Pi}_h(\varphi \boldsymbol{w}_h))).
$$

Proof. Let us first note that ϑ_h vanishes on \mathcal{R}_h and $\partial \mathcal{T}_h^c$. Therefore, defining $\mathcal{Q}_h^{dx} := \partial \mathcal{T}_h^{dx} \cap \mathcal{Q}_h$, we find after some algebraic manipulation that:

$$
a_{h,c}(\boldsymbol{w}_h, \boldsymbol{y}_h) = (\nabla \cdot (\beta w_h), y_h)_{\mathcal{T}_h} + \langle (\beta_s - \frac{1}{2}\beta \cdot n) [\boldsymbol{w}_h], y_h \rangle_{\partial \mathcal{T}_h} - \langle \frac{1}{2}\beta \cdot n [\boldsymbol{w}_h], y_h \rangle_{\partial \mathcal{T}_h} - \langle \frac{1}{2} (|\beta \cdot n| - \beta \cdot n) \varkappa_h, [\boldsymbol{y}_h] \rangle_{\partial \mathcal{E}_N \cap \mathcal{Q}_h^{dx}} + \langle \frac{1}{2} (|\beta \cdot n| - \beta \cdot n) \varkappa_h, y_h \rangle_{\partial \mathcal{E}_N \cap \mathcal{Q}_h^{dx}} + \langle \beta_s [\boldsymbol{w}_h], [\boldsymbol{y}_h] \rangle_{\mathcal{Q}_h^{dx} \setminus \mathcal{Q}_h^0} - \langle \beta_s [\boldsymbol{w}_h], y_h \rangle_{\mathcal{Q}_h^{dx} \setminus \mathcal{Q}_h^0}.
$$

Furthermore, since $(\nabla \cdot (\beta w_h), y_h)_{\mathcal{T}_h} = (\partial_t w_h, \tau_{\varepsilon} \partial_t w_h)_{\mathcal{T}_h} + (\overline{\nabla} \cdot (\overline{\beta} w_h), \tau_{\varepsilon} \partial_t w_h)_{\mathcal{T}_h}$, we find that

$$
\sum_{K \in \mathcal{T}_h} \tau_{\varepsilon} \|\partial_t w_h\|_{K}^2 = a_h(\boldsymbol{w}_h, \boldsymbol{y}_h) - a_{h,d}(\boldsymbol{w}_h, \boldsymbol{y}_h) - (\overline{\nabla} \cdot (\overline{\beta} w_h), \tau_{\varepsilon} \partial_t w_h)_{\mathcal{T}_h}
$$

$$
- \langle (\beta_s - \frac{1}{2}\beta \cdot n) [w_h], \tau_{\varepsilon} \partial_t w_h \rangle_{\partial \mathcal{T}_h} + \langle \frac{1}{2}\beta \cdot n [w_h], \tau_{\varepsilon} \partial_t w_h \rangle_{\partial \mathcal{T}_h}
$$

$$
- \langle \beta_s [w_h], \tau_{\varepsilon} [\partial_t w_h] \rangle_{\mathcal{Q}_h^{dx} \setminus \mathcal{Q}_h^0} + \langle \beta_s [w_h], \tau_{\varepsilon} \partial_t w_h \rangle_{\mathcal{Q}_h^{dx} \setminus \mathcal{Q}_h^0}
$$

$$
+ \langle \frac{1}{2} (|\beta \cdot n| - \beta \cdot n) \varkappa_h, \tau_{\varepsilon} [\partial_t w_h] \rangle_{\partial \mathcal{E}_N \cap \mathcal{Q}_h^{dx}} - \langle \frac{1}{2} (|\beta \cdot n| - \beta \cdot n) \varkappa_h, \tau_{\varepsilon} \partial_t w_h \rangle_{\partial \mathcal{E}_N \cap \mathcal{Q}_h^{dx}}
$$

$$
= a_h(\boldsymbol{w}_h, \boldsymbol{y}_h) - a_{h,d}(\boldsymbol{w}_h, \boldsymbol{y}_h) - (\overline{\nabla} \cdot (\overline{\beta} w_h), \tau_{\varepsilon} \partial_t w_h)_{\mathcal{T}_h} + T_1 + T_2 + T_3 + T_4 + T_5 + T_6.
$$

(4.30)

We will bound the last eight terms on the right hand side of the above equation. First, by eq. (3.11) , lemma [4.6,](#page-13-0) and Young's inequality, we have

$$
a_{h,d}(\boldsymbol{w}_h, \boldsymbol{y}_h) \leq c \|\|\boldsymbol{w}_h\|_{v} \|\|\boldsymbol{w}_h\|_{s} \leq c \left(\left(\sum_{\mathcal{K}\in\mathcal{T}_h} \tau_{\varepsilon} \|\partial_t w_h\|_{\mathcal{K}}^2 \right)^{1/2} + \|\|\boldsymbol{w}_h\|_{v} \right) \|\|\boldsymbol{w}_h\|_{v}
$$

$$
\leq c \left(1 + \delta^{-1}\right) \|\|\boldsymbol{w}_h\|_{v}^2 + \frac{1}{2}\delta \sum_{\mathcal{K}\in\mathcal{T}_h} \tau_{\varepsilon} \|\partial_t w_h\|_{\mathcal{K}}^2.
$$
\n
$$
(4.31)
$$

Next, using the Cauchy–Schwarz and Young's inequalities, $\delta t_{\mathcal{K}} \leq \varepsilon$ for $\mathcal{K} \in \mathcal{T}_h^{dx}$, eq. [\(3.5b\)](#page-5-2) and eq. [\(4.22\)](#page-13-1):

$$
\begin{split} (\overline{\nabla} \cdot (\overline{\beta} w_h) \cdot \tau_{\varepsilon} \partial_t w_h) \tau_h &\leq \frac{\delta}{2} \sum_{\mathcal{K} \in \mathcal{T}_h} \tau_{\varepsilon} \left\| \partial_t w_h \right\|_{\mathcal{K}}^2 + c \delta^{-1} \sum_{\mathcal{K} \in \mathcal{T}_h} \tau_{\varepsilon} \left\| \overline{\nabla} w_h \right\|_{\mathcal{K}}^2 \\ &\leq \frac{\delta}{2} \sum_{\mathcal{K} \in \mathcal{T}_h} \tau_{\varepsilon} \left\| \partial_t w_h \right\|_{\mathcal{K}}^2 + c \delta^{-1} \sum_{\mathcal{K} \in \mathcal{T}_h} \varepsilon \left\| \overline{\nabla} w_h \right\|_{\mathcal{K}}^2 + c \delta^{-1} \sum_{\mathcal{K} \in \mathcal{T}_h} \left\| w_h \right\|_{\mathcal{K}}^2. \end{split} \tag{4.32}
$$

 T_1 and T_2 can be bounded using the Cauchy–Schwarz inequality, $\frac{1}{2} |\beta \cdot n| \leq |\beta_s - \frac{1}{2}|$ $\frac{1}{2}\beta \cdot n$ for all $F \in \partial \mathcal{T}_h$, eq. [\(3.5d\)](#page-5-6), eq. [\(4.22\)](#page-13-1), and Young's inequality:

$$
T_1 + T_2 \leq \frac{1}{2} \delta \sum_{\mathcal{K} \in \mathcal{T}_h} \tau_{\varepsilon} \left\| \partial_t w_h \right\|_{\mathcal{K}}^2 + c \delta^{-1} \sum_{\mathcal{K} \in \mathcal{T}_h} \left\| \left| \beta_s - \frac{1}{2} \beta \cdot n \right|^{1/2} \left[\boldsymbol{w}_h \right] \right\|_{\partial \mathcal{K}}^2. \tag{4.33}
$$

Similarly T_3 and T_4 are bounded using the Cauchy–Schwarz inequality, lemma [3.1,](#page-5-5) eq. [\(3.5c\)](#page-5-4), eq. [\(3.9\)](#page-6-1), $\delta t_{\mathcal{K}} \leq h_K \leq \varepsilon$ for $\mathcal{K} \in \mathcal{T}_h^d$, eq. [\(4.22\)](#page-13-1), and Young's inequality. Note that we also make use of $\tilde{\varepsilon} \leq \varepsilon^{1/2} h_K^{-1/2}$ on $\mathcal{T}_h^{\,dx}$ since on $\mathcal{T}_h^{\,d}$, $\tilde{\varepsilon} = 1$ and $h_K \leq \varepsilon$ while on \mathcal{T}_h^x , $\tilde{\varepsilon} = \varepsilon^{1/2}$ and $h_K \leq 1$:

$$
T_3 + T_4 \leq c \sum_{\mathcal{K} \in \mathcal{T}_{h}^{dx}} \tau_{\varepsilon} ||\beta_s - \frac{1}{2}\beta \cdot n|^{1/2} [\boldsymbol{w}_h] ||_{\mathcal{Q}_{\mathcal{K}} \setminus \mathcal{Q}_{\mathcal{K}}^0} ((\delta t_{\mathcal{K}}^{-1} + h_K^{-1}) ||[\boldsymbol{w}_h] ||_{\mathcal{Q}_{\mathcal{K}} \setminus \mathcal{Q}_{\mathcal{K}}^0} + h_K^{-1/2} ||\partial_t w_h||_{\mathcal{K}})
$$

\n
$$
\leq c \sum_{\mathcal{K} \in \mathcal{T}_{h}^{dx}} |||\beta_s - \frac{1}{2}\beta \cdot n|^{1/2} [\boldsymbol{w}_h] ||_{\mathcal{Q}_{\mathcal{K}} \setminus \mathcal{Q}_{\mathcal{K}}^0} (\tilde{\varepsilon} ||[\boldsymbol{w}_h] ||_{\mathcal{Q}_{\mathcal{K}} \setminus \mathcal{Q}_{\mathcal{K}}^0} + \Delta t_{\mathcal{K}}^{1/2} \tilde{\varepsilon} ||\partial_t w_h||_{\mathcal{K}})
$$

\n
$$
\leq c \sum_{\mathcal{K} \in \mathcal{T}_{h}^{dx}} |||\beta_s - \frac{1}{2}\beta \cdot n|^{1/2} [\boldsymbol{w}_h] ||_{\mathcal{Q}_{\mathcal{K}} \setminus \mathcal{Q}_{\mathcal{K}}^0} (\varepsilon^{1/2} h_K^{-1/2} ||[\boldsymbol{w}_h] ||_{\mathcal{Q}_{\mathcal{K}} \setminus \mathcal{Q}_{\mathcal{K}}^0} + \tau_{\varepsilon}^{1/2} ||\partial_t w_h||_{\mathcal{K}})
$$

\n
$$
\leq \frac{1}{2} \delta \sum_{\mathcal{K} \in \mathcal{T}_h} \tau_{\varepsilon} ||\partial_t w_h||_{\mathcal{K}}^2 + c \sum_{\mathcal{K} \in \mathcal{T}_h} \varepsilon h_K^{-1} ||[\boldsymbol{w}_h] ||_{\mathcal{Q}_{\mathcal{K}}}^2 + c (1 + \delta^{-1}) \sum_{\mathcal{K} \in \mathcal{T}_h} |||\beta_s - \frac{1}{2}\beta \cdot n|^{1/2} [\boldsymbol{w}_h] ||_{\partial \mathcal{K}}^2.
$$

\n(4.34)

Similarly, to bound T_5 and T_6 , we use the Cauchy–Schwarz inequality, lemma [3.1,](#page-5-5) eq. [\(3.5c\)](#page-5-4), that $\delta t_{\mathcal{K}} \leq h_K \leq \varepsilon$ for $\mathcal{K} \in \mathcal{T}_h^d$, eq. [\(4.22\)](#page-13-1), $\tilde{\varepsilon} \leq \varepsilon^{1/2} h_K^{-1/2}$ on \mathcal{T}_h^{dx} and Young's inequality:

$$
T_5 + T_6 \leq \frac{1}{2}\delta \sum_{\mathcal{K}\in\mathcal{T}_h} \tau_{\varepsilon} \|\partial_t w_h\|_{\mathcal{K}}^2 + c \sum_{\mathcal{K}\in\mathcal{T}_h} \varepsilon h_K^{-1} \|\left[w_h\right]\|_{\mathcal{Q}_{\mathcal{K}}}^2 + c\left(1 + \delta^{-1}\right) \sum_{\mathcal{K}\in\mathcal{T}_h} \left\|\left(\frac{1}{2}\beta \cdot n\right)\right\|_{\mathcal{Q}_{\mathcal{K}}\cap\partial\mathcal{E}_N}^2.
$$
\n
$$
(4.35)
$$

Combining eqs. [\(4.30\)](#page-14-1) to [\(4.35\)](#page-15-0) and choosing $\delta = 1/5$ we obtain

$$
\sum_{\mathcal{K}\in\mathcal{T}_h} \tau_{\varepsilon} \|\partial_t w_h\|_{\mathcal{K}}^2 \le a_h(\boldsymbol{w}_h, 2\boldsymbol{y}_h) + c \|\boldsymbol{w}_h\|_{v}^2.
$$
\n(4.36)

Adding $\|\|\boldsymbol{w}_h\|_{v}^2$ to both sides of eq. [\(4.36\)](#page-15-1), the first bound in eq. [\(4.19\)](#page-11-1) yields the result. \Box

We end this section by proving eq. $(4.2b)$.

Proof of eq. [\(4.2b\)](#page-7-4). By eq. [\(4.7b\)](#page-8-8) and using that $\tau_{\varepsilon} \leq c\varepsilon$, because on \mathcal{T}_h^{dx} , $\tau_{\varepsilon} \leq \Delta t \leq c\delta t \leq c\varepsilon$ and on \mathcal{T}_h^c , $\tau_{\varepsilon} = \Delta t_{\mathcal{K}} \varepsilon \leq \varepsilon (\leq c \varepsilon)$, we find

$$
\sum_{\mathcal{K}\in\mathcal{T}_{h}}\tau_{\varepsilon}\left\|\partial_{t}\left(\Pi_{h}\left(\varphi w_{h}\right)\right)\right\|_{\mathcal{K}}^{2}\leq c\left(\sum_{\mathcal{K}\in\mathcal{T}_{h}}\tau_{\varepsilon}\left\|\partial_{t}\left(\varphi w_{h}\right)\right\|_{\mathcal{K}}^{2}+\sum_{\mathcal{K}\in\mathcal{T}_{h}}\tau_{\varepsilon}\left\|\overline{\nabla}\left(\varphi w_{h}\right)\right\|_{\mathcal{K}}^{2}\right) \n\leq c\left(c_{T}^{2}\left(\sum_{\mathcal{K}\in\mathcal{T}_{h}}\tau_{\varepsilon}\left\|\partial_{t}w_{h}\right\|_{\mathcal{K}}^{2}+c\sum_{\mathcal{K}\in\mathcal{T}_{h}}\varepsilon\left\|\overline{\nabla}w_{h}\right\|_{\mathcal{K}}^{2}\right)+\sum_{\mathcal{K}\in\mathcal{T}_{h}}\|w_{h}\|_{\mathcal{K}}^{2}\right) \n\leq c_{T}^{2}\|\left\|\left\|w_{h}\right\|_{s}^{2}.
$$

Therefore, using lemma [4.4,](#page-9-3) we conclude that

$$
\|\Pi_h(\varphi \mathbf{w}_h)\|_{s} \leq c_T \|\mathbf{w}_h\|_{s}.
$$
\n(4.37)

Equation [\(4.2b\)](#page-7-4) can now be shown to hold after combining eq. [\(4.37\)](#page-15-2) with lemmas [4.6](#page-13-0) and [4.7.](#page-14-0) \Box

4.3 The inf-sup condition with respect to $\|\cdot\|_{ss}$

Proof of theorem [4.1.](#page-7-2) We construct the test function $\kappa_h := (\kappa_h, \varsigma_h)$ such that for $K \in \mathcal{T}_h$, $\kappa_h|_K :=$ $\frac{\delta t_{\mathcal{K}}h_{\mathcal{K}}^2}{\delta t_{\mathcal{K}}+h_{\mathcal{K}}}\Pi_h(\beta \cdot \nabla w_h)$ while ς_h vanishes on all faces of \mathcal{F}_h . We first show that there exists a positive constant c_1 , independent of h_K , δt_K , ε , and T such that the following holds:

$$
\|\kappa_h\|_s \le c_1 \|w_h\|_{sd} \,. \tag{4.38}
$$

We bound each term of $||\cdot|||_s$, starting with the volume terms. Noting that $\frac{\delta t_{\mathcal{K}}h_{\mathcal{K}}^2}{\delta t_{\mathcal{K}}+h_{\mathcal{K}}} \leq 1$ and using the definition of $\lVert \cdot \rVert_{sd}$ in eq. [\(3.8c\)](#page-6-3), we have:

$$
\sum_{\mathcal{K}\in\mathcal{T}_h} \|\kappa_h\|_{\mathcal{K}}^2 \le \|w_h\|_{sd}^2.
$$
\n(4.39)

The diffusive volume term is bounded using $\varepsilon \leq 1$, $\frac{\delta t \kappa h_K^2}{\delta t \kappa + h_K} h_K^{-2} \leq 1$ and eq. [\(3.5b\)](#page-5-2):

$$
\sum_{\mathcal{K}\in\mathcal{T}_h} \varepsilon \left\| \overline{\nabla} \kappa_h \right\|_{\mathcal{K}}^2 \le c \sum_{\mathcal{K}\in\mathcal{T}_h} \left(\frac{\delta t_{\mathcal{K}} h_K^2}{\delta t_{\mathcal{K}} + h_K} \right)^2 h_K^{-2} \left\| \Pi_h \left(\beta \cdot \nabla w_h \right) \right\|_{\mathcal{K}}^2 \le c \|w_h\|_{sd}^2. \tag{4.40}
$$

For the time derivative volume term, we need eq. (4.22) and eq. $(3.5a)$:

$$
\sum_{\mathcal{K}\in\mathcal{T}_h} \tau_{\varepsilon} \|\partial_t \kappa_h\|_{\mathcal{K}}^2 \leq c \sum_{\mathcal{K}\in\mathcal{T}_h} \delta t_{\mathcal{K}} \left(\delta t_{\mathcal{K}}^{-1} + h_K^{-1}\right)^2 \left(\frac{\delta t_{\mathcal{K}} h_K^2}{\delta t_{\mathcal{K}} + h_K}\right)^2 \left\|\Pi_h \left(\beta \cdot \nabla w_h\right)\right\|_{\mathcal{K}}^2 \leq c \|w_h\|_{sd}^2. \tag{4.41}
$$

Next we turn to the facet terms. To bound the diffusive facet term, we apply eq. $(3.5c)$:

$$
\sum_{\mathcal{K}\in\mathcal{T}_{h}}\varepsilon h_{K}^{-1} \left\|[\boldsymbol{\kappa}_{h}]\right\|_{\mathcal{Q}_{\mathcal{K}}}^{2} = \sum_{\mathcal{K}\in\mathcal{T}_{h}}\varepsilon h_{K}^{-1} \left(\frac{\delta t_{\mathcal{K}} h_{K}^{2}}{\delta t_{\mathcal{K}} + h_{K}}\right)^{2} \left\|\Pi_{h}\left(\beta \cdot \nabla w_{h}\right)\right\|_{\mathcal{Q}_{\mathcal{K}}}^{2}
$$
\n
$$
\leq c \sum_{\mathcal{K}\in\mathcal{T}_{h}}\frac{\delta t_{\mathcal{K}} h_{K}^{2}}{\delta t_{\mathcal{K}} + h_{K}} \left\|\Pi_{h}\left(\beta \cdot \nabla w_{h}\right)\right\|_{\mathcal{K}}^{2} \leq c \left\|w_{h}\right\|_{sd}^{2}.
$$
\n(4.42)

We use eq. [\(3.5d\)](#page-5-6) and that $(\delta t_{\mathcal{K}}^{-1/2} + h_K^{-1/2})^2 \frac{\delta t_{\mathcal{K}} h_K}{\delta t_{\mathcal{K}} + h_I}$ $\frac{\partial t_{\mathcal{K}} h_K}{\partial t_{\mathcal{K}}+h_K} \leq 2$ to bound the advective facet term:

$$
\sum_{\mathcal{K}\in\mathcal{T}_{h}}\left\|\left|\beta_{s}-\frac{1}{2}\beta\cdot n\right|^{1/2}\left[\kappa_{h}\right]\right\|_{\partial\mathcal{K}}^{2}\leq c\sum_{\mathcal{K}\in\mathcal{T}_{h}}\left(\frac{\delta t_{\mathcal{K}}h_{\mathcal{K}}^{2}}{\delta t_{\mathcal{K}}+h_{\mathcal{K}}}\right)^{2}\left\|\Pi_{h}\left(\beta\cdot\nabla w_{h}\right)\right\|_{\partial\mathcal{K}}^{2}\n\leq c\sum_{\mathcal{K}\in\mathcal{T}_{h}}\frac{\delta t_{\mathcal{K}}h_{\mathcal{K}}^{2}}{\delta t_{\mathcal{K}}+h_{\mathcal{K}}}\left\|\Pi_{h}\left(\beta\cdot\nabla w_{h}\right)\right\|_{\mathcal{K}}^{2}\leq c\left\|w_{h}\right\|_{sd}^{2}.
$$
\n(4.43)

The Neumann boundary term vanishes since $\varsigma_h \equiv 0$. We can therefore conclude eq. [\(4.38\)](#page-16-1) from eqs. (4.39) to (4.43) .

We next show that there exists a positive constant c_2 , independent of h_K , δt_K , ε , and T such that

$$
||w_h||_{sd}^2 - c_2||w_h||_{s}||w_h||_{sd} \le a_h(\mathbf{w}_h, \kappa_h).
$$
\n(4.44)

We first write the advective part of the bilinear form as:

$$
a_{h,c}(\boldsymbol{w}_h,\boldsymbol{\kappa}_h) = (\nabla \cdot (\beta w_h), \kappa_h)_{\mathcal{T}_h} + \langle (\beta_s - \beta \cdot n) [\boldsymbol{w}_h], \kappa_h \rangle_{\partial \mathcal{T}_h} =: T_1 + T_2. \tag{4.45}
$$

We bound T_1 using the definition of the projection operator Π_h :

$$
T_{1} = \left((I - \Pi_{h}) \left(\beta \cdot \nabla w_{h} \right), \frac{\delta t_{\mathcal{K}} h_{K}^{2}}{\delta t_{\mathcal{K}} + h_{K}} \Pi_{h} \left(\beta \cdot \nabla w_{h} \right) \right)_{\mathcal{T}_{h}} + \left(\Pi_{h} \left(\beta \cdot \nabla w_{h} \right), \frac{\delta t_{\mathcal{K}} h_{K}^{2}}{\delta t_{\mathcal{K}} + h_{K}} \Pi_{h} \left(\beta \cdot \nabla w_{h} \right) \right)_{\mathcal{T}_{h}}
$$
\n
$$
= \| w_{h} \|_{sd}^{2} \,. \tag{4.46}
$$

For T_2 , we note that $|\beta_s - \beta \cdot n| \leq 2 |\beta_s - \frac{1}{2}|$ $\frac{1}{2}\beta \cdot n$ for any $F \in \mathcal{T}_h$. Then, also using eq. [\(3.5d\)](#page-5-6) and Hölder's inequality for sums,

$$
T_2 \leq c \sum_{\mathcal{K}\in\mathcal{T}_h} \left\| \beta_s - \frac{1}{2}\beta \cdot n \right\|^{1/2} \left[\boldsymbol{w}_h \right] \right\|_{\partial\mathcal{K}} \frac{\delta t_{\mathcal{K}} h_K^2}{\delta t_{\mathcal{K}} + h_K} \left\| \Pi_h \left(\beta \cdot \nabla w_h \right) \right\|_{\partial\mathcal{K}} \leq c \left\| \left\| \boldsymbol{w}_h \right\| \right\|_s \left\| w_h \right\|_{sd} . \tag{4.47}
$$

For the diffusive part of the bilinear form, we write:

$$
a_{h,d}(\boldsymbol{w}_h, \boldsymbol{\kappa}_h) = \left(\varepsilon \overline{\nabla} w_h, \overline{\nabla} \kappa_h\right)_{\mathcal{T}_h} - \left\langle \varepsilon \left[\boldsymbol{w}_h\right], \overline{\nabla}_{\bar{n}} \kappa_h \right\rangle_{\mathcal{Q}_h} - \left\langle \varepsilon \kappa_h, \overline{\nabla}_{\bar{n}} w_h \right\rangle_{\mathcal{Q}_h} + \left\langle \alpha \varepsilon h_K^{-1} \left[\boldsymbol{w}_h\right], \kappa_h \right\rangle_{\mathcal{Q}_h}
$$
\n
$$
=: I_1 + I_2 + I_3 + I_4.
$$
\n(4.48)

For I_1 , we apply the Cauchy–Schwarz inequality, eq. $(3.5b)$, and Hölder's inequality for sums:

$$
I_{1} \leq \sum_{\mathcal{K}\in\mathcal{T}_{h}} \varepsilon \|\overline{\nabla}w_{h}\|_{\mathcal{K}} \frac{\delta t_{\mathcal{K}}h_{\mathcal{K}}^{2}}{\delta t_{\mathcal{K}}+h_{\mathcal{K}}} \|\overline{\nabla}(\Pi_{h}(\beta \cdot \nabla w_{h}))\|_{\mathcal{K}}
$$
\n
$$
\leq c \sum_{\mathcal{K}\in\mathcal{T}_{h}} \varepsilon^{1/2} \|\overline{\nabla}w_{h}\|_{\mathcal{K}} \left(\frac{\delta t_{\mathcal{K}}h_{\mathcal{K}}^{2}}{\delta t_{\mathcal{K}}+h_{\mathcal{K}}}\right)^{1/2} \|\Pi_{h}(\beta \cdot \nabla w_{h})\|_{\mathcal{K}} \leq c \|\|\boldsymbol{w}_{h}\|_{s} \|w_{h}\|_{sd}.
$$
\n
$$
(4.49)
$$

For I_2 , we apply the Cauchy–Schwarz inequality, eqs. $(3.5b)$ and $(3.5c)$, and Hölder's inequality for sums:

$$
I_2 \leq \sum_{\mathcal{K}\in\mathcal{T}_h} \varepsilon \left\|[\boldsymbol{w}_h]\right\|_{\mathcal{Q}_\mathcal{K}} \frac{\delta t_\mathcal{K} h_\mathcal{K}^2}{\delta t_\mathcal{K} + h_\mathcal{K}} \left\|\overline{\nabla}\left(\Pi_h\left(\beta \cdot \nabla w_h\right)\right)\right\|_{\mathcal{Q}_\mathcal{K}} \leq c \sum_{\mathcal{K}\in\mathcal{T}_h} \varepsilon^{1/2} h_K^{-1/2} \left\|[\boldsymbol{w}_h]\right\|_{\mathcal{Q}_\mathcal{K}} \left(\frac{\delta t_\mathcal{K} h_K^2}{\delta t_\mathcal{K} + h_K}\right)^{1/2} \left\|\Pi_h\left(\beta \cdot \nabla w_h\right)\right\|_{\mathcal{K}} \leq c \left\|\|\boldsymbol{w}_h\right\|_{s}\|w_h\|_{s d}.
$$
\n(4.50)

Similarly for I_3 and I_4 , we apply the Cauchy–Schwarz inequality, eq. [\(3.5c\)](#page-5-4), and Hölder's inequality for sums:

$$
I_3 + I_4 \le c \| \boldsymbol{w}_h \|_s \| w_h \|_{sd}.
$$
\n(4.51)

Combining eqs. (4.45) to (4.51) , we conclude eq. (4.44) .

Combining eq. [\(4.44\)](#page-16-5) and eq. [\(4.38\)](#page-16-1) then yields:

$$
c_1^{-1}\left(\left\|w_h\right\|_{sd}-c_2\left\|{\boldsymbol w}_h\right\|_{s}\right)\leq \frac{a_h({\boldsymbol w}_h, \boldsymbol{\kappa}_h)}{c_1\|w_h\|_{sd}}\leq \frac{a_h({\boldsymbol w}_h, \boldsymbol{\kappa}_h)}{\left\|{\boldsymbol \kappa}_h\right\|_{s}}\leq \sup_{{\boldsymbol v}_h\in {\boldsymbol V}_h}\frac{a_h({\boldsymbol w}_h, {\boldsymbol v}_h)}{\left\|{\boldsymbol v}_h\right\|_{s}}.
$$

By combining the above with eq. [\(4.2b\)](#page-7-4),

$$
(1+(c_1^{-1}c_2+1)c_T)\sup_{{\bf v}_h\in{\bf V}_h}\frac{a_h({\bf w}_h,{\bf v}_h)}{\|{\bf v}_h\|_s}\geq c_1^{-1}\|w_h\|_{sd}+\|{\bf w}_h\|_s\geq c\|{\bf w}_h\|_{ss},
$$

proving eq. (4.1) .

 \Box

5 Error analysis

The following projection estimates for Π_h and $\Pi_h^{\mathcal{F}}$ were shown to hold for any $u|_{\mathcal{K}} \in H^{(p_t+1,p_s+1)}(\mathcal{K}),$ $K \in \mathcal{T}_h$, see Kirk et al. [\[30,](#page-30-10) Lemma 5.2], Sudirham et al. [\[42,](#page-31-2) Lemma 6.1 and Remark 6.2], and Georgoulis [\[22,](#page-30-7) Lemmas 3.13 and 3.17]:

$$
||u - \Pi_h u||_{\mathcal{K}}^2 \le c \left(h_K^{2p_s + 2} + \delta t \kappa^{2p_t + 2} \right) ||u||_{H^{(p_t + 1, p_s + 1)}(\mathcal{K})}^2,
$$
\n(5.1a)

$$
\|\overline{\nabla} (u - \Pi_h u)\|_{\mathcal{K}}^2 \le c \left(h_K^{2p_s} + \delta t_{\mathcal{K}}^{2p_t + 2} \right) \|u\|_{H^{(p_t + 1, p_s + 1)}(\mathcal{K})}^2, \tag{5.1b}
$$

$$
\|\partial_t (u - \Pi_h u)\|_{\mathcal{K}}^2 \le c \left(h_K^{2p_s} + \delta t_{\mathcal{K}}^{2p_t}\right) \|u\|_{H^{(p_t+1,p_s+1)}(\mathcal{K})}^2, \tag{5.1c}
$$

$$
\left\|\overline{\nabla}_{\bar{n}}\left(u-\Pi_{h}u\right)\right\|_{\mathcal{Q}_{\mathcal{K}}}^{2} \leq c\left(h_{K}^{2p_{s}-1}+h_{K}^{-1}\delta t_{K}^{2p_{t}+2}\right)\left\|u\right\|_{H\left(p_{t}+1,p_{s}+1\right)(K)}^{2},\tag{5.1d}
$$

$$
||u - \Pi_h u||_{\partial \mathcal{K}}^2 \le c \left(h_K^{2p_s + 1} + \delta t_K^{2p_t + 1} \right) ||u||_{H^{(p_t + 1, p_s + 1)}(\mathcal{K})}^2, \tag{5.1e}
$$

$$
\|u - \Pi_h^{\mathcal{F}} u\|_{\partial \mathcal{K}}^2 \le c \left(h_K^{2p_s + 1} + \delta t \mathcal{K}^{2p_t + 1} \right) \|u\|_{H^{(p_t + 1, p_s + 1)}(\mathcal{K})}^2.
$$
 (5.1f)

Let us define $h := \max_{\mathcal{K} \in \mathcal{T}_h} h_K$ and $\delta t := \max_{\mathcal{K} \in \mathcal{T}_h} \delta t_{\mathcal{K}}$. An immediate consequence of eq. [\(5.1\)](#page-18-1) is the following estimate.

Lemma 5.1. Let u, with $u|_K \in H^{(p_t+1,p_s+1)}(\mathcal{K})$ for all $\mathcal{K} \in \mathcal{T}_h$, and define $u := (u, u|_{\Gamma})$. Let $\mathbf{\Pi}_h \boldsymbol{u} = (\Pi_h u, \Pi_h^{\mathcal{F}} u)$. Then,

$$
\|\boldsymbol{u}-\boldsymbol{\Pi}_h\boldsymbol{u}\|^2_{ss}\leq c\left[h^{2p_s}(h+\varepsilon+\tilde{\varepsilon}\delta t)+\delta t^{2p_t}(\delta t+\varepsilon h^{-1}\delta t)\right],
$$

where the constant c depends on $\sum_{\mathcal{K}\in\mathcal{T}_h} ||u||_{H^{(p_t+1,p_s+1)}(\mathcal{K})}$.

Proof. By eq. $(5.1a)$,

$$
||u - \Pi_h u||_{\mathcal{K}}^2 \le c \left(h_K^{2p_s + 2} + \delta t_{\mathcal{K}}^{2p_t + 2} \right) ||u||_{H^{(p_t + 1, p_s + 1)}(\mathcal{K})}^2.
$$
 (5.2)

Next, eq. [\(5.1b\)](#page-18-3) gives us:

$$
\varepsilon \left\| \overline{\nabla} \left(u - \Pi_h u \right) \right\|_{\mathcal{K}}^2 \leq c \varepsilon \left(h_K^{2p_s} + \delta t_K^{2p_t + 2} \right) \left\| u \right\|_{H^{(p_t + 1, p_s + 1)}(\mathcal{K})}^2. \tag{5.3}
$$

For the advective facet terms, we use eqs. $(5.1e)$ and $(5.1f)$ and the triangle inequality:

$$
\|\beta_s - \frac{1}{2}\beta \cdot n\|^{1/2} \left((u - \Pi_h u) - (\gamma(u) - \Pi_h^{\mathcal{F}} \gamma(u)) \right)\|_{\partial \mathcal{K}}^2 + \|\mathbf{1}\|_2^2 \beta \cdot n\|^{1/2} (\gamma(u) - \Pi_h^{\mathcal{F}} \gamma(u))\|_{\partial \mathcal{K} \cap \partial \mathcal{E}_N}^2
$$

\$\leq c \left(h_K^{2p_s + 1} + \delta t_K^{2p_t + 1} \right) \|u\|_{H^{(p_t + 1, p_s + 1)}(\mathcal{K})}^2. (5.4)

Similarly, for the diffusive facet term, we again apply the triangle inequality and eqs. [\(5.1e\)](#page-18-4) and [\(5.1f\)](#page-18-5):

$$
\varepsilon h_K^{-1} \left\| (u - \Pi_h u) - (\gamma(u) - \Pi_h^{\mathcal{F}} \gamma(u)) \right\|_{\mathcal{Q}_K}^2 \leq c \varepsilon \left(h_K^{2p_s} + h_K^{-1} \delta t_K^{2p_t+1} \right) \left\| u \right\|_{H^{(p_t+1,p_s+1)}(K)}^2. \tag{5.5}
$$

For the streamline derivative term, we use eqs. [\(5.1b\)](#page-18-3) and [\(5.1c\)](#page-18-6) and that $\frac{\delta t_{\mathcal{K}}h_{K}^{2}}{\delta t_{\mathcal{K}}+h_{K}} \leq \delta t_{\mathcal{K}}h_{K}$:

$$
\frac{\delta t_{\mathcal{K}} h_K^2}{\delta t_{\mathcal{K}} + h_K} \left\| \Pi_h \left(\beta \cdot \nabla \left(u - \Pi_h u \right) \right) \right\|_{\mathcal{K}}^2 \leq c \delta t_{\mathcal{K}} h_K \left(\left\| \overline{\nabla} \left(u - \Pi_h u \right) \right\|_{\mathcal{K}}^2 + \left\| \partial_t \left(u - \Pi_h u \right) \right\|_{\mathcal{K}}^2 \right) \right\|_{\mathcal{K}}^2 \leq c \delta t_{\mathcal{K}} h_K \left(h_K^{2p_s} + \delta t_{\mathcal{K}}^{2p_t} \right) \left\| u \right\|_{H^{(p_t+1,p_s+1)}(\mathcal{K})}^2.
$$
\n(5.6)

Finally, for the time-derivative term, using eq. [\(5.1c\)](#page-18-6),

$$
\tau_{\varepsilon} \left\| \partial_{t} \left(u - \Pi_{h} u \right) \right\|_{\mathcal{K}}^{2} \leq \begin{cases} c \left(h_{K}^{2p_{s}} \delta t_{\mathcal{K}} + \delta t_{\mathcal{K}}^{2p_{t}+1} \right) \left\| u \right\|_{H^{(p_{t}+1,p_{s}+1)}(\mathcal{K})}^{2} & \text{if } \mathcal{K} \in \mathcal{T}_{h}^{d}, \\ c \varepsilon^{1/2} \left(h_{K}^{2p_{s}} \delta t_{\mathcal{K}} + \delta t_{\mathcal{K}}^{2p_{t}+1} \right) \left\| u \right\|_{H^{(p_{t}+1,p_{s}+1)}(\mathcal{K})}^{2} & \text{if } \mathcal{K} \in \mathcal{T}_{h}^{x}, \\ c \varepsilon \left(h_{K}^{2p_{s}} \delta t_{\mathcal{K}} + \delta t_{\mathcal{K}}^{2p_{t}+1} \right) \left\| u \right\|_{H^{(p_{t}+1,p_{s}+1)}(\mathcal{K})}^{2} & \text{if } \mathcal{K} \in \mathcal{T}_{h}^{c}. \end{cases} \tag{5.7}
$$
\n
$$
\leq c \tilde{\varepsilon} \left(h_{K}^{2p_{s}} \delta t_{\mathcal{K}} + \delta t_{\mathcal{K}}^{2p_{t}+1} \right) \left\| u \right\|_{H^{(p_{t}+1,p_{s}+1)}(\mathcal{K})}^{2}.
$$

 \Box

The result follows after combining eqs. [\(5.2\)](#page-18-7) to [\(5.7\)](#page-19-0) and summing over all $\mathcal{K} \in \mathcal{T}_h$.

The following lemma will be used to prove the global error estimate of theorem [5.3.](#page-20-0)

Lemma 5.2. Let u, with $u|_K \in H^{(p_t+1,p_s+1)}(\mathcal{K})$ for all $\mathcal{K} \in \mathcal{T}_h$, solve eq. [\(2.1\)](#page-2-1) and define $u := (u, \lambda)$ with $\lambda = u|_{\Gamma}$. Let $\mathbf{\Pi}_h \mathbf{u} = (\Pi_h u, \Pi_h^T u)$ and let $\mathbf{u}_h = (u_h, \lambda_h) \in \mathbf{V}_h$ be the solution to eq. [\(3.10\)](#page-6-0). The following holds:

$$
|a_h(\boldsymbol{u}-\Pi_h\boldsymbol{u},\boldsymbol{v}_h)|
$$

$$
\leq \left[c\|\boldsymbol{u}-\Pi_h\boldsymbol{u}\|_{ss}+c\||\beta_s-\frac{1}{2}\beta\cdot n|^{1/2}(\boldsymbol{u}-\Pi_h\boldsymbol{u})\|_{\partial\mathcal{T}_h}+\left(\sum_{\mathcal{K}\in\mathcal{T}_h}\varepsilon h_K\|\overline{\nabla}_{\bar{n}}(\boldsymbol{u}-\Pi_h\boldsymbol{u})\|_{\mathcal{Q}_\mathcal{K}}^2\right)^{1/2}\right]\|\boldsymbol{v}_h\|_{s}.
$$

Proof. We start with the advective part of $a_h(\cdot, \cdot)$. Writing $\zeta^+ \beta \cdot n = (\beta \cdot n + |\beta \cdot n|)/2$ and using the triangle inequality,

$$
|a_{h,c}(\mathbf{u}-\Pi_h \mathbf{u}, \mathbf{v}_h)| \leq |(\beta (u-\Pi_h u), \nabla v_h)_{\mathcal{T}_h}| + |\langle \frac{1}{2} (\beta \cdot n + |\beta \cdot n|) (\lambda - \Pi_h^{\mathcal{F}} \lambda), \mu_h \rangle_{\partial \mathcal{E}_N}|
$$

+ |\langle (\beta \cdot n) (\lambda - \Pi_h^{\mathcal{F}} \lambda) + \beta_s [\mathbf{u} - \Pi_h \mathbf{u}], [\mathbf{v}_h] \rangle_{\partial \mathcal{T}_h}| =: I_1 + I_2 + I_3.

To bound I_1 , we follow the proof of Ayuso and Marini [\[2,](#page-28-0) Theorem 5.1] by noting that if $\beta_0 = (1, \bar{\beta}_0)$ then $(\beta_0 (u - \Pi_h u), \nabla v_h)_{\mathcal{T}_h} = 0$ and $((\beta - \beta_0) (u - \Pi_h u), \nabla v_h)_{\mathcal{T}_h} = ((\overline{\beta} - \overline{\beta}_0) (u - \Pi_h u), \overline{\nabla} v_h)_{\mathcal{T}_h}$. Then, using the Cauchy–Schwarz inequality, eq. (4.4) , eq. $(3.5b)$, and Hölder's inequality for sums, we obtain

$$
I_1 \leq \sum_{\mathcal{K}\in\mathcal{T}_h} c\left\|u-\Pi_h u\right\|_{\mathcal{K}}\left\|v_h\right\|_{\mathcal{K}}.
$$

Using the Cauchy–Schwarz inequality, we bound I_2 as:

$$
I_2 \leq c \left\| \left| \frac{1}{2} \beta \cdot n \right|^{1/2} \left(\lambda - \Pi_h^{\mathcal{F}} \lambda \right) \right\|_{\partial \mathcal{E}_N} \left\| \left| \frac{1}{2} \beta \cdot n \right|^{1/2} \mu_h \right\|_{\partial \mathcal{E}_N}.
$$

With $\beta \cdot n \leq \sup |\beta \cdot n| \leq 2 (\sup |\beta \cdot n| - \frac{1}{2}\beta \cdot n)$, for all $F \in \partial \mathcal{T}_h$, and the Cauchy–Schwarz inequality, we bound I_3 as:

$$
I_3 \leq c \left| \left\langle \left(\sup \left| \beta \cdot n \right| - \frac{1}{2} \beta \cdot n \right) \left(\lambda - \Pi_h^{\mathcal{F}} \lambda + \left[\boldsymbol{u} - \Pi_h \boldsymbol{u} \right] \right), \left[v_h \right] \right\rangle_{\partial \mathcal{T}_h} \right|
$$

\$\leq c \sum_{\mathcal{K} \in \mathcal{T}_h} \left\| \left| \beta_s - \frac{1}{2} \beta \cdot n \right|^{1/2} \left(u - \Pi_h u \right) \right\|_{\partial \mathcal{K}} \left\| \left| \beta_s - \frac{1}{2} \beta \cdot n \right|^{1/2} \left[v_h \right] \right\|_{\partial \mathcal{K}}.\$

Collecting the bounds for I_1 , I_2 , and I_3 , and using Hölder's inequality for sums,

$$
|a_{h,c}(\boldsymbol{u}-\boldsymbol{\Pi}_h\boldsymbol{u},\boldsymbol{v}_h)|\leq (c||\boldsymbol{u}-\boldsymbol{\Pi}_h\boldsymbol{u}||_{ss}+c||\beta_s-\tfrac{1}{2}\beta\cdot n|^{1/2}(u-\Pi_h\boldsymbol{u})||_{\partial\mathcal{T}_h})||\boldsymbol{v}_h||_s.
$$
(5.8)

We now proceed with the diffusive part of $a_h(\cdot, \cdot)$. By the triangle inequality,

$$
|a_{h,d}(\boldsymbol{u}-\boldsymbol{\Pi}_h \boldsymbol{u}, \boldsymbol{v}_h)| \leq |(\varepsilon \overline{\nabla} (u - \Pi_h u), \overline{\nabla} v_h)_{\mathcal{T}_h}| + |\langle \varepsilon \alpha h_K^{-1} [\boldsymbol{u} - \Pi_h \boldsymbol{u}], [\boldsymbol{v}_h] \rangle_{\mathcal{Q}_h}|
$$

+ $|\langle \varepsilon [\boldsymbol{u} - \Pi_h \boldsymbol{u}], \overline{\nabla}_{\bar{n}} v_h \rangle_{\mathcal{Q}_h}| + |\langle \varepsilon \overline{\nabla}_{\bar{n}} (u - \Pi_h u), [\boldsymbol{v}_h] \rangle_{\mathcal{Q}_h}|.$

By applying the Cauchy–Schwarz inequality, the first two terms on the right-hand side can be bounded by $(1+\alpha)$ $\|\|\bm{u} - \bm{\Pi}_h\bm{u}\|$, $\|\|\bm{v}_h\|$, For the last two terms, by the Cauchy–Schwarz inequality and eq. $(3.5c)$,

$$
\begin{split} \|\langle \varepsilon\left[\boldsymbol{u}-\boldsymbol{\Pi}_h\boldsymbol{u}\right], \overline{\nabla}_{\bar{n}}v_h \rangle_{\mathcal{Q}_h} |+|\langle \varepsilon \overline{\nabla}_{\bar{n}}\left(u-\Pi_h u\right),[\boldsymbol{v}_h] \rangle_{\mathcal{Q}_h} | \\ \leq & c \sum_{\mathcal{K} \in \mathcal{T}_h} \varepsilon^{1/2} h_K^{-1/2} \left\| \left[\boldsymbol{u}-\boldsymbol{\Pi}_h \boldsymbol{u}\right] \right\|_{\mathcal{Q}_\mathcal{K}} \varepsilon^{1/2} \left\| \overline{\nabla} v_h \right\|_{\mathcal{K}} \\ & + \sum_{\mathcal{K} \in \mathcal{T}_h} \varepsilon^{1/2} h_K^{1/2} \left\| \overline{\nabla}_{\bar{n}}\left(u-\Pi_h u\right) \right\|_{\mathcal{Q}_\mathcal{K}} \varepsilon^{1/2} h_K^{-1/2} \left\| \left[\boldsymbol{v}_h\right] \right\|_{\mathcal{Q}_\mathcal{K}} . \end{split}
$$

Therefore, using Hölder's inequality for sums,

$$
|a_{h,d}(\boldsymbol{u}-\boldsymbol{\Pi}_h \boldsymbol{u}, \boldsymbol{v}_h)| \leq (c \|\boldsymbol{u}-\boldsymbol{\Pi}_h \boldsymbol{u}\|_{ss} + \left(\sum_{\mathcal{K}\in\mathcal{T}_h} \varepsilon h_K \|\overline{\nabla}_{\bar{n}} (u-\Pi_h u)\|_{\mathcal{Q}_\mathcal{K}}^2\right)^{1/2}) \|\boldsymbol{v}_h\|_{s}.
$$
 (5.9)

The result follows by combining eq. [\(5.8\)](#page-19-1) and eq. [\(5.9\)](#page-20-1).

Theorem 5.3 (Global error estimate). Let **u** and u_h be as in lemma [5.2.](#page-19-2) Then

$$
\|\boldsymbol{u}-\boldsymbol{u}_h\|_{ss}^2 \leq c_T \left[h^{2p_s}(h+\varepsilon+\tilde{\varepsilon}\delta t) + \delta t^{2p_t+1}(1+\varepsilon h^{-1})\right]
$$

where c_T depends on $\sum_{\mathcal{K} \in \mathcal{T}_h} ||u||_{H^{(p_t+1,p_s+1)}(\mathcal{K})}$.

Proof. We start by noting that Galerkin orthogonality was shown in Kirk et al. [\[30\]](#page-30-10):

$$
a_h(\mathbf{u}-\mathbf{u}_h,\mathbf{v}_h)=0 \quad \forall \mathbf{v}_h := (v_h,\mu_h) \in \mathbf{V}_h. \tag{5.10}
$$

.

By a triangle inequality, theorem [4.1,](#page-7-2) and eq. [\(5.10\)](#page-20-2) we find:

$$
\left\|\!\left\|\bm{u}_h-\bm{u}\right\|\!\right|_{ss} \leq \left\|\!\left\|\bm{u}-\bm{\Pi}_h\bm{u}\right\|\!\right|_{ss}+c_T\sup\limits_{\bm{v}_h\in\bm{V}_h}\frac{a_h(\bm{u}-\bm{\Pi}_h\bm{u},\bm{v}_h)}{\left\|\!\left\|\bm{v}_h\right\|\!\right|_s}.
$$

Using lemma [5.2,](#page-19-2)

$$
\|\boldsymbol{u}_{h} - \boldsymbol{u}\|_{ss} \leq c_{T} \|\boldsymbol{u} - \boldsymbol{\Pi}_{h} \boldsymbol{u}\|_{ss} + c_{T} \|\beta_{s} - \frac{1}{2}\beta \cdot n\|^{1/2} (u - \Pi_{h} u)\|_{\partial \mathcal{T}_{h}} + c_{T} \left(\sum_{\mathcal{K} \in \mathcal{T}_{h}} \varepsilon h_{K} \|\overline{\nabla}_{\bar{n}} (u - \Pi_{h} u)\|_{\mathcal{Q}_{\mathcal{K}}}^{2} \right)^{1/2}.
$$
 (5.11)

The second term on the right-hand side of eq. (5.11) is bounded using eq. $(5.1e)$:

$$
\sum_{\mathcal{K}\in\mathcal{T}_h} \left\| \left| \beta_s - \frac{1}{2} \beta \cdot n \right|^{1/2} \left(u - \Pi_h u \right) \right\|_{\partial\mathcal{K}_h}^2 \le c \left(h^{2p_s + 1} + \delta t^{2p_t + 1} \right) \sum_{\mathcal{K}\in\mathcal{T}_h} \left\| u \right\|_{H^{(p_t + 1, p_s + 1)}(\mathcal{K})}^2. \tag{5.12}
$$

The last term on the right-hand side of eq. [\(5.11\)](#page-20-3) is bounded using eq. [\(5.1d\)](#page-18-8):

$$
\sum_{\mathcal{K}\in\mathcal{T}_h} \varepsilon h_K \left\|\overline{\nabla}_{\bar{n}} \left(u - \Pi_h u\right)\right\|_{\mathcal{Q}_\mathcal{K}}^2 \leq c\varepsilon \left(h^{2p_s} + \delta t^{2p_t+2}\right) \sum_{\mathcal{K}\in\mathcal{T}_h} \|u\|_{H^{(p_t+1,p_s+1)}(\mathcal{K})}^2. \tag{5.13}
$$

The result follows after combining eqs. (5.11) to (5.13) and lemma [5.1.](#page-18-9)

Remark 1. The error estimate of theorem [5.3](#page-20-0) shows that if $\varepsilon < \delta t = h$ then $||u - u_h||_{ss}$ $\mathcal{O}(h^{p_s+1/2} + \delta t^{p_t+1/2}),$ while if $\delta t = h < \varepsilon$ then $\|\boldsymbol{u} - \boldsymbol{u}_h\|_{ss} = \mathcal{O}(h^{p_s} + \delta t^{p_t}).$

 \Box

 \Box

Figure 3: The spatial mesh and the ring of elements with an extra level of refinement deform over time. The solution shown is for $\varepsilon = 10^{-8}$. Plots correspond to time levels $t = 0.2, 0.5, 0.8$ from left to right.

6 Numerical example

We consider the solution of a two-dimensional rotating Gaussian pulse on a deforming domain [\[38\]](#page-31-4) to demonstrate the convergence properties of the space-time HDG method predicted by theorem [5.3.](#page-20-0) In eq. [\(2.1\)](#page-2-1) we set $\beta = (1, -4x_2, 4x_1)^T$ and $f = 0$. Defining $\tilde{x}_1 := x_1 \cos(4t) + x_2 \sin(4t)$ and $\tilde{x}_2 := x_1 \sin(4t) + x_2 \sin(4t)$ and $\tilde{x}_3 := x_1 \sin(4t) + x_2 \sin(4t)$. $\tilde{x}_2 := -x_1 \sin(4t) + x_2 \cos(4t)$, the exact solution to this problem is given by

$$
u(t, x_1, x_2) = \frac{\sigma^2}{\sigma^2 + 2\varepsilon t} \exp\left(-\frac{(\tilde{x}_1 - x_{1c})^2 + (\tilde{x}_2 - x_{2c})^2}{2\sigma^2 + 4\varepsilon t}\right),
$$

with initial and boundary conditions set appropriately. We choose $\sigma = 0.1$ and (x_{1c}, x_{2c}) $(-0.2, 0.1)$. The deforming domain $\Omega(t)$ is obtained by transforming a uniform mesh, with coordinates $(x_1^u, x_2^u) \in (-0.5, 0.5)^2$, to

$$
x_i = x_i^u + A(\frac{1}{2} - x_i^u)\sin(2\pi(\frac{1}{2} - x_i^* + t)), \quad i = 1, 2,
$$
\n(6.1)

where $(x_1^*, x_2^*) = (x_2, x_1)$ and $A = 0.1$. We consider this problem for $t \in [0, 1]$.

The space-time HDG method eq. [\(3.10\)](#page-6-0) is implemented using the finite element library deal.II [\[1\]](#page-28-1) on unstructured hexahedral space-time meshes with p4est [\[6\]](#page-28-2) to obtain distributed mesh information. We use PETSc $[5, 4, 3]$ $[5, 4, 3]$ $[5, 4, 3]$ $[5, 4, 3]$ to solve the linear systems arising at each time step (GMRES preconditioned by classical algebraic multigrid from BoomerAMG [\[24\]](#page-30-12) with an absolute solver tolerance of 10^{-12}). To create our coarsest mesh, we start with an initial mesh with elements of size $h \approx \delta t = 10^{-1}$. Space-time elements in a ring prescribed by $|((x_1^c)^2 + (x_2^c)^2)^{1/2} - 0.2| < 0.1$, where (x_1^c, x_2^c) is the spatial coordinate of the center of a space-time element, are then uniformly refined once and are of size $h \approx \delta t = 0.05$. See fig. [3](#page-21-1) for plots of the solution and spatial mesh at different time levels. The reason to consider two sets of elements is to verify that the analysis of previous sections hold on 1-irregular space-time meshes. Finer meshes are obtained by uniformly refining our coarsest mesh. In our implementation we furthermore choose the penalty parameter $\alpha = 8p_s^2$ [see, for example [39\]](#page-31-8). We show the rates of convergence for different polynomial degrees when the error is measured in $\|\cdot\|_{ss}$ in tables [1](#page-22-1) and [2](#page-22-2) for $\varepsilon = 10^{-2}$ and $\varepsilon = 10^{-8}$, respectively.

In the third row of table [1](#page-22-1) we have that $h \approx \delta t = 1.25 \times 10^{-2}$ inside the refined ring while elsewhere $h \approx \delta t = 2.5 \times 10^{-2}$. Therefore, for the first three rows in table [1,](#page-22-1) $h \approx \delta t \geq \varepsilon = 10^{-2}$ and we observe a rate of convergence of approximately $p + \frac{1}{2}$. In the following three rows we observe a drop in the rate of convergence to approximately p . This happens in two stages since there are two sets of elements in our mesh, see fig. [3.](#page-21-1) In the first stage (the fourth row of table [1\)](#page-22-1), elements in the refined ring are such that $h \approx \delta t = 6.25 \times 10^{-3} < \varepsilon$, but elsewhere $h \approx \delta t = 1.25 \times 10^{-2} > \varepsilon$. In the next stage (fifth row of table [1\)](#page-22-1), all elements satisfy $h \approx \delta t < \varepsilon$, resulting in a rate of

Table 1: The solution errors measured in $\|\cdot\|_{ss}$ and corresponding rates of convergence when using polynomial approximation $p = 1, 2, 3$ for the case $\varepsilon = 10^{-2}$.

	Cells per slab Number of slabs	$p=1$	Rates	$p=2$	Rates	$p=3$	Rates
296	10	$4.7e-2$		7.8e-3	\overline{a}	$1.3e-3$	
1100	20	$1.8e-2$	1.4	$1.6e-3$	2.4	$1.2e-4$	3.6
4372	40	7.7e-3	$1.3\,$	$3.2e-4$	2.3	1.7e-5	3.4
17572	80	3.7e-3	1.1	$7.3e-5$	2.1	$1.4e-6$	3.2
70540	160	$2.0e-3$	0.9	$2.3e-5$	1.7	$2.4e-7$	2.4
282580	320	$9.0e-4$	$1.1\,$	$4.9e-6$	2.2	$2.5e-8$	3.3

Table 2: The solution errors measured in $\|\cdot\|_{ss}$ and corresponding rates of convergence when using polynomial approximation $p = 1, 2, 3$ for the case $\varepsilon = 10^{-8}$.

convergence of p after the fifth row. In table [2](#page-22-2) we observe that $h \approx \delta t > \varepsilon = 10^{-8}$ for all cycles and the error converges at a rate of approximately $p + \frac{1}{2}$ $p + \frac{1}{2}$ $p + \frac{1}{2}$ $p + \frac{1}{2}$ $p + \frac{1}{2}$. These observations from tables 1 and 2 are in agreement with remark [1.](#page-20-5)

7 Conclusions

In this paper we analyzed the space-time HDG method for the time-dependent advection-diffusion equation on deforming domains. We proved stability of the discretization in the advection-dominated regime using a time-dependent weighted test function, and derived a priori error estimates that show a drop of 1/2 in the rate of convergence in a mesh dependent norm when transitioning from a mesh size larger than the diffusion parameter to a mesh size smaller than the diffusion parameter. A numerical example supports our theoretical analysis.

A Proof of eq. (4.7)

The following results from Sudirham [\[41,](#page-31-9) Lemma B.7] will be useful in what follows: For $\tilde{u} \in$ $H^{(k_t,k_s)}(\widetilde{\mathcal{K}}), 0 \leq \alpha_t \leq k_t, \alpha_s = (\alpha_1,\ldots,\alpha_d)$ with $|\alpha_s| \leq k_s$, and $\widetilde{F}_{\mathcal{Q}} \subset \mathcal{Q}_{\widetilde{\mathcal{K}}}$:

$$
\|\partial_{\widehat{t}}^{\alpha_t} \partial_{\widehat{x}}^{\alpha_s} \widehat{u}\|_{\widehat{\mathcal{K}}}^2 = \left(\frac{1}{2}\right)^{2(\alpha_t + |\alpha_s|) - d - 1} \delta t_{\mathcal{K}}^{2\alpha_t - 1} h_K^{2|\alpha_s| - d} \|\partial_{\widehat{t}}^{\alpha_t} \partial_{\widehat{x}}^{\alpha_s} \widetilde{u}\|_{\widetilde{\mathcal{K}}}^2, \tag{A.1a}
$$

$$
\|\partial_{\hat{t}}^{\alpha_t} \partial_{\hat{x}}^{\alpha_s} \hat{u}\|_{\widehat{F}_{\mathcal{Q}}}^2 = \left(\frac{1}{2}\right)^{2(\alpha_t + |\alpha_s|) - d} \delta t_{\mathcal{K}}^{2\alpha_t - 1} h_K^{2|\alpha_s| - d + 1} \|\partial_{\hat{t}}^{\alpha_t} \partial_{\hat{x}}^{\alpha_s} \tilde{u}\|_{\widetilde{F}_{\mathcal{Q}}}^2.
$$
 (A.1b)

To begin, we note that the projection operator on $\widehat{\mathcal{K}}$ and $\widetilde{\mathcal{K}}$ are related to the projection operator on K as follows [see [22,](#page-30-7) Definition 3.12]:

$$
\widetilde{\Pi}_h \widetilde{u} = (\Pi_h (\widetilde{u} \circ \phi_{\mathcal{K}}^{-1})) \circ \phi_{\mathcal{K}} \quad \forall \widetilde{u} \in L^2(\widetilde{\mathcal{K}}), \qquad \widehat{\Pi}\widehat{u} = (\widetilde{\Pi}_h (\widehat{u} \circ G_{\mathcal{K}}^{-1})) \circ G_{\mathcal{K}} \quad \forall \widehat{u} \in L^2(\widehat{\mathcal{K}}).
$$

Similarly, on any $F \in \partial \mathcal{K}$, we have the following relations:

$$
\widetilde{\Pi}_h^{\mathcal{F}} \widetilde{u} = (\Pi_h^{\mathcal{F}} (\widetilde{u} \circ \phi_F^{-1})) \circ \phi_F \ \forall \widetilde{u} \in L^2(\widetilde{F}), \quad \widehat{\Pi}^{\mathcal{F}} \widehat{u} = (\widetilde{\Pi}_h^{\mathcal{F}} (\widehat{u} \circ G_F^{-1})) \circ G_F \ \forall \widehat{u} \in L^2(\widehat{F}),
$$

where G_F and ϕ_F are the restrictions of G_K and ϕ_K on F, respectively.

To show eq. [\(4.7a\)](#page-8-10) we first note that the following projection estimate holds on $\hat{\mathcal{K}}$ for any $1 \leq i \leq d$ [see [22,](#page-30-7) lemma 3.7, eq. (3.12)]:

$$
\|\partial_{\widehat{x}_i} \left(\widehat{u} - \widehat{\Pi} \widehat{u} \right)\|_{\widehat{\mathcal{K}}} \leq c \|\partial_{\widehat{x}_i} \widehat{u}\|_{\widehat{\mathcal{K}}}.
$$
\n(A.2)

By the chain rule $\partial_{\tilde{x}_i} (\tilde{u} - \tilde{\Pi}_h \tilde{u}) = 2h_R^{-1} \partial_{\tilde{x}_i} ((\tilde{u} - \tilde{\Pi}_h \tilde{u}) \circ G_K) \circ G_K^{-1}$, using that $|\det G_K| = \delta t_K h_K^d 2^{-d-1}$, and using eqs. $(A.1a)$ and $(A.2)$, we find

$$
\left\|\partial_{\widetilde{x}_i} \left(\widetilde{u} - \widetilde{\Pi}_h \widetilde{u}\right)\right\|_{\widetilde{\mathcal{K}}}^2 \le c \delta t_{\mathcal{K}} h_K^{d-2} \left\|\partial_{\widehat{x}_i} \widehat{u}\right\|^2 \le c \left\|\partial_{\widetilde{x}_i} \widetilde{u}\right\|_{\widehat{\mathcal{K}}}^2.
$$
\n(A.3)

To obtain the result on the physical element, consider first that by the chain rule,

$$
\partial_{x_i}(u - \Pi_h u) = \sum_{1 \leq j \leq d} (\partial_{\widetilde{x}_j} ((u - \Pi_h u) \circ \phi_K) \circ \phi_K^{-1}) ((-1)^{i+j} (\det J_{\phi_K})^{-1} \det J_{\phi_K \setminus ij}),
$$

where we used that \tilde{t} only depends on t in ϕ^{-1}_k and that $\frac{\partial \tilde{x}_j}{\partial x_i} = (-1)^{i+j} (\det J_{\phi_K})^{-1} \det J_{\phi_K \setminus ij}$. By assumptions eq. (3.1) and eq. (3.2) , and using eq. $(A.3)$, we therefore find that:

$$
\|\partial_{x_i} (u - \Pi_h u)\|_{\mathcal{K}}^2 \leq c \sum_{1 \leq j \leq d} \|\partial_{\widetilde{x}_i} (\widetilde{u} - \widetilde{\Pi}_h \widetilde{u})\|_{\widetilde{\mathcal{K}}}^2 \leq c \sum_{1 \leq j \leq d} \|\partial_{\widetilde{x}_j} \widetilde{u}\|_{\widetilde{\mathcal{K}}}^2 \leq c \|\overline{\nabla}u\|_{\mathcal{K}}^2,
$$

where the last step reverses the scaling arguments proving eq. $(4.7a)$. The proof for eq. $(4.7b)$ is similar and therefore omitted.

For eq. [\(4.7c\)](#page-8-6), we consider a d-dimensional hypersurface $F_Q \in \mathcal{Q}_K$. We first map F_Q to the reference domain. For this, let $\phi_{F_Q} \circ G_{F_Q}(F_Q) = F_Q$, i.e., the transformation of a face from the reference domain to the physical domain. We then observe that one of the spatial coordinates, which is denoted by \hat{x}_j without loss of generality, of $\hat{F}_{\mathcal{Q}}$ is fixed. We further consider a decomposition of $\widehat{\Pi} = \widehat{\pi}_{\widehat{t}} \Pi_1 \leq i \leq d \widehat{\pi}_{\widehat{x}_i}$ where $\widehat{\pi}_{\widehat{t}}$ and $\widehat{\pi}_{\widehat{x}_i}$ are the one-dimensional L^2 -projection operators applied in the time direction and in the spatial direction \hat{x}_i , respectively. Similarly, $(\widehat{\Pi}^{\mathcal{F}})|_{\widehat{F}_{\mathcal{Q}}} = \widehat{\pi}_{\widehat{t}} \Pi_{1 \leq i \leq d, i \neq j} \widehat{\pi}_{\widehat{x}_i}$. By Georgoulis [\[22,](#page-30-7) Definitions 3.1, 3.6], we have:

$$
\|\widehat{\Pi}\widehat{u} - \widehat{\Pi}^{\mathcal{F}}\widehat{u}\|_{\widehat{F}_{\mathcal{Q}}} = \|\widehat{\pi}_{\widehat{t}}\Pi_{1 \leq i \leq d, i \neq j}\widehat{\pi}_{\widehat{x}_i}\left(\widehat{u} - \widehat{\pi}_{\widehat{x}_j}\widehat{u}\right)\|_{\widehat{F}_{\mathcal{Q}}} \leq c\|\widehat{u} - \widehat{\pi}_{\widehat{x}_j}\widehat{u}\|_{\widehat{F}_{\mathcal{Q}}} \leq c\|\partial_{\widehat{x}_j}\widehat{u}\|_{\widehat{K}},\tag{A.4}
$$

where the equality is by commutativity of $\hat{\pi}_{\hat{x}_i}$ and $\hat{\pi}_{\hat{x}_j}$ $(i \neq j)$ and the last two inequalities are due
to the boundedness of any composition of projections $\hat{\pi}$ and Coergoulis [22, Lemma 2.2]. North to the boundedness of any composition of projections $\hat{\pi}$ and Georgoulis [\[22,](#page-30-7) Lemma 3.3]. Next, using eq. $(A.1b)$, eq. $(A.4)$, and eq. $(A.1a)$,

$$
\|\widetilde{\Pi}_h\widetilde{u}-\widetilde{\Pi}_h^{\mathcal{F}}\widetilde{u}\|^2_{\widetilde{F}_{\mathcal{Q}_i}}\leq ch_K \|\partial_{\widetilde{x}_j}\widetilde{u}\|^2_{\widetilde{\mathcal{K}}}.
$$

Therefore, also using eq. [\(3.3\)](#page-3-1) and eq. [\(3.4\)](#page-3-2),

$$
\left\|\Pi_h u - \Pi_h^{\mathcal{F}} u\right\|_{F_{\mathcal{Q}}}^2 \leq c \left\|\widetilde{\Pi}_h \widetilde{u} - \widetilde{\Pi}_h^{\mathcal{F}} \widetilde{u}\right\|_{\widetilde{F}_{\mathcal{Q}}}^2 \leq ch_K \left\|\partial_{\widetilde{x}_j} \widetilde{u}\right\|_{\widetilde{\mathcal{K}}}^2 \leq ch_K \left\|\overline{\nabla}u\right\|_{\mathcal{K}}^2,
$$

where we reverse the scaling arguments in the final inequality, proving eq. $(4.7c)$.

B Proof of lemma [4.3](#page-8-11)

To prove lemma [4.3](#page-8-11) we use the following result from Sudirham [\[41,](#page-31-9) Lemma B.6]. For any $K \in \mathcal{T}_h$ and $v \in H^1(\mathcal{K})$, we consider its affine domain $\widetilde{\mathcal{K}} := \phi_{\mathcal{K}}^{-1} \circ \mathcal{K}$ (with $\mathcal{Q}_{\widetilde{\mathcal{K}}} := \phi_{\mathcal{K}}^{-1} \circ \mathcal{Q}_{\mathcal{K}}$ and $\mathcal{R}_{\widetilde{\mathcal{K}}} := \phi_{\mathcal{K}}^{-1} \circ \mathcal{R}_{\mathcal{K}}$) and $\tilde{v} := v \circ \phi_{\mathcal{K}} \in H^1(\tilde{\mathcal{K}})$. The following trace inequalities hold:

$$
\|\widetilde{v}\|_{\mathcal{Q}_{\widetilde{\mathcal{K}}}}^2 \le c \left(h_K^{-1} \|\widetilde{v}\|_{\widetilde{\mathcal{K}}}^2 + \|\widetilde{v}\|_{\widetilde{\mathcal{K}}} \|\widetilde{\nabla} \widetilde{v}\|_{\widetilde{\mathcal{K}}}\right). \tag{B.1a}
$$

$$
\|\widetilde{v}\|_{\mathcal{R}_{\widetilde{\mathcal{K}}}}^2 \le c \left(\delta t_{\mathcal{K}}^{-1} \|\widetilde{v}\|_{\widetilde{\mathcal{K}}}^2 + \|\widetilde{v}\|_{\widetilde{\mathcal{K}}} \|\partial_{\widetilde{t}} \widetilde{v}\|_{\widetilde{\mathcal{K}}}\right). \tag{B.1b}
$$

We further observe that when φ is a function of the time variable only on K, so are $\tilde{\varphi}$ and $\hat{\varphi}$ on $\tilde{\mathcal{K}}$ and $\widehat{\mathcal{K}}$ respectively. Therefore, for $w_h \in V_h^{(p_t, p_s)}$ $h^{(pt, p_s)}$, and for $1 \leq i, j \leq d, j \neq i$,

$$
\partial_{\widetilde{x}_i}^{p_s+1}(\widetilde{\varphi w_h}) = \partial_{\widetilde{x}_i}^{p_s+1} \partial_{\widetilde{x}_j}(\widetilde{\varphi w_h}) = \partial_{\widetilde{x}_i}^{p_s+1} \partial_{\widetilde{t}}(\widetilde{\varphi w_h}) = 0.
$$
\n(B.2)

The equivalent derivatives above are also zero on the reference domain $\hat{\mathcal{K}}$. Furthermore,

$$
\|\partial_{\widetilde{\mathcal{t}}}^{p_t+1}(\widetilde{\varphi w_h})\|_{\widetilde{\mathcal{K}}} \le c\delta t_{\mathcal{K}}^{-p_t} \|\widetilde{w}_h\|_{\widetilde{\mathcal{K}}},\tag{B.3a}
$$

$$
\left\|\partial_{\tilde{t}}^{p_t+1}\partial_{\tilde{x}_i}\left(\widetilde{\varphi w_h}\right)\right\|_{\tilde{\mathcal{K}}}\leq c\delta t_{\mathcal{K}}^{-p_t}h_K^{-1}\left\|\widetilde{w}_h\right\|_{\tilde{\mathcal{K}}}.\tag{B.3b}
$$

Equation [\(B.3a\)](#page-24-1) can be shown using the general Leibniz rule, that $\partial_{\tilde{t}}^{p_t+1}\widetilde{w_h} = 0$, that $\|\partial_{\tilde{t}}^{j_t}\widetilde{\varphi}\|_{\widetilde{\mathcal{K}}} \leq e$ \tilde{t} $\omega_n - \upsilon$, end ω_t for all $1 \leq j_t \leq p_t+1$, that eq. $(3.5a)$ reduces to $\|\partial_t \tilde{v}_h\|_{\tilde{\mathcal{K}}} \leq c \delta t_{\mathcal{K}}^{-1} \|\tilde{v}_h\|_{\tilde{\mathcal{K}}}$ on the axiparallel element $\tilde{\mathcal{K}}$
Less 22. Corollary 3.54] and using that $\delta t \leq 1$. Similar argum [see [22,](#page-30-7) Corollary 3.54] and using that $\delta t_K < 1$. Similar arguments can be used to show eq. [\(B.3b\)](#page-24-2).

To prove eq. [\(4.8a\)](#page-8-2) we follow the proof of Ayuso and Marini [\[2,](#page-28-0) Lemma 4.2], and apply the projection estimates in Houston et al. [\[25,](#page-30-13) Lemma 3.4] when considered on the affine domain K , eq. [\(B.2\)](#page-24-3) and eq. [\(B.3a\)](#page-24-1),

$$
\|\widetilde{\varphi w_h} - \widetilde{\Pi}_h \left(\widetilde{\varphi w_h} \right) \|_{\widetilde{\mathcal{K}}} \le c \delta t \kappa^{p_t+1} \left\| \partial_t^{p_t+1} \left(\widetilde{\varphi w_h} \right) \right\|_{\widetilde{\mathcal{K}}} \le c \delta t \kappa \left\| \widetilde{w_h} \right\|_{\widetilde{\mathcal{K}}}.
$$
\n(B.4)

A scaling argument applied to eq. [\(B.4\)](#page-24-4) from \widetilde{K} to K yields eq. [\(4.8a\)](#page-8-2).

We next prove eq. $(4.8b)$. First note,

$$
\left\| \widetilde{\overline{\nabla}} \left(\widetilde{\varphi w_h} - \widetilde{\Pi}_h \left(\widetilde{\varphi w_h} \right) \right) \right\|_{\widetilde{\mathcal{K}}}^2 \le \sum_{1 \le i \le d} ch_K^{d-2} \delta t_{\mathcal{K}} \left\| \partial_{\widehat{x}_i} \left(\widetilde{\varphi w_h} - \widehat{\Pi}_h \left(\widetilde{\varphi w_h} \right) \right) \right\|_{\widehat{\mathcal{K}}}^2. \tag{B.5}
$$

Following similar steps as in the proof of Georgoulis [\[23,](#page-30-14) Lemma 7.5], the right-hand side of eq. [\(B.5\)](#page-24-5) can be bound further using the triangle inequality, commutativity of $\partial_{\hat{x}_i}$ with $\hat{\pi}_t \Pi_{1\leq j\leq d, j\neq i} \hat{\pi}_{x_j}$,
boundedness of $\hat{\pi} \Pi$, $\hat{x}_i \in \hat{\pi}_t$, the projection estimates in Houston at al. [25, Lomma 3 boundedness of $\hat{\pi}_t \Pi_{1 \leq j \leq d, j \neq i} \hat{\pi}_{x_j}$, the projection estimates in Houston et al. [\[25,](#page-30-13) Lemma 3.4] and
Controllis [23, Lemma 7.3], or (B.2), or (A.1a), and or (B.2b) Georgoulis [\[23,](#page-30-14) Lemma 7.3], eq. [\(B.2\)](#page-24-3), eq. [\(A.1a\)](#page-22-3), and eq. [\(B.3b\)](#page-24-2),

$$
\|\partial_{\widehat{x}_i} (\widehat{\varphi w_h} - \widehat{\Pi}_h (\widehat{\varphi w_h}))\|_{\widehat{\mathcal{K}}} \leq \|\partial_{\widehat{x}_i} \widehat{\varphi w_h} - \widehat{\Pi}_h (\partial_{\widehat{x}_i} \widehat{\varphi w_h})\|_{\widehat{\mathcal{K}}} + c \|\widehat{\pi}_{x_i} (\partial_{\widehat{x}_i} \widehat{\varphi w_h}) - \partial_{\widehat{x}_i} (\widehat{\pi}_{x_i} (\widehat{\varphi w_h}))\|_{\widehat{\mathcal{K}}}
$$

$$
\leq c \|\partial_{\widehat{t}}^{p_t+1} \partial_{\widehat{x}_i} \widehat{\varphi w_h}\|_{\widehat{\mathcal{K}}} \leq c h_K^{-d/2} \delta t_{\mathcal{K}}^{1/2} \|\widetilde{w}_h\|_{\widehat{\mathcal{K}}}.
$$
(B.6)

Combining the right-hand side of eq. $(B.6)$ with eq. $(B.5)$, we find:

$$
\|\widetilde{\overline{\nabla}}\left(\widetilde{\varphi w_h} - \widetilde{\Pi}_h\left(\widetilde{\varphi w_h}\right)\right)\|_{\widetilde{\mathcal{K}}} \le c\delta t_{\mathcal{K}}h_K^{-1} \left\|\widetilde{w_h}\right\|_{\widetilde{\mathcal{K}}}.
$$
\n(B.7)

A scaling argument applied to eq. [\(B.7\)](#page-24-7) from \widetilde{K} to K yields eq. [\(4.8b\)](#page-8-5). With similar steps it can be shown that

$$
\|\partial_{\widetilde{t}}(\widetilde{\varphi w_h} - \widetilde{\Pi}_h(\widetilde{\varphi w_h}))\|_{\widetilde{\mathcal{K}}} \le c \|\widetilde{w}_h\|_{\widetilde{\mathcal{K}}}.
$$
\n(B.8)

We next prove eq. $(4.8c)$. We start with a scaling argument to transform the integral on a \mathcal{Q} -face $F_{\mathcal{Q},m}$ from the affine domain to the reference domain. Note that, without loss of generality, subscript m denotes the index of the spatial coordinate for which $\hat{x}_m \equiv 1$. Using eq. [\(A.1b\)](#page-22-4) we find

$$
\|\widetilde{\nabla}(\widetilde{\varphi w_h} - \widetilde{\Pi}_h(\widetilde{\varphi w_h}))\|_{\widetilde{F}_{\mathcal{Q},m}}^2 \leq \sum_{1 \leq i \leq d} c \delta t \chi h_K^{d-3} \|\partial_{\widehat{x}_i}(\widehat{\varphi w_h} - \widehat{\Pi}_h(\widehat{\varphi w_h}))\|_{\widehat{F}_{\mathcal{Q},m}}^2. \tag{B.9}
$$

Consider now the right hand side term. Following Georgoulis [\[23,](#page-30-14) Lemma 7.9] we consider the cases $i = m$ and $i \neq m$ separately, starting with $i = m$. Using the commutativity of $\partial_{\hat{x}_m}$ with $\hat{\pi}_t \Pi_{1 \leq j \leq d, j \neq m} \hat{\pi}_{x_j}$, the triangle inequality, and Georgoulis [\[22,](#page-30-7) Lemma 3.47] [see also [23,](#page-30-14) Lemma π s) 7.8],

$$
\|\partial_{\widehat{x}_m} (\widehat{\varphi w}_h - \widehat{\Pi}_h (\widehat{\varphi w}_h))\|_{\widehat{F}_{\mathcal{Q},m}} \leq \|\partial_{\widehat{x}_m} (\widehat{\varphi w}_h) - \widehat{\pi}_{x_m} \partial_{\widehat{x}_m} (\widehat{\varphi w}_h)\|_{\widehat{F}_{\mathcal{Q},m}} + c \|\widehat{\pi}_{x_m} \partial_{\widehat{x}_m} (\widehat{\varphi w}_h) - \widehat{\pi}_t \Pi_{1 \leq j \leq d, j \neq m} \widehat{\pi}_{x_j} (\widehat{\pi}_{x_m} \partial_{\widehat{x}_m} (\widehat{\varphi w}_h))\|_{\widehat{K}} + c \|\widehat{\pi}_t \Pi_{1 \leq j \leq d, j \neq m} \widehat{\pi}_{x_j} (\widehat{\pi}_{x_m} \partial_{\widehat{x}_m} (\widehat{\varphi w}_h) - \partial_{\widehat{x}_m} \widehat{\pi}_{x_m} (\widehat{\varphi w}_h))\|_{\widehat{K}}.
$$
\n(B.10)

The first and third terms on the right-hand side of eq. [\(B.10\)](#page-25-0) vanish by Georgoulis [\[23,](#page-30-14) Lemmas 7.2 and 7.3] and eq. [\(B.2\)](#page-24-3). The second term on the right-hand side of eq. [\(B.10\)](#page-25-0) is bounded using the same argument as in the proof of Houston et al. [\[25,](#page-30-13) Lemma 3.4] by noting that $\hat{\pi}_{x_m}$ and $\hat{\pi}_t$ are one-dimensional L^2 -projections applied in the spatial direction \hat{x}_m and time direction, respectively, the commutativity of $\hat{\pi}_{x_m}$ with $\partial_t^{p_t+1}$ $\widehat{\hat{t}}^{p_t+1}$ and $\partial_{\widehat{x}_j}^{p_s+1}$ $x_{\hat{x}_j}^{p_s+1}$ $(j \neq m)$, the boundedness of $\hat{\pi}_{x_m}$, and eq. [\(B.2\)](#page-24-3),

$$
\|\hat{\pi}_{x_m}\partial_{\hat{x}_m}\left(\widehat{\varphi w_h}\right) - \hat{\pi}_t\Pi_{1\leq j\leq d,j\neq m}\hat{\pi}_{x_j}\left(\hat{\pi}_{x_m}\partial_{\hat{x}_m}\left(\widehat{\varphi w_h}\right)\right)\|_{\widehat{\mathcal{K}}}\leq c\left\|\partial_{\widehat{t}}^{p_t+1}\partial_{\hat{x}_m}\left(\widehat{\varphi w_h}\right)\right\|_{\widehat{\mathcal{K}}},\tag{B.11}
$$

so that eq. $(B.10)$ becomes:

$$
\|\partial_{\widehat{x}_m} \left(\widehat{\varphi w_h} - \widehat{\Pi}_h \left(\widehat{\varphi w_h} \right) \right)\|_{\widehat{F}_{\mathcal{Q},m}} \leq c \|\partial_{\widehat{t}}^{p_t+1} \partial_{\widehat{x}_m} \left(\widehat{\varphi w_h} \right)\|_{\widehat{\mathcal{K}}}.
$$
\n(B.12)

We now consider the right hand side of eq. [\(B.9\)](#page-25-1) with $i \neq m$. We have by a triangle inequality

$$
\|\partial_{\widehat{x}_{i}}\left(\widehat{\varphi w_{h}}-\widehat{\Pi}_{h}\left(\widehat{\varphi w_{h}}\right)\right)\|_{\widehat{F}_{\mathcal{Q},m}} \leq \|\partial_{\widehat{x}_{i}}\left(\widehat{\varphi w_{h}}\right)-\widehat{\pi}_{x_{m}}\partial_{\widehat{x}_{i}}\left(\widehat{\varphi w_{h}}\right)\|_{\widehat{F}_{\mathcal{Q},m}} + \|\left(I-\widehat{\pi}_{t}\Pi_{1\leq j\leq d,j\neq i,j\neq m}\widehat{\pi}_{x_{j}}\right)\widehat{\pi}_{x_{m}}\partial_{\widehat{x}_{i}}\left(\widehat{\varphi w_{h}}\right)\|_{\widehat{F}_{\mathcal{Q},m}} + \|\partial_{\widehat{x}_{i}}\left(\widehat{\varphi w_{h}}\right)-\partial_{\widehat{x}_{i}}\widehat{\pi}_{x_{i}}\left(\widehat{\varphi w_{h}}\right)\|_{\widehat{F}_{\mathcal{Q},m}} + \|\left(I-\widehat{\pi}_{x_{m}}\right)\left(\partial_{\widehat{x}_{i}}\left(\widehat{\varphi w_{h}}\right)-\partial_{\widehat{x}_{i}}\widehat{\pi}_{x_{i}}\left(\widehat{\varphi w_{h}}\right)\right)\|_{\widehat{F}_{\mathcal{Q},m}} + \|\left(I-\widehat{\pi}_{t}\Pi_{1\leq j\leq d,j\neq i,j\neq m}\widehat{\pi}_{x_{j}}\right)\widehat{\pi}_{x_{m}}\left(\partial_{\widehat{x}_{i}}\left(\widehat{\varphi w_{h}}\right)-\partial_{\widehat{x}_{i}}\widehat{\pi}_{x_{i}}\left(\widehat{\varphi w_{h}}\right)\right)\|_{\widehat{F}_{\mathcal{Q},m}}.
$$
\n(B.13)

For the second and the fifth terms on the right-hand side, we observe that the functions inside the norms are polynomials in the \hat{x}_m -direction. Therefore, Georgoulis [\[23,](#page-30-14) Lemma 7.8] gives us, using similar steps used to find eq. $(B.11)$,

$$
\begin{split}\n&\| (I - \widehat{\pi}_t \Pi_{1 \le j \le d, j \ne i, j \ne m} \widehat{\pi}_{x_j}) \widehat{\pi}_{x_m} \partial_{\widehat{x}_i} \left(\widehat{\varphi w_h} \right) \right\|_{\widehat{F}_{\mathcal{Q},m}} \\
&\quad + \| (I - \widehat{\pi}_t \Pi_{1 \le j \le d, j \ne i, j \ne m} \widehat{\pi}_{x_j}) \widehat{\pi}_{x_m} \left(\partial_{\widehat{x}_i} \left(\widehat{\varphi w_h} \right) - \partial_{\widehat{x}_i} \widehat{\pi}_{x_i} \left(\widehat{\varphi w_h} \right) \right) \|_{\widehat{F}_{\mathcal{Q},m}} \\
&\le c \left(\|\partial_t^{p_t+1} \partial_{\widehat{x}_i} \left(\widehat{\varphi w_h} \right) \|_{\widehat{\mathcal{K}}} + \|\partial_{\widehat{t}} \left(\partial_{\widehat{x}_i} \left(\widehat{\varphi w_h} \right) - \partial_{\widehat{x}_i} \widehat{\pi}_{x_i} \left(\widehat{\varphi w_h} \right) \right) \|_{\widehat{\mathcal{K}}} \right) \\
&\quad + c \left(\sum_{1 \le j \le d, j \ne m, j \ne i} \| \partial_{\widehat{x}_j} \left(\partial_{\widehat{x}_i} \left(\widehat{\varphi w_h} \right) - \partial_{\widehat{x}_i} \widehat{\pi}_{x_i} \left(\widehat{\varphi w_h} \right) \right) \|_{\widehat{\mathcal{K}}} \right).\n\end{split}
$$

Next, using that $\hat{\pi}_{x_i}$ and $\partial_{\hat{x}_j}$ commute, using Georgoulis [\[23,](#page-30-14) eq.(7.5) in Lemma 7.3] and eq. [\(B.2\)](#page-24-3), we find that for any $j \neq i$ where $1 \leq i \leq d$ and $0 \leq j \leq d$ with $\widehat{x}_0 = \widehat{t}$,

$$
\|\partial_{\widehat{x}_j} (\partial_{\widehat{x}_i} (\widehat{\varphi w_h}) - \partial_{\widehat{x}_i} \widehat{\pi}_{x_i} (\widehat{\varphi w_h}))\|_{\widehat{\mathcal{K}}}^2 \leq c \|\partial_{\widehat{x}_i}^{p_s+1} \partial_{\widehat{x}_j} (\widehat{\varphi w_h})\|_{\widehat{\mathcal{K}}}^2 = 0.
$$

Therefore, the second and fifth terms on the right hand side of eq. [\(B.13\)](#page-25-3) are bounded by $c \|\partial_t^{p_t+1}\partial_{\widehat{x}_i}(\widehat{\varphi w}_h)\|_{\widehat{\mathcal{K}}}$.
All remaining terms on the right hand side of eq. (B.13) reprish by combining Congress All remaining terms on the right hand side of eq. [\(B.13\)](#page-25-3) vanish by combining Georgoulis [\[23,](#page-30-14) Lem-mas 7.2 and 7.3] eqs. [\(B.1\)](#page-24-8) and [\(B.2\)](#page-24-3) and so, for $i \neq m$,

$$
\|\partial_{\widehat{x}_i} (\widehat{\varphi w_h} - \widehat{\Pi}_h (\widehat{\varphi w_h}))\|_{\widehat{F}_{\mathcal{Q},m}} \le c \|\partial_{\widehat{t}}^{p_t+1} \partial_{\widehat{x}_i} (\widehat{\varphi w_h})\|_{\widehat{\mathcal{K}}}.
$$
\n(B.14)

Combining eqs. $(B.9)$, $(B.12)$ and $(B.14)$ and applying eqs. $(A.1a)$ and $(B.3b)$ we obtain:

$$
\|\widetilde{\nabla}(\widetilde{\varphi w_h} - \widetilde{\Pi}_h(\widetilde{\varphi w_h}))\|_{\widetilde{F}_{\mathcal{Q},m}} \le c\delta t_{\mathcal{K}} h_K^{-3/2} \|\widetilde{w}_h\|_{\widetilde{\mathcal{K}}}.
$$
\n(B.15)

A scaling argument applied to eq. [\(B.15\)](#page-26-2) from $\widetilde{\mathcal{K}}$ to \mathcal{K} yields eq. [\(4.8c\)](#page-8-7).

Equation [\(4.8d\)](#page-8-3) follows directly by combining the local trace inequality eq. [\(B.1a\)](#page-24-9) with eqs. [\(B.4\)](#page-24-4) and $(B.7)$:

$$
\|\widetilde{\varphi w_h} - \widetilde{\Pi}_h \left(\widetilde{\varphi w_h} \right) \|_{\widetilde{\mathcal{Q}}_{\mathcal{K}}} \le c h_K^{-1/2} \delta t_{\mathcal{K}} \left\| \widetilde{w_h} \right\|_{\widetilde{\mathcal{K}}},\tag{B.16}
$$

and a scaling argument applied to eq. [\(B.16\)](#page-26-3) from \widetilde{K} to K. Lastly, eq. [\(4.8e\)](#page-8-4) follows by combining the local trace inequality eq. $(B.1b)$ with eqs. $(B.4)$ and $(B.8)$:

$$
\|\widetilde{\varphi w_h} - \widetilde{\Pi}_h \left(\widetilde{\varphi w_h} \right) \|_{\widetilde{\mathcal{R}}_{\mathcal{K}}} \le c \delta t_{\mathcal{K}}^{1/2} \left\| \widetilde{w_h} \right\|_{\widetilde{\mathcal{K}}},\tag{B.17}
$$

and a scaling argument applied to eq. [\(B.17\)](#page-26-4) from \widetilde{K} to K.

C Proof of lemma [4.4](#page-9-3)

We start with volume terms in the definition of $\|\cdot\|_v$ eq. [\(3.8a\)](#page-6-4). Due to boundedness of Π_h , and using eq. $(4.7a)$,

$$
\sum_{\mathcal{K}\in\mathcal{T}_h} \|\Pi_h(\varphi w_h)\|_{\mathcal{K}}^2 + \sum_{\mathcal{K}\in\mathcal{T}_h} \varepsilon \|\overline{\nabla} \left(\Pi_h(\varphi w_h)\right)\|_{\mathcal{K}}^2 \le c\left(eT+\chi\right)^2 \|\|\boldsymbol{w}_h\|_{v}^2. \tag{C.1}
$$

Next, the diffusive facet terms are bounded using a triangle inequality, eq. [\(4.7c\)](#page-8-6), and boundedness of $\Pi_h^{\mathcal{F}}$:

$$
\sum_{\mathcal{K}\in\mathcal{T}_{h}}\varepsilon h_{K}^{-1} \left\|\Pi_{h}\left(\varphi w_{h}\right)-\Pi_{h}^{\mathcal{F}}\left(\varphi\varkappa_{h}\right)\right\|_{\mathcal{Q}_{\mathcal{K}}}^{2} \n\leq c \sum_{\mathcal{K}\in\mathcal{T}_{h}}\varepsilon h_{K}^{-1}h_{K} \left\|\overline{\nabla}\left(\varphi w_{h}\right)\right\|_{\mathcal{K}}^{2}+c \sum_{\mathcal{K}\in\mathcal{T}_{h}}\varepsilon h_{K}^{-1} \left\|\varphi\left[w_{h}\right]\right\|_{\mathcal{Q}_{\mathcal{K}}}^{2} \leq c\left(\epsilon T+\chi\right)^{2} \left\|\left(w_{h}\right)\right\|_{v}^{2}.
$$
\n(C.2)

For the Neumann boundary term in the definition of $\|\cdot\|_v$, consider first a single facet $F \in \partial \mathcal{E}_N$. Then,

$$
\|\left[\frac{1}{2}\beta \cdot n\right]^{1/2} \Pi_h^{\mathcal{F}}\left(\varphi \varkappa_h\right)\|_F
$$

\$\leq \|\left[\frac{1}{2}\beta \cdot n\right]^{1/2} \left(\Pi_h^{\mathcal{F}}\left(\varphi \varkappa_h\right) - \Pi_h\left(\varphi w_h\right)\right)\|_F + \|\left[\frac{1}{2}\beta \cdot n\right]^{1/2} \Pi_h\left(\varphi w_h\right)\|_F := I + II\$.

For term I, using that $|\beta \cdot n| \leq (\max_{(t,x)\in F} |\beta \cdot n|)$, that $\Pi_h^{\mathcal{F}} \Pi_h(\varphi w_h) = \Pi_h(\varphi w_h)$ on F, boundedness of $\Pi_h^{\mathcal{F}}$ and a triangle inequality we have:

$$
\begin{split} \|\left|\frac{1}{2}\beta \cdot n\right|^{1/2} \left(\Pi_{h}^{\mathcal{F}}\left(\varphi \varkappa_{h}\right)-\Pi_{h}\left(\varphi w_{h}\right)\right)\|_{F} &\leq \left(\frac{1}{2} \max_{(t,x)\in F} \left|\beta \cdot n\right|\right)^{1/2} \|\varphi \varkappa_{h}-\Pi_{h}\left(\varphi w_{h}\right)\|_{F} \\ &\leq \left(\frac{1}{2} \max_{(t,x)\in F} \left|\beta \cdot n\right|\right)^{1/2} \|\varphi \varkappa_{h}-\varphi w_{h}\|_{F} \\ &+ \left(\frac{1}{2} \max_{(t,x)\in F} \left|\beta \cdot n\right|\right)^{1/2} \left\|(I-\Pi_{h})\left(\varphi w_{h}\right)\right\|_{F} .\end{split}
$$

Using that $|\varphi| \leq eT + \chi$ and eq. [\(3.9\)](#page-6-1) for the first term on the right hand side, and using eq. [\(4.8d\)](#page-8-3) and that $\delta t_{\mathcal{K}} \leq h_K$ for the second term, we obtain

$$
I \leq (eT + \chi) \left\| \left(\beta_s - \frac{1}{2}\beta \cdot n\right)^{1/2} \left(\varkappa_h - w_h\right) \right\|_F + c \left\| w_h \right\|_{\mathcal{K}_F},\tag{C.3}
$$

where \mathcal{K}_F is the space-time element of which F is a facet. Next, for term II, by a triangle inequality, using eq. [\(4.8d\)](#page-8-3), that $\delta t_{\mathcal{K}} \leq h_K$, $|\varphi| \leq eT + \chi$, and eq. [\(3.9\)](#page-6-1),

$$
II \leq |||_{2}^{1} \beta \cdot n|^{1/2} (I - \Pi_{h})(\varphi w_{h})||_{F} + |||_{2}^{1} \beta \cdot n|^{1/2} \varphi(w_{h} - \varkappa_{h})||_{F} + (eT + \chi) |||_{2}^{1} \beta \cdot n|^{1/2} \varkappa_{h}||_{F}
$$

$$
\leq c ||w_{h}||_{K} + (eT + \chi) ||| \beta_{s} - \frac{1}{2} \beta \cdot n|^{1/2} (w_{h} - \varkappa_{h})||_{F} + (eT + \chi) |||_{2}^{1} \beta \cdot n|^{1/2} \varkappa_{h}||_{F}.
$$

For a facet $F \in \partial \mathcal{E}_N$ we therefore conclude that

$$
\left\| \left| \frac{1}{2} \beta \cdot n \right|^{1/2} \Pi_h^{\mathcal{F}} \left(\varphi \varkappa_h \right) \right\|_F \le c \left(eT + \chi \right) \left\| \left(\beta_s - \frac{1}{2} \beta \cdot n \right)^{1/2} \left(\varkappa_h - w_h \right) \right\|_F
$$

+ $c \left\| w_h \right\|_K + \left(eT + \chi \right) \left\| \left| \frac{1}{2} \beta \cdot n \right|^{1/2} \varkappa_h \right\|_F.$ (C.4)

We find for the Neumann term in the definition of $\|\cdot\|_{v}$:

$$
\sum_{F \in \partial \mathcal{E}_N} |||_{2}^{\frac{1}{2}} \beta \cdot n|^{1/2} \Pi_h^{\mathcal{F}} (\varphi \varkappa_h) ||_F^2 \le c ||\beta||_{L^{\infty}(\mathcal{E})} \sum_{\mathcal{K} \in \mathcal{T}_h} ||w_h||_{\mathcal{K}}^2 + (eT + \chi)^2 \left(\sum_{\mathcal{K} \in \mathcal{T}_h} ||(\beta_s - \frac{1}{2}\beta \cdot n)^{1/2} (\varkappa_h - w_h) ||_{\partial \mathcal{K}}^2 + \sum_{F \in \partial \mathcal{E}_N} |||_{2}^{\frac{1}{2}} \beta \cdot n|^{1/2} \varkappa_h||_F^2 \right). \tag{C.5}
$$

Finally, we consider the advective facet terms in the definition of $\|\cdot\|_v$. On a single facet we have:

$$
\| |\beta_s - \frac{1}{2} \beta \cdot n|^{1/2} \left(\Pi_h \left(\varphi w_h \right) - \Pi_h^{\mathcal{F}} \left(\varphi \varkappa_h \right) \right) \|_F \leq c \big(\max_{(t,x) \in F} |\beta \cdot n| \big)^{1/2} \left\| \left(\Pi_h \left(\varphi w_h \right) - \Pi_h^{\mathcal{F}} \left(\varphi \varkappa_h \right) \right) \right\|_F.
$$

Using identical steps as used to find the bound for I in eq. $(C.3)$, we find:

$$
\|\beta_s - \frac{1}{2}\beta \cdot n\|^{1/2} \left(\Pi_h \left(\varphi w_h\right) - \Pi_h^{\mathcal{F}} \left(\varphi \varkappa_h\right)\right)\|_F
$$

$$
\leq c \left(eT + \chi\right) \left\| \left(\beta_s - \frac{1}{2}\beta \cdot n\right)^{1/2} \left(\varkappa_h - w_h\right)\right\|_F + c \left\| w_h \right\|_{\mathcal{K}},
$$

so that

$$
\sum_{\mathcal{K}\in\mathcal{T}_h} ||\beta_s - \frac{1}{2}\beta \cdot n|^{1/2} \left(\Pi_h(\varphi w_h) - \Pi_h^{\mathcal{F}}(\varphi w_h)\right)\|_{\partial\mathcal{K}}^2
$$
\n
$$
\leq c\left(eT + \chi\right)^2 \sum_{\mathcal{K}\in\mathcal{T}_h} ||(\beta_s - \frac{1}{2}\beta \cdot n)^{1/2} \left(\varkappa_h - w_h\right)\|_{\partial\mathcal{K}}^2 + c\left(\sum_{\mathcal{K}\in\mathcal{T}_h} ||w_h||_{\mathcal{K}}^2\right). \quad (C.6)
$$

The result follows after collecting the bounds in eqs. $(C.1)$, $(C.2)$, $(C.5)$ and $(C.6)$.

Acknowledgements

SR gratefully acknowledges support from the Natural Sciences and Engineering Research Council of Canada through the Discovery Grant program (RGPIN-05606-2015).

This research was enabled in part by support provided by Simon Fraser University ([https://](https://www.sfu.ca/research/supercomputer-cedar) www.sfu.ca/research/supercomputer-cedar), Compute Ontario ([https://www.computeontario](https://www.computeontario.ca/). [ca/](https://www.computeontario.ca/)) and the Digital Research Alliance of Canada (<https://alliancecan.ca>). We furthermore acknowledge the support provided by the Math Faculty Computing Facility at the University of Waterloo (<https://uwaterloo.ca/math-faculty-computing-facility/>).

References

- [1] D. Arndt, W. Bangerth, M. Feder, M. Fehling, R. Gassmöller, T. Heister, L. Heltai, M. Kronbichler, M. Maier, P. Munch, J.-P. Pelteret, S. Sticko, B. Turcksin, and D. Wells. The deal.II library, version 9.4. J. Numer. Math., 30(3):231–246, 2022. doi: 10.1515/jnma-2022-0054.
- [2] B. Ayuso and L. D. Marini. Discontinuous Galerkin methods for advection-diffusion-reaction problems. SIAM J. Numer. Anal., 47(2):1391–1420, 2009. doi: 10.1137/080719583.
- [3] S. Balay, W. D. Gropp, L. C. McInnes, and B. F. Smith. Efficient management of parallelism in object oriented numerical software libraries. In E. Arge, A. M. Bruaset, and H. P. Langtangen, editors, *Modern Software Tools in Scientific Computing*, pages $163-202$. Birkhäuser Press, 1997.
- [4] S. Balay, S. Abhyankar, M. F. Adams, S. Benson, J. Brown, P. Brune, K. Buschelman, E. Constantinescu, L. Dalcin, A. Dener, V. Eijkhout, J. Faibussowitsch, W. D. Gropp, V. Hapla, T. Isaac, P. Jolivet, D. Karpeev, D. Kaushik, M. G. Knepley, F. Kong, S. Kruger, D. A. May, L. C. McInnes, R. T. Mills, L. Mitchell, T. Munson, J. E. Roman, K. Rupp, P. Sanan, J. Sarich, B. F. Smith, S. Zampini, H. Zhang, H. Zhang, and J. Zhang. PETSc/TAO users manual. Technical Report ANL-21/39 - Revision 3.19, Argonne National Laboratory, 2023.
- [5] S. Balay, S. Abhyankar, M. F. Adams, S. Benson, J. Brown, P. Brune, K. Buschelman, E. M. Constantinescu, L. Dalcin, A. Dener, V. Eijkhout, J. Faibussowitsch, W. D. Gropp, V. Hapla, T. Isaac, P. Jolivet, D. Karpeev, D. Kaushik, M. G. Knepley, F. Kong, S. Kruger, D. A. May, L. C. McInnes, R. T. Mills, L. Mitchell, T. Munson, J. E. Roman, K. Rupp, P. Sanan, J. Sarich, B. F. Smith, S. Zampini, H. Zhang, H. Zhang, and J. Zhang. PETSc Web page. <https://petsc.org/>, 2023. URL <https://petsc.org/>.
- [6] W. Bangerth, C. Burstedde, T. Heister, and M. Kronbichler. Algorithms and data structures for massively parallel generic adaptive finite element codes. ACM Trans. Math. Software, 38 (2):14:1–14:28, 2011. doi: 10.1145/2049673.2049678.
- [7] A. N. Brooks and T. J. R. Hughes. Streamline upwind/Petrov–Galerkin formulations for convection dominated flows with particular emphasis on the incompressible Navier–Stokes equations. Comput. Methods Appl. Mech. Engrg., 32(1-3):199–259, 1982. doi: 10.1016/ 0045-7825(82)90071-8.
- [8] E. Burman. A unified analysis for conforming and nonconforming stabilized finite element methods using interior penalty. SIAM J. Numer. Anal., 43(5):2012–2033, 2005. doi: 10.1137/ S0036142903437374.
- [9] E. Burman. Consistent SUPG-method for transient transport problems: Stability and convergence. Comput. Methods Appl. Mech. Engrg., 199(17-20):1114–1123, 2010. doi: 10.1016/j.cma.2009.11.023.
- [10] E. Burman and M. A. Fernández. Finite element methods with symmetric stabilization for the transient convection-diffusion-reaction equation. Comput. Methods Appl. Mech. Engrg., 198 (33-36):2508–2519, 2009. doi: 10.1016/j.cma.2009.02.011.
- [11] E. Burman and P. Hansbo. Edge stabilization for Galerkin approximations of convectiondiffusion-reaction problems. Comput. Methods Appl. Mech. Engra., 193(15-16):1437-1453, 2004. doi: 10.1016/j.cma.2003.12.032.
- [12] E. Burman and B. Stamm. Minimal stabilization for discontinuous Galerkin finite element methods for hyperbolic problems. J. Sci. Comput., 33:183–208, 2007. doi: 10.1007/ s10915-007-9149-5.
- [13] Y. Chen and B. Cockburn. Analysis of variable-degree HDG methods for convection-diffusion equations. Part I: general nonconforming meshes. IMA J. Numer. Anal., 32(4):1267–1293, 2012. doi: 10.1093/imanum/drr058.
- [14] B. Cockburn and C.-W. Shu. The local discontinuous Galerkin method for time-dependent convection-diffusion systems. SIAM J. Numer. Anal., 35(6):2440–2463, 1998. doi: 10.1137/ S0036142997316712.
- [15] B. Cockburn and C.-W. Shu. Runge–Kutta discontinuous Galerkin methods for convectiondominated problems. J. Sci. Comput., 16:173–261, 2001. doi: 10.1023/A:1012873910884.
- [16] B. Cockburn, J. Gopalakrishnan, and R. Lazarov. Unified hybridization of discontinuous Galerkin, mixed, and continuous Galerkin methods for second order elliptic problems. SIAM J. Numer. Anal., 47(2):1319–1365, 2009. doi: 10.1137/070706616.
- [17] J. de Frutos, B. García-Archilla, and J. Novo. Local error estimates for the SUPG method applied to evolutionary convection-reaction-diffusion equations. J. Sci. Comput., $66:528-554$, 2016. doi: 10.1007/s10915-015-0035-2.
- [18] H. Egger and J. Schöberl. A hybrid mixed discontinuous Galerkin finite-element method for convection-diffusion problems. IMA J. Numer. Anal., 30(4):1206–1234, 2010. doi: 10.1093/ imanum/drn083.
- [19] A. Ern and J.-L. Guermond. Discontinuous Galerkin methods for Friedrichs' systems. I. General theory. SIAM J. Numer. Anal., 44(2):753–778, 2006. doi: 10.1137/050624133.
- [20] M. Feistauer, V. Kučera, K. Najzar, and J. Prokopová. Analysis of space-time discontinuous Galerkin method for nonlinear convection-diffusion problems. Numer. Math., 117:251–288, 2011. doi: 10.1007/s00211-010-0348-x.
- [21] G. Fu, W. Qiu, and W. Zhang. An analysis of HDG methods for convection-dominated diffusion problems. ESAIM Math. Model. Numer. Anal., 49(1):225–256, 2015. doi: 10.1051/ m2an/2014032.
- [22] E. H. Georgoulis. Discontinuous Galerkin Methods on Shape-Regular and Anisotropic Meshes. PhD thesis, University of Oxford, 2003.
- [23] E. H. Georgoulis. hp-version interior penalty discontinuous Galerkin finite element methods on anisotropic meshes. Int. J. Numer. Anal. Model., 3(1):52–79, 2006.
- [24] V. E. Henson and U. M. Yang. BoomerAMG: A parallel algebraic multigrid solver and preconditioner. Appl. Numer. Math., 41(1):155–177, 2002. doi: 10.1016/S0168-9274(01)00115-5.
- [25] P. Houston, C. Schwab, and E. Süli. Discontinuous hp -finite element methods for advectiondiffusion-reaction problems. SIAM J. Numer. Anal., 39(6):2133–2163, 2002. doi: 10.1137/ S0036142900374111.
- [26] T. J. R. Hughes, L. P. Franca, and M. Mallet. A new finite element formulation for computational fluid dynamics: VI. Convergence analysis of the generalized SUPG formulation for linear time-dependent multidimensional advective-diffusive systems. Comput. Methods Appl. Mech. Engrg., 63(1):97–112, 1987. doi: 10.1016/0045-7825(87)90125-3.
- [27] P. Jamet. Galerkin-type approximations which are discontinuous in time for parabolic equations in a variable domain. SIAM J. Numer. Anal., 15(5):912–928, 1978. doi: 10.1137/0715059.
- [28] P. Jamet and B. Bonnerot. Numerical solution of the Eulerian equations on compressible flow by a finite element method which follows the free boundary and the interfaces. J. Comput. Phys., 18(1):21–45, 1975. doi: 10.1016/0021-9991(75)90100-X.
- [29] V. John and J. Novo. Error analysis of the SUPG finite element discretization of evolutionary convection-diffusion-reaction equations. SIAM J. Numer. Anal., 49(3):1149–1176, 2011. doi: 10.1137/100789002.
- [30] K. L. A. Kirk, T. L. Horvath, A. Cesmelioglu, and S. Rhebergen. Analysis of a spacetime hybridizable discontinuous Galerkin method for the advection-diffusion problem on timedependent domains. SIAM J. Numer. Anal., 57(4):1677–1696, 2019. doi: 10.1137/18M1202049.
- [31] T. A. Manteuffel, J. Ruge, and B. S. Southworth. Nonsymmetric algebraic multigrid based on local approximate ideal restriction (ℓAIR). SIAM J. Sci. Comput., 40(6):A4105-A4130, 2018. doi: 10.1137/17M1144350.
- [32] T. A. Manteuffel, S. Münzenmaier, J. Ruge, and B. S. Southworth. Nonsymmetric reductionbased algebraic multigrid. SIAM J. Sci. Comput., 41(5):S242–S268, 2019. doi: 10.1137/ 18M1193761.
- [33] J. R. Munkres. Analysis on Manifolds. Addison-Wesley, Redwood City, CA, 1991.
- [34] N. C. Nguyen, J. Peraire, and B. Cockburn. An implicit high-order hybridizable discontinuous Galerkin method for linear convection-diffusion equations. J. Comput. Phys., 228(9):3232-3254, 2009. doi: 10.1016/j.jcp.2009.01.030.
- [35] D. A. D. Pietro and A. Ern. Mathematical Aspects of Discontinuous Galerkin Methods, volume 69 of Mathématiques et Applications. Springer-Verlag Berlin Heidelberg, 2012.
- [36] W. H. Reed and T. R. Hill. Triangular mesh methods for the neutron transport equation. Technical report, Los Alamos Scientific Laboratory, Tech. Report LA-UR-73-479, 1973.
- [37] S. Rhebergen and B. Cockburn. A space-time hybridizable discontinuous Galerkin method for incompressible flows on deforming domains. J. Comput. Phys., 231(11):4185–4204, 2012. doi: 10.1016/j.jcp.2012.02.011.
- [38] S. Rhebergen and B. Cockburn. Space-time hybridizable discontinuous Galerkin method for the advection-diffusion equation on moving and deforming meshes. In C. A. de Moura and C. S. Kubrusly, editors, The Courant–Friedrichs–Lewy (CFL) condition, 80 years after its discovery, pages 45–63. Birkhäuser Science, 2013. doi: 10.1007/978-0-8176-8394-8.4.
- [39] B. Rivière. Discontinuous Galerkin Methods for Solving Elliptic and Parabolic Equations, volume 35 of Frontiers in Applied Mathematics. Society for Industrial and Applied Mathematics, Philadelphia, 2008.
- [40] A. A. Sivas, B. S. Southworth, and S. Rhebergen. AIR algebraic multigrid for a space-time hybridizable discontinuous Galerkin discretization of advection(-diffusion). SIAM J. Sci. Comput., 43(5):A3393–A3416, 2021. doi: 10.1137/20M1375103.
- [41] J. J. Sudirham. Space-time discontinuous Galerkin methods for convection-diffusion problems: Application to wet-chemical etching. PhD thesis, University of Twente, 2005.
- [42] J. J. Sudirham, J. J. W. van der Vegt, and R. M. J. van Damme. Space-time discontinuous Galerkin method for advection-diffusion problems on time-dependent domains. Appl. Numer. Math., 56(12):1491–1518, 2006. doi: 10.1016/j.apnum.2005.11.003.
- [43] J. J. W. van der Vegt and H. van der Ven. Space-time discontinuous Galerkin finite element method with dynamic grid motion for inviscid compressible flows: I. General formulation. J. Comput. Phys., 182(2):546–585, 2002. doi: 10.1006/jcph.2002.7185.
- [44] G. N. Wells. Analysis of an interface stabilized finite element method: the advection-diffusionreaction equation. SIAM J. Numer. Anal., 49(1):87–109, 2011. doi: 10.1137/090775464.