Objects in JWST's mirrors are closer than they appear

Stephen Serjeant¹, Tom J.L.C. Bakx²

¹ School of Physical Sciences, The Open University, Milton Keynes, MK7 6AA, UK ² Department of Space, Earth, & Environment, Chalmers University of Technology, Chalmersplatsen 4 412 96 Gothenburg, Sweden

The James Webb Space Telescope (JWST) has revealed extremely distant galaxies at unprecedentedly early cosmic epochs from its deep imaging using the technique of photometric redshift estimation (e.g., ¹), with its subsequent spectroscopy confirming their redshifts unambiguously (e.g., ²), demonstrating the ability of JWST to probe the earliest galaxies, one of its major scientific goals. However, as larger samples continue to be followed up spectroscopically, it has become apparent that nearly all photometric redshifts at these epochs are biased high with confidence >>99%, for as yet unclear reasons. Here we show that this is the same statistical effect that was predicted in different contexts by Sir Arthur Eddington³ in 1913, in that there exist more lower redshift galaxies to be scattered upwards than the reverse. The bias depends on the shape of the intrinsic redshift distribution, but as an approximate heuristic, all ultra-high photometric redshift estimates must be corrected downwards by up to one standard deviation.

In this work we use a recent spectroscopic redshift compilation⁴ comprising 26 galaxies with redshifts of z_{spec} =8.61 to 13.20. This is the largest and most comprehensive ultra-high-redshift spectroscopic sample to date and includes the spectroscopic confirmations of the four galaxies in Ref.¹. The galaxies in this compilation were initially selected via the inferred presence of the Lyman absorption break, in which the intervening neutral intergalactic medium absorbs essentially all photons at energies above the *n*=2-1 atomic hydrogen transition. These redshifts are well within the reionization epoch, before population III stars, OB stars and active nuclei reionized their surroundings, so the overwhelming preponderance of neutral gas is predicted to generate a very strong Lyman decrement. We include the galaxy GN-z11, with a photometric redshift estimate⁶ of $z_{phot} = 11.09^{+0.08}_{-0.12}$ from grism spectroscopy binned to a coarse wavelength resolution similar to those of Hubble Space Telescope (HST) medium-width passbands. We exclude one galaxy from the compilation, GS+53.11243-27.77461, because it was discovered in blind spectroscopy rather than being pre-selected photometrically, but add the spectroscopic confirmation⁵ of the galaxy GLASS-z12.

In Figure 1, we compare the spectroscopic redshifts with the original photometrically-estimated redshifts⁴. The photometric redshifts are subtly but strikingly systematically overestimating the accurate spectroscopic measurements. The photometric redshift uncertainties are often

asymmetric; in about half of our sample the lower error is worse, while in the rest of the sample the upper error is equal or worse. Of the 26 galaxies in this sample, 21 show photometric redshift overprediction. We can reject the null hypothesis of a p=0.5 binomial distribution at 99.8% confidence. The mean of the histogram in Figure 1 is offset from zero with a significance 4.1 σ , equivalent to 99.9998% confidence. An Anderson-Darling test rejects a Gaussian with zero mean and unit variance at a significance level of ~2×10⁻⁸.

Eddington³ argued for a systematic bias in the number counts of stars arising from the effect of measurement uncertainties. This bias is well-known to affect faint submillimetre galaxy surveys. The same principle applies here. The luminosity function evolves steeply with redshift at these epochs, not only with a $(1+z)^4$ decline in luminosity density and comoving volume elements dV/dz also scaling inversely approximately as (1+z), but also with a steep bright-end luminosity function slope⁷, and with luminosities at a given flux density also scaling approximately as $(1+z)^{1.5}$ it is likely that each ultra-high-redshift galaxy is sampled from a population with extremely steep intrinsic number counts. The observed compilation in Ref.⁴, however, is a heterogeneous compilation with a diverse range of selection effects between objects, so the observed N(z) in Ref.⁴ need not reflect the steepness of the underlying number counts or luminosity function in an obvious way; indeed, the largest predictions at *z*-13 and *z*-15 for each survey in Ref.⁸ imply intrinsic $dN/dz \propto (1+z)^{-k}$ with $k = 19 \pm 6$.




Figure 1: The left panel compares the photometric redshift estimates against the precise spectroscopic measurements in our ultra high-redshift sample, including the four galaxies in Ref.^{*z*1}. The red dashed line shows the one-to-one relation; note that the data trends above this line. The right panel shows the histogram of photometric redshift overestimation divided by the uncertainties in photometric redshift, with the red dashed line again indicating no bias; note the striking offset to the left.

Each ultra-high-redshift galaxy will therefore be selected from an intrinsic number count dN/dz that has been convolved with the photometric redshift probability distribution function. An exact analytic

form does not exist, but we find the mean offset of photometric and exact spectroscopic redshifts can be fairly well approximated by

 $\langle z_{spec} - z_{phot} \rangle \approx -0.949 \, k^{1.03} z_{phot}^{-1.01} \sigma^{2.03}$ (1)

where the intrinsic dN/dz is parameterised as being a $(1+z)^{-k}$ power law, and σ is a symmetrical uncertainty in photometric redshift. This analytic prediction agrees to within 20% of the full numerical calculation for $7.5 \le z_{phot} \le 26.5$, $0.1 \le \sigma \le 0.75$ and $5 \le k \le 24$. This approximate relation is roughly linear in *k* but has a surprisingly strong dependence on σ . A Schechter luminosity function would further curve the redshift distribution downwards at the highest redshifts, making the Eddington bias stronger, similarly to how magnification bias preferentially affects the number counts of the brightest submillimetre galaxies.

Furthermore, the photometric redshift uncertainties are themselves likely to be understated due to limitations in the templates used. The presence or absence of the Ly α emission line results in a mean shift of 0.07 in photometric redshift⁹, while the discovery of a Ly α absorption damping wing (e.g. Ref.¹⁰) can shift the estimates by 0.06 in the opposite direction. At lower redshifts, template-based systematics can easily approach¹¹ 0.02(1+z). Depending on how likelihoods or posteriors are implemented, there may also be stochastic biases¹³ at a similar level to the random errors driven by the asymmetric uncertainties. In partial support of our argument that photometric redshift estimates tend to subtly understate their uncertainties, the JWST spectrum¹² of Ly α in GN-z11 is curiously inconsistent with the apparent position of the spectral break in the earlier HST grism data⁶, suggesting either underestimated uncertainties or systematics in the difficult subtraction of foreground contaminants in the HST grism data, and/or insufficient signal-to-noise per spaxel to detect the extended Ly α , and/or insufficient diversity in templates to capture the possible spatio-spectral variation in continuum, HI opacity and Ly α emission.

As an illustrative exercise, we find that the distribution of $(z_{spec}-z_{phot})/\sigma$ can be made consistent with a Gaussian distribution with zero mean and unit variance if the z_{phot} values are corrected assuming k=19 (consistent with Ref.⁸) and a template-based additional systematic uncertainty^{9,10} of $\sigma(z)=\pm 0.1$. We conclude that Eddington bias, together with a subtle underestimate in photometric redshift uncertainties, can entirely account for the systematic shift. Our recommendation is that future photometric redshift works incorporate a model intrinsic dN/dz into Bayesian priors, with both biases and uncertainties tested against simulations.

The authors declare no competing interests.

References

 Robertson, B.E., et al., Identification and properties of intense star-forming galaxies at redshifts z > 10. Nature Astronomy, 7, 611–621 (2023)

- Curtis-Lake, E., et al., Spectroscopic confirmation of four metal-poor galaxies at z = 10.3-13.2. *Nature Astronomy*, **7**, 622-632 (2023)
- 3. Eddington, A.S., On a formula for correcting statistics for the effects of a known error of observation. *Mon. Not. R. Astr. Soc.*, **73**, 359-360 (1913)
- 4. Harikane, Y., et al., Pure Spectroscopic Constraints on UV Luminosity Functions and

Cosmic Star Formation History From 25 Galaxies at zspec=8.61-13.20 Confirmed with

JWST/NIRSpec. Preprint at https://arxiv.org/abs/2304.06658 (2023)

- 5. Bakx, T.J.L.C., et al., Deep ALMA redshift search of a z~12 GLASS-JWST galaxy candidate. *Mon. Not. R. Astr. Soc.*, **519**, 5076-5085 (2023)
- 6. Oesch, P.A., et al., A Remarkably Luminous Galaxy at z=11.1 Measured with Hubble Space Telescope Grism Spectroscopy. *Astrophys. J.*, **819**, 129 (2016)
- Bouwens, R.J., et al., Evolution of the UV LF from z~15 to z~8 Using New JWST NIRCam Medium-Band Observations over the HUDF/XDF. *Mon. Not. R. Astr. Soc.*, **523**, 1036-1055 (2023)
- Harikane, Y., et al., A Search for H-Dropout Lyman Break Galaxies at z ~ 12–16, Astrophys. J., 929, 1 (2022)
- Hainline, K.N., et al., The Cosmos in its Infancy: JADES Galaxy Candidates at z > 8 in GOODS-S and GOODS-N. Preprint at <u>https://arxiv.org/abs/2306.02468</u> (2023)
- 10. Heintz., K.E., et al., Extreme damped Lyman- α absorption in young star-forming galaxies at

z=9-11. Preprint at <u>https://arxiv.org/abs/2306.00647</u> (2023)

- 11. Molino, A., et al., Assessing the photometric redshift precision of the S-PLUS survey: the Stripe-82 as a test-case. *Mon. Not. R. Astr. Soc.*, **499**, 3884-3908 (2020)
- Bunker, A.J., et al., JADES NIRSpec Spectroscopy of GN-z11: Lyman-α emission and possible enhanced nitrogen abundance in a z = 10.60 luminous galaxy. Preprint at <u>https://arxiv.org/abs/2302.07256</u> (2023)
- 13. Sellentin, E., et al., The skewed weak lensing likelihood: why biases arise, despite data and theory being sound. *Mon. Not. R. Astr. Soc.*, **477**, 4879-4895 (2018)