

# ON AN ADJUSTMENT TO THE FOWLER EQUATION

BORYS ÁLVAREZ-SAMANIEGO

*Núcleo de Investigadores Científicos*

*Facultad de Ciencias*

*Universidad Central del Ecuador (UCE)*

*Quito, Ecuador*

WILSON P. ÁLVAREZ-SAMANIEGO

*Núcleo de Investigadores Científicos*

*Facultad de Ciencias*

*Universidad Central del Ecuador (UCE)*

*Quito, Ecuador*

KEVIN LLOACANA-UNDA

*Núcleo de Investigadores Científicos*

*Facultad de Ciencias*

*Universidad Central del Ecuador (UCE)*

*Quito, Ecuador*

**ABSTRACT.** Following closely the analysis performed by Andrew C. Fowler to derive the first canonical equation for nonlinear dune dynamics, but considering some appropriate changes of variables, suitable scalings, and by neglecting higher order terms, we obtain an adaptation of the aforementioned equation, which contains an additional term, to describe dune morphodynamics.

---

*E-mail addresses:* balvarez@uce.edu.ec, borys\_yamil@yahoo.com,  
wpalvarez@uce.edu.ec, alvarezwilson@hotmail.com, kpaul\_rap@hotmail.com.

*Date:* July 18, 2023.

*2020 Mathematics Subject Classification.* 47J35; 35Q35.

*Key words and phrases.* Fowler equation; nonlocal evolution equation.

## 1. INTRODUCTION

Dunes are landforms of sand which are formed by the erosive effect of some mobile medium above an erodible substrate. There are several types of dunes, such as Barchan, transverse, star, parabolic, and others ([2, 3, 4]).

A. C. Fowler, relatively recently, proposed the first canonical equation, given by (2.1) below, to model nonlinear dune formation ([2, 3, 4]). Then, B. Álvarez-Samaniego, and P. Azerad showed, in [1], the existence of travelling-waves, and local well-posedness, in a subspace of  $C_b^1(\mathbb{R})$ , of the aforesaid equation, which is called the Fowler equation.

At the beginning of Section 2, we mention some previous results that can be found in [1, 2, 3, 4]. More specifically, we first write the Exner equation, given in (2.2), which describes the conservation of bed material ([1, 4]). Next, by taking into account an additional force due to gravity, an approximation for the net stress is written in (2.3), which was obtained from a suitable analysis considering the Orr-Sommerfeld equation from a Saint-Venant type turbulent model ([4]). Also, an adequate estimate for the bedload transport is exhibited in (2.4), see [1, 2, 3].

Subsequently, by performing a convenient change of variable, (2.12), and a proper scaling of the dependent variable, (2.23), we obtain a suited approximation, (2.24), to analyze sand dune dynamics. Then, by neglecting the terms of order  $O\left(\frac{\varepsilon}{\delta^{\frac{1}{3}}}\right)$ , with  $0 < \varepsilon \ll 1 \ll \delta$ , where  $\varepsilon$  and  $\delta$  are typical physical scales associated with the amplitude, and the length of the dune, respectively, we get (2.26). Finally, by making a suitable change of variable, (2.29), and a handy scaling, (2.37), an adaptation of the Fowler equation is achieved in (2.39), which is the main result of this manuscript.

## 2. DEDUCTION OF THE EQUATION

We begin this section by exposing some previous results, given in [1, 2, 3, 4]. The Fowler equation for nonlinear dune formation is given by

$$\partial_2 u(x, t) + \partial_1 \left[ \frac{u^2}{2} + \int_0^{+\infty} \xi^{-\frac{1}{3}} \partial_1 u(\cdot - \xi, \cdot) d\xi - \partial_1 u \right] (x, t) = 0, \quad (2.1)$$

for all  $(x, t) \in \mathbb{R} \times [0, +\infty)$ , where  $u$  represents the dune amplitude. The derivation of equation (2.1) follows from the Exner equation, namely,

$$\partial_2 u(x, t) + \partial_1(q \circ \tau)(x, t) = 0, \quad (2.2)$$

for all  $(x, t) \in \mathbb{R} \times [0, +\infty)$ , where  $q$  is the bedload transport, and  $\tau$  is the stress exerted by the fluid on the erodible bed. Moreover, the stress,  $\tau$ , can be approximated by the expression

$$\tau(x, t) \approx 1 + u(x, t) + \int_0^{+\infty} \xi^{-\frac{1}{3}} \partial_1 u(x - \xi, t) d\xi + u^2(x, t) - \partial_1 u(x, t), \quad (2.3)$$

for all  $(x, t) \in \mathbb{R} \times [0, +\infty)$ . By considering function  $q$ , proposed by Meyer-Peter, and Müller, a Taylor expansion is performed, around the point 1, up to order 2, which is represented by the following approximation

$$q(\tau) \approx q(1) + q'(1)(\tau - 1) + \frac{q''(1)}{2}(\tau - 1)^2, \quad (2.4)$$

for all real number  $\tau$  belonging to some neighborhood of 1. Let  $(x, t) \in \mathbb{R} \times [0, +\infty)$ . By using the Chain Rule, we have that

$$\partial_1(q \circ \tau)(x, t) = q'(\tau(x, t))\partial_1\tau(x, t). \quad (2.5)$$

Now, by employing (2.4), and (2.3), we get

$$\begin{aligned} q'(\tau(x, t)) &\approx q'(1) + q''(1)(\tau(x, t) - 1) \\ &\approx q'(1) + q''(1) \left( u(x, t) + u^2(x, t) - \partial_1 u(x, t) \right. \\ &\quad \left. + \int_0^{+\infty} \xi^{-\frac{1}{3}} \partial_1 u(x - \xi, t) d\xi \right). \end{aligned} \quad (2.6)$$

It follows from (2.3) that

$$\begin{aligned} \partial_1\tau(x, t) &\approx \partial_1 u(x, t) + 2u(x, t)\partial_1 u(x, t) - \partial_1^2 u(x, t) \\ &\quad + \partial_1 \left( \int_0^{+\infty} \xi^{-\frac{1}{3}} \partial_1 u(\cdot - \xi, \cdot) d\xi \right)(x, t). \end{aligned} \quad (2.7)$$

By using (2.6), and (2.7), we obtain that

$$\begin{aligned} &q'(\tau(x, t))\partial_1\tau(x, t) \\ &\approx [q'(1) + q''(1)(\tau(x, t) - 1)]\partial_1\tau(x, t) \\ &= q'(1)\partial_1\tau(x, t) + q''(1)(\tau(x, t) - 1)\partial_1\tau(x, t) \\ &\approx q'(1)\partial_1 u(x, t) + q'(1) \left[ 2u(x, t)\partial_1 u(x, t) - \partial_1^2 u(x, t) \right. \\ &\quad \left. + \partial_1 \left( \int_0^{+\infty} \xi^{-\frac{1}{3}} \partial_1 u(\cdot - \xi, \cdot) d\xi \right)(x, t) \right] + q''(1)(\tau(x, t) - 1)\partial_1\tau(x, t). \end{aligned} \quad (2.8)$$

From (2.3), and (2.7), we derive that

$$\begin{aligned}
& (\tau(x, t) - 1)\partial_1\tau(x, t) \\
& \approx \left[ u(x, t) + u^2(x, t) + \int_0^{+\infty} \xi^{-\frac{1}{3}}\partial_1 u(x - \xi, t) d\xi - \partial_1 u(x, t) \right] \left[ \partial_1 u(x, t) \right. \\
& \quad \left. + 2u(x, t)\partial_1 u(x, t) - \partial_1^2 u(x, t) + \partial_1 \left( \int_0^{+\infty} \xi^{-\frac{1}{3}}\partial_1 u(\cdot - \xi, \cdot) d\xi \right)(x, t) \right] \\
& = u(x, t)\partial_1 u(x, t) + 2u^2(x, t)\partial_1 u(x, t) - u(x, t)\partial_1^2 u(x, t) \\
& \quad + u(x, t)\partial_1 \left( \int_0^{+\infty} \xi^{-\frac{1}{3}}\partial_1 u(\cdot - \xi, \cdot) d\xi \right)(x, t) + u^2(x, t)\partial_1 u(x, t) \\
& \quad + 2u^3(x, t)\partial_1 u(x, t) - u^2(x, t)\partial_1^2 u(x, t) + u^2(x, t)\partial_1 \left( \int_0^{+\infty} \xi^{-\frac{1}{3}}\partial_1 u(\cdot - \xi, \cdot) d\xi \right)(x, t) \\
& \quad + \partial_1 u(x, t) \left( \int_0^{+\infty} \xi^{-\frac{1}{3}}\partial_1 u(x - \xi, t) d\xi \right) + 2u(x, t)\partial_1 u(x, t) \left( \int_0^{+\infty} \xi^{-\frac{1}{3}}\partial_1 u(x - \xi, t) d\xi \right) \\
& \quad + \left( \int_0^{+\infty} \xi^{-\frac{1}{3}}\partial_1 u(x - \xi, t) d\xi \right) \cdot \partial_1 \left( \int_0^{+\infty} \xi^{-\frac{1}{3}}\partial_1 u(\cdot - \xi, \cdot) d\xi \right)(x, t) \\
& \quad - \partial_1^2 u(x, t) \left( \int_0^{+\infty} \xi^{-\frac{1}{3}}\partial_1 u(x - \xi, t) d\xi \right) - (\partial_1 u(x, t))^2 - 2u(x, t)(\partial_1 u(x, t))^2 \\
& \quad + \partial_1 u(x, t)\partial_1^2 u(x, t) - \partial_1 u(x, t)\partial_1 \left( \int_0^{+\infty} \xi^{-\frac{1}{3}}\partial_1 u(\cdot - \xi, \cdot) d\xi \right)(x, t). \tag{2.9}
\end{aligned}$$

By substituting (2.9) into (2.8), it follows from (2.2), and (2.5) that

$$\begin{aligned}
& \partial_2 u(x, t) + q'(1)\partial_1 u(x, t) + q'(1) \left[ 2u(x, t)\partial_1 u(x, t) - \partial_1^2 u(x, t) \right. \\
& \quad \left. + \partial_1 \left( \int_0^{+\infty} \xi^{-\frac{1}{3}}\partial_1 u(\cdot - \xi, \cdot) d\xi \right)(x, t) \right] + q''(1)u(x, t)\partial_1 u(x, t) \\
& \quad + q''(1) \left[ 3u^2(x, t)\partial_1 u(x, t) - u(x, t)\partial_1^2 u(x, t) - u^2(x, t)\partial_1^2 u(x, t) \right. \\
& \quad \left. + 2u^3(x, t)\partial_1 u(x, t) + u(x, t)\partial_1 \left( \int_0^{+\infty} \xi^{-\frac{1}{3}}\partial_1 u(\cdot - \xi, \cdot) d\xi \right)(x, t) \right]
\end{aligned}$$

$$\begin{aligned}
& + u^2(x, t) \partial_1 \left( \int_0^{+\infty} \xi^{-\frac{1}{3}} \partial_1 u(\cdot - \xi, \cdot) d\xi \right) (x, t) + \partial_1 u(x, t) \left( \int_0^{+\infty} \xi^{-\frac{1}{3}} \partial_1 u(x - \xi, t) d\xi \right) \\
& - (\partial_1 u(x, t))^2 + 2u(x, t) \partial_1 u(x, t) \left( \int_0^{+\infty} \xi^{-\frac{1}{3}} \partial_1 u(x - \xi, t) d\xi \right) \\
& + \left( \int_0^{+\infty} \xi^{-\frac{1}{3}} \partial_1 u(x - \xi, t) d\xi \right) \cdot \partial_1 \left( \int_0^{+\infty} \xi^{-\frac{1}{3}} \partial_1 u(\cdot - \xi, \cdot) d\xi \right) (x, t) \\
& - \partial_1^2 u(x, t) \left( \int_0^{+\infty} \xi^{-\frac{1}{3}} \partial_1 u(x - \xi, t) d\xi \right) - \partial_1 u(x, t) \partial_1 \left( \int_0^{+\infty} \xi^{-\frac{1}{3}} \partial_1 u(\cdot - \xi, \cdot) d\xi \right) (x, t) \\
& + \partial_1 u(x, t) \partial_1^2 u(x, t) - 2u(x, t) (\partial_1 u(x, t))^2 \Big] \approx 0. \tag{2.10}
\end{aligned}$$

By employing the Chain Rule, we obtain from (2.10) that

$$\begin{aligned}
& \partial_2 u(x, t) + q'(1) \partial_1 u(x, t) + [2q'(1) + q''(1)] u(x, t) \partial_1 u(x, t) - q'(1) \partial_1^2 u(x, t) \\
& + q'(1) \partial_1 \left( \int_0^{+\infty} \xi^{-\frac{1}{3}} \partial_1 u(\cdot - \xi, \cdot) d\xi \right) (x, t) + 2q''(1) u^3(x, t) \partial_1 u(x, t) \\
& - q''(1) u(x, t) \partial_1^2 u(x, t) + q''(1) \partial_1 u(x, t) \partial_1^2 u(x, t) - q''(1) (\partial_1 u(x, t))^2 \\
& + q''(1) \partial_1 \left[ u \left( \int_0^{+\infty} \xi^{-\frac{1}{3}} \partial_1 u(\cdot - \xi, \cdot) d\xi \right) \right] (x, t) \\
& + q''(1) \partial_1 \left[ u^2 \left( \int_0^{+\infty} \xi^{-\frac{1}{3}} \partial_1 u(\cdot - \xi, \cdot) d\xi \right) \right] (x, t) \\
& - q''(1) \partial_1 \left[ \partial_1 u \left( \int_0^{+\infty} \xi^{-\frac{1}{3}} \partial_1 u(\cdot - \xi, \cdot) d\xi \right) \right] (x, t) \\
& + q''(1) \partial_1 \left[ \frac{1}{2} \left( \int_0^{+\infty} \xi^{-\frac{1}{3}} \partial_1 u(\cdot - \xi, \cdot) d\xi \right)^2 \right] (x, t) \\
& + 3q''(1) u^2(x, t) \partial_1 u(x, t) - q''(1) u^2(x, t) \partial_1^2 u(x, t) \\
& - 2q''(1) u(x, t) (\partial_1 u(x, t))^2 \approx 0. \tag{2.11}
\end{aligned}$$

Taking into account some typical physical scales of the model (see Figure 1), namely, the amplitude,  $0 < \varepsilon \ll 1$ , and the length of the dune,  $1 \ll \delta$ , we get  $\varepsilon \ll \delta$ .




FIGURE 1. Various parameters of the physical model.

We now consider the following function

$$v(y, s) := u(\delta y, \delta s), \quad (2.12)$$

for all  $(y, s) \in \mathbb{R} \times [0, +\infty)$ . Let  $(y, s) \in \mathbb{R} \times [0, +\infty)$ . Then,

$$\partial_1 u(\delta y, \delta s) = \frac{1}{\delta} \partial_1 v(y, s), \quad (2.13)$$

$$\partial_1^2 u(\delta y, \delta s) = \frac{1}{\delta^2} \partial_1^2 v(y, s), \quad (2.14)$$

and

$$\partial_2 u(\delta y, \delta s) = \frac{1}{\delta} \partial_2 v(y, s). \quad (2.15)$$

For all  $x \in \mathbb{R}$ , we write,  $\varphi(x) := \chi_{(0,+\infty)}(x) x^{-\frac{1}{3}}$ , where  $\chi_{(0,+\infty)} : \mathbb{R} \rightarrow \mathbb{R}$  is the characteristic function of the interval  $(0, +\infty)$ . Thus,

$$\begin{aligned} \int_0^{+\infty} \xi^{-\frac{1}{3}} \partial_1 u(\delta y - \xi, \delta s) d\xi &= \int_{-\infty}^{+\infty} \chi_{(0,+\infty)}(\xi) \xi^{-\frac{1}{3}} \partial_1 u(\delta y - \xi, \delta s) d\xi \\ &= (\varphi * \partial_1 u(\cdot, \delta s))(\delta y) \\ &=: (\varphi * \partial_1 u(\cdot, \cdot))(\delta y, \delta s). \end{aligned} \quad (2.16)$$

It follows from (2.13), and (2.16) that

$$\partial_1 \left( \int_0^{+\infty} \xi^{-\frac{1}{3}} \partial_1 u(\cdot - \xi, \cdot) d\xi \right)(\delta y, \delta s)$$

$$\begin{aligned}
&:= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \left( \int_0^{+\infty} \xi^{-\frac{1}{3}} \partial_1 u(\cdot - \xi, \cdot) d\xi \right) (\delta y + h, \delta s) \right. \\
&\quad \left. - \left( \int_0^{+\infty} \xi^{-\frac{1}{3}} \partial_1 u(\cdot - \xi, \cdot) d\xi \right) (\delta y, \delta s) \right] \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \int_0^{+\infty} \xi^{-\frac{1}{3}} \partial_1 u(\delta y + h - \xi, \delta s) d\xi - \int_0^{+\infty} \xi^{-\frac{1}{3}} \partial_1 u(\delta y - \xi, \delta s) d\xi \right] \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{1}{\delta} \int_0^{+\infty} \xi^{-\frac{1}{3}} \partial_1 v \left( y + \frac{h}{\delta} - \frac{\xi}{\delta}, s \right) d\xi - \frac{1}{\delta} \int_0^{+\infty} \xi^{-\frac{1}{3}} \partial_1 v \left( y - \frac{\xi}{\delta}, s \right) d\xi \right] \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \frac{1}{\delta^{\frac{1}{3}}} \int_0^{+\infty} \zeta^{-\frac{1}{3}} \partial_1 v \left( y + \frac{h}{\delta} - \zeta, s \right) d\zeta - \frac{1}{\delta^{\frac{1}{3}}} \int_0^{+\infty} \zeta^{-\frac{1}{3}} \partial_1 v(y - \zeta, s) d\zeta \right] \\
&= \frac{1}{\delta^{\frac{4}{3}}} \lim_{h \rightarrow 0} \frac{1}{h} \left[ \int_0^{+\infty} \zeta^{-\frac{1}{3}} \partial_1 v \left( y + \frac{h}{\delta} - \zeta, s \right) d\zeta - \int_0^{+\infty} \zeta^{-\frac{1}{3}} \partial_1 v(y - \zeta, s) d\zeta \right] \\
&= \frac{1}{\delta^{\frac{4}{3}}} \lim_{a \rightarrow 0} \frac{1}{a} \left[ \int_0^{+\infty} \zeta^{-\frac{1}{3}} \partial_1 v(y + a - \zeta, s) d\zeta - \int_0^{+\infty} \zeta^{-\frac{1}{3}} \partial_1 v(y - \zeta, s) d\zeta \right] \\
&= \frac{1}{\delta^{\frac{4}{3}}} \lim_{a \rightarrow 0} \frac{1}{a} \left[ \left( \int_0^{+\infty} \zeta^{-\frac{1}{3}} \partial_1 v(\cdot - \zeta, \cdot) d\zeta \right)(y + a, s) - \left( \int_0^{+\infty} \zeta^{-\frac{1}{3}} \partial_1 v(\cdot - \zeta, \cdot) d\zeta \right)(y, s) \right] \\
&=: \frac{1}{\delta^{\frac{4}{3}}} \partial_1 \left( \int_0^{+\infty} \zeta^{-\frac{1}{3}} \partial_1 v(\cdot - \zeta, \cdot) d\zeta \right)(y, s) \\
&=: \frac{1}{\delta^{\frac{4}{3}}} \partial_1 (\varphi * \partial_1 v(\cdot, \cdot))(y, s).
\end{aligned} \tag{2.17}$$

By using (2.13), (2.12), and (2.17), we have that

$$\begin{aligned}
&\partial_1 \left[ u \left( \int_0^{+\infty} \xi^{-\frac{1}{3}} \partial_1 u(\cdot - \xi, \cdot) d\xi \right) \right] (\delta y, \delta s) \\
&= \partial_1 u(\delta y, \delta s) \left( \int_0^{+\infty} \xi^{-\frac{1}{3}} \partial_1 u(\delta y - \xi, \delta s) d\xi \right) \\
&\quad + u(\delta y, \delta s) \partial_1 \left( \int_0^{+\infty} \xi^{-\frac{1}{3}} \partial_1 u(\cdot - \xi, \cdot) d\xi \right) (\delta y, \delta s)
\end{aligned}$$

$$\begin{aligned}
&= \frac{1}{\delta^2} \partial_1 v(y, s) \left( \int_0^{+\infty} \xi^{-\frac{1}{3}} \partial_1 v \left( y - \frac{\xi}{\delta}, s \right) d\xi \right) \\
&\quad + \frac{1}{\delta^{\frac{4}{3}}} v(y, s) \partial_1 \left( \int_0^{+\infty} \zeta^{-\frac{1}{3}} \partial_1 v(\cdot - \zeta, \cdot) d\zeta \right) (y, s) \\
&= \frac{1}{\delta^{\frac{4}{3}}} \partial_1 v(y, s) \left( \int_0^{+\infty} \zeta^{-\frac{1}{3}} \partial_1 v(y - \zeta, s) d\zeta \right) \\
&\quad + \frac{1}{\delta^{\frac{4}{3}}} v(y, s) \partial_1 \left( \int_0^{+\infty} \zeta^{-\frac{1}{3}} \partial_1 v(\cdot - \zeta, \cdot) d\zeta \right) (y, s) \\
&= \frac{1}{\delta^{\frac{4}{3}}} \partial_1 \left[ v \cdot (\varphi * \partial_1 v(\cdot, \cdot)) \right] (y, s), \tag{2.18}
\end{aligned}$$

$$\begin{aligned}
&\partial_1 \left[ u^2 \left( \int_0^{+\infty} \xi^{-\frac{1}{3}} \partial_1 u(\cdot - \xi, \cdot) d\xi \right) \right] (\delta y, \delta s) \\
&= 2u(\delta y, \delta s) \partial_1 u(\delta y, \delta s) \left( \int_0^{+\infty} \xi^{-\frac{1}{3}} \partial_1 u(\delta y - \xi, \delta s) d\xi \right) \\
&\quad + u^2(\delta y, \delta s) \partial_1 \left( \int_0^{+\infty} \xi^{-\frac{1}{3}} \partial_1 u(\cdot - \xi, \cdot) d\xi \right) (\delta y, \delta s) \\
&= \frac{2}{\delta^2} v(y, s) \partial_1 v(y, s) \left( \int_0^{+\infty} \xi^{-\frac{1}{3}} \partial_1 v \left( y - \frac{\xi}{\delta}, s \right) d\xi \right) \\
&\quad + \frac{1}{\delta^{\frac{4}{3}}} v^2(y, s) \partial_1 \left( \int_0^{+\infty} \zeta^{-\frac{1}{3}} \partial_1 v(\cdot - \zeta, \cdot) d\zeta \right) (y, s) \\
&= \frac{1}{\delta^{\frac{4}{3}}} 2v(y, s) \partial_1 v(y, s) \left( \int_0^{+\infty} \zeta^{-\frac{1}{3}} \partial_1 v(y - \zeta, s) d\zeta \right) \\
&\quad + \frac{1}{\delta^{\frac{4}{3}}} v^2(y, s) \partial_1 \left( \int_0^{+\infty} \zeta^{-\frac{1}{3}} \partial_1 v(\cdot - \zeta, \cdot) d\zeta \right) (y, s) \\
&= \frac{1}{\delta^{\frac{4}{3}}} \partial_1 \left[ v^2 \cdot (\varphi * \partial_1 v(\cdot, \cdot)) \right] (y, s), \tag{2.19}
\end{aligned}$$

and

$$\begin{aligned}
& \partial_1 \left[ \frac{1}{2} \left( \int_0^{+\infty} \xi^{-\frac{1}{3}} \partial_1 u(\cdot - \xi, \cdot) d\xi \right)^2 \right] (\delta y, \delta s) \\
&= \left( \int_0^{+\infty} \xi^{-\frac{1}{3}} \partial_1 u(\delta y - \xi, \delta s) d\xi \right) \cdot \partial_1 \left( \int_0^{+\infty} \xi^{-\frac{1}{3}} \partial_1 u(\cdot - \xi, \cdot) d\xi \right) (\delta y, \delta s) \\
&= \frac{1}{\delta^{\frac{7}{3}}} \left( \int_0^{+\infty} \xi^{-\frac{1}{3}} \partial_1 v \left( y - \frac{\xi}{\delta}, s \right) d\xi \right) \cdot \partial_1 \left( \int_0^{+\infty} \zeta^{-\frac{1}{3}} \partial_1 v(\cdot - \zeta, \cdot) d\zeta \right) (y, s) \\
&= \frac{1}{\delta^{\frac{5}{3}}} \left( \int_0^{+\infty} \zeta^{-\frac{1}{3}} \partial_1 v(y - \zeta, s) d\zeta \right) \cdot \partial_1 \left( \int_0^{+\infty} \zeta^{-\frac{1}{3}} \partial_1 v(\cdot - \zeta, \cdot) d\zeta \right) (y, s) \\
&= \frac{1}{\delta^{\frac{5}{3}}} \partial_1 \left[ \frac{1}{2} (\varphi * \partial_1 v(\cdot, \cdot))^2 \right] (y, s).
\end{aligned} \tag{2.20}$$

Moreover, by employing (2.14), (2.13), and (2.17), we get

$$\begin{aligned}
& \partial_1 \left[ \partial_1 u \left( \int_0^{+\infty} \xi^{-\frac{1}{3}} \partial_1 u(\cdot - \xi, \cdot) d\xi \right) \right] (\delta y, \delta s) \\
&= \partial_1^2 u(\delta y, \delta s) \left( \int_0^{+\infty} \xi^{-\frac{1}{3}} \partial_1 u(\delta y - \xi, \delta s) d\xi \right) \\
&\quad + \partial_1 u(\delta y, \delta s) \partial_1 \left( \int_0^{+\infty} \xi^{-\frac{1}{3}} \partial_1 u(\cdot - \xi, \cdot) d\xi \right) (\delta y, \delta s) \\
&= \frac{1}{\delta^3} \partial_1^2 v(y, s) \left( \int_0^{+\infty} \xi^{-\frac{1}{3}} \partial_1 v \left( y - \frac{\xi}{\delta}, s \right) d\xi \right) \\
&\quad + \frac{1}{\delta^{\frac{7}{3}}} \partial_1 v(y, s) \partial_1 \left( \int_0^{+\infty} \zeta^{-\frac{1}{3}} \partial_1 v(\cdot - \zeta, \cdot) d\zeta \right) (y, s) \\
&= \frac{1}{\delta^{\frac{7}{3}}} \partial_1^2 v(y, s) \left( \int_0^{+\infty} \zeta^{-\frac{1}{3}} \partial_1 v(y - \zeta, s) d\zeta \right) \\
&\quad + \frac{1}{\delta^{\frac{7}{3}}} \partial_1 v(y, s) \partial_1 \left( \int_0^{+\infty} \zeta^{-\frac{1}{3}} \partial_1 v(\cdot - \zeta, \cdot) d\zeta \right) (y, s) \\
&= \frac{1}{\delta^{\frac{7}{3}}} \partial_1 \left[ \partial_1 v \cdot (\varphi * \partial_1 v(\cdot, \cdot)) \right] (y, s).
\end{aligned} \tag{2.21}$$

From (2.15), (2.13), (2.14), (2.12), (2.17), and (2.18)-(2.21), the approximation (2.11) may be rewritten as

$$\begin{aligned} & \frac{1}{\delta} \partial_2 v(y, s) + \frac{q'(1)}{\delta} \partial_1 v(y, s) + \frac{[2q'(1) + q''(1)]}{\delta} v(y, s) \partial_1 v(y, s) - \frac{q'(1)}{\delta^2} \partial_1^2 v(y, s) \\ & + \frac{q'(1)}{\delta^{\frac{4}{3}}} \partial_1 (\varphi * \partial_1 v(\cdot, \cdot))(y, s) + \frac{3q''(1)}{\delta} v^2(y, s) \partial_1 v(y, s) + \frac{2q''(1)}{\delta} v^3(y, s) \partial_1 v(y, s) \\ & - \frac{q''(1)}{\delta^2} v(y, s) \partial_1^2 v(y, s) - \frac{q''(1)}{\delta^2} v^2(y, s) \partial_1^2 v(y, s) + \frac{q''(1)}{\delta^3} \partial_1 v(y, s) \partial_1^2 v(y, s) \\ & - \frac{q''(1)}{\delta^2} (\partial_1 v(y, s))^2 - \frac{2q''(1)}{\delta^2} v(y, s) (\partial_1 v(y, s))^2 \\ & + \frac{q''(1)}{\delta^{\frac{4}{3}}} \partial_1 \left[ v \cdot (\varphi * \partial_1 v(\cdot, \cdot)) \right](y, s) + \frac{q''(1)}{\delta^{\frac{4}{3}}} \partial_1 \left[ v^2 \cdot (\varphi * \partial_1 v(\cdot, \cdot)) \right](y, s) \\ & - \frac{q''(1)}{\delta^{\frac{7}{3}}} \partial_1 \left[ \partial_1 v \cdot (\varphi * \partial_1 v(\cdot, \cdot)) \right](y, s) + \frac{q''(1)}{\delta^{\frac{5}{3}}} \partial_1 \left[ \frac{1}{2} (\varphi * \partial_1 v(\cdot, \cdot))^2 \right](y, s) \approx 0. \end{aligned} \quad (2.22)$$

Now, we define the function

$$w(z, r) := \frac{1}{\varepsilon} v(z, r), \quad (2.23)$$

for all  $(z, r) \in \mathbb{R} \times [0, +\infty)$ . Let  $(z, r) \in \mathbb{R} \times [0, +\infty)$ . It follows from the approximation (2.22) that

$$\begin{aligned} & \frac{\varepsilon}{\delta} \partial_2 w(z, r) + \frac{q'(1)\varepsilon}{\delta} \partial_1 w(z, r) + \frac{[2q'(1) + q''(1)]\varepsilon^2}{\delta} w(z, r) \partial_1 w(z, r) - \frac{q'(1)\varepsilon}{\delta^2} \partial_1^2 w(z, r) \\ & + \frac{q'(1)\varepsilon}{\delta^{\frac{4}{3}}} \partial_1 (\varphi * \partial_1 w(\cdot, \cdot))(z, r) + \frac{3q''(1)\varepsilon^3}{\delta} w^2(z, r) \partial_1 w(z, r) \\ & + \frac{2q''(1)\varepsilon^4}{\delta} w^3(z, r) \partial_1 w(z, r) - \frac{q''(1)\varepsilon^2}{\delta^2} w(z, r) \partial_1^2 w(z, r) - \frac{q''(1)\varepsilon^3}{\delta^2} w^2(z, r) \partial_1^2 w(z, r) \\ & + \frac{q''(1)\varepsilon^2}{\delta^3} \partial_1 w(z, r) \partial_1^2 w(z, r) - \frac{q''(1)\varepsilon^2}{\delta^2} (\partial_1 w(z, r))^2 - \frac{2q''(1)\varepsilon^3}{\delta^2} w(z, r) (\partial_1 w(z, r))^2 \\ & + \frac{q''(1)\varepsilon^2}{\delta^{\frac{4}{3}}} \partial_1 \left[ w \cdot (\varphi * \partial_1 w(\cdot, \cdot)) \right](z, r) + \frac{q''(1)\varepsilon^3}{\delta^{\frac{4}{3}}} \partial_1 \left[ w^2 \cdot (\varphi * \partial_1 w(\cdot, \cdot)) \right](z, r) \\ & - \frac{q''(1)\varepsilon^2}{\delta^{\frac{7}{3}}} \partial_1 \left[ \partial_1 w \cdot (\varphi * \partial_1 w(\cdot, \cdot)) \right](z, r) + \frac{q''(1)\varepsilon^2}{\delta^{\frac{5}{3}}} \partial_1 \left[ \frac{1}{2} (\varphi * \partial_1 w(\cdot, \cdot))^2 \right](z, r) \approx 0. \end{aligned}$$

Hence,

$$\begin{aligned} & \partial_2 w(z, r) + q'(1) \partial_1 w(z, r) + [2q'(1) + q''(1)]\varepsilon w(z, r) \partial_1 w(z, r) - \frac{q'(1)}{\delta} \partial_1^2 w(z, r) \\ & + \frac{q'(1)}{\delta^{\frac{1}{3}}} \partial_1 (\varphi * \partial_1 w(\cdot, \cdot))(z, r) + 3q''(1)\varepsilon^2 w^2(z, r) \partial_1 w(z, r) \\ & + 2q''(1)\varepsilon^3 w^3(z, r) \partial_1 w(z, r) - \frac{q''(1)\varepsilon}{\delta} w(z, r) \partial_1^2 w(z, r) - \frac{q''(1)\varepsilon^2}{\delta} w^2(z, r) \partial_1^2 w(z, r) \end{aligned}$$

$$\begin{aligned}
& + \frac{q''(1)\varepsilon}{\delta^2} \partial_1 w(z, r) \partial_1^2 w(z, r) - \frac{q''(1)\varepsilon}{\delta} (\partial_1 w(z, r))^2 - \frac{2q''(1)\varepsilon^2}{\delta} w(z, r) (\partial_1 w(z, r))^2 \\
& + \frac{q''(1)\varepsilon}{\delta^{\frac{1}{3}}} \partial_1 \left[ w \cdot (\varphi * \partial_1 w(\cdot, \cdot)) \right] (z, r) + \frac{q''(1)\varepsilon^2}{\delta^{\frac{1}{3}}} \partial_1 \left[ w^2 \cdot (\varphi * \partial_1 w(\cdot, \cdot)) \right] (z, r) \\
& - \frac{q''(1)\varepsilon}{\delta^{\frac{4}{3}}} \partial_1 \left[ \partial_1 w \cdot (\varphi * \partial_1 w(\cdot, \cdot)) \right] (z, r) \\
& + \frac{q''(1)\varepsilon}{\delta^{\frac{2}{3}}} \partial_1 \left[ \frac{1}{2} (\varphi * \partial_1 w(\cdot, \cdot))^2 \right] (z, r) \approx 0. \tag{2.24}
\end{aligned}$$

In what follows, we neglect the terms of the form  $O\left(\frac{\varepsilon}{\delta^{\frac{1}{3}}}\right)$ , as  $\varepsilon \rightarrow 0^+$ , and  $\delta \rightarrow +\infty$ , more precisely, we remove the terms containing the following factors

$$\varepsilon^2, \quad \varepsilon^3, \quad \frac{\varepsilon}{\delta}, \quad \frac{\varepsilon^2}{\delta}, \quad \frac{\varepsilon}{\delta^2}, \quad \frac{\varepsilon^2}{\delta^{\frac{1}{3}}}, \quad \frac{\varepsilon}{\delta^{\frac{4}{3}}}, \quad \text{and} \quad \frac{\varepsilon}{\delta^{\frac{2}{3}}}, \tag{2.25}$$

when  $\varepsilon \rightarrow 0^+$ , and  $\delta \rightarrow +\infty$ . Therefore,

$$\begin{aligned}
& \partial_2 w(z, r) + q'(1) \partial_1 w(z, r) - \frac{q'(1)}{\delta} \partial_1^2 w(z, r) + [2q'(1) + q''(1)] \varepsilon w(z, r) \partial_1 w(z, r) \\
& + \frac{q'(1)}{\delta^{\frac{1}{3}}} \partial_1 (\varphi * \partial_1 w(\cdot, \cdot)) (z, r) + q''(1) \frac{\varepsilon}{\delta^{\frac{1}{3}}} \partial_1 \left[ w \cdot (\varphi * \partial_1 w(\cdot, \cdot)) \right] (z, r) \approx 0. \tag{2.26}
\end{aligned}$$

The bedload transport function, proposed by Meyer-Peter, and Müller ([2, 3, 4]), is given by

$$q(\tau) = C ([\tau - \tau_c]_+)^{\frac{3}{2}}, \tag{2.27}$$

for all  $\tau \in \mathbb{R}$ , where  $C > 0$ ,  $[x]_+ = \max\{x, 0\}$ , for all  $x \in \mathbb{R}$ , and  $\tau_c = 0.047$  is a yield stress, called the *Shields stress*.

**Lemma 2.1.** *Let  $q : \mathbb{R} \rightarrow \mathbb{R}$  be the function given by (2.27). Then,  $q'(1) \neq 0$ .*

*Proof.* We see that

$$\begin{aligned}
q'_+(\tau_c) &= \lim_{h \rightarrow 0^+} \frac{q(\tau_c + h) - q(\tau_c)}{h} = \lim_{h \rightarrow 0^+} \frac{C h^{\frac{3}{2}}}{h} \\
&= C \lim_{h \rightarrow 0^+} h^{\frac{1}{2}} = 0,
\end{aligned}$$

and

$$q'_-(\tau_c) = \lim_{k \rightarrow 0^-} \frac{q(\tau_c + k) - q(\tau_c)}{k} = \lim_{k \rightarrow 0^-} \frac{0}{k} = 0.$$

So, there exists

$$q'(\tau_c) := \lim_{l \rightarrow 0} \frac{q(\tau_c + l) - q(\tau_c)}{l} = 0.$$

Therefore,

$$q'(\tau) = \begin{cases} \frac{3}{2}C(\tau - \tau_c)^{\frac{1}{2}}, & \text{if } \tau > \tau_c, \\ 0, & \text{if } \tau \leq \tau_c, \end{cases} =: \frac{3}{2}C([\tau - \tau_c]_+)^{\frac{1}{2}}. \quad (2.28)$$

□

We now consider the function

$$\phi(e, l) := w\left(\frac{e}{\delta} + \frac{l}{\delta}, \frac{l}{q'(1)\delta}\right), \quad (2.29)$$

for all  $(e, l) \in \mathbb{R} \times [0, +\infty)$ . Let  $(e, l) \in \mathbb{R} \times [0, +\infty)$ . By using (2.29), and the Chain Rule, we get

$$\partial_1 w\left(\frac{e}{\delta} + \frac{l}{\delta}, \frac{l}{q'(1)\delta}\right) = \delta \partial_1 \phi(e, l), \quad (2.30)$$

$$\partial_1^2 w\left(\frac{e}{\delta} + \frac{l}{\delta}, \frac{l}{q'(1)\delta}\right) = \delta^2 \partial_1^2 \phi(e, l), \quad (2.31)$$

and

$$\partial_2 w\left(\frac{e}{\delta} + \frac{l}{\delta}, \frac{l}{q'(1)\delta}\right) = q'(1) \delta \partial_2 \phi(e, l) - q'(1) \delta \partial_1 \phi(e, l). \quad (2.32)$$

From (2.30), we obtain that

$$\begin{aligned} (\varphi * \partial_1 w(\cdot, \cdot))\left(\frac{e}{\delta} + \frac{l}{\delta}, \frac{l}{q'(1)\delta}\right) &:= \left( \int_0^{+\infty} \zeta^{-\frac{1}{3}} \partial_1 w(\cdot - \zeta, \cdot) d\zeta \right) \left(\frac{e}{\delta} + \frac{l}{\delta}, \frac{l}{q'(1)\delta}\right) \\ &= \int_0^{+\infty} \zeta^{-\frac{1}{3}} \partial_1 w\left(\frac{e}{\delta} + \frac{l}{\delta} - \zeta, \frac{l}{q'(1)\delta}\right) d\zeta \\ &= \delta \int_0^{+\infty} \zeta^{-\frac{1}{3}} \partial_1 \phi(e - \delta\zeta, l) d\zeta \\ &= \delta^{\frac{1}{3}} \int_0^{+\infty} \mu^{-\frac{1}{3}} \partial_1 \phi(e - \mu, l) d\mu \\ &=: \delta^{\frac{1}{3}} (\varphi * \partial_1 \phi(\cdot, \cdot))(e, l), \end{aligned} \quad (2.33)$$

and

$$\begin{aligned}
& \partial_1 (\varphi * \partial_1 w(\cdot, \cdot)) \left( \frac{e}{\delta} + \frac{l}{\delta}, \frac{l}{q'(1)\delta} \right) \\
&:= \partial_1 \left( \int_0^{+\infty} \zeta^{-\frac{1}{3}} \partial_1 w(\cdot - \zeta, \cdot) d\zeta \right) \left( \frac{e}{\delta} + \frac{l}{\delta}, \frac{l}{q'(1)\delta} \right) \\
&:= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \int_0^{+\infty} \zeta^{-\frac{1}{3}} \partial_1 w \left( \frac{e}{\delta} + \frac{l}{\delta} + h - \zeta, \frac{l}{q'(1)\delta} \right) d\zeta \right. \\
&\quad \left. - \int_0^{+\infty} \zeta^{-\frac{1}{3}} \partial_1 w \left( \frac{e}{\delta} + \frac{l}{\delta} - \zeta, \frac{l}{q'(1)\delta} \right) d\zeta \right] \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \delta \int_0^{+\infty} \zeta^{-\frac{1}{3}} \partial_1 \phi(e + \delta h - \delta \zeta, l) d\zeta - \delta \int_0^{+\infty} \zeta^{-\frac{1}{3}} \partial_1 \phi(e - \delta \zeta, l) d\zeta \right] \\
&= \lim_{h \rightarrow 0} \frac{1}{h} \left[ \delta^{\frac{1}{3}} \int_0^{+\infty} \mu^{-\frac{1}{3}} \partial_1 \phi(e + \delta h - \mu, l) d\mu - \delta^{\frac{1}{3}} \int_0^{+\infty} \mu^{-\frac{1}{3}} \partial_1 \phi(e - \mu, l) d\mu \right] \\
&= \delta^{\frac{4}{3}} \lim_{h \rightarrow 0} \frac{1}{\delta h} \left[ \int_0^{+\infty} \mu^{-\frac{1}{3}} \partial_1 \phi(e + \delta h - \mu, l) d\mu - \int_0^{+\infty} \mu^{-\frac{1}{3}} \partial_1 \phi(e - \mu, l) d\mu \right] \\
&= \delta^{\frac{4}{3}} \lim_{k \rightarrow 0} \frac{1}{k} \left[ \int_0^{+\infty} \mu^{-\frac{1}{3}} \partial_1 \phi(e + k - \mu, l) d\mu - \int_0^{+\infty} \mu^{-\frac{1}{3}} \partial_1 \phi(e - \mu, l) d\mu \right] \\
&=: \delta^{\frac{4}{3}} \partial_1 \left( \int_0^{+\infty} \mu^{-\frac{1}{3}} \partial_1 \phi(\cdot - \mu, \cdot) d\mu \right) (e, l) \\
&=: \delta^{\frac{4}{3}} \partial_1 (\varphi * \partial_1 \phi(\cdot, \cdot))(e, l).
\end{aligned} \tag{2.34}$$

By employing (2.30), (2.33), (2.29), and (2.34), we see that

$$\begin{aligned}
& \partial_1 \left[ w \cdot (\varphi * \partial_1 w(\cdot, \cdot)) \right] \left( \frac{e}{\delta} + \frac{l}{\delta}, \frac{l}{q'(1)\delta} \right) \\
&= \partial_1 w \left( \frac{e}{\delta} + \frac{l}{\delta}, \frac{l}{q'(1)\delta} \right) (\varphi * \partial_1 w(\cdot, \cdot)) \left( \frac{e}{\delta} + \frac{l}{\delta}, \frac{l}{q'(1)\delta} \right) + w \left( \frac{e}{\delta} + \frac{l}{\delta}, \frac{l}{q'(1)\delta} \right) \\
&\quad \cdot \partial_1 (\varphi * \partial_1 w(\cdot, \cdot)) \left( \frac{e}{\delta} + \frac{l}{\delta}, \frac{l}{q'(1)\delta} \right) \\
&= \delta^{\frac{4}{3}} \partial_1 \phi(e, l) (\varphi * \partial_1 \phi(\cdot, \cdot))(e, l) + \delta^{\frac{4}{3}} \phi(e, l) \partial_1 (\varphi * \partial_1 \phi(\cdot, \cdot))(e, l) \\
&= \delta^{\frac{4}{3}} \partial_1 \left[ \phi \cdot (\varphi * \partial_1 \phi(\cdot, \cdot)) \right] (e, l).
\end{aligned} \tag{2.35}$$

By using (2.29)-(2.32), (2.34), and (2.35), we get from (2.26) that

$$\begin{aligned} & q'(1) \delta \partial_2 \phi(e, l) - q'(1) \delta \partial_1 \phi(e, l) + q'(1) \delta \partial_1 \phi(e, l) - q'(1) \delta \partial_1^2 \phi(e, l) \\ & + [2q'(1) + q''(1)] \varepsilon \delta \phi(e, l) \partial_1 \phi(e, l) + q'(1) \delta \partial_1 (\varphi * \partial_1 \phi(\cdot, \cdot))(e, l) \\ & + q''(1) \varepsilon \delta \partial_1 \left[ \phi \cdot (\varphi * \partial_1 \phi(\cdot, \cdot)) \right](e, l) \approx 0. \end{aligned}$$

So,

$$\begin{aligned} & \partial_2 \phi(e, l) - \partial_1^2 \phi(e, l) + \frac{[2q'(1) + q''(1)]\varepsilon}{q'(1)} \phi(e, l) \partial_1 \phi(e, l) + \partial_1 (\varphi * \partial_1 \phi(\cdot, \cdot))(e, l) \\ & + \frac{q''(1)\varepsilon}{q'(1)} \partial_1 \left[ \phi \cdot (\varphi * \partial_1 \phi(\cdot, \cdot)) \right](e, l) \approx 0. \end{aligned} \quad (2.36)$$

It follows from (2.28) that

$$\frac{q'(1)}{2q'(1) + q''(1)} = \frac{1.4295}{3(0.953) + 0.75} \approx 0.3961.$$

We now define the function

$$\psi(x, t) := \beta \phi(x, t), \quad (2.37)$$

for all  $(x, t) \in \mathbb{R} \times [0, +\infty)$ , where

$$\beta := \frac{[2q'(1) + q''(1)]}{q'(1)} \cdot \varepsilon.$$

From the approximation (2.36), for all  $(x, t) \in \mathbb{R} \times [0, +\infty)$ ,

$$\begin{aligned} & \frac{1}{\beta} \partial_2 \psi(x, t) - \frac{1}{\beta} \partial_1^2 \psi(x, t) + \frac{1}{\beta} \psi(x, t) \partial_1 \psi(x, t) + \frac{1}{\beta} \partial_1 (\varphi * \partial_1 \psi(\cdot, \cdot))(x, t) \\ & + \frac{q''(1)\varepsilon}{q'(1)\beta^2} \partial_1 \left[ \psi \cdot (\varphi * \partial_1 \psi(\cdot, \cdot)) \right](x, t) \approx 0. \end{aligned}$$

We denote

$$\eta := \frac{q''(1)}{2q'(1) + q''(1)}. \quad (2.38)$$

Finally, we obtain the key result of this paper

$$\partial_2 \psi(x, t) + \partial_1 \left[ \frac{\psi^2}{2} - \partial_1 \psi + (\varphi * \partial_1 \psi(\cdot, \cdot)) + \eta \psi \cdot (\varphi * \partial_1 \psi(\cdot, \cdot)) \right](x, t) \approx 0, \quad (2.39)$$

for all  $(x, t) \in \mathbb{R} \times [0, +\infty)$ .

## REFERENCES

- [1] B. Álvarez-Samaniego, and P. Azerad, *Existence of travelling-wave solutions and local well-posedness of the Fowler equation*, Discrete and Continuous Dynamical Systems. Series B. A Journal Bridging Mathematics and Sciences **12** (2009), No. 4, 671-692.
- [2] A. C. Fowler, *Dunes and drumlins*, in “Geomorphological Fluid Mechanics” (eds. N. J. Balmforth, and A. Provenzale), Springer-Verlag, (2001), 430-454.

- [3] A. C. Fowler, *Evolution equations for dunes and drumlins*, Mathematics and Environment (Spanish) (Paris, 2002), RACSAM Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. **96** (2002), No. 3, 377-387.
- [4] A. C. Fowler, *Mathematics and the Environment*, Lecture Notes, 2004, (<https://people.maths.ox.ac.uk/fowler/courses/mathenvo.html>).