

Noncommutative differential geometry on crossed product algebras

Andrea Sciandra^{*,1} and Thomas Weber^{†,2,3}

¹*Dipartimento di Matematica "Giuseppe Peano", Università degli Studi di Torino
Via Carlo Alberto 10, 10123 Torino, Italy*

²*Dipartimento di Matematica, Università di Bologna
Piazza di Porta S. Donato 5, I-40126 Bologna, Italy*

³*Istituto Nazionale di Fisica Nucleare, Sezione di Bologna
Via Bertini Pichat 6/2, I-40127 Bologna, Italy*

Abstract

We provide a differential structure on arbitrary cleft extensions $B := A^{\text{co}H} \subseteq A$ for an H -comodule algebra A . This is achieved by constructing a covariant calculus on the corresponding crossed product algebra $B \#_{\sigma} H$ from the data of a bicovariant calculus on the structure Hopf algebra H and a calculus on the base algebra B , which is compatible with the 2-cocycle and measure of the crossed product. The result is a quantum principal bundle with canonical strong connection and we describe the induced bimodule covariant derivatives on associated bundles of the crossed product. It is proven that connections of the quantum principal bundle are in bijection with connection 1-forms. All results specialize to trivial extensions and smash product algebras $B \# H$ and we give a characterization of the smash product calculus in terms of the differentials of the cleaving map $j: H \rightarrow A$ and the inclusion $B \hookrightarrow A$. The construction is exemplified for pointed Hopf algebras. In particular, the case of Radford Hopf algebras $H_{(r,n,q)}$ is spelled out in full detail.

Contents

1	Introduction	2
2	Preliminaries	3
2.1	Equivariant and covariant modules	3
2.2	Crossed product algebras	4
2.3	Noncommutative differential calculi	7
2.4	Quantum principal bundles and connections	9
3	The crossed product calculus	12
3.1	Crossed product 1-forms	12
3.2	A classification of the smash product calculus	16
3.3	Higher order forms	19
3.4	Quantum principal bundle and connections	21
3.5	Connection 1-forms and fundamental vector fields	25
4	Construction for pointed Hopf algebras	27
5	References	30

*andrea.sciandra@unito.it

†thomasmartin.weber@unibo.it

1 Introduction

Without doubt, principal bundles play a decisive role in differential geometry, topology and gauge theory. An ingenious generalization to quantum principal bundles on noncommutative algebras was proposed by Schneider [23] and Brzeziński–Majid [7]. The central idea is to employ Hopf–Galois extensions [14], where an H -comodule algebra A encodes the total space, the Hopf algebra H replaces the structure group and the base is given by the subalgebra $B = A^{\text{co}H} \subseteq A$ of H -coinvariant elements. Principality of such a quantum bundle is then characterized by the invertibility of the Hopf–Galois map $A \otimes_B A \rightarrow A \otimes H$. In order to ensure an appropriate geometric behaviour one commonly assumes faithful flatness of the Hopf–Galois extension, thus resulting in a categorical equivalence of B -modules and A - H -Hopf modules [23]. Throughout the last decades the notion of quantum principal bundle has been featured in numerous articles [8, 12, 15], just to name a few, establishing it as common knowledge in noncommutative differential geometry.

A class of particular interest is given by cleft extensions, i.e., H -comodule algebras A which admit a cleaving map $j: H \rightarrow A$. The latter is a convolution invertible comodule map and it provides an explicit inverse of the Hopf–Galois map. It was shown by Doi–Takeuchi [10] that cleft extensions are precisely the Hopf–Galois extensions with the normal basis property, meaning that A is isomorphic (as comodule algebra) to the crossed product algebra $B \#_{\sigma} H$ with $B = A^{\text{co}H}$ and, conversely, every crossed product algebra gives rise to a cleft extension. A crossed product algebra is determined by a 2-cocycle $\sigma: H \otimes H \rightarrow B$ and a measure (this is a weak version of an action) of H on a σ -twisted H -module algebra B . In case of the trivial 2-cocycle we recover smash product algebras $B \# H$, which correspond to trivial extensions $B = A^{\text{co}H} \subseteq A$. The isomorphism $A \cong B \#_{\sigma} H$ provides a certain triviality of the bundle, in the sense that the total space algebra can be expressed through the base algebra and structure Hopf algebra, however, we would like to stress that the algebra structure of $B \#_{\sigma} H$ is nontrivial, as it involves the measure and 2-cocycle. In particular, $B \#_{\sigma} H$ is in general not isomorphic to the tensor product algebra $B \otimes H$. It follows that cleft extensions are automatically faithfully flat if the antipode of the Hopf algebra is invertible and one considers vector spaces over a base field [6].

While in classical differential geometry there is a canonical differential structure on every manifold, namely the de Rham calculus, in noncommutative geometry there are typically many differential calculi on a given noncommutative algebra. For this reason, a noncommutative differential structure requires additional information, for example a spectral triple in the framework of noncommutative geometry [9], or an ideal in the kernel of the counit in case of a covariant calculus on a Hopf algebra [26]. The main contribution of this paper is to provide a canonical covariant calculus on arbitrary cleft extensions $B = A^{\text{co}H} \subseteq A$ (or equivalently on arbitrary crossed product algebras $B \#_{\sigma} H$) from a bicovariant calculus $(\Omega^1(H), d_H)$ on the structure Hopf algebra H and a calculus $(\Omega^1(B), d_B)$ on the base algebra B , which is compatible with the 2-cocycle and measure of the crossed product. We call it the crossed product calculus. The construction (Section 3.1) is in the spirit of the normal basis property, where the 1-forms $\Omega^1(A)$ on A are build as the direct sum $(\Omega^1(B) \otimes H) \oplus (B \otimes \Omega^1(H))$ and the A -bimodule actions mimic the crossed product multiplication; the differential is the sum $d_A = d_B + d_H$. In a similar way, higher order forms are constructed (Section 3.3) and we prove that the crossed product calculus forms a quantum principal bundle (Section 3.4) in the sense of [7], meaning that there is a short exact sequence

$$0 \longrightarrow \Omega^1(B) \otimes H \hookrightarrow \Omega^1(A) \xrightarrow{\text{ver}} A \otimes {}^{\text{co}H}\Omega^1(H) \longrightarrow 0,$$

where ver denotes the vertical map and ${}^{\text{co}H}\Omega^1(H)$ are the left H -coinvariant 1-forms on H . This allows us to identify $\Omega^1(B) \otimes H$ as the horizontal forms and $B \otimes \Omega^1(H)$ as the vertical forms of the quantum principal bundle. It follows that the quantum principal bundle is strong (i.e., $A\Omega^1(B) = \Omega^1(B)A$) and regular in the sense of [3] (it descends from the universal quantum principal bundle with strong connection). We describe the induced bimodule covariant derivatives on all associated bundles. As an application, we construct higher order vertical maps and exact sequences for the crossed product calculus and show that the noncommutative de Rham cohomology $H_{\text{dR}}^{\bullet}(A)$ can be computed from the tensor product of cohomologies $H_{\text{dR}}^{\bullet}(B) \otimes H_{\text{dR}}^{\bullet}(H)$ via the Leray–Serre spectral sequence. We further give a bijection between connections and connection 1-forms on the crossed product calculus (Section 3.5), using the notion of fundamental vector field.

For the convenience of the reader we include preliminaries (Section 2), recalling the notions of Hopf algebra theory and noncommutative differential geometry which are relevant for the crossed product calculus, rendering the presentation self-contained. To further justify the importance of the crossed product calculus we conclude the paper with explicit examples (Section 4) arising from pointed Hopf algebras, particularly from the Radford Hopf algebras $H_{(r,n,q)}$. We would like to stress that the resulting calculus on a pointed Hopf algebra H is covariant with respect to the group algebra of G/N , where G and N are the groups of group-like elements of H and of the indecomposable component of H containing the unit, respectively. In particular, it is not covered by the Woronowicz classification [26].

It is important to note that part of this paper is inspired by the work [18] of Pflaum–Schauenburg. They construct a canonical covariant calculus on smash product algebras $B \# H$ or, equivalently, on trivial extensions $B = A^{\text{co}H} \subseteq A$. The results of this paper generalize the findings in [18], in the sense that we consider cleft extensions or, equivalently, crossed product algebras $B \#_{\sigma} H$, with arbitrary 2-cocycles $\sigma: H \otimes H \rightarrow B$ with

values in B . The generalization is not straightforward, since, for example, the necessary compatibility of σ with the differential of B which we provide is automatic in case of the trivial 2-cocycle. We further give a classification of the smash product calculus in Section 3.2, in terms of the differentials of the cleaving and inclusion maps. There are interesting classes of cleft extensions which are not trivial (see Example 2.10 and Section 4), in particular in the context of pointed Hopf algebras, which further justify the relevance of this generalization. Let us also point out the reference [2], where differential calculi on double smash products are considered. A generalization to double crossed products might be feasible and we leave this as a future project. To avoid confusion let us clarify that the authors of [2] use the name cross product instead of smash product, while we follow the convention of [17], where crossed products allow for non-trivial 2-cocycles and measures.

Conventions and notations. We fix a field \mathbb{k} and all vector spaces are understood to be \mathbb{k} -vector spaces unless otherwise specified. By a linear map we mean a \mathbb{k} -linear map and the unadorned tensor product \otimes is the one of vector spaces. All linear maps whose domain is a tensor product will usually be defined on generators and understood to be extended by linearity. Algebras over \mathbb{k} will be associative and unital. Our general references for Hopf algebra theory are [24] and [17], where one can find the notions of coalgebra, bialgebra, Hopf algebra and (co)modules. We denote the multiplication and the unit of an algebra by m and u , while the comultiplication and the counit of a coalgebra are denoted by Δ and ε ; S will denote the antipode of a Hopf algebra, which we always assume invertible in the following. We will use Sweedler's notation for calculations involving the comultiplication, meaning we shall write $\Delta(h) = h_1 \otimes h_2$ for any $h \in H$ (with the usual summation convention, where $h_1 \otimes h_2$ stands for the finite sum $\sum_i h_1^i \otimes h_2^i$). Given a coalgebra C and a right (resp. left) C -comodule M we use Sweedler's like notation for calculations involving the coaction $\rho: M \rightarrow M \otimes C$ (resp. $\lambda: M \rightarrow C \otimes M$), namely we omit finite summation and write $\rho(m) = m_0 \otimes m_1$ (resp. $\lambda(m) = m_{-1} \otimes m_0$), for any $m \in M$. In this work H will always denote a bialgebra, while A and B are associative unital algebras. A map $f: V^\bullet \rightarrow W^\bullet$ between \mathbb{N} -graded vector spaces $V^\bullet = \bigoplus_{n \in \mathbb{N}} V^n$, $W^\bullet = \bigoplus_{n \in \mathbb{N}} W^n$ is of degree $k \in \mathbb{Z}$ if $f(V^n) \subseteq W^{n+k}$ for all $n \in \mathbb{N}$, where we set $W^{-n} = \{0\}$ by default. The tensor product $V^\bullet \otimes W^\bullet$ of \mathbb{N} -graded vector spaces V^\bullet , W^\bullet becomes an \mathbb{N} -graded vector space with $(V^\bullet \otimes W^\bullet)^n := \bigoplus_{k=0}^n V^k \otimes W^{n-k}$ for all $n \in \mathbb{N}$.

2 Preliminaries

In this section we recall some basic notions and results concerning (co)action compatible modules (Section 2.1), crossed product algebras (Section 2.2), noncommutative differential calculi (Section 2.3), as well as quantum principal bundles and their connections (Section 2.4). The reader who is familiar with these concepts can proceed to Section 3, which contains the first original contribution of this paper.

2.1 Equivariant and covariant modules

Fix a bialgebra H . It is well-known that the categories of left (right, bi) H -modules ${}_H\mathfrak{M}$ (\mathfrak{M}_H , ${}_H\mathfrak{M}_H$) are monoidal categories. Recall that, for every M and N in ${}_H\mathfrak{M}$, the tensor product $M \otimes N$ has a left H -action given by $h \cdot (m \otimes n) = h_1 \cdot m \otimes h_2 \cdot n$ for every $h \in H$, $m \in M$ and $n \in N$ (called the *diagonal left action*), where we denote the actions on M and N by the same symbol. The unit object is the base field \mathbb{k} with left H -action given by $h \cdot k = \varepsilon(h)k$ for every $h \in H$ and $k \in \mathbb{k}$ (called the *trivial left action*).

Dually, the categories of right (left, bi) H -comodules \mathfrak{M}^H (${}^H\mathfrak{M}$, ${}^H\mathfrak{M}^H$) are monoidal. Namely, for every M and N in \mathfrak{M}^H the tensor product $M \otimes N$ is an object in \mathfrak{M}^H with $\rho_{M \otimes N}(m \otimes n) = m_0 \otimes n_0 \otimes m_1 n_1$ for every $m \in M$ and $n \in N$ (called the *diagonal right coaction*). The unit object is \mathbb{k} , with right H -coaction $\rho_{\mathbb{k}}(k) = k \otimes 1_H$ for every $k \in \mathbb{k}$ (called the *trivial right coaction*). Since these categories of modules and comodules are monoidal we can consider categories of monoids (algebras) and comonoids (coalgebras) in them.

Definition 2.1. *A left H -module algebra is an algebra object in the category ${}_H\mathfrak{M}$, while a right H -comodule algebra is an algebra object in the category \mathfrak{M}^H . Explicitly, an algebra (A, m, u) is:*

- i.) *a left H -module algebra if A is a left H -module such that $m: A \otimes A \rightarrow A$ and $u: \mathbb{k} \rightarrow A$ are left H -linear maps, i.e., for all $h \in H$ and $a, b \in A$ we have*

$$h \cdot (ab) = (h_1 \cdot a)(h_2 \cdot b) \quad \text{and} \quad h \cdot 1_A = \varepsilon(h)1_A.$$

- ii.) *a right H -comodule algebra if A is a right H -comodule such that $m: A \otimes A \rightarrow A$ and $u: \mathbb{k} \rightarrow A$ are right H -colinear maps, i.e., for all $a, b \in A$ we have*

$$\rho(ab) = a_0 b_0 \otimes a_1 b_1 \quad \text{and} \quad \rho(1_A) = 1_A \otimes 1_H.$$

Morphisms of left H -module algebras are left H -linear algebra morphisms, while morphisms of right H -comodule algebras are right H -colinear algebra morphisms.

Similarly, right H -module algebras, H -bimodule algebras, left H -comodule algebras and H -bicomodule algebras are defined. Given an algebra object A in the category ${}_H\mathfrak{M}$, we can consider the category of left (right, bi) A -modules in ${}_H\mathfrak{M}$, denoted by ${}_A({}_H\mathfrak{M})$ ($({}_H\mathfrak{M})_A$, ${}_A({}_H\mathfrak{M})_A$). Analogously, if A is an algebra object in the category \mathfrak{M}^H we can consider the category of left (right, bi) A -modules in the category \mathfrak{M}^H , denoted by ${}_A(\mathfrak{M}^H)$ $(\mathfrak{M}^H)_A$, ${}_A(\mathfrak{M}^H)_A$, even removing the parentheses. Thus, arriving at the following definition.

Definition 2.2. *i.) Let A be a left H -module algebra. A left H -equivariant left A -module is an object in the category ${}_A({}_H\mathfrak{M})$. Explicitly, it is a left A -module M with action $A \otimes M \ni a \otimes m \mapsto am \in M$ which is also a left H -module with action $H \otimes M \ni h \otimes m \mapsto h \cdot m \in M$ such that*

$$h \cdot (am) = (h_1 \cdot a)(h_2 \cdot m) \quad \text{for all } h \in H, a \in A \text{ and } m \in M. \quad (1)$$

A morphism of left H -equivariant left A -modules is a left H -linear and left A -linear map. Similarly left H -equivariant right A -modules and A -bimodules are defined.

ii.) Let A be a right H -comodule algebra. A right H -covariant left A -module is an object in the category ${}_A\mathfrak{M}^H$. Explicitly, it is a left A -module M with action $A \otimes M \ni a \otimes m \mapsto am \in M$ which is also a right H -comodule with coaction $\rho : M \rightarrow M \otimes H$ such that

$$\rho(am) = a_0 m_0 \otimes a_1 m_1 \quad \text{for all } a \in A \text{ and } m \in M. \quad (2)$$

Morphisms of right H -covariant left A -modules are right H -colinear and left A -linear maps. Similarly right H -covariant right A -modules and A -bimodules are defined.

We give some canonical examples of (co)module algebras and their corresponding bimodules.

Example 2.3. *i.) If H is a Hopf algebra it becomes a left H -module algebra with respect to the adjoint action $H \otimes H \rightarrow H$, $h \otimes h' \mapsto h_1 h' S(h_2)$. Given a Hopf algebra H and an arbitrary left H -module algebra A , the algebra of endomorphisms $\text{End}_{\mathbb{k}}(A)$ is a left H -module algebra with respect to the adjoint action*

$$H \otimes \text{End}_{\mathbb{k}}(A) \rightarrow \text{End}_{\mathbb{k}}(A), \quad h \otimes \phi \mapsto \left(h \cdot \phi : a \mapsto (h \cdot \phi)(a) := h_1 \cdot \phi(S(h_2) \cdot a) \right).$$

ii.) Any bialgebra H is a right (left and bi) H -comodule algebra with coaction given by the coproduct $\Delta : H \rightarrow H \otimes H$. Given a vector space V one can construct a right H -covariant right H -module $V \otimes H$ with (co)actions the ones on the second tensor factor. If H is a Hopf algebra, every object M in \mathfrak{M}_H^H is (isomorphic to a module) of this form, with $V = M^{\text{co}H} := \{m \in M \mid \rho(m) = m \otimes 1_H\}$ the vector subspace of coinvariants and $M \xrightarrow{\cong} M^{\text{co}H} \otimes H$, $m \mapsto m_0 \cdot S(m_1) \otimes m_2$. This is commonly known as the Fundamental Theorem of Hopf modules [24]. In case M is an object in ${}_H^H\mathfrak{M}_H^H$ the above becomes an isomorphism in ${}_H^H\mathfrak{M}_H^H$ and $M^{\text{co}H}$ becomes a Yetter-Drinfel'd module, leading to a (braided) monoidal equivalence of the categories of H -bicovariant H -bimodules and Yetter-Drinfel'd modules [22].

iii.) For a Hopf algebra surjection $\pi : A \rightarrow H$ one can structure A as a right H -comodule algebra with right H -coaction $\rho := (\text{Id} \otimes \pi) \circ \Delta : A \rightarrow A \otimes H$. The vector subspace of coinvariants $B := A^{\text{co}H}$ is a subalgebra of A . If the canonical map $A \otimes_B A \rightarrow A \otimes H$, $a \otimes_B a' \mapsto aa'_0 \otimes a'_1$ is invertible and A is faithfully flat as a left B -module there is a categorical equivalence ${}_B\mathfrak{M}^H \cong {}_B\mathfrak{M}_B^A$ where an object V in ${}_B\mathfrak{M}^H$ is assigned to the cotensor product $V \square_H A$. This is known as Takeuchi equivalence [25]. For $H = \mathbb{k}$ and $\pi = \varepsilon$ we obtain $B = A$ and recover the Fundamental Theorem of Hopf modules.

2.2 Crossed product algebras

In this subsection H denotes a Hopf algebra and B an algebra. We say that H measures B if there is a linear map $H \otimes B \rightarrow B$, $h \otimes b \mapsto h \cdot b$, such that $h \cdot 1_B = \varepsilon(h)1_B$ and $h \cdot (bb') = (h_1 \cdot b)(h_2 \cdot b')$ for all $h \in H$ and $b, b' \in B$.

Definition 2.4. *If H measures B and $\sigma : H \otimes H \rightarrow B$ is a linear map which is convolution invertible, i.e., there exists a linear map $\sigma^{-1} : H \otimes H \rightarrow B$ such that $\sigma(h_1 \otimes h'_1)\sigma^{-1}(h_2 \otimes h'_2) = \varepsilon(hh')1_B = \sigma^{-1}(h_1 \otimes h'_1)\sigma(h_2 \otimes h'_2)$ for every $h, h' \in H$, then the crossed product $B \#_{\sigma} H$ of B with H is the vector space $B \otimes H$ endowed with the linear map*

$$(B \otimes H) \otimes (B \otimes H) \rightarrow B \otimes H, \quad (b \otimes h) \otimes (b' \otimes h') \mapsto b(h_1 \cdot b')\sigma(h_2 \otimes h'_1) \otimes h_3 h'_2 \quad (3)$$

for all $b, b' \in B$ and $h, h' \in H$.

Lemma 2.5 ([17, Lemma 7.1.2]). *The crossed product $B\#_\sigma H$ is an associative algebra with multiplication given by the assignment (3) and unit $1_B \otimes 1_H$ if and only if the following two conditions are satisfied:*

i.) B is a σ -twisted left H -module, i.e., $1_H \cdot b = b$ for all $b \in B$ and

$$h \cdot (h' \cdot b) = \sigma(h_1 \otimes h'_1)(h_2 h'_2 \cdot b) \sigma^{-1}(h_3 \otimes h'_3) \quad (4)$$

for all $h, h' \in H$ and $b \in B$.

ii.) σ is a 2-cocycle with values in B , that is, $\sigma(h \otimes 1_H) = \varepsilon(h)1_B = \sigma(1_H \otimes h)$ for all $h \in H$ and

$$(h_1 \cdot \sigma(h'_1 \otimes h''_1))\sigma(h_2 \otimes h'_2 h''_2) = \sigma(h_1 \otimes h'_1)\sigma(h_2 h'_2 \otimes h''_2) \quad (5)$$

for all $h, h', h'' \in H$.

It is straightforward to prove that, in case the assumptions of Lemma 2.5 are satisfied, $B\#_\sigma H$ is a right H -comodule algebra with right H -coaction $\rho := \text{Id}_B \otimes \Delta: B\#_\sigma H \rightarrow (B\#_\sigma H) \otimes H$. Note that $(B\#_\sigma H)^{\text{co}H} = B \otimes \mathbb{k}1_H \cong B$. In the following we refer to $B\#_\sigma H$ with the associative product (3) as the *crossed product algebra*. The product will be denoted by

$$(b \otimes h) \cdot (b' \otimes h') = b(h_1 \cdot b')\sigma(h_2 \otimes h'_1) \otimes h_3 h'_2$$

for $b \otimes h, b' \otimes h' \in B\#_\sigma H$.

Remark 2.6. *A particular class of crossed product algebras is obtained in the following way: every left H -module algebra B is a σ -twisted left H -module with respect to the trivial 2-cocycle $h \otimes h' \mapsto \varepsilon(hh')1_B$. The corresponding product is $(b \otimes h) \cdot (b' \otimes h') = b(h_1 \cdot b') \otimes h_2 h'$ and one obtains $B\#H$, the smash product algebra of B and H .*

We recall that crossed product algebras classify cleft extensions, with smash product algebras corresponding to trivial extensions. Let A be a right H -comodule algebra and consider $B := A^{\text{co}H}$ the subalgebra of right H -coinvariant elements.

Definition 2.7. *We call $B \subseteq A$ a cleft extension if there is a convolution invertible morphism $j: H \rightarrow A$ of right H -comodules, the so-called cleaving map. If $j: H \rightarrow A$ is a morphism of right H -comodule algebras we call $B \subseteq A$ a trivial extension.*

By definition, every trivial extension is a cleft extension. The convolution inverse of a cleaving map $j: H \rightarrow A$ is commonly denoted by $j^{-1}: H \rightarrow A$. Note that, if $j: H \rightarrow A$ is a cleaving map, we obtain another cleaving map $j': H \rightarrow A$ by $j'(h) := j^{-1}(1_H)j(h)$ for all $h \in H$. It follows that j' and its convolution inverse are unital. For this reason we can always assume a cleaving map to be unital with unital convolution inverse. If in the following we mention a cleaving map we refer to the unital convention.

Remark 2.8. *Note that every cleaving map j is injective. Indeed, supposing $j(h) = 0$ for an $h \in H$ gives $0 = j(h)_0 \otimes j(h)_1 = j(h_1) \otimes h_2$ and hence $j(h_1) \otimes h_2 \otimes h_3 = 0$. Thus $0 = j(h_1)j^{-1}(h_2) \otimes h_3 = \varepsilon(h_1)1_A \otimes h_2 = 1_A \otimes h$, which implies $h = 0$.*

The following theorem shows that crossed product algebras are in 1:1-correspondence with cleft extensions, where smash product algebras correspond to trivial extensions.

Theorem 2.9 ([17, Theorem 7.2.2]). *Any crossed product algebra $B\#_\sigma H$ is a cleft extension $B \subseteq B\#_\sigma H$ with cleaving map $j: H \rightarrow B\#_\sigma H$, $h \mapsto 1_B \otimes h$. Conversely, given a cleft extension $B \subseteq A$ with cleaving map $j: H \rightarrow A$, define a measure on B*

$$\cdot: H \otimes B \rightarrow B, \quad h \otimes b \mapsto h \cdot b := j(h_1)bj^{-1}(h_2)$$

and a 2-cocycle with values in B

$$\sigma: H \otimes H \rightarrow B, \quad h \otimes h' \mapsto \sigma(h \otimes h') := j(h_1)j(h'_1)j^{-1}(h_2 h'_2).$$

Then $A \cong B\#_\sigma H$ are isomorphic as right H -comodule algebras via $\theta: a \mapsto a_0 j^{-1}(a_1) \otimes a_2$, with inverse $\theta^{-1}: b \otimes h \mapsto bj(h)$. A cleft extension is a trivial extension if and only if the corresponding crossed product algebra is a smash product algebra.

We recall some examples of crossed product algebras and cleft extensions.

Example 2.10. i.) *Every group algebra $\mathbb{k}G$ can be seen as a crossed product algebra of the form $\mathbb{k}G = \mathbb{k}N\#_\sigma \mathbb{k}[G/N]$, given a normal subgroup N of G . Denoting $\bar{G} := G/N$ and choosing a coset representative $\gamma(\bar{x})$ for each coset $\bar{x} \in \bar{G}$, we have $\bar{x} \cdot n = \gamma(\bar{x})n\gamma(\bar{x})^{-1}$ and $\sigma(\bar{x} \otimes \bar{y}) = \gamma(\bar{x})\gamma(\bar{y})\gamma(\bar{x}\bar{y})^{-1}$.*

- ii.) The universal enveloping algebra $U(\mathfrak{g})$ of any Lie algebra \mathfrak{g} is isomorphic to the crossed product algebra $U(\mathfrak{h})\#_{\sigma}U(\bar{\mathfrak{g}})$ for any Lie ideal \mathfrak{h} of \mathfrak{g} and $\bar{\mathfrak{g}} := \mathfrak{g}/\mathfrak{h}$. Indeed, $U(\mathfrak{g})$ is a $U(\bar{\mathfrak{g}})$ -comodule algebra with coinvariants $U(\mathfrak{h})$ and $U(\mathfrak{h}) \subseteq U(\mathfrak{g})$ is a cleft extension with cleaving map $j : U(\bar{\mathfrak{g}}) \rightarrow U(\mathfrak{g})$ given by $j(\bar{x}_1 \cdots \bar{x}_n) := x_1 \cdots x_n$. For details we refer to [17, Corollary 7.2.8].
- iii.) Given fields $F \subseteq E$ containing \mathbb{k} , such that E/F is a finite Galois extension with Galois group G , we obtain $E \cong F\#_{\sigma}(\mathbb{k}G)^*$. Details can be found in [17, Example 7.1.11].
- iv.) Consider a Hopf algebra H , together with a convolution invertible normalized 2-cocycle $\gamma : H \otimes H \rightarrow \mathbb{k}$, meaning that for all $a, b, c \in H$ we have $\gamma(1 \otimes a) = \varepsilon(a) = \gamma(a \otimes 1)$,

$$\gamma(a_1 \otimes b_1)\gamma(a_2 b_2 \otimes c) = \gamma(b_1 \otimes c_1)\gamma(a \otimes b_2 c_2)$$

and that there is a map $\gamma^{-1} : H \otimes H \rightarrow \mathbb{k}$ such that $\gamma(a_1 \otimes b_2)\gamma^{-1}(a_2 \otimes b_2) = \varepsilon(ab) = \gamma^{-1}(a_1 \otimes b_2)\gamma(a_2 \otimes b_2)$. There is a Hopf algebra H_{γ} , the twisted Hopf algebra, which coincides with H as a coalgebra, but is endowed with the product and antipode

$$a \cdot_{\gamma} b := \gamma(a_1 \otimes b_1)a_2 b_2 \gamma^{-1}(a_3 \otimes b_3), \quad S_{\gamma}(a) := U(a_1)S(a_2)U^{-1}(a_3)$$

for $a, b \in H$, where $U : H \rightarrow \mathbb{k}$, $a \mapsto U(a) := \gamma(a_1 \otimes S(a_2))$ is a convolution invertible map with inverse $U^{-1} : H \rightarrow \mathbb{k}$, cf. [16, p.61]. Furthermore, $H_{\bullet_{\gamma}} := (H, \bullet_{\gamma})$ with product $a \bullet_{\gamma} b := a_1 b_1 \gamma^{-1}(a_2 \otimes b_2)$ becomes a right H_{γ} -comodule algebra with coaction given by the coproduct and, following [1, Example 6.3], we obtain a cleft extension $\mathbb{k} \subseteq H_{\bullet_{\gamma}}$ with cleaving map $j = \text{Id}_H : H_{\gamma} \rightarrow H_{\bullet_{\gamma}}$. Note that this cleft extension is not trivial in general, since H_{γ} and $H_{\bullet_{\gamma}}$ are not always isomorphic as algebras. Observe that, if (H, \mathcal{R}) is a coquasitriangular Hopf algebra, then \mathcal{R} is a convolution invertible normalized 2-cocycle and we obtain a cleft extension $\mathbb{k} \subseteq H_{\bullet_{\mathcal{R}}}$, which is not trivial in general. In this case, as it is shown in [13, Example 12, page 355], we have that $H_{\mathcal{R}}$ is the opposite Hopf algebra H^{op} .

- v.) We give an explicit example of iv.), by considering the cotriangular Sweedler's Hopf algebra. First, let us recall some details. Consider the free \mathbb{k} -algebra H generated by two elements g and x modulo the relations

$$g^2 = 1, \quad x^2 = 0 \quad \text{and} \quad xg = -gx,$$

where 1 denotes the unity of H . It becomes a Hopf algebra with comultiplication, counit and antipode determined by $\Delta(g) = g \otimes g$, $\Delta(x) = x \otimes 1 + g \otimes x$, $\varepsilon(g) = 1$, $\varepsilon(x) = 0$, $S(g) = g$ and $S(x) = -gx$, respectively. In particular, H is neither commutative nor cocommutative. We recall from [16, Example 2.2.6] that H has a cotriangular structure given by

$$\mathcal{R} \begin{pmatrix} 1 \otimes 1 & 1 \otimes g & 1 \otimes x & 1 \otimes xg \\ g \otimes 1 & g \otimes g & g \otimes x & g \otimes xg \\ x \otimes 1 & x \otimes g & x \otimes x & x \otimes xg \\ xg \otimes 1 & xg \otimes g & xg \otimes x & xg \otimes xg \end{pmatrix} = \begin{pmatrix} 1 & 1 & 0 & 0 \\ 1 & -1 & 0 & 0 \\ 0 & 0 & \alpha & -\alpha \\ 0 & 0 & \alpha & \alpha \end{pmatrix}, \quad (6)$$

for $\alpha \in \mathbb{k}$, with convolution inverse $\mathcal{R}^{-1} = \mathcal{R}^{\text{op}}$ given by the transpose matrix

$$\mathcal{R}^{-1} \begin{pmatrix} 1 \otimes 1 & 1 \otimes g & 1 \otimes x & 1 \otimes xg \\ g \otimes 1 & g \otimes g & g \otimes x & g \otimes xg \\ x \otimes 1 & x \otimes g & x \otimes x & x \otimes xg \\ xg \otimes 1 & xg \otimes g & xg \otimes x & xg \otimes xg \end{pmatrix} = \begin{pmatrix} 1 & 1 & 0 & 0 \\ 1 & -1 & 0 & 0 \\ 0 & 0 & \alpha & \alpha \\ 0 & 0 & -\alpha & \alpha \end{pmatrix}. \quad (7)$$

As discussed in iv.) we have $H_{\mathcal{R}} = H^{\text{op}}$. Now we build $H_{\bullet_{\mathcal{R}}}$, which will be a right H^{op} -comodule algebra, providing a cleft extension $\mathbb{k} \subseteq H_{\bullet_{\mathcal{R}}}$. As a vector space $H_{\bullet_{\mathcal{R}}}$ equals $H = \mathbb{k}1 \oplus \mathbb{k}g \oplus \mathbb{k}x \oplus \mathbb{k}xg$. We obtain

$$g \bullet_{\mathcal{R}} g = g_1 g_1 \mathcal{R}^{-1}(g_2 \otimes g_2) = g^2 \mathcal{R}^{-1}(g \otimes g) = -1$$

$$x \bullet_{\mathcal{R}} x = x_1 x_1 \mathcal{R}^{-1}(x_2 \otimes x_2) = x^2 \mathcal{R}^{-1}(1 \otimes 1) + xg \mathcal{R}^{-1}(1 \otimes x) + gx \mathcal{R}^{-1}(x \otimes 1) + g^2 \mathcal{R}^{-1}(x \otimes x) = \alpha 1$$

$$x \bullet_{\mathcal{R}} g = x_1 g_1 \mathcal{R}^{-1}(x_2 \otimes g_2) = xg \mathcal{R}^{-1}(1 \otimes g) + g^2 \mathcal{R}^{-1}(x \otimes g) = xg$$

and similarly $g \bullet_{\mathcal{R}} x = gx$. Hence, $H_{\bullet_{\mathcal{R}}}$ can be considered as the free \mathbb{k} -algebra generated by g and x modulo the relations

$$g^{\bullet_{\mathcal{R}^2}} = -1, \quad x^{\bullet_{\mathcal{R}^2}} = \alpha 1 \quad \text{and} \quad x \bullet_{\mathcal{R}} g = -g \bullet_{\mathcal{R}} x$$

and then it is clear that $H_{\bullet_{\mathcal{R}}}$ and H^{op} are not isomorphic as algebras. Thus $\mathbb{k} \subseteq H_{\bullet_{\mathcal{R}}}$ is a cleft extension which is not trivial.

2.3 Noncommutative differential calculi

In this subsection we discuss differential structures on noncommutative algebras. We recall the definition of a (covariant) differential calculus, together with some well-known classification results. The original source of this material is [26] and we refer the reader to [13, 21] for more details.

Recall that a *differential graded algebra* (A^\bullet, \wedge, d) is an \mathbb{N} -graded vector space $A^\bullet = \bigoplus_{n \in \mathbb{N}} A^n$, endowed with a degree 0 map $\wedge : A^\bullet \otimes A^\bullet \rightarrow A^\bullet$, the *wedge product*, a degree 1 map $d : A^\bullet \rightarrow A^{\bullet+1}$, the *differential*, and an element $1 \in A^0$, such that \wedge is associative and unital with respect to 1 , and such that $d^2 := d \circ d = 0$, as well as the *Leibniz rule*

$$d(\omega \wedge \eta) = d\omega \wedge \eta + (-1)^n \omega \wedge d\eta \quad (8)$$

hold for all $\omega \in A^n$ and $\eta \in A^\bullet$. Note that (A^\bullet, \wedge) is an algebra in the monoidal category $\text{Vec}_{\mathbb{N}}$ of \mathbb{N} -graded vector spaces and it is equipped with a differential d which satisfies $d^2 = 0$ and the Leibniz rule. A morphism $\Phi : (A^\bullet, \wedge, d) \rightarrow (A'^\bullet, \wedge', d')$ between two differential graded algebras is a linear map $\Phi : A^\bullet \rightarrow A'^\bullet$ of degree 0 such that $\Phi(\omega \wedge \eta) = \Phi(\omega) \wedge' \Phi(\eta)$ and $\Phi \circ d = d' \circ \Phi$ for all $\omega, \eta \in A^\bullet$. Furthermore, Φ is an isomorphism if it is invertible.

Definition 2.11. *A differential calculus (DC) on an algebra A is a differential graded algebra $(\Omega^\bullet(A), \wedge, d)$ with $\Omega^0(A) = A$ such that, for all $n \geq 1$, the vector space $\Omega^n(A)$ is spanned by elements of the form $a^0 da^1 \wedge \cdots \wedge da^n$ with $a^i \in A$. A morphism Φ from a DC $(\Omega^\bullet(A), \wedge, d)$ on an algebra A to a DC $(\Omega^\bullet(A'), \wedge', d')$ on an algebra A' is a morphism of the underlying differential graded algebras.*

Given $(\Omega^\bullet(A), \wedge, d)$ and $(\Omega^\bullet(A'), \wedge', d')$ DCi on algebras A and A' respectively, we call an algebra map $f : A \rightarrow A'$ *differentiable* if there exists a morphism $\Omega^\bullet(f) : \Omega^\bullet(A) \rightarrow \Omega^\bullet(A')$ of differential graded algebras with $\Omega^0(f) = f$. Thus, $\Omega^\bullet(f) : \bigoplus_{n \geq 0} \Omega^n(A) \rightarrow \bigoplus_{n \geq 0} \Omega^n(A')$ is an algebra map defined by a collection of linear maps $\Omega^n(f) : \Omega^n(A) \rightarrow \Omega^n(A')$ such that $d' \circ \Omega^n(f) = \Omega^{n+1}(f) \circ d$. Observe that $\Omega^\bullet(f)$ is completely determined by f since $\Omega^n(f)(a^0 da^1 \wedge \cdots \wedge da^n) = f(a^0) d'(f(a^1)) \wedge \cdots \wedge d'(f(a^n))$ for all $a^i \in A$.

Further recall that an \mathbb{N} -graded vector space $V = \bigoplus_{n \in \mathbb{N}} V^n$ is a *graded right H -comodule* if there is a right H -coaction $\rho : V \rightarrow V \otimes H$ such that $\rho(V^n) \subseteq V^n \otimes H$ for all $n \geq 0$. Since $\text{Vec}_{\mathbb{N}}$ is a monoidal category one can consider H in $\text{Comon}(\text{Vec}_{\mathbb{N}})$, i.e., a \mathbb{N} -graded coalgebra and form the category $\text{Vec}_{\mathbb{N}}^H$ whose objects V are \mathbb{N} -graded vector spaces and right H -comodules via $\rho : V \rightarrow V \otimes H$ such that $\rho(V^n) \subseteq V^n \otimes H$. Here we are considering H with trivial grading. It follows that, for any $n \geq 0$, the vector space V^n is a right H -comodule via the restriction and corestriction $\rho^n := \rho|_{V^n} : V^n \rightarrow V^n \otimes H$, so we can write $\rho = \bigoplus_{n \in \mathbb{N}} \rho^n$. A linear map $\phi : V \rightarrow W$ of degree k between graded right H -comodules (V, ρ_V) and (W, ρ_W) is said to be right H -colinear if $\rho_W^{n+k} \circ \phi^n = (\phi^n \otimes \text{Id}_H) \circ \rho_V^n$ for all $n \geq 0$, where $\phi^n := \phi|_{V^n} : V^n \rightarrow W^{n+k}$. If H is a bialgebra and then an object in $\text{Bimon}(\text{Vec}_{\mathbb{N}})$ considered with trivial grading, then $\text{Vec}_{\mathbb{N}}^H$ is monoidal. An object V in $\text{Mon}(\text{Vec}_{\mathbb{N}}^H)$ is called *graded right H -comodule algebra*. It is a graded algebra which is also a graded right H -comodule and the algebra structure is compatible with the right H -coaction in the usual sense. The notion of graded comodule algebra is essential for the following definition.

Definition 2.12. *A differential calculus $(\Omega^\bullet(A), \wedge, d)$ on a right H -comodule algebra (A, ρ_A) is called right H -covariant if $(\Omega^\bullet(A), \wedge)$ is a graded right H -comodule algebra with graded coaction $\rho : \Omega^\bullet(A) \rightarrow \Omega^\bullet(A) \otimes H$ such that $\rho^0 = \rho_A$ and the differential $d : \Omega^\bullet(A) \rightarrow \Omega^\bullet(A)$ is right H -colinear.*

Similarly, left H -covariant and H -bicovariant DC are defined on left H -comodule algebras and H -bicomodule algebras, respectively. In case of the comodule algebra H with coaction given by the coproduct we call an H -covariant calculus covariant, i.e., we omit the reference to H . We also use the expression (Ω^\bullet, d) for a DC on an algebra A when the latter is clear from the context.

Example 2.13. *i.) Every algebra A admits the trivial DC $\Omega^n := \{0\}$ for $n > 0$.*

ii.) On every algebra (A, m) one can construct the universal DC (Ω_u^\bullet, d_u) in the following way: for $n > 0$ we set $\Omega_u^n := \bigcap_{k=1}^n \ker m^k$, where $m^k := \text{Id}_{A^{\otimes(k-1)}} \otimes m \otimes \text{Id}_{A^{\otimes(n-k)}} : A^{\otimes(n+1)} \rightarrow A^{\otimes n}$ denotes the multiplication of the k^{th} and $(k+1)^{\text{th}}$ tensor factors. Then Ω_u^n becomes an A -bimodule with left multiplication on the first and right multiplication on the last tensor factor, respectively. The corresponding wedge product is determined by

$$(a^0 \otimes \cdots \otimes a^n) \wedge_u (b^0 \otimes \cdots \otimes b^m) := a^0 \otimes \cdots \otimes a^n b^0 \otimes \cdots \otimes b^m$$

for $a^0, \dots, a^n, b^0, \dots, b^m \in A$.¹ The differential is given by

$$d_u(a^0 \otimes \cdots \otimes a^n) := \sum_{k=0}^{n+1} (-1)^k a^0 \otimes \cdots \otimes a^{k-1} \otimes 1 \otimes a^k \otimes \cdots \otimes a^n.$$

¹Here and below we denote an arbitrary element of Ω_u^n by $a^0 \otimes \cdots \otimes a^n$, omitting the finite summation $\sum_i a^{0,i} \otimes \cdots \otimes a^{n,i}$ with $a^{k,i} \in A$, such that $\sum_i a^{0,i} \cdots a^{n,i} = 0$.

One can show that (Ω_u^\bullet, d_u) is a DC on A , cf. [3, Theorem 1.33]. If A is a right H -comodule algebra, then (Ω_u^\bullet, d_u) is right H -covariant. Moreover, every DC on A is a quotient of (Ω_u^\bullet, d_u) , see e.g. [11, Appendix B]. The latter means that, if (Ω^\bullet, d) is an arbitrary DC on A , there is a surjective morphism $(\Omega_u^\bullet, d_u) \rightarrow (\Omega^\bullet, d)$ of differential graded algebras.

iii.) Given a DC (Ω^\bullet, d) on A and a DC (Ω'^\bullet, d') on another algebra A' there is a DC $(\Omega_\otimes^\bullet, d_\otimes)$ on the tensor product algebra $A \otimes A'$. Here, the graded module $\Omega_\otimes^n := \bigoplus_{k=0}^n \Omega^k \otimes \Omega'^{n-k}$ becomes a graded algebra with multiplication determined on homogeneous elements $\omega, \eta \in \Omega^\bullet$ and $\omega', \eta' \in \Omega'^\bullet$ by

$$(\omega \otimes \omega') \wedge_\otimes (\eta \otimes \eta') := (-1)^{|\omega'| |\eta|} (\omega \wedge \eta) \otimes (\omega' \wedge \eta').$$

The differential on Ω_\otimes^\bullet is given on homogeneous elements by

$$d_\otimes(\omega \otimes \omega') := d\omega \otimes \omega' + (-1)^{|\omega|} \omega \otimes d'\omega'.$$

It is straightforward to show that $(\Omega_\otimes^\bullet, d_\otimes)$ is a DC on $A \otimes A'$. We call it the tensor product calculus. Note that this construction implies that DCi form a monoidal category with monoidal unit given by the trivial calculus on \mathbb{k} . By i.) the category of differential calculi over a fixed algebra has a terminal object and by ii.) the same category has an initial object.

Remark 2.14. Let us spell out explicitly that, given a bialgebra H , a bicovariant DC on H is a graded H -bicomodule algebra $\Omega^\bullet(H)$ with graded coactions $\lambda : \Omega^\bullet(H) \rightarrow H \otimes \Omega^\bullet(H)$ and $\rho : \Omega^\bullet(H) \rightarrow \Omega^\bullet(H) \otimes H$ equipped with a differential d that is H -bilinear. In particular, we have that $\Omega^n(H)$ is an H -bicomodule for every $n \in \mathbb{N}$. In addition, for every $\gamma, \gamma' \in \Omega^\bullet(H)$ we also have the equations

$$(\gamma \wedge \gamma')_0 \otimes (\gamma \wedge \gamma')_1 = \gamma_0 \wedge \gamma'_0 \otimes \gamma_1 \gamma'_1 \quad \text{and} \quad (\gamma \wedge \gamma')_{-1} \otimes (\gamma \wedge \gamma')_0 = \gamma_{-1} \gamma'_{-1} \otimes \gamma_0 \wedge \gamma'_0.$$

It is proven in [21, Lemma 5.4] that Δ is differentiable if and only if $\Omega^\bullet(H)$ is a differential graded bialgebra with $\Omega^0(H) = H$ as bialgebras. Thus, in this case, we have $\Omega^\bullet(\Delta) : \Omega^\bullet(H) \rightarrow \Omega^\bullet(H \otimes H)$ where $\Omega^\bullet(H \otimes H)$ is the tensor product DC on $H \otimes H$. For an arbitrary $\gamma \in \Omega^\bullet(H)$ we write $\Omega^\bullet(\Delta)(\gamma) = \gamma_{[1]} \otimes \gamma_{[2]}$ and, denoting by $\pi^{kl} : \Omega^{k+l}(H \otimes H) \rightarrow \Omega^k(H) \otimes \Omega^l(H)$ the canonical projection. For a homogeneous element $\gamma \in \Omega^j(H)$ and $k+l=j$ we write $\pi^{kl}\Omega^j(\Delta)(\gamma) =: \gamma_{\langle 1, k \rangle} \otimes \gamma_{\langle 2, l \rangle}$, so that $\Omega^j(\Delta)(\gamma) = \sum_{k+l=j} \gamma_{\langle 1, k \rangle} \otimes \gamma_{\langle 2, l \rangle}$ in accordance with [18]. In terms of this notation, the fact that $\Omega^\bullet(\Delta)$ is an algebra map reads

$$(\gamma \wedge \gamma')_{[1]} \otimes (\gamma \wedge \gamma')_{[2]} = \sum_{\substack{k+l=j \\ k'+l'=j'}} (-1)^{lk'} \gamma_{\langle 1, k \rangle} \wedge \gamma'_{\langle 1, k' \rangle} \otimes \gamma_{\langle 2, l \rangle} \wedge \gamma'_{\langle 2, l' \rangle} \quad (9)$$

for every $\gamma \in \Omega^j(H)$ and $\gamma' \in \Omega^{j'}(H)$. Applying the projection $\pi^{0, k, l}$ to the equation $(\text{Id} \otimes \Omega^\bullet(\Delta))\Omega^\bullet(\Delta)(\gamma) = (\Omega^\bullet(\Delta) \otimes \text{Id})\Omega^\bullet(\Delta)(\gamma)$ gives

$$\gamma_{-1} \otimes \gamma_{0, \langle 1, k \rangle} \otimes \gamma_{0, \langle 2, l \rangle} = \gamma_{\langle 1, k \rangle, -1} \otimes \gamma_{\langle 1, k \rangle, 0} \otimes \gamma_{\langle 2, l \rangle} \quad (10)$$

and we have a similar formula for the projection $\pi^{k, l, 0}$.

In many situations the knowledge of $(\Omega^1, d|_A)$ is sufficient to describe the full differential calculus on A , e.g. via "maximal prolongation" (see [3, Lemma 1.32]) or via a braided exterior algebra construction (see [26, Theorem 4.1]). This motivates the importance of such "first order" DC data and leads to the following definition.

Definition 2.15. A first order differential calculus (FODC) on an algebra A is a tuple $(\Omega^1(A), d_A)$, where:

- i.) $\Omega^1(A)$ is an A -bimodule.
- ii.) $d_A : A \rightarrow \Omega^1(A)$ is a linear map which satisfies the Leibniz rule $d_A(ab) = (d_A a)b + a d_A b$ for all $a, b \in A$.
- iii.) $\Omega^1(A) = \text{Ad}_A A := \text{span}_{\mathbb{k}}\{a d_A b \mid a, b \in A\}$ (surjectivity condition).

A morphism of FODC $(\Omega^1(A), d_A) \rightarrow (\Omega^1(B), d_B)$ is a pair (f, F) , such that $f : A \rightarrow B$ is a morphism of algebras, $F : \Omega^1(A) \rightarrow \Omega^1(B)$ is an A -bimodule map, where f defines the A -bimodule structure on $\Omega^1(B)$ and $F \circ d_A = d_B \circ f$. An isomorphism of FODCi $(\Omega^1(A), d_A)$ and $(\Omega^1(B), d_B)$ is a morphism $(f, F) : (\Omega^1(A), d_A) \rightarrow (\Omega^1(B), d_B)$, such that f and F are invertible.

Given $(\Omega^1(A), d_A)$ and $(\Omega^1(B), d_B)$ FODCi, an algebra map $f : A \rightarrow B$ is called *differentiable*, if there exists a map $F : \Omega^1(A) \rightarrow \Omega^1(B)$ such that the pair (f, F) is a morphism of FODCi. Observe that in this case F is uniquely determined by f via $F(ad_A b) = f(a)d_B(f(b))$. Thus, we sometimes write $F = \widehat{f}$.

Remark 2.16. Let $(f, F): (\Omega^1(A), d_A) \rightarrow (\Omega^1(B), d_B)$ be an isomorphism of FODCi. Then the inverse algebra morphism $f^{-1}: B \rightarrow A$ of f is automatically differentiable with $\widehat{f^{-1}} = \widehat{f}^{-1}: \Omega^1(B) \rightarrow \Omega^1(A)$, the inverse A -module morphism of \widehat{f} . Clearly \widehat{f}^{-1} is an isomorphism of B -modules through f^{-1} and, since $\widehat{f} \circ d_A = d_B \circ f$, we obtain $d_A = \widehat{f}^{-1} \circ \widehat{f} \circ d_A = \widehat{f}^{-1} \circ d_B \circ f$, which implies $d_A \circ f^{-1} = f^{-1} \circ d_B$. Hence, by uniqueness, $\widehat{f^{-1}} = \widehat{f}^{-1}$.

We further discuss the first order of covariant DC on comodule algebras and, dually, FODCi which are invariant under a Hopf algebra action.

Definition 2.17. Let H be a Hopf algebra.

- i.) Consider a right (left) H -comodule algebra A and a FODC $(\Omega^1(A), d_A)$. The latter is said to be right (left) H -covariant if $\Omega^1(A)$ is in ${}^A\mathfrak{M}_A^H$ (resp. ${}^H\mathfrak{M}_A$) and d_A is in \mathfrak{M}^H (resp. ${}^H\mathfrak{M}$). Given an H -bicomodule algebra A , we say that $(\Omega^1(A), d_A)$ is H -bicovariant if $\Omega^1(A)$ is in ${}^H\mathfrak{M}_A^H$ and d_A is in ${}^H\mathfrak{M}^H$.
- ii.) Let A be a left (right) H -module algebra. A FODC $(\Omega^1(A), d_A)$ is called a left (right) H -module FODC if $\Omega^1(A)$ is in ${}_{A(H\mathfrak{M})_A}$ (resp. ${}_{A(\mathfrak{M}_H)_A}$) and d_A is in ${}_{H\mathfrak{M}}$ (resp. \mathfrak{M}_H). If A is an H -bimodule algebra, $(\Omega^1(A), d_A)$ is an H -bimodule FODC if $\Omega^1(A)$ is in ${}_{A(H\mathfrak{M}_H)_A}$ and d_A is in ${}_{H\mathfrak{M}_H}$.

As in the more general case of DC, we usually omit the reference to H in case of the H -comodule H itself with coaction given by the coproduct, i.e., we speak of *right/left or bicovariant FODC* in this setting. We recall from [3, Theorem 2.26] the classification theorem due to Woronowicz (for the original source of this result see [26, Theorem 1.8]). Recall that a Hopf algebra H is a right H -comodule via the *adjoint right coaction* $\text{Ad}_R: H \rightarrow H \otimes H$, $h \mapsto h_2 \otimes S(h_1)h_3$ and a left H -comodule via the *adjoint left coaction* $\text{Ad}_L: H \rightarrow H \otimes H$, $h \mapsto h_1 S(h_3) \otimes h_2$. We also denote the kernel of the counit by $H^+ := \ker \varepsilon$.

Theorem 2.18. Let H be a Hopf algebra. For any right ideal $I \subseteq H$ with $I \subseteq H^+$ we obtain a left covariant FODC $(\Omega^1(H), d_H)$ on H by $\Omega^1(H) := H \otimes (H^+/I)$ and $d_H h = (\text{Id} \otimes \pi)(\Delta(h) - h \otimes 1_H)$, where $\Omega^1(H)$ is an H -bimodule via

$$h \cdot (h' \otimes [g]) = hh' \otimes [g], \quad (h \otimes [g]) \cdot h' = hh' \otimes [gh'_2],$$

where $h, h' \in H$, $g \in H^+$ and $\pi: H^+ \rightarrow H^+/I$, $g \mapsto \pi(g) = [g]$ denotes the quotient map. The left H -coaction on $\Omega^1(H)$ is $\lambda := \Delta \otimes \text{Id}_{H^+/I}$. If $\text{Ad}_R(I) \subseteq I \otimes H$ the FODC $(\Omega^1(H), d_H)$ is bicovariant with right H -coaction

$$\rho(h \otimes [g]) = (h_1 \otimes [g_2]) \otimes h_2 S(g_1)g_3$$

for $h \in H$, $g \in H^+$. Moreover, every left/bi covariant FODC on H is isomorphic to a FODC of this form.

Remark 2.19. We also formulate the previous result in terms of right covariant calculi and left ideals. Then, if $I \subseteq H$ is a left ideal with $I \subseteq H^+$ we obtain a right covariant calculus on H by $\Omega^1(H) := (H^+/I) \otimes H$ and $d_H h = (\pi \otimes \text{Id})(\Delta(h) - 1_H \otimes h)$ where the left and right H -actions are

$$h \cdot ([g] \otimes h') = [h_1 g] \otimes h_2 h', \quad ([g] \otimes h') \cdot h = [g] \otimes h' h$$

for $h, h' \in H$, $g \in H^+$ and $\pi: H^+ \rightarrow H^+/I$, $g \mapsto \pi(g) = [g]$ denotes the quotient map. The right H -coaction on $\Omega^1(H)$ is $\rho := \text{Id}_{H^+/I} \otimes \Delta$. If $\text{Ad}_L(I) \subseteq H \otimes I$ the FODC is bicovariant with left H -coaction

$$\lambda([g] \otimes h) = g_1 S(g_3)h_1 \otimes ([g_2] \otimes h_2)$$

and every right/bi covariant FODC on H is isomorphic to a FODC of this form.

2.4 Quantum principal bundles and connections

We recall from [7] the definition of a quantum principal bundle and of a connection on it, see also [3, Section 5.4]. In this subsection H denotes a Hopf algebra, A is a right H -comodule algebra and we write $B := A^{\text{co}H} \subseteq A$ for the subalgebra of coinvariants. Given a FODC $(\Omega^1(A), d_A)$ we always consider the *pullback calculus* on B : this is the FODC $(\Omega^1(B), d_B)$ given by the restrictions $\Omega^1(B) := B d_A B \subseteq \Omega^1(A)$, $d_B := d_A|_B: B \rightarrow \Omega^1(B)$, and it is easy to prove that this gives a FODC on B . We further define the *horizontal forms* as the A -bimodule generated by $\Omega^1(B)$, i.e., $\Omega^1_{\text{hor}}(A) := A\Omega^1(B)A \subseteq \Omega^1(A)$.

Definition 2.20. Given a right H -comodule algebra A , a quantum principal bundle (QPB) on A is the data of

- i.) a right H -covariant FODC $(\Omega^1(A), d_A)$ on A ,
- ii.) a bicovariant FODC $(\Omega^1(H), d_H)$ on H ,

such that the vertical map

$$\text{ver} : \Omega^1(A) \rightarrow A \otimes {}^{\text{co}H}\Omega^1(H), \quad \text{ad}_A a' \mapsto aa'_0 \otimes S(a'_1) d_H a'_2$$

is well-defined and the Atiyah sequence

$$0 \longrightarrow \Omega^1_{\text{hor}}(A) \xrightarrow{i} \Omega^1(A) \xrightarrow{\text{ver}} A \otimes {}^{\text{co}H}\Omega^1(H) \longrightarrow 0 \quad (11)$$

is exact in ${}_A\mathfrak{M}$, where i is the inclusion.

Suppose that the vertical map is well-defined. Then a general element $\alpha \in \Omega^1_{\text{hor}}(A)$ is of the form $\alpha = a_i (d_A b^i) a^i = a_i d_A (b^i a^i) - a_i b^i d_A a^i$ with $a_i, a^i \in A$ and $b^i \in B$, where the summation symbol is omitted. Thus

$$\text{ver}(\alpha) = a_i b^i a^i_0 \otimes S(b^i_1 a^i_1) d_H (b^i_2 a^i_2) - a_i b^i a^i_0 \otimes S(a^i_1) d_H a^i_2 = a_i b^i a^i_0 \otimes S(a^i_1) d_H a^i_2 - a_i b^i a^i_0 \otimes S(a^i_1) d_H a^i_2 = 0$$

shows that $\Omega^1_{\text{hor}}(A) \subseteq \ker(\text{ver})$. Thus, given i.) and ii.) in the above definition, we have a QPB if ver is well-defined and surjective such that $\ker(\text{ver}) \subseteq \Omega^1_{\text{hor}}(A)$. We discuss a criterion for ver to be well-defined.

Remark 2.21. Let A be a right H -comodule algebra with a FODC $(\Omega^1(A), d_A)$ on A and a bicovariant FODC $(\Omega^1(H), d_H)$ on H . By [18, Theorem 2.8] we have that $\rho : A \rightarrow A \otimes H$ is differentiable if and only if $(\Omega^1(A), d_A)$ is right H -covariant and there is a map $\pi : \Omega^1(A) \rightarrow A \square_H \Omega^1(H)$ in ${}_A\mathfrak{M}_A^H$ such that $(\text{Id}_A \otimes d_H) \circ \rho = \pi \circ d_A$. Using the isomorphism $\phi : A \square_H \Omega^1(H) \rightarrow A \otimes {}^{\text{co}H}\Omega^1(H)$, $a \otimes \gamma \mapsto a \otimes S(\gamma_{-1})\gamma_0$ we obtain

$$\phi \circ \pi(\text{ad}_A a') = a_0 a'_0 \otimes S((a_1 d_H a'_1)_{-1}) (a_1 d_H a'_1)_0 = a_0 a'_0 \otimes S(a_1 a'_1) a_2 d_H a'_2 = a_0 a'_0 \otimes S(a'_1) \varepsilon(a_1) d_H a'_2 = \text{ver}(\text{ad}_A a').$$

Thus, if we assume to have a bicovariant FODC $(\Omega^1(H), d_H)$ and a FODC $(\Omega^1(A), d_A)$, ρ being differentiable is equivalent to $(\Omega^1(A), d_A)$ being right H -covariant and ver being well-defined (see also [3, page 425]). If ρ is differentiable then $\pi := \pi_2 \circ \hat{\rho}$, where $\pi_2 : \Omega^1(A \otimes H) \rightarrow A \otimes \Omega^1(H)$ is the canonical projection and so $\text{ver} = \phi \circ \pi_2 \circ \hat{\rho}$. Moreover, $\rho' := \pi_1 \circ \hat{\rho}$ is the right H -coaction of $\Omega^1(A)$, where $\pi_1 : \Omega^1(A \otimes H) \rightarrow \Omega^1(A) \otimes H$ is the other canonical projection.

Note that the inclusion $i : \Omega^1_{\text{hor}}(A) \rightarrow \Omega^1(A)$ is a morphism in ${}_A\mathfrak{M}^H$ and that $A \otimes {}^{\text{co}H}\Omega^1(H)$ is in ${}_A\mathfrak{M}^H$ with left A -action given by the multiplication on the first tensorand and right H -coaction given by the diagonal one. Indeed, the left A -action is clearly right H -colinear as $(aa')_0 \otimes \gamma_0 \otimes (aa')_1 \gamma_1 = a_0 a'_0 \otimes \gamma_0 \otimes a_1 a'_1 \gamma_1$ for all $a, a' \in A$ and $\gamma \in {}^{\text{co}H}\Omega^1(H)$. If ver is well-defined then it is automatically right H -colinear, indeed

$$\begin{aligned} \text{ver}(\text{ad}_A a'_0) \otimes \text{ver}(\text{ad}_A a'_1) &= (aa'_0)_0 \otimes (S(a'_1) d_H a'_2)_0 \otimes (aa'_1)_1 (S(a'_1) d_H a'_2)_1 \\ &= a_0 a'_0 \otimes S(a'_2)_1 (d_H a'_3)_0 \otimes a_1 a'_1 S(a'_2)_2 (d_H a'_3)_1 \\ &= a_0 a'_0 \otimes S(a'_3) d_H a'_4 \otimes a_1 a'_1 S(a'_2) a'_5 \\ &= a_0 a'_0 \otimes S(a'_1) d_H a'_2 \otimes a_1 a'_3 = \text{ver}(a_0 d_A a'_0) \otimes a_1 a'_1 \\ &= \text{ver}((\text{ad}_A a')_0) \otimes (\text{ad}_A a')_1. \end{aligned}$$

Thus, if ver is well-defined, the exactness of the sequence (11) in ${}_A\mathfrak{M}$ coincides with the exactness in ${}_A\mathfrak{M}^H$.

Definition 2.22. A connection is a splitting of (11), i.e., a morphism $c : A \otimes {}^{\text{co}H}\Omega^1(H) \rightarrow \Omega^1(A)$ in ${}_A\mathfrak{M}^H$ such that $\text{ver} \circ c = \text{Id}$.

Remark 2.23. Note that in [3, page 426] a connection is defined as a morphism $\pi : \Omega^1(A) \rightarrow \Omega^1(A)$ in ${}_A\mathfrak{M}^H$ such that $\pi \circ \pi = \pi$ and $\ker(\pi) = \Omega^1_{\text{hor}}(A)$ and π is called strong if $(\text{Id} - \pi) d_A A \subseteq \Omega^1(B)A$. Moreover, it is shown in [3, Proposition 5.41] that there is a 1-1 correspondence between connections in the previous sense and connections as in Definition 2.22. To each π is associated c such that $c(a \otimes \gamma) = \pi \text{ver}^{-1}(a \otimes \gamma)$ for all $a \in A$ and $\gamma \in {}^{\text{co}H}\Omega^1(H)$ and to each c is associated $\pi = c \circ \text{ver}$.

Example 2.24 ([3, Lemma 5.7]). Consider the universal FODC $(\Omega^1_u(A), d_A^u)$ and $(\Omega^1_u(H), d_H^u)$. This is a QPB if and only if $B = A^{\text{co}H} \subseteq A$ is a Hopf-Galois extension, i.e., if and only if the canonical map

$$\text{can} : A \otimes_B A \rightarrow A \otimes H, \quad a \otimes_B a' \mapsto aa'_0 \otimes a'_1$$

is bijective. In the literature this is sometimes referred to as the universal QPB. Connections on the universal QPB are given by unital right H -colinear² maps $\omega : H \rightarrow A \otimes A$, such that

$$\text{can} \circ \pi \circ \omega = 1_A \otimes \text{Id}_H,$$

²Here we endow H with the right H -coaction Ad_R and $A \otimes A$ with the diagonal right coaction.

where $\pi: A \otimes A \rightarrow A \otimes_B A$ is the quotient map. We also remark that, when the Hopf algebra H has invertible antipode, the right H -coaction ρ on A converts to a left H -coaction $\lambda(a) = S^{-1}(a_1) \otimes a_0$ for all $a \in A$, making A a left H^{op} -comodule algebra. Then $\text{Id}_A \otimes \rho, \lambda \otimes \text{Id}_A$ make $A \otimes A$ into an H -bicomodule. So, when H has bijective antipode, strong connections on the universal QPB correspond to unital bicomodule maps $\omega: H \rightarrow A \otimes A$ such that $m_A \omega(h) = \varepsilon(h)1_A$ for all $h \in H$ (see [3, Lemma 5.8]). Moreover, when H has bijective antipode, the existence of a strong connection is equivalent to faithfully flatness of A as a left B -module and to the right H -equivariant projectivity of A , i.e., to the fact that there exists a left B -linear and right H -colinear map $s: A \rightarrow B \otimes A$ such that $m \circ s = \text{Id}_A$, where m is the restriction of m_A to $B \otimes A$ [6, Section 6.4]. If $B \subseteq A$ is a cleft extension there exists such an s , namely $s(a) = a_0 j^{-1}(a_1) \otimes j(a_2)$, where j is the cleaving map.

In [3] a QPB is called *regular* if it descends from the universal QPB (i.e., from a Hopf–Galois extension) such that the latter has a strong connection. Hence, a QPB on a cleft extension is always regular in this sense. Moreover, a QPB is called *strong* if $A\Omega^1(B) = \Omega^1(B)A$ as subspaces of $\Omega^1(A)$.

We finish this subsection by discussing noncommutative vector fields, introduced in [4, 5]. In noncommutative geometry one needs to choose a left or right-handedness to express the duality between vector fields and 1-forms. We follow the convention of [18, Definition 4.3], where vector fields are paired from the right with differential forms, since this seems to be more common in the study of QPBs. Given an algebra A and an A -bimodule M , the *left dual* of M is the A -bimodule ${}^*M := {}_A\text{Hom}(M, A)$ of left A -linear maps from M to A , with bimodule structure $(a \cdot f \cdot b)(m) = f(m \cdot a)b$ for $a, b \in A, m \in M$ and $f \in {}^*M$.

Definition 2.25 ([4]). *Given a FODC $(\Omega^1(A), d_A)$ on an algebra A we call*

$$\mathfrak{X}(A) := {}^*\Omega^1(A) = \{X: \Omega^1(A) \rightarrow A \mid X \text{ is left } A\text{-linear}\}$$

the A -bimodule of vector fields on A . We further consider the A -bilinear pairing $\langle \cdot, \cdot \rangle: \Omega^1(A) \otimes_A \mathfrak{X}(A) \rightarrow A$ given by the evaluation and the Lie derivative $\mathcal{L}: \mathfrak{X}(A) \rightarrow \text{End}_{\mathbb{k}}(A)$, which is defined for all $X \in \mathfrak{X}(A)$ by

$$\mathcal{L}_X: a \mapsto \mathcal{L}_X(a) := \langle da, X \rangle.$$

For a Hopf algebra H and a left covariant FODC $(\Omega^1(H), d_H)$ we call a vector field $X: \Omega^1(H) \rightarrow H$ *left-invariant* if $(\text{Id}_H \otimes X) \circ \lambda = \Delta \circ X$, where $\lambda: \Omega^1(H) \rightarrow H \otimes \Omega^1(H)$ is the left H -coaction of $\Omega^1(H)$.

Note that from the right A -linearity of the pairing $\langle \cdot, \cdot \rangle$ we obtain $\mathcal{L}_{X \cdot a}(b) = \mathcal{L}_X(b)a$ and from the Leibniz rule of d_A it follows that \mathcal{L}_X is a noncommutative derivation of A for all $X \in \mathfrak{X}(A)$, in the sense that

$$\mathcal{L}_X(ab) = a\mathcal{L}_X(b) + \mathcal{L}_{b \cdot X}(a)$$

for all $a, b \in A$.

We recall from [26] (see also [3, Lemma 6.1]) that the vector space of left-invariant vector fields on a Hopf algebra H can be identified with the *quantum tangent space*

$$T_\varepsilon H := \{\alpha: \Omega^1(H) \rightarrow \mathbb{k} \mid \alpha(h\gamma) = \varepsilon(h)\alpha(\gamma) \text{ for all } h \in H, \gamma \in \Omega^1(H)\},$$

where $(\Omega^1(H), d_H)$ denotes the corresponding left-covariant FODC on H .

Lemma 2.26. *Let $(\Omega^1(H), d_H)$ be a left-covariant calculus on a Hopf algebra H . Then, there is a 1-1 correspondence between left-invariant vector fields $X: \Omega^1(H) \rightarrow H$ and elements of the quantum tangent space $T_\varepsilon H$ given by*

$$\begin{aligned} \{X \in \mathfrak{X}(H) \mid X \text{ is left-invariant}\} &\xleftarrow{1:1} T_\varepsilon H \\ X &\longmapsto \varepsilon \circ X, \\ (\text{Id}_H \otimes \alpha) \circ \lambda &\longleftarrow \alpha, \end{aligned}$$

where $\lambda: \Omega^1(H) \rightarrow H \otimes \Omega^1(H)$ is the left H -coaction of $\Omega^1(H)$. Moreover, if the antipode of H is bijective then $T_\varepsilon H = \text{Hom}_{\mathbb{k}}({}^{\text{co}H}\Omega^1(H), \mathbb{k})$.

Note that, if ${}^{\text{co}H}\Omega^1(H)$ is finite-dimensional and the antipode of H is bijective, we can structure $T_\varepsilon H = \text{Hom}_{\mathbb{k}}({}^{\text{co}H}\Omega^1(H), \mathbb{k})$ as a right H -comodule via $T_\varepsilon H \rightarrow (T_\varepsilon H) \otimes H, \alpha \mapsto \alpha_0 \otimes \alpha_1$, where $\alpha_0(\gamma)\alpha_1 = \alpha(\gamma_0)S^{-1}(\gamma_1)$ for any $\gamma \in {}^{\text{co}H}\Omega^1(H)$, see e.g. [18, Definition 4.4].

Remark 2.27. *Given a right H -comodule algebra A with differentiable right H -coaction $\rho: A \rightarrow A \otimes H$, for any $\alpha \in T_\varepsilon H$ there is a vector field on A given by $(\text{Id}_A \otimes \alpha) \circ \pi_2 \circ \widehat{\rho}: \Omega^1(A) \rightarrow A$, where $\pi_2: \Omega^1(A \otimes H) \rightarrow A \otimes \Omega^1(H)$ is the canonical projection, cf. [3, Corollary 6.2 and page 490]. In the case of a QPB with vertical map*

ver : $\Omega^1(A) \rightarrow A \otimes {}^{\text{co}H}\Omega^1(H)$, given $\alpha \in T_\varepsilon H$ (the antipode of H is assumed bijective), one then obtains a vector field on A

$$\bar{\alpha} := (\text{Id}_A \otimes \alpha) \circ \text{ver} : \Omega^1(A) \rightarrow A,$$

which is vertical in the sense that it vanishes on $\Omega^1(A) \otimes H$. In the next section we will show that these vector fields are exactly the vertical ones which satisfy an additional property in the framework of the crossed product algebra $B\#_\sigma H$.

3 The crossed product calculus

Given a left H -module algebra B , there is a canonical FODC on the smash product algebra $B\#H$ which is right H -covariant, starting from a bicovariant FODC on H and an H -module FODC on B (see [18, Theorem 2.7]). In Section 3.1 we generalize this construction to the case of the crossed product algebra $B\#_\sigma H$, when B is a σ -twisted left H -module algebra. A characterization of the smash product calculus follows in Section 3.2. Higher order forms for the crossed product algebra are constructed in Section 3.3, the QPB structure of crossed product forms and corresponding bimodule covariant derivatives on associated bundles are discussed in Section 3.4, while we prove the 1:1 correspondence of connections and connection 1-forms of the crossed product calculus in Section 3.5.

3.1 Crossed product 1-forms

In this subsection H is a Hopf algebra and B a σ -twisted left H -module algebra. We introduce a right H -covariant FODC on the crossed product algebra $B\#_\sigma H$. First, we define a twisted analog of a B -bimodule.

Definition 3.1. Let M be in ${}_B\mathfrak{M}_B$. We say that M is a σ -twisted B -bimodule if there exists a linear map $\cdot : H \otimes M \rightarrow M$ such that for all $m \in M$, $h, h' \in H$ and $b, b' \in B$

$$i.) \quad 1_H \cdot m = m$$

$$ii.) \quad h \cdot (bmb') = (h_1 \cdot b)(h_2 \cdot m)(h_3 \cdot b')$$

$$iii.) \quad h \cdot (h' \cdot m) = \sigma(h_1 \otimes h'_1)(h_2 h'_2 \cdot m) \sigma^{-1}(h_3 \otimes h'_3)$$

If σ is the trivial 2-cocycle we recover the notion of left H -equivariant B -bimodule (see Definition 2.2). Indeed, in this case B becomes a left H -module algebra and in the above definition the first and third equations imply that M is in ${}_H\mathfrak{M}$ and the second equation confers that the left and right B -actions of M are left H -linear, so that M is in ${}_B({}_H\mathfrak{M})_B$.

Lemma 3.2. Let $(\Omega^1(B), d_B)$ be a FODC on B such that there exists a linear map $\cdot : H \otimes \Omega^1(B) \rightarrow \Omega^1(B)$ with

$$h \cdot (bd_B b') = (h_1 \cdot b)(h_2 \cdot d_B b') \tag{12}$$

$$d_B(h \cdot b) = h \cdot d_B b \tag{13}$$

$$d_B \circ \sigma = 0 \tag{14}$$

for every $h \in H$ and $b, b' \in B$. Then $\Omega^1(B)$ is a σ -twisted B -bimodule.

Proof. Since every $\beta \in \Omega^1(B)$ is a finite sum $\beta = b^i d_B b'_i$ with $b^i, b'_i \in B$, it is sufficient to show *i.*, *ii.*) and *iii.*) of Definition 3.1 for $bd_B b'$, with $b, b' \in B$. By assumption $\Omega^1(B)$ is a B -bimodule and we obtain *i.*, since

$$1_H \cdot (bd_B b') \stackrel{(12)}{=} (1_H \cdot b)(1_H \cdot d_B b') \stackrel{(13)}{=} bd_B(1_H \cdot b') = bd_B b'.$$

Furthermore, we have that for all $b, b', b'' \in B$ and $h \in H$

$$\begin{aligned} h \cdot ((bd_B b')b'') &= h \cdot (b((d_B b')b'')) = h \cdot (bd_B(b'b'') - bb'd_B b'') \\ &\stackrel{(12)}{=} (h_1 \cdot b)(h_2 \cdot d_B(b'b'')) - (h_1 \cdot bb')(h_2 \cdot d_B b'') \\ &\stackrel{(13)}{=} (h_1 \cdot b)d_B(h_2 \cdot b'b'') - (h_1 \cdot b)(h_2 \cdot b')(h_3 \cdot d_B b'') \\ &\stackrel{(13)}{=} (h_1 \cdot b)d_B((h_2 \cdot b')(h_3 \cdot b'')) - (h_1 \cdot b)(h_2 \cdot b')d_B(h_3 \cdot b'') \\ &= (h_1 \cdot b)d_B(h_2 \cdot b')(h_3 \cdot b'') \stackrel{(13)}{=} (h_1 \cdot b)(h_2 \cdot d_B b')(h_3 \cdot b'') \\ &\stackrel{(12)}{=} (h_1 \cdot bd_B b')(h_2 \cdot b''), \end{aligned}$$

which implies

$$\begin{aligned}
h \cdot (b(b'd_B b'')b''') &= h \cdot ((bb'd_B b'')b''') = (h_1 \cdot bb'd_B b'')(h_2 \cdot b''') \\
&= (h_1 \cdot bb')(h_2 \cdot d_B b'')(h_3 \cdot b''') \\
&= (h_1 \cdot b)(h_2 \cdot b')(h_3 \cdot d_B b'')(h_4 \cdot b''') \\
&= (h_1 \cdot b)(h_2 \cdot b'd_B b'')(h_3 \cdot b'''),
\end{aligned}$$

i.e., *ii.*) holds. Finally, we show *iii.*). Observe that, given (14), we have

$$\begin{aligned}
d_B(\sigma^{-1}(h \otimes h')) &= \varepsilon(h_1 h'_1) d_B(\sigma^{-1}(h_2 \otimes h'_2)) = \sigma^{-1}(h_1 \otimes h'_1) \sigma(h_2 \otimes h'_2) d_B(\sigma^{-1}(h_3 \otimes h'_3)) \\
&= \sigma^{-1}(h_1 \otimes h'_1) d_B(\sigma(h_2 \otimes h'_2)) \sigma^{-1}(h_3 \otimes h'_3) + \sigma^{-1}(h_1 \otimes h'_1) \sigma(h_2 \otimes h'_2) d_B(\sigma^{-1}(h_3 \otimes h'_3)) \\
&= \sigma^{-1}(h_1 \otimes h'_1) d_B(\sigma(h_2 \otimes h'_2) \sigma^{-1}(h_3 \otimes h'_3)) = \sigma^{-1}(h_1 \otimes h'_1) d_B(\varepsilon(h_2 h'_2) 1_B) = 0.
\end{aligned}$$

Thus, we obtain that

$$\begin{aligned}
h \cdot (h' \cdot b d_B b') &= (h_1 \cdot (h'_1 \cdot b))(h_2 \cdot (h'_2 \cdot d_B b')) = (h_1 \cdot (h'_1 \cdot b)) d_B(h_2 \cdot (h'_2 \cdot b')) \\
&= \sigma(h_1 \otimes h'_1) (h_2 h'_2 \cdot b) \sigma^{-1}(h_3 \otimes h'_3) d_B(\sigma(h_4 \otimes h'_4) (h_5 h'_5 \cdot b') \sigma^{-1}(h_6 \otimes h'_6)) \\
&= \sigma(h_1 \otimes h'_1) (h_2 h'_2 \cdot b) \sigma^{-1}(h_3 \otimes h'_3) \sigma(h_4 \otimes h'_4) d_B(h_5 h'_5 \cdot b') \sigma^{-1}(h_6 \otimes h'_6) \\
&= \sigma(h_1 \otimes h'_1) (h_2 h'_2 \cdot b) d_B(h_3 h'_3 \cdot b') \sigma^{-1}(h_4 \otimes h'_4) \\
&= \sigma(h_1 \otimes h'_1) (h_2 h'_2 \cdot b) (h_3 h'_3 \cdot d_B b') \sigma^{-1}(h_4 \otimes h'_4) \\
&= \sigma(h_1 \otimes h'_1) (h_2 h'_2 \cdot (b d_B b')) \sigma^{-1}(h_3 \otimes h'_3),
\end{aligned}$$

where in the fourth equation we used the Leibniz rule and (14). \square

Note that, if the map $\cdot : H \otimes \Omega^1(B) \rightarrow \Omega^1(B)$ in Lemma 3.2 exists, then it is unique, since by (12) and (13) it is completely determined by $h \cdot (b d_B b') = (h_1 \cdot b) d_B(h_2 \cdot b')$ for all $h \in H$ and $b, b' \in B$.

Definition 3.3. Let $(\Omega^1(B), d_B)$ be a FODC on a σ -twisted left H -module algebra B . We say that it is a σ -twisted H -module differential calculus if there exists a linear map $\cdot : H \otimes \Omega^1(B) \rightarrow \Omega^1(B)$ such that (12), (13) and (14) of Lemma 3.2 are satisfied. In particular, $\Omega^1(B)$ is a σ -twisted B -bimodule.

Remark 3.4. To explain the nomenclature of the previous definition, recall that in [18, Definition 2.5] an H -module differential calculus on a left H -module algebra B is a FODC $(\Omega^1(B), d_B)$, where $\Omega^1(B)$ is in ${}_B(H\mathfrak{M})$ and d_B is in ${}_H\mathfrak{M}$; it follows that $\Omega^1(B)$ is in ${}_B(H\mathfrak{M})_B$. In Definition 3.3 we recover this case if σ is the trivial 2-cocycle; as already mentioned, Definition 3.1 implies that $\Omega^1(B)$ is in ${}_B(H\mathfrak{M})_B$, while (13) of Lemma 3.2 implies that d_B is left H -linear. Observe that (14) is trivially satisfied when σ is trivial, while we will show that, when σ is not trivial, it is a necessary condition in order to construct the FODC on the crossed product algebra $B \#_\sigma H$.

Lemma 3.5. If M is a σ -twisted B -bimodule and N is in ${}_H^H\mathfrak{M}_H^H$, then $(M \otimes H) \oplus (B \otimes N)$ is in ${}_{B \#_\sigma H} \mathfrak{M}_{B \#_\sigma H}^H$ with respect to the $B \#_\sigma H$ -actions determined for all $b, b' \in B$, $h, h' \in H$, $\beta \in M$ and $\gamma \in N$ by

$$(b' \otimes h') \otimes (\beta \otimes h + b \otimes \gamma) \mapsto b'(h'_1 \cdot \beta) \sigma(h'_2 \otimes h_1) \otimes h'_3 h_2 + b'(h'_1 \cdot b) \sigma(h'_2 \otimes \gamma_{-1}) \otimes h'_3 \gamma_0 \quad (15)$$

$$(\beta \otimes h + b \otimes \gamma) \otimes (b' \otimes h') \mapsto \beta(h_1 \cdot b') \sigma(h_2 \otimes h'_1) \otimes h_3 h'_2 + b(\gamma_{-2} \cdot b') \sigma(\gamma_{-1} \otimes h'_1) \otimes \gamma_0 h'_2 \quad (16)$$

and right H -coaction

$$\rho' : C \rightarrow C \otimes H, \quad \beta \otimes h + b \otimes \gamma \mapsto \beta \otimes h_1 \otimes h_2 + b \otimes \gamma_0 \otimes \gamma_1. \quad (17)$$

Proof. Let $C := (M \otimes H) \oplus (B \otimes N)$ and fix $b, b', b'' \in B$, $h, h', h'' \in H$, $\beta \in M$ and $\gamma \in N$. We first show that C is a $B \#_\sigma H$ -bimodule. Indeed, using the properties of the σ -twisted B -bimodule M and that B is a σ -twisted left H -module algebra we obtain

$$\begin{aligned}
(b'' \otimes h'') \cdot ((b' \otimes h') \cdot (\beta \otimes h)) &= (b'' \otimes h'') \cdot (b'(h'_1 \cdot \beta) \sigma(h'_2 \otimes h_1) \otimes h'_3 h_2) \\
&= b''(h''_1 \cdot b'(h'_1 \cdot \beta) \sigma(h'_2 \otimes h_1)) \sigma(h''_2 \otimes h'_3 h_2) \otimes h''_3 h'_4 h_3 \\
&= b''(h''_1 \cdot b')(h''_2 \cdot (h'_1 \cdot \beta)) (h''_3 \cdot \sigma(h'_2 \otimes h_1)) \sigma(h''_4 \otimes h'_3 h_2) \otimes h''_5 h'_4 h_3 \\
&= b''(h''_1 \cdot b') \sigma(h''_2 \otimes h'_1) (h''_3 h'_2 \cdot \beta) \sigma^{-1}(h''_4 \otimes h'_3) (h''_5 \cdot \sigma(h'_4 \otimes h_1)) \sigma(h''_6 \otimes h'_5 h_2) \otimes h''_7 h'_6 h_3 \\
&= b''(h''_1 \cdot b') \sigma(h''_2 \otimes h'_1) (h''_3 h'_2 \cdot \beta) \sigma^{-1}(h''_4 \otimes h'_3) \sigma(h''_5 \otimes h'_4) \sigma(h''_6 h'_5 \otimes h_1) \otimes h''_7 h'_6 h_2 \\
&= b''(h''_1 \cdot b') \sigma(h''_2 \otimes h'_1) (h''_3 h'_2 \cdot \beta) \sigma(h''_4 h'_3 \otimes h_1) \otimes h''_5 h'_4 h_2 \\
&= (b''(h''_1 \cdot b') \sigma(h''_2 \otimes h'_1) \otimes h''_3 h'_2) \cdot (\beta \otimes h) \\
&= ((b'' \otimes h'') (b' \otimes h')) \cdot (\beta \otimes h)
\end{aligned}$$

and

$$(1_B \otimes 1_H) \cdot (\beta \otimes h) = 1_B(1_H \cdot \beta)\sigma(1_H \otimes h_1) \otimes 1_H h_2 = \beta \varepsilon(h_1) \otimes h_2 = \beta \otimes h.$$

Similarly it follows that

$$\begin{aligned} (b'' \otimes h'') \cdot ((b' \otimes h') \cdot (b \otimes \gamma)) &= (b'' \otimes h'') \cdot (b'(h'_1 \cdot b)\sigma(h'_2 \otimes \gamma_{-1}) \otimes h'_3 \gamma_0) \\ &= b''(h'_1 \cdot b')(h'_1 \cdot b)\sigma(h'_2 \otimes \gamma_{-1})\sigma(h''_2 \otimes (h'_3 \gamma_0)_{-1}) \otimes h''_3 (h'_3 \gamma_0)_0 \\ &= b''(h'_1 \cdot b')(h''_2 \cdot (h'_1 \cdot b))(h''_3 \cdot \sigma(h'_2 \otimes \gamma_{-2}))\sigma(h''_4 \otimes h'_3 \gamma_{-1}) \otimes h''_5 h'_4 \gamma_0 \\ &= b''(h'_1 \cdot b')\sigma(h''_2 \otimes h'_1)(h''_3 h'_2 \cdot b)\sigma^{-1}(h''_4 \otimes h'_3)(h''_5 \cdot \sigma(h'_4 \otimes \gamma_{-2}))\sigma(h''_6 \otimes h'_5 \gamma_{-1}) \otimes h''_7 h'_6 \gamma_0 \\ &= b''(h'_1 \cdot b')\sigma(h''_2 \otimes h'_1)(h''_3 h'_2 \cdot b)\sigma^{-1}(h''_4 \otimes h'_3)\sigma(h''_5 \otimes h'_4)\sigma(h''_6 h'_5 \otimes \gamma_{-1}) \otimes h''_7 h'_6 \gamma_0 \\ &= b''(h'_1 \cdot b')\sigma(h''_2 \otimes h'_1)(h''_3 h'_2 \cdot b)\sigma(h''_4 h'_3 \otimes \gamma_{-1}) \otimes h''_5 h'_4 \gamma_0 \\ &= (b''(h'_1 \cdot b')\sigma(h''_2 \otimes h'_1) \otimes h''_3 h'_2) \cdot (b \otimes \gamma) \\ &= ((b'' \otimes h'')(b' \otimes h')) \cdot (b \otimes \gamma) \end{aligned}$$

and

$$(1_B \otimes 1_H) \cdot (b \otimes \gamma) = 1_B(1_H \cdot b)\sigma(1_H \otimes \gamma_{-1}) \otimes 1_H \gamma_0 = b \varepsilon(\gamma_{-1}) \otimes \gamma_0 = b \otimes \gamma.$$

Thus, C is a left $B\#_\sigma H$ -module. The fact that C is a right $B\#_\sigma H$ -module follows in complete analogy. Moreover, the two actions are compatible, indeed

$$\begin{aligned} ((b' \otimes h') \cdot (\beta \otimes h)) \cdot (b'' \otimes h'') &= (b'(h'_1 \cdot \beta)\sigma(h'_2 \otimes h_1) \otimes h'_3 h_2) \cdot (b'' \otimes h'') \\ &= b'(h'_1 \cdot \beta)\sigma(h'_2 \otimes h_1)(h'_3 h_2 \cdot b'')\sigma(h'_4 h_3 \otimes h''_1) \otimes h'_5 h_4 h''_2 \\ &= b'(h'_1 \cdot \beta)\sigma(h'_2 \otimes h_1)(h'_3 h_2 \cdot b'')\sigma^{-1}(h'_4 \otimes h_3)\sigma(h'_5 \otimes h_4)\sigma(h'_6 h_5 \otimes h''_1) \otimes h'_7 h_6 h''_2 \\ &= b'(h'_1 \cdot \beta)\sigma(h'_2 \otimes h_1)(h'_3 h_2 \cdot b'')\sigma^{-1}(h'_4 \otimes h_3)(h'_5 \cdot \sigma(h_4 \otimes h''_1))\sigma(h'_6 \otimes h_5 h''_2) \otimes h'_7 h_6 h''_3 \\ &= b'(h'_1 \cdot \beta)(h'_2 \cdot (h_1 \cdot b''))(h'_3 \cdot \sigma(h_2 \otimes h''_1))\sigma(h'_4 \otimes h_3 h''_2) \otimes h'_5 h_4 h''_3 \\ &= b'(h'_1 \cdot \beta)(h_1 \cdot b'')\sigma(h_2 \otimes h''_1)\sigma(h'_2 \otimes h_3 h''_2) \otimes h'_3 h_4 h''_3 \\ &= (b' \otimes h') \cdot (\beta(h_1 \cdot b'')\sigma(h_2 \otimes h''_1) \otimes h_3 h''_2) \\ &= (b' \otimes h') \cdot ((\beta \otimes h) \cdot (b'' \otimes h'')) \end{aligned}$$

and

$$\begin{aligned} ((b' \otimes h') \cdot (b \otimes \gamma)) \cdot (b'' \otimes h'') &= (b'(h'_1 \cdot b)\sigma(h'_2 \otimes \gamma_{-1}) \otimes h'_3 \gamma_0) \cdot (b'' \otimes h'') \\ &= b'(h'_1 \cdot b)\sigma(h'_2 \otimes \gamma_{-1})((h'_3 \gamma_0)_{-2} \cdot b'')\sigma((h'_3 \gamma_0)_{-1} \otimes h''_1) \otimes (h'_3 \gamma_0)_0 h''_2 \\ &= b'(h'_1 \cdot b)\sigma(h'_2 \otimes \gamma_{-3})(h'_3 \gamma_{-2} \cdot b'')\sigma(h'_4 \gamma_{-1} \otimes h''_1) \otimes h'_5 \gamma_0 h''_2 \\ &= b'(h'_1 \cdot b)\sigma(h'_2 \otimes \gamma_{-5})(h'_3 \gamma_{-4} \cdot b'')\sigma^{-1}(h'_4 \otimes \gamma_{-3})\sigma(h'_5 \otimes \gamma_{-2})\sigma(h'_6 \gamma_{-1} \otimes h''_1) \otimes h'_7 \gamma_0 h''_2 \\ &= b'(h'_1 \cdot b)\sigma(h'_2 \otimes \gamma_{-5})(h'_3 \gamma_{-4} \cdot b'')\sigma^{-1}(h'_4 \otimes \gamma_{-3})(h'_5 \cdot \sigma(\gamma_{-2} \otimes h''_1))\sigma(h'_6 \otimes \gamma_{-1} h''_2) \otimes h'_7 \gamma_0 h''_3 \\ &= b'(h'_1 \cdot b)(h'_2 \cdot (\gamma_{-3} \cdot b''))(h'_3 \cdot \sigma(\gamma_{-2} \otimes h''_1))\sigma(h'_4 \otimes \gamma_{-1} h''_2) \otimes h'_5 \gamma_0 h''_3 \\ &= b'(h'_1 \cdot b(\gamma_{-2} \cdot b''))\sigma(\gamma_{-1} \otimes h''_1)\sigma(h'_2 \otimes (\gamma_0 h''_2)_{-1}) \otimes h'_3 (\gamma_0 h''_2)_0 \\ &= (b' \otimes h') \cdot (b(\gamma_{-2} \cdot b'')\sigma(\gamma_{-1} \otimes h''_1) \otimes \gamma_0 h''_2) \\ &= (b' \otimes h') \cdot ((b \otimes \gamma) \cdot (b'' \otimes h'')). \end{aligned}$$

Hence, we have that C is a $B\#_\sigma H$ -bimodule. Recall that $B\#_\sigma H$ is a right H -comodule algebra with $\rho : B\#_\sigma H \rightarrow (B\#_\sigma H) \otimes H$, $b \otimes h \mapsto b \otimes h_1 \otimes h_2$. Clearly, C is a right H -comodule via

$$\rho' : C \rightarrow C \otimes H, \quad \beta \otimes h + b \otimes \gamma \mapsto \beta \otimes h_1 \otimes h_2 + b \otimes \gamma_0 \otimes \gamma_1 \quad (18)$$

and we can show that there is compatibility between the right H -coaction and the two $B\#_\sigma H$ -actions, so that C is in ${}_{B\#_\sigma H}\mathfrak{M}_{B\#_\sigma H}^H$. In fact, we have

$$\begin{aligned} \rho'((\beta \otimes h + b \otimes \gamma) \cdot (b' \otimes h')) &= \rho'(\beta(h_1 \cdot b')\sigma(h_2 \otimes h'_1) \otimes h_3 h'_2 + b(\gamma_{-2} \cdot b')\sigma(\gamma_{-1} \otimes h'_1) \otimes \gamma_0 h'_2) \\ &= \beta(h_1 \cdot b')\sigma(h_2 \otimes h'_1) \otimes h_3 h'_2 \otimes h_4 h'_3 + b(\gamma_{-2} \cdot b')\sigma(\gamma_{-1} \otimes h'_1) \otimes \gamma_0 h'_2 \otimes \gamma_1 h'_3 \\ &= (\beta \otimes h_1) \cdot (b' \otimes h'_1) \otimes h_2 h'_2 + (b \otimes \gamma_0) \cdot (b' \otimes h'_1) \otimes \gamma_1 h'_2 \end{aligned}$$

and

$$\begin{aligned} \rho'((b' \otimes h') \cdot (\beta \otimes h + b \otimes \gamma)) &= \rho'(b'(h'_1 \cdot \beta)\sigma(h'_2 \otimes h_1) \otimes h'_3 h_2 + b'(h'_1 \cdot b)\sigma(h'_2 \otimes \gamma_{-1}) \otimes h'_3 \gamma_0) \\ &= b'(h'_1 \cdot \beta)\sigma(h'_2 \otimes h_1) \otimes h'_3 h_2 \otimes h'_4 h_3 + b'(h'_1 \cdot b)\sigma(h'_2 \otimes \gamma_{-1}) \otimes h'_3 \gamma_0 \otimes h'_4 \gamma_1 \\ &= (b' \otimes h'_1) \cdot (\beta \otimes h_1) \otimes h'_2 h_2 + (b' \otimes h'_1) \cdot (b \otimes \gamma_0) \otimes h'_2 \gamma_1. \end{aligned}$$

and the thesis follows. \square

Corollary 3.6. Given $(\Omega^1(H), d_H)$ a bicovariant FODC on H and $(\Omega^1(B), d_B)$ a σ -twisted H -module FODC on B , $(\Omega^1(B) \otimes H) \oplus (B \otimes \Omega^1(H))$ is in ${}_{B\#_\sigma H} \mathfrak{M}_{B\#_\sigma H}^H$ with left and right $B\#_\sigma H$ -actions given by (15) and (16), respectively and right H -coaction given by (17).

Finally, we construct the announced right H -covariant FODC on $B\#_\sigma H$, starting from a bicovariant FODC on H and a σ -twisted H -module FODC on B . We call it the *crossed product calculus* on $B\#_\sigma H$.

Theorem 3.7. Given $(\Omega^1(H), d_H)$ a bicovariant FODC on H and $(\Omega^1(B), d_B)$ a σ -twisted H -module FODC on B we obtain a right H -covariant FODC $(\Omega^1(B\#_\sigma H), d_{\#_\sigma})$ on $B\#_\sigma H$, where

$$\Omega^1(B\#_\sigma H) := (\Omega^1(B) \otimes H) \oplus (B \otimes \Omega^1(H)) \text{ and } d_{\#_\sigma} : B\#_\sigma H \rightarrow \Omega^1(B\#_\sigma H), \quad b \otimes h \mapsto d_B b \otimes h + b \otimes d_H h.$$

Moreover, the right H -coaction on $B\#_\sigma H$ is differentiable with respect to this calculus.

Proof. In Corollary 3.6 we have shown that $\Omega^1(B\#_\sigma H) := (\Omega^1(B) \otimes H) \oplus (B \otimes \Omega^1(H))$ is in ${}_{B\#_\sigma H} \mathfrak{M}_{B\#_\sigma H}^H$. In a next step we prove that $d_{\#_\sigma}$, as defined above, satisfies the Leibniz rule. Indeed, we have

$$\begin{aligned} (d_{\#_\sigma}(b \otimes h)) \cdot (b' \otimes h') + (b \otimes h) \cdot (d_{\#_\sigma}(b' \otimes h')) & \\ &= (d_B b \otimes h + b \otimes d_H h) \cdot (b' \otimes h') + (b \otimes h) \cdot (d_B b' \otimes h' + b' \otimes d_H h') \\ &= (d_B b)(h_1 \cdot b') \sigma(h_2 \otimes h'_1) \otimes h_3 h'_2 + b((d_H h)_{-2} \cdot b') \sigma((d_H h)_{-1} \otimes h'_1) \otimes (d_H h)_0 h'_2 \\ &\quad + b(h_1 \cdot d_B b') \sigma(h_2 \otimes h'_1) \otimes h_3 h'_2 + b(h_1 \cdot b') \sigma(h_2 \otimes (d_H h')_{-1}) \otimes h_3 (d_H h')_0 \\ &= (d_B b)(h_1 \cdot b') \sigma(h_2 \otimes h'_1) \otimes h_3 h'_2 + b(h_1 \cdot b') \sigma(h_2 \otimes h'_1) \otimes (d_H h_3) h'_2 \\ &\quad + b d_B (h_1 \cdot b') \sigma(h_2 \otimes h'_1) \otimes h_3 h'_2 + b(h_1 \cdot b') \sigma(h_2 \otimes h'_1) \otimes h_3 d_H h'_2 \\ &= b(h_1 \cdot b') \sigma(h_2 \otimes h'_1) \otimes d_H (h_3 h'_2) + d_B (b(h_1 \cdot b')) \sigma(h_2 \otimes h'_1) \otimes h_3 h'_2 \\ &= b(h_1 \cdot b') \sigma(h_2 \otimes h'_1) \otimes d_H (h_3 h'_2) + d_B (b(h_1 \cdot b')) \sigma(h_2 \otimes h'_1) \otimes h_3 h'_2 \\ &= d_{\#_\sigma}((b \otimes h)(b' \otimes h')), \end{aligned}$$

where we used the fact that d_H is left H -colinear and (13) in the third equality, Leibniz rules of d_H and d_B in the fourth and (14) in the fifth.

Furthermore, $\Omega^1(B\#_\sigma H)$ is generated by $d_{\#_\sigma}(B\#_\sigma H)$ as left $B\#_\sigma H$ -module. In order to prove this it is sufficient to show that the left $B\#_\sigma H$ -submodule of $\Omega^1(B\#_\sigma H)$ generated by $d_{\#_\sigma}(B\#_\sigma H)$ contains $bd_B b' \otimes h$ and $b \otimes hd_H h'$ for every $b, b' \in B$ and $h, h' \in H$. The first element is obtained by

$$\begin{aligned} (b \otimes 1_H) \cdot d_{\#_\sigma}(b' \otimes h) - (bb' \otimes 1_H) \cdot d_{\#_\sigma}(1_B \otimes h) & \\ &= (b \otimes 1_H) \cdot (d_B b' \otimes h + b' \otimes d_H h) - (bb' \otimes 1_H) \cdot (1_B \otimes d_H h) \\ &= b(1_H \cdot d_B b') \sigma(1_H \otimes h_1) \otimes 1_H h_2 + b(1_H \cdot b') \sigma(1_H \otimes (d_H h)_{-1}) \otimes 1_H (d_H h)_0 \\ &\quad - bb'(1_H \cdot 1_B) \sigma(1_H \otimes (d_H h)_{-1}) \otimes 1_H (d_H h)_0 \\ &= bd_B b' \varepsilon(h_1) \otimes h_2 = bd_B b' \otimes h, \end{aligned}$$

while for the second element we have that

$$\begin{aligned} (b\sigma^{-1}(h_1 \otimes h'_1) \otimes h_2) \cdot d_{\#_\sigma}(1_B \otimes h'_2) &= (b\sigma^{-1}(h_1 \otimes h'_1) \otimes h_2) \cdot (1_B \otimes d_H h'_2) \\ &= b\sigma^{-1}(h_1 \otimes h'_1)(h_2 \cdot 1_B) \sigma(h_3 \otimes (d_H h'_2)_{-1}) \otimes h_4 (d_H h'_2)_0 \\ &= b\sigma^{-1}(h_1 \otimes h'_1) \varepsilon(h_2) \sigma(h_3 \otimes h'_2) \otimes h_4 d_H h'_3 \\ &= b\sigma^{-1}(h_1 \otimes h'_1) \sigma(h_2 \otimes h'_2) \otimes h_3 d_H h'_3 \\ &= b\varepsilon(h_1) \varepsilon(h'_1) \otimes h_2 d_H h'_2 = b \otimes hd_H h', \end{aligned}$$

using again that d_H is left H -colinear. Hence, the surjectivity condition is shown and $(\Omega^1(B\#_\sigma H), d_{\#_\sigma})$ is a FODC on the crossed product algebra $B\#_\sigma H$.

Now, denoting by ρ the right H -coaction of $B\#_\sigma H$, define $\widehat{\rho} : \Omega^1(B\#_\sigma H) \rightarrow \Omega^1((B\#_\sigma H) \otimes H)$ as $\beta \otimes h + b \otimes \gamma \mapsto \beta \otimes h_1 \otimes h_2 + b \otimes \gamma_0 \otimes \gamma_1 + b \otimes \gamma_{-1} \otimes \gamma_0$. This is a morphism in ${}_{B\#_\sigma H} \mathfrak{M}_{B\#_\sigma H}$, since

$$\begin{aligned} \widehat{\rho}(b' \otimes h') \cdot (\beta \otimes h + b \otimes \gamma) &= \widehat{\rho}(b'(h'_1 \cdot \beta) \sigma(h'_2 \otimes h_1) \otimes h'_3 h_2 + b'(h'_1 \cdot b) \sigma(h'_2 \otimes \gamma_{-1}) \otimes h'_3 \gamma_0) \\ &= b'(h'_1 \cdot \beta) \sigma(h'_2 \otimes h_1) \otimes h'_3 h_2 \otimes h'_4 h_3 + b'(h'_1 \cdot b) \sigma(h'_2 \otimes \gamma_{-1}) \otimes h'_3 \gamma_0 \otimes h'_4 \gamma_1 \\ &\quad + b'(h'_1 \cdot b) \sigma(h'_2 \otimes \gamma_{-2}) \otimes h'_3 \gamma_{-1} \otimes h'_4 \gamma_0 \\ &= (b' \otimes h'_1)(\beta \otimes h_1) \otimes h'_2 h_2 + (b' \otimes h'_1)(b \otimes \gamma_0) \otimes h'_2 \gamma_1 + (b' \otimes h'_1)(b \otimes \gamma_{-1}) \otimes h'_2 \gamma_0 \\ &= (b' \otimes h') \cdot \widehat{\rho}(\beta \otimes h + b \otimes \gamma) \end{aligned}$$

and, analogously, $\widehat{\rho}((\beta \otimes h + b \otimes \gamma) \cdot (b' \otimes h')) = \widehat{\rho}(\beta \otimes h + b \otimes \gamma) \cdot (b' \otimes h')$. Furthermore, we have

$$\begin{aligned}\widehat{\rho}d_{\#_\sigma}(b \otimes h) &= d_B b \otimes h_1 \otimes h_2 + b \otimes (d_H h)_0 \otimes (d_H h)_1 + b \otimes (d_H h)_{-1} \otimes (d_H h)_0 \\ &= d_B b \otimes h_1 \otimes h_2 + b \otimes d_H h_1 \otimes h_2 + b \otimes h_1 \otimes d_H h_2 = d_{\#_\sigma}(b \otimes h_1) \otimes h_2 + b \otimes h_1 \otimes d_H h_2 \\ &= d_{\otimes} \rho(b \otimes h),\end{aligned}$$

hence ρ is differentiable. Note that, from Remark 2.21, we automatically have that $(\Omega^1(B\#_\sigma H), d_{\#_\sigma})$ is right H -covariant such that, denoting by $\pi_1 : \Omega^1((B\#_\sigma H) \otimes H) \rightarrow \Omega^1(B\#_\sigma H) \otimes H$ the canonical projection, the right H -coaction of $\Omega^1(B\#_\sigma H)$ is exactly $\rho' := \pi_1 \circ \widehat{\rho} : \beta \otimes h + b \otimes \gamma \mapsto \beta \otimes h_1 \otimes h_2 + b \otimes \gamma_0 \otimes \gamma_1$. On the other hand, the right H -covariance of the calculus also follows explicitly from the computation

$$\begin{aligned}\rho'd_{\#_\sigma}(b \otimes h) &= \rho'(d_B b \otimes h + b \otimes d_H h) = d_B b \otimes h_1 \otimes h_2 + b \otimes (d_H h)_0 \otimes (d_H h)_1 = d_B b \otimes h_1 \otimes h_2 + b \otimes d_H h_1 \otimes h_2 \\ &= d_{\#_\sigma}(b \otimes h_1) \otimes h_2 = (d_{\#_\sigma} \otimes \text{Id}_H)\rho(b \otimes h),\end{aligned}$$

using that d_H is right H -colinear. \square

As one easily verifies, the above crossed product calculus $(\Omega^1(B\#_\sigma H), d_{\#_\sigma})$ reduces to the smash product calculus $(\Omega^1(B\#H), d_{\#})$ of [18, Theorem 2.7] in case of a trivial extension, where σ is the trivial 2-cocycle.

We conclude this subsection by showing that the condition $d_B \circ \sigma = 0$ is necessary in case we want $(\Omega^1(B\#_\sigma H), d_{\#_\sigma})$ to be a FODC on $B\#_\sigma H$ starting from a bicovariant FODC $(\Omega^1(H), d_H)$ on H and a FODC $(\Omega^1(B), d_B)$ on B such that $\Omega^1(B)$ is a σ -twisted B -bimodule.

Proposition 3.8. *Let $(\Omega^1(B), d_B)$ be a FODC on B such that $\Omega^1(B)$ is a σ -twisted B -bimodule and $(\Omega^1(H), d_H)$ a bicovariant FODC on H such that $(\Omega^1(B\#_\sigma H), d_{\#_\sigma})$, as defined in Theorem 3.7, is a FODC on $B\#_\sigma H$. Then $d_B \circ \sigma = 0$.*

Proof. Since $\Omega^1(B)$ is σ -twisted B -bimodule and $\Omega^1(H)$ is in ${}^H_H\mathfrak{M}_H^H$, then $\Omega^1(B\#_\sigma H)$ is in ${}_{B\#_\sigma H}{}^H\mathfrak{M}_{B\#_\sigma H}^H$ by Lemma 3.5. Since $d_{\#_\sigma}$ satisfies the Leibniz rule by assumption we obtain

$$\begin{aligned}d_B(\sigma(h_1 \otimes h'_1)) \otimes h_2 h'_2 + \sigma(h_1 \otimes h'_1) \otimes d_H(h_2 h'_2) &= d_{\#_\sigma}(\sigma(h_1 \otimes h'_1) \otimes h_2 h'_2) = d_{\#_\sigma}((1_B \otimes h)(1_B \otimes h')) \\ &= (d_{\#_\sigma}(1_B \otimes h))(1_B \otimes h') + (1_B \otimes h)d_{\#_\sigma}(1_B \otimes h') \\ &= \sigma(h_1 \otimes h'_1) \otimes (d_H h_2)h'_2 + \sigma(h_1 \otimes h'_1) \otimes h_2 d_H h'_2 \\ &= \sigma(h_1 \otimes h'_1) \otimes d_H(h_2 h'_2),\end{aligned}$$

so that $d_B(\sigma(h_1 \otimes h'_1)) \otimes h_2 h'_2 = 0$ and then $d_B(\sigma(h \otimes h')) = 0$, for every $h, h' \in H$. \square

3.2 A classification of the smash product calculus

In this subsection we consider a right H -comodule algebra A such that $B := A^{\text{co}H} \subseteq A$ is a trivial extension. We denote the algebra inclusion by $\iota : B \rightarrow A$ and the cleaving map by $j : H \rightarrow A$. In the following, we characterize the calculi on $B\#H$ which are isomorphic to the smash product calculus $(\Omega^1(B\#H), d_{\#})$ via the differential of the algebra isomorphism $A \cong (B\#H)$. Recall that the calculus $(\Omega^1(B\#H), d_{\#})$ is obtained from Theorem 3.7 if one considers the trivial 2-cocycle.

Lemma 3.9. *The algebra morphisms $\iota_1 : B \rightarrow B\#H$, $b \mapsto b \otimes 1_H$ and $\iota_2 : H \rightarrow B\#H$, $h \mapsto 1_B \otimes h$ are differentiable.*

Proof. Clearly ι_1 and ι_2 are morphisms of algebras. We consider the morphisms $\widehat{\iota}_1 : \Omega^1(B) \rightarrow \Omega^1(B\#H)$, $\beta \mapsto \beta \otimes 1_H$ and $\widehat{\iota}_2 : \Omega^1(H) \rightarrow \Omega^1(B\#H)$, $\gamma \mapsto 1_B \otimes \gamma$. Now $\widehat{\iota}_1(b\beta b') = b\beta b' \otimes 1_H = (b \otimes 1_H) \cdot (\beta \otimes 1_H) \cdot (b' \otimes 1_H) = b\widehat{\iota}_1(\beta)b'$, so that $\widehat{\iota}_1$ is a morphism of B -bimodules and $\widehat{\iota}_2(h\gamma h') = 1_B \otimes h\gamma h' = (1_B \otimes h) \cdot (1_B \otimes \gamma) \cdot (1_B \otimes h') = h\widehat{\iota}_2(\gamma)h'$, so that $\widehat{\iota}_2$ is a morphism of H -bimodules. Furthermore, $\widehat{\iota}_1 d_B b = d_B b \otimes 1_H = d_{\#}\iota_1 b$ and $\widehat{\iota}_2 d_H h = 1_B \otimes d_H h = d_{\#}\iota_2 h$. \square

Note that $\widehat{\iota}_1$ and $\widehat{\iota}_2$ are injective. Observe also that, in case of a pure crossed product $B\#_\sigma H$ (i.e., with σ not trivial), ι_2 is not a morphism of algebras in general.

Remark 3.10. *Consider a FODC $(\Omega^1(A), d_A)$ on A and the corresponding pullback calculus $(\Omega^1(B), d_B)$ on B . In particular, the inclusion $\iota : B \rightarrow A$ is differentiable. By Theorem 2.9 the smash product algebra $B\#H$ is isomorphic to A as right H -comodule algebras through $\theta : A \rightarrow B\#H$. In order to build the right H -covariant FODC on $B\#H$ given by Theorem 3.7 we need that $(\Omega^1(B), d_B)$ is an H -module FODC on B . If $(\Omega^1(B), d_B)$ is an H -module FODC on B , the corresponding left action $\cdot : H \otimes \Omega^1(B) \rightarrow \Omega^1(B)$ satisfies*

$$\begin{aligned}h \cdot (bd_B b') &= (h_1 \cdot b)(h_2 \cdot d_B b') = (h_1 \cdot b)d_B(h_2 \cdot b') = j(h_1)bj^{-1}(h_2)d_B(j(h_3)b'j^{-1}(h_4)) \\ &= j(h_1)bb'd_A(j^{-1}(h_2)) + j(h_1)bd_B(b')j^{-1}(h_2) + j(h_1)bj^{-1}(h_2)d_A(j(h_3))b'j^{-1}(h_4) \\ &= j(h_1)bd_B(b')j^{-1}(h_2) + j(h_1)b(b'd_A(j^{-1}(h_2)) + j^{-1}(h_2)d_A(j(h_3))b'j^{-1}(h_4))\end{aligned}$$

for all $h \in H$, $b, b' \in B$, where we recall that $d_B := d_A|_B : B \rightarrow \Omega^1(B) \subseteq \Omega^1(A)$.

We are now prepared to give a classification of the smash product calculus in terms of the embedding $\iota: B \rightarrow A$, the cleaving map $j: H \rightarrow A$ and a "commutativity condition" on the differential d_A .

Theorem 3.11. *Let $B \subseteq A$ be a trivial Hopf-Galois extension with cleaving map $j: H \rightarrow A$. Consider an H -module FODC $(\Omega^1(B), d_B)$ on B and a bicovariant FODC $(\Omega^1(H), d_H)$ on H . Then, a FODC $(\Omega^1(A), d_A)$ on A with $\Omega^1(A)$ torsion-free as left and right A -module such that $(\Omega^1(B), d_B)$ is the corresponding pullback calculus on B is isomorphic to $(\Omega^1(B\#H), d_{\#})$ through θ if and only if:*

- (1) j is differentiable with \widehat{j} injective.
- (2) $\widehat{\iota}(\Omega^1(B))j(H) \cap \iota(B)\widehat{j}(\Omega^1(H)) = \{0\}$.
- (3) $d_A(j(h_1))bj^{-1}(h_2) = -j(h_1)bd_A(j^{-1}(h_2))$ for all $h \in H$, $b \in B$.

If these conditions are satisfied then $(\Omega^1(A), d_A)$ is a right H -covariant FODC on A .

Proof. Assume that $(\Omega^1(A), d_A) \cong (\Omega^1(B\#H), d_{\#})$ as FODCi through $\theta: A \xrightarrow{\cong} B\#H$. Then θ and θ^{-1} are differentiable with $\widehat{\theta^{-1}} = \widehat{\theta}^{-1}: \Omega^1(B\#H) \rightarrow \Omega^1(A)$ by Remark 2.16. Furthermore, we know that ι_2 is differentiable by Lemma 3.9. Then, since $j = \theta^{-1} \circ \iota_2$, we obtain that j is differentiable with $\widehat{j} = \widehat{\theta^{-1}} \circ \widehat{\iota_2}$. In particular, the diagram

$$\begin{array}{ccccccc} \Omega^1(H) & \xrightarrow{\widehat{\iota_2}} & \Omega^1(B\#H) & \xrightarrow{\widehat{\theta^{-1}}} & \Omega^1(A) & \xleftarrow{\widehat{\iota}} & \Omega^1(B) \\ d_H \uparrow & & \uparrow d_{\#} & & d_A \uparrow & & \uparrow d_B \\ H & \xrightarrow{\iota_2} & B\#H & \xrightarrow{\theta^{-1}} & A & \xleftarrow{\iota} & B \end{array}$$

commutes. Moreover, $\widehat{\iota_2}$ and $\widehat{\theta^{-1}}$ are injective, so \widehat{j} is injective and thus (1) holds.

We give the explicit expression of $\widehat{\theta^{-1}}$ which is uniquely determined by θ^{-1} .

$$\begin{aligned} \widehat{\theta^{-1}}(bd_B b' \otimes h) &= \widehat{\theta^{-1}}((b \otimes 1_H)d_{\#}(b' \otimes h) - (bb' \otimes 1_H)d_{\#}(1_B \otimes h)) \\ &= \theta^{-1}(b \otimes 1_H)d_A(\theta^{-1}(b' \otimes h)) - \theta^{-1}(bb' \otimes 1_H)d_A(\theta^{-1}(1_B \otimes h)) \\ &= bd_A(b'j(h)) - bb'd_A(j(h)) = bb'd_A(j(h)) + bd_A(b')j(h) - bb'd_A(j(h)) \\ &= bd_A(b')j(h) \end{aligned}$$

and so $\widehat{\theta^{-1}}(\Omega^1(B) \otimes H) = \widehat{\iota}(\Omega^1(B))j(H)$, while

$$\widehat{\theta^{-1}}(b \otimes hd_H h') = \widehat{\theta^{-1}}((b \otimes h)d_{\#}(1_B \otimes h')) = \theta^{-1}(b \otimes h)d_A(\theta^{-1}(1_B \otimes h')) = bj(h)d_A(j(h')) = \widehat{j}(hd_H h')$$

and then $\widehat{\theta^{-1}}(B \otimes \Omega^1(H)) = \iota(B)\widehat{j}(\Omega^1(H))$. Thus, we obtain

$$\begin{aligned} \widehat{\theta}(\widehat{\iota}(\Omega^1(B))j(H) \cap \iota(B)\widehat{j}(\Omega^1(H))) &\subseteq \widehat{\theta}(\widehat{\theta^{-1}}(\Omega^1(B) \otimes H)) \cap \widehat{\theta}(\widehat{\theta^{-1}}(B \otimes \Omega^1(H))) \\ &= (\Omega^1(B) \otimes H) \cap (B \otimes \Omega^1(H)) \\ &= \{0\} \end{aligned}$$

from which $\widehat{\iota}(\Omega^1(B))j(H) \cap \iota(B)\widehat{j}(\Omega^1(H)) = \{0\}$ follows, since $\widehat{\theta}$ is injective. Thus, (2) holds.

Furthermore, recalling that $j^{-1}(h)_0 \otimes j^{-1}(h)_1 = j^{-1}(h_2) \otimes S(h_1)$ and that $j^{-1}(hh') = j^{-1}(h')j^{-1}(h)$ for all $h, h' \in H$, we obtain

$$\begin{aligned} \widehat{\theta}(d_A(j(h_1))bj^{-1}(h_2)) &= d_{\#}(\theta(j(h_1)))\theta(bj^{-1}(h_2)) \\ &= d_{\#}(j(h_1)_0 j^{-1}(j(h_1)_1) \otimes j(h_1)_2)(b_0 j^{-1}(h_2)_0 j^{-1}(b_1 j^{-1}(h_2)_1) \otimes b_2 j^{-1}(h_2)_2) \\ &= d_{\#}(1_A \otimes h_1)(bj^{-1}(h_4)j^{-1}(S(h_3)) \otimes S(h_2)) = (1_A \otimes d_H h_1)(bj^{-1}(S(h_3)h_4) \otimes S(h_2)) \\ &= (1_A \otimes d_H h_1)(b \otimes S(h_2)) = ((d_H h_1)_{-1} \cdot b) \otimes (d_H h_1)_0 S(h_2) = (h_1 \cdot b) \otimes d_H(h_2)S(h_3) \\ &= -(h_1 \cdot b) \otimes h_2 d_H(S(h_3)) = -((h_1 \cdot b) \otimes h_2)(1_A \otimes d_H(S(h_3))) \\ &= -(j(h_1)bj^{-1}(h_2) \otimes h_3)d_{\#}(1_A \otimes S(h_4)) = -\theta(j(h_1)b)d_{\#}(\theta(j^{-1}(h_2))) \\ &= -\widehat{\theta}(j(h_1)bd_A(j^{-1}(h_2))) \end{aligned}$$

and from this, since $\widehat{\theta}$ is injective, we obtain that (3) holds true.

On the other hand, suppose that (1), (2) and (3) are satisfied. First note that, since (3) holds, by Remark 3.10 we have $h \cdot (bd_B b') = j(h_1)bd_B(b')j^{-1}(h_2)$ for all $h \in H$ and $b, b' \in B$. Thus, we obtain

$$h \cdot \beta = j(h_1)\beta j^{-1}(h_2), \quad \text{for all } h \in H \text{ and } \beta \in \Omega^1(B). \quad (19)$$

Next, we show that θ^{-1} is differentiable, i.e., that there exists a $B\#H$ -bimodule morphism $\widehat{\theta^{-1}} : \Omega^1(B\#H) \rightarrow \Omega^1(A)$ such that $\widehat{\theta^{-1}} \circ d_{\#} = d_A \circ \theta^{-1}$. We define

$$\widehat{\theta^{-1}}(\beta \otimes h + b \otimes \gamma) := \widehat{\iota}(\beta)j(h) + b\widehat{j}(\gamma), \quad \text{for } \beta \otimes h + b \otimes \gamma \in \Omega^1(B\#H).$$

Then

$$\widehat{\theta^{-1}}d_{\#}(b \otimes h) = \widehat{\theta^{-1}}(d_B b \otimes h + b \otimes d_H h) = \widehat{\iota}(d_B b)j(h) + b\widehat{j}(d_H h) = d_A(b)j(h) + b d_A(j(h)) = d_A(bj(h)) = d_A \theta^{-1}(b \otimes h),$$

since j is differentiable. Furthermore, $\widehat{\theta^{-1}}$ is a morphism of $B\#H$ -bimodules. Indeed, we can compute

$$\begin{aligned} \widehat{\theta^{-1}}((b' \otimes h') \cdot (\beta \otimes h + b \otimes \gamma)) &= \widehat{\theta^{-1}}(b'(h'_1 \cdot \beta) \otimes h'_2 h + b'(h'_1 \cdot b) \otimes h'_2 \gamma) = b'(h'_1 \cdot \beta)j(h'_2 h) + b'(h'_1 \cdot b)\widehat{j}(h'_2 \gamma) \\ &\stackrel{(19)}{=} b'j(h'_1)\beta j^{-1}(h'_2)j(h'_3)j(h) + b'j(h'_1)b j^{-1}(h'_2)\widehat{j}(h'_3 \gamma) \\ &= b'j(h')\beta j(h) + b'j(h'_1)b j^{-1}(h'_2)j(h'_3)\widehat{j}(\gamma) = b'j(h')\beta j(h) + b'j(h')\widehat{j}(\gamma) \\ &= b'j(h')(\beta j(h) + b\widehat{j}(\gamma)) = \theta^{-1}(b' \otimes h')(\widehat{\iota}(\beta)j(h) + b\widehat{j}(\gamma)) \\ &= (b' \otimes h') \cdot \widehat{\theta^{-1}}(\beta \otimes h + b \otimes \gamma), \end{aligned}$$

as well as

$$\begin{aligned} \widehat{\theta^{-1}}((\beta \otimes h) \cdot (b' \otimes h')) &= \widehat{\theta^{-1}}(\beta(h_1 \cdot b') \otimes h_2 h') = \beta(h_1 \cdot b')j(h_2 h') = \beta j(h_1)b'j^{-1}(h_2)j(h_3)j(h') = \beta j(h)b'j(h') \\ &= \widehat{\theta^{-1}}(\beta \otimes h) \cdot (b' \otimes h') \end{aligned}$$

and

$$\begin{aligned} \widehat{\theta^{-1}}((b \otimes hd_H h') \cdot (b' \otimes h'')) &= \widehat{\theta^{-1}}(b((hd_H h')_{-1} \cdot b') \otimes (hd_H h')_0 h'') = \widehat{\theta^{-1}}(b(h_1 h'_1 \cdot b') \otimes h_2 d_H(h'_2) h'') \\ &= b(h_1 h'_1 \cdot b')\widehat{j}(h_2 d_H(h'_2) h'') = b j(h_1 h'_1) b' j^{-1}(h_2 h'_2) j(h_3) d_A(j(h'_3)) j(h'') \\ &= b j(h) j(h'_1) b' j^{-1}(h'_2) d_A(j(h'_3)) j(h'') = -b j(h) j(h'_1) b' d_A(j^{-1}(h'_2)) j(h'_3) j(h'') \\ &\stackrel{(3)}{=} b j(h) d_A(j(h'_1)) b' j^{-1}(h'_2) j(h'_3) j(h'') = b j(h) d_A(j(h')) b' j(h'') = b\widehat{j}(hd_H h') \theta^{-1}(b' \otimes h'') \\ &= \widehat{\theta^{-1}}(b \otimes hd_H h') \cdot (b' \otimes h''). \end{aligned}$$

Hence, we have that θ^{-1} is differentiable with $\widehat{\theta^{-1}}$ given by $\mu_{\Omega^1(A)}^r \circ (\widehat{\iota} \otimes j) + \mu_{\Omega^1(A)}^l \circ (\iota \otimes \widehat{j})$. Now we show that $\widehat{\theta^{-1}}$ is bijective in order to obtain that it is an isomorphism of $B\#H$ -bimodules. The right and left A -module actions $\mu_{\Omega^1(A)}^r$ and $\mu_{\Omega^1(A)}^l$ on $\Omega^1(A)$ are injective since $\Omega^1(A)$ is a right and left torsion-free A -module. Moreover, ι , $\widehat{\iota}$, j and \widehat{j} are injective, and thus $\mu_{\Omega^1(A)}^r \circ (\widehat{\iota} \otimes j)$ and $\mu_{\Omega^1(A)}^l \circ (\iota \otimes \widehat{j})$ are injective. It follows that $\widehat{\theta^{-1}}$ is injective, since $\widehat{\iota}(\Omega^1(B))j(H) \cap \iota(B)\widehat{j}(\Omega^1(H)) = \{0\}$. Therefore, it only remains to prove the surjectivity of $\widehat{\theta^{-1}}$. Let $ad_A a' \in \Omega^1(A)$ be arbitrary and consider $(a_0 j^{-1}(a_1) \otimes a_2) d_{\#}(a'_0 j^{-1}(a'_1) \otimes a'_2) \in \Omega^1(B\#H)$. Thus, we have that

$$\begin{aligned} (a_0 j^{-1}(a_1) \otimes a_2) (d_B(a'_0 j^{-1}(a'_1)) \otimes a'_2 + a'_0 j^{-1}(a'_1) \otimes d_H a'_2) \\ = a_0 j^{-1}(a_1) (a_2 \cdot d_B(a'_0 j^{-1}(a'_1))) \otimes a_3 a'_2 + a_0 j^{-1}(a_1) (a_2 \cdot a'_0 j^{-1}(a'_1)) \otimes a_3 d_H a'_2 \\ \stackrel{(19)}{=} a_0 j^{-1}(a_1) j(a_2) d_B(a'_0 j^{-1}(a'_1)) j^{-1}(a_3) \otimes a_4 a'_2 + a_0 j^{-1}(a_1) j(a_2) a'_0 j^{-1}(a'_1) j^{-1}(a_3) \otimes a_4 d_H a'_2 \\ = a_0 d_B(a'_0 j^{-1}(a'_1)) j^{-1}(a_1) \otimes a_2 a'_2 + a_0 a'_0 j^{-1}(a'_1) j^{-1}(a_1) \otimes a_2 d_H a'_2, \end{aligned}$$

which implies

$$\begin{aligned} \widehat{\theta^{-1}}((a_0 j^{-1}(a_1) \otimes a_2) d_{\#}(a'_0 j^{-1}(a'_1) \otimes a'_2)) &= \widehat{\theta^{-1}}(a_0 d_B(a'_0 j^{-1}(a'_1)) j^{-1}(a_1) \otimes a_2 a'_2 + a_0 a'_0 j^{-1}(a'_1) j^{-1}(a_1) \otimes a_2 d_H a'_2) \\ &= a_0 d_A(a'_0 j^{-1}(a'_1)) j^{-1}(a_1) j(a_2) j(a'_2) + a_0 a'_0 j^{-1}(a'_1) j^{-1}(a_1) j(a_2) d_A(j(a'_2)) \\ &= a d_A(a'_0 j^{-1}(a'_1)) j(a'_2) + a a'_0 j^{-1}(a'_1) d_A(j(a'_2)) \\ &= a (d_A(a'_0 j^{-1}(a'_1)) j(a'_2) + a'_0 j^{-1}(a'_1) d_A(j(a'_2))) \\ &= a d_A(a'_0 j^{-1}(a'_1) j(a'_2)) = a d_A(a'_0 \varepsilon(a'_1)) \\ &= a d_A a'. \end{aligned}$$

We have shown that $\widehat{\theta^{-1}}$ is also surjective and consequently an isomorphism of $B\#H$ -bimodules. Hence, $(\Omega^1(A), d_A) \cong (\Omega^1(B\#H), d_{\#})$ and the converse implication follows.

If $(\Omega^1(A), d_A) \cong (\Omega^1(B\#H), d_{\#})$ are isomorphic as FODCi through θ we define $\rho : \Omega^1(A) \rightarrow \Omega^1(A) \otimes H$ by $\rho := (\widehat{\theta^{-1}} \otimes \text{Id}_H) \circ \rho' \circ \widehat{\theta}$, where ρ' is the right H -coaction of $\Omega^1(B\#H)$. If we show that ρ is a right H -coaction, then $\widehat{\theta}$ is automatically right H -colinear. Using that $\widehat{\theta}$ is left A -linear and that the left $B\#H$ -action of $\Omega^1(B\#H)$, θ and $d_{\#}$ are right H -colinear we obtain

$$\begin{aligned} (\widehat{\theta^{-1}} \otimes \text{Id}_H) \rho' \widehat{\theta}(ad_A a') &= (\widehat{\theta^{-1}} \otimes \text{Id}_H) \rho'(\theta(a) d_{\#}(\theta(a'))) = (\widehat{\theta^{-1}} \otimes \text{Id}_H)(\theta(a)_0 d_{\#}(\theta(a'))_0 \otimes \theta(a)_1 d_{\#}(\theta(a'))_1) \\ &= (\widehat{\theta^{-1}} \otimes \text{Id}_H)(\theta(a_0) d_{\#}(\theta(a')_0) \otimes a_1 \theta(a')_1) = (\widehat{\theta^{-1}} \otimes \text{Id}_H)(\theta(a_0) d_{\#}(\theta(a'_0)) \otimes a_1 a'_1) \\ &= \theta^{-1}(\theta(a_0)) d_A(\theta^{-1}(\theta(a'_0))) \otimes a_1 a'_1 = a_0 d_A a'_0 \otimes a_1 a'_1. \end{aligned}$$

This is clearly a right H -coaction, since

$$(\rho \otimes \text{Id}_H) \rho(ad_A a') = a_0 d_A a'_0 \otimes a_1 a'_1 \otimes a_2 a'_2 = (\text{Id}_{\Omega^1(A)} \otimes \Delta) \rho(ad_A a') \text{ and } (a_0 d_A a'_0) \varepsilon(a_1 a'_1) = ad_A a'.$$

Then $\Omega^1(A)$ is in ${}_A \mathfrak{M}_A^H$ since A is a right H -comodule algebra and d_A is right H -colinear. \square

3.3 Higher order forms

In this subsection H denotes a Hopf algebra and B a σ -twisted left H -module algebra with measure $\cdot : H \otimes B \rightarrow B$ and 2-cocycle $\sigma : H \otimes H \rightarrow B$. In the following we introduce higher order forms on $B\#_{\sigma}H$. By generalizing Definition 3.3 we give the following:

Definition 3.12. Let $(\Omega^{\bullet}(B), d_B)$ be a DC on B . We say that it is a σ -twisted H -module DC if there exist linear maps $\cdot : H \otimes \Omega^n(B) \rightarrow \Omega^n(B)$ for all $n \geq 1$, such that

$$\begin{aligned} h \cdot (b^0 d_B b^1 \wedge \cdots \wedge d_B b^n) &= (h_1 \cdot b^0)(h_2 \cdot d_B b^1) \wedge \cdots \wedge (h_{n+1} \cdot d_B b^n), \\ h \cdot d_B b &= d_B(h \cdot b), \\ d_B \circ \sigma &= 0 \end{aligned}$$

for all $b, b^0, \dots, b^n \in B$ and $h \in H$.

Let us discuss some immediate consequences of the above definition.

Remark 3.13. From the first two equations in Definition 3.12 we obtain that, if they exist, the linear maps $\cdot : H \otimes \Omega^n(B) \rightarrow \Omega^n(B)$ are uniquely determined by

$$h \cdot (b^0 d_B b^1 \wedge \cdots \wedge d_B b^n) = (h_1 \cdot b^0) d_B(h_2 \cdot b^1) \wedge \cdots \wedge d_B(h_{n+1} \cdot b^n).$$

Clearly $1_H \cdot \beta = \beta$ for all $\beta \in \Omega^{\bullet}(B)$ and, as in the proof of Lemma 3.2, one shows that

$$h \cdot (h' \cdot \beta) = \sigma(h_1 \otimes h'_1)(h_2 h'_2 \cdot \beta) \sigma^{-1}(h_3 \otimes h'_3) \text{ for all } \beta \in \Omega^{\bullet}(B) \text{ and } h, h' \in H. \quad (20)$$

In addition,

$$h \cdot (\beta \wedge \beta') = (h_1 \cdot \beta) \wedge (h_2 \cdot \beta') \text{ for all } h \in H, \beta \in \Omega^n(B) \text{ and } \beta' \in \Omega^m(B) \quad (21)$$

holds true for arbitrary $n, m \geq 0$.

We now define higher order forms on $B\#_{\sigma}H$ by generalizing Theorem 3.7.

Theorem 3.14. Let $(\Omega^{\bullet}(B), d_B)$ be a σ -twisted H -module DC on B and $(\Omega^{\bullet}(H), d_H)$ a bicovariant DC on H . Let us define

$$\Omega^n(B\#_{\sigma}H) := \bigoplus_{i=0}^n \Omega^{n-i}(B) \otimes \Omega^i(H) \quad \text{for all } n \geq 0$$

and set

$$\begin{aligned} (\omega \otimes \gamma) \wedge (\omega' \otimes \gamma') &:= (-1)^{jk} (\omega \wedge (\gamma_{-2} \cdot \omega') \sigma(\gamma_{-1} \otimes \gamma'_{-1})) \otimes (\gamma_0 \wedge \gamma'_0), \\ d_{\#_{\sigma}}(\omega \otimes \gamma) &:= d_B \omega \otimes \gamma + (-1)^i \omega \otimes d_H \gamma \end{aligned}$$

for $\omega \in \Omega^i(B)$, $\gamma \in \Omega^j(H)$ and $\omega' \in \Omega^k(B)$. Then $(\Omega^{\bullet}(B\#_{\sigma}H), d_{\#_{\sigma}})$ is a right H -covariant DC on $B\#_{\sigma}H$ with respect to which the right H -coaction $\rho : B\#_{\sigma}H \rightarrow (B\#_{\sigma}H) \otimes H$ is differentiable.

Proof. First, note that \wedge is a map of degree 0, since this is the case for the products of $\Omega^\bullet(B)$ and $\Omega^\bullet(H)$, $\lambda(\Omega^n(H)) \subseteq H \otimes \Omega^n(H)$ and $H \cdot \Omega^n(B) \subseteq \Omega^n(B)$ for all $n \geq 0$. Now, we show the associativity of the product. Given $\omega \in \Omega^i(B)$, $\omega' \in \Omega^{i'}(B)$, $\omega'' \in \Omega^{i''}(B)$, $\gamma \in \Omega^j(H)$, $\gamma' \in \Omega^{j'}(H)$ and $\gamma'' \in \Omega^{j''}(H)$, we obtain

$$\begin{aligned}
& ((\omega \otimes \gamma) \wedge (\omega' \otimes \gamma')) \wedge (\omega'' \otimes \gamma'') = (-1)^{ji'} (\omega \wedge (\gamma_{-2} \cdot \omega') \sigma(\gamma_{-1} \otimes \gamma'_{-1}) \otimes \gamma_0 \wedge \gamma'_0) \wedge (\omega'' \otimes \gamma'') \\
& = (-1)^{j(i'+j)+i''} \omega \wedge (\gamma_{-2} \cdot \omega') \sigma(\gamma_{-1} \otimes \gamma'_{-1}) \wedge ((\gamma_0 \wedge \gamma'_0)_{-2} \cdot \omega'') \sigma((\gamma_0 \wedge \gamma'_0)_{-1} \otimes \gamma''_{-1}) \otimes (\gamma_0 \wedge \gamma'_0)_0 \wedge \gamma''_0 \\
& = (-1)^{j(i'+i'')+j''} \omega \wedge (\gamma_{-4} \cdot \omega') \sigma(\gamma_{-3} \otimes \gamma'_{-3}) \wedge (\gamma_{-2} \gamma'_{-2} \cdot \omega'') \sigma(\gamma_{-1} \gamma'_{-1} \otimes \gamma''_{-1}) \otimes \gamma_0 \wedge \gamma'_0 \wedge \gamma''_0 \\
& = (-1)^{j(i'+i'')+j''} \omega \wedge (\gamma_{-6} \cdot \omega') \wedge \sigma(\gamma_{-5} \otimes \gamma'_{-5}) (\gamma_{-4} \gamma'_{-4} \cdot \omega'') \sigma^{-1}(\gamma_{-3} \otimes \gamma'_{-3}) \sigma(\gamma_{-2} \otimes \gamma'_{-2}) \sigma(\gamma_{-1} \gamma'_{-1} \otimes \gamma''_{-1}) \\
& \quad \otimes \gamma_0 \wedge \gamma'_0 \wedge \gamma''_0 \\
& \stackrel{(20)}{=} (-1)^{j(i'+i'')+j''} \omega \wedge (\gamma_{-4} \cdot \omega') \wedge (\gamma_{-3} \cdot (\gamma'_{-3} \cdot \omega'')) (\gamma_{-2} \cdot \sigma(\gamma'_{-2} \otimes \gamma''_{-2})) \sigma(\gamma_{-1} \otimes \gamma'_{-1} \gamma''_{-1}) \otimes \gamma_0 \wedge \gamma'_0 \wedge \gamma''_0 \\
& \stackrel{(21)}{=} (-1)^{j(i'+i'')+j''} \omega \wedge (\gamma_{-2} \cdot (\omega' \wedge (\gamma'_{-2} \cdot \omega'')) \sigma(\gamma'_{-1} \otimes \gamma''_{-1})) \sigma(\gamma_{-1} \otimes (\gamma'_0 \wedge \gamma''_0)_{-1}) \otimes \gamma_0 \wedge (\gamma'_0 \wedge \gamma''_0)_0 \\
& = (\omega \otimes \gamma) \wedge ((-1)^{j''} \omega' \wedge (\gamma'_{-2} \cdot \omega'') \sigma(\gamma'_{-1} \otimes \gamma''_{-1}) \otimes \gamma'_0 \wedge \gamma''_0) \\
& = (\omega \otimes \gamma) \wedge ((\omega' \otimes \gamma') \wedge (\omega'' \otimes \gamma'')).
\end{aligned}$$

Furthermore, $d_{\#_\sigma}$ satisfies the Leibniz rule, indeed

$$\begin{aligned}
& d_{\#_\sigma}(\omega \otimes \gamma) \wedge (\omega' \otimes \gamma') + (-1)^{i+j} (\omega \otimes \gamma) \wedge d_{\#_\sigma}(\omega' \otimes \gamma') \\
& = (d_B \omega \otimes \gamma) \wedge (\omega' \otimes \gamma') + (-1)^i (\omega \otimes d_H \gamma) \wedge (\omega' \otimes \gamma') + (-1)^{i+j} (\omega \otimes \gamma) \wedge (d_B \omega' \otimes \gamma') + (-1)^{i+j+i'} (\omega \otimes \gamma) \wedge (\omega' \otimes d_H \gamma') \\
& = (-1)^{ji'} d_B \omega \wedge (\gamma_{-2} \cdot \omega') \sigma(\gamma_{-1} \otimes \gamma'_{-1}) \otimes \gamma_0 \wedge \gamma'_0 + (-1)^{i+(j+i')i''} \omega \wedge ((d_H \gamma)_{-2} \cdot \omega') \sigma((d_H \gamma)_{-1} \otimes \gamma'_{-1}) \otimes (d_H \gamma)_0 \wedge \gamma'_0 \\
& \quad + (-1)^{i+j+j(i'+1)} \omega \wedge (\gamma_{-2} \cdot d_B \omega') \sigma(\gamma_{-1} \otimes \gamma'_{-1}) \otimes \gamma_0 \wedge \gamma'_0 + (-1)^{i+j+i'+j''} \omega \wedge (\gamma_{-2} \cdot \omega') \sigma(\gamma_{-1} \otimes (d_H \gamma')_{-1}) \otimes \gamma_0 \wedge (d_H \gamma')_0 \\
& = (-1)^{ji'} d_B \omega \wedge (\gamma_{-2} \cdot \omega') \sigma(\gamma_{-1} \otimes \gamma'_{-1}) \otimes \gamma_0 \wedge \gamma'_0 + (-1)^{j''+i+i'} (\omega \wedge (\gamma_{-2} \cdot \omega') \sigma(\gamma_{-1} \otimes \gamma'_{-1}) \otimes d_H \gamma_0 \wedge \gamma'_0) \\
& \quad + (-1)^{i+j''} \omega \wedge d_B((\gamma_{-2} \cdot \omega') \sigma(\gamma_{-1} \otimes \gamma'_{-1})) \otimes \gamma_0 \wedge \gamma'_0 + (-1)^{i+j+i'+j''} \omega \wedge (\gamma_{-2} \cdot \omega') \sigma(\gamma_{-1} \otimes \gamma'_{-1}) \otimes \gamma_0 \wedge d_H \gamma'_0 \\
& = (-1)^{ji'} (d_B \omega \wedge (\gamma_{-2} \cdot \omega') \sigma(\gamma_{-1} \otimes \gamma'_{-1}) \otimes \gamma_0 \wedge \gamma'_0 + (-1)^i \omega \wedge d_B((\gamma_{-2} \cdot \omega') \sigma(\gamma_{-1} \otimes \gamma'_{-1})) \otimes \gamma_0 \wedge \gamma'_0) \\
& \quad + (-1)^{j''+i+i'} (\omega \wedge (\gamma_{-2} \cdot \omega') \sigma(\gamma_{-1} \otimes \gamma'_{-1}) \otimes d_H \gamma_0 \wedge \gamma'_0 + (-1)^j \omega \wedge (\gamma_{-2} \cdot \omega') \sigma(\gamma_{-1} \otimes \gamma'_{-1}) \otimes \gamma_0 \wedge d_H \gamma'_0) \\
& = (-1)^{ji'} (d_B(\omega \wedge (\gamma_{-2} \cdot \omega') \sigma(\gamma_{-1} \otimes \gamma'_{-1})) \otimes \gamma_0 \wedge \gamma'_0) + (-1)^{j''+i+i'} (\omega \wedge (\gamma_{-2} \cdot \omega') \sigma(\gamma_{-1} \otimes \gamma'_{-1}) \otimes d_H(\gamma_0 \wedge \gamma'_0)) \\
& = (-1)^{ji'} (d_B(\omega \wedge (\gamma_{-2} \cdot \omega') \sigma(\gamma_{-1} \otimes \gamma'_{-1})) \otimes \gamma_0 \wedge \gamma'_0 + (-1)^{i+i'} \omega \wedge (\gamma_{-2} \cdot \omega') \sigma(\gamma_{-1} \otimes \gamma'_{-1}) \otimes d_H(\gamma_0 \wedge \gamma'_0)) \\
& = (-1)^{ji'} d_{\#_\sigma}(\omega \wedge (\gamma_{-2} \cdot \omega') \sigma(\gamma_{-1} \otimes \gamma'_{-1}) \otimes \gamma_0 \wedge \gamma'_0) = d_{\#_\sigma}((\omega \otimes \gamma) \wedge (\omega' \otimes \gamma'))
\end{aligned}$$

where in the third equality we use second and third equations of Definition 3.12, while the fact that $d_{\#_\sigma}^2 = 0$ follows in complete analogy to the tensor product DC of Example 2.13 iii). Hence $(\Omega^\bullet(B\#_\sigma H), \wedge, d_{\#_\sigma})$ is a differential graded algebra such that $\Omega^0(B\#_\sigma H) = B\#_\sigma H$. Moreover, from Theorem 3.7 we know that $\Omega^1(B\#_\sigma H)$ is spanned by elements of the form $(b \otimes h) d_{\#_\sigma}(b' \otimes h')$, for $b, b' \in B$ and $h, h' \in H$. Now, suppose that for a fixed $n > 1$ $\Omega^{n-1}(B\#_\sigma H)$ is spanned by elements of the form $(b^0 \otimes h^0) d_{\#_\sigma}(b^1 \otimes h^1) \wedge \dots \wedge d_{\#_\sigma}(b^{n-1} \otimes h^{n-1})$ for $b^i \in B$ and $h^i \in H$. A generic element in $\Omega^n(B\#_\sigma H)$ is a sum of elements in $\Omega^k(B) \otimes \Omega^l(H)$ with $k, l \in \{0, \dots, n\}$ such that $k+l = n$ and then, by using the "surjectivity condition" of $\Omega^k(B)$ and $\Omega^l(H)$, of the form $(b^0 d_B b^1 \wedge d_B b^2 \wedge \dots \wedge d_B b^k) \otimes (h^0 d_H h^1 \wedge \dots \wedge d_H h^l)$. Then, we have that

$$\begin{aligned}
& (b^0 d_B b^1 \otimes 1_H) \wedge ((d_B b^2 \wedge \dots \wedge d_B b^k) \otimes (h^0 d_H h^1 \wedge \dots \wedge d_H h^l)) \\
& = b^0 d_B b^1 \wedge (1_H \cdot d_B b^2 \wedge \dots \wedge d_B b^k) \sigma(1_H \otimes (h^0 d_H h^1 \wedge \dots \wedge d_H h^l)_{-1}) \otimes 1_H (h^0 d_H h^1 \wedge \dots \wedge d_H h^l)_0 \\
& = (b^0 d_B b^1 \wedge d_B b^2 \wedge \dots \wedge d_B b^k) \otimes (h^0 d_H h^1 \wedge \dots \wedge d_H h^l)
\end{aligned}$$

and $(d_B b^2 \wedge \dots \wedge d_B b^k) \otimes (h^0 d_H h^1 \wedge \dots \wedge d_H h^l)$ is in $\Omega^{n-1}(B\#_\sigma H)$, which is by hypothesis a finite sum of elements of the aforementioned form. Since $b^0 d_B b^1 \otimes 1_H = (b^0 \otimes 1_H) d_{\#_\sigma}(b^1 \otimes 1_H)$, we can conclude by induction that $\Omega^n(B\#_\sigma H)$ is spanned by elements of the form $(b^0 \otimes h^0) d_{\#_\sigma}(b^1 \otimes h^1) \wedge \dots \wedge d_{\#_\sigma}(b^n \otimes h^n)$ with $b^i \in B$ and $h^i \in H$. Thus, we have obtained that $(\Omega^\bullet(B\#_\sigma H), d_{\#_\sigma})$ is a DC on $B\#_\sigma H$ and it is right H -covariant since the DC $(\Omega^\bullet(H), d_H)$ is right covariant.

Finally, we show that the algebra map $\rho : B\#_\sigma H \rightarrow (B\#_\sigma H) \otimes H$, $b \otimes h \mapsto b \otimes h_1 \otimes h_2$ is differentiable. Recalling Remark 2.14, we define $\Omega^\bullet(\rho) : \Omega^\bullet(B\#_\sigma H) \rightarrow \Omega^\bullet((B\#_\sigma H) \otimes H)$ as $\Omega^\bullet(\rho)(\omega \otimes \gamma) = \omega \otimes \gamma_{[1]} \otimes \gamma_{[2]}$

such that $\Omega^0(\rho) = \rho$. Then $\Omega^\bullet(\rho)$ is an algebra map: indeed, given $\omega, \omega', \gamma, \gamma'$ as before, we obtain

$$\begin{aligned}
\Omega^\bullet(\rho)((\omega \otimes \gamma) \wedge (\omega' \otimes \gamma')) &= (-1)^{j i'} \Omega^\bullet(\rho)(\omega \wedge (\gamma_{-2} \cdot \omega') \sigma(\gamma_{-1} \otimes \gamma'_{-1}) \otimes \gamma_0 \wedge \gamma'_0) \\
&= (-1)^{j i'} \omega \wedge (\gamma_{-2} \cdot \omega') \sigma(\gamma_{-1} \otimes \gamma'_{-1}) \otimes (\gamma_0 \wedge \gamma'_0)_{[1]} \otimes (\gamma_0 \wedge \gamma'_0)_{[2]} \\
&\stackrel{(9)}{=} \sum_{\substack{k+l=j \\ k'+l'=j'}} (-1)^{j i' + l k'} \omega \wedge (\gamma_{-2} \cdot \omega') \sigma(\gamma_{-1} \otimes \gamma'_{-1}) \otimes (\gamma_{0, <1, k>} \wedge \gamma'_{0, <1, k'>}) \otimes (\gamma_{0, <2, l>} \wedge \gamma'_{0, <2, l'>}) \\
&= \sum_{\substack{k+l=j \\ k'+l'=j'}} (-1)^{k i' + l(i' + k')} \omega \wedge (\gamma_{<1, k>, -2} \cdot \omega') \sigma(\gamma_{<1, k>, -1} \otimes \gamma'_{<1, k'>, -1}) \otimes (\gamma_{<1, k>, 0} \wedge \gamma'_{<1, k'>, 0}) \otimes (\gamma_{<2, l>} \wedge \gamma'_{<2, l'>}) \\
&= \sum_{\substack{k+l=j \\ k'+l'=j'}} (-1)^{l(i' + k')} (\omega \otimes \gamma_{<1, k>}) \wedge (\omega' \otimes \gamma'_{<1, k'>}) \otimes (\gamma_{<2, l>} \wedge \gamma'_{<2, l'>}) \\
&= \sum_{\substack{k+l=j \\ k'+l'=j'}} (\omega \otimes \gamma_{<1, k>} \otimes \gamma_{<2, l>}) \tilde{\wedge} (\omega' \otimes \gamma'_{<1, k'>} \otimes \gamma'_{<2, l'>}) \\
&= \Omega^\bullet(\rho)(\omega \otimes \gamma) \tilde{\wedge} \Omega^\bullet(\rho)(\omega' \otimes \gamma'),
\end{aligned}$$

where we have used (10) for γ and γ' in the fourth equation and the product

$$\begin{aligned}
\tilde{\wedge}: (\Omega^\bullet(B \#_\sigma H) \otimes \Omega^\bullet(H))^{\otimes 2} &\rightarrow \Omega^\bullet(B \#_\sigma H) \otimes \Omega^\bullet(H) \\
(\vartheta \otimes \gamma) \otimes (\vartheta' \otimes \gamma') &\mapsto (-1)^{|\vartheta'| |\gamma|} (\vartheta \wedge \vartheta') \otimes (\gamma \wedge \gamma')
\end{aligned}$$

in the sixth equation. Furthermore, from $d_\otimes \circ \Omega^\bullet(\Delta) = \Omega^\bullet(\Delta) \circ d_H$, we obtain that

$$\begin{aligned}
\Omega^\bullet(\rho) d_{\#_\sigma}(\omega \otimes \gamma) &= \Omega^\bullet(\rho)(d_B \omega \otimes \gamma + (-1)^{|\omega|} \omega \otimes d_H \gamma) = d_B \omega \otimes \gamma_{[1]} \otimes \gamma_{[2]} + (-1)^{|\omega|} \omega \otimes (d_H \gamma)_{[1]} \otimes (d_H \gamma)_{[2]} \\
&= d_B \omega \otimes \gamma_{[1]} \otimes \gamma_{[2]} + (-1)^{|\omega|} \omega \otimes d_\otimes(\gamma_{[1]} \otimes \gamma_{[2]}) = d_\otimes(\omega \otimes \gamma_{[1]} \otimes \gamma_{[2]}) \\
&= d_\otimes \Omega^\bullet(\rho)(\omega \otimes \gamma),
\end{aligned}$$

hence $\Omega^\bullet(\rho) \circ d_{\#_\sigma} = d_\otimes \circ \Omega^\bullet(\rho)$. Thus ρ is differentiable. \square

3.4 Quantum principal bundle and connections

Now we study some constructions related to the FODC on the crossed product algebra $B \#_\sigma H$. In this subsection we show that there is a (regular and strong) QPB on $B \#_\sigma H$ with respect to this calculus. Afterwards, we study bimodule covariant derivatives on associated bundles of the crossed product algebra. Furthermore, we compute the noncommutative de Rham cohomology of $B \#_\sigma H$ from the cohomologies of B and H via the Leray–Serre spectral sequence. The latter employs higher order vertical maps and Atiyah sequences.

We consider $B \#_\sigma H$ as right H -comodule algebra via $\rho := \text{Id}_B \otimes \Delta$ and the right H -covariant FODC $(\Omega^1(B \#_\sigma H), d_{\#_\sigma})$ on it given in Theorem 3.7, when $(\Omega^1(B), d_B)$ is a σ -twisted H -module FODC on B and $(\Omega^1(H), d_H)$ is a bicovariant FODC on H . Recalling that $(B \#_\sigma H)^{\text{co}H} = B \otimes \mathbb{k}1_H$, the induced pullback calculus on $B \otimes \mathbb{k}1_H$ is given by

$$\Omega^1(B \otimes \mathbb{k}1_H) = \text{span}_{\mathbb{k}}\{(b \otimes 1_H) d_{\#_\sigma}(b' \otimes 1_H) \mid b, b' \in B\} = \text{span}_{\mathbb{k}}\{b d_B b' \otimes 1_H \mid b, b' \in B\} = \Omega^1(B) \otimes \mathbb{k}1_H$$

with differential given by the restriction and corestriction of $d_{\#_\sigma}$. The horizontal forms are

$$\Omega^1(B \#_\sigma H)_{\text{hor}} := (B \#_\sigma H)(\Omega^1(B) \otimes \mathbb{k}1_H)(B \#_\sigma H) = \Omega^1(B) \otimes H.$$

The last equality holds, since clearly $\Omega^1(B \#_\sigma H)_{\text{hor}} \subseteq \Omega^1(B) \otimes H$ and, given $b d_B b' \otimes h$ for $b, b' \in B$ and $h \in H$

$$((b \otimes 1_H) \cdot (d_B b' \otimes 1_H)) \cdot (1_B \otimes h) = (b d_B b' \otimes 1_H) \cdot (1_B \otimes h) = b d_B b' \varepsilon(h_1) \otimes h_2 = b d_B b' \otimes h.$$

We prove that the horizontal forms fit into a short exact sequence, forming a QPB. This QPB is automatically regular, in the sense that it descends from a universal QPB with a strong connection.

Theorem 3.15. *Given $(\Omega^1(H), d_H)$ a bicovariant FODC on H and $(\Omega^1(B), d_B)$ a σ -twisted H -module FODC on B , there is a regular and strong QPB on the crossed product algebra $B \#_\sigma H$. Explicitly,*

$$0 \longrightarrow \Omega^1(B) \otimes H \xrightarrow{\iota_1} \Omega^1(B \#_\sigma H) \xrightarrow{\text{ver}} (B \#_\sigma H) \otimes {}^{\text{co}H} \Omega^1(H) \longrightarrow 0 \quad (22)$$

is an exact sequence in $B \#_\sigma H \mathfrak{M}$, where ι_1 is the inclusion and $(\Omega^1(B \#_\sigma H), d_{\#_\sigma})$ is the right H -covariant FODC on $B \#_\sigma H$, constructed in Theorem 3.7.

Proof. Since we have shown in Theorem 3.7 that the right H -coaction $\rho : B\#_\sigma H \rightarrow (B\#_\sigma H) \otimes H$ is differentiable we have that $\text{ver} = \phi \circ \pi'_2 \circ \widehat{\rho}$ is well-defined by Remark 2.21, where

$$\begin{aligned}\widehat{\rho} : \Omega^1(B\#_\sigma H) &\rightarrow \Omega^1((B\#_\sigma H) \otimes H), \\ \beta \otimes h + b \otimes \gamma &\mapsto \beta \otimes h_1 \otimes h_2 + b \otimes \gamma_0 \otimes \gamma_1 + b \otimes \gamma_{-1} \otimes \gamma_0,\end{aligned}$$

$\pi'_2 : \Omega^1((B\#_\sigma H) \otimes H) \rightarrow (B\#_\sigma H) \otimes \Omega^1(H)$ is the canonical projection and

$$\phi : (B\#_\sigma H) \square_H \Omega^1(H) \rightarrow (B\#_\sigma H) \otimes {}^{\text{co}H}\Omega^1(H), \quad b \otimes h \otimes \gamma \mapsto b \otimes h \otimes S(\gamma_{-1})\gamma_0$$

is an isomorphism. Thus, we obtain $\text{ver}(\beta \otimes h + b \otimes \gamma) = b \otimes \gamma_{-2} \otimes S(\gamma_{-1})\gamma_0$, which we can rewrite as $\text{ver} = p \circ \pi_2$, where $\pi_2 : \Omega^1(B\#_\sigma H) \rightarrow B \otimes \Omega^1(H)$ is the canonical projection and $p : B \otimes \Omega^1(H) \rightarrow (B\#_\sigma H) \otimes {}^{\text{co}H}\Omega^1(H)$ is the linear map defined by $p(b \otimes \gamma) := b \otimes \gamma_{-2} \otimes S(\gamma_{-1})\gamma_0$. Observe that p is left $B\#_\sigma H$ -linear, since we have

$$\begin{aligned}p((b' \otimes h') \cdot (b \otimes \gamma)) &= p(b'(h'_1 \cdot b)\sigma(h'_2 \otimes \gamma_{-1}) \otimes h'_3\gamma_0) \\ &= b'(h'_1 \cdot b)\sigma(h'_2 \otimes \gamma_{-3}) \otimes h'_3\gamma_{-2} \otimes S(h'_4\gamma_{-1})h'_5\gamma_0 \\ &= b'(h'_1 \cdot b)\sigma(h'_2 \otimes \gamma_{-3}) \otimes h'_3\gamma_{-2} \otimes S(\gamma_{-1})\gamma_0 \\ &= (b' \otimes h')(b \otimes \gamma_{-2}) \otimes S(\gamma_{-1})\gamma_0\end{aligned}$$

for all $b, b' \in B$, $h' \in H$ and $\gamma \in \Omega^1(H)$.

We show that p is bijective in order to obtain that ver is surjective and that $\ker(\text{ver}) = \ker(\pi_2) = \Omega^1(B) \otimes H$. We define the linear map $g : (B\#_\sigma H) \otimes {}^{\text{co}H}\Omega^1(H) \rightarrow B \otimes \Omega^1(H)$ by $g(b \otimes h \otimes \gamma) := b \otimes h\gamma$. The morphism g is the inverse of p . Indeed, given $b \otimes h \otimes \gamma \in (B\#_\sigma H) \otimes {}^{\text{co}H}\Omega^1(H)$, we have

$$\begin{aligned}pg(b \otimes h \otimes \gamma) &= p(b \otimes h\gamma) = b \otimes (h\gamma)_{-2} \otimes S((h\gamma)_{-1})(h\gamma)_0 = b \otimes h_1\gamma_{-2} \otimes S(h_2\gamma_{-1})h_3\gamma_0 \\ &= b \otimes h_1 \otimes S(h_2)h_3\gamma = b \otimes h \otimes \gamma\end{aligned}$$

since γ is in ${}^{\text{co}H}\Omega^1(H)$ and, given $b \otimes \gamma \in B \otimes \Omega^1(H)$, we obtain

$$gp(b \otimes \gamma) = g(b \otimes \gamma_{-2} \otimes S(\gamma_{-1})\gamma_0) = b \otimes \gamma_{-2}S(\gamma_{-1})\gamma_0 = b \otimes \varepsilon(\gamma_{-1})\gamma_0 = b \otimes \gamma.$$

Thus, p is bijective and the sequence (22) is exact in ${}_{B\#_\sigma H}\mathfrak{M}$, hence we have a QPB on the crossed product algebra $B\#_\sigma H$. Regularity of the QPB follows from the discussion in Example 2.24, since crossed product algebras correspond to cleft extensions. Finally, observe that $(\Omega^1(B) \otimes H)^{\text{co}H} = \Omega^1(B) \otimes H^{\text{co}H} = \Omega^1(B) \otimes \mathbb{k}1_H$ and then, using [3, Corollary 5.53], the QPB is strong (the bijectivity of the antipode of H is needed here). \square

Remark 3.16. Note that clearly $(\Omega^1(B) \otimes \mathbb{k}1_H)(B\#_\sigma H) = \Omega^1(B) \otimes H$ since every $\beta \otimes h \in \Omega^1(B) \otimes H$ can be written as $(\beta \otimes 1_H) \cdot (1_B \otimes h)$. Since the QPB of Theorem 3.15 is strong we also obtain $(B\#_\sigma H)(\Omega^1(B) \otimes \mathbb{k}1_H) = \Omega^1(B) \otimes H$.

We show that the above QPB exhibits a canonical strong connection. Applying Definition 2.22 and Remark 2.23 to the present framework, we recall that a connection is a section $c : (B\#_\sigma H) \otimes {}^{\text{co}H}\Omega^1(H) \rightarrow \Omega^1(B\#_\sigma H)$ in ${}_{B\#_\sigma H}\mathfrak{M}^H$ of the vertical map ver and it is a strong connection if $(\text{Id} - c \circ \text{ver})d_{\#_\sigma}(B\#_\sigma H) \subseteq \Omega^1(B)(B\#_\sigma H)$.

Proposition 3.17. The QPB on the crossed product algebra $B\#_\sigma H$ of Theorem 3.15 admits a strong connection

$$c : (B\#_\sigma H) \otimes {}^{\text{co}H}\Omega^1(H) \rightarrow \Omega^1(B\#_\sigma H), \quad (b \otimes h) \otimes \gamma \mapsto b \otimes h\gamma.$$

Proof. Let us define $c := \iota_2 \circ g$, where $\iota_2 : B \otimes \Omega^1(H) \rightarrow \Omega^1(B\#_\sigma H)$ is the canonical inclusion and $g : (B\#_\sigma H) \otimes {}^{\text{co}H}\Omega^1(H) \rightarrow B \otimes \Omega^1(H)$, $b \otimes h \otimes \gamma \mapsto b \otimes h\gamma$. We have already noted that ι_2 is a morphism in ${}_{B\#_\sigma H}\mathfrak{M}^H$. Moreover, g is left $B\#_\sigma H$ -linear, since

$$\begin{aligned}g((b' \otimes h') \cdot (b \otimes h \otimes \gamma)) &= g(b'(h'_1 \cdot b)\sigma(h'_2 \otimes h_1) \otimes h'_3h_2 \otimes \gamma) = b'(h'_1 \cdot b)\sigma(h'_2 \otimes h_1) \otimes h'_3h_2\gamma \\ &= b'(h'_1 \cdot b)\sigma(h'_2 \otimes h_1\gamma_{-1}) \otimes h'_3h_2\gamma_0 = b'(h'_1 \cdot b)\sigma(h'_2 \otimes (h\gamma)_{-1}) \otimes h'_3(h\gamma)_0 \\ &= (b' \otimes h') \cdot g(b \otimes h \otimes \gamma)\end{aligned}$$

for all $b, b' \in B$, $h, h' \in H$ and $\gamma \in {}^{\text{co}H}\Omega^1(H)$ and right H -colinear, indeed

$$g(b \otimes h \otimes \gamma)_0 \otimes g(b \otimes h \otimes \gamma)_1 = b \otimes (h\gamma)_0 \otimes (h\gamma)_1 = b \otimes h_1\gamma_0 \otimes h_2\gamma_1 = (g \otimes \text{Id}_H)(b \otimes h_1 \otimes \gamma_0 \otimes h_2\gamma_1).$$

Thus, g is a morphism in ${}_{B\#_\sigma H}\mathfrak{M}^H$ and then also $\iota_2 \circ g$ is in ${}_{B\#_\sigma H}\mathfrak{M}^H$. Since $\text{ver} = p \circ \pi_2$, where $p : B \otimes \Omega^1(H) \rightarrow (B\#_\sigma H) \otimes {}^{\text{co}H}\Omega^1(H)$, $b \otimes \gamma \mapsto b \otimes \gamma_{-2} \otimes S(\gamma_{-1})\gamma_0$ is the inverse of g , we obtain

$$\text{ver} \circ \iota_2 \circ g = p \circ \pi_2 \circ \iota_2 \circ g = p \circ g = \text{Id},$$

proving that $c = \iota_2 \circ g$ is a connection. It remains to prove that c is strong.

On elements $b \otimes h \in B \#_\sigma H$ the strong condition $(\text{Id} - c \circ \text{ver})d_{\#_\sigma}(B \#_\sigma H) \subseteq \Omega^1(B)(B \#_\sigma H)$ reads $d_B b \otimes h + b \otimes d_H h - c(b \otimes h_1 \otimes S(h_2)d_H h_3) \in \Omega^1(B)(B \#_\sigma H)$. The latter is satisfied, since

$$d_B b \otimes h + b \otimes d_H h - c(b \otimes h_1 \otimes S(h_2)d_H h_3) = d_B b \otimes h + b \otimes d_H h - b \otimes h_1 S(h_2) d_H h_3 = d_B b \otimes h = (d_B b \otimes 1_H)(1_B \otimes h)$$

holds. \square

Using the strong connection of Proposition 3.17 we can construct covariant derivatives of associated bundles of the crossed product QPB. Recall that, given a FODC $(\Omega^1(B), d_B)$ on an algebra B , a *covariant derivative* on a left B -module E is a linear map $\nabla_E: E \rightarrow \Omega^1(B) \otimes_B E$ obeying the left Leibniz rule

$$\nabla_E(b \cdot e) = d_B b \otimes_B e + b \cdot \nabla_E e,$$

for $e \in E$ and $b \in B$. In the notation of [3] we denote by $B\xi$ the category whose objects are given by pairs (E, ∇_E) where E is a left B -module and ∇_E is a covariant derivative on E and morphisms are left B -linear maps intertwining ∇ . More explicitly, a morphism between two objects (E, ∇_E) and (K, ∇_K) in $B\xi$ is a left B -linear map $T: E \rightarrow K$ such that $\nabla_K \circ T = (\text{Id} \otimes_B T) \circ \nabla_E$. If E is a B -bimodule, a covariant derivative $\nabla_E: E \rightarrow \Omega^1(B) \otimes_B E$ on E is called *bimodule covariant derivative* if it is equipped with a B -bimodule map $\sigma_E: E \otimes_B \Omega^1(B) \rightarrow \Omega^1(B) \otimes_B E$ obeying $\nabla_E(e \cdot b) = \sigma_E(e \otimes_B d_B b) + (\nabla_E e) \cdot b$, for $e \in E$ and $b \in B$. Note that σ_E is not an additional data as, if it exists, it is uniquely determined by the stated condition. Hence, being a bimodule covariant derivative or not is a property of a covariant derivative ∇_E on a given bimodule E . A particular class of interest is given by the *associated bundles* $E := (A \otimes V)^{\text{co}H}$, where $B := A^{\text{co}H} \subseteq A$ is a regular QPB and V is a right H -comodule. Here E is a B -bimodule with left and right B -actions given by multiplication on A . Those admit associated covariant derivatives, see [3, Proposition 5.48]. We give their explicit expression for the regular QPB on the crossed product algebra of Theorem 3.15: they correspond to the differential on the base algebra.

Proposition 3.18. *Given V a right H -comodule, then the B -bimodule $E := ((B \#_\sigma H) \otimes V)^{\text{co}H}$, with actions induced from the multiplication of $B \#_\sigma H$, acquires an associated covariant derivative $\nabla_E: E \rightarrow \Omega^1(B) \otimes_B E$ given by*

$$\nabla_E(b \otimes h \otimes v) = (\text{Id} - c \circ \text{ver})d_{\#_\sigma}(b \otimes h) \otimes v = d_B b \otimes h \otimes v,$$

where c is the canonical strong connection of Proposition 3.17. Moreover, ∇_E is a bimodule covariant derivative with associated $\sigma_E: E \otimes_B \Omega^1(B) \rightarrow \Omega^1(B) \otimes_B E$ defined by $\sigma_E(b \otimes h \otimes v \otimes_B \beta) = b(h_1 \cdot \beta) \otimes h_2 \otimes v$.

Proof. The first part follows by applying [3, Proposition 5.48] as we have discussed before. It remains to prove that σ_E is a B -bimodule map obeying $\nabla_E(e \cdot b) = \sigma_E(e \otimes_B d_B b) + (\nabla_E e) \cdot b$, for $e \in E$ and $b \in B$. We compute

$$\sigma_E(b' \cdot (b \otimes h \otimes v \otimes_B \beta)) = \sigma_E(b' b \otimes h \otimes v \otimes_B \beta) = b' b (h_1 \cdot \beta) \otimes h_2 \otimes v = b' \cdot \sigma_E(b \otimes h \otimes v \otimes_B \beta),$$

as well as

$$\begin{aligned} \sigma_E((b \otimes h \otimes v \otimes_B \beta) \cdot b') &= \sigma_E(b \otimes h \otimes v \otimes_B \beta b') = b(h_1 \cdot \beta b') \otimes h_2 \otimes v \\ &= b(h_1 \cdot \beta)(h_2 \cdot b') \otimes h_3 \otimes v = \sigma_E(b \otimes h \otimes v \otimes_B \beta) \cdot b' \end{aligned}$$

for all $b \otimes h \otimes v \in E$, $b' \in B$ and $\beta \in \Omega^1(B)$. Thus σ_E is a B -bimodule map. Moreover, we have

$$\begin{aligned} \nabla_E((b \otimes h \otimes v) \cdot b') &= \nabla_E(b(h_1 \cdot b') \otimes h_2 \otimes v) = d_B(b(h_1 \cdot b')) \otimes h_2 \otimes v \\ &= b d_B(h_1 \cdot b') \otimes h_2 \otimes v + (d_B b)(h_1 \cdot b') \otimes h_2 \otimes v \\ &= b(h_1 \cdot d_B b') \otimes h_2 \otimes v + (d_B b \otimes h \otimes v) \cdot b' \\ &= \sigma_E(b \otimes h \otimes v \otimes_B d_B b') + \nabla_E(b \otimes h \otimes v) \cdot b' \end{aligned}$$

and then the thesis follows. \square

We close this subsection by studying another application of the crossed product QPB.

Leray–Serre spectral sequence. The *noncommutative de Rham cohomology* of a DC (Ω^\bullet, d) on an algebra A is the graded algebra

$$H_{\text{dR}}^n(A) := \frac{\ker(d|_{\Omega^n})}{\text{Im}(d|_{\Omega^{n-1}})},$$

where we understand $d|_{\Omega^n} = 0$ if $n < 0$. In the following we show that the noncommutative de Rham cohomology of a crossed product algebra $B \#_\sigma H$ is determined by the noncommutative de Rham cohomology of the base

algebra B and of the structure Hopf algebra H . Namely, we prove that there is a spectral sequence with second page $E_2^{p,q} = H_{\text{dR}}^p(B) \otimes H_{\text{dR}}^q(H)$, which converges to $H_{\text{dR}}(B\#_\sigma H)$.

Recall that a *spectral sequence* of degree $r_0 \geq 0$ is a collection $(E_r^{\bullet,\bullet}, D_r)_{r \geq r_0}$ of \mathbb{Z} -bigraded vector spaces $E_r^{\bullet,\bullet} = \bigoplus_{k,\ell \in \mathbb{Z}} E_r^{k,\ell}$ (where $E_r^{k,\ell}$ is assumed to be zero if either $k < 0$ or $\ell < 0$) and linear maps $D_r: E_r^{k,\ell} \rightarrow E_r^{k+r,\ell-r+1}$ such that $D_r \circ D_r = 0$ and $E_{r+1}^{k,\ell} \cong H^{k,\ell}(E_r, D_r)$, where

$$H^{k,\ell}(E_r, D_r) := \frac{\ker D_r}{\text{Im } D_r|_{E_r^{k-r,\ell+r-1}}}.$$

One calls $(E_r^{\bullet,\bullet}, D_r)$ the r^{th} page of the spectral sequence. So the above means that the $(r+1)^{\text{th}}$ page is isomorphic to the cohomology of the r^{th} page.

Now we proceed by constructing a filtration of $\Omega^\bullet(B\#_\sigma H)$, which will then provide the announced spectral sequence. Recall from Theorem 3.14 that the right H -coaction $\rho: B\#_\sigma H \rightarrow (B\#_\sigma H) \otimes H$ is differentiable with differential $\Omega^\bullet(\rho): \Omega^\bullet(B\#_\sigma H) \rightarrow \Omega^\bullet((B\#_\sigma H) \otimes H)$ and let us denote the graded components by $\rho^{n-k,k}: \Omega^n(B\#_\sigma H) \rightarrow \Omega^{n-k}(B\#_\sigma H) \otimes \Omega^k(H)$. We define for $m, n \geq 0$

$$F^m(\Omega^n(B\#_\sigma H)) := \begin{cases} \bigcap_{k > n-m} \ker \rho^{n-k,k} & \text{if } m \leq n \\ 0 & \text{otherwise} \end{cases}$$

a filtration of $\Omega^\bullet(B\#_\sigma H)$. The corresponding spectral sequence $(E_r^{\bullet,\bullet}, D_r)_{r \geq 1}$ of degree 1 is constructed in the following way. Its first page is the \mathbb{Z} -bigraded vector space $E_1^{\bullet,\bullet}$, where

$$E_1^{p,q} := \frac{\ker \left(d_{\#_\sigma}: \frac{F^p(\Omega^{p+q}(B\#_\sigma H))}{F^{p+1}(\Omega^{p+q}(B\#_\sigma H))} \rightarrow \frac{F^p(\Omega^{p+q+1}(B\#_\sigma H))}{F^{p+1}(\Omega^{p+q+1}(B\#_\sigma H))} \right)}{\text{Im} \left(d_{\#_\sigma}: \frac{F^p(\Omega^{p+q-1}(B\#_\sigma H))}{F^{p+1}(\Omega^{p+q-1}(B\#_\sigma H))} \rightarrow \frac{F^p(\Omega^{p+q}(B\#_\sigma H))}{F^{p+1}(\Omega^{p+q}(B\#_\sigma H))} \right)}.$$

The differential $D_1: E_1^{p,q} \rightarrow E_1^{p+1,q}$ is the one induced from $d_{\#_\sigma}$ on the quotient and higher pages are defined as the cohomology of the previous pages. According to [3, Lemma 5.58] the spectral sequence $(E_r^{\bullet,\bullet}, D_r)_{r \geq 1}$ converges to $H_{\text{dR}}(B\#_\sigma H)$.

In the rest of this subsection we show that the above spectral sequence coincides with famous Leray–Serre spectral sequence (see [3, Theorem 4.66]). In order to do this we have to introduce higher order analogues of the vertical map. We recall once more from Theorem 3.14 that the right H -coaction $\rho: B\#_\sigma H \rightarrow (B\#_\sigma H) \otimes H$ is differentiable with

$$\begin{aligned} \Omega^\bullet(\rho): \Omega^\bullet(B\#_\sigma H) &\rightarrow \Omega^\bullet((B\#_\sigma H) \otimes H), \\ \omega \otimes \gamma &\mapsto \omega \otimes \gamma_{[1]} \otimes \gamma_{[2]}, \end{aligned}$$

where $\Omega^\bullet(\Delta)(\gamma) = \gamma_{[1]} \otimes \gamma_{[2]}$ is the differential of the coproduct $\Delta: H \rightarrow H \otimes H$ with graded components $\Omega^n(\Delta)(\gamma) = \sum_{k+l=n} \gamma_{\langle 1,k \rangle} \otimes \gamma_{\langle 2,l \rangle} = \sum_m \gamma_{\langle 1,m \rangle} \otimes \gamma_{\langle 2,n-m \rangle}$ for $\gamma \in \Omega^n(H)$. We denote the graded components of crossed product forms by $\omega^i \otimes \gamma^j \in \Omega^i(B) \otimes \Omega^j(H)$. Following [3, page 449] we define for all $n > 0$ and $n \geq m \geq 0$

$$\begin{aligned} \text{ver}^{m,n-m}: \Omega^n(B\#_\sigma H) &\rightarrow \Omega^m(B\#_\sigma H) \otimes {}^{\text{co}H}\Omega^{n-m}(H) \\ \sum_{i+j=n} \omega^i \otimes \gamma^j &\mapsto \sum_{i+j=n} \omega^i \otimes \gamma_{\langle 1,m \rangle}^j \otimes S((\gamma_{\langle 2,n-m \rangle}^j)_{-1})(\gamma_{\langle 2,n-m \rangle}^j)_0, \end{aligned}$$

where we recall that $\gamma_{\langle 1,m \rangle}^j \otimes \gamma_{\langle 2,n-m \rangle}^j = 0$ if $m > j$ or $n-m > j$. In other words, $\text{ver}^{m,n-m}$ is defined as the composition $(\text{Id} \otimes \phi^{n-m}) \circ \pi^{m,n-m} \circ \Omega^n(\rho)$, where $\pi^{m,n-m}: \Omega^n((B\#_\sigma H) \otimes H) \rightarrow \Omega^m(B\#_\sigma H) \otimes \Omega^{n-m}(H)$ is the canonical projection and $\phi^{n-m}: \Omega^{n-m}(H) \rightarrow {}^{\text{co}H}\Omega^{n-m}(H)$, $\gamma \mapsto S(\gamma_{-1})\gamma_0$ is the projection on left H -coinvariant elements.

Note that $\text{ver}^{0,1} = \text{ver}: \Omega^1(B\#_\sigma H) \rightarrow (B\#_\sigma H) \otimes {}^{\text{co}H}\Omega^1(H)$ is the classical vertical map. More in general, for $n > 0$, the vertical map $\text{ver}^{0,n}: \Omega^n(B\#_\sigma H) \rightarrow (B\#_\sigma H) \otimes {}^{\text{co}H}\Omega^n(H)$ is given for $\omega^i \in \Omega^i(B)$, $\gamma^i \in \Omega^i(H)$, $b \in B$ and $h \in H$ by

$$\text{ver}^{0,n}(\omega^n \otimes h + \omega^{n-1} \otimes \gamma^1 + \dots + \omega^1 \otimes \gamma^{n-1} + b \otimes \gamma^n) = b \otimes \gamma_{-2}^n \otimes S(\gamma_{-1}^n)\gamma_0^n.$$

We have shown in Theorem 3.15 that $\text{ver} = \text{ver}^{0,1}$ fits into a short exact sequence, the Atiyah sequence (22). It turns out that also $\text{ver}^{0,n}$ for $n > 0$ give rise to short exact sequences and, under a flatness assumption, the same is true for all higher vertical maps $\text{ver}^{m,n}$.

Proposition 3.19. *The sequence*

$$0 \longrightarrow \Omega^1(B) \wedge \Omega^{n-1}(B\#_\sigma H) \xrightarrow{i} \Omega^n(B\#_\sigma H) \xrightarrow{\text{ver}^{0,n}} (B\#_\sigma H) \otimes^{\text{co}H} \Omega^n(H) \longrightarrow 0 \quad (23)$$

is exact for all $n > 0$. If $\Omega^m(B)$ is flat as a right B -module for all $m > 0$ then

$$0 \longrightarrow \Omega^{m+1}(B) \wedge \Omega^{n-1}(B\#_\sigma H) \xrightarrow{i} \Omega^m(B) \wedge \Omega^n(B\#_\sigma H) \xrightarrow{\text{ver}^{m,n}} \Omega^m(B) \otimes_B (B\#_\sigma H) \otimes^{\text{co}H} \Omega^n(H) \longrightarrow 0 \quad (24)$$

is exact for all $n > 0$ and $m \geq 0$.

Proof. Since i is injective, for the exactness of (23) it remains to prove that the map $\text{ver}^{0,n}$ is surjective with $\ker(\text{ver}^{0,n}) = \Omega^1(B) \wedge \Omega^{n-1}(B\#_\sigma H)$ for every $n > 0$. Observe that $\text{ver}^{0,n} = p^n \circ \pi_n$ where $\pi_n : \Omega^n(B\#_\sigma H) \rightarrow B \otimes \Omega^n(H)$ is the canonical projection and $p^n : B \otimes \Omega^n(H) \rightarrow (B\#_\sigma H) \otimes^{\text{co}H} \Omega^n(H)$, $b \otimes \gamma^n \mapsto b \otimes \gamma_{-2}^n \otimes S(\gamma_{-1}^n) \gamma_0^n$ for all $\gamma^n \in \Omega^n(H)$. The inverse of p^n is given by $g^n : (B\#_\sigma H) \otimes^{\text{co}H} \Omega^n(H) \rightarrow B \otimes \Omega^n(H)$, $b \otimes h \otimes \gamma^n \mapsto b \otimes h \gamma^n$ as one easily verifies. Thus, $\text{ver}^{0,n}$ is surjective and $\ker(\text{ver}^{0,n}) = \ker(\pi_n) = \bigoplus_{k=0}^{n-1} \Omega^{n-k}(B) \otimes \Omega^k(H) = \Omega^1(B) \wedge \Omega^{n-1}(B\#_\sigma H)$. Then, under the flatness assumption, the exactness of (24) follows directly from [3, Lemma 5.60]. \square

The above proposition allows us to apply [3, Corollary 5.62], which states that the spectral sequence $(E_r^{\bullet, \bullet}, D_r)_{r \geq 1}$ is the Leray–Serre spectral sequence and, in case $\mathbb{H}_{\text{dR}}(H)$ is left-invariant, we obtain an explicit description of the second page in terms of the cohomology of the base algebra and structure Hopf algebra.

Proposition 3.20. *Let $(\Omega^\bullet(B), d_B)$ be a σ -twisted H -module DC on B and $(\Omega^\bullet(H), d_H)$ a bicovariant DC on H . If $\Omega^m(B)$ is flat as a right B -module for all $m > 0$ and $\mathbb{H}_{\text{dR}}(H)$ is left-invariant, then we get a spectral sequence with second page $E_2^{p,q} = \mathbb{H}_{\text{dR}}^p(B) \otimes \mathbb{H}_{\text{dR}}^q(H)$ which converges to $\mathbb{H}_{\text{dR}}(B\#_\sigma H)$.*

3.5 Connection 1-forms and fundamental vector fields

In this subsection we fix a Hopf algebra H with invertible antipode and a bicovariant FODC $(\Omega^1(H), d_H)$ on H , as well as a σ -twisted left H -module algebra B and a σ -twisted H -module FODC $(\Omega^1(B), d_B)$ on B . Moreover, we assume that ${}^{\text{co}H}\Omega^1(H)$ is finite-dimensional and consider $T_\varepsilon H = \text{Hom}_{\mathbb{k}}({}^{\text{co}H}\Omega^1(H), \mathbb{k})$. Recalling Remark 2.27, to any $\alpha \in T_\varepsilon H$ corresponds a vector field

$$\bar{\alpha} := (\text{Id} \otimes \alpha) \circ \text{ver} : \Omega^1(B\#_\sigma H) \rightarrow B\#_\sigma H,$$

which is vertical, since $\bar{\alpha}(\Omega^1(B) \otimes H) = 0$ and satisfies $\bar{\alpha}(1_B \otimes \gamma) = \alpha(\gamma)1_B \otimes 1_H$ for all $\gamma \in {}^{\text{co}H}\Omega^1(H)$. It is the unique vector field with these two properties (since $\bar{\alpha}$ is left $B\#_\sigma H$ -linear and every $b \otimes \gamma \in B \otimes \Omega^1(H)$ can be written as $(b \otimes \gamma_{-2}) \cdot (1_B \otimes S(\gamma_{-1})\gamma_0)$) and thus, following [18, Definition 4.6], we call $\bar{\alpha}$ the *fundamental vector field* corresponding to $\alpha \in T_\varepsilon H$.

Suppose that $\dim({}^{\text{co}H}\Omega^1(H)) = n$ and let $\{x_j\}_{j=1}^n$ be a basis of $T_\varepsilon H$ and $\{x^j\}_{j=1}^n$ be the dual basis of ${}^{\text{co}H}\Omega^1(H)$, which can be identified with $(T_\varepsilon H)^*$.

Definition 3.21. *A connection 1-form is an element $\phi = \sum_{j=1}^n x_j \otimes \phi_j \in (T_\varepsilon H \otimes \Omega^1(B\#_\sigma H))^{\text{co}H}$, satisfying $\sum_{j=1}^n x_j \otimes \bar{\alpha}(\phi_j) = \alpha \otimes 1_B \otimes 1_H$, for all $\alpha \in T_\varepsilon H$.*

This definition of connection 1-form is in complete analogy to [18, Definition 4.10], however in case of the crossed product, rather than the smash product calculus. The following lemma and proposition are the corresponding analogues of [18, Lemma 4.12 and Theorem 4.13]. We report here their proofs for the sake of completeness.

Lemma 3.22. *An element $\phi \in (T_\varepsilon H \otimes \Omega^1(B\#_\sigma H))^{\text{co}H}$ is a connection 1-form if and only if $(\text{Id} \otimes \pi_2)(\phi) = \sum_{j=1}^n x_j \otimes 1_B \otimes x^j$, where $\pi_2 : \Omega^1(B\#_\sigma H) \rightarrow B \otimes \Omega^1(H)$ is the canonical projection.*

Proof. First note that, given an element $\phi = \sum_{j=1}^n x_j \otimes \phi_j \in (T_\varepsilon H \otimes \Omega^1(B\#_\sigma H))^{\text{co}H}$, we can write $\phi_j = \phi'_j + \phi''_j$ where $\phi'_j \in \Omega^1(B) \otimes H$ and $\phi''_j \in B \otimes \Omega^1(H)$, for every $j = 1, \dots, n$. But now $\sum_{j=1}^n x_j \otimes \bar{\alpha}(\phi_j) = \sum_{j=1}^n x_j \otimes \bar{\alpha}(\phi''_j)$ for all $\alpha \in T_\varepsilon H$ and furthermore $(\text{Id} \otimes \pi_2)(\phi) = \sum_{j=1}^n x_j \otimes \phi''_j$. Thus, the claim of the lemma follows if we prove that $\sum_{j=1}^n x_j \otimes \bar{\alpha}(\phi''_j) = \alpha \otimes 1_B \otimes 1_H$ holds for all $\alpha \in T_\varepsilon H$ if and only if $\sum_{j=1}^n x_j \otimes \phi''_j = \sum_{j=1}^n x_j \otimes 1_B \otimes x^j$. Clearly, if we have that $\sum_{j=1}^n x_j \otimes \phi''_j = \sum_{j=1}^n x_j \otimes 1_B \otimes x^j$, then

$$\sum_{j=1}^n x_j \otimes \bar{\alpha}(\phi''_j) = \sum_{j=1}^n x_j \otimes (\text{Id} \otimes \alpha) \text{ver}(1_B \otimes x^j) = \sum_{j=1}^n \alpha(x^j) x_j \otimes 1_B \otimes 1_H = \alpha \otimes 1_B \otimes 1_H$$

for all $\alpha \in T_\varepsilon H$. On the other hand, suppose that $\sum_{j=1}^n x_j \otimes \bar{\alpha}(\phi_j'') = \alpha \otimes 1_B \otimes 1_H$ for all $\alpha \in T_\varepsilon H$. As a consequence, given $k, i \in \{1, \dots, n\}$, we have $\sum_{j=1}^n x_j(x^i) \bar{x}_k(\phi_j'') = x_k(x^i) 1_B \otimes 1_H$. We know that $\phi_j'' \in B \otimes \Omega^1(H)$ and that $g : (B \#_\sigma H) \otimes^{\text{co}H} \Omega^1(H) \rightarrow B \otimes \Omega^1(H)$, $b \otimes h \otimes \gamma \mapsto b \otimes h \gamma$ and $p : B \otimes \Omega^1(H) \rightarrow (B \#_\sigma H) \otimes^{\text{co}H} \Omega^1(H)$, $b \otimes \gamma \mapsto b \otimes \gamma_{-2} \otimes S(\gamma_{-1}) \gamma_0$ are inverse isomorphisms in $B \#_\sigma H \mathfrak{M}^H$. Thus, for all $j \in \{1, \dots, n\}$, we can write $\phi_j'' = \sum_{i=1}^n g(b_{i,j} \otimes h_{i,j} \otimes x^i) = \sum_{i=1}^n b_{i,j} \otimes h_{i,j} x^i$. Hence, we can compute

$$\begin{aligned} b_{i,j} \otimes h_{i,j} &= \sum_{l=1}^n b_{i,l} \otimes h_{i,l} x_j(x^l) = \sum_{l=1}^n (\text{Id} \otimes x_j) p g(b_{i,l} \otimes h_{i,l} \otimes x^l) = \sum_{l=1}^n (\text{Id} \otimes x_j) p(b_{i,l} \otimes h_{i,l} x^l) \\ &= \sum_{l=1}^n (\text{Id} \otimes x_j) \text{ver}(b_{i,l} \otimes h_{i,l} x^l) = \sum_{l=1}^n \bar{x}_j(b_{i,l} \otimes h_{i,l} x^l) = \sum_{k=1}^n \sum_{l=1}^n x_k(x^i) \bar{x}_j(b_{k,l} \otimes h_{k,l} x^l) \\ &= \sum_{k=1}^n x_k(x^i) \bar{x}_j \left(\sum_{l=1}^n b_{k,l} \otimes h_{k,l} x^l \right) = \sum_{k=1}^n x_k(x^i) \bar{x}_j(\phi_k'') = x_j(x^i) 1_B \otimes 1_H, \end{aligned}$$

for all $i, j \in \{1, \dots, n\}$. Thus, if $i \neq j$ we obtain that $b_{i,j} \otimes h_{i,j} = 0$ while if $i = j$ we obtain that $b_{j,j} \otimes h_{j,j} = 1_B \otimes 1_H$ and then $\phi_j'' = 1_B \otimes x^j$ for every $j = 1, \dots, n$. Hence $\sum_{j=1}^n x_j \otimes \phi_j'' = \sum_{j=1}^n x_j \otimes 1_B \otimes x^j$. \square

Remark 3.23. Given a connection 1-form $\phi = \sum_{j=1}^n x_j \otimes \phi_j$, we then have that

$$\phi = \sum_{j=1}^n x_j \otimes \phi_j' + \sum_{j=1}^n x_j \otimes 1_B \otimes x^j$$

where $\phi_j' \in \Omega^1(B) \otimes H$ for $j = 1, \dots, n$. Thus, for all $\gamma \in^{\text{co}H} \Omega^1(H)$, we obtain that

$$\sum_{j=1}^n x_j(\gamma) \phi_j = \sum_{j=1}^n x_j(\gamma) \phi_j' + \sum_{j=1}^n x_j(\gamma) 1_B \otimes x^j = \sum_{j=1}^n x_j(\gamma) \phi_j' + 1_B \otimes \gamma$$

and then

$$\sum_{j=1}^n x_j(\gamma) \text{ver}(\phi_j) = \sum_{j=1}^n x_j(\gamma) \text{ver}(\phi_j') + \text{ver}(1_B \otimes \gamma) = 1_B \otimes 1_H \otimes \gamma. \quad (25)$$

Observe also that the coevaluation map $\mathbb{k} \rightarrow (T_\varepsilon H) \otimes (T_\varepsilon H)^*$, $1_{\mathbb{k}} \mapsto \sum_{j=1}^n x_j \otimes x^j$ is right H -colinear. Thus, since $\sum_{j=1}^n x_{j0} \otimes 1_B \otimes x_0^j \otimes x_{j1} x_1^j = \sum_{j=1}^n x_j \otimes 1_B \otimes x^j \otimes 1_H$, the fact that ϕ is coinvariant is expressed by $\sum_{j=1}^n x_{j0} \otimes \phi_{j0}' \otimes x_{j1} \phi_{j1}' = \sum_{j=1}^n x_j \otimes \phi_j' \otimes 1_H$.

Proposition 3.24. For the crossed product calculus there is a bijection between connections and connection 1-forms.

Proof. First suppose to have a connection $c : (B \#_\sigma H) \otimes^{\text{co}H} \Omega^1(H) \rightarrow \Omega^1(B \#_\sigma H)$, i.e., a morphism in $B \#_\sigma H \mathfrak{M}^H$ such that $\text{ver} \circ c = \text{Id}$ and define $\phi_c := \sum_{j=1}^n x_j \otimes c(1_B \otimes 1_H \otimes x^j) \in T_\varepsilon H \otimes \Omega^1(B \#_\sigma H)$. Observe that $\phi_c \in (T_\varepsilon H \otimes \Omega^1(B \#_\sigma H))^{\text{co}H}$, indeed

$$\begin{aligned} \sum_{j=1}^n x_{j0} \otimes c(1_B \otimes 1_H \otimes x^j)_0 \otimes x_{j1} c(1_B \otimes 1_H \otimes x^j)_1 &= \sum_{j=1}^n x_{j0} \otimes c(1_B \otimes 1_H \otimes x_0^j) \otimes x_{j1} x_1^j \\ &= \sum_{j=1}^n x_j \otimes c(1_B \otimes 1_H \otimes x^j) \otimes 1_H \end{aligned}$$

since c is right H -colinear as well as the map $\mathbb{k} \in 1_{\mathbb{k}} \mapsto \sum_{j=1}^n x_j \otimes x^j \in T_\varepsilon H \otimes (T_\varepsilon H)^*$. Furthermore, we have

$$\sum_{j=1}^n x_j \otimes \bar{\alpha}(c(1_B \otimes 1_H \otimes x^j)) = \sum_{j=1}^n x_j \otimes (\text{Id} \otimes \alpha) \text{ver}(c(1_B \otimes 1_H \otimes x^j)) = \sum_{j=1}^n \alpha(x^j) x_j \otimes 1_B \otimes 1_H = \alpha \otimes 1_B \otimes 1_H,$$

thus ϕ_c is a connection 1-form.

On the other hand, if we have a connection 1-form $\phi = \sum_{j=1}^n x_j \otimes \phi_j \in (T_\varepsilon H \otimes \Omega^1(B \#_\sigma H))^{\text{co}H}$, we can define $c_\phi : (B \#_\sigma H) \otimes^{\text{co}H} \Omega^1(H) \rightarrow \Omega^1(B \#_\sigma H)$ as the morphism in $B \#_\sigma H \mathfrak{M}^H$ given by $c_\phi(1_B \otimes 1_H \otimes \gamma) := \sum_{j=1}^n x_j(\gamma) \phi_j$, so that $c_\phi(b \otimes h \otimes \gamma) = (b \otimes h) \cdot c_\phi(1_B \otimes 1_H \otimes \gamma) = \sum_{j=1}^n x_j(\gamma) ((b \otimes h) \cdot \phi_j)$. We also have that c_ϕ is right

H -colinear, indeed

$$\begin{aligned}
c_\phi(b \otimes h \otimes \gamma)_0 \otimes c_\phi(b \otimes h \otimes \gamma)_1 &= \sum_{j=1}^n x_j(\gamma)((b \otimes h) \cdot \phi_j)_0 \otimes ((b \otimes h) \cdot \phi_j)_1 = \sum_{j=1}^n x_j(\gamma)((b \otimes h_1) \cdot \phi_{j0}) \otimes h_2 \phi_{j1} \\
&= \sum_{j=1}^n x_j(\gamma_0)((b \otimes h_1) \cdot \phi_{j0}) \otimes h_2 \varepsilon(\gamma_1) \phi_{j1} = \sum_{j=1}^n ((b \otimes h_1) \cdot \phi_{j0}) \otimes h_2 \gamma_2 x_j(\gamma_0) S^{-1}(\gamma_1) \phi_{j1} \\
&= \sum_{j=1}^n x_{j0}(\gamma_0)((b \otimes h_1) \cdot \phi_{j0}) \otimes h_2 \gamma_1 x_{j1} \phi_{j1} = \sum_{j=1}^n x_j(\gamma_0)((b \otimes h_1) \cdot \phi_j) \otimes h_2 \gamma_1 \\
&= c(b \otimes h_1 \otimes \gamma_0) \otimes h_2 \gamma_1,
\end{aligned}$$

using that the left $B \#_\sigma H$ -action of $\Omega^1(B \#_\sigma H)$ is right H -colinear and that ϕ is a coinvariant element. Thus, c_ϕ is a morphism in ${}_{B \#_\sigma H} \mathfrak{M}^H$ and, since ver is left $B \#_\sigma H$ -linear, we also have

$$\text{ver} c_\phi(b \otimes h \otimes \gamma) = (b \otimes h) \cdot \sum_{j=1}^n x_j(\gamma) \text{ver}(\phi_j) \stackrel{(25)}{=} (b \otimes h) \cdot (1_B \otimes 1_H \otimes \gamma) = b \otimes h \otimes \gamma,$$

so that c_ϕ is a connection.

It remains to prove that we have a bijection. If we start with a connection c with corresponding connection 1-form $\phi_c = \sum_{j=1}^n x_j \otimes c(1_B \otimes 1_H \otimes x^j)$ and then consider the connection c_{ϕ_c} , we obtain

$$c_{\phi_c}(b \otimes h \otimes \gamma) = \sum_{j=1}^n x_j(\gamma)((b \otimes h) \cdot c(1_B \otimes 1_H \otimes x^j)) = \sum_{j=1}^n x_j(\gamma) c(b \otimes h \otimes x^j) = c(b \otimes h \otimes \gamma),$$

i.e., $c_{\phi_c} = c$. On the other hand, if we start with a connection 1-form $\phi = \sum_{j=1}^n x_j \otimes \phi_j$ with corresponding connection c_ϕ and then consider the connection 1-form $\phi_{c_\phi} = \sum_{j=1}^n x_j \otimes c_\phi(1_B \otimes 1_H \otimes x^j)$, we obtain $\phi_{c_\phi} = \phi$, since $c_\phi(1_B \otimes 1_H \otimes x^j) = \sum_{i=1}^n x_i(x^j) \phi_i = \phi_j$ and thus the thesis follows. \square

4 Construction for pointed Hopf algebras

We apply the construction of the FODC on $B \#_\sigma H$ developed in Theorem 3.7 to build examples of FODCi on pointed Hopf algebras.

Pointed Hopf algebras as crossed products. Recall that a coalgebra C is called *simple* if it is not 0 and if it has no non-zero proper subcoalgebras and that C is called *pointed* if all its simple subcoalgebras are 1-dimensional. A bialgebra (Hopf algebra) is pointed if it is pointed as a coalgebra. Since the set of group-like elements is in 1-1 correspondence with the set of 1-dimensional subcoalgebras via $g \mapsto \mathbb{k}g$ (see e.g. [24, Lemma 8.0.1 e)], C is pointed if and only if the coradical C_0 of C , i.e., the subcoalgebra of C given by the sum of all its simple subcoalgebras, is the group algebra $\mathbb{k}[G(C)]$. Among the classical examples of pointed Hopf algebras are group algebras, universal enveloping algebras and the q -deformations of the universal enveloping algebras of semisimple Lie algebras. It is known that pointed Hopf algebras can be seen as crossed products as we recall in the following.

Proposition 4.1 ([19, Theorem 15.2.4]). *Every pointed Hopf algebra H is isomorphic to $H_1 \#_\sigma \mathbb{k}[G/N]$, where H_1 is the indecomposable component of H containing $\mathbb{k}1_H$ and $\mathbb{k}[G/N]$ is the group Hopf algebra over G/N , with G and N the groups of group-like elements of H and H_1 , respectively.*

Let us recall some details of this result. Given $\{g_i\}_{i \in \mathcal{I}}$ a complete set of representatives for the left cosets of N in G , we have that $H = \bigoplus_{i \in \mathcal{I}} H_{g_i} = \bigoplus_{i \in \mathcal{I}} H_1 g_i$ by [19, Proposition 15.2.3]. Let $\bar{g}_i = g_i N$ for all $i \in \mathcal{I}$. The measure $\cdot : \mathbb{k}[G/N] \otimes H_1 \rightarrow H_1$ is given by $\bar{g}_i \cdot a := g_i a g_i^{-1}$, while the 2-cocycle $\sigma : \mathbb{k}[G/N] \otimes \mathbb{k}[G/N] \rightarrow H_1$ is defined in the following way. Given $i, j \in \mathcal{I}$, then $\bar{g}_i \bar{g}_j = \bar{g}_l$ for a unique $l \in \mathcal{I}$. Hence there is a unique $n_{i,j} \in N \subseteq H_1$, such that $g_i g_j = n_{i,j} g_l$. One then defines σ by $\sigma(\bar{g}_i \otimes \bar{g}_j) := n_{i,j}$. Moreover, the isomorphism of Hopf algebras $f : H \rightarrow H_1 \#_\sigma \mathbb{k}[G/N]$ is given by $f(a g_i) = a \otimes \bar{g}_i$ for all $a \in H_1$ and $i \in \mathcal{I}$, where $H_1 \#_\sigma \mathbb{k}[G/N]$ has the tensor product coalgebra structure.

Differential calculus on pointed Hopf algebras. Let $H \cong H_1 \#_\sigma \mathbb{k}[G/N]$ be a pointed Hopf algebra with notation as before. In the following we build a right $\mathbb{k}[G/N]$ -covariant calculus on H . Starting from a σ -twisted $\mathbb{k}[G/N]$ -module FODC on H_1 and a bicovariant FODC on $\mathbb{k}[G/N]$, we then use the construction of Theorem 3.7. The bicovariant FODCi on $\mathbb{k}[G/N]$ are described by Theorem 2.18. In particular, we know that every

bicovariant FODC $(\Omega^1(\mathbb{k}[G/N]), d_{\mathbb{k}[G/N]})$ on $\mathbb{k}[G/N]$ is such that $\Omega^1(\mathbb{k}[G/N]) := \mathbb{k}[G/N] \otimes (\mathbb{k}[G/N]^+ / I)$, where $I \subseteq \mathbb{k}[G/N]^+$ is a right ideal of $\mathbb{k}[G/N]$ and $d_{\mathbb{k}[G/N]}\bar{g}_i = (\text{Id} \otimes \pi)(\bar{g}_i \otimes \bar{g}_i - \bar{g}_i \otimes \bar{1}) = \bar{g}_i \otimes [\bar{g}_i - \bar{1}]$, where $\pi : \mathbb{k}[G/N]^+ \rightarrow \mathbb{k}[G/N]^+ / I$, $\alpha \mapsto [\alpha]$ is the quotient map. Note that the condition $\text{Ad}_R(I) \subseteq I \otimes \mathbb{k}[G/N]$ is automatically satisfied. Observe also that $\mathbb{k}[G/N]^+$ consists exactly of those elements $\sum_{i \in \mathcal{I}} k_i \bar{g}_i$ in $\mathbb{k}[G/N]$ such that $\sum_{i \in \mathcal{I}} k_i = 0$, that we denote by $k_i \bar{g}_i$, summation understood. Left and right $\mathbb{k}[G/N]$ -actions are defined as

$$\bar{g}_l \cdot (\bar{g}_j \otimes [k_i \bar{g}_i]) := \bar{g}_l \bar{g}_j \otimes [k_i \bar{g}_i] \quad \text{and} \quad (\bar{g}_j \otimes [k_i \bar{g}_i]) \cdot \bar{g}_l := \bar{g}_j \bar{g}_l \otimes [k_i \bar{g}_i \bar{g}_l],$$

while left and right $\mathbb{k}[G/N]$ -coactions are given by

$$\lambda(\bar{g}_j \otimes [k_i \bar{g}_i]) := \bar{g}_j \otimes \bar{g}_j \otimes [k_i \bar{g}_i] \quad \text{and} \quad \rho(\bar{g}_j \otimes [k_i \bar{g}_i]) := \bar{g}_j \otimes [k_i \bar{g}_i] \otimes \bar{g}_j.$$

We also need a σ -twisted $\mathbb{k}[G/N]$ -module FODC on H_1 , i.e., a FODC $(\Omega^1(H_1), d_{H_1})$ such that there exists a linear map $\cdot : \mathbb{k}[G/N] \otimes \Omega^1(H_1) \rightarrow \Omega^1(H_1)$ which satisfies (12), (13) and (14) of Lemma 3.2. In particular, from (12) and (13) we know that this linear map, if it exists, is uniquely determined by

$$\bar{g}_i \cdot (\text{ad}_{H_1} a') = (\bar{g}_i \cdot a) d_{H_1}(\bar{g}_i \cdot a') = g_i a g_i^{-1} d_{H_1}(g_i a' g_i^{-1}) \quad (26)$$

for every $a, a' \in H_1$ and $\bar{g}_i \in \mathbb{k}[G/N]$. Thus, one has to prove that (26) is well-defined. Furthermore, we need $d_{H_1}(n_{i,j}) = 0$ for every $n_{i,j} \in N$ and $i, j \in \mathcal{I}$ in order to satisfy (14). In particular, (14) holds true if $d_{H_1}|_N = 0$. We summarize the previous discussion in the following statement.

Proposition 4.2. *Given a pointed Hopf algebra $H \cong H_1 \#_{\sigma} \mathbb{k}[G/N]$, a right ideal $I \subseteq \mathbb{k}[G/N]^+$ and a σ -twisted $\mathbb{k}[G/N]$ -module FODC $(\Omega^1(H_1), d_{H_1})$ on H_1 we obtain a right $\mathbb{k}[G/N]$ -covariant FODC $(\Omega^1(H), d_H)$ on H .*

Observe that the above calculus on H is not right H -covariant in general and thus it does not fit the classification Theorem 2.18 (Remark 2.19) of Woronowicz.

Let us consider some special cases of the construction of Proposition 4.2.

Remark 4.3. *i.) If H is a cocommutative pointed Hopf algebra then $N = \{1\}$ by [19, Proposition 15.3.1] and so $G/N = G$. Hence, the measure becomes $g \cdot a = g a g^{-1}$ for all $g \in G$, $a \in H_1$ and the 2-cocycle is such that $\sigma(g \otimes g') = 1_H$ for all $g, g' \in G$. As a consequence, H is isomorphic to the smash product $H_1 \# \mathbb{k}G$ as it is said in [19, Theorem 15.3.2]. In this case, given a right ideal $I \subseteq \mathbb{k}[G]^+$ and a $\mathbb{k}[G]$ -module FODC on H_1 , by Proposition 4.2 we obtain a right $\mathbb{k}[G]$ -covariant FODC on H .*

ii.) Two classical examples of pointed Hopf algebras are the group Hopf algebra $\mathbb{k}[\tilde{G}]$ for a group \tilde{G} and the universal enveloping algebra $U(\mathfrak{g})$ for a Lie algebra \mathfrak{g} , which are both cocommutative, thus special cases of i.). Furthermore, for these two examples, the isomorphism with the smash product is trivial. Indeed, $(\mathbb{k}[\tilde{G}])_1 = \mathbb{k}1_{\tilde{G}}$ since $\mathbb{k}[\tilde{G}] = \bigoplus_{g \in \tilde{G}} \mathbb{k}g$ as coalgebras and $G(\mathbb{k}[\tilde{G}]) = \tilde{G}$, while $G(U(\mathfrak{g})) = 1$ and $U(\mathfrak{g})_1 = U(\mathfrak{g})$, since $U(\mathfrak{g})$ is irreducible (see [19, Corollary 7.6.5]) and the irreducible components coincide with the indecomposable ones in the cocommutative case (see [19, Proposition 15.3.1]). In the first case $\Omega^1(\mathbb{k}1_{\tilde{G}} \# \mathbb{k}[\tilde{G}]) = \mathbb{k}1_{\tilde{G}} \otimes \Omega^1(\mathbb{k}[\tilde{G}])$ and then the right $\mathbb{k}[\tilde{G}]$ -covariant calculus on $\mathbb{k}[\tilde{G}]$ is simply the starting bicovariant FODC on $\mathbb{k}[\tilde{G}]$. In the second case $\Omega^1(U(\mathfrak{g}) \# \mathbb{k}1) = \Omega^1(U(\mathfrak{g})) \otimes \mathbb{k}1$ and then the right covariant FODC on $U(\mathfrak{g})$ is simply the starting calculus on $U(\mathfrak{g})$ where $\Omega^1(U(\mathfrak{g}))$ is equipped with trivial coaction.

Now we consider a class of examples of pointed Hopf algebras not cocommutative in order to preserve the pure crossed product in the isomorphism and we build the corresponding right covariant FODC explicitly.

The pointed Hopf algebras $H_{(r,n,q)}$. Consider the pointed Hopf algebras $H_{(r,n,q)}$ given in [19, page 477] (see also [20] where they were introduced). Thus, let $r, n \geq 1$ and $M = rn$ and suppose that \mathbb{k} has a primitive M^{th} root of unity q . Let $H = H_{(r,n,q)}$ be the Hopf algebra over \mathbb{k} , sometimes referred to as *Radford Hopf algebras*, described as follows. As an algebra H is generated by a and x subject to the relations

$$a^M = 1, \quad x^n = 0, \quad xa = qax$$

and the coalgebra structure of H is given by

$$\Delta(a) = a \otimes a, \quad \Delta(x) = 1 \otimes x + x \otimes a^r.$$

In this case, using the notation above, $G = (a)$ and $N = (a^r)$, so that $G/N = \{\bar{a}^i \mid i = 0, \dots, r-1\} \cong C_r$, where C_r is the cyclic group with r elements. Hence $H_{(r,n,q)}$ has r distinct indecomposable components: $H_1, H_1 a, \dots, H_1 a^{r-1}$. Observe that, considering $r = 1$, we obtain the Taft algebra, which is indecomposable and has $G = N = (a)$. We know that $H_{(r,n,q)}$ has a basis given by $\{a^l x^m \mid 0 \leq l < M, 0 \leq m < n\}$, so that $H_{(r,n,q)} = \bigoplus_{0 \leq l < M, 0 \leq m < n} \mathbb{k} a^l x^m$ as vector spaces. Thus, it is not difficult to deduce that

$$H_1 = \bigoplus_{0 \leq m < n} \mathbb{k} x^m \oplus \bigoplus_{0 \leq m < n} \mathbb{k} a^r x^m \oplus \dots \oplus \bigoplus_{0 \leq m < n} \mathbb{k} a^{r(n-1)} x^m$$

and we know that $H \cong H_1 \#_{\sigma} \mathbb{k}[C_r]$ by Proposition 4.1. So, an element $\chi \in H_1$ is of the form $\chi = \sum_{0 \leq m, l < n} \alpha_{l,m} a^{lr} x^m$, with $\alpha_{l,m} \in \mathbb{k}$ and then the measure on H_1 is given, for all $i \in \{0, \dots, r-1\}$, by

$$\overline{a^i} \cdot \chi = a^i \chi a^{M-i} = \sum_{0 \leq m, l < n} \alpha_{l,m} a^i a^{lr} x^m a^{M-i} = \sum_{0 \leq m, l < n} \alpha_{l,m} q^{m(M-i)} a^i a^{lr} a^{M-i} x^m = \sum_{0 \leq m, l < n} \alpha_{l,m} q^{m(M-i)} a^{lr} x^m$$

and thus we simply change the coefficients of χ . Consider a FODC $(\Omega^1(H_1), d_{H_1})$ on H_1 and two elements $a^{lr} x^m$ and $a^{kr} x^s$ of the basis of H_1 for indexes $0 \leq l, k, m, s < n$. Then $a^{lr} x^m d_{H_1}(a^{kr} x^s) \in \Omega^1(H_1)$ and we obtain

$$\begin{aligned} \overline{a^i} \cdot (a^{lr} x^m d_{H_1}(a^{kr} x^s)) &= a^i a^{lr} x^m a^{M-i} d_{H_1}(a^i a^{kr} x^s a^{M-i}) = q^{m(M-i)} a^{lr} x^m d_{H_1}(q^{s(M-i)} a^{kr} x^s) \\ &= q^{(M-i)(m+s)} a^{lr} x^m d_{H_1}(a^{kr} x^s). \end{aligned}$$

This shows that (26) is well-defined for every FODC on H_1 . Furthermore, note that for $l, k \in \{0, \dots, r-1\}$ we have $0 \leq l+k \leq 2r-2$. Thus, if $0 \leq l+k \leq r-1$ we have that $a^l a^k = 1 a^{l+k}$ for unique $1 \in N$, while if $r \leq l+k \leq 2r-2$ we have that $a^l a^k = a^r a^{l+k-r}$ with $0 \leq l+k-r \leq r-2$, for unique $a^r \in N$. This implies that, in order to satisfy (14), we have to require $d_{H_1}(a^r) = 0$. So, if the FODC on H_1 satisfies $d_{H_1}(a^r) = 0$, assuming that $d_{H_1}(a^{kr}) = 0$ for k an arbitrary positive integer, we obtain

$$d_{H_1}(a^{(k+1)r}) = d_{H_1}(a^{kr} a^r) = a^{kr} d_{H_1}(a^r) + d_{H_1}(a^{kr}) a^r = 0$$

so that, by induction, $d_{H_1}(N) = 0$. For a generic element $\chi = \sum_{0 \leq m, l < n} \alpha_{l,m} a^{lr} x^m \in H_1$ this gives

$$d_{H_1}(\chi) = \sum_{0 \leq m, l < n} \alpha_{l,m} a^{lr} d_{H_1}(x^m) + \sum_{0 \leq m, l < n} \alpha_{l,m} d_{H_1}(a^{lr}) x^m = \sum_{0 \leq m, l < n} \alpha_{l,m} a^{lr} d_{H_1}(x^m). \quad (27)$$

Thus, we have obtained the following result:

Proposition 4.4. *Given the pointed Hopf algebra $H_{(r,n,q)} \cong (H_{(r,n,q)})_1 \#_{\sigma} \mathbb{k}[C_r]$, a right ideal $I \subseteq \mathbb{k}[C_r]^+$ and a FODC on $(H_{(r,n,q)})_1$ which satisfies (27) we obtain a right $\mathbb{k}[C_r]$ -covariant FODC on $H_{(r,n,q)}$.*

Let us consider a more specific example in order to show all details of the right covariant FODC.

Example 4.5. *Let $r = 2$ and $n = 2$. The generators a and x are subject to the relations*

$$a^4 = 1, \quad x^2 = 0, \quad xa = qax,$$

where q is a primitive fourth root of unity. In this case we have two indecomposable components, i.e., H_1 and $H_1 a$, where

$$H_1 = \bigoplus_{0 \leq m < 2} \mathbb{k} x^m \oplus \bigoplus_{0 \leq m < 2} \mathbb{k} a^2 x^m = \mathbb{k} 1 \oplus \mathbb{k} x \oplus \mathbb{k} a^2 \oplus \mathbb{k} a^2 x.$$

Therefore, for $H = H_{(2,2,q)}$ an element $\chi \in H_1$ is of the form $\chi = \alpha 1 + \beta x + \gamma a^2 + \delta a^2 x$ for $\alpha, \beta, \gamma, \delta \in \mathbb{k}$ and $G/N = \{\bar{1}, \bar{a}\} \cong C_2$. The σ -twisted $\mathbb{k}[C_2]$ -module FODC on H_1 is determined by $d_{H_1} x$, so that

$$d_{H_1}(\chi) = \alpha d_{H_1} 1 + \beta d_{H_1} x + \gamma d_{H_1}(a^2) + \delta d_{H_1}(a^2 x) = \beta d_{H_1} x + \delta a^2 d_{H_1} x.$$

The only right ideals I of $\mathbb{k}[C_2]$ which are contained in $\mathbb{k}[C_2]^+ = \mathbb{k}(\bar{a} - \bar{1})$ are $I = \{0\}$ and $I = \mathbb{k}(\bar{a} - \bar{1})$. Since the latter choice gives a trivial quotient we elaborate on the first one, which corresponds to the universal FODC. For $I = \{0\}$ we have that $\Omega^1(\mathbb{k}[C_2]) := \mathbb{k}[C_2] \otimes \mathbb{k}(\bar{a} - \bar{1})$ and $d_{\mathbb{k}[C_2]} \bar{a} = \bar{a} \otimes (\bar{a} - \bar{1})$. Thus, $d_{\mathbb{k}[C_2]}(e\bar{1} + f\bar{a}) = f\bar{a} \otimes (\bar{a} - \bar{1})$, for $e, f \in \mathbb{k}$ and the right $\mathbb{k}[C_2]$ -covariant FODC $(\Omega^1(H_1 \#_{\sigma} \mathbb{k}[C_2]), d_{\#_{\sigma}})$ on $H_1 \#_{\sigma} \mathbb{k}[C_2]$, where $\Omega^1(H_1 \#_{\sigma} \mathbb{k}[C_2]) = (\Omega^1(H_1) \otimes \mathbb{k}[C_2]) \oplus (H_1 \otimes \mathbb{k}[C_2] \otimes \mathbb{k}(\bar{a} - \bar{1}))$, is given by

$$\begin{aligned} d_{\#_{\sigma}}(\chi \otimes (e\bar{1} + f\bar{a})) &= d_{H_1}(\chi) \otimes (e\bar{1} + f\bar{a}) + \chi \otimes d_{\mathbb{k}[C_2]}(e\bar{1} + f\bar{a}) \\ &= (\beta d_{H_1} x + \delta a^2 d_{H_1} x) \otimes (e\bar{1} + f\bar{a}) + (\alpha 1 + \beta x + \gamma a^2 + \delta a^2 x) \otimes f\bar{a} \otimes (\bar{a} - \bar{1}) \\ &= \beta e d_{H_1} x \otimes \bar{1} + \beta f d_{H_1} x \otimes \bar{a} + \delta e a^2 d_{H_1} x \otimes \bar{1} + \delta f a^2 d_{H_1} x \otimes \bar{a} \\ &\quad + \alpha f 1 \otimes \bar{a} \otimes (\bar{a} - \bar{1}) + \beta f x \otimes \bar{a} \otimes (\bar{a} - \bar{1}) + \gamma f a^2 \otimes \bar{a} \otimes (\bar{a} - \bar{1}) + \delta f a^2 x \otimes \bar{a} \otimes (\bar{a} - \bar{1}) \end{aligned}$$

for all $\chi \otimes (e\bar{1} + f\bar{a}) \in H_1 \#_{\sigma} \mathbb{k}[C_2]$.

Acknowledgements

We would like to thank P. Aschieri for valuable suggestions throughout all phases of this work. In particular, he conjectured the results of Section 3.2. This paper was written while the first author was member of the ‘‘National Group for Algebraic and Geometric Structures and their Applications’’ (GNSAGA-INdAM). He was partially supported by MIUR within the National Research Project PRIN 2017. The second author was partially supported by HORIZON-MSCA-2021-SE-01-101086123 CaLIGOLA and by INFN Sezione di Bologna (Gast).

5 References

- [1] P. Aschieri, G. Landi, C. Pagani, *Braided Hopf algebras and gauge transformations*. Preprint arXiv:2203.13811.
- [2] R. Aziz, S. Majid, *Quantum differentials on cross product Hopf algebras*. J. Algebra, **563**, 1 (2020) 303-351.
- [3] E.J. Beggs, S. Majid, *Quantum Riemannian geometry*. Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], **355**. Springer, Cham, 2020.
- [4] A. Borowiec, *Cartan pairs*. Czech. J. Phys., **46** (1996) 1197-1202.
- [5] A. Borowiec, *Vector Fields and differential operators: noncommutative case*. Czech. J. Phys., **47** (1997) 1097-1100.
- [6] T. Brzeziński, G., Janelidze, T. Maszczyk, *Galois structures*, in P.M. Hajac (ed.), *Lecture Notes on Noncommutative Geometry and Quantum Groups*. <http://www.mimuw.edu.pl/~pwit/toknotes/toknotes>
- [7] T. Brzeziński, S. Majid, *Quantum group gauge theory on quantum spaces*. Comm. Math. Phys., **157** (1993) 591-638.
- [8] T. Brzeziński, B. Zieliński, *Quantum principal bundles over quantum real projective spaces*. J. Geom. Phys., **62**, 5 (2012) 1097-1107.
- [9] A. Connes, *Noncommutative Geometry*. Academic Press, Inc., San Diego, CA, 1994.
- [10] Y. Doi, M. Takeuchi, *Cleft comodule algebras for a bialgebra*. Commun. Algebra, **14**, 5 (1986) 801-817.
- [11] M. Durdević, *Geometry of quantum principal bundles I*. Comm. Math. Phys., **175**, 3 (1996) 457-520.
- [12] P.M. Hajac, *Strong connections on quantum principal bundles*. Commun. Math. Phys., **182** (1996) 579-617.
- [13] A. Klimyk, K. Schmüdgen, *Quantum Groups and Their Representations*. Texts and Monog. in Physics, Springer-Verlag Berlin, 1997.
- [14] H.F. Kreimer, M. Takeuchi, *Hopf Algebras and Galois Extensions of an Algebra*. Indiana Univ. Math. J., **30**, 5 (1981) 675-692.
- [15] A. Krutov, R. Ó Buachalla, K. Strung, *Nichols Algebras and Quantum Principal Bundles*. International Mathematics Research Notices, rnac366 (2023).
- [16] S. Majid, *Foundations of Quantum Group Theory*. Cambridge University Press, 1995.
- [17] S. Montgomery, *Hopf algebras and their actions on rings*. CBMS Regional Conference Series in Math, vol. **82**, AMS, 1994.
- [18] M. Pflaum, P. Schauenburg, *Differential calculi on noncommutative bundles*. Z Phys C-Particles and Fields, **76** (1997) 733-744.
- [19] D. E. Radford, *Hopf algebras*. Series on Knots and Everything, **49**. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2012.
- [20] D. E. Radford, *On Kauffman's knot invariants arising from finite-dimensional Hopf algebras*. Advances in Hopf algebras (Chicago, IL, 1992), 205-266, Lecture Notes in Pure and Appl. Math., **158**, Dekker, New York, 1994.
- [21] P. Schauenburg, *Differential-Graded Hopf Algebras and Quantum Group Differential Calculi*. J. Algebra, **180** (1996) 239-286.
- [22] P. Schauenburg, *Hopf Modules and Yetter-Drinfel'd Modules*. J. Algebra, **169** (1994) 874-890.
- [23] H-J. Schneider, *Principal homogeneous spaces for arbitrary Hopf algebras*. Isr. J. Math., **72** (1990) 167-195.
- [24] M.E. Sweedler, *Hopf algebras*. Mathematics Lecture Note Series W. A. Benjamin, Inc., New York, 1969.
- [25] M. Takeuchi, *Relative Hopf modules-equivalences and freeness conditions*. J. Algebra, **60** (1979) 452-471.
- [26] S.L. Woronowicz, *Differential calculus on compact matrix pseudogroups (quantum groups)*. Comm. Math. Phys., **122**, 1 (1989) 125-170.