

Accelerating self-modulated nonlinear waves in weakly and strongly magnetized relativistic plasmas

Felipe A. Asenjo^{1,*}

¹*Facultad de Ingeniería y Ciencias, Universidad Adolfo Ibáñez, Santiago 7491169, Chile.*

(Dated: September 13, 2023)

It is known that a nonlinear Schrödinger equation describes the self-modulation of a large amplitude circularly polarized wave in relativistic electron-positron plasmas in the weakly and strongly magnetized limits. Here, we show that such equation can be written as a modified second Painlevé equation, producing accelerated propagating wave solutions for those nonlinear plasmas. This solution even allows the plasma wave to reverse its direction of propagation. The acceleration parameter depends on the plasma magnetization. This accelerating solution is different to the usual soliton solution propagating at constant speed.

PACS numbers:

Keywords:

I. INTRODUCTION

One of the nonlinear effects present in relativistic hot magnetized electron-positron plasma is the self-modulation of circularly polarized electromagnetic waves or Alfvén waves. The large-amplitude of the electromagnetic wave, and a background magnetic field, can modify the relativistic motion of particles in a significant way. Thus, the self-modulation depends on the magnetization of the plasma. This nonlinear process has been thoroughly studied for the weakly magnetized plasma and strongly magnetized plasma cases in Refs. [1] and [2], respectively.

Interestingly, the self-modulation gives origin to soliton plasma wave solutions, propagating at constant speed [1, 2]. However, there exist other kind of accelerating plasma wave solution that deserved to be explored for relativistic plasmas. This work is devoted to show that there are nonlinear plasma waves that accelerate due to the self-modulation of the magnetized plasma system.

Relativistic plasma wave modes, presenting accelerating behavior, have been recently introduced in Refs. [3–5]. For the case of self-modulation of a circularly polarized electromagnetic or Alfvén wave in a weakly or strongly magnetized relativistic electron-positron plasma with finite temperature, the nonlinear Schrödinger equation that model such effect has been found to be [1, 2]

$$i\frac{\partial a}{\partial t} + P\frac{\partial^2 a}{\partial z^2} + Q|a|^2 a = 0, \quad (1)$$

where $a = a(t, z)$ is the time and space dependent complex modulational amplitude of the circularly polarized electromagnetic wave, under the approximation of a slowly time-varying modulation [1, 2]. Here, $P = c^2/(2\omega)$, where c is the speed of light, and ω is the frequency of the wave. Differently, Q depend on the limit

case of magnetization of the plasma. In the weakly magnetized case we have that [1]

$$Q = \frac{3\lambda\omega_p^2\Omega_c^2}{\omega^3 f^5}, \quad (2)$$

where $\lambda = e^2/m^2c^4$ (with the electron charge e and mass m), ω_p is the plasma frequency of the electron-positron plasma, Ω_c is the cyclotron frequency, and f is the thermodynamic function relating the plasma density enthalpy per unit of mass and unit of number density (being function of temperature). On the other hand, for the strongly magnetized plasma case we find that [2]

$$Q = \frac{f\lambda\omega_p^2\omega^3}{4\Omega_c^4}. \quad (3)$$

It is clear that is the background magnetic field, through the cyclotron frequency, the physical quantity that induces the nonlinear behavior of the waves.

II. NON-ACCELERATING SOLITON

Eq. (1) is usually solved in terms of a soliton amplitude propagating with constant speed v . This solution can be found by requiring that the amplitude of the electromagnetic wave has the form $|a(t, z)| = |a(z - vt)|$. Straightforwardly, it is found that such solution has the form of a soliton [1, 2]

$$a(t, z) = \operatorname{sech}\left(\sqrt{\frac{Q}{2P}}(z - vt)\right) \exp\left(i\frac{v}{2P}z - i\left(\frac{v^2}{4P} - \frac{Q}{2}\right)t\right). \quad (4)$$

III. ACCELERATING SOLUTION

However, Eq. (1) can be also solved for an accelerating wave, i.e, we look for solution with amplitude in the form

*Electronic address: felipe.asenjo@uai.cl

$|a(t, z)| = |a(z + vt - \beta t^2/2)|$, where where v and β play the role of an initial velocity and the acceleration of the propagation.

Similar to the case of previous section, this kind of solution can be studied by assuming the form of the plasma wavepacket as

$$a(t, z) = f(\xi) \exp(i\eta(t, z)), \quad (5)$$

where f is a function to be determined through Eq. (1), and depending on the accelerated argument

$$\xi = \sqrt{\frac{Q}{2P}} \left(z + vt - \frac{\beta}{2} t^2 \right). \quad (6)$$

Notice that we have assumed, in analogy with the constant velocity soliton solution, the same factor $\sqrt{Q/2P}$ for the argument. Also, the phase function $\eta(t, z)$ must be determined.

Using (5) in Eq. (1) we can determine that when the acceleration is

$$\beta = Q\sqrt{\frac{QP}{2}}, \quad (7)$$

and the phase is

$$\eta(t, z) = \frac{Q}{2} \sqrt{\frac{Q}{2P}} \left(tz - \frac{v}{Q} \sqrt{\frac{2}{QP}} z - \frac{v^2}{Q^2} \sqrt{\frac{Q}{2P}} t + vt^2 - \frac{Q}{3} \sqrt{\frac{QP}{2}} t^3 \right), \quad (8)$$

then, the function f fulfills a modified second Painlevé equation [6]

$$\frac{d^2 f}{d\xi^2} - \xi f + 2f^3 = 0. \quad (9)$$

Solutions of this equation can be generally studied numerically, but they are no longer solitons. For the current case, they describe an accelerated propagation of a electromagnetic plasma wave train, with non-constant amplitude, and with acceleration $\beta = \sqrt{Q^3 P}/2$ along its direction of propagation. The acceleration of this nonlinear plasma wave depends on the magnetization of the plasma through Q , given in Eqs (2) and (3). We emphasize that this accelerated behavior for the plasma is only possible in the weakly and strongly magnetized limits.

In order to show the accelerating behavior of this nonlinear plasma solution, we display in Fig. 1 the density plot for the numerical solution of Eq. (9) for $f(\xi = 0) = 1$, $d_\xi f(\xi) = 0$, and with $v = 0$. This plot shows the magnitude of function f in the $t - z$ space, in terms of normalized time $t' = Qt/\sqrt{2}$, and normalized distance $z' = \sqrt{Q/2P}z$. The solution shows curved (parabolic) trajectories for any part of the wave (maxima or minima). This can be explicitly seen through the red dashed lines, that are used as examples. Those lines correspond




FIG. 1: Density plot for $f(\xi)$, with $f(\xi = 0) = 1$ and $d_\xi f(\xi) = 0$, in terms of time t' and distance z' , for $v = 0$. Red dashed lines corresponds to parabolic trajectories $z' - t'^2/2 = \xi_0$, with $\xi_0 = -10, -4, 0, 5, 10$.

to $\xi = z' - t'^2/2 = \xi_0$, for $\xi_0 = -10, -4, 0, 5, 10$. All the red dashed parabolic curves coincide with the dynamics of the plasma wave. Therefore, we conclude that the whole plasma wave propagates with acceleration β given by (7).

In Fig. 2 we show what occur when the initial velocity v is considered. For the normalized case $v' = v/\sqrt{PQ} = 2$, we display the numerical behavior of f in a density plot in terms of the normalized variables t' and z' . For this case, because its accelerating nature, the whole wavepacket change its initial direction of propagation. This occur for any part of the wave, as it is shown with the parabolic curved in $t - z$ space (in red dashed lines), for $\xi = z' + v't' - t'^2/2 = \xi_0$, with $\xi_0 = -4, 0, 5, 10, 15$. The acceleration, therefore, allow to this nonlinear plasma solution to reverse its propagation direction, in a time equal to

$$t = \frac{v}{\beta}. \quad (10)$$

Furthermore, although a numerical study of the whole solution is possible, analytical properties of the solution for f can be found in the case when $\xi \rightarrow -\infty$. In this limit, the solution of the modified second Painlevé equation (9) behaves as [6]

$$f(\xi) \approx \kappa \text{Ai}(\xi), \quad (11)$$

where Ai is the Airy function, and κ is an arbitrary constant. As the accelerating propagating properties of this electromagnetic plasma wave are present for any value




FIG. 2: Density plot for $f(\xi)$, with $f(\xi = 0) = 1$ and $d_\xi f(\xi) = 0$, in terms of time t' , distance z' , and normalized velocity $v' = 2$. Red dashed lines corresponds to parabolic trajectories $z' + v't' - t'^2/2 = \xi_0$, with $\xi_0 = -4, 0, 5, 10, 15$.

of ξ , this solution pertains to the same family of other accelerating Airy solutions already found in optics and plasmas [3–5, 7–18].

IV. FINAL REMARK

We have presented a new nonlinear plasma solution with accelerating properties. As the initial velocity of the argument (6) is arbitrary, a whole set of new kind of different propagation can be obtained. This is achieved in a relativistic plasma regime, depending on how magnetized the plasma is.

Finally, the Painlevé equation has been explored in different realms of plasma physics [19–23]. Therefore, this work contributes to show that the Painlevé equation is also a straightforward consequence of accelerating solutions in relativistic nonlinear plasmas.

Acknowledgments

FAA thanks to FONDECYT grant No. 1230094 that partially supported this work.

-
- [1] F. A. Asenjo, F. A. Borotto, A. C.-L. Chian, V. Muñoz and J. A. Valdivia, *Phys. Rev. E* **85**, 046406 (2012).
 - [2] R. A. López, F. A. Asenjo, V. Muñoz, A. C.-L. Chian and J. A. Valdivia, *Phys. Rev. E* **88**, 023105 (2013).
 - [3] H. Li, X. Li and J. Wang, *J. Plasma Phys.* **82**, 905820103 (2016).
 - [4] A. E. Minovich *et al.*, *Laser Photon. Rev.* **8**, 221 (2014).
 - [5] M. A. Winkler, C. Vásquez-Wilson and F. A. Asenjo, *Eur. Phys. J. D* **77**, 97 (2023).
 - [6] P. A. Clarkson, *Journal of Computational and Applied Mathematics* **153**, 127 (2003).
 - [7] A. Mahalov and S. K. Suslov, *Phys. Lett. A*, **377**, 33 (2012).
 - [8] N. Wiersma, N. Marsal, M. Sciamanna and D. Wolfersberger, *Sci. Rep.* **6**, 35078 (2016).
 - [9] Y. Jiang, K. Huang, and X. Lu, *Opt. Express* **20**, 18579 (2012).
 - [10] D. Abdollahpour *et al.*, *Phys. Rev. Lett.* **105**, 253901 (2010).
 - [11] T. Bouchet *et al.* *Sci Rep* **12**, 9064 (2022).
 - [12] A. Chong *et al.*, *Nature Photon* **4**, 103 (2010).
 - [13] I. Kaminer *et al.*, *Phys. Rev. Lett.* **108**, 163901 (2012).
 - [14] P. Panagiotopoulos *et al.*, *Nat Commun* **4**, 2622 (2013).
 - [15] H. Esat Kondakci and A. F. Abouraddy, *Phys. Rev. Lett.* **120**, 163901 (2018).
 - [16] N. K. Efremidis *et al.*, *Optica* **6**, 686 (2019).
 - [17] S. Chávez-Cerda *et al.*, *Opt. Exp.* **19**, 16448 (2011).
 - [18] J. Baumgartl, M. Mazilu and K. Dholakia, *Nature Photon* **2**, 675 (2008).
 - [19] C. Rogers and P. A. Clarkson, *Jour. Nonlinear Math. Phys.* **25**, 247 (2018).
 - [20] S. Kumar *et al.*, *Phys. Scr.* **97** 035201 (2022).
 - [21] A. H. Khater, D. K. Callebaut, A. B. Shamardan and R. S. Ibrahim, *Phys. Plasmas* **4**, 3910 (1997).
 - [22] A. H. Khater, D. K. Callebaut and R. S. Ibrahim, *Phys. Plasmas* **5**, 395 (1998).
 - [23] R. S. Ibrahim, *IMA Journal of Applied Mathematics* **68**, 523 (2003).