Minimum Width for Deep, Narrow MLP: A Diffeomorphism Approach

Geonho Hwang

^aCenter for AI and Natural Sciences, Korea Institute for Advanced Study, 85, Hoegi-ro, Dongdaemun-gu, 02455, Seoul, Republic of Korea

Abstract

Recently, there has been a growing focus on determining the minimum width requirements for achieving the universal approximation property in deep, narrow Multi-Layer Perceptrons (MLPs). Among these challenges, one particularly challenging task is approximating a continuous function under the uniform norm, as indicated by the significant disparity between its lower and upper bounds. To address this problem, we propose a framework that simplifies finding the minimum width for deep, narrow MLPs into determining a purely geometrical function denoted as $w(d_x, d_y)$. This function relies solely on the input and output dimensions, represented as d_x and d_y , respectively. Two key steps support this framework. First, we demonstrate that deep, narrow MLPs, when provided with a small additional width, can approximate a C^2 -diffeomorphism. Subsequently, using this result, we prove that $w(d_x, d_y)$ equates to the optimal minimum width required for deep, narrow MLPs to achieve universality. By employing the aforementioned framework and the Whitney embedding theorem, we provide an upper bound for the minimum width, given by $\max(2d_x+1, d_y) + \alpha(\sigma)$, where $0 \leq \alpha(\sigma) \leq 2$ represents a constant depending on the activation function. Furthermore, we provide a lower bound of 4 for the minimum width in cases where the input and output dimensions are both equal to two.

Keywords: Universal Approximation Theorem, Deep Narrow Network, Multilayer Perceptron, Invertible Neural Network, Whitney Embedding Theorem

1. Introduction

The universal approximation property (UAP) refers to the capability of neural networks to approximate a wide range of functions. As this property forms the foundation for the efficacy of neural networks, it has garnered significant interest within the research community.

Initial research focused mainly on two-layered Multilayer Perceptrons (MLPs). Cybenko (1989) demonstrated that two-layered MLPs with sigmoidal activation functions possess the UAP for approximating continuous functions. Later, Leshno et al. (1993) expanded the scope of activation functions to more general ones. In addition to two-layered MLPs, extensive investigation has been conducted into the UAP of *deep*, *narrow MLPs*. These MLPs have a constrained width and an arbitrary number of layers. Given the common use of MLPs with relatively modest widths and more than two layers in practical scenarios, the UAP of deep, narrow MLPs has attracted significant interest.

In this regard, a series of studies have been undertaken to determine the *minimum width*, which is the necessary and sufficient width for the UAP. The minimum width depends on factors such as the input dimension d_x , the output dimension d_y , the activation function, and the type of norm employed. For instance, Lu et al. (2017) demonstrated that deep, narrow MLPs with ReLU activation functions possess the UAP, leading to further research that narrowed down the minimum width range. Hanin and Sellke (2017) extended the study to encompass arbitrary output dimensions d_y . Johnson (2018) showed that a width of d_x is insufficient to achieve the UAP in continuous function spaces, while Kidger and Lyons (2020) proved that a dimension of $d_x + d_y + 2$ is sufficient. On the other hand, Park et al. (2020) presented the optimal minimum width for deep, narrow MLPs with ReLU activation functions in L_p space. Furthermore, Cai (2022) explored the lower bound of the minimum width for arbitrary activation functions.

In this paper, we concentrate on the universal approximation of continuous functions under the uniform norm. The previous results concerning uniform approximation are organized in Table 1. So far, research on the minimum width for approximations under the uniform norm using continuous activation functions has suggested that the minimum width lies between $\max(d_x + 1, d_y)$ and $d_x + d_y$. Recently, Li et al. (2023) claimed that the upper bound could be reduced to $\max(d_x + 1, d_y) + \mathbf{1}_{d_x+1=d_y}$. On the other hand, Kim et al. (2023) proved that the lower bound equals or exceeds $d_y + 1$ if d_y is less than or equal to $2d_x$. This leads to the contradiction $d_y + 1 \le w_{min} \le d_y$ for $d_x + 2 \le d_y \le 2d_x$. Therefore, there should be a more rigorous proof of the minimum width for the uniform approximation of continuous functions.

In this context, we provide rigorous upper and lower bounds for the minimum width required for deep, narrow MLPs to possess the UAP. It is substantiated by proving that the minimum width for deep, narrow MLPs with Leaky-ReLU activation function is equal to a geometrical function denoted as $w(d_x, d_y)$. $w(d_x, d_y)$ is the required dimension of diffeomorphisms for approximating arbitrary continuous functions with d_x -dimensional input and d_y dimensional output. This is built upon the concept of the UAP of invertible neural networks. Specifically, we employ the result of Teshima et al. (2020), which demonstrated that approximating arbitrary C^2 -diffeomorphisms is equivalent to approximating arbitrary single-coordinate transformations. We prove that deep, narrow MLPs are capable of approximating single-coordinate transformations, thereby confirming their capability to approximate C^2 -diffeomorphisms. Using the above statement, we provide some upper and lower bounds. By leveraging classical results from topological geometry, we establish that any continuous function can be approximated by MLPs with width $\max(2d_x +$ $1, d_{y}$). Moreover, we provide the non-trivial lower bound 4 for the case of input and output dimensions two, which provides a necessity of the framework.

Our contributions are as follows:

- We prove that deep, narrow MLPs of width d with Leaky-ReLU activation function can approximate any C^2 -diffeomorphisms on \mathbb{R}^d . For more general activation functions, we prove that deep, narrow MLPs of width d + 1 and d + 2 with ReLU and general activation function, respectively, can approximate any C^2 -diffeomorphisms on \mathbb{R}^d .
- We suggest the purely topological indicator $w(d_x, d_y)$, which is equal to the optimal minimum width for the UAP of deep, narrow MLP with Leaky-ReLU activation function.
- Building on the above results, we prove that deep, narrow MLP with width $\max(2d_x + 1, d_y) + \alpha(\sigma)$ can approximate any continuous function in $C(\mathbb{R}^{d_x}, \mathbb{R}^{d_y})$ on a compact domain, where $0 \leq \alpha(\sigma) \leq 2$ is the constant depending on the activation function.
- We prove that width 4 is the optimal minimum width that deep, narrow MLP to approximate arbitrary continuous function from a compact set $K \subset \mathbb{R}^2$ to \mathbb{R}^2 .

1.1. Organization

The remainder of the paper is structured as follows: In Section 2, we provide the notations and definitions that will be used. Section 3 addresses the key theorems in this paper. Subsection 3.1 formulated the core problem. Subsection 3.2 explores the theorems and their corresponding proofs related to approximating a diffeomorphism using deep, narrow MLPs. Subsection 3.3 introduces a geometrical invariant that establishes a necessary and sufficient condition for the universal approximation of continuous functions. Subsection 3.4 presents the proof of the universal approximation theorem for deep, narrow MLPs with a width of $\max(2d_x+1, d_y)$ and offers an alternative proof for the result by Kidger and Lyons (2020). Subsection 3.5 establishes a minimum width lower bound of 4 for the specific case when $d_x = d_y = 2$. Section 4 concludes the study.

2. Notation and Definition

In this section, we introduce notations and definitions used throughout this paper.

- \mathbb{R} represents the set of real numbers.
- \mathbb{R}_+ denotes the set of positive real numbers.
- \mathbb{N} is the set of natural numbers, and $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$.
- For $a, b \in \mathbb{R}$, [a, b] and (a, b) represent the closed and open intervals from a to b, respectively.
- $M_{n,m}$ is the set of $n \times m$ real matrices.
- $GL(n) \subset M_{n,n}$ is the set of invertible matrices.
- Aff_{*n,m*} and IAff_{*n*} are sets of affine transformations from \mathbb{R}^n to \mathbb{R}^m and invertible affine transformations from \mathbb{R}^n to \mathbb{R}^n , respectively.
- For a *d*-dimensional vector $x \in \mathbb{R}^d$, x_i will denote the *i*-th component of x; that is, $x = (x_1, x_2, \ldots, x_d)$. And, $x_{i:j}$ will represent the (j i + 1)-dimensional vector $(x_i, x_{i+1}, \ldots, x_j)$.

Reference	Domain	Activation σ	Upper / lower bounds
Hanin and Sellke (2017)	$C(K, \mathbb{R}^{d_y})$	ReLU	$d_x + 1 \le w_{min} \le d_x + d_y$
Johnson (2018)	$C(K,\mathbb{R})$	uniformly conti. [†]	$w_{min} \ge d_x + 1$
Kidger and Lyons	$C(K, \mathbb{R}^{d_y})$	conti. nonpoly [‡]	$w_{min} \le d_x + d_y + 1$
(2020)	$C(K, \mathbb{R}^{d_y})$	nonaffine poly	$w_{min} \le d_x + d_y + 2$
Park et al. (2020)	$C([0,1],\mathbb{R}^2)$	ReLU	$w_{min} = 3 > \max\{d_x + 1, d_y\}$
	$C(K, \mathbb{R}^{d_y})$	ReLU+STEP	$w_{min} = \max\{d_x + 1, d_y\}$
Cai (2022)	$C(K, \mathbb{R}^{d_y})$	Arbitrary	$w_{\min} \ge \max(d_x, d_y)$
	$C(K, \mathbb{R}^{d_y})$	ReLU+FLOOR	$w_{\min} = \max(d_x, d_y, 2)$
	$C(K, \mathbb{R}^{d_y})$	UOE+FLOOR	$w_{\min} = \max(d_x, d_y)$
	$C([0,1],\mathbb{R}^{d_y})$	UOE	$w_{\min} = d_y$
Kim et al. (2023)	$C(K, \mathbb{R}^{d_y})$	uniformly conti. [†]	$w_{\min} \ge d_y + 1_{d_x < d_y \le 2d_x}$
Ours	$C(K, \mathbb{R}^{d_y})$	Leaky-ReLU	$w_{\min} \le \max(2d_x + 1, d_y)$
	$C(K, \mathbb{R}^{d_y})$	ReLU	$w_{\min} \le \max(2d_x + 1, d_y) + 1$
	$C(K, \mathbb{R}^{d_y})$	conti. nonpoly [‡]	$w_{\min} \le \max(2d_x + 1, d_y) + 2$
	$C([0,1]^2,\mathbb{R}^2)$	ReLU	$w_{\min} = 4$
	$C([0,1]^2,\mathbb{R}^2)$	Leaky-ReLU	$w_{\min} = 4$
	$C([0,1]^2,\mathbb{R}^2)$	uniformly conti. [†]	$w_{\min} \ge 4$

Table 1: A summary of known results on minimum width for universal approximation of continuous functions. *K* denotes a compact domain, and "Conti." is short for continuous.

[†] requires that σ is uniformly approximated by a sequence of one-to-one functions.

[†] requires that σ is continuously differentiable at at least one point (say z), with $\sigma'(z) \neq 0$.

2.1. Compact Approximation

C(X,Y) represents the set of continuous functions from X to Y. For a function $f \in C(X,Y)$ and a set $X' \subset X$, $f|_{X'}$ denotes a restriction of the function to the domain X'. For a set of functions $\mathcal{A} \subset C(X,Y)$, $\mathcal{A}|_{X'}$ is defined as $\{f|_{X'}|f \in \mathcal{A}\}$. We are concerned with the uniform approximation of a continuous function on a compact set, defined as follows:

Definition 1. For two function spaces, $\mathcal{A}, \mathcal{B} \subset C(\mathbb{R}^n, \mathbb{R}^m)$, we say that \mathcal{A} compactly approximates \mathcal{B} if for any $f \in \mathcal{B}$, a compact set $K \subset \mathbb{R}^n$, and $\epsilon > 0$, there exists $g \in \mathcal{A}$ such that

$$||f - g||_{\infty,K} := \sup_{x \in K} ||f(x) - g(x)||_{\infty} < \epsilon.$$
(1)

This is denoted as $\mathcal{A} \succ \mathcal{B}$ or $\mathcal{B} \prec \mathcal{A}$.

The compact approximation relation is transitive: if $\mathcal{A} \succ \mathcal{B}$, and $\mathcal{B} \succ \mathcal{C}$, then, $\mathcal{A} \succ \mathcal{C}$.

Proof Consider an arbitrary $f \in C$ and a compact set $K \subset \mathbb{R}^m$. Because $\mathcal{B} \succ C$, there exists $g \in \mathcal{B}$ such that $\|f - g\|_{\infty,K} < \frac{\epsilon}{2}$. Because $\mathcal{A} \succ \mathcal{B}$, there exists $h \in \mathcal{A}$ such that $\|g - h\|_{\infty,K} < \frac{\epsilon}{2}$. Then, $\|f - h\|_{\infty,K} < \|f - g\|_{\infty,K} + \|g - h\|_{\infty,K} < \epsilon$, and the relation $\mathcal{A} \succ C$ holds.

We also use the notation $f \prec \mathcal{A}$ to indicate that $\{f\} \prec \mathcal{A}$. For a set of functions $\mathcal{A} \subset C(X,Y)$, $\overline{\mathcal{A}}$ is the closure with respect to the uniform norm.

2.2. Activation Function

We follow the commonly used condition for activation functions as proposed by Kidger and Lyons (2020).

Condition 1. An activation function σ is a C^1 -function near $\alpha \in \mathbb{R}$, with $\sigma'(\alpha) \neq 0$.

We define several activation functions that satisfy Condition 1.

• ReLU: ReLU(x) :=
$$\begin{cases} x & \text{if } x \ge 0\\ 0 & \text{if } x < 0 \end{cases}$$

• Leaky-ReLU :
$$LR_{\beta}(x) := \begin{cases} x & \text{if } x \ge 0\\ \beta x & \text{if } x < 0 \end{cases}$$

Activation functions applied to vectors function as componentwise operators. For $x \in \mathbb{R}^d$,

$$\sigma(x) := (\sigma(x_1), \dots, \sigma_{x_d}). \tag{2}$$

2.3. Deep, Narrow MLP

A set of MLPs, denoted as $\mathcal{N}_{d_0,d_1,\ldots,d_N}^{\sigma}$, is defined as follows:

$$\mathcal{N}_{d_0,d_1,\dots,d_N}^{\sigma} := \left\{ f : \mathbb{R}^{d_0} \to \mathbb{R}^{d_N} \middle| W_i \in \operatorname{Aff}_{d_{i-1},d_i}, f(x) = W_N \circ \sigma \circ \dots \circ \sigma \circ W_1 \right\}.$$
(3)

For Leaky-ReLU, an additional parameter β can vary for each layer, resulting in the set $\mathcal{N}_{d_0,d_1,\ldots,d_N}^{LR}$:

$$\mathcal{N}_{d_0,d_1,\dots,d_N}^{\mathrm{LR}} := \left\{ W_N \circ \mathrm{LR}_{\beta_{N-1}} \circ \dots \circ \mathrm{LR}_{\beta_1} \circ W_1 : \frac{\mathbb{R}^{d_0}}{\rightarrow \mathbb{R}^{d_N}} \middle| W_i \in \mathrm{Aff}_{d_{i-1},d_i}, \beta_i \in \mathbb{R}_+, \right\}$$
(4)

We can define a set of deep, narrow MLPs with input dimension d_x , output dimension d_y , and at most n intermediate dimensions as follows:

$$\mathcal{N}^{\sigma}_{d_x,d_y,n} := \bigcup_{N \in \mathbb{N}_0} \bigcup_{1 \le d_1, d_2, \dots, d_N \le n} \mathcal{N}^{\sigma}_{d_x, d_1, d_2, \dots, d_N, d_y}.$$
 (5)

For natural numbers $n \geq m \in \mathbb{N}$, we define the natural projection $p_{n,m}$: $\mathbb{R}^n \to \mathbb{R}^m$ and the inclusion $q_{m,n}$ as follows:

$$p_{n,m}: (x_1, \dots, x_n) \mapsto (x_1, \dots, x_m), \tag{6}$$

and

$$q_{m,n}: (x_1, \dots, x_m) \mapsto (x_1, \dots, x_m, 0, \dots, 0).$$
 (7)

We can check that for any $f \in \mathcal{N}_{d_r,d_n,n}^{\sigma}$, f can be decomposed as:

$$f = p_{n,d_y} \circ g \circ q_{d_x,n},\tag{8}$$

where $g \in \mathcal{N}_{n,n,n}^{\sigma}$.

2.4. Subsets of Diffeomorphisms

In this section, we define several subsets of the set of diffeomorphisms.

Definition 2 (Invertible Neural Networks). For any natural number d, let \mathcal{G} be a subset of invertible functions from \mathbb{R}^d to \mathbb{R}^d . Then, $\text{INN}_{\mathcal{G}}$ is defined as:

$$INN_{\mathcal{G}} := \{ W_1 \circ g_1 \circ \dots \circ W_n \circ g_n \circ W_{n+1} | n \in \mathbb{N}, g_i \in \mathcal{G}, W_i \in IAff_d \}$$
(9)

Note that the approximation capability of $INN_{\mathcal{G}}$ remains unchanged even if $IAff_d$, in the definition, is replaced with $Aff_{d,d}$.

Definition 3 (Diffeomorphism: $\mathcal{D}^r(U)$). Let $U \subset \mathbb{R}^d$ be an open subset, and let r be a non-negative integer or infinity. $\mathcal{D}^r(U)$ is the set of C^r diffeomorphisms from U to \mathbb{R}^d .

Definition 4 (Compactly supported diffeomorphism: $\operatorname{Diff}_{c}^{r}(\mathbb{R}^{d})$). A diffeomorphism $f : \mathbb{R}^{d} \to \mathbb{R}^{d}$ is compactly supported if there exists a compact subset $K \subset \mathbb{R}^{d}$ such that for any $x \notin K$, f(x) = x. $\operatorname{Diff}_{c}^{r}(\mathbb{R}^{d})$ is the set of all compactly supported C^{r} -diffeomorphisms from \mathbb{R}^{d} to \mathbb{R}^{d} . **Definition 5 (Single-coordinate transformations:** $S_c^r(\mathbb{R}^d)$). $S_c^r(\mathbb{R}^d)$ is the set of all compactly supported C^r -diffeomorphisms defined as follows:

$$\mathcal{S}_{c}^{r}(\mathbb{R}^{d}) := \left\{ \tau \in \operatorname{Diff}_{c}^{r}(\mathbb{R}^{d}) \, \middle| \, \tau(x) = (x_{1}, \dots, x_{d-1}, \tau_{d}(x)), \tau_{d} \in C(\mathbb{R}^{d}, \mathbb{R}) \right\}.$$
(10)

Definition 6 (Single-coordinate affine coupling flows). ACF_d is the set of all single-coordinate affine coupling flows defined as follows:

$$ACF_d := \left\{ \left(x_1, \dots, x_{d-1}, exp(s(x_{1:d-1})x_d + t(x_{1:d-1}))) \middle| s, t \in C(\mathbb{R}^{d-1}, \mathbb{R}) \right\},$$
(11)

3. Main Theorem

3.1. Problem Formulation

Our primary objective is to determine the minimum width $w_{\min} \in \mathbb{N}$ such that for any compact set $K \subset \mathbb{R}^n$, a continuous function $f \in C(K, \mathbb{R}^m)$ can be uniformly approximated by $\mathcal{N}_{n,m,w_{\min}}^{\sigma}$. In other words, we want to find the value $w_{\min}(n, m, \sigma)$ such that

$$w_{min}(n,m,\sigma) := \min\left\{l \in \mathbb{N} \left| C(\mathbb{R}^n,\mathbb{R}^m) \prec \mathcal{N}_{n,m,l}^{\sigma} \right\}.$$
 (12)

3.2. Approximating Diffeomorphisms

In this subsection, we initially establish the capability of deep, narrow MLPs to approximate diffeomorphisms.

Theorem 7. Let σ be a continuous function that satisfies Condition 1. Then, for a natural number $d \in \mathbb{N}$, the set $\mathcal{N}_{d,d,d+\alpha(\sigma)}^{\sigma}$ compactly approximates $\mathcal{D}^2(\mathbb{R}^d)$, where

$$\alpha(\sigma) = \begin{cases} 0 & \text{if } \sigma = Leaky-ReLU \\ 1 & \text{if } \sigma = ReLU \\ 2 & \text{if } \sigma = otherwise \end{cases}$$
(13)

In other words, we have the relation

$$\mathcal{D}^2(\mathbb{R}^d) \prec \mathcal{N}^{\sigma}_{d,d,d+\alpha(\sigma)}.$$
(14)

To prove the theorem, we introduce a lemma that suggests we can focus on approximating $\mathcal{S}_{c}^{\infty}(\mathbb{R}^{d})$ to achieve the approximation of diffeomorphisms. **Lemma 8.** The following relation holds:

$$INN_{\mathcal{S}^{\infty}_{c}(\mathbb{R}^{d})} \succ \mathcal{D}^{2}(\mathbb{R}^{d}).$$
(15)

Proof This result is a direct consequence of Theorem 1(B) of Teshima et al. (2020). Since $\mathcal{S}_{c}^{\infty}(\mathbb{R}^{d})$ is locally bounded, which is due to its continuity and invertible, it satisfies the conditions of Theorem 1. Given that $\text{INN}_{\mathcal{S}_{c}^{\infty}(\mathbb{R}^{d})} \succ \mathcal{S}_{c}^{\infty}(\mathbb{R}^{d})$, we can conclude that $\text{INN}_{\mathcal{S}_{c}^{\infty}(\mathbb{R}^{d})} \succ \mathcal{D}^{2}(\mathbb{R}^{d})$.

Proof [Proof of Theorem 7] According to Lemma 8, it suffices to prove that the set of neural networks can serve as an approximator for \mathcal{S}_c^{∞} : $\mathcal{N}_{d,d,d+\alpha(\sigma)}^{\sigma} \succ \mathcal{S}_c^{\infty}(\mathbb{R}^d)$.

For σ = Leaky-ReLU, we need to prove that $\mathcal{N}_{d,d,d}^{\sigma} \succ \mathcal{S}_{c}^{\infty}(\mathbb{R}^{d})$. We can accomplish this by employing Lemma 9.

In the case of σ = ReLU, by Theorem 1 in Hanin and Sellke (2017), for $f(x) = (x_1, \ldots, x_d, \tau(x))$, we have $f \prec \mathcal{N}^{\sigma}_{d,d+1,d+1}$. Therefore, $(x_1, \ldots, x_{d-1}, \tau(x)) \in \mathcal{N}^{\sigma}_{d,d,d+1}$, implying that $\mathcal{S}^{\infty}_{c}(\mathbb{R}^d) \prec \mathcal{N}^{\sigma}_{d,d,d+1}$.

For other continuous activation functions σ , Proposition 4.2 of Kidger and Lyons (2020) demonstrates that for $f(x) = (x_1, \ldots, x_d, \tau(x))$, we have $f \prec \mathcal{N}^{\sigma}_{d,d+1,d+2}$. Consequently, $(x_1, \ldots, x_{d-1}, \tau(x)) \in \mathcal{N}^{\sigma}_{d,d,d+2}$, concluding that $\mathcal{S}^{\infty}_{c}(\mathbb{R}^d) \prec \mathcal{N}^{\sigma}_{d,d,d+2}$.

It is important to note that although the original statements in Theorem 1 of Hanin and Sellke (2017) and Proposition 4.2 of Kidger and Lyons (2020) do not explicitly state that the approximated function has the form $(x_1, \ldots, x_d, \tau(x))$, their proofs implicitly make use of this form.

Now, the remaining task is to prove the following lemma for the Leaky-ReLU case.

Lemma 9 (Single-Coordinate Transformations to Leaky-ReLU). The following relation holds:

$$\mathcal{N}_{d,d,d}^{LR} \succ \mathcal{S}_c^{\infty}(\mathbb{R}^d).$$
(16)

The proof of this lemma involves a series of lemmas and corollaries that gradually extend the scope of functions that can be approximated using Leaky-ReLU.

The following lemma implies that any increasing function can be approximated by composing Leaky-ReLUs and affine transformations. Lemma 10 (Increasing Functions to Leaky-ReLU). Define the sets as follows:

$$U_0 := \{ ax + b | a \in \mathbb{R}_+, b \in \mathbb{R} \}, \qquad (17)$$

$$U_{n+1} := \left\{ aLR_{\beta}(f) + b | a, \beta \in \mathbb{R}_{+}, b \in \mathbb{R}, f \in U_{n} \right\},$$
(18)

$$U := \bigcup_{n=0}^{\infty} U_n.$$
(19)

Then, for any continuous, increasing activation function $\sigma : \mathbb{R} \to \mathbb{R}$,

$$\sigma \prec U. \tag{20}$$

The proof of Lemma 10 is provided in Appendix Appendix A.1. The lemma directly implies the subsequent corollary: deep, narrow MLPs with Leaky-ReLU activation function can approximate a deep, narrow MLP with an increasing activation function and the same width.

Corollary 11 (Generalization of Activation). For any continuous, increasing activation function σ , the following relation holds:

$$\mathcal{N}_{d,d,d}^{\sigma} \prec \mathcal{N}_{d,d,d}^{LR}.$$
 (21)

Using the above corollary, we can prove that any ACF can be approximated by Leaky-ReLU deep, narrow MLPs.

Lemma 12 (ACF to Leaky-ReLU). The following relation holds:

$$INN_{ACF_d} \prec \mathcal{N}_{d.d.d.}^{LR}$$
 (22)

Proof of Lemma 12 is provided in Appendix Appendix A.2. Next, we establish a technical lemma. This lemma serves as the multidimensional counterpart of Lemma 10. For a multidimensional function from \mathbb{R}^d to \mathbb{R} increasing with a coordinate x_d , we can freely change the value when x_d is large, while the value remains unaffected when x_d is small.

Lemma 13. Consider a compact set $K = [0, 1]^d \subset \mathbb{R}^d$, two distinct real values $\alpha_1 < \alpha_2$, and a single-coordinate transformation $F = (x_1, \ldots, x_{d-1}, f(x)) \in S_c^r$. The function f(x) satisfies the following relation:

$$f(x) \le 0 \quad \text{if } x_d < \alpha_1, \quad \text{and } f(x) = 0 \quad \text{if } x_d = \alpha_1. \tag{23}$$

Assuming that $F \in \overline{\mathcal{N}_{d,d,d}^{LR}}_{K}$, for a continuous function $b : \mathbb{R}^{d-1} \to \mathbb{R}$ such that $b(x_{1:d-1}) > 0$ for all $x \in K$, there exists a single-coordinate transformation $G = (x_1, \ldots, x_{d-1}, g(x)) \in \overline{\mathcal{N}_{d,d,d}^{LR}}_{K}$ that satisfies the following relation:

$$g(x) := \begin{cases} f(x) & \text{if } x_d \le \alpha_1 \\ f(x)b(x_{1:d-1}) & \text{if } x_d = \alpha_2 \end{cases}.$$
 (24)

The proof of Lemma 13 is provided in Appendix Appendix A.3. With the help of this lemma, we can prove Lemma 9.

Proof [Proof of Lemma 9] Consider an arbitrary single-coordinate transformation $F(x) = (x_1, \ldots, x_{d-1}, \tau(x_1, \ldots, x_d))$ and a compact set $K \subset \mathbb{R}^d$. Without loss of generality, we assume that $K = [0, 1]^d$. In the case where Kis not $[0, 1]^d$, we can rescale the domain of the function to fit within $[0, 1]^d$ and continuously extend the domain to $[0, 1]^d$. Additionally, assume that τ is strictly increasing with respect to x_d .

Because τ is a continuous function defined on a compact set, for an arbitrary $\epsilon > 0$, a natural number $N \in \mathbb{N}$ exists such that if $||x - x'|| < \frac{1}{N}$, then $|\tau(x) - \tau(x')| < \epsilon$. Now, define $u_i : \mathbb{R}^{d-1} \to \mathbb{R}$ as follows:

$$u_i(x_{1:d-1}) := F\left(x_{1:d-1}, \frac{i}{N}\right).$$
 (25)

If there exists a single-coordinate transformation $G = (x_1, \ldots, x_{d-1}, g(x_{1:d})) \prec \mathcal{N}_{d,d,d}^{\mathrm{LR}}$ such that $u_i(x_{1:d-1}) = g\left(x_{1:d-1}, \frac{i}{N}\right)$ for $x \in K$, then $F \prec \mathcal{N}_{d,d,d}^{\mathrm{LR}}$. We will demonstrate the existence of a sequence $\{G_n\}_{n=1}^{\infty} \subset \overline{\mathcal{N}}_{d,d,d}^{\mathrm{LR}}|_K$ such that $G_n(x_{1:d-1}, \frac{i}{N}) = u_i(x_{1:d-1})$ for $1 \leq i \leq n$ through mathematical induction.

By Lemma 13, there exists a single-coordinate transformation $G_0 = (x_1, \ldots, x_{d-1}, g_0(x))$ such that $g_0(x_{1:d-1}, 0) = u_0(x_{1:d-1})$. Assume that the induction hypothesis holds for some $n = n_0$, meaning that there exists a single-coordinate transformation $G_{n_0} = (x_1, \ldots, x_{d-1}, g_{n_0}(x))$ such that $g_{n_0}(x_{1:d-1}, \frac{i}{N}) = u_i(x_{1:d-1})$ for $1 \leq i \leq n_0$. Then, by Lemma 12, we can construct $G'_{n_0} := (x_1, \ldots, x_{d-1}, g_{n_0}(x) - u_{n_0}(x_{1:d-1})) \in \overline{\mathcal{N}_{d,d,d}^{\mathrm{LR}}}_{K}$. Notably, G'_{n_0} satisfies the assumptions of Lemma 13 with $\alpha_1 = \frac{n_0}{N}$ and $\alpha_2 = \frac{n_0+1}{N}$. By applying Lemma 13 with $b(x_{1:d-1}) = \frac{u_{n_0+1}(x_{1:d-1})-u_{n_0}(x_{1:d-1})}{g_{n_0}(x_{1:d-1}, \frac{n_0+1}{N})-u_{n_0}(x_{1:d-1})}$, we obtain a single-coordinate transformation $G''(n_0) = (x_1, \ldots, x_{d-1}, g''_{n_0}(x))$ such that $g''_{n_0}(x_{1:d-1}, \frac{i}{N}) = u_i(x_{1:d-1}) - u_{n_0}(x_{1:d-1})$ for $i \leq n_0 + 1$. Finally, by Lemma 12, we can get $G_{n_0+1} := (x_1, \ldots, x_{d-1}, g''_{n_0}(x) + u_{n_0}(x_{1:d-1})) \in \overline{\mathcal{N}_{d,d,d}^{\mathrm{LR}}}_{K}$. As

a result, the induction hypothesis is satisfied, and this completes the proof. \blacksquare

3.3. Diffeormorphism to Continuous Function

In this subsection, we aim to prove that any continuous function can be approximated by composing linear transformations and diffeomorphisms and to determine the required width w(n,m) for approximation. Moreover, we will prove that the network-independently defined value w(n,m) equals the minimum width of deep, narrow, Leaky-ReLU MLPs.

Let $\operatorname{Emb}(X, Y)$ be the set of smooth embeddings from X to Y. Let $\operatorname{Emb}_{p.l.}(X, Y)$ be the set of smooth embeddings from X to Y. For natural numbers $d_1 \geq d_2$, let $p_{d_1,d_2} : \mathbb{R}^{d_1} \to \mathbb{R}^{d_2}$ be a projection to the first d_2 coordinates. Define w(n,m) as

$$w(n,m) := \min\left\{l \in \mathbb{N}_0 \left| p_{l,m}\left(\overline{\operatorname{Emb}([0,1]^n, \mathbb{R}^l)}\right) = C([0,1]^n, \mathbb{R}^m)\right\}.$$
 (26)

Intuitively, w(n, m) is the least width required to approximate an arbitrary continuous function with diffeomorphism.

Remark 14. We remark that interval [0,1] can be replaced with interval [a,b] for a < b. And $\operatorname{Emb}([0,1]^n, \mathbb{R}^l)$ can be replaced with any dense subset of $\overline{\operatorname{Emb}([0,1]^n, \mathbb{R}^l)}$, such as $\operatorname{Emb}_{p.l.}([0,1]^n, \mathbb{R}^l)$, the set of piecewise linear embedding from $[0,1]^n$ to \mathbb{R}^l (Munkres, 1960).

We will prove that w(n,m) has a similar value to $w(n,m,\sigma)$ and the same value to w(n,m,Leaky-ReLU). The next lemma demonstrates that any smooth embedding can be represented by composition of inclusion and smooth diffeomorphism.

Lemma 15 (Theorem C of Palais (1960)). Consider natural numbers nand m where $n \leq m$, and an embedding $f : K = [0,1]^n \to \mathbb{R}^m$. Then, there exists a smooth diffeomorphism $F : \mathbb{R}^m \to \mathbb{R}^m$ such that the following equation holds:

$$F \circ q_{n,m} = f. \tag{27}$$

Then, the following theorem is satisfied.

Theorem 16. Let σ be a continuous function satisfying Condition 1. Then, $\mathcal{N}_{n,m,w(n,m)+\alpha(\sigma)}^{\sigma}$ compactly approximates $C(\mathbb{R}^n, \mathbb{R}^m)$, where

$$\alpha(\sigma) = \begin{cases} 0 & \text{if } \sigma = Leaky-ReLU \\ 1 & \text{if } \sigma = ReLU \\ 2 & \text{if } \sigma = otherwise \end{cases}$$
(28)

In other words,

$$C(\mathbb{R}^n, \mathbb{R}^m) \prec \mathcal{N}^{\sigma}_{n,m,w(n,m)+\alpha(\sigma)}.$$
(29)

Proof Without loss of generality, assume that $K = [0, 1]^n$. In other cases, we can continuously extend the function to a cube containing K and rescale. By the definition of w(n, m), for arbitrary $f \in C([0, 1]^n, \mathbb{R}^m)$ and $\epsilon > 0$, there exists an embedding $g \in \text{Emb}([0, 1]^n, \mathbb{R}^{w(n,m)})$ such that

$$\|f - p_{w(n,m),n} \circ g\|_{\infty,[0,1]^n} < \epsilon.$$
(30)

Because $w(n,m) \ge n$, by Lemma 15, for $q_{n,w(n,m)} : (x_1, \ldots, x_n) \mapsto (x_1, \ldots, x_n, 0, \ldots, 0)$, there exists a smooth diffeomorphism G such that $g = G \circ q_{n,w(n,m)}$. By Theorem 7, there exists an $H \in \mathcal{N}^{\sigma}_{w(n,m),w(n,m),w(n,m)+\alpha(\sigma)}$ such that

$$\|G - H\|_{\infty, K \times [0,1]^{w(n,m)-n}} < \epsilon.$$

$$(31)$$

Then,

$$\|p_{w(n,m),n} \circ H \circ q_{n,w(n,m)} - p_{w(n,m),n} \circ G \circ q_{n,w(n,m)}\|_{\infty,[0,1]^n} < \epsilon.$$
(32)

Therefore,

$$\|f - p_{w(n,m),m} \circ H \circ q_{n,w(n,m)}\|_{\infty,K} < 2\epsilon.$$
(33)

 $p_{w(n,m),m} \circ H \circ q_{n,w(n,m)} \in \mathcal{N}^{\sigma}_{w(n,m),w(n,m),w(n,m)+\alpha(\sigma)}$, and it complete the proof.

Furthermore, we can give the lower bound of the minimum width for the universal approximation.

Proposition 17. Let σ be an increasing, continuous activation function. For $n, m \in \mathbb{N}, \mathcal{N}_{n,m,w(n,m)-1}^{\sigma}$ does not compactly approximate $C(\mathbb{R}^n, \mathbb{R}^m)$. In other words, the following relation holds:

$$C(\mathbb{R}^n, \mathbb{R}^m) \not\prec \mathcal{N}^{\sigma}_{n,m,w(n,m)-1}.$$
(34)

Proof For an increasing continuous activation function σ , there exist smooth, strictly increasing activation functions σ_n such that uniformly converge to σ . Therefore, $\mathcal{N}_{d,d,d}^{\sigma} \prec \bigcup_{n \in \mathbb{N}} \mathcal{N}_{d,d,d}^{\sigma_n} \prec \mathcal{D}^{\infty}(\mathbb{R}^d)$, and it is sufficient to consider only smooth, strictly increasing activation function σ .

For $f \in \mathcal{N}_{n,m,w(n,m)-1}^{\sigma}$, f can be decomposed as

$$f = p_{w(n,m)-1,m} \circ g \circ q_{n,w(n,m)-1},$$
(35)

where $g \in \mathcal{N}_{w(n,m)-1,w(n,m)-1,w(n,m)-1}^{\sigma}$. Because $\mathcal{N}_{w(n,m)-1,w(n,m)-1,w(n,m)-1}^{\sigma} \prec \mathcal{D}^{\infty}(\mathbb{R}^{w(n,m)-1}), g \circ q_{n,w(n,m)-1}|_{[0,1]^n} \in \overline{\mathrm{Emb}([0,1]^n, \mathbb{R}^{w(n,m)-1})}$. Therefore,

$$f|_{[0,1]^n} \in p_{w(n,m)-1,m}\left(\overline{\operatorname{Emb}([0,1]^n, \mathbb{R}^{w(n,m)-1})}\right),$$
 (36)

and because $f \in \mathcal{N}^{\sigma}_{n,m,w(n,m)-1}$ is arbitrary, we have

$$\mathcal{N}_{n,m,w(n,m)-1}^{\sigma}\Big|_{[0,1]^n} \subset p_{w(n,m)-1,m}\left(\overline{\mathrm{Emb}([0,1]^n,\mathbb{R}^{w(n,m)-1})}\right).$$
(37)

Because w(n,m) - 1 < w(n,m), by the definition of w(n,m),

$$p_{w(n,m)-1,m}\left(\overline{\operatorname{Emb}([0,1]^n,\mathbb{R}^{w(n,m)-1})}\right) \not\supseteq C([0,1]^n,\mathbb{R}^m),\tag{38}$$

and

$$\mathcal{N}^{\sigma}_{n,m,w(n,m)-1} \not\supseteq C([0,1]^n, \mathbb{R}^m).$$
(39)

Thus, we can conclude that $C(\mathbb{R}^n, \mathbb{R}^m) \not\prec \mathcal{N}^{\sigma}_{n,m,w(n,m)-1}$.

By combining Theorem 16 and Proposition 17 together, we can conclude that the minimum width $w_{\min}(n, m, \text{Leaky-ReLU})$ is equal to w(n, m) for Leaky-ReLU and can get tight inequality for general increasing activation functions.

Corollary 18. The following equation holds:

$$w_{\min}(n, m, Leaky-ReLU) = w(n, m)$$
(40)

For a general increasing activation function σ , which satisfies Condition 1, the following inequality holds:

$$w(n,m) \le w_{\min}(n,m,\sigma) \le w(n,m) + \alpha(\sigma), \tag{41}$$

where

$$\alpha(\sigma) = \begin{cases} 1 & \text{if } \sigma = ReLU \\ 2 & \text{if } \sigma = otherwise \end{cases}$$
(42)

3.4. Some Observation about Upper bound of w(n,m)

In the previous subsection, we demonstrated that the minimum width of the deep, narrow MLP is fundamentally correlated with w(n,m). In this subsection, we will present the sufficient condition for the w(n,m) to be equal to m. The following lemma demonstrates that a continuous function can be approximated by a smooth embedding when the output dimension is larger than twice the input dimension:

Lemma 19. Consider natural numbers n and m where m > 2n. Let $f : K = [0,1]^n \subset \mathbb{R}^n \to \mathbb{R}^m$ be a continuous function. Then, for $\epsilon \in \mathbb{R}_+$, there exists a smooth embedding $g : K \to \mathbb{R}^m$ such that

$$\|f - g\|_{\infty,K} < \epsilon. \tag{43}$$

Proof Consider a connected, open subset U of \mathbb{R}^n such that $K \subset U \subset \mathbb{R}^n$. Since K is compact, there exists a continuous extension f_0 of f such that

$$f_0|_K = f. \tag{44}$$

As U is a manifold, the assumptions of Theorem 3.17 and 3.18 of Persson (2014) are satisfied. Therefore, there exists an injective immersion g such that

$$\|f - g\|_{\infty, U} < \epsilon. \tag{45}$$

Consequently, the restriction $g|_K$ defined on the compact set K becomes a smooth embedding.

Theorem 20. Let σ be a continuous function that satisfies Condition 1. Then, for any natural numbers $n, m \in \mathbb{N}$, the set $\mathcal{N}_{n,m,\max(2n+1,m)+\alpha(\sigma)}^{\sigma}$ compactly approximates $C(\mathbb{R}^n, \mathbb{R}^m)$, where

$$\alpha(\sigma) = \begin{cases} 0 & \text{if } \sigma = Leaky-ReLU \\ 1 & \text{if } \sigma = ReLU \\ 2 & \text{if } \sigma = otherwise \end{cases}$$
(46)

In other words, we have the relation

$$C(\mathbb{R}^n, \mathbb{R}^m) \prec \mathcal{N}^{\sigma}_{n,m,\max(2n+1,m)+\alpha(\sigma)}.$$
(47)

Proof Lemma 19 implies that $w(n,m) \leq \max(2n+1,m)$. By Theorem 16, we can immediately get the conclusion.

Remark 21. As previously mentioned in the introduction, Kim et al. (2023) demonstrated that the minimum width $w_{\min}(d_x, d_y, \sigma)$ satisfies the relation $w_{\min} \ge d_y + \mathbf{1}_{d_x < d_y \le 2d_x}$ for an increasing activation function. It indicates that when the output dimension d_y is twice the input dimension d_x , and the activation function is Leaky-ReLU, $w_{\min}(d_x, 2d_x, \sigma)$ is equal to or larger than $d_y + 1$. In the same configuration, according to Theorem 20, we can get the following relation: $w_{\min} \le 2d_x + 1 = d_y + 1$. By combining these two results, we arrive at the optimal minimum width $w_{\min} = d_y + 1 = 2d_x + 1$.

Besides the above relation, there is an obvious upper bound for w(n, m):

$$w(n,m) \le n+m,\tag{48}$$

for all $n, m \in \mathbb{N}$. It reproofs the result of Hanin and Sellke (2017) for the Leaky-ReLU case:

$$w_{min}(n, m, \text{Leaky-ReLU}) \le n + m,$$
 (49)

and slightly worse results for ReLU (Hanin and Sellke, 2017) and other general activation functions Kidger and Lyons (2020):

$$w_{\min}(n, m, \text{Leaky-ReLU}) \le n + m + \alpha(\sigma),$$
 (50)

where $\alpha(\sigma) = 1$ for $\sigma = \text{ReLU}$, and $\alpha(\sigma) = 2$ for other activation functions.

3.5. Lower Bound of w(n,m)

In this subsection, we provide a nontrivial example of minimum width using the concept of w(n, m). In particular, we will prove that w(2, 2) = 4using some algebraic topological techniques.

We use the following lemma, which implies that the homology of the level set of a function is robust to the perturbation.

Lemma 22 (Theorem 2 of Bendich et al. (2010)). Let \mathbb{X} be a compact topological space. For a continuous function $f : \mathbb{X} \to \mathbb{R}$ and $f_a : \mathbb{X} \to \mathbb{R}$ defined as $f_a(x) := |f(x) - a|$, define \mathbb{X}_r as follows:

$$X_r(f_a) = f_a^{-1}[0, r]$$
(51)

For $h : \mathbb{X} \to \mathbb{R}$, such that $||f - h||_{\infty,\mathbb{X}} < r$, $h^{-1}(a)$ is included in $\mathbb{X}_r(f_a)$:

$$h^{-1}(a) \hookrightarrow \mathbb{X}_r(f_a),$$
 (52)

and the inclusion induces the homomorphism of the homology:

$$\mathbf{j}_h : \mathbf{H}_n\left(h^{-1}(a)\right) \to \mathbf{H}_n\left(\mathbb{X}_r\left(f_a\right)\right)$$
(53)

In addition, as $f^{-1}(a-r)$ and $f^{-1}(a+r)$ are also included in $\mathbb{X}_r(f_a)$, the inclusion

$$\iota^{0}: f^{-1}(a+r) \hookrightarrow \mathbb{X}_{r}(f_{a}), \qquad (54)$$

and

$$\iota^{1}: f^{-1}(a-r) \hookrightarrow \mathbb{X}_{r}(f_{a}), \qquad (55)$$

induce the homomorphisms ι^0_* and ι^1_* of the homology:

$$\iota^{0}_{*}: \mathrm{H}_{n}\left(f^{-1}(a+r)\right) \to \mathrm{H}_{n}\left(\mathbb{X}_{r}\left(f_{a}\right)\right),$$
(56)

and

$$u_*^1: \mathrm{H}_n\left(f^{-1}(a-r)\right) \to \mathrm{H}_n\left(\mathbb{X}_r\left(f_a\right)\right).$$
(57)

Define $B_{0,r}$ and $B_{1,r}$ as the images of two homomorphisms: $B_{0,r} := \iota^0_* (H_n (f^{-1}(a+r)))$ and $B_{1,r} := \iota^1_* (H_n (f^{-1}(a-r)))$. Define $U_n(r)$ as

$$U_n(r) = \bigcap_{\|h-f\|_{\infty,\mathbb{X}} \le r} \operatorname{im}(j_h).$$
(58)

Then, the following equation holds:

$$U_n(r) = B_{0,r} \cap B_{1,r}.$$
 (59)

We employ the well-known theorem as a lemma.

Lemma 23 (Hurewicz Theorem (Theorem 2A.1 of Hatcher (2000))). By regarding loops as singular 1-cycles, we obtain a homomorphism h: $\pi_1(X, x_0) \rightarrow H_1(X)$. If X is path-connected, then h is surjective and has kernel the commutator subgroup of $\pi_1(X)$, so h induces an isomorphism from the abelianization of $\pi_1(X)$ onto $H_1(X)$. **Definition 24 (Winding Number).** For a closed curve $c : [0,1] \rightarrow \mathbb{R}^2 - O$, consider c as the element of the fundamental group:

$$[c] \in \pi_1(\mathbb{R}^2 - O, x_0) = \mathbb{Z},\tag{60}$$

where the fundamental group $\pi_1(\mathbb{R}^2 - O, x_0)$ is generated by the curve $\omega_1 = (\cos(2\pi\theta), \sin(2\pi\theta))$. Then, a winding number of c is the natural number [c] as an element of $\pi_1(\mathbb{R}^2 - O, x_0) = \mathbb{Z}$.

Lemma 25. For any closed curve $c: S^1 \to \{(x, y) \in \mathbb{R}^2 | 1 < x^2 + y^2 < 2\}$ in the annulus with a winding number larger than 1, c is not injective.

Proof Assume that c is an injective curve. By the Jordan Curve Theorem (See Proposition 2B.1 of Hatcher (2000) for details), an injective curve bounds a region homeomorphic to the disk. Therefore, there exists an embedding $C: D^2 \to \mathbb{R}^2$ such that its restriction to the boundary is equal to c.

$$C|_{S^1} = c.$$
 (61)

Because $S^1 \hookrightarrow D^2 - \{O\}$ induces an isomorphism of the fundamental group, the degree should be 1 or -1. Therefore, a curve with a winding number larger than 1 is not injective.

Using the lemmas, we prove the following theorem.

Theorem 26. w(2,2) = 4.

Proof of Theorem 26 is provided in Appendix Appendix A.4.

Corollary 27.

$$w_{min}(2, 2, ReLU) = w_{min}(2, 2, Leaky-ReLU) = 4.$$
 (62)

Proof The lower bound $w_{min}(2, 2, \text{ReLU}) = w(2, 2) \ge 4$ is the exact consequence of Theorem 26 and Corollary 18. The upper bound is by Hanin and Sellke (2017).

4. Conclusion

In this paper, we have introduced a novel upper bound for the minimum width of a deep, narrow MLP necessary for achieving universal approximation within continuous function spaces. While our derived upper bound demonstrates optimality under specific conditions, we propose that the approach of approximating arbitrary functions through diffeomorphisms could lead to achieving optimality across all cases. Investigating this perspective presents an intriguing avenue for future research. Furthermore, we anticipate that analyzing the quantitative approximation capacity of general MLPs from the standpoint of diffeomorphisms may yield valuable insights.

Acknowldegement

This work was supported by a KIAS Individual Grant [AP092801] via the Center for AI and Natural Sciences at Korea Institute for Advanced Study.

Appendix A. Proofs

Appendix A.1. Proof of Lemma 10

Proof Because increasing piecewise linear functions are dense in the space of increasing continuous functions defined on a compact interval, it is sufficient to prove that for an arbitrary natural number $n \in \mathbb{N}$ and an increasing piecewise linear function f with n breakpoints, we have $f \in U_n$. We will use mathematical induction on n. For n = 0, there is nothing to prove. Now, assume that the induction hypothesis is satisfied for some $n = n_0$. Consider the case of $n = n_0+1$, where we have an increasing piecewise linear function f with n_0+1 breakpoints, denoted as $\alpha_1 < \alpha_2 < \cdots < \alpha_{n_0+1}$. The function f is affine on each of the intervals $(-\infty, \alpha_1], [\alpha_1, \alpha_2], \ldots, [\alpha_{n_0}, \alpha_{n_0+1}], [\alpha_{n_0+1}, \infty)$. Now, let f has values as follows:

$$f(x) = \begin{cases} f(\alpha_{n_0+1}) + \gamma_1(x - \alpha_{n_0+1}) & \text{if } x \in [\alpha_{n_0}, \alpha_{n_0+1}] \\ f(\alpha_{n_0+1}) + \gamma_2(x - \alpha_{n_0+1}) & \text{if } x \in [\alpha_{n_0+1}, \infty) \end{cases}.$$
 (A.1)

Consider the function f_0 defined as follows:

$$f_0(x) := \begin{cases} f(x) & \text{if } x \in (-\infty, \alpha_{n_0+1}] \\ f(\alpha_{n_0+1}) + \gamma_1(x - \alpha_{n_0+1}) & \text{if } x \in [\alpha_{n_0+1}, \infty) \end{cases}.$$
(A.2)

The function f_0 coincides with f on the interval $(-\infty, \alpha_{n_0+1}]$ and is affine on the interval $[\alpha_{n_0}, \infty)$. This means that the affine function on the interval $[\alpha_{n_0}, \alpha_{n_0+1}]$ naturally extends to the interval $[\alpha_{n_0}+1, \infty)$ with the same slope. Therefore, f_0 has n_0 breakpoints, and by the induction hypothesis, $f_0 \in U_{n_0}$. We can express f in terms of f_0 as follows:

$$f(x) = \frac{\gamma_2}{\gamma_1} \operatorname{LR}_{\frac{\gamma_1}{\gamma_2}} \left(f_0(x) - f(\alpha_{n_0+1}) \right) + f(\alpha_{n_0+1}).$$
(A.3)

Then, $f \in U_{n_0+1}$, and the induction hypothesis is satisfied for $n = n_0 + 1$. It completes the proof.

Appendix A.2. Proof of Lemma 12

Proof For $\beta \in \mathbb{R}_+$, $a, c \in \mathbb{R}$ and $b \in \mathbb{R}^{d-1}$, we define the function g as follows:

$$g: (x_1, x_2, \dots, x_d) \mapsto (x_1, x_2, \dots, x_{d-1}, x_d + a \operatorname{LR}_{\beta}(b \cdot x_{1:d-1} + c)).$$
(A.4)

Then, we will prove that $g \prec \mathcal{N}_{d,d,d}^{\text{LR}}$. If b is the zero vector, g is a constant adding function, and the statement is satisfied. If b is not the zero vector, for $b = (b_1, \ldots, b_{d-1})$, there exists an index $1 \leq i \leq d-1$ such that $b_i \neq 0$. We can define an invertible affine transformation $W \in \text{IAff}_d$ as follows:

$$W: (x_1, x_2, \dots, x_d) \mapsto (x_1, x_2, \dots, x_{i-1}, b \cdot x_{1:d-1} + c, x_{i+1}, \dots, x_d).$$
(A.5)

Because b_i is nonzero, W is invertible. Then, by applying LR_{β} to the *i*-th component, we get

$$(x_1, \dots, x_{i-1}, \operatorname{LR}_{\beta}(b \cdot x_{1:d-1} + c), x_{i+1}, \dots, x_d) \prec \mathcal{N}_{d,d,d}^{\operatorname{LR}}.$$
 (A.6)

By adding a times the i-th component to the last component, we have

$$(x_1, \dots, \operatorname{LR}_{\beta}(b \cdot x_{1:d-1} + c), \dots, x_d + a \operatorname{LR}_{\beta}(b \cdot x_{1:d-1}) + c) \prec \mathcal{N}_{d,d,d}^{\operatorname{LR}}.$$
 (A.7)

By applying $LR_{\frac{1}{q}}$ to the *i*-th component and applying W^{-1} , we get

$$(x_1, \dots, x_{d-1}, x_d + a \operatorname{LR}_{\beta}(b \cdot x_{1:d-1} + c)) \prec \mathcal{N}_{d,d,d}^{\operatorname{LR}}.$$
 (A.8)

Next, we will prove that for the function h defined as

$$h: (x_1, x_2, \dots, x_d) \mapsto (x_1, x_2, \dots, x_{d-1}, x_d + t(x_1, \dots, x_{d-1})),$$
(A.9)

 $h \prec \mathcal{N}_{d,d,d}^{\mathrm{LR}}$. By the UAP of two-layered neural networks (Leshno et al., 1993), for arbitrary $\epsilon > 0$ and a compact set $K \subset \mathbb{R}^{d-1}$, there exist $\beta \in \mathbb{R}_+$, $a_i, c_i \in \mathbb{R}$, and $b_i \in \mathbb{R}^{d-1}$ such that the following inequality holds:

$$\left\| t(x_{1:d-1}) - \sum_{i=1}^{n} a_i \mathrm{LR}_{\beta}(b_i \cdot x_{1:d-1} + c_i) \right\|_{\infty,K} < \epsilon.$$
 (A.10)

By composing Eq (A.8) for n different a_i, b_i , and c_i , we obtain

$$\left(x_1, \dots, x_{d-1}, x_d + \sum_{i=1}^n a_i \operatorname{LR}_\beta \left(b_i \cdot x_{1:d-1} + c_i\right)\right) \prec \mathcal{N}_{d,d,d}^{\operatorname{LR}}.$$
 (A.11)

Thus, $h \prec \mathcal{N}_{d,d,d}^{\mathrm{LR}}$.

Finally, by compositing operations described so far, we demonstrate that any ACF can be approximated by $\mathcal{N}_{d,d,d}^{\text{LR}}$. It is achieved by combining the following four operations:

- Apply the logarithm to the last component.
- Add $\log(s(x_1, \ldots, x_{d-1}))$ to the last component.
- Apply the exponential function to the last component.
- Add $t(x_1, \ldots, x_{d-1})$ to the last component.

This results in the following transformation:

$$(x_1, x_2, \dots, x_d) \mapsto (x_1, x_2, \dots, x_{d-1}, \exp(\log(x_d) + \log(s)) + t) = (x_1, x_2, \dots, x_{d-1}, sx_d + t) \prec \mathcal{N}_{d,d,d}^{\mathrm{LR}}.$$
 (A.12)

It completes the proof.

Appendix A.3. Proof of Lemma 13

Proof We begin by observing that it is sufficient to consider only those functions b that satisfy $b(x_{1:d-1}) \ge 1$ for all $x \in K$. Let's define β as $\beta := \inf_{x \in K} b(x_{1:d-1})$. We introduce a function $\widetilde{F}(x) := (x_1, \ldots, x_{d-1}, \widetilde{f}(x))$ defined as follows:

$$\overline{f}(x) := \beta \operatorname{LR}_{\frac{1}{\beta}}(f(x))).$$
(A.13)

If $F \in \overline{\mathcal{N}_{d,d,d}^{\mathrm{LR}}|_{K}}$, then $\widetilde{F} \in \overline{\mathcal{N}_{d,d,d}^{\mathrm{LR}}|_{K}}$. The value of $\widetilde{f}(x)$ can be calculated as:

$$\widetilde{f}(x) = \begin{cases} f(x) & \text{if } x_d \le \alpha_1\\ \beta f(x) & \text{if } x_d > \alpha_1 \end{cases}.$$
(A.14)

This ratio $\frac{g(x)}{\tilde{f}(x)} = \frac{b(x_{1:d-1})}{\beta} \ge 1$ for all $x \in K$, and \tilde{F} also satisfied all the assumptions of the lemma. Therefore, we only need to consider functions b that satisfy $b \ge 1$.

Next, we will inductively construct a sequence $\{G_i = (x_1, \ldots, x_{d-1}, g_i(x))\}_{i=1}^{\infty} \subset \overline{\mathcal{N}_{d,d,d}^{\mathrm{LR}}}_{K}$ that uniformly converges to G when $x_d = \alpha_2$. We start with $g_0(x) := f(x)$. Define $b_i : \mathbb{R}^{d-1} \to \mathbb{R}$ as

$$b_i(x_{1:d-1}) := \frac{g_i(x_{1:d-1}, \alpha_2)}{f(x_{1:d-1}, \alpha_2)},$$
(A.15)

for all $x \in K$. Define $\gamma_i \in \mathbb{R}$ as follows:

$$\gamma_i := \sup\left\{ \frac{b(x_{1:d-1})}{b_i(x_{1:d-1})} \middle| x \in K \right\}.$$
 (A.16)

Now, define two mutually exclusive sets, $L_{i,0}$ and $L_{i,1}$:

$$L_{i,0} = \left\{ x_{1:d-1} \in [0,1]^{d-1} \middle| 1 \le \frac{b(x_{1:d-1})}{b_i(x_{1:d-1})} \le \gamma_i^{\frac{1}{3}} \right\}.$$
 (A.17)

$$L_{i,1} = \left\{ x_{1:d-1} \in [0,1]^{d-1} | \gamma_i^{\frac{2}{3}} \le \frac{b(x_{1:d-1})}{b_i(x_{1:d-1})} \le \gamma_i \right\}.$$
 (A.18)

Define a distance metric D as:

$$D(x,C) := \inf_{y \in C} \|x - y\|_2, \tag{A.19}$$

and then define $\phi_i : \mathbb{R}^{d-1} \to \mathbb{R}$ as:

$$\phi_i(x) := \frac{D(x, L_{i,0})}{D(x, L_{i,0}) + D(x, L_{i,1})}.$$
(A.20)

The function ϕ_i satisfies the inequality $0 \leq \phi_i(x_{1:d-1}) \leq 1$ for all $x \in K$, has value zero on $L_{i,0}$, and has value one on $L_{i,1}$. Define $h_i : \mathbb{R}^{d-1} \to \mathbb{R}$ as follows:

$$h_i(x_{1:d-1}) := (1 - \phi_i(x_{1:d-1}))g_i(x_{1:d-1}, \alpha_2)$$
(A.21)

Then, $0 \leq h_i(x_{1:d-1}) \leq g_i(x_{1:d-1}, \alpha_2)$ for all $x \in K$, has a value of zero on $L_{i,1}$ and has a value of $g_i(\underline{x_{1:d-1}, \alpha_2})$ on $L_{i,0}$. Now, we define $g_{i+1}(x) \in \overline{\mathcal{N}_{d,d,d}^{\mathrm{LR}}|_{K}}$ as follows:

$$g_{i+1}(x) := \gamma_i^{\frac{1}{3}} \operatorname{LR}_{\gamma_i^{-\frac{1}{3}}}(g_i(x) - h_i(x_{1:d-1})) + h_i(x_{1:d-1}).$$
(A.22)

We have

$$g_{i+1}(x) \begin{cases} = g_i(x) = 0 & \text{if } x_d \le \alpha_1 \\ = g_i(x) & \text{if } x_d = \alpha_2 \text{ and } x_{1:d-1} \in L_{i,0} \\ = \gamma_i^{\frac{1}{3}} g_i(x) & \text{if } x_d = \alpha_2 \text{ and } x_{1:d-1} \in L_{i,1} \\ \le \gamma_i^{\frac{1}{3}} g_i(x) & \text{if } x_d = \alpha_2 \text{ and } x_{1:d-1} \notin L_{i,0} \cup L_{i,1} \end{cases}$$
(A.23)

Thus, for x that $x_d = \alpha_2$ and $x_{1:d-1} \in L_{i,0}$, we have

$$\frac{g(x)}{g_{i+1}(x)} = \frac{g(x)}{g_i(x)} = \frac{b(x_{1:d-1})}{b_i(x_{1:d-1})}.$$
(A.24)

As $1 \leq \frac{b(x_{1:d-1})}{b_i(x_{1:d-1})} \leq \gamma_i^{\frac{1}{3}}$ for $x_{1:d-1} \in L_{i,0}$, we get $1 \leq \frac{g(x)}{g_{i+1}(x)} \leq \gamma_i^{\frac{1}{3}}$. For x that $x_d = \alpha_2$ and $x_{1:d-1} \in L_{i,1}$,

$$\frac{g(x)}{g_{i+1}(x)} = \frac{g(x)}{\gamma_i^{\frac{1}{3}}g_i(x)} = \gamma_i^{-\frac{1}{3}}\frac{b(x_{1:d-1})}{b_i(x_{1:d-1})}.$$
(A.25)

As $\gamma_i^{\frac{2}{3}} \leq \frac{b(x_{1:d-1})}{b_i(x_{1:d-1})} \leq \gamma_i$ for $x_{1:d-1} \in L_{i,1}$, we get $1 \leq \gamma_i^{\frac{1}{3}} \leq \frac{g(x)}{g_{i+1}(x)} \leq \gamma_i^{\frac{2}{3}}$. For x that $x_d = \alpha_2$ and $x_{1:d-1} \notin L_{i,0} \cup L_{i,1}$,

$$\gamma_i^{-\frac{1}{3}} \frac{b(x_{1:d-1})}{b_i(x_{1:d-1})} = \frac{g(x)}{\gamma_i^{\frac{1}{3}} g_i(x)} \le \frac{g(x)}{g_{i+1}(x)} \le \frac{g(x)}{g_i(x)} = \frac{b(x_{1:d-1})}{b_i(x_{1:d-1})}.$$
 (A.26)

As $\gamma_i^{\frac{1}{3}} \leq \frac{b(x_{1:d-1})}{b_i(x_{1:d-1})} \leq \gamma_i^{\frac{2}{3}}$ for $x_{1:d-1} \notin L_{i,0} \cup L_{i,1}$, we get $1 \leq \frac{g(x)}{g_{i+1}(x)} \leq \gamma_i^{\frac{2}{3}}$. We obtain the following results: for all $x \in K$, where $x_d = \alpha_2$, we have $1 \leq \frac{g(x)}{g_{i+1}(x)} = \frac{b(x_{1:d-1})}{b_{i+1}(x_{1:d-1})} \leq \gamma_i^{\frac{2}{3}}$. This implies $1 \leq \gamma_{i+1} \leq \gamma_i^{\frac{2}{3}}$. Consequently, as *i* tends towards infinity, γ_i converges to one. Therefore, $\frac{g(x_{1:d-1},\alpha_2)}{g_{i+1}(x_{1:d-1},\alpha_2)}$ uniformly converges to one as *i* increases, implying that G_i converges to *G*. As a

result, there exists a function $G = (x_1, \ldots, x_{d-1}, g(x_{1:d})) \in \overline{\mathcal{N}_{d,d,d}^{\mathrm{LR}}}_{d,d,d}|_K$ such that $g(x_{1:d-1}, \alpha_2) = b(x_{1:d-1})f(x_{1:d-1}, \alpha_2).$

To check that G is a single-coordinate transformation, we can observe that:

$$g_{i+1}(x_{1:d-1}, x_d) - g_{i+1}(x_{1:d-1}, x'_d) > g_i(x_{1:d-1}, x_d) - g_i(x_{1:d-1}, x'_d), \quad (A.27)$$

for $x_d > x'_d$, which implies that $g(x_{1:d-1}, x_d) - g(x_{1:d-1}, x'_d) > g_0(x_{1:d-1}, x_d) - g_0(x_{1:d-1}, x'_d) > 0$ for all $x \in K$. Therefore, g satisfies the strictly increasing condition, and G becomes a single-coordinate transformation.

Appendix A.4. Proof of Theorem 26

Proof It is obvious that $w(2,2) \leq 4 = 2+2$. Therefore, it is sufficient to prove that $w(2,2) \geq 4$. Assume that $w(2,2) \leq 3$. Then, for an <u>arbitrary continuous</u> function f in $C([-2,2]^2, \mathbb{R}^2)$, f is contained in $p_{3,2} \circ \overline{\text{Emb}}([-2,2]^2, \mathbb{R}^3) = p_{3,2} \circ \overline{\text{Emb}}_{p,l.}([-2,2]^2, \mathbb{R}^3)$. Consider a piecewise linear map $f : [-2,2]^2 \to \mathbb{R}^2$ defined as follows:

$$f(x_1, x_2) := \begin{cases} \begin{pmatrix} 1 & -1 \\ 0 & 2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} & \text{if } 0 \le x_2 \le x_1 \\ \begin{pmatrix} 1 & -1 \\ 2 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} & \text{if } 0 \le x_1 \le x_2 \\ -f(x_2, -x_1) & \text{if } x_1 \le 0 \text{ and } 0 \le x_2 \\ f(-x_1, -x_2) & \text{if } x_2 \le 0 \end{cases}$$
(A.28)

We can check that f is the piecewise linear double-winding function. By the assumption, there exists a piecewise linear embedding $G \in \text{Emb}_{p.l.}([-2,2]^2,\mathbb{R}^3)$ such that

$$||f - p_{3,2} \circ G||_{\infty,[-2,2]^2} < \frac{1}{4}.$$
 (A.29)

Let $\Sigma : \mathbb{R}^2 \to \mathbb{R}$ be defined as

$$\Sigma: (x_1, x_2) \mapsto |x_1| + |x_2|. \tag{A.30}$$

We can observe that f conserves the level of Σ : that is, $(\Sigma \circ f)(x) = \Sigma(x)$ for all $x \in \mathbb{R}^2$. Therefore,

$$(\Sigma \circ f)^{-1}(1) = \Sigma^{-1}(1) = \{ (x_1, x_2) \in \mathbb{R}^2 | |x_1| + |x_2| = 1 \},$$
 (A.31)

which is homeomorphic to a circle S^1 . Similarly,

$$(\Sigma \circ f)^{-1} \left(\frac{1}{2}\right) = \Sigma^{-1} \left(\frac{1}{2}\right) = \left\{ \left(x_1, x_2\right) \in \mathbb{R}^2 \left| |x_1| + |x_2| = \frac{1}{2} \right\}, \quad (A.32)$$

and

$$(\Sigma \circ f)^{-1} \left(\frac{3}{2}\right) = \Sigma^{-1} \left(\frac{3}{2}\right) = \left\{ \left(x_1, x_2\right) \in \mathbb{R}^2 \left| \left|x_1\right| + \left|x_2\right| = \frac{3}{2} \right\}, \quad (A.33)$$

are homeomorphic to S^1 , and

$$(\Sigma \circ f)^{-1} \left(\left[\frac{1}{2}, \frac{3}{2} \right] \right) = \Sigma^{-1} \left(\left[\frac{1}{2}, \frac{3}{2} \right] \right) = \left\{ \left(x_1, x_2 \right) \in \mathbb{R}^2 \left| \frac{1}{2} \le |x_1| + |x_2| \le \frac{3}{2} \right\} \right.$$
(A.34)

is homeomorphic to a closed annulus $S^1 \times [0, 1]$.

Define g as $g := p_{3,2} \circ G$. Because $||f - g||_{\infty, [-2,2]^2} < \frac{1}{4}$, we have

$$|\Sigma \circ f - \Sigma \circ g| < \frac{1}{2}.$$
 (A.35)

We will apply Lemma 22 to $\Sigma \circ f$. Because $(\Sigma \circ f)^{-1} \left(\frac{1}{2}\right) = \Sigma^{-1}(\frac{1}{2})$ and $(\Sigma \circ f)^{-1} \left(\frac{3}{2}\right) = \Sigma^{-1}(\frac{3}{2})$ are a deformation retract of $(\Sigma \circ f)^{-1} \left(\left[\frac{1}{2}, \frac{3}{2}\right]\right)$, the following equation holds:

$$U_{1}\left(\frac{1}{2}\right) = H_{1}\left(B_{0,\frac{1}{2}}\right) = H_{1}\left(B_{1,\frac{1}{2}}\right) =$$
$$= H_{1}\left(\left(\Sigma \circ f\right)^{-1}\left(\left[\frac{1}{2},\frac{3}{2}\right]\right)\right) = H_{1}\left(\Sigma^{-1}\left(1\right)\right) = \mathbb{Z} \quad (A.36)$$

Thus, $U_1(\frac{1}{2}) = \mathbb{Z}$. Recall that $j_g : H_1((\Sigma \circ g)^{-1}(1)) \to U_1(\frac{1}{2})$ is surjective. Because g and Σ are piecewise linear, $(\Sigma \circ g)^{-1}(1)$ consists of finite con-

Because g and Σ are piecewise linear, $(\Sigma \circ g)^{-1}(1)$ consists of finite connected components A_1, \ldots, A_k . Then, the first homology $H_1((\Sigma \circ g)^{-1}(1))$ is decomposed as

$$H_1\left(\left(\Sigma \circ g\right)^{-1}(1)\right) = \bigoplus_{i=1}^k H_1\left(A_i\right), \qquad (A.37)$$

and j_g can be decomposed as the sum of homomorphisms $j_g^i : H_1(A_i) \to U_1(\frac{1}{2})$:

$$j_g(x) = \sum_{i=1}^k j_g^i(x_i),$$
 (A.38)

for $x = \bigoplus_{i=1}^{k} x_i$. As j_g is surjective, we can choose an index i_0 such that $j_g^{i_0}$ is a nonzero homomorphism. Set any basepoint $x_0 \in A_{i_0}$. By Lemma 23, there exists a surjective Hurewicz homomorphism $h_1 : \pi_1(A_{i_0}, x_0) \to H_1(A_{i_0})$. We know that Hurewicz homomorphism $h_2 : \pi_1((\Sigma \circ f)^{-1}([\frac{1}{2}, \frac{3}{2}]), x_0) \to$ $H_1((\Sigma \circ f)^{-1}([\frac{1}{2}, \frac{3}{2}]))$ is an isomorphism. By compositing homomorphisms as follows,

$$\pi_1 \left(A_{i_0}, x_0 \right) \xrightarrow{h_1} \operatorname{H}_1 \left(A_{i_0} \right)$$

$$\xrightarrow{j_g^{i_0}} \operatorname{H}_1 \left(\left(\Sigma \circ f \right)^{-1} \left(\left[\frac{1}{2}, \frac{3}{2} \right] \right) \right) = \operatorname{U}_1 \left(\frac{1}{2} \right) \xrightarrow{h_2^{-1}} \pi_1 \left(\left(\Sigma \circ f \right)^{-1} \left(\left[\frac{1}{2}, \frac{3}{2} \right] \right), x_0 \right),$$

$$(A.39)$$

we get the nonzero homomorphism $h_2^{-1} \circ j_g^{i_0} \circ h_1$:

$$h_2^{-1} \circ j_g^{i_0} \circ h_1 : \pi_1(A_{i_0}, x_0) \to \pi_1\left((\Sigma \circ f)^{-1}\left(\left[\frac{1}{2}, \frac{3}{2}\right]\right), x_0\right).$$
 (A.40)

We can observe that

$$h_2^{-1} \circ j_q^{i_0} \circ h_1 = \iota_*, \tag{A.41}$$

where ι_* is the homomorphism of the fundamental group induced by the inclusion $\iota: A_{i_0} \hookrightarrow (\Sigma \circ f)^{-1} \left(\left[\frac{1}{2}, \frac{3}{2} \right] \right)$. Now, we will prove that there exists a simple closed curve $\gamma: S^1 \to \mathbb{C}$

Now, we will prove that there exists a simple closed curve $\gamma : S^1 \to (\Sigma \circ g)^{-1}(1)$, which is homotopic to the cycle $\omega_1 : \theta \mapsto (\cos(2\pi\theta), \sin(2\pi\theta))$. Because g and Σ are piecewise linear, $(\Sigma \circ g)^{-1}(1)$ can be realized by simplicial complex, and we can assume that $\pi_1((\Sigma \circ g)^{-1}(1), x_0)$ is generated by curves consisting of finite segments, and all self-intersection points are breakpoints of curves. Choose $\gamma_0 \in \pi_1((\Sigma \circ g)^{-1}(1), x_0) = \pi_1(A_{i_0}, x_0)$ such that $\iota_*([\gamma_0])$ is nonzero. We will inductively construct a closed curve γ_i until γ_i has no self-intersection point. Assume that γ_i has self-intersection point $a \neq b$: $\gamma_i(a) = \gamma_i(b)$. Let γ_i^+ be defined as $\gamma_i|_{[a,b]}$ and γ_i^- be defined as $\gamma_i|_{S^1-(a,b)}$. Then, γ_i^+ and γ_i^- become two closed curves again with a strictly smaller number of segments than γ_i . Because the winding number of γ_i is equal to the sum of those of γ_i^+ and γ_i^- , at least one of γ_i^+ or γ_i^- has a nonzero winding number, and we set γ_{i+1} as the one with a nonzero winding number. Each γ_i has a strictly smaller number of segments as *i* increases and has a winding number not equal to zero. Because γ_0 has finite segments, this process stops in a finite sequence. Therefore, we can get a non-self-intersecting curve $\gamma := \gamma_n$ with a nonzero winding number. If γ has a winding number of which the absolute value is larger than one, by Lemma 25, it has a self-intersection point. Thus, γ has a winding number 1 or -1. Reverse reparametrization yields a curve with winding number one.

Because g is homotopic to f through the linear interpolation and γ is homotopic to ω_1 , the compositions of the two functions are homotopic, which implies the same winding number between $g \circ h$ and $f \circ \omega_1$. Therefore, the winding number of $g \circ \gamma : S^1 \to S^1 = \Sigma^{-1}(1)$ becomes two. Now consider G. Because G is an embedding, it is injective. Therefore, $G|_{\gamma(S^1)} : \gamma(I) \to S^1 \times \mathbb{R}$ is injective. As the image $G(\gamma(S^1))$ is compact, the image in $S^1 \times \mathbb{R}$ can be embedded in the annuls $\{(x_1, x_2) \in \mathbb{R} | 1 - \epsilon \leq |x_1| + |x_2| \leq 1 + \epsilon\}$. And the map $G \circ \gamma$ has winding number two. However, by Lemma 25, any map with winding number two is not injective, and it becomes a contradiction.

References

- Bendich, P., Edelsbrunner, H., Morozov, D., Patel, A., 2010. The robustness of level sets, in: European symposium on algorithms, Springer. pp. 1–10.
- Cai, Y., 2022. Achieve the minimum width of neural networks for universal approximation, in: The Eleventh International Conference on Learning Representations.
- Cybenko, G., 1989. Approximation by superpositions of a sigmoidal function. Mathematics of control, signals and systems 2, 303–314.
- Hanin, B., Sellke, M., 2017. Approximating continuous functions by relu nets of minimal width. arXiv preprint arXiv:1710.11278.
- Hatcher, A., 2000. Algebraic topology.
- Johnson, J., 2018. Deep, skinny neural networks are not universal approximators. arXiv preprint arXiv:1810.00393.

- Kidger, P., Lyons, T., 2020. Universal approximation with deep narrow networks, in: Conference on learning theory, PMLR. pp. 2306–2327.
- Kim, N., Min, C., Park, S., 2023. Minimum width for universal approximation using relu networks on compact domain. arXiv preprint arXiv:2309.10402
- Leshno, M., Lin, V.Y., Pinkus, A., Schocken, S., 1993. Multilayer feedforward networks with a nonpolynomial activation function can approximate any function. Neural networks 6, 861–867.
- Li, L., Duan, Y., Ji, G., Cai, Y., 2023. Minimum width of leaky-ReLU neural networks for uniform universal approximation, in: Proceedings of the 40th International Conference on Machine Learning, PMLR. pp. 19460–19470. URL: https://proceedings.mlr.press/v202/li23g.html.
- Lu, Z., Pu, H., Wang, F., Hu, Z., Wang, L., 2017. The expressive power of neural networks: A view from the width. Advances in neural information processing systems 30.
- Munkres, J., 1960. Obstructions to the smoothing of piecewise-differentiable homeomorphisms. Annals of Mathematics , 521–554.
- Palais, R.S., 1960. Extending diffeomorphisms. Proceedings of the American Mathematical Society 11, 274–277.
- Park, S., Yun, C., Lee, J., Shin, J., 2020. Minimum width for universal approximation. arXiv preprint arXiv:2006.08859.
- Persson, M., 2014. The whitney embedding theorem.
- Teshima, T., Ishikawa, I., Tojo, K., Oono, K., Ikeda, M., Sugiyama, M., 2020. Coupling-based invertible neural networks are universal diffeomorphism approximators. Advances in Neural Information Processing Systems 33, 3362–3373.