ERDŐS–KO–RADO TYPE RESULTS FOR PARTITIONS VIA SPREAD APPROXIMATIONS

ANDREY KUPAVSKII

ABSTRACT. In this paper, we address several Erdős–Ko–Rado type questions for families of partitions. Two partitions of $[n]$ are *t*-intersecting if they share at least t parts, and are partially t -intersecting if some of their parts intersect in at least t elements. The question of what is the largest family of pairwise t-intersecting partitions was studied for several classes of partitions: Peter Erdős and Székely studied partitions of $[n]$ into ℓ parts of unrestricted size; Ku and Renshaw studied unrestricted partitions of [n]; Meagher and Moura, and then Godsil and Meagher studied partitions into ℓ parts of equal size. We improve and generalize the results proved by these authors.

Meagher and Moura, following the work of Erdős and Székely, introduced the notion of partially t-intersecting partitions, and conjectured, what should be the largest partially t-intersecting family of partitions into ℓ parts of equal size k. The main result of this paper is the proof of their conjecture for all t, k , provided ℓ is sufficiently large.

All our results are applications of the spread approximation technique, introduced by Zakharov and the author. In order to use it, we need to refine some of the theorems from the original paper. As a byproduct, this makes the present paper a self-contained presentation of the spread approximation technique for t-intersecting problems.

1. INTRODUCTION

For a positive integer n, we use notation $[n] = \{1, \ldots, n\}$ and, more generally, $[a, b] =$ $\{a, a+1, \ldots, b\}$. For a set X, the notation 2^X and $\binom{X}{k}$ stand for the collection of all subsets and all k-element subsets of the set X , respectively. A *family* is a collection of subsets of X for some X .

This paper deals with intersection theorems, which constitute a large and important class of results in extremal combinatorics. A family of sets is intersecting if any two sets in the family have non-empty intersection, and is *t-intersecting* if any two sets intersect in at least t elements. In their seminal paper, Erdős, Ko and Rado $[6]$ determined the largest size of an intersecting family in $2^{[n]}$ and $\binom{[n]}{k}$ $\binom{n}{k}$ for all n, k . Later, after a series of papers by Frankl [\[8\]](#page-19-1), Wilson [\[18\]](#page-19-2), and Frankl and Füredi [\[9\]](#page-19-3), Ahlswede and Khachatrian [\[1\]](#page-19-4) confirmed Frankl's conjecture on the size of the largest *t*-intersecting family in $\binom{[n]}{k}$ $\binom{n}{k}$ for any n, k, t .

Intersecting questions were investigated for many other structures. See a great survey paper by Ellis [\[4\]](#page-19-5) on the subject. The first result in this direction was due to Deza and Frankl [\[3\]](#page-19-6), who determined the largest size of an intersecting family of permutations. We call two permutations σ, π intersecting if for some element x we have $\sigma(x) = \pi(x)$. Deza and Frankl also found the largest family of 2- and 3-intersecting permutations when n and $n-1$ is

a prime number, respectively. They also made a conjecture concerning the size of the largest t-intersecting family of permutations. The progress on this problem was rather difficult. After a series of papers, Ellis, Friedgut and Pilpel [\[5\]](#page-19-7) managed to solve it for any constant t, provided $n > n_0(t)$. Recently, the problem was resolved for any n and t that satisfy $n > Ct \log^2 t$ by Zakharov and the author [\[12\]](#page-19-8). This was later improved to $n > Ct$ by Keller, Lifshitz, Minzer and Sheinfeld [\[11\]](#page-19-9) and then to $n > (1 + \epsilon)t$ for any $\epsilon > 0$ by Kupavskii [?]. Early approaches to this question were algebraic, based on Hoffman-Delsarte type bounds and representation theory. The approach of [\[5\]](#page-19-7) combines junta approximations, coming from Boolean Analysis, with representation theory. Zakharov and the author introduced a combinatorial technique of spread approximations that is based on the breakthrough in the Erdős–Rado sunflower problem due to Alweiss, Lovett, Wu and Zhang [\[2\]](#page-19-10). The approach of [\[11\]](#page-19-9) is based on hypercontractivity, an important tool from Analysis.

Another class of EKR-type problems with algebraic flavour deals with different classes of partitions. Until now, it was approached either using the Delta-system method or algebraic tools, based on the Delsarte–Hoffman bounds for suitable graphs. In this paper, we show that the technique of spread approximations (an essentially combinatorial approach) allows to gain significant progress in these questions.

Consider two partitions $P = (P_1, \ldots, P_{\ell_1})$ and $Q = (Q_1, \ldots, Q_{\ell_2})$ of [n]. We say that P and Q t-intersect if they share at least t parts, that is, $P_{i_s} = Q_{j_s}$ for two sets of t indices $\{i_1, \ldots, i_t\}$ and $\{j_1, \ldots, j_t\}$. We say that P and Q partially t-intersect if there are parts P_i, Q_j , such that $|P_i \cap Q_j| \geq t$. A family of partitions is *(partially) t-intersecting* if any two partitions from the family (partially) t-intersect.

1.1. General partitions. Let \mathcal{B}_n be the family of all partitions of [n]. Recall that $|\mathcal{B}_n|$ is the *n*-th Bell number B_n (for general reference, see [\[19\]](#page-19-11)). A natural example of t intersecting family in \mathcal{B}_n is the family of all partitions that have $\{1\}, \{2\}, \ldots, \{t\}$ as parts. The size of this family is B_{n-t} . Ku and Renshaw [\[13\]](#page-19-12) proved the following result.

Theorem 1 (Ku and Renshaw [\[13\]](#page-19-12)). Let $n \geq n_0(t)$ and assume that $\mathcal{F} \subset \mathcal{B}_n$ is tintersecting. Then $|\mathcal{F}| \leq \mathcal{B}_{n-t}$, with equality only possible if $\mathcal F$ is a family of all partitions with t fixed singletons.

In their proof, they require $t \leq c \log n$. The theorem below gives a much better dependence between the parameters.

Theorem 2. The conclusion of Theorem [1](#page-1-0) is valid for $n \geq Ct \log^2 t$ with some absolute constant C.

1.2. **Partitions with** k **parts.** Let \mathcal{P}_n^{ℓ} be the family of all partitions of [n] into ℓ parts. Recall that $|\mathcal{P}_n^{\ell}| = \begin{cases} n \\ \ell \end{cases}$ ℓ , where the expression on the right is the Stirling number of the second kind (for general reference, see [\[20\]](#page-19-13)). Peter Erdős and Lászlo Székely [\[7\]](#page-19-14) proved the following result.

Theorem 3 (Erdős and Székely [\[7\]](#page-19-14)). Let $n \ge n_0(\ell)$ be large enough. Assume that $\mathcal{F} \subset \mathcal{P}_n^{\ell}$ is t-intersecting. Then $|\mathcal{F}| \leq \begin{cases} n-t \end{cases}$ $\ell-t$ $\Big\}.$

Here, we improve their result as follows.

Theorem 4. Let n, ℓ, t be integers such that $t \leq \ell - 2$, $n \geq 2\ell \log_2 n$ and $n \geq 48$. Assume that $\mathcal{F} \subset \mathcal{P}_n^{\ell}$ is t-intersecting. Then $|\mathcal{F}| \leq \begin{cases} n-t \\ \ell \end{cases}$ $\ell-t$ $\Big\}$. Moreover, if F is not contained in a family of all partitions with t fixed singletons, then $|\mathcal{F}| \leq \frac{1}{2}$ $\int n-t$ $\ell-t$ $\Big\} .$

Note that the question is trivial for $t = \ell - 1$, since if two k-partitions have $\ell - 1$ common parts then the last part is also common, so they coincide.

1.3. Partitions with fixed profile. Let $P = (k_1, \ldots, k_\ell)$ be a non-decreasing sequence of positive integers, called a *profile*, and put $n = \sum_{i=1}^{\ell} k_i$. Consider the family of partitions U_P of n into ℓ parts, where the *i*-th part has size k_i (partitions with profile P). A canonical *t*-intersecting family of such partitions is a family $\mathcal{A}_{P,t}^X$, defined by a *t*-tuple X of disjoint sets X_1, \ldots, X_t , where $|X_i| = k_i$. It consists of all partitions that contain each X_i as one of its parts.

A (k, ℓ) -partition is a particular type of profiled partitions, when $k_1 = \ldots = k_\ell =: k$. In other words, it is a partition of $\left[\mathcal{k}\ell\right]$ into exactly ℓ blocks, each of size k. Let $\mathcal{U}_{k,\ell}$ stand for the family of all (k, ℓ) -partitions, and put $u_{k,\ell} := |\mathcal{U}_{k,\ell}|$.

Meagher and Moura [\[14\]](#page-19-15) proved the following theorems.

Theorem 5 ([\[14\]](#page-19-15)). Fix positive integers k, ℓ . Let $\mathcal{F} \subset \mathcal{U}_{k,\ell}$ be an intersecting family of (k, ℓ) -partitions. Then $|\mathcal{F}| \leq u_{k,\ell-1}$, and the equality is only possible for canonical intersecting families.

Theorem 6 ([\[14\]](#page-19-15)). Fix positive integers k, ℓ , t. Suppose that either $(k \geq k_0(\ell, t))$ or $(k \geq$ $t+2$ and $\ell \geq \ell_0(k,t)$). Let $\mathcal{F} \subset \mathcal{U}_{k,\ell}$ be a t-intersecting family of (k,ℓ) -partitions. Then $|\mathcal{F}| \leq u_{k,\ell-t}$, and the equality is only possible for canonical t-intersecting families.

In the theorem below, we extend their results to a large class of profiled partitions, significantly improve the dependence between the parameters, and give a strong stability result.

Theorem 7. Let t, ℓ be positive integers and $P = (k_1, \ldots, k_\ell)$ be a profile. Assume that the following holds: $\ell \ge 600$; $t \le \ell/2$; $k_{t+1} \ge 2$. If $\mathcal{F} \subset \mathcal{U}_P$ is t-intersecting then $|\mathcal{F}| \le |\mathcal{A}_{P,t}^X|$. Moreover, if F is not contained in $\mathcal{A}_{P,t}^X$ for some X then $|\mathcal{A}| \leq \frac{1}{2} |\mathcal{A}_{P,t}^X|$.

We did not try to optimize constant 600 in the statement. The point is that the bound is reasonable.

1.4. Partially *t*-intersecting partitions. In the notation of the previous subsection, let $P = (k_1, \ldots, k_\ell)$ be a profile that additionally satisfies $k := k_\ell \geq t$. Put $n = \sum_i k_i$. What is the largest partially *t*-intersecting subfamily of U_P ? A natural candidate is a *canonically* partially t-intersecting family $\mathcal{C}_P^T \subset \mathcal{U}_P$, where T is a set of size t, and \mathcal{C}_P^T consists of all partitions from \mathcal{U}_P that fully contain T in one of its parts. Similarly, for the case of (k, ℓ) -partitions, a *canonically partially t-intersecting partition* is a family $\mathcal{C}_{k,\ell}^T \subset \mathcal{U}_{k,\ell}$ that consists of all (k, ℓ) -partitions with a part that contains a fixed set T, $|T| = t$. Meagher and Moura [\[14\]](#page-19-15) conjectured the following.

Conjecture 1 (Meagher and Moura, [\[14\]](#page-19-15)). Let ℓ, k, t be integers and $\mathcal{F} \subset \mathcal{U}_{k,\ell}$ be a partially t-intersecting family. Then, for a t-element $T \subset [k\ell],$

$$
|\mathcal{F}| \leq |\mathcal{C}_{k,\ell}^T|.
$$

Note that any two partitions are partially 1-intersecting, and thus $\mathcal{U}_{k,\ell}$ itself is a partially 1-intersecting family. The Meagher–Moura conjecture was proved by Godsil and Meagher [\[10\]](#page-19-16) for $k = t = 2$ and all ℓ . Meagher, Shirazi and Stevens [\[15\]](#page-19-17) proved it for $t = 2$ and any k, provided ℓ is sufficiently large. The approach in the first paper is algebraic, while the second paper uses Delta-systems.

We prove the following theorem, which is the main result of this paper.

Theorem 8. The Meagher–Moura conjecture is true for any k, t, provided ℓ is sufficiently large. Moreover, the only extremal examples have the form $\mathcal{C}_{k,\ell}^T$ for some T.

This could be generalized to a much larger class of profiled partitions.

Theorem 9. For any k, t there exists ℓ such that for any P as above every largest partially t-intersecting family of \mathcal{U}_P is of the form \mathcal{C}^T_P for some $T.$

The proof that we present for Theorem [8](#page-3-0) goes almost verbatim for Theorem [9,](#page-3-1) except for the more complicated notation and some slightly different estimates. Thus, we decided to omit it. We also note that in both settings the proofs give a stability result that, informally speaking, tells that any family that is within constant factor of the size of the extremal family must be very close to one of the extremal families.

1.5. The structure of the remainder of the paper. In order to prove the theorems, we use the spread approximation method. We need to make some improvements in the main ingredients of the method, relaxing the conditions that we impose on the families, as compared to $[12]$. For this reason, we present the proofs of the results from $[12]$, taking into account the requirements needed for this paper. The proof of the first theorem stays essentially the same, while the second theorem is simplified and has an improved bound as compared to [\[12\]](#page-19-8). This makes this paper a self-contained presentation of the version of the spread approximation technique for t -intersecting problems. We also note that, while working in the spread approximation framework gives us a huge lever, each of the problems poses its own difficulties that we need to overcome.

In the next section we give an outline of the proofs of our theorems. In Section [3,](#page-5-0) we present the parts of the spread approximation method that we use in this paper. In Section [4,](#page-9-0) we give the proofs of our main results. We note that in Section [4](#page-9-0) we prove an almost sharp lower bound on the ratio of consecutive Bell numbers, as well as the lower bound for the number of partitions with parts of sizes at least 2, which may be of some independent interest.

2. Sketches of the proofs

As we have already mentioned, the proofs of our Theorems are based on the spread approximation method, introduced by Zakharov and the author [\[12\]](#page-19-8). The steps of the proof are summarized below.

- (1) Reformulate the problem in terms of families of sets. The problems we deal with are for families of partitions, and the method of spread approximation is designed for families of sets. Thus, we need to give an adequate set interpretation to the problems. Theorems [2,](#page-1-1) [4,](#page-2-0) and [7](#page-2-1) are based on a more straightforward interpretation: each possible part in a partition is treated as a singleton, and thus partitions become sets of size at most n or exactly ℓ , depending on the question. Theorems [8](#page-3-0) and [9](#page-3-1) require a more complicated interpretation. There, each pair of elements from the original ground set becomes a singleton in the new ground set, and each partition is turned into a set of all pairs of singletons that lie in the same part.
- (2) Prove that the set family, corresponding to the ambient family of partitions, is sufficiently spread (this is a certain quasirandomness notion that is crucial for the spread approximations method and that we introduce in the next section). This is again more or less complicated for different settings. For example, in the proof of Theorem [2,](#page-1-1) in order to lower bound spreadness, we need to lower bound the ratio $B(n)/B(n-1)$.
- (3) Take the extremal intersecting family $\mathcal F$ (depending on the setting) and apply the spread approximation theorem. Get a lower-uniformity family S , which encodes a family of "partial" partitions and that covers most of our family, as well as a small remainder $\mathcal{F}' \subset \mathcal{F}$ that is not covered by S. The approximation is given by Theorem [13.](#page-6-0)
- (4) Show that this lower-uniformity family S is trivial for the extremal family: it consists of a single t-element set T (corresponding to different "partial" partitions depending on the setting). This is done using Theorem [14.](#page-7-0)
- (5) Show that, for an extremal family \mathcal{F} , the remainder \mathcal{F}' must be empty. This is more of an ad-hoc argument, which requires a significant amount of effort in some cases. In particular, for the case of $\mathcal{B}(n)$, we need to lower bound the number of partitions with parts of sizes at least 2, and for the case of partially t-intersecting families, we need to lower bound the probability that two random partitions of the type as in Theorems [8,](#page-3-0) [9,](#page-3-1) have a not so small probability to not partially t-intersect. Such bounds allow us to say that what we "lose" because of having some partitions in \mathcal{F}' cannot be compensated by what we "gain" by adding sets from \mathcal{F}' (keeping in mind that we have upper bounds on the size of \mathcal{F}').

For Theorems [4](#page-2-0) and [7,](#page-2-1) steps 3 and 5 are omitted, since these families are sufficiently spread so that we can apply Theorem [14](#page-7-0) directly to $\mathcal F$ instead of $\mathcal S$. Thanks to that, we get the strongest stability results in these cases. We note that Theorem [14](#page-7-0) alone can be seen as a strengthening of one of the important parts of the Delta-system method.

In the third and fourth step we need to do some refinement of the method, proposed by Kupavskii and Zakharov. The fourth step is simplified and improved as compared to [\[12\]](#page-19-8).

3. Spread approximations

For a set $X \subset [n]$ and a family $\mathcal{G} \subset 2^{[n]}$ we use the following standard notation:

$$
\mathcal{G}(X) := \{ A \setminus X : G \in \mathcal{G}, X \subset A \},
$$

$$
\mathcal{G}(\bar{X}) := \{ A : A \in \mathcal{G}, A \cap X = \emptyset \}.
$$

We think of $\mathcal{G}(X), \mathcal{G}(\bar{X})$ as of subfamilies of $2^{[n]\setminus X}$. Let $r > 1$ be some real number. We say that a family F is r-spread if, for any set X, $|\mathcal{F}(X)| \leq r^{-|X|}|\mathcal{F}|$. We denote by $\|\mathcal{F}\|$ the average size of a set sampled from \mathcal{F} : $\|\mathcal{F}\| := \frac{1}{|\mathcal{F}|} \sum_{S \in \mathcal{F}} |S|$. Note that $\|\mathcal{F}\|$ is at most the size of the largest set in F. We say that W is a p-random subset of $[n]$ if each element of $[n]$ is included in W with probability p and independently of others.

The following statement is a variant due to Tao [\[16\]](#page-19-18) of the breakthrough result that was proved by Alweiss, Lowett, Wu and Zhang [\[2\]](#page-19-10).

Theorem 10 ([\[2\]](#page-19-10), a sharpening due to [\[16\]](#page-19-18)). If for some $n, r \ge 1$ a family $\mathcal{F} \subset 2^{[n]}$ is r-spread and W is an $(m\delta)$ -random subset of $[n]$, then

$$
\Pr[\exists F \in \mathcal{F} \; : \; F \subset W] \ge 1 - \left(\frac{5}{\log_2(r\delta)}\right)^m ||\mathcal{F}||.
$$

Recall that an ℓ -sunflower is a collection of ℓ sets F_1, \ldots, F_ℓ such that for any $i \neq j$ we have $F_i \cap F_j = \bigcap_{i=1}^{\ell} F_i$. (In particular, ℓ pairwise disjoint sets form an ℓ -sunflower.) The set $\cap F_i$ is called the *core*. The theorem implies an important strengthening on the size of the family that guarantees the existence of an ℓ -sunflower. Namely, it implies that any family F of k-sets with

$$
|\mathcal{F}| > (C\ell \log_2(k\ell))^k
$$

contains an ℓ -sunflower, where C is an absolute constant and can be taken to be 2^{10} .

To construct spread approximations, we will need the following easy observation.

Observation 11. Given $r > 1$ and a family $\mathcal{F} \subset 2^{[n]}$, let X be an inclusion-maximal set that satisfies $|\mathcal{F}(X)| \geq r^{-|X|}|\mathcal{F}|$. Then $\mathcal{F}(X)$ is r-spread as a family in $2^{[n]\setminus X}$.

Proof. Indeed, by maximality, for any $B \supsetneq X$ of size b we have

$$
|\mathcal{F}(B)| \le r^{-b}|\mathcal{F}| \le r^{-b+|X|}|\mathcal{F}(X)|.
$$

 \Box

To relate size and spreadness, we use the following observation.

Observation 12. If for some $\alpha > 1$ and $\mathcal{F} \subset \binom{[n]}{k}$ $\binom{n}{k}$ we have $|\mathcal{F}| > \alpha^k$ then F contains an α -spread subfamily of the form $\mathcal{F}(X)$ for some set X of size strictly smaller than k.

Proof. If F is α -spread then we are done. Otherwise, take the largest X that violates the α -spreadness of F. Clearly, $|X| < k$ and $\mathcal{F}(X)$ satisfies the requirements of the observation. \square

We note that this observation together with Theorem [10](#page-5-1) implies bound [\(1\)](#page-5-2).

The next theorem allows to construct low-uniformity approximations for sufficiently spread families.

Theorem 13. Let $n, k, t \geq 1$ be some integers and $A \subset 2^{[n]}$ be a family. Consider a family $\mathcal{F} \subset \mathcal{A} \cap \binom{[n]}{$ $\binom{[n]}{\leq k}$ that is t-intersecting. Let $q, r, r_0 \geq 1$ satisfy the following: $r > 2^{12} \log_2(2k)$, $r \geq 2q$ and $\overline{r}_0 > r$. Assume that A is r_0 -spread.

Then there exists a t-intersecting family S of sets of size at most q (a spread approximation of \mathcal{F}) and a 'remainder' $\mathcal{F}' \subset \mathcal{F}$ such that

- (i) We have $\mathcal{F} \setminus \mathcal{F}' \subset \mathcal{A}[\mathcal{S}];$
- (ii) for any $B \in \mathcal{S}$ there is a family $\mathcal{F}_B \subset \mathcal{F}$ such that $\mathcal{F}_B(B)$ is r-spread;
- (iii) $|\mathcal{F}'| \le (r_0/r)^{-q-1} |\mathcal{A}|$.

The crucial difference with [\[12,](#page-19-8) Theorem 11] is that we only require r-spreadness from A, instead of (r, t) -sreadness^{[1](#page-6-1)}. This is crucial in our application to Theorem [8,](#page-3-0) because the set interpretation of $\mathcal{U}_{k,\ell}$ is an r-spread, but not an (r, t) -spread, family. The proof of this theorem is essentially the same as in [\[12\]](#page-19-8). We present it here for completeness.

Proof. The theorem is obtained using the following procedure. For $i = 1, 2, ...$ with $\mathcal{F}^1 :=$ $\mathcal F$ we do the following steps.

- (1) Find an inclusion-maximal set S_i such that $|\mathcal{F}^i(S_i)| \geq r^{-|S_i|}|\mathcal{F}^i|$;
- (2) If $|S_i| > q$ or $\mathcal{F}^i = \emptyset$ then stop. Otherwise, put $\mathcal{F}^{i+1} := \mathcal{F}^i \setminus \mathcal{F}^i[S_i].$

The family $\mathcal{F}^i(S_i)$ is r-spread by Observation [11.](#page-5-3)

Let m be the step of the procedure for F at which we stop. Put $S := \{S_1, \ldots, S_{m-1}\}.$ Clearly, $|S_i| \leq q$ for each $i \in [m-1]$. The family \mathcal{F}_B promised in (ii) is defined to be $\mathcal{F}^i[S_i]$ for $B = S_i$. Next, note that if \mathcal{F}^m is non-empty, then

$$
|\mathcal{F}^m| \le r^{|S_m|} |\mathcal{F}^m(S_m)| \le r^{|S_m|} |\mathcal{A}(S_m)| \le (r/r_0)^{|S_m|} |\mathcal{A}|,
$$

where in the last inequality we used the r_0 -spreadness of A. We put $\mathcal{F}' := \mathcal{F}^m$. Since either $|S_m| > q$ or $\mathcal{F}' = \emptyset$, we have $|\mathcal{F}'| \le (r_0/r)^{-q-1} |\mathcal{A}|$.

The only thing left to verify is the t -intersection property. Take any (not necessarily distinct) $S_i, S_j \in \mathcal{S}$ and assume that $|S_i \cap S_j| < t$. Recall that $\mathcal{G}_i := \mathcal{F}^i(S_i)$ and $\mathcal{G}_j := \mathcal{F}^j(S_j)$ are both r-spread.

$$
|\mathcal{G}_j(\bar{S}_i)| \geq |\mathcal{G}_j| - \sum_{x \in S_i \setminus S_j} |\mathcal{G}_j[\{x\}]| \geq \Big(1 - \frac{|S_i|}{r}\Big) |\mathcal{G}_j| \geq \frac{1}{2} |\mathcal{G}_j|.
$$

 $1(r, t)$ -sreadness means that any subfamily of the form $\mathcal{A}(T)$ with $|T| \leq t$ must be r-spread)

In the last inequality we used that $|S_i| \leq q$ and that $r \geq 2q$. The same is valid for $\mathcal{G}_i(\bar{S}_j)$. Note that both $\mathcal{G}'_j := \mathcal{G}_j(\bar{S}_i)$ and $\mathcal{G}'_i := \mathcal{G}_i(\bar{S}_j)$ are subfamilies of $2^{[n]\setminus (S_i \cup S_j)}$. Because of the last displayed inequality and the trivial inclusion $\mathcal{G}'_j(Y) \subset \mathcal{G}_j(Y)$, valid for any Y, both \mathcal{G}'_i and \mathcal{G}'_j are $\frac{r}{2}$ -spread, where $\frac{r}{2} > 2^{11} \log_2(2k)$. Indeed, we have

$$
2|\mathcal{G}'_j| \geq |\mathcal{G}_j| \geq r^{|X|}|\mathcal{G}_j(X)| \geq r^{|X|}|\mathcal{G}'_j(X)|.
$$

We are about to apply Theorem [10.](#page-5-1) Let us put $m = \log_2(2k)$, $\delta = (2\log_2(2k))^{-1}$, and let $r/2$ play the role of r. Note that $m\delta = \frac{1}{2}$ and $\frac{r}{2}\delta > 2^{10}$ by our choice of r. Theorem [10](#page-5-1) implies that a $\frac{1}{2}$ -random subset W of $[n] \setminus (S_i \cup S_j)$ contains a set from \mathcal{G}'_j with probability strictly bigger than

$$
1 - \left(\frac{5}{\log_2 2^{10}}\right)^{\log_2 2k} k = 1 - 2^{-\log_2 2k} k = \frac{1}{2}.
$$

Consider a random partition of $[n] \setminus (S_i \cup S_j)$ into 2 parts U_i, U_j , including each element with probability $1/2$ in each of the parts. Then both $U_{\ell}, \ell \in \{i, j\}$, are distributed as W above. Thus, the probability that there is $F_{\ell} \in \mathcal{G}'_{\ell}$ such that $F_{\ell} \subset U_{\ell}$ is strictly bigger than $\frac{1}{2}$. Using the union bound, we conclude that, with positive probability, it holds that there are such $F_{\ell}, F_{\ell} \subset U_{\ell}$, for each $\ell \in \{i, j\}$. Fix such choices of U_{ℓ} and $F_{\ell}, \ell \in \{i, j\}$. Then, on the one hand, both $F_i \cup S_i$ and $F_j \cup S_j$ belong to F and, on the other hand, $|(F_i \cup S_i) \cap (F_j \cup S_j)| = |S_i \cap S_j| < t$, a contradiction with F being t-intersecting. \square

An important second step is to show that the approximation family S is trivial for an extremal family. In [\[12,](#page-19-8) Theorem 12], the authors worked with (r, t) -spread families, which we cannot afford in the proof of Theorem [8](#page-3-0) for the family $\mathcal{U}_{k,\ell}$, and thus have to find ways around it. Let us say that a family $A \subset 2^{[n]}$ is *weakly* (r, t) -spread if there exists a set $T \subset [n]$ of size t such that for any nonnegative integer s and a set $U \subset [n]$ of size $t + s$ we have $|\mathcal{A}(U)| \leq r^{-s} |\mathcal{A}(T)|$. Informally speaking, this akin to r-spreadness for the family $\mathcal{A}(T)$, where T of size t is chosen so that $|\mathcal{A}(T)|$ is maximal.

Recall that a *t*-intersecting family S is non-trivial if $| \bigcap_{F \in \mathcal{S}} F | < t$.

Theorem 14. Let $\varepsilon \in (0,1], n,r,q,t \geq 1$ be such that $\varepsilon r \geq 24q$. Let $\mathcal{A} \subset 2^{[n]}$ be a weakly (r, t) -spread family and let $S \subset \binom{[n]}{\leq q}$ $\begin{bmatrix} [n] \ \leq q \end{bmatrix}$ be a non-trivial t-intersecting family. Then there exists a t-element set T such that $|\overline{A}[S]|\leq \varepsilon |\mathcal{A}[T]|$.

The proof again follows closely the proof of [\[12,](#page-19-8) Theorem 12], with a few changes. Again, we present it here in full for completeness. In what follows, assume that T is a set of size t that maximizes $|\mathcal{A}(T)|$. For the proof, we will need the following simple observation.

Observation 15. For any positive integers n, p, a family $A \subset 2^{[n]}$ and a t-intersecting family $S \subset \binom{[n]}{n}$ $\begin{array}{c} \Gamma^{[n]} \ \leq_P \end{array}$ there exists a t-intersecting family $\mathcal{T} \subset \bigl(\frac{[n]}{\leq_P} \bigr)$ $\{S^{[n]}_{\leq p}\}$ such that $\mathcal{A}[\mathcal{S}]\subset \mathcal{A}[\mathcal{T}]$ and for any $T \in \overline{\mathcal{T}}$ and any proper subset $X \subsetneq T$ there exists $T^{\overline{\prime}} \in \mathcal{T}$ such that $|X \cap T'| < t$.

One natural way to choose such $\mathcal T$ is to repeatedly replace sets in $\mathcal S$ by their proper subsets while preserving the t-intersecting property.

In terms of Theorem [14,](#page-7-0) let us iteratively define the following series of families.

- (1) Let \mathcal{T}_0 be a family given by Observation [15](#page-7-1) when applied to A and S with $p = q$.
- (2) For $i = 0, \ldots, q t$ we put $\mathcal{W}_i = \mathcal{T}_i \cap \binom{[n]}{q}$ $\begin{bmatrix} |n| \\ q-i \end{bmatrix}$ and let \mathcal{T}_{i+1} be the family given by Observation [15](#page-7-1) when applied to the families A (playing the role of A) and $\mathcal{T}_i \setminus \mathcal{W}_i$ playing the role of S with $p = q - i - 1$.

Remark that \mathcal{T}_i is t-intersecting for each $i = 0, \ldots, q - t$ by definition. We summarize the properties of these series of families in the following lemma.

Lemma 16. The following properties hold for each $i = 0, \ldots, q - t$.

- (i) All sets in \mathcal{T}_i have size at most $q i$.
- (ii) We have $\mathcal{A}[\mathcal{T}_{i-1}] \subset \mathcal{A}[\mathcal{T}_{i}] \cup \mathcal{A}[\mathcal{W}_{i-1}].$
- (iii) The family \mathcal{T}_i does not have a subfamily \mathcal{Y} and a set X such that $\mathcal{Y}(X)$ satisfies $|\mathcal{Y}(X)| > 1$ and is $> (q - i - t + 1)$ -spread.
- (iv) We have $|\mathcal{W}_i| \leq (6(q i))^{q i t}$.
- (v) If \mathcal{T}_i consists of a single t-element set X and this is not the case for \mathcal{T}_{i-1} then $|\mathcal{A}[\mathcal{T}_{i-1} \setminus \mathcal{W}_{i-1}]| \leq \frac{q}{r} |\mathcal{A}[T]|.$

The lemma is similar [\[12,](#page-19-8) Lemma 14], with two major differences. First, we replace the use of sunflowers by the use of spread families, which improves the bound in (iv). Second, in (v) , where we replace a concrete X with a "universal" T on the right-hand side.

Proof. (i) This easily follows by induction on i from the fact that all sets in S have size at most q and the definition of \mathcal{T}_i .

(ii) We have $\mathcal{A}[\mathcal{T}_{i-1}] = \mathcal{A}[\mathcal{T}_{i-1} \setminus \mathcal{W}_{i-1}] \cup \mathcal{A}[\mathcal{W}_{i-1}]$ and, by the definition of \mathcal{T}_i , we have $\mathcal{A}[\mathcal{T}_i] \supset \mathcal{A}[\mathcal{T}_{i-1} \setminus \mathcal{W}_{i-1}].$

(iii) Assume such Y and X exist. Assume that a set $T' \in \mathcal{T}_i$ intersects X in $t - j$ elements, $j > 0$. Then T' intersects each set in $\mathcal{Y}(X)$ in at least j elements. For any set Z disjoint with X we have $|Y(X \cup Z)| < (q - i - t + 1)^{-|Z|}|\mathcal{Y}|$. There are, however, only $\binom{T' \setminus X}{i}$ $\binom{X}{j} \leq \binom{q-i-t+j}{j}$ $j^{-(t+j)}$ subsets of $T' \setminus X$ of size j, and thus the part of $\mathcal{Y}(X)$ that intersects T' in at least j elements has size strictly smaller than

$$
\binom{q-i-t+j}{j}(q-i-t+1)^{-j}|\mathcal{Y}|\leq (q-i-t+1)^{-j}\prod_{a=1}^j\frac{q-i-t+a}{a}|\mathcal{Y}|\leq |\mathcal{Y}|.
$$

This is a contradiction, ad thus $j = 0$. But then this contradicts the minimality of \mathcal{T}_i : we could have replaced $\mathcal{T}_i(X)$ with X.

(iv) This is trivial for $i = q - t$ since \mathcal{T}_{q-t} contains at most 1 set. In what follows, we assume that $i < q - t$. Take any set $Y \in \mathcal{W}_i$. Since \mathcal{T}_i is t-intersecting, there is a t-element subset $X \subset Y$ such that $|\mathcal{W}_i| \leq {q-i \choose t}$ $\binom{-i}{t}|{\cal W}_i(X)| = \binom{q-i}{q-i-1}$ $\frac{q-i}{q-i-t}$ | $\mathcal{W}_i(X)$ |. Next, $\mathcal{W}_i(X)$ is $(q - i - t)$ -unform and does not contain a subfamily $\mathcal Y$ and a set X such that $\mathcal Y(X)$ is

 $>(q-i-t+1)$ -spread. Using Observation [12,](#page-6-2) we conclude that $|\mathcal{W}_i(X)| \leq (q-i-t+1)^{q-i-t}$.

$$
|\mathcal{W}_i| \leq {q-i \choose q-i-t} (q-i-t+1)^{q-i-t}
$$

$$
\leq \left(\frac{e(q-i)}{q-i-t}\right)^{q-i-t} (2(q-i-t))^{q-i-t}
$$

$$
\leq 6(q-i)^{q-i-t}
$$

(v) Let us assume that $\mathcal{T}_i = \{X\}$ for some t-element set X. Note that all sets in \mathcal{T}_{i-1} have size at least $t + 1$. Otherwise, if there is $X' \in \mathcal{T}_{i-1}$ of size t then X' is a proper subset of all other sets from \mathcal{T}_{i-1} , which contradicts the property of \mathcal{T}_{i-1} guaranteed by Observation [15.](#page-7-1) Thus, the sets in $\mathcal{T}' := \mathcal{T}_{i-1} \setminus \mathcal{W}_{i-1}$, if any, have size at least $t + 1$ and all contain X. Recall that, for a family F, the covering number $\tau(F)$ is the size of the smallest cover: a set Y such that $Y \cap F \neq \emptyset$ for each $F \in \mathcal{F}$. Assume that $\tau(\mathcal{T}'(X)) > q$. Each set in \mathcal{W}_{i-1} either contains X or intersects every set from $\mathcal{T}'(X)$. In the latter case, it has size at least $\tau(\mathcal{T}'(X))$, which is impossible because each set in W_{i-1} has size at most q. Thus, all sets from \mathcal{W}_{i-1} contain X, implying that all sets from T_{i-1} contain X, a contradiction. Therefore, $\tau(\mathcal{T}'(X)) \leq q$. Recall that $T, |T| = t$, is such that $|\mathcal{A}(T)|$ is maximal. If $\{x_1, \ldots, x_q\}$ is a covering of $\mathcal{T}'(X)$ then we have

$$
|\mathcal{A}[\mathcal{T}']| \leq |\mathcal{A}[X \cup \{x_1\}]| + \ldots + |\mathcal{A}[X \cup \{x_q\}]| \leq \frac{q}{r} |\mathcal{A}[T]|,
$$

where in the last inequality we used the definition of T and the weak (r, t) -spreadness. \Box

Proof of Theorem [14.](#page-7-0) Fix i as in Lemma [16](#page-8-0) (v). Note that by (i) such a choice always exists. Let T be a t-element set such that $|\mathcal{A}[T]|$ is maximal. By the weak (r, t) -spreadness, for any $j < i$ and any $W \in \mathcal{W}_j$ we have $|\mathcal{A}[W]| \leq r^{-(q-j-t)}|\mathcal{A}[T]|$. By (iv) and the union bound, we get $|\mathcal{A}[\mathcal{W}_j]| \leq r^{-(q-j-t)} (6q)^{(q-j-t)} |\mathcal{A}[T]|$. Using this and (v) we obtain

$$
|\mathcal{A}[\mathcal{S}]| \stackrel{(ii)}{\leq} |\mathcal{A}[\mathcal{T}_{i-1}]| + \sum_{j=0}^{i-1} |\mathcal{A}[\mathcal{W}_j]| \stackrel{(iv), (v)}{\leq} \left(\frac{q}{r} + \sum_{j=1}^{\infty} r^{-j} (6q)^j\right) |\mathcal{A}[T]|
$$

$$
\leq \left(\frac{\epsilon}{2} + \sum_{j=1}^{\infty} \left(\frac{\epsilon}{4}\right)^j\right) |\mathcal{A}[T]| \leq \epsilon |\mathcal{A}[T]|,
$$

where in the third inequality we used the condition on r and the bound on C_0 .

4. Proofs

4.1. General partitions. Proof of Theorem [2.](#page-1-1)

Lemma 17. For any $n \geq 2$ we have $\frac{B_{n+1}}{B_n} \geq \frac{n}{2 \log n}$ $\frac{n}{2 \log_e n}$.

Proof. We use the following remarkable explicit formula for B_n (see [\[17\]](#page-19-19)):

(2)
$$
B_n = \frac{1}{e} \sum_{s=0}^{\infty} \frac{s^n}{s!}.
$$

Let us compare the $(s - 1)$ -th and s-th terms in the summation. We have

(3)
$$
\frac{\frac{(s-1)^n}{(s-1)!}}{\frac{s^n}{s!}} = s\left(1 - \frac{1}{s}\right)^n \le s e^{-n/s} < e^{-1},
$$

provided $\frac{n}{s} \geq 1 + \log_e s$, that is, $n \geq s + s \log_e s$. Let us denote $s_0(n)$ the largest integer s that satisfies this inequality, and note that $s_0(n) \geq \frac{n}{\log n}$ $\frac{n}{\log_e n}$ for $n \ge 16$. (It is easy to verify by direct substitution, using that $e^e < 16$.)

Using (3) , we conclude that the terms in (2) grow faster than a geometric progression with base e until at least s_0 , and so, using the formula for the sum of a geometric progression, we get $\sum_{s=0}^{s_0-1} \frac{s^n}{s!} \leq \frac{1}{e-1}$ $\frac{s_0^n}{s_0!}$. We can thus bound B_n as follows:

$$
B_n \leq \frac{e}{e-1} \sum_{s=s_0}^{\infty} \frac{s^n}{s!}.
$$

(We note that, of course, with a bit more care and for relatively large n , the fraction in front should be essentially 1.) Let us now bound the ratio.

$$
\frac{B_{n+1}}{B_n} \ge \frac{\sum_{s=s_0(n)}^{\infty} \frac{s^{n+1}}{s!}}{\frac{e}{e-1} \sum_{s=s_0(n)}^{\infty} \frac{s^n}{s!}} \ge \frac{s_0(e-1)}{e} \ge \frac{n}{2 \log_e n}
$$

.

This completes the proof for $n \geq 16$. We have $B_n \geq 2B_{n-1}$ since, for each partition P of $[n-1]$, element *n* can be either made a separate part or adjoined to one of the parts in *P*. Similarly, $B_n \geq 3B_{n-1}$ for $n \geq 5$ because for each partition P but the single partition with 1 part it can be extended by n in at least 3 ways and, moreover, there are partitions that can be extended in at least 4 ways. We are left to note that $n/(2 \log_e n) \leq 2$ for $2 \leq n \leq 8$ and $n/(2\log_e n) \leq 3$ for $9 \leq n \leq 15$.

Given a partition $P = (P_1, \ldots, P_\ell) \in \mathcal{P}_n$, we denote by $\mathcal{D}_n \langle P \rangle \subset \mathcal{B}_n$ the family of all partitions from \mathcal{B}_n that do not contain any of P_i as a part. We call this family Pderangements.

Lemma 18. For any partition $P \in \mathcal{B}_n$ we have $|\mathcal{D}_n \langle P \rangle| \geq c' e^{-\frac{3}{2} \log_e^2 n} B_n$.

Following our approach with more care, the constant in front of $\log_e^2 s$ can be improved to $\frac{1}{2} + \epsilon$, provided $n \ge n_0(\epsilon)$ is large enough.

Proof. The first step, inspired by a compression operation from Ku and Renshaw [\[13\]](#page-19-12), is to show that $|\mathcal{D}_n \langle P \rangle|$ is minimized for a partition that consists of n singletons. To do so, for any given partition P that has a part P_i with $|P_i| \geq 2$, we introduce a new partition P' that is the same as P except that it replaces part P_i with two parts Q_i, Q'_i , where these are non-empty disjoint sets such that $Q_i \sqcup Q'_i = P_i$. Take any partition $U \in \mathcal{D}_n \langle P' \rangle \setminus \mathcal{D}_n \langle P \rangle$. Partitions P and P' mostly coincide, and the only part that the former has and the latter does not is P_i . Thus, U must contain P_i as a part. Define $f(U)$ to be a partition that coincides with U except it replaces P_i with Q_i, Q'_i . Then $f(U) \in \mathcal{D}_n \langle P \rangle \setminus \mathcal{D}_n \langle P' \rangle$. Moreover, f is an injection. Thus, $|\mathcal{D}_n \langle P' \rangle| \leq |\mathcal{D}_n \langle P \rangle|$.

Repeatedly applying the splitting operation, we arrive at the all-singleton partition S , for which the number of derangements is thus minimized. Following [\[13\]](#page-19-12), we denote

$$
\tilde{B_n}=|\mathcal{D}_n\langle S\rangle|.
$$

Note that $\mathcal{D}_n\langle S\rangle$ is the family of all partitions with parts of size ≥ 2 . We have the following recurrence relations:

$$
B_{n+1} = \sum_{i=0}^{n} \binom{n}{i} B_i
$$

$$
\tilde{B}_{n+1} = \sum_{i=2}^{n-1} \binom{n}{i} \tilde{B}_i.
$$

Let us prove by induction on $s \geq 2$ that we have

(4)
$$
\tilde{B}_s \geq \frac{1}{2} \prod_{i=2}^{s-1} \left(1 - \frac{2 \log_e(i+1)}{i+1} \right) \left(1 - \frac{2i+2}{3^i} \right) \cdot B_s.
$$

We have $B_{i+1} \geq 2B_i$ if $i \geq 1$, and thus $1 = B_0 = B_1$ and $B_i \geq 2^{i-1}$ for $i \geq 2$. Thus,

$$
\frac{\sum_{i=2}^s \binom{s}{i} B_i}{\sum_{i=0}^s \binom{s}{i} B_i} = 1 - \frac{s+1}{\frac{1}{2} + \frac{1}{2} \sum_{i=0}^s 2^i \binom{s}{i}} \ge 1 - \frac{2s+2}{3^s}.
$$

Returning to [\(4\)](#page-11-0), it is true for $s = 2$ because $\tilde{B}_2 = 1$ and $B_2 = 2$. We have (using induction in the first inequality below, the last displayed inequality in the third inequality below, and Lemma [17](#page-9-2) in the fourth inequality)

$$
\tilde{B}_{s+1} = \sum_{i=2}^{s-1} {s \choose i} \tilde{B}_i \ge \frac{1}{2} \sum_{i=2}^{s-1} {s \choose i} \prod_{j=2}^{i-1} \left(1 - \frac{2 \log_e(j+1)}{j+1}\right) \left(1 - \frac{2j+2}{3^j}\right) B_i
$$
\n
$$
\ge \frac{1}{2} \prod_{j=2}^{s-1} \left(1 - \frac{2 \log_e(j+1)}{j+1}\right) \left(1 - \frac{2j+2}{3^j}\right) \sum_{i=2}^{s-1} {s \choose i} B_i
$$
\n
$$
\ge \frac{1}{2} \prod_{j=2}^{s-1} \left(1 - \frac{2 \log_e(j+1)}{j+1}\right) \prod_{j=2}^{s} \left(1 - \frac{2j+2}{3^j}\right) \sum_{i=0}^{s-1} {s \choose i} B_i
$$
\n
$$
= \frac{1}{2} \prod_{j=2}^{s-1} \left(1 - \frac{2 \log_e(j+1)}{j+1}\right) \prod_{j=2}^{s} \left(1 - \frac{2j+2}{3^j}\right) (B_{s+1} - B_s)
$$
\n
$$
\ge \frac{1}{2} \prod_{j=2}^{s} \left(1 - \frac{2 \log_e(j+1)}{j+1}\right) \left(1 - \frac{2j+2}{3^j}\right) B_{s+1}.
$$

This proves [\(4\)](#page-11-0). Let us now obtain a bound on \tilde{B}_n . First, we note that $\prod_{i=2}^{s} \left(1 - \frac{2i+2}{3^i}\right)$ $\frac{i+2}{3^i}\Big)\geq$ 1 $\frac{1}{3}\left(1-\sum_{i=3}^{s}\frac{2i+2}{3^{i}}\right)$ $\frac{i+2}{3^i}\big) \geq \frac{1}{3}$ $\frac{1}{3}(1-\frac{8}{27}-2\cdot\frac{10}{81})\geq\frac{1}{9}$ $\frac{1}{9}$. Next,

$$
\prod_{i=3}^{s} \left(1 - \frac{2 \log_e i}{i} \right) \le ce^{-\sum_{i=3}^{s} \frac{3 \log_e i}{i}} \le ce^{-\frac{3}{2} \log_e^2 s}
$$

for some positive constant c. In the first inequality we used the bound $(1 - \frac{1}{r})$ $(\frac{1}{x}) = (1 +$ 1 $\frac{1}{x-1}$)⁻¹ $\geq e^{-\frac{1}{x-1}}$. Thus, we get

$$
\tilde{B}_n \ge \frac{1}{18} c e^{-\frac{3}{2} \log_e^2 n} B_n = c' e^{-\frac{3}{2} \log_e^2 n} B_n,
$$

$$
c'.
$$

for some positive constant c'

Proof of Theorem [2.](#page-1-1) We give the following set interpretation to B_n . Consider the ground set $2^{[n]}$, and let $P \in \mathcal{B}_n$ be mapped into a $\leq n$ -element set A on $2^{[n]}$, where each element of A corresponds to a part from the partition P. In what follows, we think of \mathcal{B}_n as a family of sets. Note that, using Lemma [17,](#page-9-2) for any set $X \in \binom{2^{[n]}}{s}$ $(s^{(n)}_s), s \leq n$, such that $\mathcal{B}_n(X)$ is non-empty, we get

$$
\left(\frac{|\mathcal{B}_n(X)|}{|\mathcal{B}_n|}\right)^{1/s} \ge \left(\frac{B_{n-s}}{B_n}\right)^{1/s} \ge \left(\prod_{i=n-s}^{n-1} \frac{i}{2\log_e i}\right)^{1/s} \ge \left(\frac{(n-1)!}{(2\log_e (n-1))^{n-1}}\right)^{1/(n-1)} \ge
$$

$$
\frac{n/e}{2\log_e (n-1)} \ge \frac{n}{6\log_e n},
$$

where for the last inequality we need n to be somewhat large (e.g., $n \geq 50$ is sufficent).

From the above, we get that the family \mathcal{B}_n is $r_0 = \frac{n}{6 \log_e n}$ spread. Moreover, it is weakly $\left(\frac{n}{12 \log_e n}, t\right)$ -spread for any $t \leq n/2$. Indeed, it is sufficient to consider the number of all partitions that fix t singletons (there are B_{n-t} of them) and compare that with the number of partitions that fix some $t + s$ parts, $s \ge 1$ (there are at most B_{n-t-s} of those for any choice of the parts to fix). To bound the ratio, we reuse the bounds displayed above.

We are now ready to prove the theorem. Take a *t*-intersecting family $\mathcal{F} \subset \mathcal{B}_n$. First, we apply Theorem [13](#page-6-0) with $r = r_0/2$ and $q = 2^{-10} \frac{n}{\log_2 n}$. We get a *t*-intersecting family S of sets of size at most q (corresponding to collections of q pairwise disjoint sets) and a remainder family \mathcal{F}' such that

$$
|\mathcal{F}'| \le 2^{-q-1} B_n \le n^{-q/\log_2 n} B_n \le n^{-t-4\log_e n} B_n \le n^{-4\log_e n} B_{n-t}.
$$

Here we used a bound $B_n/B_{n-1} \leq n$, which is valid since any partition of $n-1$ elements can be prolonged in at most n ways to a partition on n elements, as well as the bound $t + 4\log_e n \leq \frac{q}{\log_e n}$ $\frac{q}{\log_2 n}$, valid for our choice of parameters.

Next, we apply Theorem [14](#page-7-0) to the family S with $\epsilon = 1/2$. The family \mathcal{B}_n is weakly $(r_0/2, t)$ -spread, and we have $r_0/2 \geq 48q$, and thus the inequality on the parameters is satisfied. We conclude that either $|\mathcal{F}| \leq \frac{1}{2}B_{n-t} + |\mathcal{F}'| \leq 0.6B_{n-t}$, or S consists of one

t-element set S. Moreover, in order for $\mathcal F$ to be maximal, this set in the partition language must clearly correspond to a collection of t singletons.

Finally, let us show that $\mathcal{F}'\backslash\mathcal{B}_n[S]$ must be empty. Indeed, assume $P \in \mathcal{F}'\backslash\mathcal{B}_n[S]$. Then, by Lemma [18,](#page-10-1) the number of partitions in $\mathcal{B}_n(S)$ that are derangements with respect to P (induced on the complement of S in the partition language), is at least $c'e^{-\frac{3}{2}\log_e^2(n-t)}B_{n-t}$, which is larger than $|\mathcal{F}'|$ by the last displayed formula. We get that if \mathcal{F}' is non-empty, then $\mathcal F$ cannot be extremal. This completes the proof of the theorem.

4.2. Partitions with k parts. Proof of Theorem [4.](#page-2-0) We will need the following relation between Stirling coefficients of the second kind.

Lemma 19. For each
$$
n \ge 1 + 2\ell \log_2 n
$$
, $\ell \ge 2$ we have $\begin{Bmatrix} n \\ \ell \end{Bmatrix} \ge n^2 \begin{Bmatrix} n-1 \\ \ell-1 \end{Bmatrix}$.

Proof. Stirling numbers obey the following recurrence relation:

$$
\begin{Bmatrix} n \\ \ell \end{Bmatrix} = \begin{Bmatrix} n-1 \\ \ell-1 \end{Bmatrix} + \ell \begin{Bmatrix} n-1 \\ \ell \end{Bmatrix}.
$$

From here, we see that $\begin{cases} n \\ 0 \end{cases}$ ℓ $\} \geq \ell \left\{ \frac{n-1}{\ell} \right\}$ ℓ). Next, let us compare $\begin{cases} n-1 \\ 0 \end{cases}$ ℓ $\left\{\right. \text{and} \left\{\right. \begin{array}{c} n-1 \\ 0 \end{array}\right.$ $\ell-1$ $\big\}$. Consider a bipartite graph between ℓ -partitions and $(\ell-1)$ -partitions of $[n-1]$, where two partitions are connected by an edge if one is a refinement of the other. Let us count the degrees in this graph. The neighbors of an ℓ -partition X are the $(\ell-1)$ -partitions obtained by merging two parts of X. Thus, the degree of X is $\binom{\ell}{2}$ $\binom{\ell}{2}$. The neighbors of an $(\ell-1)$ partition Y are those that can be obtained by subdividing one of the parts of Y into nonempty parts. Assume that the parts Y_1, \ldots, Y_ℓ have sizes $k_1, \ldots, k_\ell, \sum_{i=1}^\ell k_i = n$. There are $2^{\ell_i-1}-1$ ways to subdivide Y_i . Thus, the degree of Y is $\sum_{i=1}^{\ell} (2^{k_i-1}-1) \geq \ell 2^{-1+(n-1)/\ell} - \ell$ by convexity. The last expression is at least $ln^2/2-\ell$ by our assumption on n. Concluding, the degree of any element in the ℓ -partitions part is $\geq n^2/\ell$ times smaller than the degree of any element in the $(\ell-1)$ -partitions part. Double counting the number of edges, we get that $\begin{cases} n-1 \\ 1 \end{cases}$ ℓ $\Big\}\geq n^2/\ell\Big\{\frac{n-1}{2}\Big\}$ $\ell-1$ in our assumptions. Combining it with the displayed formula, we get that $\begin{cases} n \\ 0 \end{cases}$ ℓ $\geq n^2 \begin{cases} n-1 \\ 0 \end{cases}$ $\ell-1$ <u>}</u> .

Proof of Theorem [4.](#page-2-0) We interpret \mathcal{P}_n^{ℓ} as a subfamily of \mathcal{B}_n . That is, the ground set is $2^{[n]}$, and each $X \in \mathcal{P}_n^{\ell}$ is mapped into an ℓ -element set on $2^{[n]}$, where each element of the set corresponds to a part from the partition X. In what follows, we think of \mathcal{P}_n^{ℓ} as a family of ℓ -element sets.

Let us derive that \mathcal{P}_n^{ℓ} is weakly $(\frac{n^2}{2})$ $\frac{b^2}{2}$, t)-spread for each $t \leq \ell - 2$. We take a collection of t distinct singletons as the set T in the definition of a weakly spread partition. The number of ℓ -partitions extending it is $\begin{cases} n-t \\ 0 \end{cases}$ $\ell-t$ The number of partitions with any $t+s$ fixed parts

is at most
$$
\begin{cases} n-t-s \\ \ell-t-s \end{cases}
$$
. Using Lemma 19, we have

$$
\begin{cases} n-t \\ \ell-t \end{cases} / \begin{cases} n-t-s \\ \ell-t-s \end{cases} \ge (n-t-s)^{2s} \ge (n^2/2)^s,
$$

provided $\ell - t - s \geq 1$. If $\ell - t - s = 0$ then we combine it with the bound $\begin{cases} n - t - s + 2 \end{cases}$ 2 $\}$ = $2^{n-t-s+1} \geq 2^{n/2} \geq n^4 = n^4 \begin{cases} n-t-s \end{cases}$ $\ell-t-s$ (the last inequality is valid due to our choice of n). Thus, the last displayed inequality is always true, and we conclude that \mathcal{P}_n^{ℓ} is weakly $\left(\frac{n^2}{2}\right)$ $\frac{1^2}{2}$, t)-spread for each $t \leq \ell - 2$.

Apply Theorem [14](#page-7-0) to F with $\varepsilon = 1/2$, ℓ playing the role of q and $r = n^2/2$. We need to check if $r \geq 48\ell$. The inequality clearly holds for n satisfying $n \geq 2\ell \log_2 n$ and $n \geq 48$. We conclude that either $\mathcal F$ is a family of partitions extending a fixed set of t singletons, or $|\mathcal{F}| \leq \frac{1}{2}$ $\int n-t$ $\ell-t$. This concludes the proof of the theorem. \Box

4.3. Profiled t-intersecting partitions. Proof of Theorem [7.](#page-2-1) For this proof, we also interpret U_P as a subfamily of the family corresponding to \mathcal{B}_n . This way, we have $\mathcal{U}_P \subset \bigl(\frac{2^{[n]}}{\ell}$ ^{[*u*}]. In this setup, we also can directly apply Theorem [14](#page-7-0) with $\epsilon = 1/2$ and get the desired conclusion, provided that we can show that $\mathcal{U}_{k,\ell}$ is a weakly (r, t) -spread family with $r \geq 48\ell$. We will show this below. Let a_t be the number of partitions that contain fixed parts of sizes k_1, \ldots, k_t , and let b_U be the number of partitions that contain fixed parts of sizes $k_i, i \in U$. Note that, for any w and U with $|U| = w$ we have $a_w \ge b_U$. Let U be a subset of [ℓ] of size $t + s$. For shorthand, let us denote $n_j = \sum_{i=j+1}^{\ell} k_i$ and note that for $j > j'$ we have $\frac{n_{j'}}{n_{j'}}$ $\frac{n_{j'}}{n_j} \geq \frac{\ell - j'}{\ell - j}$ $\frac{\ell-j}{\ell-j}$. Also note that in our assumptions we have $n_t \ge k_{t+1}(\ell-t) \ge \max{\{\ell, 2s\}}.$

$$
\frac{a_t}{b_U} \ge \frac{a_t}{a_{t+s}} = \frac{\frac{n_t}{\prod_{i=t+1}^{\ell} k_i!}}{\frac{n_{t+s}!}{\prod_{i=t+s+1}^{\ell} k_i!}} = \frac{n_t!}{n_{t+s}! \prod_{i=t+1}^{t+s} k_i!} \ge \frac{\prod_{i=n_{t+s}+1}^{n_t} i}{2^{s-1}((n_t - n_{t+s} - 2(s-1))!}.
$$

 $n.1$

In the last inequality, to bound the product of factorials in the denominator we used that for $k \geq k'$ we have $k!k' \leq (k+1)!(k-1)!$, as well as that $k_i \geq 2$ for $i \geq t+1$. We iteratively applied it to show that, in the conditions $k_i \geq 2$ and $\sum_{i=t+1}^{t+s} k_i = n_t - n_{t+s}$, the denominator is maximized when the first $s - 1$ k_i's are equal to 2. Below, we use that $\prod_{i=n_t-2s+3}^{n_t} i = (n_t)!/(n_t-2s+2)! \ge ((n_t)!)^{(2s-2)/n_t} \ge (n_t/e)^{2s-2}$ and that $n_t \ge \ell$. The last displayed expression is equal to

$$
\frac{\prod_{i=n_t-2s+3}^{n_t} i}{2^{s-1}} \frac{\prod_{i=n_{t+s}+1}^{n_t-2s+2} i}{((n_t - n_{t+s} - 2(s-1)))!} \ge \frac{(n_t/e)^{2s-2}}{2^{s-1}} \cdot \binom{n_t-2s+2}{n_{t+s}} \ge (\ell/6)^{2s-2} \binom{n_t-2s+2}{n_{t+s}}.
$$

If $s \geq \ell/4$, then it is an easy calculation to check $(\ell/6)^{2s-2} \geq (\ell/12)^{2s}$. If $s < \ell/4$ then $\binom{n_t-2s+2}{n_t}$ $\binom{-2s+2}{n_{t+s}} > \binom{\ell/2}{2}$ $\binom{2}{2}$. In any case, we can conclude that the RHS of the last displayed

inequality is at least $(\ell/12)^{2s}$. We get that the family \mathcal{U}_P is weakly $(\ell^2/12, t)$ -spread. Since $\ell^2/12 > 48\ell$ for $\ell > 600$, this is sufficient for our application of Theorem [14.](#page-7-0)

4.4. Partially *t*-intersecting partitions. Proof of Theorem [8.](#page-3-0) In what follows, we assume that $k \geq 3$. First, we interpret $\mathcal{U}_{k,\ell}$ as a family of sets. Put $N := \binom{k\ell}{2}$ $_{2}^{\mathfrak{c}\ell})$ and correspond to each partition P in $\mathcal{U}_{k,\ell}$ the set of all pairs of elements $x_1, x_2 \in [k\ell]$ such that x_1 and x_2 belong to the same part in P. Note that, in graph terms, P is a collection of ℓ *k*-cliques. As a set, each P has size $\ell \binom{k}{2}$ $\binom{k}{2}$. In what follows, we often work with partitions in this set form and think of $\mathcal{U}_{k,\ell}$ as a family in $\binom{[N]}{\binom{k}{2}\ell}$.

We shall show that, as a set family, $\mathcal{U}_{k,\ell}$ is sufficiently spread, provided ℓ is large enough. For that, we need to analyze the numbers of different partitions. Consider a collection $X = \{X_1, \ldots, X_a\}$ of disjoint sets in $[k\ell]$, such that $2 \leq |X_i| \leq k$. We call X a subpartition. Put $m(X) = \sum_{i=1}^{d} (|X_i| - 1)$. We say that a (k, ℓ) -partition $P = (P_1, \ldots, P_\ell)$ extends a subpartition X if for each i there is j such that $X_i \subset P_i$. Assume first that $m(X) \leq k\ell/3$. We claim that the number $u_{k,\ell}(X)$ of partitions that extend X is at most $\left(\frac{9}{\ell}\right)$ $\left(\frac{9}{\ell}\right)^m u_{k,\ell}$. Indeed, we have the following bound:

$$
\frac{u_{k,\ell}(X)}{u_{k,\ell}} = \frac{\frac{(k\ell - m(X) - a)!}{(\ell - a)!(k!)^{\ell - a} \prod_{i=1}^{a} (k - |X_i|)!}}{\frac{(k\ell)!}{\ell!(k!)^{\ell}}}\leq
$$

$$
\frac{k^{m(X) + a\ell a}}{(k\ell - m(X) - a)^{m(X) + a}} \leq \frac{k^{m(X) + a\ell a}}{(k\ell/3)^{m(X) + a}} \leq \frac{3^{m(X) + a}}{\ell^{m(X)}} \leq \frac{9^{m(X)}}{\ell^{m(X)}},
$$

where we used twice that $m(X) \ge a$. Note that this holds for subpartitions X that fix at most $k\ell/3$ elements and fix at least 2 elements in each "active" block. If a partition fixes more elements (but also has at least 2 elements in each block), then we can simply take a subpartition of size $k\ell/3$ and apply the bound to that subpartition. Thus, we get the following bound for larger subpartitions.

(5)
$$
\frac{u_{k,\ell}(X)}{u_{k,\ell}} \le \frac{9^{k\ell/3}}{\ell^{k\ell/3}} \le \frac{9^{m(X)/3}}{\ell^{m(X)/3}}.
$$

Note that the last bound is valid for any partition that fixes at least 2 elements in each block.

Let us now analyze, how is restricting to partitions that extend X is expressed in set terms. The family of partitions that extend X can be expressed in different ways.^{[2](#page-15-0)} Actually, it is necessary and sufficient for each $i \in [a]$ to fix a subgraph on X_i that is connected, and add no other edges. Thus, the largest number of edges we can fix is $\sum_{i=1}^{a} {\binom{|X_i|}{2}}$ $\sum_{i=1}^a$ $\frac{|X_i|(|X_i|-1)}{2} \leq \frac{k}{2}$ $\frac{k}{2}\sum_{i=1}^{a}(|X_i|-1) = \frac{km(X)}{2}$. Put differently, if we have fixed x edges, then the corresponding X satisfies $m(X) \geq \lceil \frac{2x}{k} \rceil$. We also note that, in the set interpretation, the corresponding partition cannot fix exactly 1 element in a part, and thus the bound [\(5\)](#page-15-1)

²In what follows, we refer to elements of $[N]$ as edges in the graph sense. It is natural since they correspond to pairs of elements of the original ground set.

is valid for all subpartitions that may arise this way. From here, we get that the family $\mathcal{U}_{k,\ell}$ is $(\frac{\ell}{9})^{2/3k}$ -spread.

If ℓ is sufficiently large (say, $\ell > k^{Ck}$ with some large constant C), this spreadness is sufficient to apply the spread approximation machinery. But before we do so, let us analyze, what happens with the t -intersecting property when passing to the set interpretation. If two (k, l) -partitions partially *t*-intersect, then they have $\binom{l}{2}$ $_{2}^{t}$) edges in common. In what follows, we will be working with this $\binom{t}{2}$ $_{2}^{t}$)-intersection property for set families. However, there is a complication that we have to overcome: there are obviously many other ways for two sets to have intersection of size $\geq \binom{t}{2}$ $_{2}^{t}$, without the corresponding partitions necessary being partially t-intersecting. Luckily, being partially t-intersecting is the most "economical" way, which allows us to overcome this complication.

Consider a $\binom{t}{2}$ ^t₂)-intersecting (in the set sense) family $\mathcal{F} \subset \mathcal{U}_{k,\ell}$. Apply Theorem [13](#page-6-0) with $r = (\frac{\ell}{9})^{1/3k}, r_0 = (\frac{\ell}{9})^{2/3k}, q = k^{10}$. The uniformity $\ell {k \choose 2}$ $\binom{k}{2}$ plays the role of k from the theorem. We need $(\ell/9)^{1/3k} > 2k^{10}$ and $(\frac{\ell}{9})^{2/k} > 2^{12} \log_2(2\ell(\frac{k}{2}))$ $\binom{k}{2}$). Both inequalities on r are valid, provided, say, $\ell > k^{100k}$. This allows us to apply Theorem [13](#page-6-0) and get a family S of sets of size at most q that cover most partitions in \mathcal{F} , and a remainder family $\mathcal{F}' \subset \mathcal{F}$ satisfying

(6)
$$
|\mathcal{F}'| \le (r/r_0)^{q+1} u_{k,\ell} \le \left(\left(\frac{9}{\ell}\right)^{1/3k}\right)^{k^{10}} u_{k,\ell} \le \ell^{-k^7} u_{k,\ell}.
$$

The next step is to apply Theorem [14](#page-7-0) to our approximation S with $\binom{t}{2}$ $_{2}^{t}$) playing the role of t. Before we do so, we need to show that $\mathcal{U}_{k,\ell}$ possess the weak $(r', \begin{bmatrix} \overline{t} \\ 2 \end{bmatrix})$ $_{2}^{t})$)-spreadness property with a sufficiently large r' . In order to do so, we need to return to the analysis of the subpartitions. Let E be a collection of $\binom{t}{2}$ $\binom{t}{2}$ edges and let $X = (X_1, \ldots, X_a)$ be the corresponding subpartition. Then $\binom{t}{2}$ $\binom{t}{2} = \sum_{i=1}^{a} {\binom{|X_i|}{2}}$ $\sum_{i=1}^{X_i}$). Recall that $m(X) = \sum_{i=1}^{a} (|X_i| - 1)$.

Observation 20. (i) We have $m(X) = t - 1$ if and only if $a = 1$, and the subpartition consists of just 1 set X_1 of size t.

(*ii*) If E is a collection of $\binom{t}{2}$ $\binom{t}{2}$ + s edges, then the corresponding partition $X(E)$ satisfies $m(X(E)) \geq t - 1 + \frac{s}{k}.$

Proof. (i) We have $\sum_{i=1}^{a} \binom{|X_i|}{2}$ $\binom{X_i}{2} \leq \max |X_i| \frac{m(X)}{2} \leq \binom{t}{2}$ $_{2}^{t}$, where the last inequality turns into equality iff $\max |X_i| = t$.

(ii) We know that for a set $E' \subset E$ of $\binom{t}{2}$ $\binom{t}{2}$ edges the corresponding partition $X(E')$ satisfies $m(X(E')) \geq t-1$. Next, assuming E is a collection of $\binom{t}{2}$ $\binom{t}{2} + s$ edges as in the proposition and that extends E' , we have

$$
s = |E| - |E'| = \sum_{i=1}^{a} \binom{|X_i(E)|}{2} - \binom{|X_i(E')|}{2}
$$

$$
\leq \sum_{i=1}^{a} \frac{|X_i(E)|^2 - |X_i(E')|^2}{2} \leq k \sum_{i=1}^{a} |X_i(E)| - |X_i(E')| = k(m(X(E)) - m(X(E'))).
$$

combining with $m(X(E')) \geq t - 1$, we get the result.

Choose a collection T of $\binom{t}{2}$ $\binom{t}{2}$ edges that corresponds to a subpartition of 1 t-element set (in other words, a t-clique). Once we are equipped with the property that for any collection of edges E of size $\binom{t}{2}$ $\binom{t}{2} + s$ the corresponding partition X satisfies $m(X) \geq t - 1 + \frac{s}{k}$, it is easy to do similar calculations as for $u_{k,\ell}$ and $u_{k,\ell}(X)$ above, and obtain that $\mathcal{U}_{n,k}(T)$ is r'-spread with $r' = \left(\frac{\ell-1}{9}\right)$ $\frac{(-1)}{9}$ ^{1/3k}. We note that this weak (r, t') -spreadness property is subtle in this application: the family $\mathcal{U}_{k,\ell}$ only possesses it for $t' = \begin{pmatrix} t \\ 2 \end{pmatrix}$ $t₂$ for integer t.

The next step is to apply Theorem [14](#page-7-0) to our approximation S. Let $\binom{t}{2}$ $_{2}^{t}$) play the role of t, $\epsilon = 1/2$, r' playing the role of r, and $q = k^{10}$, as above. Again, we can see that the inequality on r' is valid, provided ℓ is large enough (again $\ell > k^{Ck}$ is sufficient). The conclusion is that S must consist of a single set T of size $\binom{t}{2}$ \mathcal{L}_2^t , otherwise the size of $\mathcal{U}_{k,\ell}[S]$ is at least twice smaller than $\mathcal{U}_{k,\ell}[T]$ for the largest T. Moreover, T must correspond to a subpartition consisting of one t-element set.

At this point, we have proved a rough version of the conjecture, along with stability: if the size of a partially t-intersecting family of (k, ℓ) -partitions is at least $0.51|\mathcal{C}_{k,\ell}^T|$, then it is contained in some $\mathcal{C}_{k,\ell}^T$, with an exception of at most $\ell^{-k^7}u_{k,\ell}$ sets. These exceptional sets form the family \mathcal{F}' from above, and we next aim to show that this family is empty. (In what follows, we assume that $\mathcal{F}' \cap \mathcal{C}_{k,\ell}^T = \emptyset$, since otherwise we could move these sets to $\mathcal{F} \setminus \mathcal{F}'$.)

In order to show that $\mathcal{F}' = \emptyset$ for an extremal \mathcal{F} , we need to get some understanding on how often do random (k, ℓ) -partitions partially *t*-intersect.

Lemma 21. Let $t \geq 2$. For a given set T of size t and a (k, ℓ) -partition $Y = (Y_1, \ldots, Y_\ell) \notin$ $\mathcal{C}_{k,\ell}^T$, the number of other (k,ℓ) -partitions from $\mathcal{C}_{k,\ell}^T$ that do not partially t-intersect it is at least $\ell^{-2k^2}u_{k,\ell}$.

Proof. Fix some partition $X = (X_1, \ldots, X_\ell) \in C_{k,\ell}^T$ and consider a random permutation $\sigma : [k\ell] \setminus T \to [k\ell] \setminus T$. We will prove that

 $Pr[\sigma(X)]$ partially t-intersects $Y] \leq 1 - \ell^{-k^2}$.

Given that $|\mathcal{C}_{k,\ell}^T| \geq \ell^{-k^2} u_{k,\ell}$ and that a uniform random permutation σ generates a uniformly random element of $\mathcal{C}_{k,\ell}^T$, the statement of the lemma follows from the displayed formula.

We shall expose σ block by block, where the *i*-th block describes where is X_i mapped. Let us denote by A_i the event that $|\sigma(X_i) \cap Y_j| \geq t$ for some j. We suppose that $T \subset X_1$. Let us first deal with the first "exceptional" event A_1 . There are two possible ways for it to occur. First, it is possible that for some j we have $|Y_i \cap T| = t - 1$ and $\sigma(X_1 \setminus T) \cap Y_i \neq \emptyset$. Note, that there are at most 2 such j (at most 1 for $t \geq 3$). To bound this probability, we simply look at the probability that these sets intersect. The second possibility is covered by the event that $|\sigma(X_1 \setminus T) \cap Y_j| \geq 2$ for some j. This is the event that the sets of edges of $X_1 \setminus T$ and of the partition Y intersect. Thus, we can provide the following simple bound:

(7)
$$
\Pr[\sigma(X_1) \cap Y_j \ge t \text{ for some } j] \le \frac{2k|X_1 \setminus T|}{k\ell - t} + \frac{\binom{k}{2}\ell\binom{|X_1 \setminus T|}{2}}{\binom{k\ell - t}{2}} \le \frac{1}{2}
$$

Next, we are going to bound the probability that A_i happens, given that none of the A_1, \ldots, A_{i-1} happened. Each of these events is included in the event that $|\sigma(X_i) \cap Y_i| \geq 2$ for some j. This is the event that the sets of edges of $X_1 \backslash T$ and of the partition Y intersect. At this point, we are working with the partition that Y induces on $[k\ell] \setminus \sigma(X_1 \cup \ldots \cup X_{i-1})$. The latter set has size $k(\ell - i + 1)$. Restricted to it, Y is a partition into parts of size at most k each. Thus, the number of edges in Y is at most $\binom{k}{2}$ $\binom{k}{2}(\ell-i+1)$, and we have the bound

$$
(8) \quad \Pr[A_i | \bar{A}_1, \ldots, \bar{A}_{i-1}] \leq \frac{{k \choose 2} \sum_{j=1}^{\ell} { |Y_j \setminus (\sigma(X_1 \cup \ldots \cup X_{i-1}))| \choose 2}}{k \choose 2} \leq \frac{{k \choose 2}^2 (\ell-i+1)}{\binom{k(\ell-i+1)}{2}} \leq \frac{k^2}{2(\ell-i)}.
$$

We will use this bound up to $i = \ell - k^2$. The remaining parts, that is, $R := \sigma(X_{\ell-k^2+1} \cup$ $\dots \cup X_{\ell}$ form a set of size k^3 . The partition Y induced on R again consists of sets of size at most k, and clearly there is at least 1 choice of σ so that each X_i does not intersect each part of Y induced on R in more than 1 element. At the same time, the number of different partitions of R into k^2 parts of size k is at most k^{3k^3} , and thus

(9)
$$
\Pr[\bar{A}_{\ell-k^2+1} \cap \ldots \cap \bar{A}_{\ell} | \bar{A}_1, \ldots, \bar{A}_{\ell-k^2}] \geq k^{-3k^3}.
$$

Combining the bounds (7) , (8) , (9) , we get

$$
\Pr[\cap_{i=1}^{\ell} \bar{A}_i] \ge \frac{1}{2} k^{-3k^3} \prod_{i=1}^{\ell-k^2} \left(1 + \frac{k^2}{2(\ell-i) - k^2}\right)^{-1} \ge \frac{1}{2} k^{-3k^3} e^{-\sum_{i=1}^{\ell-k^2} \frac{k^2}{2(\ell-i) - k^2}} \ge
$$

$$
\frac{1}{2} k^{-3k^3} e^{-\frac{k^2}{2} \log_e \ell} = 12k^{-3k^3} \ell^{-\frac{k^2}{2}} \ge \ell^{-k^2}.
$$

This completes the proof of the lemma.

Returning to the extremal family \mathcal{F} , assume that \mathcal{F}' is non-empty and thus contains some partition Y. Lemma [21](#page-17-0) implies that $|\mathcal{C}_{k,\ell}^T \setminus \mathcal{F}| \geq \ell^{-2k^2} u_{k,\ell}$. But [\(6\)](#page-16-0) implies that $|\mathcal{F}'| \leq \ell^{-k^7}$. We conclude that $|\mathcal{F}'| \ll |\mathcal{C}_{k,\ell}^T \setminus \mathcal{F}|$, and thus \mathcal{F}' cannot be extremal unless \mathcal{F}' is empty. This concludes the proof of the theorem.

5. Acknowledgements

The research of the author was supported by the grant of the Russian Science Foundation N 24-71-10021.

.

REFERENCES

- [1] R. Ahlswede and L.H. Khachatrian, The Complete Intersection Theorem for Systems of Finite Sets, European Journal of Combinatorics. 18 (1997), 125–136.
- [2] R. Alweiss, S. Lovett, K. Wu, and J. Zhang, Improved bounds for the sunflower lemma, [arXiv:1908.08483](http://arxiv.org/abs/1908.08483) (2019)
- [3] M. Deza and P. Frankl, On the maximum number of permutations with given maximal or minimal distance, J. Combin. Theory, Ser. A 22 (1977), 352–360.
- [4] D. Ellis, Intersection Problems in Extremal Combinatorics: Theorems, Techniques and Questions Old and New (2021). https://doi.org/10.48550/arXiv.2107.06371.
- [5] D. Ellis, E. Friedgut, and H. Pilpel, Intersecting Families of Permutations, J. Amer. Math. Soc. 24 (2011), 649–682.
- [6] P. Erdős, C. Ko, and R. Rado, *Intersection theorems for systems of finite sets*, The Quart. J. Math. 12 (1961), N1, 313–320.
- [7] P.L. Erdős, L.A. Székely, Erdős-Ko-Rado Theorems of Higher Order, in: I. Althöfer, N. Cai, G. Dueck, L. Khachatrian, M.S. Pinsker, A. Sárközy, I. Wegener, Z. Zhang (Eds.), Numbers, Information and Complexity, Springer US, Boston, MA (2000), 117–124. https://doi.org/10.1007/978-1-4757-6048-4_11
- [8] P. Frankl, *The Erdős-Ko-Rado theorem is true for n=ckt*, Combinatorics (Proc. Fifth Hungarian Colloq., Keszthely, 1976), Vol. I, 365–375, Colloq. Math. Soc. János Bolyai, 18, North-Holland.
- [9] P. Frankl and Z. Füredi, *Beyond the Erdos-Ko-Rado theorem*, J. Combin. Theory Ser. A 56 (1991) N2, 182–194.
- [10] Chris Godsil and Karen Meagher, An algebraic proof of the Erdős–Ko–Rado theorem for intersecting families of perfect matchings, Ars Math. Contemp. 12 (2017), N2, 205–217.
- [11] N. Keller, N. Lifshitz, D. Minzer, O. Sheinfeld, On t-Intersecting Families of Permutations, arXiv.2303.15755 (2023).
- [12] A. Kupavskii and D. Zakharov, Spread approximations for forbidden intersections problems, [arXiv:2203.13379](http://arxiv.org/abs/2203.13379)
- [13] C.Y. Ku, D. Renshaw, *Erdős–Ko–Rado theorems for permutations and set partitions*, Journal of Combinatorial Theory, Series A. 115 (2008), 1008–1020. https://doi.org/10.1016/j.jcta.2007.12.004.
- [14] K. Meagher, L. Moura, *Erdős–Ko–Rado theorems for uniform set-partition systems*, The Electronic Journal of Combinatorics (2005), R40. https://doi.org/10.37236/1937.
- [15] K. Meagher, M.N. Shirazi, B. Stevens, An Extension of the Erdős–Ko–Rado Theorem to uniform set partitions (2021). https://arxiv.org/abs/2108.07692v1
- [16] T. Tao, The sunflower lemma via shannon entropy, https://terrytao.wordpress.com/2020/07/20/the-sunflower-lemm
- [17] H.S. Wilf, *generatingfunctionology*, CRC press (2005).
- [18] R.M. Wilson, *The exact bound in the Erdős–Ko–Rado theorem*, Combinatorica 4 (1984) 247–257.
- [19] https://en.wikipedia.org/wiki/Bell_number
- [20] https://en.wikipedia.org/wiki/Stirling_numbers_of_the_second_kind

Moscow Institute of Physics and Technology, Russia, St. Petersburg State University; Email: kupavskii@ya.ru