# AN EXTENSION OF LUCAS THEOREM

ZUBEYIR CINKIR AND AYSEGUL OZTURKALAN

ABSTRACT. We give elementary proofs of some congruence criteria to compute binomial coefficients in modulo a prime. These criteria are analogues to the symmetry property of binomial coefficients. We give extended version of Lucas Theorem by using those criteria. We give applications of these criteria by describing a method to derive identities and congruences involving sums of binomial coefficients.

## 1. INTRODUCTION

It is because of the beauty and the richness of number theory, and mathematics in general, that you can come up with new formulations and discoveries even in the basic and fundamental subjects.

Pascal's triangle and its properties have been investigated for a long time and it is a source of entertainment even for non-mathematicians. Its building blocks, binomial coefficients, are subject to numerous studies. For example, Lucas Theorem gives a remarkable method to compute binomial coefficients in modulo a prime. Namely,

**Theorem 1.1** (Lucas Theorem, [11]). For a given prime number n, let  $m \ge n$  be an integer and let k be an integer with  $0 \le k \le m$ . Suppose  $m = \sum_{i=0}^{r} m_i n^i$  and  $k = \sum_{i=0}^{r} k_i n^i$  are expansions of m and k in base n so that r is an integer of appropriate size and  $0 \le m_i < n$ and  $0 \le k_i < n$  are integers for each  $i = 0, 1, \dots, r$ . Then we have

$$\binom{m}{k} \equiv \binom{m_r}{k_r} \binom{m - m_r n^r}{k - k_r n^r} \mod n.$$

Moreover,

$$\binom{m}{k} \equiv \binom{m_r}{k_r} \binom{m_{r-1}}{k_{r-1}} \cdots \binom{m_1}{k_1} \binom{m_0}{k_0} \mod n.$$

Although, Lucas theorem is very powerful in reducing the computational complexity of  $\binom{m}{k}$  in modulo n when n is a prime and  $m \ge n$ , it is of no use when m < n in which case  $\binom{m}{k}$  can be very big comparing to n. Moreover, the application of Lucas Theorem comes down to the computation of some other binomial coefficients of numbers that are smaller than n. For example,

$$\binom{15683463}{10824637} \equiv \binom{248}{171} \binom{235}{205} \binom{230}{11} \mod 251$$

We still need to compute  $\binom{248}{171}$ ,  $\binom{235}{205}$  and  $\binom{230}{11}$ .

Key words and phrases. Binomial Coefficients, Pascal's Triangle, Lucas Theorem, Summation Identities, Cogruences of Binomial Coefficients.

The symmetry property  $\binom{m}{k} = \binom{m}{m-k}$  is also useful. Namely, it will be enough to compute the values of  $\binom{m}{k}$  only for those k in  $\{0, 1, \dots, \lfloor \frac{m}{2} \rfloor\}$ .

In this paper, we give elementary proofs of some other symmetry criteria so that the computations of binomial coefficients  $\binom{m}{k}$  in modulo a prime number will be much more simpler. In §2 (see Theorem 2.2 and Theorem 2.4), we obtained the following equivalences, where n is a prime, k and s are integers with  $0 \le s \le k < n$ :

(1) 
$$(-1)^{k+s} \binom{n-1-s}{n-1-k} = (-1)^{k+s} \binom{n-1-s}{k-s} \equiv \binom{k}{s} \\ = \binom{k}{k-s} \equiv (-1)^s \binom{n-1-k+s}{s} = (-1)^s \binom{n-1-k+s}{n-1-k} \mod n.$$

We can use (1) to derive the following extended version of Lucas theorem (by using (3)):

**Theorem 1.2** (Extended Lucas Theorem). For a given prime number n, let  $m \ge n$  be an integer and let k be an integer with  $0 \le k \le m$ . Suppose  $m = \sum_{i=0}^{r} m_i n^i$  and  $k = \sum_{i=0}^{r} k_i n^i$  are expansions of m and k in base n so that r is an integer of appropriate size and  $0 \le m_i < n$  and  $0 \le k_i < n$  are integers for each  $i = 0, 1, \dots, r$ . We set  $t_i = \min\{k_i, m_i - k_i\}$  for each  $i = 0, 1, \dots, r$ . Let us define  $\binom{m_i}{t_i}_n$  as follows

$$\binom{m_i}{t_i}_n := \begin{cases} (-1)^{t_i} \binom{n-1-m_i+t_i}{n-1-m_i}, & \text{if } m_i+t_i > n-1\\ (-1)^{t_i} \binom{n-1-m_i+t_i}{t_i}, & \text{if } m_i+t_i \le n-1 \text{ and } 2m_i > n-1+t_i\\ \binom{m_i}{t_i}, & \text{otherwise }. \end{cases}$$

Then we have

$$\binom{m}{k} \equiv \binom{m_r}{t_r}_n \binom{m - m_r n^r}{k - k_r n^r} \mod n.$$

Moreover,

$$\binom{m}{k} \equiv \binom{m_r}{t_r}_n \binom{m_{r-1}}{t_{r-1}}_n \cdots \binom{m_1}{t_1}_n \binom{m_0}{t_0}_n \mod n.$$

Observations given in (1) are not only helpful for the computation of binomial coefficients in modulo a prime number, but also crucial to devise a new method of proving identities and congruences that involve summations of binomial coefficients. In §3, we use (1) either to give new proofs of a known identity or to derive new identities. Similarly, we use (1) to establish (new) congruences in §4.

Having completed our paper, we noticed that symmetries in Pascal's triangle are studied and used in various articles such as [5, Section 4], [6], [8], [9], [13], [14], [15]. Almost all of them consider Pascal's triangle in modulo  $n^k$  for a prime n, and they consider Pascal's triangle with  $n^k$  or more rows. Many of them uses Lucas Theorem as a tool for their study. However, we focus only on the first n-1 rows, and work in modulo n. We use the symmetries to extend Lucas theorem and to devise a method of finding identities and congruences that involve Binomial coefficients.

# 2. Congruences of Binomial Coefficients

In this section, we establish some congruence criteria that we call semi-symmetry properties of binomial coefficients in modulo a prime. These are given in Theorem 2.2 and Theorem 2.4. We illustrate these criteria on Pascal's triangle. Then we describe an algorithm to apply these criteria to get a simplified equivalent form of a given binomial coefficients. This allows us to extend Lucas theorem. Then we give some examples.

The following simple equality will be very helpful. For each integer n, k and s with  $0 \le s \le k \le n$ , we have the cancellation identity [1, pg 29]:

(2) 
$$\binom{n}{k}\binom{k}{s} = \binom{n}{s}\binom{n-s}{k-s}.$$

Let us recall the following well known fact.

**Theorem 2.1.** A positive integer n is prime iff  $\binom{n-1}{i} \equiv (-1)^i \mod n$  for all  $0 \le i \le n-1$ .

In fact, only if part in Theorem 2.1 was proved by Lucas [10] in 1879.

Next, we give the first of our congruence criteria.

**Theorem 2.2.** Let n be a prime. For each integers k and s with  $0 \le s \le k < n$ , we have

$$\binom{k}{s} \equiv (-1)^{k+s} \binom{n-1-s}{k-s} = (-1)^{k+s} \binom{n-1-s}{n-1-k} \mod n.$$

*Proof.* By equation (2), we have

$$\binom{n-1}{k}\binom{k}{s} = \binom{n-1}{s}\binom{n-1-s}{k-s}.$$

Since *n* is a prime, we have  $\binom{n-1}{k} \equiv (-1)^k \mod n$ , and similarly  $\binom{n-1}{s} \equiv (-1)^s \mod n$ . The result follows from these congruences and the equality.

We noticed that the congruence given in Theorem 2.2 is obtained via different method in [14, Corollary 7]. The following result was proved in [13, Lemma 4] by induction.

**Corollary 2.3.** Let n be a prime. For  $1 \le i \le n$  and  $0 \le k \le n-i$  the following congruence holds:

$$\binom{n-i}{k} \equiv (-1)^k \binom{i-1+k}{i-1} \mod n.$$

*Proof.* With the given conditions on i and k, we have  $i - 1 + k \le n - 1$  and  $0 \le i - 1 < n$ . Thus, we apply Theorem 2.2 to obtain

$$\binom{i-1+k}{i-1} \equiv (-1)^{i-1+k+i-1} \binom{n-1-(i-1)}{i-1+k-(i-1)} = (-1)^k \binom{n-i}{k} \mod n.$$

This completes the proof.

Another result with the same flavor can be given as follows:

**Theorem 2.4.** Let n be a prime. For each integers k and s with  $0 \le s \le k < n$ , we have

$$\binom{k}{s} \equiv (-1)^s \binom{n-1-k+s}{s} = (-1)^s \binom{n-1-k+s}{n-1-k} \mod n$$

*Proof.* By the symmetry property,  $\binom{k}{s} = \binom{k}{k-s}$ . Then by Theorem 2.2,

$$\binom{k}{k-s} \equiv (-1)^{2k-s} \binom{n-1-k+s}{s} \equiv (-1)^s \binom{n-1-k+s}{s} \mod n.$$

This gives what we want, since the equality in the theorem follows from the symmetry.  $\Box$ 



FIGURE 1.

Next we give some examples. By Theorem 2.2 and the symmetry, we have

$$\binom{248}{171} = \binom{248}{77} = \binom{250-2}{79-2} \equiv (-1)^{81} \binom{79}{2} \equiv -3081 \equiv 182 \mod 251.$$

Similarly,

$$\binom{235}{205} = \binom{235}{30} = \binom{250 - 15}{45 - 15} \equiv (-1)^{60} \binom{45}{15} = 344867425584 \equiv 27 \mod 251,$$

and

$$\binom{230}{11} = \binom{250 - 20}{31 - 20} \equiv (-1)^{51} \binom{31}{20} = -\binom{31}{11} = -84672315 \equiv 25 \mod 251.$$

We can use Lucas Theorem along with our extensions:

$$\binom{15683463}{10824637} \equiv \binom{248}{171} \binom{235}{205} \binom{230}{11} \equiv \binom{79}{2} \binom{45}{15} \binom{31}{11} \\ \equiv 182 \cdot 27 \cdot 25 = 122850 \equiv 111 \mod 251.$$

The figure on the left in Figure 1 shows the Symmetry Property  $\binom{m}{k} = \binom{m}{m-k}$  and the symmetry line  $S_0$  in the Pascal's Triangle, and the figure on the right in Figure 1 shows the Semi-Symmetry Property in Theorem 2.2,  $\binom{k}{s} \equiv (-1)^{k+s} \binom{n-1-s}{n-1-k}$  and the symmetry line  $S_1$  in the Pascal's Triangle in modulo a prime n.

The figure on the left in Figure 2 shows the Semi-Symmetry Property described by Theorem 2.4,  $\binom{k}{s} \equiv (-1)^s \binom{n-1-k+s}{s}$  and the symmetry line  $S_2$  in the Pascal's Triangle in modulo a prime n.

The figure on the right in Figure 2 shows the symmetry lines and some certain regions of Pascal's triangle. These regions are critical when we apply symmetry and semi-symmetry properties. In this figure,

Binomial coefficients  $\binom{m}{k}$  on the symmetry line  $S_0$ ,  $S_1$  and  $S_2$  are given by the conditions k = m - k, m + k = n - 1 and 2m = n - 1 + k, respectively. Therefore,  $\binom{m}{k}$  is in the region A if  $k \leq m - k$  and  $2m \leq n - 1 + k$ . Likewise,  $\binom{m}{k}$  is in the region of B if  $k \leq m - k$ ,  $m + k \leq n - 1$  and  $2m \geq n - 1 + k$ . Moreover,  $\binom{m}{k}$  is in the region of C if  $k \leq m - k$ ,  $m + k \geq n - 1$ .



FIGURE 2.

**Question:** Given a binomial coefficient  $\binom{m}{k}$  and a prime n with  $0 \le k \le m < n$ , how can we apply symmetry and semi-symmetry properties effectively to get a simpler equivalent form of  $\binom{m}{k}$  in modulo n?

One can do this in various ways. Here is one possible method:

**Step 1:** If k > m - k, apply the symmetry property, i.e., replace  $\binom{m}{k}$  by  $\binom{m}{m-k}$ .

**Step 2:** We are given  $\binom{m}{k}$  with  $k \leq m-k$  as step 1 is applied. If m+k > n-1, apply the semi-symmetry property given in Theorem 2.2, i.e., replace  $\binom{m}{k}$  by  $(-1)^{m+k}\binom{n-1-k}{n-1-m}$ .

That is, if a binomial coefficient lies in the region C of the Pascal's triangle in the second figure of Figure 2, applying step 2 moves it to the region  $A \cup B$ . Note that the conditions  $k \leq m-k$  and m+k > n-1 ensures that n-1-m < (n-1-k) - (n-1-m).

**Step 3:** We are given  $\binom{m}{k}$  with  $k \leq m-k$  and  $m+k \leq n-1$  as step 1 and step 2 are applied. If 2m > n-1+k, apply the semi-symmetry property given in Theorem 2.4, i.e., replace  $\binom{m}{k}$  by  $(-1)^k \binom{n-1-m+k}{k}$ .

That is, if a binomial coefficient lies in the region B of the Pascal's triangle in the second figure of Figure 2, applying step 3 moves it to the region A. Hence, the region A is the fundamental region in Pascal's triangle modulo a prime.

Note that if step 2 is applied, then step 3 must be applied afterwards. To see this, one can check that output of step 2 (under the conditions of step 2) satisfies conditions of step 3. If step 2 is not applied, we may need to apply step 3 or not. This will lead to the following shorter answer to the question above:

Given  $\binom{m}{k}$ , set  $t := \min\{k, m-k\}$ . Then we have the following congruence in modulo n:

(3) 
$$\binom{m}{k} \equiv \begin{cases} (-1)^t \binom{n-1-m+t}{n-1-m}, & \text{if } m+t > n-1 \\ (-1)^t \binom{n-1-m+t}{t}, & \text{if } m+t \le n-1 \text{ and } 2m > n-1+t \\ \binom{m}{t}, & \text{otherwise (i.e. } m+t \le n-1 \text{ and } 2m \le n-1+t ). \end{cases}$$

Next, we give some examples:

Compute  $\binom{33}{20}$  in modulo 37. By step 1,  $\binom{33}{20} = \binom{33}{13}$ . Then

#### ZUBEYIR CINKIR AND AYSEGUL OZTURKALAN

1											1												1										
1	1										1	1											1	1									
1	2	1									1	2	1										1	2	1								
1	3	3	1								1	3	3	1									1	3	3	1							
1	4	6	4	1							1	4	6	4	1								1	4	6	4	1						
1	5	10	10	5	1						1	5	10	10	5	1							1	5	10	10	5	1					
1	6	4	9	4	6	1					1	6	4	9	4	6		L					1	6	4	9	4	6	1				
1	7	10	2	2	10	7	1				1	7	10	2	2	1	Э Т	7	1				1	7	10	2	2	10	7	1			
1	8	6	1	4	1	6	8	1			1	8	6	1	4	1	. (	5	8	1			1	8	6	1	4	1	6	8	1		
1	9	3	7	5	5	7	3	9	1		1	9	3	7	5	5	-	7	3	9	1		1	9	3	7	5	5	7	3	9	1	
1	10	1	10	1	10	1	10	1	10	1	1	10	1	10	) 1	1	0	L	10	1	10	1	1	10	1	10	1	10	1	10	1	10	1

### FIGURE 3.

1	1	1
1 1	1 1	1 1
1 <b>2</b> 1	1 <b>2</b> 1	1 <b>2</b> 1
<b>1 3</b> 3 1	<b>1 3</b> 3 1	<b>1 3</b> 3 1
1 4 6 4 1	<b>1 4 6</b> 4 1	1 4 6 4 1
1 5 10 10 5 1	1 5 10 10 5 1	<b>1</b> 5 10 10 5 1
<b>1 6 4 9</b> 4 6 1	<b>1 6 4 9</b> 4 6 1	1 6 <b>4 9</b> 4 6 1
<b>1 7 10 2</b> 2 10 7 1	<b>1 7 10 2</b> 2 10 7 1	1 7 10 2 2 10 7 1
1 8 6 1 <b>4</b> 1 6 8 1	<b>1 8 6</b> 1 4 1 6 8 1	1 8 6 1 4 1 6 8 1
<b>1 9 3 7 5</b> 5 7 3 9 1	<b>19</b> 37557391	1 9 3 7 5 5 7 3 9 1
1 10 1 10 1 <b>10</b> 1 10 1 10 1	<b>1</b> 10 1 10 1 10 1 10 1 10 1	1 10 1 10 1 10 1 10 1 10 1

FIGURE 4.

$$\binom{33}{13} \equiv (-1)^{33+13} \binom{23}{3} \mod 37, \text{ by step 2 as } 13 < 33 - 13 \text{ and } 33 + 13 > 37 - 1.$$
$$\equiv (-1)^3 \binom{16}{3} \equiv 32 \mod 37, \text{ by step 3 as } 2 \cdot 23 \ge 37 - 1 + 3.$$

Compute  $\binom{87}{40}$  in modulo 101.

$$\binom{87}{40} \equiv (-1)^{87+40} \binom{60}{13} \mod 101, \text{ by step 2 as } 40 < 87 - 40 \text{ and } 87 + 40 > 101 - 1.$$
$$\equiv -(-1)^{13} \binom{53}{13} \equiv 40 \mod 101, \text{ by step 3 as } 2 \cdot 60 \ge 101 - 1 + 13.$$

When n = 11, the symmetry line  $S_0$  and the semi-symmetry lines  $S_1$ ,  $S_2$  are illustrated in Figure 3. Furthermore, applications of above steps in this case are illustrated in Figure 4.

We used [12] to prepare Figure 3 and Figure 4.

The rest of the paper is about applications of Theorem 2.2 and Theorem 2.4. The identities shown in §3 and the congruences given in §4 are for the illustration purposes only. One can apply our method to various other cases.

# 3. Obtaining identities

In this section, we illustrate a method as an application of Theorem 2.2 and Theorem 2.4. It is about proving the summation identities involving binomial coefficients. Given such an identity, we apply any of these two theorems with a prime p to rewrite the identity. This

 $\mathbf{6}$ 

gives a congruence in modulo p. If we can choose the prime in such a way that the terms in the congruence are independent of p, the congruence is in fact an identity. Of course, the identity derived can be a known identity or a new identity. In any case, this gives a new effective method. Actually, this method shows that there is a duality between such identities.

The first equality in the following theorem was known (set x = n in the equation 3.50 in [2]). Here, we derive it with a new proof.

**Theorem 3.1.** Let n be a nonnegative integer. Then we have

$$\sum_{j=0}^{n} (-1)^{j} \binom{n}{j} \binom{2n-j}{n} = 1, \quad \sum_{j=0}^{n} (-1)^{j} \binom{2n-j}{j} \binom{2n-2j}{n-j} = 1$$

*Proof.* By Vandermonde identity, for any m > n we have  $\binom{m-1}{n} = \sum_{j=0}^{n} \binom{n}{j} \binom{m-1-n}{n-j}$ . When m is a prime, we have  $\binom{m-1}{n} \equiv (-1)^n \mod m$ . On the other hand, we use Theorem 2.2 to derive  $(-1)^{m+j} \binom{m-1-n}{n-j} \equiv (-1)^{n+1} \binom{2n-j}{n} \mod m$  if m is a prime. Then we obtain

(4) 
$$(-1)^n \equiv \sum_{j=0}^n (-1)^{m+n+j+1} \binom{n}{j} \binom{2n-j}{n} \mod m.$$

We can rewrite the congruence (4) as follows:

(5) 
$$1 \equiv \sum_{j=0}^{n} (-1)^{j} \binom{n}{j} \binom{2n-j}{n} \mod m$$

Note that the congruence (5) is independent of m. By choosing m large enough, we see that the congruence (5) is in fact an equality. This gives the first equality in the theorem.

The second equality in the theorem follows from the first one if we use the cancellation identity  $\binom{2n-j}{n}\binom{n}{j} = \binom{2n-j}{j}\binom{2n-2j}{n-j}$ .

**Theorem 3.2.** Let s and m be any positive integers. Then we have

$$\sum_{j=0}^{s} (-1)^{j} \binom{m+s+j}{s} \binom{s}{j} = (-1)^{s}, \quad \sum_{j=0}^{s} (-1)^{j} \binom{m+s+j}{j} \binom{m+s}{s-j} = (-1)^{s}.$$

*Proof.* We use the following equality [4, pg 169] (consider first equality at pg. 169 and set n = 0) for any integers m, s and l:

$$\sum_{j=0}^{s} \binom{l}{m+j} \binom{s}{j} = \binom{l+s}{l-m}.$$

Next, for a prime p, we take l = p - 1 - s and rewrite this equation to obtain:

(6) 
$$\sum_{j=0}^{s} \binom{p-1-s}{m+j} \binom{s}{j} = \binom{p-1}{p-1-s-m}$$

On the other hand, we have  $\binom{p-1}{p-1-s-m} \equiv (-1)^{p-1-s-m} = (-1)^{s+m} \mod p$  by Theorem 2.1 and  $\binom{p-1-s}{m+j} = \binom{p-1-s}{m+j+s-s} \equiv (-1)^{m+j+2s} \binom{m+s+j}{s} \mod p$  by Theorem 2.2. We use these in (6) to obtain:

(7) 
$$\sum_{j=0}^{s} (-1)^{j} \binom{m+s+j}{s} \binom{s}{j} \equiv (-1)^{s} \mod p.$$

Note that the congruence (7) is independent of p. By choosing p large enough, we see that the congruence (7) is in fact an equality. This gives the first equality in the theorem.

The second equality in the theorem follows from the first one if we use the cancellation identity  $\binom{m+s+j}{s}\binom{s}{j} = \binom{m+s+j}{j}\binom{m+s}{s-j}$ .

Note that the identity given in Theorem 3.2 is closely related to the identity with number 3.47 in [2] (consider the case r = n in 3.47).

**Theorem 3.3.** Let s and m be any positive integers. Then we have

$$\sum_{j=0}^{m} (-1)^j \frac{\binom{m}{j}}{\binom{s+j}{s}} = \frac{s}{s+m}.$$

*Proof.* We use the following equality [4, pg 173] for any integers  $n \ge m \ge 0$ :

$$\sum_{j=0}^{m} \frac{\binom{m}{j}}{\binom{n}{j}} = \frac{n+1}{n+1-m}.$$

Given positive integers m and s, take a prime p such that  $p - 1 - s \ge m$ . Next, we rewrite the equation above for m and n = p - 1 - s:

(8) 
$$\sum_{j=0}^{m} \frac{\binom{m}{j}}{\binom{p-1-s}{j}} = \frac{p-s}{p-s-m}.$$

On the other hand, we have  $\binom{p-1-s}{j} = \binom{p-1-s}{j+s-s} \equiv (-1)^{j+2s} \binom{s+j}{s} \mod p$  by Theorem 2.2. We use these in (8) to obtain:

(9) 
$$\sum_{j=0}^{m} (-1)^j \frac{\binom{m}{j}}{\binom{s+j}{s}} \equiv \frac{s}{s+m} \mod p.$$

Note that the congruence (9) is independent of p. By choosing p large enough, we see that the congruence (9) is in fact an equality. This gives what we want.

Note that the identity given in Theorem 3.3 was listed as the identity with number 1.42 and 4.30 in [2].

**Theorem 3.4.** Let s be any positive integer. Then we have

$$\sum_{j=0}^{s} (-1)^{j} j \binom{s+j}{s} \binom{s}{j} = (-1)^{s} (s+1)s$$

*Proof.* We use the following equality [4, pg 181] for any integers s and  $n \ge 0$ :

$$\sum_{j=0} j\binom{n}{j}\binom{s}{j} = s\binom{n+s-1}{n-1}.$$

Given a positive integer s, take an odd prime p such that  $p - 1 - s \ge 0$ . Next, we rewrite the equation above for s and n = p - 1 - s:

(10) 
$$\sum_{j=0}^{s} j \binom{p-1-s}{j} \binom{s}{j} = s \binom{p-2}{s}.$$

On the other hand, we have  $\binom{p-2}{s} \equiv (-1)^s \binom{s+1}{1} \mod p$  and that  $\binom{p-1-s}{j} = \binom{p-1-s}{j+s-s} \equiv (-1)^{j+2s} \binom{s+j}{s} \mod p$  by Theorem 2.2. We use these in (10) to obtain:

(11) 
$$\sum_{j=0}^{s} (-1)^{j} j \binom{s+j}{j} \binom{s}{j} \equiv (-1)^{s} (s+1)s \mod p.$$

Note that the congruence (11) is independent of p. By choosing p large enough, we see that the congruence (11) is in fact an equality. This gives what we want.

**Theorem 3.5.** Let s and n be positive integers. Then we have

$$\sum_{j=1}^{n} (-1)^{j} \binom{s-1}{j-1} \frac{1}{j} = (-1)^{n} \frac{1}{s} \binom{s-1}{n} - \frac{1}{s}$$

*Proof.* We use the identity [3, Identity 8.13 at pg 48]

$$\sum_{j=1}^{n} \binom{j+r-1}{r} \frac{1}{j} = \frac{1}{r} \binom{n+r}{r} - \frac{1}{r}$$

For any given positive integers n and s with s > n, take r = p - s for some prime p > s. Then we notice that  $\binom{s-1}{j-1} \equiv (-1)^{j-1} \binom{p-1-s+j}{j-1} \mod p$  by Theorem 2.4. Therefore,

(12)  

$$\sum_{j=1}^{n} (-1)^{j} {\binom{s-1}{j-1}} \frac{s}{j} \equiv \sum_{j=1}^{n} {\binom{p-1-s+j}{j-1}} \frac{p-s}{j} \mod p$$

$$= \sum_{j=1}^{n} {\binom{p-1-s+j}{p-s}} \frac{p-s}{j}$$

$$\equiv {\binom{n+p-s}{p-s}} - 1 \mod p.$$

For the right hand side,  $\binom{n+p-s}{p-s} = \binom{p-1-(s-1)+n}{n} \equiv (-1)^n \binom{s-1}{n} \mod p$  by Theorem 2.4 when  $s-1 \ge n$ . Thus, we can rewrite (12) as follows:

(13) 
$$\sum_{j=1}^{n} (-1)^{j} {\binom{s-1}{j-1}} \frac{s}{j} \equiv (-1)^{n} {\binom{s-1}{n}} - 1 \mod p.$$

As the congruence (13) is independent of p, it must be an equality. Hence, the theorem is proved when s > n.

If  $s \leq n$ , we only need to show that  $\sum_{j=1}^{s} (-1)^j {s-1 \choose j-1} = -1$ . Using  ${s \choose j} = \frac{s}{j} {s-1 \choose j-1}$  when  $j \neq 0$ , we need  $\sum_{j=1}^{s} (-1)^j {s \choose j} = -1$  which clearly holds.

# 4. Obtaining congruences

In this section, we give examples of obtaining congruences in modulo a prime by applying Theorem 2.2 and Theorem 2.4. This is interesting, because it shows the connections between seemingly unrelated congruences.

9

**Theorem 4.1.** Let n be a prime. For any nonnegative integer s < n - 1 we have

$$\sum_{k=1}^{n-1-s} \frac{1}{k} \binom{k+s}{s} \equiv -\sum_{k=1}^{n-1-s} \frac{1}{k} \mod n.$$

In particular, when s = 0 we have,  $\sum_{k=1}^{n-1} \frac{1}{k} \equiv 0 \mod n$  if n > 2.

*Proof.* We have the following equality (see [2, identity 1.45]), for any integer m:

$$\sum_{k=1}^{m} (-1)^{k-1} \frac{1}{k} \binom{m}{k} = \sum_{k=1}^{m} \frac{1}{k}.$$

We take m = n - 1 - s and applying Theorem 2.2, we derive the main congruence. The second congruence follows from the first one if we set s = 0. Note that the second congruence was obtained in modulo  $p^2$  by J. Wolstenholme in 1862.

Next, we have another example.

**Theorem 4.2.** Let p be an odd prime and let r be an integer such that  $0 \le r \le \frac{p-1}{2}$ . Then we have

$$\sum_{j=r}^{\frac{p-1}{2}} \binom{2j}{j} \binom{j}{r} 2^{p-1-2k} \equiv \begin{cases} 0 \mod p, & \text{if } r < \frac{p-1}{2} \\ (-1)^{\frac{p-1}{2}} \mod p, & \text{if } r = \frac{p-1}{2}. \end{cases}$$

*Proof.* We use the following equality [2, pg. 45] for any integers r and n:

$$\sum_{j=0}^{\lfloor \frac{n}{2} \rfloor} (-1)^j \binom{n-j}{j} \binom{j}{r} 2^{n-2k} = (-1)^r \binom{n+1}{2r+1}.$$

Taking n = p - 1 for a odd prime p in this equality gives

(14) 
$$\sum_{j=0}^{\frac{p-1}{2}} (-1)^j \binom{p-1-j}{j} \binom{j}{r} 2^{p-1-2k} = (-1)^r \binom{p}{2r+1}.$$

Since we have  $\binom{p-1-j}{j} \equiv (-1)^j \binom{2j}{j} \mod p$  by Theorem 2.2, the result follows from (14) by noting that  $\binom{p}{2r+1} \equiv 0 \mod p$  if  $r < \frac{p-1}{2}$  and  $\binom{p}{2r+1} \equiv 1 \mod p$  if  $r = \frac{p-1}{2}$ .

The following example shows that one can prove many congruences easily by applying Theorem 2.2 and Theorem 2.4.

**Theorem 4.3.** We have the following congruences in modulo an odd prime p:

$$F_{p} \equiv \sum_{0 \le i, j \le p-1} (-1)^{i+j} {\binom{i+j}{i}}^{2} \equiv \sum_{j=0}^{\frac{p-1}{2}} (-1)^{j} {\binom{2j}{j}} \equiv \begin{cases} 1, & \text{if } p \equiv \pm 1 \mod 5\\ -1, & \text{if } p \equiv \pm 2 \mod 5. \end{cases}$$
$$F_{p+1} \equiv \sum_{0 \le i, j \le p} (-1)^{i+j} {\binom{i+j-1}{j}} {\binom{i+j}{i}} \equiv \sum_{j=0}^{\frac{p-1}{2}} (-1)^{j} {\binom{2j-1}{j}} \equiv \begin{cases} 1, & \text{if } p \equiv \pm 1 \mod 5\\ 0, & \text{if } p \equiv \pm 2 \mod 5. \end{cases}$$

#### AN EXTENSION OF LUCAS THEOREM

*Proof.* We use the following equality [7, pg. 1570] for any positive integer n:

(15) 
$$F_{n+1} = \sum_{0 \le i, j \le n} (-1)^i \binom{n-i}{j} \binom{i+j}{i}.$$

Taking n = p - 1 for an odd prime p and using  $\binom{p-1-i}{j} \equiv (-1)^j \binom{i+j}{j} = (-1)^j \binom{i+j}{i} \mod p$  by Theorem 2.2 in (15) gives

(16) 
$$F_p \equiv \sum_{0 \le i, j \le p-1} (-1)^{i+j} \binom{i+j}{i}^2 \mod p.$$

Similarly, taking n = p for an odd prime p and using  $\binom{p-i}{j} \equiv (-1)^j \binom{i+j-1}{j} \mod p$  by Theorem 2.4 in (15) gives

(17) 
$$F_{p+1} \equiv \sum_{0 \le i, j \le p} (-1)^{i+j} \binom{i+j-1}{j} \binom{i+j}{i} \mod p.$$

On the other hand, we have

(18) 
$$F_n = \sum_{j=0}^{\lfloor \frac{n-1}{2} \rfloor} \binom{n-1-j}{j}.$$

In particular, for an odd prime p

(19) 
$$F_p = \sum_{j=0}^{\frac{p-1}{2}} {p-1-j \choose j}.$$

We have  $\binom{p-1-j}{j} \equiv (-1)^j \binom{2j}{j} \mod p$  by Theorem 2.2. Thus, the following congruence is obtained from (19):

(20) 
$$F_p \equiv \sum_{j=0}^{\frac{p-1}{2}} (-1)^j \binom{2j}{j} \mod p.$$

Again, for an odd prime p it follows from (18) that

(21) 
$$F_{p+1} = \sum_{j=0}^{\frac{p-1}{2}} {p-j \choose j}.$$

We have  $\binom{p-j}{j} \equiv (-1)^j \binom{2j-1}{j} \mod p$  by Theorem 2.4. Thus, the following congruence is obtained from (21):

(22) 
$$F_{p+1} \equiv \sum_{j=0}^{\frac{p-1}{2}} (-1)^j \binom{2j-1}{j} \mod p.$$

Finally, we have  $F_p \equiv 1 \mod p$  and  $F_{p+1} \equiv 1 \mod p$  if  $p \equiv \pm 1 \mod 5$ . We also have  $F_p \equiv -1 \mod p$  and  $F_{p+1} \equiv 0 \mod p$  if  $p \equiv \pm 2 \mod 5$  [16, pg 78-79]. This completes the proof.

### ZUBEYIR CINKIR AND AYSEGUL OZTURKALAN

**Declaration of competing interest:** The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

#### References

- D. Bressoud and S. Wagon, A Course in Computational Number Theory. John Wiley & Sons, NJ, (2000).
- [2] H. W. Gould, *Combinatorial Identities*. Morgantown Printing and Binding Co., Morgantown W. Va., 1972.
- [3] H. W. Gould, Combinatorial Identities Volume 1. Can be found at http://www.math.wvu.edu/~hgould/Vol.1.PDF, 2010.
- [4] R. Graham, D. Knuth and O. Patashnik, *Concrete Mathematics*. Addison-Wesley Publishing Company, New York, 1989.
- [5] A. Granville, Zaphod Beeblebrox's brain and the fifty-ninth row of Pascal's triangle, The American Mathematical Monthly, 99, no. 4, (1992), 318–331.
- [6] N. E. Myerson-Jain, S. Liu, W. Ji, C. Xu and S. Vijay, *Pascal's Triangle Fractal Symmetries*, Phys. Rev. Lett., Vol 128, Issue 11, (2022), 115301–115307.
- [7] E. Kılıç and T. Arıkan, Double binomial sums and double sums related with certain linear recurrences of various order, Chiang Mai Journal of Science, 45, (2018), 1569–1577.
- [8] R. Kubelka, Self-similarity and symmetries of Pascal's triangles and simplices mod p, Fibonacci Quarterly, 42, (2004), 70–75.
- [9] C. T. Long, Pascal's triangle modulo p, Fibonacci Quart., 19, No. 5, (1981), 458-463.
- [10] E. Lucas, Amer. Jour. Math., 1, (1879), 229–230.
- [11] E. Lucas, Sur les congruences des nombres eulériens et les coefficients différentiels des fonctions trigonométriques, suivant un module premier, Bull. Soc. Math. France 6 (1877–1878), 49–54.
- [12] Wolfram Research, Inc., Mathematica, Version 13.1, Wolfram Research Inc., Champaign, IL., (2022).
- [13] M. Prunescu, Symmetries in the Pascal triangle: p-adic valuation, sign-reduction modulo p and the last non-zero digit, Bull. Math. Soc. Sci. Math. Roumanie, Tome 65 (113), No. 4, (2022), 431-447.
- [14] Eric Rowland, Lucas' Theorem Modulo p<sup>2</sup>, The American Mathematical Monthly, 129:9, (2022), 846– 855.
- [15] K. M. Shannon, M. J. Bardzell, Searching for patterns in Pascal's Triangle With a Twist, Journal of Online Mathematics and its Applications, 3, (2003).

https://maa.org/press/periodicals/loci/joma/patterns-in-pascals-triangle-with-a-twist

[16] S. Vajda, Fibonacci and Lucas Numbers and The Golden Section, Ellis Horwood Limited, Chichester, England, 1989.

ZUBEYIR CINKIR, DEPARTMENT OF INDUSTRIAL ENGINEERING, ABDULLAH GUL UNIVERSITY, 38100, KAYSERI, TURKEY,

Email address: zubeyir.cinkir@agu.edu.tr

AYSEGUL OZTURKALAN, DEPARTMENT OF ENGINEERING SCIENCES, ABDULLAH GUL UNIVERSITY, 38100, KAYSERI, TURKEY,

Email address: aysegul.ozturkalan@agu.edu.tr