Intuitionistic Gödel-Löb logic, à la Simpson: labelled systems and birelational semantics

Anupam Das ⊠

University of Birmingham, United Kingdom

Iris van der Giessen ⊠ University of Birmingham, United Kingdom

Sonia Marin ⊠ University of Birmingham, United Kingdom

— Abstract

We derive an intuitionistic version of Gödel-Löb modal logic (GL) in the style of Simpson, via proof theoretic techniques. We recover a labelled system, ℓIGL , by restricting a non-wellfounded labelled system for GL to have only one formula on the right. The latter is obtained using techniques from cyclic proof theory, sidestepping the barrier that GL's usual frame condition (converse wellfoundedness) is not first-order definable. While existing intuitionistic versions of GL are typically defined over only the box (and not the diamond), our presentation includes both modalities.

Our main result is that ℓIGL coincides with a corresponding semantic condition in birelational semantics: the composition of the modal relation and the intuitionistic relation is conversely well-founded. We call the resulting logic IGL. While the soundness direction is proved using standard ideas, the completeness direction is more complex and necessitates a detour through several intermediate characterisations of IGL.

2012 ACM Subject Classification Theory of computation \rightarrow Proof theory

Keywords and phrases provability logic, proof theory, intuitionistic modal logic, cyclic proofs, non-wellfounded proofs, proof search, cut-elimination, labelled sequents

 ${\sf Funding}~{\rm This}$ work was partially supported by a UKRI Future Leaders Fellowship, 'Structure vs Invariants in Proofs', project reference MR/S035540/1.

Acknowledgements We would like to thank Marianna Girlando and Jan Rooduijn for several valuable discussions on the topic. We also thank Marianna for her significant input during our reading group sessions on intuitionistic modal logic that led to preliminary ideas that have resulted in this paper.

1 Introduction

Gödel-Löb logic (GL) originates in the provability reading of modal logic: \Box is interpreted as "it is provable that", in an arithmetical theory with suitable coding capacity, inducing its corresponding provability logic. Löb formulated a set of necessary conditions on the provability logic of Peano Arithmetic (PA), giving rise to GL, extending basic modal logic K by what we now call Löb's axiom: $\Box(\Box A \to A) \to \Box A$. Somewhat astoundingly GL turns out to be complete for PA's provability logic, a celebrated result of Solovay [40]: all that PA can prove about its own provability is already a consequence of a relatively simple (and indeed decidable) propositional modal logic.

Proof theoretically Löb's axiom represents a form of *induction*. Indeed, at the level of modal logic semantics, GL enjoys a correspondence with transitive relational structures that are *terminating*¹ [36]. Duly, in computer science, Löb's axiom has inspired several

¹ Other authors refer to this property as *Noetherian* or *conversely well-founded*, but we opt for this simpler nomenclature.

variants of modal type theories that extend simply-typed lambda calculus with some form of recursion. These range, for instance, from the seminal work of Nakano [30], to more recent explorations into guarded recursion [6] and intensional recursion [23]. Based in type theories, these developments naturally cast GL in a *constructive* setting, but little attention was given to analysing the induced intuitionistic modal logics. Indeed, for this reason, [10] has proposed a more foundational basis for studying computational interpretations of GL, by way of a sequent calculus for the logic of the topos of trees.

Returning to the provability reading of modal logic, several constructive variants of GL have been independently proposed (see [26, 16] for overviews). An important such logic is iGL (and its variants) which now enjoys a rich mathematical theory, from semantics (e.g., [26]) to proof theory (e.g., [17, 19]). Interestingly, while iGL is known to be sound for the provability logic of Heyting Arithmetic (HA), the intuitionistic counterpart of PA, it is not complete. The provability logic of HA has been recently announced by [29], currently under review.

One shortfall of all the above mentioned approaches is that they do not allow us to recover a bona fide computational interpretation of classical GL along, say, the Gödel-Gentzen negative translation (GG), a standard way of lifting interpretations of intuitionistic logics to their classical counterparts. Indeed it was recently observed that the 'i' (or 'constructive') traditions of intuitionistic modal logic are too weak to validate the GG translation [12, 13].

On the other hand modal logic's relational semantics effectively renders it a fragment of usual first-order predicate logic (FOL), the so-called standard translation. Interpreting this semantics in an intuitionistic meta-theory defines a logic that *does* validate GG, for the same reason that intuitionistic FOL GG-interprets classical FOL. This is the approach taken (and considerably developed) by Simpson [38], building on earlier work of Fischer Servi [15] and Plotkin and Stirling [33]. The resulting logic IK (and friends) enjoys a remarkably robust proof theory by way of *labelled systems*, which may be duly seen as a fragment of Gentzen's systems for intuitionistic FOL by way of the standard translation.

Contribution

In this work we develop an intuitionistic version of GL, following Simpson's methodology [38]. In particular, while logics such as iGL are defined over only the \Box , our logic is naturally formulated over both the \Box and the \diamond . A key stumbling block to this end, as noticed already in the classical setting by Negri in [31], is that GL's correspondence to terminating relations cannot seemingly be inlined within a standard labelled system: termination is not even FOL-definable. To side-step this barrier we draw from a now well-developed proof-theoretic approach to (co)induction: non-wellfounded proofs (e.g. [32, 9, 39, 3, 14, 11]). Such proofs allow infinite branches, and so are (typically) equipped with a progress condition that ensures sound reasoning. Starting from a standard labelled system for transitive relations, K4, we recover a labelled calculus ℓGL for GL by allowing non-wellfounded derivations under a typical progress condition. In fact, our progress condition is precisely the one from Simpson's Cyclic Arithmetic [39, 11]. Following (the same) Simpson, we duly recover an intuitionistic version ℓ IGL of GL syntactically in a standard way: we restrict ℓ GL to one formula on the right.

At the same time we can recover an intuitionistic version of GL semantically by suitably adapting the *birelational semantics* of intuitionistic modal logics, which combine the partial order \leq of intuitionistic semantics with the accessibility relation R of modal semantics. Our semantic formulation \mathscr{B} IGL is duly obtained by reading the termination criterion of classical GL's semantics intuitionistically: the composition $\leq R$ must be terminating. Our main result is that these two characterisations, ℓIGL and $\mathcal{B}IGL$, are indeed equivalent, and we duly dub the resulting logic IGL.



Figure 1 Summary of our main results, where arrows \rightarrow denote inclusions of modal logics.

The soundness direction is proved via standard techniques from intuitionistic modal logic and non-wellfounded proof theory. The completeness direction, on the other hand, is more cumbersome. To this end we exercise an intricate combination of proof theoretic techniques, necessitating two further (and ultimately equivalent) characterisations of IGL: semantically \mathscr{P} IGL, essentially a class of intuitionistic FOL structures, and syntactically m ℓ IGL, a multisuccedent variant of ℓ IGL. These formulations facilitate a countermodel construction from failed proof search, inspired by Takeuti's for LJ [41]. Due to the non-wellfoundedness of proofs we employ a determinacy principle to organise the construction, a standard technique in non-wellfounded and cyclic proof theory (e.g. [32]). For the reduction to ℓ IGL we devise a form of continuous cut-elimination, building on more recent ideas in non-wellfounded proof theory, cf. [3, 14]. Our 'grand tour' of results is visualised in Figure 1, also indicating the organisation of this paper. Due to space constraints, proofs are relegated to appendices.

Other related work

The proof theory of GL is (in)famously complex. The first sequent calculus of GL was considered in [24] but its cut-elimination property was only finally settled (positively) in [18] after several attempts [43, 35, 28]. Intuitionistic versions of sequent calculi for GL are developed in [17, 19] and provide calculi for iGL. Labelled calculi [31] and *nested calculi* [34] have also been proposed. However all of these calculi are arguably unsatisfactory: the modality introduction rules are non-standard (which is atypical for labelled calculi) and, in particular, modal rules may change the polarity of formula occurrences, due to the inductive nature of Löb's axiom.

It is possible to recover a system for GL by admitting non-wellfounded proofs in a sequent calculus for K4, as observed by Shamkanov [37]. An intuitionistic version of this system has been studied by Iemhoff [21]. In these works both the base calculus and the corresponding correctness criterion are bespoke, rather than 'recovered' from established foundations.

Non-wellfounded proofs originate in the study of modal logics with fixed points, in particular Niwinski and Walukiewicz's seminal work on the μ -calculus [32]. These ideas were later inlined into the proof theory of FOL with certain inductive definitions by Brotherston and Simpson (e.g. [9]), a source of inspiration for the present work. As already mentioned, recent extensions of these ideas to PA [39, 11] and advances on cut-elimination [3, 14] are quite relevant to our development.

2 Preliminaries on (classical) modal logic

Throughout this work we work with a set Pr of **propositional symbols**, written p, q, etc., which we simultaneously construe as unary predicate symbols when working in predicate logic. For the latter we also assume a single binary relation symbol R and a countable set Var of (individual) variables, written x, y, etc.

2.1 Language and semantics

The formulas of modal logic are generated by the following grammar:

 $A \quad ::= \quad p \in \mathsf{Pr} \ \mid \ \bot \ \mid \ A \land A \ \mid \ A \lor A \ \mid \ A \to A \ \mid \ \Box A \ \mid \ \Diamond A$

As usual we write $\neg A := A \rightarrow \bot$, and employ standard bracketing conventions.

Modal formulas are interpreted over relational frames $\mathcal{F} = (W, R^{\mathcal{F}})$ formed of a non empty set of worlds W equipped with an accessibility relation $R^{\mathcal{F}} \subseteq W \times W$.² A relational model $\mathcal{M} = (W, R^{\mathcal{M}}, V)$ is a structure where $(W, R^{\mathcal{M}})$ is a frame with a valuation $V : W \to \mathcal{P}(\mathsf{Pr})$. Let $\mathcal{M} = (W, R^{\mathcal{M}}, V)$ be a relational model. For worlds $w \in W$ and formulas A we define the satisfaction judgement $\mathcal{M}, w \models A$ as follows:

- $= \mathcal{M}, w \vDash p \text{ if } p \in V(w); \qquad = \mathcal{M}, w \vDash A \land B \text{ if } \mathcal{M}, w \vDash A \text{ and } \mathcal{M}, w \vDash B;$
- $= \mathcal{M}, w \nvDash \bot; \qquad = \mathcal{M}, w \vDash A \lor B \text{ if } \mathcal{M}, w \vDash A \text{ or } \mathcal{M}, w \vDash B;$
- $= \mathcal{M}, w \vDash A \to B \text{ if } \mathcal{M}, w \vDash A \text{ then } \mathcal{M}, w \vDash B;$
- $\blacksquare \mathcal{M}, w \vDash \Box A$ if for all v such that $wR^{\mathcal{M}}v$ we have $\mathcal{M}, v \vDash A$;
- $= \mathcal{M}, w \vDash \Diamond A$ if there exists v such that $wR^{\mathcal{M}}v$ and $\mathcal{M}, v \vDash A;$

If $\mathcal{M}, w \models A$ for all $w \in W$ we simply write $\mathcal{M} \models A$, and if $(W, R^{\mathcal{M}}, V) \models A$ for all valuations V based on frame $\mathcal{F} = (W, R^{\mathcal{M}})$, we simply write $\mathcal{F} \models A$.

2.2 Axiomatisations and Gödel-Löb logic

Turning to syntax, let us now build up the modal logics we are concerned with axiomatically, before relating them to the semantics just discussed. The modal logic K is axiomatised by all the theorems of classical propositional logic (CPL) together with axiom (k) and closed under the rules (mp) (modus ponens) and nec (necessitation) from Figure 2. The logic K4 is defined in the same way by further including the axiom (4) from Figure 2.

Referring back to the earlier semantics, the following characterisation is well-known [7]:

▶ Theorem 1 (K/K4 characterisation). $K \vdash A$ (resp. $K4 \vdash A$) iff A is satisfied in all relational frames (resp. with transitive accessibility relation).

The main subject matter of this work is an extension of K by the Löb axiom (löb), in Figure 2, a sort of induction principle:

▶ **Definition 2** (Gödel-Löb logic). GL *is defined by extending* K4 *by the Löb axiom* (löb) *and is closed under* (mp) *and* (nec).³

Semantically ($l\ddot{o}b$) says that, as long as worlds satisfy A whenever all its successors satisfy A, then A holds universally. This amounts to a 'reverse' induction principle for the accessibility relation. Indeed we have an associated characterisation for GL just like that for K4.

We call a frame is **terminating** if its accessibility relation has no infinite path.

² We parameterise the relations by the frame or models to be able distinguish it from the fixed binary relation symbol R.

 $^{^{3}}$ Alternatively, GL can be axiomatised by adding the (löb) axiom to K, as (4) can be derived from (löb).

$$(\mathsf{k}): \Box(A \to B) \to (\Box A \to \Box B) \qquad (\mathsf{mp}) \frac{A \to B \quad A}{B} \qquad (\mathsf{nec}) \frac{A}{\Box A}$$
$$(4): \Box A \to \Box \Box A \qquad (\mathsf{l\"ob}): \Box(\Box A \to A) \to \Box A$$

Figure 2 Some modal axioms and rules.

Theorem 3 (GL characterisation). $GL \vdash A$ iff all transitive terminating frames satisfy A.

2.3 Labelled calculi and the standard translation

The relational semantics of modal logic may be viewed as a bona fide fragment of predicate logic. Recalling the predicate language we fixed at the start of the section, the **standard translation** is defined as follows: for individual variables x and modal formulas A we define the predicate formula x : A by:

x:p is $p(x);$	$ = x : \diamond A \text{ is } \exists y (xRy \land (y : A)) ; $
$x : \perp$ is \perp ;	$ = x : \Box A \text{ is } \forall y(xRy \to (y:A)). $
$x:A\star B \text{ is } (x:A)\star (x:B) \text{ for } \star \in \{\lor,\land,\rightarrow\};$	

This induces a well-behaved proof theory as a fragment of usual first-order predicate systems [31], adaptable to many extensions under correspondence theorems such as Theorem 1.

A (labelled) sequent is an expression $\mathbf{R}, \Gamma \Rightarrow \Delta$, where \mathbf{R} is a set of relational atoms, i.e. formulas of form xRy, and Γ and Δ are multisets of labelled formulas, i.e. formulas of form x : A. We sometimes refer to the variable x in x : A as a label. We write Var(S) for the subset of labels/variables that occur in a given sequent S. Similarly so for $Var(\mathbf{R})$, etc.

Notationally, we have identified labelled formulas with the standard translation at the beginning of this section. This is entirely suggestive, as we can now easily distil systems for modal logics of interest by appealing to the sequent calculus LK for first-order predicate logic. In this vein, the labelled calculus ℓK for modal logic K is given in Figure 3. From here, under the aforementioned correspondence between K4 and transitive frames, we have a system $\ell K4$ for K4 that extends ℓK by the **transitivity rule**:

tr
$$\frac{\mathbf{R}, xRy, yRz, xRz, \Gamma \Rightarrow \Delta}{\mathbf{R}, xRy, yRz, \Gamma \Rightarrow \Delta}$$

For a labelled system S we write $S \vdash \mathbf{R}, \Gamma \Rightarrow \Delta$, if there is a proof of $\mathbf{R}, \Gamma \Rightarrow \Delta$ using the rules from S. We write $S \vdash x : A$, or even $S \vdash A$, to mean $S \vdash \emptyset \Rightarrow x : A$. Almost immediately from Theorem 1 and metatheorems for predicate logic, we have:

▶ **Proposition 4** (Soundness and completeness). $\ell K \vdash x : A$ (resp. $\ell K4 \vdash x : A$) iff $K \vdash A$ (resp. $K4 \vdash A$).

Naturally systems for many other modal logics can be readily obtained, when they correspond to simple frame properties, by adding further relational (structural) rules [31]. Indeed, let us call a labelled calculus *standard* if it does not extend ℓK by any new logical or (non-relational) structural rules, nor any new logical axioms. This terminology is suggestive, since it forces the left and right introduction rules for modalities to coincide with those induced by the *standard* translation. As remarked by Negri in [31], typical standard calculi cannot be complete GL, as termination of a relation is not even first-order definable. We shall sidestep this barrier in the next section by making use of *non-wellfounded* systems.

Identity and cut:

$$\mathsf{id}\,\frac{}{\mathbf{R},x:p\Rightarrow x:p}\qquad\mathsf{cut}\,\frac{\mathbf{R},\Gamma\Rightarrow\Delta,x:A\quad\mathbf{R},\Gamma',x:A\Rightarrow\Delta}{\mathbf{R},\Gamma,\Gamma'\Rightarrow\Delta,\Delta'}$$

Structural rules:

Relational structural rule:

$$\begin{split} & \mathsf{w}\text{-}l\,\frac{\mathbf{R},\Gamma\Rightarrow\Delta}{\mathbf{R},\Gamma,x:A\Rightarrow\Delta} & \mathsf{c}\text{-}l\,\frac{\mathbf{R},\Gamma,x:A,x:A\Rightarrow\Delta}{\mathbf{R},\Gamma,x:A\Rightarrow\Delta} & \mathsf{th}\,\frac{\mathbf{R},\Gamma\Rightarrow\Delta}{\mathbf{R},\mathbf{R}',\Gamma\Rightarrow\Delta} \\ & \mathsf{w}\text{-}r\,\frac{\mathbf{R},\Gamma\Rightarrow\Delta}{\mathbf{R},\Gamma\Rightarrow\Delta,x:A} & \mathsf{c}\text{-}r\,\frac{\mathbf{R},\Gamma\Rightarrow\Delta,x:A,x:A}{\mathbf{R},\Gamma\Rightarrow\Delta,x:A} \end{split}$$

Propositional logical rules:

$$\begin{split} & \perp -l \frac{\mathbf{R}, r : \perp, \Gamma \Rightarrow \Delta}{\mathbf{R}, r : \perp, \Gamma \Rightarrow \Delta} & (\text{no right rule for } \perp) \\ & \rightarrow -l \frac{\mathbf{R}, \Gamma \Rightarrow \Delta, x : A - \mathbf{R}, \Gamma', x : B \Rightarrow \Delta'}{\mathbf{R}, \Gamma, \Gamma', x : A \to B \Rightarrow \Delta, \Delta'} & \rightarrow -r \frac{\mathbf{R}, \Gamma, x : A \Rightarrow \Delta, x : B}{\mathbf{R}, \Gamma \Rightarrow \Delta, x : A \to B} \\ & \wedge -l \frac{\mathbf{R}, \Gamma, x : A_i \Rightarrow \Delta}{\mathbf{R}, \Gamma, x : A_0 \land A_1 \Rightarrow \Delta} i \in \{0, 1\} & \vee -l \frac{\mathbf{R}, \Gamma, x : A \Rightarrow \Delta - \mathbf{R}, \Gamma, x : B \Rightarrow \Delta}{\mathbf{R}, \Gamma, x : A \lor B \Rightarrow \Delta} \\ & \vee -r \frac{\mathbf{R}, \Gamma \Rightarrow \Delta, x : A_i}{\mathbf{R}, \Gamma \Rightarrow \Delta, x : A_0 \lor A_1} i \in \{0, 1\} & \wedge -r \frac{\mathbf{R}, \Gamma \Rightarrow \Delta, x : A - \mathbf{R}}{\mathbf{R}, \Gamma \Rightarrow \Delta, x : A \land B} \end{split}$$

Modal logical rules:

$$\diamond -l \, \frac{\mathbf{R}, xRy, \Gamma, y : A \Rightarrow \Delta}{\mathbf{R}, \Gamma, x : \diamond A \Rightarrow \Delta} \, y \text{ fresh} \quad \diamond -r \, \frac{\mathbf{R}, xRy, \Gamma \Rightarrow \Delta, y : A}{\mathbf{R}, xRy, \Gamma \Rightarrow \Delta, x : \diamond A} \\ \Box -r \, \frac{\mathbf{R}, xRy, \Gamma \Rightarrow \Delta, y : A}{\mathbf{R}, \Gamma \Rightarrow \Delta, x : \Box A} \, y \text{ fresh} \quad \Box -l \, \frac{\mathbf{R}, xRy, \Gamma, y : A \Rightarrow \Delta}{\mathbf{R}, xRy, \Gamma, x : \Box A \Rightarrow \Delta}$$

Figure 3 The standard labelled calculus ℓK for modal logic K.

3 Recovering a proof theoretic account for GL

In another branch of the proof theory literature, structural treatments of induction and well-foundedness have been developed in the guise of *non-wellfounded* and *cyclic* proofs, e.g. [32, 9, 3, 5, 39, 11]. Here non-wellfoundedness in proofs allows for inductive reasoning, and soundness is ensured by some global correctness condition. By incorporating these ideas into modal proof theory, one can design a non-wellfounded proof system for GL [37]. In this section we recover a *standard* labelled calculus for GL, in the sense of the preceding discussion. Our presentation is based on the correctness condition for *Cyclic Arithmetic* in [39, 11].

3.1 A standard calculus for GL, via non-wellfounded proofs

In what follows, we consider systems S that will typically be some fragment of the system $\ell\mathsf{K4}$ given earlier. The definitions we give apply to all 'non-wellfounded' systems of this work.

▶ **Definition 5** (Preproofs). A preproof in a system S is a possibly infinite derivation generated from the rules of S.

$$\begin{array}{c} \vdots \\ \operatorname{tr} \frac{zRz, x: \Box(\Box p \to p) \Rightarrow z: p}{xRy, x: \Box(\Box p \to p) \Rightarrow z: p} \bullet [z/y] \\ \to -l \frac{zRy, x: \Box(\Box p \to p) \Rightarrow y: \Box p}{xRy, x: \Box(\Box p \to p), y: \Box p \to p \Rightarrow y: p} \\ \Box -l \frac{zRy, x: \Box(\Box p \to p), x: \Box(\Box p \to p) \Rightarrow y: p}{zRy, x: \Box(\Box p \to p), x: \Box(\Box p \to p) \Rightarrow y: p} \bullet \\ \operatorname{c-l} \frac{xRy, x: \Box(\Box p \to p), x: \Box(\Box p \to p) \Rightarrow y: p}{zRy, x: \Box(\Box p \to p), x: \Box(\Box p \to p) \Rightarrow y: p} \bullet \\ \to -r \frac{xRy, x: \Box(\Box p \to p) \Rightarrow x: \Box p}{zRy, x: \Box(\Box p \to p) \Rightarrow y: p} \bullet \\ \to -r \frac{xRy, x: \Box(\Box p \to p) \Rightarrow y: p}{zRy, x: \Box(\Box p \to p) \Rightarrow y: p} \bullet \\ \to -r \frac{xRy, x: \Box(\Box p \to p) \Rightarrow y: p}{zRy, x: \Box(\Box p \to p) \Rightarrow y: p} \bullet \\ \end{array}$$

Figure 4 An ∞ -proof in ℓ K4 of Löb's axiom, left, and its contraposition, right. Identity steps on y: p above the $\rightarrow -l$ step, left, and the $\wedge -r$ step, right, are omitted for space considerations.

As preproofs may, in particular, be non-wellfounded, they may conclude fallacious theorems, so we require a correctness criterion. Appealing to the correspondence of GL over transitive terminating relations, we may directly import a 'trace' condition first used in [39].

▶ Definition 6 (Traces and proofs). Fix a preproof P and an infinite branch $(S_i)_{i<\omega}$. A trace along $(S_i)_{i<\omega}$ is a sequence of variables $(x_i)_{i<\omega}$ such that either: 1. $x_{i+1} = x_i$; or, 2. $x_i R x_{i+1}$ appears in S_i (then i is a **progress point** of $(x_i)_{i<\omega}$). A trace is **progressing** if it is not ultimately constant, i.e. if 2 above applies infinitely often. A branch $(S_i)_{i<\omega}$ is progressing if it has a progressing trace, and a preproof P is progressing, or simply is a ∞-**proof**, if each infinite branch has a progressing trace. We write $S \vdash^{\infty} \mathbf{R}, \Gamma \Rightarrow \Delta$ if there is a ∞-proof in S of the sequent $\mathbf{R}, \Gamma \Rightarrow \Delta$.

▶ **Example 7** (Löb and contra-Löb). An example of a ∞ -proof of (löb) in ℓ K4 is given in Figure 4, left. Here we have used bullets • to identify identical subproofs, up to the indicated renamings of variables. The preproof is indeed progressing: it has only one infinite branch, whose progress points are coloured red and trigger at each iteration of the •-loop. Notice that each sequent has only one formula on the right-hand side (RHS).

Another example of a ∞ -proof in ℓ K4 is the *contraposition* of (löb), given in Figure 4, right. We have merged several steps, and omitted some routine structural steps and initial sequents, a convention we shall continue to employ throughout this work. Again the preproof is indeed progressing, by the same argument as before. Notice, this time, that there are two formulas in the RHS of the premiss of \bullet . Indeed, by basic inspection of proof search, there is no cut-free ∞ -proof avoiding this feature.

▶ Remark 8 (On regularity). In non-wellfounded proof theory, special attention is often paid to the subset of *regular* preproofs, which may be written as finite (possibly cyclic) graphs, as in Example 7 above. Nonetheless these will play no role in the present work, as we are purely concerned with logical and proof theoretic investigations, not with effectivity.

The progress condition is invariant under expansion of relational contexts in a preproof: We may omit consideration of the thinning rule th (Figure 3) when reasoning about ∞ -proofs:

▶ **Observation 9.** th *is eliminable* ∞ *-proofs of* S.

3.2 Soundness and completeness

Let us now argue that our notion of ∞ -proof for $\ell K4$ is sound and complete for GL. The most interesting part is soundness, comprising a contradiction argument by infinite descent as is common in non-wellfounded proof theory, relying on the characterisation result of Theorem 3:

▶ **Proposition 10** (Soundness). *If* $\ell K4 \vdash^{\infty} x : A$ *then* $GL \vdash A$.

On the other hand, thanks to Example 7 and the known completeness of $\ell K4$ for K4, we can use cut-rules to derive:

▶ **Proposition 11** (Completeness). *If* $\mathsf{GL} \vdash A$ *then* $\ell \mathsf{K4} \vdash^{\infty} x : A$.

These results motivate the following notation:

▶ **Definition 12.** ℓ GL *is the class of* ∞ *-proofs of* ℓ K4.

Henceforth for a class of ∞ -proofs **P** we may write simply $\mathbf{P} \vdash S$ if **P** contains a ∞ -proof of the sequent S. For instance, writing $\ell \mathsf{GL} \vdash S$ is the same as $\ell \mathsf{K4} \vdash^{\infty} S$.

4 Recovering intuitionistic versions of GL from syntax and semantics

In this section we propose two intuitionistic versions of GL, in the style of Simpson, respectively by consideration of the proof theory and semantics of GL discussed in the previous sections. Later sections are then devoted to proving the equivalence of these two notions.

4.1 An intuitionistic GL, via syntax

Following Gentzen, it is natural to define intuitionistic calculi based on their classical counterparts by restricting sequents to one formula on the RHS. However, when implementing this restriction to different starting calculi for modal logic K (based on sequents, nested sequents, labelled sequents) one ends up with different intuitionistic variants of K (see e.g. [12]). The restriction of the ordinary sequent calculus for K defines a logic which is not compatible with the standard translation in the way we described for K. On the other hand labelled calculi are well designed for this purpose.

▶ **Definition 13.** The system ℓ IK (resp. ℓ IK4) is the restriction of ℓ K (resp. ℓ K4) to sequents in which exactly one formula occurs on the RHS.

Note that the rules w-r and c-r cannot be used in ℓ IK4, by the singleton restriction on the RHS of a sequent. In the style of Simpson [38], we can from here duly recover a standard intuitionistic analogue of GL, by restricting ℓ GL to ∞ -proofs with only singleton RHSs.

▶ **Definition 14.** We write ℓ IGL for the class of ∞-proofs of ℓ IK4.

▶ Example 15 (Löb, revisited). Recalling Example 7 earlier, note that the ∞-proof of Löb's axiom in Figure 4, left, indeed satisfies the singleton RHS restriction, and so ℓ IGL \vdash (löb). On the other hand its contraposition, right, does not satisfy this restriction. We will see shortly that it is indeed *not* a theorem of ℓ IGL.

4.2 An intuitionistic GL, via semantics

To give our semantic version of intuitionistic GL, we must first recall models of intuitionistic modal logic. Birelational semantics include models \mathcal{B} with two relations, the intuitionistic \leq and the modal $R^{\mathcal{B}}$ (parameterised with \mathcal{B} to distinguish from the fixed predicate symbol R used in this paper), which duly gives it the capacity to model intuitionistic modal logics.

▶ **Definition 16** (Birelational semantics). A *birelational frame* \mathcal{F} *is a triple* $(W, \leq, R^{\mathcal{F}})$, where W is a nonempty set of worlds equipped with a partial order \leq and an accessibility relation $R^{\mathcal{F}} \subset W \times W$. We require the following frame conditions:

(F1) If $w \leq w'$ and $wR^{\mathcal{F}}v$, then there exists v' such that $v \leq v'$ and $w'R^{\mathcal{F}}v'$.

(F2) If $wR^{\mathcal{F}}v$ and v < v', then there exists w' such that w < w' and $w'R^{\mathcal{F}}v'$.

A birelational model is a tuple $(W, \leq, R^{\mathcal{B}}, V)$, where $(W, \leq, R^{\mathcal{B}})$ is a birelational frame and V is a valuation $W \to \mathcal{P}(\mathsf{Pr})$ that is monotone in \leq , i.e., $w \leq w'$ implies $V(w) \subseteq V(w')$.

Let $\mathcal{B} = (W, \leq, \mathbb{R}^{\mathcal{B}}, V)$ be a birelational model. For worlds $w \in W$ and formulas A we define the satisfaction judgement $\mathcal{B}, w \vDash A$ as follows:

- $\blacksquare \mathcal{B}, w \vDash p \text{ if } p \in V(w);$ $\blacksquare \mathcal{B}, w \vDash A \land B \text{ if } \mathcal{B}, w \vDash A \text{ and } \mathcal{B}, w \vDash B;$ $\begin{array}{c} \mathcal{B}, w \nvDash \bot; \\ \mathcal{B}, w \vDash A \to B \text{ if for all } w' \ge w, \text{ if } \mathcal{B}, w \vDash A \lor B \text{ if } \mathcal{B}, w \vDash A \text{ or } \mathcal{B}, w \vDash B; \\ \end{array}$
- $= \mathcal{B}, w \models \Box A \text{ if for all } w' \ge w \text{ and for all } v \text{ such that } w' R^{\mathcal{B}} v \text{ we have } \mathcal{B}, v \models A;$
- $\blacksquare \mathcal{B}, w \vDash \Diamond A \text{ if there exists } v \text{ such that } w R^{\mathcal{B}} v \text{ and } \mathcal{B}, v \vDash A;$

We write $\mathcal{B} \vDash A$ if $\mathcal{B}, w \vDash A$ for all $w \in W$.

▶ Lemma 17 (Monotonicity lemma, [38]). Let $\mathcal{B} = (W, \leq, R^{\mathcal{B}}, V)$ be a birelational model. For any formula A and $w, w' \in W$, if $w \leq w'$ and $\mathcal{B}, w \models A$, then $\mathcal{B}, w' \models A$.

A soundness and completeness theorem is recovered in [38], similarly to the classical case (Proposition 4 and Theorem 1).

▶ Theorem 18 ([38]). ℓ |K $\vdash x : A$ (resp. ℓ |K4 $\vdash x : A$) iff A is satisfied in all birelational models (resp. with transitive accessibility relation)

We want to introduce a birelational counterpart of the transitive and terminating models of GL. The class of models that we will introduce here is different from the birelational semantics known for logic iGL [26, 1] which only require termination of the accessibility relation. Due to the global nature of \Box -evaluation in Definition 16, we here require the *composition* of \leq and $R^{\mathcal{B}}$ to be terminating.

- ▶ **Definition 19.** \mathscr{B} IGL is the class of birelational models $\mathcal{B} = (W, \leq, R^{\mathcal{B}}, V)$ such that:
- $\blacksquare R^{\mathcal{B}}$ is transitive: and,
- $= (\leq; R^{\mathcal{B}})$ is terminating, i.e., there are no infinite paths $x_1 \leq y_1 R^{\mathcal{B}} x_2 \leq y_2 R^{\mathcal{B}} x_3 \dots$

For a formula A, we write $\mathscr{B}\mathsf{IGL} \vDash A$ to mean that $\mathcal{B} \vDash A$ for all $\mathcal{B} \in \mathscr{B}\mathsf{IGL}$.

Example 20 (Contra-Löb, revisited). Recalling Examples 7 and 15 we indeed have that the contraposition of Löb's axiom, $\Diamond p \to \Diamond (p \land \Box \neg p)$, is not valid in $\mathscr{B}\mathsf{IGL}$. It is falsified at world w_1 of the following $\mathscr{B}\mathsf{IGL}$ -model \mathcal{B} where we assign p to all worlds. In the picture we omit transitive (and reflexive) edges of $R^{\mathcal{B}}$ (and \leq).

$$\begin{array}{c} v_1 \xrightarrow{v_1} v_2 \xrightarrow{v_2} v_3 \\ \leq \uparrow \qquad \stackrel{R^{\mathcal{B}}}{=} \leq \uparrow \qquad \stackrel{R^{\mathcal{B}}}{=} v_2 \\ w_1 \xrightarrow{w_2} w_2 \end{array}$$

5 Soundness

The main result of this section is the soundness of ℓIGL with respect to $\mathscr{B}IGL$ (Theorem 25 and Corollary 26). First we must extend the notion of satisfaction in birelational models to labelled sequents.

Let us fix a sequent $S = (\mathbf{R}, \Gamma \Rightarrow x : A)$ and a birelational model $\mathcal{B} = (W, \leq, R^{\mathcal{B}}, V)$ for the remainder of this section. An **interpretation** of S into \mathcal{B} is a function $\mathcal{I} : \mathsf{Var}(S) \to W$ such that $\mathcal{I}(x)R^{\mathcal{B}}\mathcal{I}(y)$ whenever $xRy \in \mathbf{R}$. We write

 $\mathcal{B}, \mathcal{I} \vDash \mathbf{R}, \Gamma \Rightarrow x : A \quad \text{if} \quad \mathcal{B}, \mathcal{I}(x) \vDash A \text{ when } \mathcal{B}, \mathcal{I}(y) \vDash B \text{ for all } y : B \in \Gamma.$

If $\mathcal{B}, \mathcal{I} \vDash S$ for all interpretations \mathcal{I} of S into \mathcal{B} , we simply write $\mathcal{B} \vDash S$, and if $\mathcal{B} \vDash S$ for all models $\mathcal{B} \in \mathscr{B}\mathsf{IGL}$, we simply write $\mathscr{B}\mathsf{IGL} \vDash S$. See Appendix B for a proof of the following.

▶ **Observation 21.** $\mathcal{B} \vDash \emptyset, x : B_1, \dots, x : B_n \Rightarrow x : A$ iff $\mathcal{B} \vDash (B_1 \land \dots \land B_n) \to A$.

In order to prove local soundness of $\ell IK4$ rules, we use a lifting lemma similarly to the one in [38] whose proof relies on the tree-like structure of **R**.

▶ Definition 22 ((Quasi-)tree-like). **R** is a tree if there is $x_0 \in Var(\mathbf{R})$ such that for each $x \in Var(\mathbf{R})$, $x \neq x_0$, there is a unique sequence $x_0Rx_1, x_1Rx_2, \ldots, x_mRx \in \mathbf{R}$. **R** is a quasi-tree if there is some $\mathbf{R}_0 \subseteq \mathbf{R} \subseteq \mathbf{R}_0^+$ where \mathbf{R}_0 is a tree and \mathbf{R}_0^+ denotes the transitive closure of \mathbf{R}_0 . Sequent S is (quasi-)tree-like if either $\mathbf{R} = \emptyset$ and $Var(\Gamma) \subseteq \{x\}$, or **R** is a (resp., quasi-)tree and $Var(\Gamma) \cup \{x\} \subseteq Var(\mathbf{R})$.

▶ Lemma 23 (Lifting lemma). Suppose S is quasi-tree-like. Let \mathcal{I} be interpretation of S into \mathcal{B} , $x \in Var(S)$ and $w \geq \mathcal{I}(x)$. There is an interpretation \mathcal{I}' of S into \mathcal{B} such that $\mathcal{I}'(x) = w$ and for all $y \in Var(S)$ we have $\mathcal{I}'(y) \geq \mathcal{I}(y)$.

Let us employ some conventions on ∞ -proofs. Note that for every inference rule of $\ell |\mathsf{K4}$, except for th and cut, the premiss(es) are quasi-tree-like whenever the conclusion is. By Observation 9 we shall duly assume that th is not used. Note that this forces relational contexts to be *growing*, bottom-up: the relational context of a sequent always contains those below it. If a cut-formula z : C has label z that does not occur in the conclusion, we may safely rename z to a variable that does. Thus we may assume that any ∞ -proof in $\ell |\mathsf{IGL}|$ of x : A has only quasi-tree-like sequents in it.

▶ Proposition 24 (Local soundness). Suppose S is quasi-tree-like Let I be an interpretation of S into B such that $\mathcal{B}, \mathcal{I} \nvDash S$. For any inference step of $\ell \mathsf{IK4} \setminus \{\mathsf{th}\}$ that S concludes, there is a premiss S' and an interpretation I' of S' into B such that $\mathcal{B}, \mathcal{I} \nvDash S'$ and $\mathcal{I}'(z) \ge \mathcal{I}(z)$ for all $z \in \mathsf{Var}(S)$.

The proof uses some direct calculations in most cases and the lifting lemma (Lemma 23) to handle rules $\rightarrow -r$ and \Box -r. We give some cases in Appendix B. From here, as in the classical setting for ℓ GL, we can employ a contradiction argument by infinite descent to conclude:

- ▶ Theorem 25 (Soundness). Suppose S is quasi-tree-like. Then $\ell \mathsf{IGL} \vdash S$ implies $\mathscr{B} \mathsf{IGL} \models S$.
- ▶ Corollary 26. If $\ell | \mathsf{GL} \vdash x : A \text{ then } \mathscr{B} | \mathsf{GL} \models A$.

The remainder of this work is devoted to proving the converse result. The sections that follow structure the proof into the three parts according to the arrows indicated in Figure 1.

6 From birelational models to Kripke predicate models

Towards our countermodel construction in the next section we turn to *predicate models*, essentially via the standard translation, whose internal structure is richer than that of birelational models, thus providing useful invariants for the sequel.

▶ Definition 27 (Predicate models). A Kripke structure is a tuple

- \blacksquare W is a non-empty set of worlds partially ordered by \leq ;
- $= \{D_w\}_{w \in W}$ is a family of non-empty **domains**, such that $D_w \subseteq D_{w'}$ whenever $w \leq w'$;
- = $\{\mathsf{Pr}_w\}_{w\in W}$ is a family of mappings $\mathsf{Pr}_w : \mathsf{Pr} \to \mathcal{P}(D_w)$ such that for each $p \in \mathsf{Pr}$, $\mathsf{Pr}_w(p) \subseteq \mathsf{Pr}_{w'}(p)$ whenever $w \leq w'$;
- $= \{R_w\}_{w \in W} \text{ is a family of relations } R_w \subseteq D_w \times D_w \text{ such that } R_w \subseteq R_{w'} \text{ whenever } w \leq w'.$

▶ **Definition 28** (Environment). Let \mathcal{K} be a Kripke structure. A w-environment is a function ρ : Var $\rightarrow D_w$.

Note that a *w*-environment is also a w'-environment for any $w' \ge w$.

▶ **Definition 29** (Satisfaction). For modal formula A, structure \mathcal{K} and w-environment ρ , we inductively define the judgement $\mathcal{K}, w \models^{\rho} x : A$ as follows, where $\rho[x := d]$ is the map that sends variable x to d and agrees with ρ on all other variables:

 $\begin{aligned} &= \mathcal{K}, w \models^{\rho} x : p \text{ if } \rho(x) \in \mathsf{Pr}_{w}(p); \\ &= \mathcal{K}, w \nvDash^{\rho} \bot; \\ &= \mathcal{K}, w \models^{\rho} x : A \land B \text{ if } \mathcal{K}, w \models^{\rho} x : A \text{ and } \mathcal{K}, w \models^{\rho} x : B; \\ &= \mathcal{K}, w \models^{\rho} x : A \lor B \text{ if } \mathcal{K}, w \models^{\rho} x : A \text{ or } \mathcal{K}, w \models^{\rho} x : B; \\ &= \mathcal{K}, w \models^{\rho} x : A \to B \text{ if for all } w' \ge w, \text{ if } \mathcal{K}, w' \models^{\rho} x : A \text{ then } \mathcal{K}, w' \models^{\rho} x : B; \\ &= \mathcal{K}, w \models^{\rho} x : \Box A \text{ if for all } w' \ge w \text{ and } d \in D_{w'}, \text{ if } \rho(x) R_{w'} d, \text{ then } \mathcal{K}, w' \models^{\rho[y:=d]} y : A; \end{aligned}$

 $= \mathcal{K}, w \models^{\rho} x : \Diamond A \text{ if there exists } d \in D_w \text{ such that } \rho(x) R_w d \text{ and } \mathcal{K}, w \models^{\rho[y:=d]} y : A.$

We write $\mathcal{K} \vDash x : A$ if $\mathcal{K}, w \vDash^{\rho} x : A$ for all worlds w and w-environments ρ .

The monotonicity lemma also holds in Kripke structures.

▶ Lemma 30 (Monotonicity lemma). Let \mathcal{K} be a Kripke structure. If $w \leq w'$ and $\mathcal{K}, w \models^{\rho} x : A$, then $\mathcal{K}, w' \models^{\rho} x : A$.

To capture transitivity in Kripke structures, it is sufficient to require each R_w to be transitive, which can be considered as a local condition on Kripke structures. However interpreting termination requires us to consider the interactions between \leq and R_w similarly to the previous section.

Let \mathcal{K} be a Kripke structure. We write D_W for the set of ordered pairs of the form (w, d)with $w \in W$ and $d \in D_w$. We define the two binary relations $\leq_{D_W}, R_{D_W} \subseteq D_W \times D_W$ as:

■ $(w, d) \leq_{D_W} (w', d')$ iff $w \leq w'$ and d = d'; ■ $(w, d)R_{D_W}(w', d')$ iff w = w' and $dR_w d'$.

▶ Definition 31. Write *P*IGL for the class of Kripke structures K satisfying the following:

- for all $w \in W$, R_w is transitive; and
- relation $(\leq_{D_W}; R_{D_W})$ is terminating, i.e., there are no infinite paths $(w_1, d_1) \leq_{D_W} (w_2, d_1) R_{D_W}(w_2, d_2) \leq_{D_W} (w_3, d_2) R_{D_W}(w_3, d_3) \dots$

We write $\mathscr{P}\mathsf{IGL} \vDash A$ to mean that $\mathcal{K} \vDash x : A$ for all $\mathcal{K} \in \mathscr{P}\mathsf{IGL}$ and any $x \in \mathsf{Var}$.

The same construction converting a predicate structure into a birelational model from [38, Section 8.1.1] can be used to prove the following result.

▶ **Proposition 32.** *If* \mathscr{B} IGL $\vDash A$ *then* \mathscr{P} IGL $\vDash A$.

7 Completeness of a multi-succedent calculus via determinacy

Towards completeness we perform a countermodel construction using a proof search strategy based on an intuitionistic *multi-succedent* calculus. This is inspired by analogous arguments for intuitionistic predicate logic, e.g. in [41], but adapted to a non-wellfounded setting.

▶ Definition 33 (Multi-succedent intuitionistic calculus). The system $m\ell IK4$ is the restriction of $\ell K4$ where the \Box -right and \rightarrow -right rules must have exactly one formula on the RHS. We write $m\ell IGL$ for the class of ∞ -proofs of $m\ell IK4$.

The remainder of this section is devoted to proving the following completeness result:

▶ Theorem 34. If \mathscr{P} IGL $\vDash A$ then m ℓ IGL $\vdash x : A$.

Here we informally describe the construction of the proof search tree. For a more formal treatment of parts below we also refer to Appendix D. During bottom-up proof search we will always proceed according to the three following phases in order of priority: applications of rule tr, applications of invertible rules (other than tr), and application of non-invertible rules. The first two together we call the **invertible phase**, the other the **non-invertible phase**. In fact, invertibility of all rules except \Box -r and \rightarrow -r is guaranteed by applying suitable contractions at the same time. For instance, we apply the following derivable 'macro' rules for \rightarrow on the left and \diamond on the right:

$$\rightarrow -l \, \frac{\mathbf{R}, \Gamma, x: A \rightarrow B \Rightarrow \Delta, x: A \quad \mathbf{R}, \Gamma, x: A \rightarrow B, x: B \Rightarrow \Delta}{\mathbf{R}, \Gamma, x: A \rightarrow B \Rightarrow \Delta} \quad \diamondsuit r \, \frac{\mathbf{R}, xRy, \Gamma \Rightarrow \Delta, x: \diamondsuit A, y: A}{\mathbf{R}, xRy, \Gamma \Rightarrow \Delta, x: \diamondsuit A}$$

By furthermore building weakening into the identity, i.e. allowing initial sequents of form $\mathbf{R}, \Gamma, x : p \Rightarrow \Delta, x : p$, structural rules become redundant for proof search. In the invertible phase we can apply the rules in any order. The non-invertible phase creates predecessor nodes for each possible rule instance of $\rightarrow -r$ and $\Box -r$.

In order to carry out our countermodel construction to show completeness of $\mathfrak{m}\ell \mathsf{IGL}$, we rely on two features of the proof search space that digress from usual countermodel constructions in intuitionistic predicate logic [41]. The first important feature of this strategy is that the invertible phase is always finite and ends in so-called *saturated sequents*, i.e., sequents for which any bottom-up rule application (other than id and \perp -l) yields a premiss that is the same sequent, up to multiplicities (see Appendix D for formal definition). Looking ahead to the countermodel, worlds w are defined on the basis of invertible phases and will as a result all have a finite domain D_w .

▶ Lemma 35. Following the proof search strategy described above, each invertible phase constructs a finite subtree that has saturated sequents at its leaves.

Secondly, and perhaps more importantly, we employ a technique from non-wellfounded proof theory to help us organise the countermodel constructed from a failed proof search: we appeal to determinacy of a *proof search game*. To understand the motivation here, a classical countermodel-from-failed-proof search argument proceeds (very roughly) as follows:

(1) assume a formula is not provable; (2) for each rule instance there must be an unprovable premiss; (3) continue in this way to construct an (infinite) 'unprovable' branch; (4) extract a countermodel from this branch. In our setting we will need the branch obtained through the process above to be *not progressing* in order to deduce that the structure we extract is indeed one of $\mathscr{P}IGL$. However the local nature of the process above does not at all guarantee that this will be the case. For this, we rely on *(lightface) analytic determinacy*, which is equivalent to the existence of 0^{\sharp} over ZFC [20], of the corresponding proof search game. As we only use it as a tool and it is not the main focus of our work, we refer to Appendix D for more details.

▶ **Proposition 36.** Given an unprovable sequent S there is a subtree T of the proof search space rooted at S, closed under bottom-up non-invertible rule application⁴ such that each infinite branch of T is not progressing.

The properties in Lemma 35 and Proposition 36 enable us to construct a countermodel:

▶ **Theorem 37** (Countermodel construction). If $m\ell IGL \not\vdash \mathbf{R}, \Gamma \Rightarrow \Delta$, then there is a structure $\mathcal{K} \in \mathscr{P}IGL$ with w-environment ρ such that for labelled formulas x : A we have

■ *if* $x : A \in \Gamma$, *then* $\mathcal{K}, w \models^{\rho} x : A$, *and*, ■ *if* $x : A \in \Delta$, *then* $\mathcal{K}, w \nvDash^{\rho} x : A$.

From here Theorem 34 easily follows. Notice that we did not use rule cut in the proof search strategy so we can actually conclude a stronger result:

▶ Corollary 38. mℓIGL is cut-free complete over 𝒫IGL.

8 Completeness of *l*IGL via (partial) cut-elimination

To obtain completeness of ℓ IGL for \mathscr{B} IGL, we will simulate $m\ell$ IGL using cuts in an extension of ℓ IGL that allows reasoning over disjunctions of labelled formulas. We apply a (partial) cut-elimination procedure to eliminate these disjunctions, and then note that any resulting proof is already one of ℓ IGL.

We shall use metavariables φ, ψ etc. to vary over *disjunctions* of labelled formulas. I.e. $\varphi, \psi, \ldots ::= (x : A) \mid \varphi \lor \psi$.

▶ **Definition 39.** The system $\forall \ell \mathsf{IK4}$ is the extension of $\ell \mathsf{IK4}$ by duly adapting identity, cut, structural and \forall rules to allow for φ -formulas. In particular it has the following \forall rules:

$$\vee -l \, \frac{\mathbf{R}, \Gamma, \varphi_0 \Rightarrow \psi \quad \mathbf{R}, \Gamma, \varphi_1 \Rightarrow \psi}{\mathbf{R}, \Gamma, \varphi_0 \lor \varphi_1 \Rightarrow \psi} \qquad \vee -r \, \frac{\mathbf{R}, \Gamma \Rightarrow \varphi_i}{\mathbf{R}, \Gamma \Rightarrow \varphi_0 \lor \varphi_1} \, i \in \{0, 1\}$$

The **degree** of a formula $(x_1 : A_1) \lor \cdots \lor (x_d : A_d)$ is d. The degree of a cut is the degree d of its cut-formula, in which case we say it is a d-cut. The degree of a preproof is the maximum degree of its cuts (when this is well-defined).

▶ **Proposition 40.** If m ℓ IK4 has a cut-free ∞ -proof of x : A, $\forall \ell$ IK4 has one of bounded degree.

Moreover, immediately from definitions, we have:

▶ **Observation 41.** $A \lor \ell I K4 \infty$ -proof containing only labelled formulas is a $\ell I K4 \infty$ -proof.

⁴ I.e. if $S_0 \in T$ concludes some non-invertible step with premiss S_1 , then also $S_1 \in T$.

Thus, to conclude completeness of ℓIGL for $\mathscr{B}IGL$, it suffices to eliminate the use of disjunctions of labelled formulas in $\lor \ell IK4 \infty$ -proofs. We will prove this by a partial cutelimination procedure, reducing cuts over disjunctions of labelled formulas until they are on labelled formulas.

For the remainder of this section we work only with preproofs without thinning th, by Observation 9. Recall that this means that relational contexts are *growing*, bottom-up. For a sequent S write \mathbf{R}_S for its relational context. I.e. if S is $\mathbf{R}, \Gamma \Rightarrow \varphi$, then $\mathbf{R}_S := \mathbf{R}$.

▶ Lemma 42 (Invertibility). If $\forall \ell \mathsf{IK4}$ has a ∞ -proof P of $\mathbf{R}, \Gamma, \varphi_0 \lor \varphi_1 \Rightarrow \psi$ then it also has ∞ -proofs P_i of $\mathbf{R}, \Gamma, \varphi_i \Rightarrow \psi$, for $i \in \{0, 1\}$. Moreover, for each branch $(S_i)_{i < \omega}$ of P_i there is a branch $(S'_i)_{i < \omega}$ of P such that $\mathbf{R}_{S_i} \subseteq \mathbf{R}_{S'_i}$ for all $i < \omega$.

From here the key cut-reduction for φ -formulas is simply,

$$\underset{\mathsf{cut}}{\overset{\vee}{\mathbf{R},\Gamma\Rightarrow\chi_{i}}} \underbrace{\frac{\mathbf{R},\Gamma\Rightarrow\chi_{i}}{\mathbf{R},\Gamma\Rightarrow\chi_{0}\vee\chi_{1}}}_{\mathbf{R},\Gamma,\Gamma'\Rightarrow\varphi} \underbrace{\overset{Q}{\mathbf{Q}}}_{\mathbf{R},\Gamma,\chi_{0}\vee\chi_{1}\Rightarrow\varphi} \qquad \qquad \underset{\mathsf{cut}}{\overset{\varphi}{\mathbf{R},\Gamma\Rightarrow\chi_{i}}} \underbrace{\frac{\mathbf{R},\Gamma\Rightarrow\chi_{i}}{\mathbf{R},\Gamma',\chi_{i}\Rightarrow\varphi}}_{\mathbf{R},\Gamma,\Gamma'\Rightarrow\varphi}$$

where Q_i is obtained by Lemma 42 above. Note that this reduction 'produces' a cut of lower complexity. Commutative cut-reduction cases, where the cut-formula is not principal on the left, are standard and always produce. Note that, thanks to invertibility, we do not consider commutations over the right premiss of a φ -cut. However commutative cases may *increase* the heights of progress points; for instance when commuting over a \Box -l step,

$$\begin{array}{c} \square -l \\ \overbrace{\mathbf{R}, xRy, \Gamma, y : A \Rightarrow \chi} & \overbrace{\mathbf{R}, xRy, \Gamma', \chi \Rightarrow \varphi} \\ \mathsf{cut} \\ \overbrace{\mathbf{R}, xRy, \Gamma, x : \square A \Rightarrow \chi} & \overbrace{\mathbf{R}, xRy, \Gamma', \chi \Rightarrow \varphi} \\ \overbrace{\mathbf{R}, xRy, \Gamma, \Gamma', x : \square A \Rightarrow \varphi} & \sim \end{array} \\ \begin{array}{c} \mathsf{cut} \\ \overbrace{\mathbf{R}, xRy, \Gamma, y : A \Rightarrow \chi} & \overbrace{\mathbf{R}, xRy, \Gamma', \chi \Rightarrow \varphi} \\ \square -l \\ \overbrace{\mathbf{R}, xRy, \Gamma, \Gamma', x : \square A \Rightarrow \varphi} \end{array} \\ \end{array}$$

observe that progress points in Q have been raised. Thus, to show that the limit of cutreduction is progressing we will need appropriate invariants, requiring additional notions.

A **bar** of a preproof is a (necessarily finite, by König's Lemma) antichain intersecting each infinite branch. Each bar *B* induces a (necessarily finite) subtree $\lfloor B \rfloor$ of nodes beneath (and including) it. Given a cut-reduction $P \rightsquigarrow_r P'$ we associate to each bar *B* of *P* a bar r(B)of *P'* in the natural way. In particular we have that r(B) satisfies the following properties:

▶ Lemma 43 (Trace preservation). If $P \sim_{\mathsf{r}} P'$ and B a bar of P, for any sequent $S' \in \mathsf{r}(B)$ there is a sequent $S \in B$ such that $\mathbf{R}_{S'} \supseteq \mathbf{R}_S$.

This lemma allows us to keep track of a fixed amount of progress information during cutelimination, sidestepping the issue that commutative cases raise progress points. Note that we really need the $\mathbf{R}_{S'} \supseteq \mathbf{R}_S$ due to the $\diamondsuit l$ commutative case:

$$\diamond -l \frac{\mathbf{R}, xRy, \Gamma, y : A \Rightarrow \chi}{\operatorname{cut} \frac{\mathbf{R}, \Gamma, x : \diamond A \Rightarrow \chi}{\mathbf{R}, \Gamma, \Gamma', x : \diamond A \Rightarrow \varphi}} \sum_{\mathbf{R}, \Gamma', \chi \Rightarrow \varphi} \operatorname{cut} \frac{\mathbf{R}, xRy, \Gamma, y : A \Rightarrow \chi}{\operatorname{cut} \frac{\mathbf{R}, xRy, \Gamma, y : A \Rightarrow \chi}{\mathbf{R}, \Gamma, \Gamma', y : A \Rightarrow \varphi}} \sum_{\mathbf{R}, \Gamma, \Gamma', x : \diamond A \Rightarrow \varphi}$$

where the preproof xRy, Q is obtained from Q by prepending xRy to the LHS of each sequent. From here we can effectively reduce infinitary cut-elimination to finitary cut-elimination, by eliminating cuts beneath higher and higher bars. The step case is given by:

▶ Lemma 44 (Productivity). For any ∞-proof P and bar B there is a sequence of cutreductions $P = P_0 \rightsquigarrow_{r_1} \cdots \rightsquigarrow_{r_n} P_n$ such that $r_n \cdots r_1(B)$ has no d-cuts beneath it in P_n .

The argument proceeds in a relatively standard way: by induction on the number of d-cuts in $\lfloor B \rfloor$, with a subinduction on the multiset of the distances of topmost d-cuts in $\lfloor B \rfloor$ from B, where the distance is the length of the shortest path from the cut to B. Applying Lemma 44 to higher and higher bars allows us to reduce cut-degrees as required:

▶ **Proposition 45** (Degree-reduction). For each ∞-proof of $\forall \ell IK4$ of x : A of degree d, there is one of degree < d.

Importantly here, the preservation of progress in the limit crucially relies on Lemma 43, under König's Lemma. Finally by induction on degree we have:

▶ Corollary 46 (Partial cut-elimination). If $\forall \ell IK4$ has a bounded-degree ∞ -proof of x : A, then it has one containing only labelled formulas.

From here we have our desired converse to Theorem 25, following from Observation 41 and Propositions 32 and 40, Theorem 34 and Corollary 46:

▶ **Theorem 47** (Completeness). If \mathscr{B} IGL $\vDash A$ then ℓ IGL $\vdash x : A$.

9 Conclusions

We have recovered several intuitionistic formulations of GL, both syntactically and semantically, in the tradition of Simpson, on intuitionistic modal logic [38], and Simpson, on non-wellfounded proofs [39]. We proved the equivalence of all these formulations, cf. Figure 1, motivating the following definition:

▶ Definition 48. IGL is the modal logic given by any/all of the nodes of Figure 1.

Thanks to the methodology we followed, IGL satisfies Simpson's *requirements* from [38]. IGL also interprets (classical) GL along the Gödel-Gentzen negative translation.

It would be interesting to examine IGL as a logic of provability, returning to the origins of GL. In particular IGL (even IK) has a *normal* \diamond , distributing over \lor , so let us point out that it is not sound to interpret \diamond as consistency $\neg \Box \neg$. At the same time (effective) model-theoretic readings of the \diamond stumble on the consequence (already of IK) $\Box(A \rightarrow B) \rightarrow \diamond A \rightarrow \diamond B$. We expect IGL to rather correspond to the provability logic of some *model* of HA.

To this end it would be pertinent to develop a bona fide *axiomatisation* of IGL. As far as we know it is possible that IGL is simply the extension of IK4 by Löb's axiom but, similarly to other 'I' logics, we expect that further axioms involving \diamond will be necessary. On a related note, we believe that a full cut-elimination result holds for ℓ IGL, in particular by extending our argument for $\vee \ell$ IGL. Crucial here is that no cut-reductions *delete* progress points, since we do not cut on relational atoms (cf. a key \Box reduction). A full development of this is beyond the scope (and allocated space) of this work.

— References

- 1 Mohammad Ardeshir and Motjaba Mojtahedi. The Σ_1 -provability logic of HA. Annals of Pure and Applied Logic, 169(10):997–1043, 2018.
- 2 Arnold Avron. On modal systems having arithmetical interpretations. *The Journal of Symbolic Logic*, 49(3):935–942, 1984.
- 3 David Baelde, Amina Doumane, and Alexis Saurin. Infinitary proof theory: the multiplicative additive case. In Jean-Marc Talbot and Laurent Regnier, editors, 25th EACSL Annual Conference on Computer Science Logic, CSL 2016, August 29 - September 1, 2016, Marseille, France, volume 62 of LIPIcs, pages 42:1–42:17. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2016.
- 4 Gianluigi Bellin, Valeria de Paiva, and Eike Ritter. Extended Curry-Howard correspondence for a basic constructive modal logic. In *Proceedings of methods for modalities*, volume 2, 2001.
- 5 Stefano Berardi and Makoto Tatsuta. Equivalence of inductive definitions and cyclic proofs under arithmetic. In 32nd Annual ACM/IEEE Symposium on Logic in Computer Science LICS, pages 1–12. IEEE, 2017.
- 6 Lars Birkedal, Rasmus Ejlers Møgelberg, Jan Schwinghammer, and Kristian Støvring. First steps in synthetic guarded domain theory: step-indexing in the topos of trees. Logical Methods in Computer Science, 8, 2012.
- 7 Patrick Blackburn, Maarten De Rijke, and Yde Venema. *Modal logic*, volume 53. Cambridge University Press, 2001.
- 8 Milan Božić and Kosta Došen. Models for normal intuitionistic modal logics. Studia Logica, 43(3):217–245, 1984.
- 9 James Brotherston and Alex Simpson. Sequent calculi for induction and infinite descent. Journal of Logic and Computation, 21(6):1177–1216, 2011.
- 10 Ranald Clouston and Rajeev Goré. Sequent calculus in the topos of trees. In Foundations of Software Science and Computation Structures: 18th International Conference, FOSSACS 2015, pages 133–147. Springer, 2015.
- 11 Anupam Das. On the logical complexity of cyclic arithmetic. Logical Methods in Computer Science, 16, 2020
- 12 Anupam Das and Sonia Marin. Brouwer meets Kripke: constructivising modal logic, 2022. Post on *The Proof Theory Blog* (accessed 2 August 2023). https://prooftheory.blog/2022/ 08/19/brouwer-meets-kripke-constructivising-modal-logic/
- 13 Anupam Das and Sonia Marin. On intuitionistic diamonds (and lack thereof), 2023. Accepted to TABLEAUX '23.
- 14 Anupam Das and Damien Pous. Non-wellfounded proof theory for (Kleene+action)(algebras+lattices). In Dan R. Ghica and Achim Jung, editors, 27th EACSL Annual Conference on Computer Science Logic, CSL 2018, September 4-7, 2018, Birmingham, UK, volume 119 of LIPIcs, pages 19:1–19:18. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2018.
- 15 Gisèle Fischer Servi. On modal logic with an intuitionistic base. Studia Logica, 36:141–149, 1977.
- 16 Iris van der Giessen, Uniform Interpolation and Admissible Rules. Proof-theoretic investigations into (intuitionistic) modal logics. PhD thesis, Utrecht University, 2022.
- 17 Iris van der Giessen and Rosalie Iemhoff. Sequent calculi for intuitionistic Gödel–Löb logic. Notre Dame Journal of Formal Logic, 62(2):221–246, 2021.
- 18 Rajeev Goré and Revantha Ramanayake. Valentini's cut-elimination for provability logic resolved. In Carlos Areces and Robert Goldblatt, editors, *Proceedings of the 7th conference on Advances in Modal Logic*, pages 67–86. College Publications, 2008.
- 19 Rajeev Goré and Ian Shillito. Direct elimination of additive-cuts in GL4ip: verified and extracted. In Advances in Modal Logic 14, papers from the fourteenth conference on "Advances in Modal Logic", 22-26 August 2022, 2022.

- 20 Leo Harrington. Analytic determinacy and 0[#]. The Journal of Symbolic Logic, 43(4):685–693, 1978.
- 21 Rosalie Iemhoff. Reasoning in circles. In Jan van Eijck, Joost J. Joosten, and Rosalie Iemhoff, editors, *Liber Amicorum Alberti. A Tribute to Albert Visser*, pages 165–178. College Publications, 2016
- 22 Rosalie Iemhoff. Terminating sequent calculi for two intuitionistic modal logics. Journal of Logic and Computation, 28(7):1701–1712, 2018.
- 23 G. Alex Kavvos. Intensionality, intensional recursion and the Gödel-Löb axiom. FLAP, 8(8):2287–2312, 2021.
- 24 Daniel Leivant. On the proof theory of the modal logic for arithmetic provability. *The Journal* of *Symbolic Logic*, 46(3):531–538, 1981.
- 25 Daniel Leivant. Absoluteness of Intuitionistic Logic. PhD thesis, University of Amsterdam, 1975.
- 26 Tadeusz Litak. Constructive modalities with provability smack. In Guram Bezhanishvili, editor, Leo Esakia on Duality in Modal and Intuitionistic Logics, volume 4 of Outstanding Contributions to Logic, pages 187–216. Springer Netherlands, 2014.
- 27 Michael Mendler and Valeria de Paiva. Constructive CK for contexts. *Context Representation and Reasoning (CRR-2005)*, 13, 2005.
- 28 Anders Moen. The proposed algorithms for eliminating cuts in the provability calculus GLS do not terminate. In *The 13th Nordic Workshop in Programming Theory*, 2001.
- 29 Mojtaba Mojtahedi: On provability logic of HA. Eprint 2206.00445, arXiv (2022). https: //doi.org/10.48550/arXiv.2206.00445
- 30 Hiroshi Nakano. A modality for recursion. In Proceedings Fifteenth Annual IEEE Symposium on Logic in Computer Science (Cat. No. 99CB36332), pages 255–266. IEEE, 2000.
- 31 Sara Negri. Proof analysis in modal logic. Journal of Philosophical Logic, 34:507–544, 2005
- 32 Damian Niwiński and Igor Walukiewicz. Games for the μ-calculus. Theoretical Computer Science, 163(1-2):99–116, 1996.
- 33 Gordon Plotkin and Colin Stirling. A framework for intuitionistic modal logics. In Proceedings of the 1st Conference on Theoretical Aspects of Reasoning about Knowledge (TARK), pages 399–406, 1986.
- 34 Francesca Poggiolesi. A purely syntactic and cut-free sequent calculus for the modal logic of provability. The Review of Symbolic Logic, 2(4):593–611, 2009.
- 35 Katsumi Sasaki. Löb's axiom and cut-elimination theorem. Academia Mathematical Sciences and Information Engineering Nanzan University, 1:91–98, 2001.
- 36 Krister Segerberg. Results in non-classical propositional logic. Uppsala: Filosofiska Föreningen och Filosofiska Institutionen vid Uppsala Universitet, 1971.
- 37 Daniyar S. Shamkanov. Circular proofs for the Gödel-Löb provability logic. Mathematical Notes, 96:575–585, 2014.
- 38 Alex K. Simpson. The Proof Theory and Semantics of Intuitionistic Modal Logic. PhD thesis, University of Edinburgh, 1994.
- 39 Alex K. Simpson. Cyclic arithmetic is equivalent to Peano arithmetic. In Foundations of Software Science and Computation Structures Proceedings, volume 10203 of Lecture Notes in Computer Science, pages 283–300, 2017.
- 40 Robert M. Solovay. Provability interpretations of modal logic. Israel Journal of Mathematics, 25:287–304, 1976.
- 41 Gaisi Takeuti. *Proof Theory*. New York, N.Y., U.S.A.: Sole distributors for the U.S.A. and Canada, Elsevier Science Pub. Co., 1975.
- 42 Aldo Ursini. A modal calculus analogous to K4W, based on intuitionistic propositional logic, I°. Studia Logica, 38(3):297–311, 1979.
- 43 Silvio Valentini. The modal logic of provability: cut-elimination. Journal of Philosophical logic, pages 471–476, 1983.

44 Frank Wolter and Michael Zakharyaschev. On the relation between intuitionistic and classical modal logics. Algebra and Logic, 36:73–92, 1997.

A Appendix for Section 3

Proof of Observation 9. Delete every thinning step and replace each relational context \mathbf{R} with the union of all relational contexts beneath it.

Proof of Proposition 10. First, given a model $\mathcal{M} = (W, R^{\mathcal{M}}, V)$, an $(\mathcal{M}$ -)assignment for a sequent $S = \mathbf{R}, \Gamma \Rightarrow \Delta$ is a function $\rho : \mathsf{Var}(S) \to W$ such that whenever xRy occurs in \mathbf{R} we have $\rho(x)R^{\mathcal{M}}\rho(y)$. We write $\mathcal{M}, \rho \models S$ if: $\mathcal{M}, \rho(x) \models A$ for some x : A in Δ , whenever $\mathcal{M}, \rho(y) \nvDash B$ for all y : B in Γ .

Suppose otherwise. Let π be a ∞ -proof of x : A in $\ell \mathsf{K4}$, and let $\mathcal{M} = (W, R^{\mathcal{M}}, V)$ be a structure in which $R^{\mathcal{M}}$ is transitive yet $\mathcal{M}, w \nvDash A$. We shall show that $R^{\mathcal{M}}$ is not terminating on W.

Set $\rho_0(x) = w$. By local soundness of the rules, construct an infinite branch $(S_i)_{i < \omega}$ and assignments $\rho_i : \operatorname{Var}(S_i) \to W$ such that $\mathcal{M}, \rho_i \nvDash S_i$ always satisfying: $\rho_i(z) = \rho_{i+1}(z)$, whenever $z \in \operatorname{Var}(S_i) \cap \operatorname{Var}(S_{i+1})$.

Now, let $(x_i)_{i < \omega}$ be a progressing trace along $(S_i)_{i < \omega}$. By inspection of the rules we have that, at any progress point x_i , necessarily $\rho_i(x_i) R^{\mathcal{M}} \rho_{i+1}(x_{i+1})$, otherwise $x_i = x_{i+1}$ and so $\rho_i(x_i) = \rho_{i+1}(x_{i+1})$ by above. Thus $(x_i)_{i < \omega}$ induces an infinite path along $R^{\mathcal{M}}$, and so $R^{\mathcal{M}}$ is not terminating.

Proof of Proposition 11. It is well-known that ℓ K4 is complete for K4, and so derives all its axioms [31]. Together with the derivation of Löb's axiom in Example 7, completeness follows by closure of ℓ GL under necessitation (simply by \Box -r) and modus ponens (by cut).

B Appendix for Section 5

Proof of Observation 21. For left to right, suppose $v \ge w$ such that $\mathcal{B}, v \models \bigwedge_i B_i$. Then for $\mathcal{I}(x) = v$ we have $\mathcal{B}, \mathcal{I}(x) \models B_i$ for all $i \le n$. So by assumption, $\mathcal{B}, v \models A$ as desired. For the other direction, let \mathcal{I} be an interpretation, for which its domain only consists of x. Let $\mathcal{B}, \mathcal{I}(x) \models B_i$ for all $i \le n$. Then, $\mathcal{B}, \mathcal{I}(x) \models \bigwedge_i B_i$ and so by assumption $\mathcal{B}, \mathcal{I}(x) \models A$.

Proof of Lemma 23. In our proof we rely on the lifting lemma in [38] (Lemma 8.1.3), where S is assumed to be a tree-like sequent. We rely on the fact that each quasi-tree \mathbf{R} is supported by a unique tree \mathbf{R}' , i.e., \mathbf{R}' is a tree and $\mathsf{Var}(\mathbf{R}') = \mathsf{Var}(\mathbf{R})$. Consider sequent $S = (\mathbf{R}', \Gamma \Rightarrow x : A)$. Note that \mathcal{I} is also an interpretation of S', so we can now apply Lemma 8.1.3 from [38] to obtain an interpretation \mathcal{I}' of S' such that $\mathcal{I}'(x) = w$ and for all $y \in \mathsf{Var}(S)$ we have $\mathcal{I}'(y) \geq \mathcal{I}(y)$. Since \mathcal{B} is transitive, \mathcal{I}' is also an interpretation for S, which concludes the proof.

Proof of Proposition 24. Let us show the proofs for the rules $\rightarrow -r, \rightarrow -l, \Box -l, \Box -r, \diamond -l, \diamond -r$. w-l, and tr. For $\rightarrow -r, \Box -r$ we use the lifting lemma (Lemma 23).

For rule $\rightarrow -r$, by assumption, $\mathcal{B}, \mathcal{I} \nvDash \mathbf{R}, \Gamma \Rightarrow x : A \rightarrow B$. In other words, $\mathcal{B}, \mathcal{I}(y) \vDash D$ for all $y : D \in \Gamma$ and $\mathcal{B}, \mathcal{I}(x) \nvDash A \rightarrow B$. So there exists $w \ge \mathcal{I}(x)$ such that $\mathcal{B}, w \vDash A$ and $\mathcal{B}, w \nvDash B$. By the lifting lemma (Lemma 23), there is an interpretation \mathcal{I}' of S such that $\mathcal{I}'(x) = w$ and $\mathcal{I}'(z) \ge \mathcal{I}(z)$ for all $z \in \mathsf{Var}(S)$. By monotonicity (Lemma 17) we have that $\mathcal{B}, \mathcal{I}'(y) \vDash D$ for all $y : D \in \Gamma$. Also $\mathcal{B}, \mathcal{I}'(x) \vDash A$ and $\mathcal{B}, \mathcal{I}'(x) \nvDash B$. Therefore, $\mathcal{B}, \mathcal{I}' \nvDash \mathbf{R}, \Gamma, x : A \Rightarrow x : B$ as desired.

For rule $\rightarrow -l$, by assumption $\mathcal{B}, \mathcal{I} \nvDash \mathbf{R}, \Gamma_1, \Gamma_2, x : A \rightarrow B \Rightarrow z : C$. So, $\mathcal{B}, \mathcal{I}(y) \vDash D$ for all $y : D \in \Gamma_1 \cup \Gamma_2, \mathcal{B}, \mathcal{I}(x) \vDash A \rightarrow B$, and $\mathcal{B}, \mathcal{I}(z) \nvDash C$. Either $\mathcal{B}, \mathcal{I}(x) \vDash A$ or $\mathcal{B}, \mathcal{I}(x) \nvDash A$. In the latter case, $\mathcal{B}, \mathcal{I} \nvDash \mathbf{R}, \Gamma_1 \Rightarrow x : A$. In the former case, since $\mathcal{B}, \mathcal{I}(x) \vDash A \rightarrow B$, also $\mathcal{B}, \mathcal{I}(x) \vDash x : B$, and hence $\mathcal{B}, \mathcal{I} \nvDash \mathbf{R}, \Gamma_2, x : B \Rightarrow z : C$.

Consider rule \Box -l. By assumption we have that $\mathcal{B}, \mathcal{I} \nvDash \mathbf{R}, xRy, \Gamma, x : \Box A \Rightarrow z : C$. As \mathcal{I} is an interpretation we have $\mathcal{I}(x)R^{\mathcal{B}}\mathcal{I}(y)$. From $\mathcal{B}, \mathcal{I}(x) \vDash \Box A$ it follows that $\mathcal{B}, \mathcal{I}(y) \vDash A$. Hence $\mathcal{B}, \mathcal{I} \nvDash \mathbf{R}, xRy, \Gamma, y : A \Rightarrow z : C$.

For rule \Box -r, we have $\mathcal{B}, \mathcal{I} \nvDash \mathbf{R}, \Gamma \Rightarrow x : \Box A$. So there exist worlds w and v such that $\mathcal{I}(x) \leq wR^{\mathcal{B}}v$ and $\mathcal{B}, v \nvDash A$. By the lifting lemma we know that there is an interpretation \mathcal{I}' of S satisfying $\mathcal{I}'(x) = w$ and $\mathcal{I}'(z) \geq \mathcal{I}(z)$ for all $z \in \mathsf{Var}(S)$. Let us define an interpretation \mathcal{J} of premises $\mathbf{R}, xRy, \Gamma \Rightarrow y : A$ with fresh label y, that agrees with \mathcal{I}' on all labels from $\mathbf{R} \cup \{x\}$, and $\mathcal{J}(y) = v$. This is a well-defined interpretation by the observation that $\mathcal{J}(x)R^{\mathcal{B}}\mathcal{J}(y)$. By monotonicity (Lemma 17), we have $\mathcal{B}, \mathcal{J}' \nvDash \mathbf{R}, xRy, \Gamma \Rightarrow y : A$.

For rule \diamond -*l*, we have $\mathcal{B}, \mathcal{I} \nvDash \mathbf{R}, \Gamma, x : \diamond A \Rightarrow z : C$. So there exists a *v* such that $\mathcal{I}(x)R^{\mathcal{B}}v$ and $\mathcal{B}, v \vDash A$. We define a new interpretation \mathcal{J} of tree $\mathbf{R} \cup \{xRy\}$ with fresh label *y*, that agrees with \mathcal{I} on all labels from \mathbf{R} , and $\mathcal{J}(y) = v$. We have $\mathcal{B}, \mathcal{J} \nvDash \mathbf{R}, xRy, \Gamma, y : A \Rightarrow z : C$.

For rule \diamond -*r* we have $\mathcal{B}, \mathcal{I} \nvDash \mathbf{R}, xRy, \Gamma \Rightarrow x : \diamond A$. Since, $\mathcal{I}(x)R^{\mathcal{B}}\mathcal{I}(y)$ and $\mathcal{B}, \mathcal{I}(x) \nvDash \diamond A$, we have $\mathcal{B}, \mathcal{I} \nvDash \mathbf{R}, xRy, \Gamma \Rightarrow y : A$.

For rule w-l we have $\mathcal{B}, \mathcal{I} \nvDash \mathbf{R}, \Gamma, x : A \Rightarrow z : C$. Since sequent $\mathbf{R}, \Gamma, x : A \Rightarrow z : C$ is assumed to be quasi-tree-like, $\mathbf{R}, \Gamma \Rightarrow z : C$ will contain the same labels. So \mathcal{I} will be an interpretation for $\mathbf{R}, \Gamma \Rightarrow \Delta$ and $\mathcal{B}, \mathcal{I} \nvDash \mathbf{R}, \Gamma \Rightarrow z : C$.

Finally, let us consider rule tr. We have $\mathcal{B}, \mathcal{I} \nvDash \mathbf{R}, xRy, yRz, \Gamma \Rightarrow z : C$. It follows immediately from transitivity of \mathcal{B} that $\mathcal{B}, \mathcal{I} \nvDash \mathbf{R}, xRy, yRz, xRz, \Gamma \Rightarrow z : C$.

Proof of Theorem 25. We proceed by contradiction and we use the fact that rule th is admissible in ℓ IGL. Suppose that ℓ IGL $\vdash \mathbf{R}, \Gamma \Rightarrow x : A$, but suppose that $\mathcal{B}, \mathcal{I} \nvDash \mathbf{R}, \Gamma \Rightarrow x : A$ for some model $\mathcal{B} \in \mathscr{B}$ IGL with interpretation \mathcal{I} of \mathbf{R} into \mathcal{B} . Let π be the ∞ -proof of $\mathbf{R}, \Gamma \Rightarrow x : A$ in ℓ IGL without the use of the thinning rule th. By the local soundness of the rules (Proposition 24), we can construct a (possibly) infinite sequence of labelled sequents $(\mathbf{R}_i, \Gamma_i \Rightarrow x_i : A_i)_i$ and interpretations \mathcal{I}_i of \mathbf{R}_i into \mathcal{B} such that

- 1. $(\mathbf{R}_0, \Gamma_0 \Rightarrow x_0 : A_0) = (\mathbf{R}, \Gamma \Rightarrow x : A)$ and $\mathcal{I}_0 = \mathcal{I}$;
- 2. $\mathbf{R}_{i+1}, \Gamma_{i+1} \Rightarrow x_{i+1} : A_{i+1}$ is a premise of the rule in π with conclusion $\mathbf{R}_i, \Gamma_i \Rightarrow x_i : A_i$;
- **3.** $\mathcal{B}, \mathcal{I}_i \nvDash \mathbf{R}_i; \Gamma_i \Rightarrow x_i : A_i;$ and
- **4.** $\mathcal{I}_{i+1}(z) \geq \mathcal{I}_i(z)$ for all $z \in \text{Dom}(\mathcal{I}_i)$.

If the sequence is finite, we end in a sequent \mathbf{R}_n ; $\Gamma_n \Rightarrow x_n : A_n$ derived by id or $\perp l$ which is true in every model by definition, deriving a contradiction with $\mathcal{B}, \mathcal{I}_n \nvDash \mathbf{R}_n; \Gamma_n \Rightarrow x_n : A_n$. If the sequence is infinite, it corresponds to an infinite branch in π which has a progressing trace $(y_i)_{i>k}$ by definition. So,

■ $y_i = y_{i+1}$ in which case $\mathcal{I}_i(y_i) \leq \mathcal{I}_{i+1}(y_{i+1})$ by construction, or; ■ $y_i R y_{i+1}$ appears in \mathbf{R}_i , which means that $\mathcal{I}_i(y_i) R^{\mathcal{B}} \mathcal{I}_i(y_{i+1}) \leq \mathcal{I}_{i+1}(y_{i+1})$.

The second property occurs infinitely often, which creates a sequence of worlds $(\mathcal{I}_i(y_i))_{i \geq k}$ in model \mathcal{B} which is increasing in the order $(\leq; R^{\mathcal{B}})$, which contradicts the fact that $(\leq; R^{\mathcal{B}})$ is terminating.

C Appendix for Section 6

Proof of Proposition 32. Let $\mathcal{K} \in \mathscr{P}$ IGL. Following the construction from Section 8.1.1 in [38], we convert \mathcal{K} into a birelational model $\mathcal{B}_{\mathcal{K}} = (W', \leq', R', V')$ as follow where unary predicate symbol $p \in \mathsf{Pr}$ is considered to be a propositional variable:

$$W' = D_W;$$

$$\leq' = \leq_{D_W};$$

$$R' = R_{D_W};$$

$$V'((w,d)) = \{p \mid d \in \mathsf{Pr}_w(p)\}.$$

Indeed, \leq' is a partial order and V' is monotone. Also the frame properties (F1) and (F2) are easy to check [38]. Model $\mathcal{B}_{\mathcal{K}}$ is in \mathscr{B} IGL, because R' is transitive as each R_w is transitive by definition of \mathcal{K} and therefore R_{D_W} is transitive. Moreover, $(\leq'; R')$ is terminating simply because $(\leq_{D_W}; R_{D_w})$ is required to be terminating.

Let $w \in W$ and let ρ be a *w*-environment. By a straightforward induction on A we can prove that $\mathcal{K}, w \models^{\rho[x:=d]} x : A$ if and only if $\mathcal{B}_{\mathcal{K}}, (w, d) \models A$. Therefore, since $\mathscr{B}\mathsf{IGL} \models A$ we have that $\mathcal{B}_{\mathcal{K}}, (w, d) \models A$ for all $(w, d) \in D_W$ and so $\mathcal{K}, w \models^{\rho[x:=d]} x : A$ for all $(w, d) \in D_W$. Hence $\mathcal{K}, w \models^{\rho} x : A$.

D Appendix for Section 7

We start this appendix by setting up the proof search strategy in a more formal manner than explained in the main text.

▶ Definition 49. A rule of the form

$$\frac{S_1 \quad \dots \quad S_n}{S}$$

is **invertible**, if for every instance of the rule, for all *i*, the instance of S_i is provable whenever the instance of *S* is provable.

One can show by coinduction that all rules with built-in contraction and built-in weakening, except $\rightarrow -r$ and \Box -r, are invertible.

In order to structure the proof search strategy we define saturated sequents.

▶ Definition 50 (Saturation). Let $\mathbf{R}, \Gamma \Rightarrow \Delta$ be a sequent that contains labelled formula x : A. We say that x : A is saturated in $\mathbf{R}, \Gamma \Rightarrow \Delta$ if the following conditions hold based on the form of A:

- A equals \perp or p for some $p \in \mathsf{Pr}$;
- if $A = A_1 \land A_2$ and $x : A \in \Gamma$, then $x : A_1 \in \Gamma$ and $x : A_2 \in \Gamma$;
- if $A = A_1 \land A_2$ and $x : A \in \Delta$, then $x : A_1 \in \Delta$ or $x : A_2 \in \Delta$;
- if $A = A_1 \lor A_2$ and $x : A \in \Gamma$, then $x : A_1 \in \Gamma$ or $x : A_2 \in \Gamma$;
- if $A = A_1 \lor A_2$ and $x : A \in \Delta$, then $x : A_1 \in \Delta$ and $x : A_2 \in \Delta$;
- if $A = A_1 \rightarrow A_2$ and $x : A \in \Gamma$, then $x : A_1 \in \Delta$ or $x : A_2 \in \Gamma$;
- if $A = \Box A_1$ and $x : A \in \Gamma$, then for all y with $xRy \in \mathbf{R}$, $y : A_1 \in \Gamma$;
- if $A = \Diamond A_1$ and $x : A \in \Gamma$, there exists y s.t. $xRy \in \mathbf{R}$ and $y : A_1 \in \Gamma$;
- if $A = \Diamond A_1$ and $x : A \in \Delta$, then for all y with $xRy \in \mathbf{R}$, $y : A_1 \in \Delta$.

A sequent $\mathbf{R}; \Gamma \Rightarrow \Delta$ is saturated if all its formulas are saturated and \mathbf{R} is transitively closed.

Now we define the proof search strategy. During bottom-up proof search we will always proceed according to the three following phases in order of priority: applications of rule tr, applications of invertible rules other than tr, and application of non-invertible rules. The first two together we call the invertible phase, the other the non-invertible phase. More precisely, when possible, we apply rule tr. If not, if possible, we apply another invertible rule for which the principal formula is not saturated and create predecessor node(s) labelled by the premiss(s). We will argue below that each invertible phase is finite and ends in saturated sequents. When $\mathbf{R}, \Gamma \Rightarrow \Delta$ is saturated, we create predecessor nodes for each possible rule instance of $\rightarrow -r$ and $\Box -r$, i.e., for each $x : A \rightarrow B \in \Delta$ we create predecessor $\mathbf{R}, \Gamma, x : A \Rightarrow x : B$, and for each $x : \Box A \in \Delta$ we create predecessor node $\mathbf{R}, xRy, \Gamma \Rightarrow y : A$, where y is fresh.

Now we prove Lemma 35 stated in the main text, i.e., we show that given a proof search tree as defined above, each invertible phase is a finite subtree with saturated sequents in its leaves.

Proof of Lemma 35. By construction of the invertible phase, each leaf is saturated. Let $\mathbf{R}, \Gamma \Rightarrow \Delta$ be the start sequent of the invertible phase. We argue that the invertible phase terminates. First observe that, bottom-up, the number of labelled formulas in a sequent increase for each rule because we only apply a rule in the invertible phase when the concerned principal formulas is not saturated. To ensure termination, we show that the number of labelled formulas of each sequent in the invertible phase is bounded.

Let $\mathbf{R}', \Gamma' \Rightarrow \Delta'$ be a sequent in the invertible phase. We first show that the number of different labels in Γ' is bounded. A new label in Γ' can only be created via the rules \Box -l and \diamond -l. Note that rule \rightarrow -r can also create a new label in Γ' , but this rule is not part of the invertible phase. Therefore, each new formula in Γ' is a subformula of another formula in Γ' . As there are only finitely many subformulas, and we only apply the rules to non-saturated sequents, one can only apply rules \Box -l and \diamond -l finitely many times. Hence, there are only finitely many extra labels in Γ' .

Now for each label x, again by saturation and the fact that new formulas in the sequent are subformulas of the sequent, one can only create finitely many formulas labelled by x. Since the number of labels is bounded, the number of formulas occurring in $\mathbf{R}', \Gamma' \Rightarrow \Delta'$ is bounded.

Now we set up the proof of Proposition 36 using techniques from non-wellfounded proof theory. As mentioned in the main text, a classical countermodel-from-failed-proof search argument proceeds (very roughly) as follows: (1) Assume a formula is not provable, (2) For each rule instance there must be an unprovable premiss, (3) Continue in this way to construct an (infinite) 'unprovable' branch, (4) Extract a countermodel from this branch. In our setting we will need the branch obtained through the process above to be *not progressing* in order to deduce that the structure we extract is indeed one of \mathscr{P} IGL. However the local nature of the process above does not at all guarantee that this will be the case.

In the setting of classical GL, the invertibility of the rules of ℓGL allow us to nonetheless carry out a naïve such argument. For the intuitionistic setting, where invertibe and non-invertible rules may interact in a nontrivial way, we must use more powerful machinery.

▶ Definition 51 (Proof search game). For a system S, the proof search game is a game between two players, Prover (P) and Denier (D), played as follows:

- \blacksquare The game initialises on a sequent S of S.
- \blacksquare P chooses a rule instance of S that has S as its conclusion.
- D chooses a premiss of that rules instance, and they repeat the process.

A **play** of the game is a maximal sequence of moves as described above. An infinite play of the game is **won** by P (and **lost** by D) just if it has a progressing thread. (If deadlock is reached, the player with no valid move loses.)

It is not hard to see that P-strategies from S are *just* S-preproofs of S, and that *winning* P-strategies from S are just S-proofs of S. What is more difficult is to show that winning D-strategies induce countermodels; this is what we do in the next subsection.

Before that, we better state the following:

▶ **Proposition 52** (Determinacy, $\exists 0^{\#}$). The proof search game for S is determined.

This is a consequence of *analytic determinacy* as the progressing condition is indeed (lightface) Σ_1^1 : "there *exists* an infinite thread that progresses infinitely often" [20].

One can view the proof search space of $m\ell IGL$ as a collection of plays of the proof search game (Definition 51), where each branch represents a play. We use determinacy of the game to extract a 'failed branch' and construct the countermodel from that.

Proof of Proposition 36. If there is no proof of sequent $S = \mathbf{R}, \Gamma \Rightarrow \Delta$, we know by determinacy of the game (Proposition 52) that there is a winning strategy for Denier (D) for $\mathbf{R}, \Gamma \Rightarrow \Delta$. So, when navigating the proof search space from bottom-up, whenever we must make a choice at a branching rule in an invertible phase, we can simply follow the choice given by the winning strategy for D. Starting from the bottom sequent, we perform this strategy and for each non-invertible branching that we encounter, we continue the same process for each branch. As we follow the winning strategy of D, no infinite branch can be progressing.

Proof of Theorem 37. We rely on the properties from Lemma 35 and Proposition 36 for $S = (\mathbf{R}, \Gamma \Rightarrow \Delta)$. By Lemma 35, the subtree *T* found in Proposition 36 consists of countably many finite segments of the form $S_1 \dots S_n$ where S_1 is the start sequent or premises of a non-invertible rule, and S_n is saturated. We assume that the segments are indexed from index set *I*.

To define the countermodel, we use the fact that for each segment *i* of the form $S_1^i ldots S_{n_i}^i$, we know that all its sequents are contained in $S_n^i = \mathbf{R}_i, \Gamma_i \Rightarrow \Delta_i$. So for each such segment *i*, consider $S_{n_i}^i$. We define the structure $\mathcal{K} = (W, \leq, \{D_w\}_{w \in W}, \{\mathsf{Pr}_w\}_{w \in W}, \{R_w\}_{w \in W})$ as follows:

$$\begin{split} W &= \{w_i \mid i \in I\};\\ w_i \leq w_j & \text{iff } i = j \text{ or } \mathbf{R}_i, \Gamma_i \Rightarrow \Delta_i \text{ becomes below } \mathbf{R}_j; \Gamma_j \Rightarrow \Delta_j \text{ in the proof search tree};\\ D_{w_i} &= \mathsf{Var}(\mathbf{R}_i);\\ xR_{w_i}y & \text{iff } xRy \in \mathbf{R}_i;\\ \mathsf{Pr}_{w_i}(p) &= \{x \in \mathsf{Var}(\mathbf{R}_i) \mid x : p \in \Gamma_i\}. \end{split}$$

It remains to show that \mathcal{K} is in $\mathscr{P}\mathsf{IGL}$ and that it satisfies the conditions from the theorem.

We first prove that \mathcal{K} is an intuitionistic structure. By construction of the rules in the countermodel construction, for each segments *i* that is below a segment *j* we have $\mathbf{R}_i \subseteq \mathbf{R}_j$ and $\Gamma_i \subseteq \Gamma_j$. Therefore, for each $w_i \leq w_j$, it holds that $D_{w_i} \subseteq D_{w_j}$, $\mathsf{Pr}_{w_i}(p) \subseteq \mathsf{Pr}_{w_j}(p)$ for each $p \in \mathsf{Pr}$, and $R_{w_i} \subseteq R_{w_j}$. Also note that \leq is a partial order.

Each R_{w_i} is transitive because sequents $\mathbf{R}_i, \Gamma_i \Rightarrow \Delta_i$ are saturated by construction. In addition, relation $(\leq_{D_W}; R_{D_W})$ as defined in Section 6 is terminating as proved as follows. Suppose we have an infinite branch in the model:

$$(w^1, y^1) \leq_{D_W} (w^2, y^1) R_{D_W}(w^2, y^2) \leq_{D_W} (w^3, y^2) R_{D_W}(w^3, y^3) \dots$$

This means that there is an infinite branch in the strategy of D, say $(\mathbf{R}^k, \Gamma^k \Rightarrow x^k : A^k)_k$ such that $y^k R y^{k+1} \in \mathbf{R}^k$ for infinitely many k. But this means that this branch has a progressing trace, which cannot be the case in the game strategy of D, a contradiction. Therefore, $(\leq_{D_W}; R_{D_W})$ is terminating, and therefore $\mathcal{K} \in \mathscr{P}\mathsf{IGL}$.

Now we show that for all w_i -environments ρ_{w_i} with $\rho_{w_i}(x) = x$ for $x \in Var(\mathbf{R}_i)$, we have for all labelled formulas x : A and all $i \in I$:

if $x : A \in \Gamma_i$, then $\mathcal{K}, w_i \models^{\rho_{w_i}} x : A$, and, if $x : A \in \Delta_i$, then $\mathcal{K}, w_i \nvDash^{\rho_{w_i}} x : A$.

We do so by induction on A. We show some cases and leave the remaining cases to the reader.

For A = p, if $x : p \in \Gamma_i$, then $\mathcal{K}, w_i \models^{\rho_{w_i}} x : p$ by definition of the structure and definition of ρ .

If $x : p \in \Delta_i$ we observe that $x : p \notin \Gamma_i$, otherwise sequent $\mathbf{R}_i, \Gamma_i \Rightarrow \Delta_i$ would be provable which cannot be the case for a sequent in D's strategy. Hence, $\mathcal{K}, w_i \nvDash^{\rho_{w_i}} x : p$.

- For $A = A_1 \wedge A_2$, if $x : A_1 \wedge A_2 \in \Gamma_i$, then by saturation, $x : A_1 \in \Gamma_i$ and $x : A_2 \in \Gamma_i$. By induction hypothesis we have $\mathcal{K}, w_i \models^{\rho_{w_i}} x : A_1$ and $\mathcal{K}, w_i \models^{\rho_{w_i}} x : A_2$. Hence $\mathcal{K}, w_i \models^{\rho_{w_i}} x : A_1 \wedge A_2$. Similar reasoning applies when $x : A_1 \wedge A_2 \in \Delta_i$.
- For $A = A_1 \to A_2$, if $x : A_1 \to A_2 \in \Gamma_i$, let $w_j \ge w_i$ such that $\mathcal{K}, w_j \models^{\rho_{w_i}} x : A_1$. By the proof search, $\Gamma_i \subseteq \Gamma_j$, therefore $x : A_1 \to A_2 \in \Gamma_j$. By saturation we have $x : A_1 \in \Delta_j$ or $x : A_2 \in \Gamma_j$. The first cannot occur as by induction hypothesis we would have $\mathcal{K}, w_j \nvDash^{\rho_{w_j}} x : A_1$ which is in contradiction to the assumption. (Note that $x : A_1$ has no other free variable than x and indeed $\rho_{w_i}(x) = x = \rho_{w_j}(x)$.) By applying the induction hypothesis to the second we obtain $\mathcal{K}, w_j \vDash^{\rho_{w_i}} x : A_2$ as desired.

If $x : A_1 \to A_2 \in \Delta_i$, then for an immediate segment j above segment i in the proof search tree we have $x : A_1 \in \Gamma_j$ and $x : A_2 \in \Delta_j$. By induction hypothesis we have $\mathcal{K}, w_j \models^{\rho_{w_j}} x : A_1$ and $\mathcal{K}, w_j \nvDash^{\rho_{w_j}} x : A_2$ (and by similar reasoning from above $\mathcal{K}, w_j \models^{\rho_{w_i}} x : A_1$ and $\mathcal{K}, w_j \nvDash^{\rho_{w_i}} x : A_2$). So, since $w_i \leq w_j$, we have $\mathcal{K}, w_i \nvDash^{\rho_{w_i}} x : A_1 \to A_2$.

For $A = \Box A'$, if $x : \Box A' \in \Gamma_i$, let $w_j \ge w_i$ and let $y \in D_{w_j} = \mathsf{Var}(\mathbf{R}_j)$. Suppose $R_{w_j}(\rho_{w_i}(x), y)$. Note that $\rho_{w_i}(x) = x$, and thus $xRy \in \mathbf{R}_j$. Note that $x : \Box A' \in \Gamma_j$. By saturation, $y : A' \in \Gamma_j$. By induction hypothesis we conclude $\mathcal{K}, w_j \models^{\rho_{w_j}} y : A'$. Since $\rho_{w_i}(y) = y$, we conclude $\mathcal{K}, w_j \models^{\rho_{w_i}[y:=y]} y : A'$.

If $x : \Box A' \in \Delta_i$, then for an immediate segment j above i in the proof search tree we have $xRy \in \mathbf{R}_j$ and $y : A' \in \Delta_j$. So by induction hypothesis $\mathcal{K}, w_j \not\models^{\rho_{w_j}} y : A'$. Since $w_i \leq w_j$ and $xR_{w_i}y$, and since $\rho_{w_i}(y) = y$ we have $\mathcal{K}, w_i \not\models^{\rho_{w_i}[y:=y]} y : \Box A'$.

◀

E Appendix for Section 8

Proof of Proposition 40. Replace the RHS of any $m\ell IK4 \propto$ -proof by the disjunction of its labelled formulas (without repetitions), and locally simulate any right step by deriving

the corresponding implication and cutting against it. Since cuts are inserted locally on only logical formulas, not interfering with relational atoms, the resulting preproof is indeed progressing.

For Lemmas 43 and 44, we describe r(B) along with the cut-reduction cases in the next subsection, whence the arguments follow by inspection of the cases.

Proof of Lemma 42. Replace $\varphi_0 \lor \varphi_1$ and all its direct ancestors by φ_i . The only critical cases are when it is principal, in which case simply take the *i*-premiss (and choose the corresponding branch). Formally, the argument is structured as a coinduction on P.

Proof of Proposition 45. Starting with a ∞ -proof P, write B_1, B_2, \ldots for its bars of heights $1, 2, \ldots$ respectively. Define a sequence of cut-reductions $P_0 \sim_{\vec{r}_1}^* P_1 \sim_{\vec{r}_2}^* \cdots$ as follows:

$$\blacksquare$$
 Set $P_0 := P$.

 \vec{r}_{n+1} is obtained by applying Lemma 43 to P_n and $\vec{r}_n \cdots \vec{r}_1(B_{n+1})$.

So, e.g., \vec{r}_1 is obtained by applying Lemma 43 to P_0 and B_1 ; \vec{r}_2 is obtained by applying Lemma 43 to P_1 and $\vec{r}_1(B_2)$; and so on. Since cut-reductions leave nodes beneath untouched, this process must approach a limit preproof, say P_{ω} .

We must now show that P_{ω} is progressing. Consider an infinite branch $(S_i)_{i<\omega}$ of P_{ω} and write $\mathbf{R} := \bigcup_{i<\omega} \mathbf{R}_{S_i}$. Notice that, by Lemma 43, \mathbf{R} must contain the relational contexts of arbitrarily high sequents from P. Specifically, the maximal prefix of $(S_i)_{i<\omega}$ consistent with P_{n+1} ends at $\vec{r}_n \cdots \vec{r}_1(B_{n+1})$ and so at a sequent whose relational context contains that of some sequent in B_{n+1} , i.e. at height n, by Lemma 43.

Finally, since relational contexts are growing, bottom-up, notice that the set of sequents of P whose relational contexts are contained in \mathbf{R} forms a subtree T of P. Since, as just discussed, T contains arbitrarily high nodes of P, it is infinite and so has an infinite branch by König's Lemma. Thus T has a progressing trace and so by definition \mathbf{R} has an infinite path. Again by definition, this means $(S_i)_{i < \omega}$ has a progressing trace.

E.1 Cut-reduction cases and r(B)

E.1.1 Key case

This is the key cut-reduction on φ -formulas again:



where Q_i is obtained by Lemma 42 above.

Here, if B intersects P and Q before reduction then r(B) consists of the corresponding points after reduction. If B intersects the premiss of the cut or \vee -r before reduction then r(B) consists of the conclusion of P after reduction, as well as the corresponding part of Q before the reduction. Note in all these cases that the degree below the bar has decreased. If B is below (or including) the conclusion of the cut before reduction, then r(B) is the corresponding bar after reduction, i.e. the cut-reduction leaves the bar unchanged.

The definition of r(B) is similar for all the commutative cases below, always designed to decrease the distance of a topmost *d*-cut from the bar.

E.1.2 **Commutative cases**

This is the commutation over a $\rightarrow -l$ step,

$$\rightarrow -l \frac{\overbrace{\mathbf{R}, \Gamma \Rightarrow x : A \quad \mathbf{R}, \Gamma', x : B \Rightarrow \chi}^{P}}{\operatorname{cut} \frac{\mathbf{R}, \Gamma, \Gamma', x : A \Rightarrow B \Rightarrow \chi}{\mathbf{R}, \Gamma, \Gamma', \Gamma'', x : A \Rightarrow B \Rightarrow \varphi} \xrightarrow{R} \rightarrow -l \frac{\overbrace{\mathbf{R}, \Gamma \Rightarrow x : A}^{P} \operatorname{cut} \frac{\mathbf{R}, \Gamma', x : B \Rightarrow \chi}{\mathbf{R}, \Gamma', T'', x : B \Rightarrow \chi} \xrightarrow{R} (\Gamma'', \chi) \xrightarrow{R} (\Gamma', \chi) \xrightarrow{R}$$

This is the commutation over a \lor -l step,

$$\rightarrow -l \frac{\mathbf{R}, \Gamma \Rightarrow x : A}{\mathbf{R}, \Gamma, \Gamma', \Gamma'', x : B \Rightarrow \chi} \frac{\mathbf{R}, \Gamma', x : B \Rightarrow \chi}{\mathbf{R}, \Gamma', \Gamma'', \chi \Rightarrow \varphi} \frac{\mathbf{R}, \Gamma', \chi \Rightarrow \varphi}{\mathbf{R}, \Gamma, \Gamma', \Gamma'', x : B \Rightarrow \varphi}$$

$$\operatorname{cut} \frac{\mathbf{R}, \Gamma, \psi_{0} \Rightarrow \chi}{\vee -l} \frac{\mathbf{R}, \Gamma', \chi \Rightarrow \varphi}{\mathbf{R}, \Gamma', \psi_{0} \Rightarrow \varphi} \operatorname{cut} \frac{\mathbf{R}, \Gamma, \psi_{1} \Rightarrow \chi}{\mathbf{R}, \Gamma, \Gamma', \psi_{1} \Rightarrow \varphi} \frac{\mathbf{R}, \Gamma, \chi \Rightarrow \varphi}{\mathbf{R}, \Gamma, \Gamma', \psi_{1} \Rightarrow \varphi}$$

This is the commutation over a \diamond -l step:

$$\diamond -l \frac{\mathbf{R}, xRy, \Gamma, y : A \Rightarrow \chi}{\operatorname{cut} \frac{\mathbf{R}, \Gamma, x : \diamond A \Rightarrow \chi}{\mathbf{R}, \Gamma', x : \diamond A \Rightarrow \varphi}} \underset{\mathsf{Cut}}{\overset{\mathcal{Q}}{\underset{\mathsf{R}, \Gamma, r', x : \diamond A \Rightarrow \varphi}{\operatorname{cut} \frac{\mathbf{R}, xRy, \Gamma, y : A \Rightarrow \chi}{\mathbf{R}, \Gamma, r', y : A \Rightarrow \varphi}} \underset{\sim}{\overset{\mathcal{R}, xRy, \Gamma, \Gamma', y : A \Rightarrow \chi}{\operatorname{cut} \frac{\mathbf{R}, xRy, \Gamma, \Gamma', y : A \Rightarrow \varphi}{\diamond -l \frac{\mathbf{R}, xRy, \Gamma, \Gamma', y : A \Rightarrow \varphi}{\mathbf{R}, \Gamma, \Gamma', x : \diamond A \Rightarrow \varphi}}$$

where xRy, Q is obtained from Q by prepending xRy to the LHS of each sequent. This is the commutation over a \Box -*l* step:

$$\Box -l \frac{\mathbf{R}, xRy, \Gamma, y : A \Rightarrow \chi}{\mathbf{R}, xRy, \Gamma, x : \Box A \Rightarrow \chi} \qquad \overbrace{\mathbf{R}, xRy, \Gamma', \chi \Rightarrow \varphi}_{\mathbf{R}, xRy, \Gamma, \Gamma', x : \Box A \Rightarrow \varphi} \sim$$



This is the commutation over a c-l step:

$$\operatorname{c-l} \frac{\mathbf{R}, \Gamma, \psi, \psi \Rightarrow \chi}{\operatorname{cut} \frac{\mathbf{R}, \Gamma, \psi, \psi \Rightarrow \chi}{\mathbf{R}, \Gamma, \psi, \Gamma' \Rightarrow \varphi}} \underset{\operatorname{cut}}{\overset{Q}{\mathbf{R}, \Gamma, \psi, \psi \Rightarrow \chi}} \underset{\operatorname{R}, \Gamma, \psi, \Gamma' \Rightarrow \varphi}{\overset{Q}{\mathbf{R}, \Gamma, \psi, \psi \Rightarrow \chi}} \underset{\operatorname{R}, \Gamma, \psi, \psi, \Gamma' \Rightarrow \varphi}{\overset{Q}{\mathbf{R}, \Gamma, \psi, \psi, \Gamma' \Rightarrow \varphi}} \underset{\operatorname{R}, \Gamma, \psi, \psi, \Gamma' \Rightarrow \varphi}{\overset{Q}{\mathbf{R}, \Gamma, \psi, \psi, \Gamma' \Rightarrow \varphi}}$$

The commutations over w-l and \wedge -l are similar.