# Finite element approximation for a delayed generalized Burgers-Huxley equation with weakly singular kernels: Part I Well-posedness, Regularity and Conforming approximation

SUMIT MAHAJAN<sup>†</sup> ARBAZ KHAN<sup>‡</sup>, AND MANIL T. MOHAN§

Department of Mathematics, Indian Institute of Technology Roorkee, Roorkee 247667, India

[Received on 6 September 2023]

The analysis of a delayed generalized Burgers-Huxley equation (a non-linear advection-diffusionreaction problem) with weakly singular kernels is carried out in this work. Moreover, numerical approximations are performed using the conforming finite element method (CFEM). The existence, uniqueness and regularity results for the continuous problem have been discussed in detail using the Faedo-Galerkin approximation technique. For the numerical studies, we first propose a semi-discrete conforming finite element scheme for space discretization and discuss its error estimates under minimal regularity assumptions. We then employ a backward Euler discretization in time and CFEM in space to obtain a fully-discrete approximation. Additionally, we derive a prior error estimates for the fully-discrete approximated solution. Finally, we present computational results that support the derived theoretical results.

Keywords: Burgers-Huxley equation, Galerkin approximation, weak solution, A priori error analysis,

Conforming finite element method.

### 1. Introduction

The generalized Burgers-Huxley equation (GBHE) with weakly singular kernels describes a prototype model for describing the interaction between reaction mechanisms, convection effects and diffusion transports with delays. The GBHE has various applications in different fields such as traffic flows, nuclear waste disposal, the mobility of fish populations, motion of the domain wall of a ferroelectric material in an electric field, etc. In this work, we consider a delayed generalized Burgers-Huxley equation with weakly singular kernels defined on  $\Omega \times (0,T)$ , where  $\Omega \subset \mathbb{R}^d$  (d = 2,3) be an open bounded and simply connected convex domain with Lipschitz boundary  $\partial \Omega$ . The delayed GBHE under our consideration is given by

$$\begin{cases} \frac{\partial u(x,t)}{\partial t} + \alpha u(x,t)^{\delta} \sum_{i=1}^{d} \frac{\partial u(x,t)}{\partial x_{i}} - \nu \Delta u(x,t) - \eta \int_{0}^{t} K(t-s) \Delta u(x,s) \, \mathrm{d}s \\ &= \beta u(x,t)(1-u(x,t)^{\delta})(u(x,t)^{\delta} - \gamma) + f(x,t), \, (x,t) \in \Omega \times (0,T), \\ u(x,t) = 0, \, x \in \partial \Omega, \, t \in [0,T), \\ u(x,0) = u_{0}(x), \, x \in \Omega, \end{cases}$$
(1.1)

<sup>†</sup>Email: sumit\_m@ma.iitr.ac.in

<sup>‡</sup>Corresponding author. Email: arbaz@ma.iitr.ac.in

<sup>§</sup>Email: maniltmohan@ma.iitr.ac.in, maniltmohan@gmail.com.

where K(t) is a weakly singular kernel,  $f(\cdot, \cdot)$  is an external forcing,  $\alpha, \beta, \nu, \delta, \eta$  are parameters such that  $\alpha > 0$  is the advection coefficient,  $\beta > 0$ ,  $\delta \ge 1$  is the retardation time,  $\eta \ge 0$  is the relaxation time,  $\gamma \in (0,1)$  and d = 2,3. The integration term is used to study the delayed effect.

For  $\eta = 0$ , the equation (1.1) becomes

$$\frac{\partial u}{\partial t} + \alpha u^{\delta} \sum_{i=1}^{d} \frac{\partial u}{\partial x_{i}} - \nu \Delta u = \beta u (1 - u^{\delta}) (u^{\delta} - \gamma) + f$$

which is the generalized Burgers-Huxley equation (GBHE). A wide literature is available on the GBHE and related models. Solitary and travelling wave solutions of the generalized Burgers-Huxley equation using a relevant non-linear transformation is obtained (Wang *et al.* (see 1990); Ablowitz *et al.* (see 1987); El-Danaf (see 2007); Wazwaz (see 2005), etc.). The global solvability results for the stochastic generalized Burgers-Huxley (SGBH) equation have also been proved in Kumar & Mohan (2022). Many researchers have established numerical solutions of the GBHE by using various numerical techniques such as Hashim *et al.* (2006) used an Adomian decomposition method, Khattak (2009) provided a numerical solution of GBHE based on the collocation method using mesh-less radial basis functions, Sari *et al.* (2011) performed numerical studies using a fourth-order Runge-Kutta (RK4) scheme in time combined with up to tenth-order finite difference scheme in space, Kumar *et al.* (2011) used a three-step Taylor–Galerkin finite element scheme to approximate the solution of the singularly perturbed GBHE.

For  $\delta = 1$ ,  $\eta = 0$ ,  $\alpha \neq 0$  and  $\beta \neq 0$ , the equation (1.1) is known as the *Burgers-Huxley equation* (see Wang *et al.* (1990); Wazwaz (2009), etc.). For  $\alpha = 0$ ,  $\eta = 0$  and  $\delta = 1$ , the equation (1.1) takes the form which is known as the *Huxley equation* which describes the nerve pulse propagation in nerve fibres and wall motion in liquid crystals (see Wang, 1985). For  $\beta = 0$ ,  $\delta = 1$ , and  $\alpha = 1$ , the equation (1.1) reduces to the well-known viscous Burgers' equation which represents the mathematical modelling of the turbulence phenomena (see Burgers, 1948), far field of wave propagation in non-linear dissipative systems, shock wave theory, vorticity transportation, heat conduction, wave processes in thermoelastic medium, dispersion in porous media, hydrodynamic turbulence, elasticity, gas dynamics, etc.

The kernel function  $K(\cdot)$  is *weakly singular* such that  $K \in L^1(0,T)$ . Moreover, we assume that the function  $K(\cdot)$  is a *positive kernel*, that is, if for any T > 0, we have

$$\int_0^T \int_0^t K(t-s)u(s)u(t)ds \, \mathrm{d}t \ge 0, \quad \text{for all } u \in \mathrm{L}^2(0,T).$$

$$(1.2)$$

One primary example of the weakly singular kernel is given by

$$K(t) = \frac{1}{\Gamma(\alpha)} \frac{1}{t^{1-\alpha}}, \quad 0 < \alpha < 1,$$

where  $\Gamma(\alpha) = \int_0^\infty t^{\alpha-1} e^{-t} dt$  is the *Euler Gamma function*. The convolution integral can be interpreted as a fractional integral of order  $\alpha$  (see, e.g., Adolfsson *et al.*, 2003; Oldham & Spanier, 1974). By varying the fractional integral exponent, we obtain a link between parabolic behavior (in the limit  $\alpha = 0$ , the kernel *K* approaches to the delta distribution) and hyperbolic behavior (specializing  $\alpha = 1$ ). The global solvability and numerical studies, for the one dimensional GBHE and the two-dimensional stationary generalized Burgers- Huxley equation, have been recently established in Mohan & Khan (2021) and Khan *et al.* (2021), respectively. The literature relevant to the finite element analysis of integrodifferential equations of parabolic type is vast. The time discretization for integro-differential equations with a smooth kernel was introduced in Sloan & Thomée (1986). A prior error estimates for linear

parabolic equations can be found in Cannon & Lin (1987) and Pani & Peterson (1996), and for both parabolic and hyperbolic equations in Yanik & Fairweather (1988). Error estimates for the semi-discrete finite element method (FEM) have been discussed in Thomée & Zhang (1989). Time discretization and error estimation using the Crank-Nicolson scheme and Padé-type scheme are discussed in Cannon & Lin (1988) and Zhang (1993), respectively.

Fully-discrete FEM for parabolic integro-differential equations with a weakly singular kernel using positive quadrature rule has been discussed in Chen *et al.* (1992) and McLean & Thomée (1993). A second-order accurate numerical method for a semi-linear integro-differential equation using the Crank-Nicolson scheme with a weakly singular kernel is discussed in Mustapha & Mustapha (2010). Recently, Zhou *et al.* (2019) used the weak Galerkin finite element scheme for the parabolic integro-differential equation with a weakly singular kernel to obtain the optimal convergence order in  $L^2$ .

To the best of the authors' knowledge, this work appears to be the first in the field of theoretical and numerical partial differential equations to investigate the solvability of the model generalized Burgers-Huxley equation with memory (1.1). This research article makes significant contributions from both theoretical and numerical perspectives:

- Well-posedness results: From theoretical point of view, the well-posedness of the generalized Burgers-Huxley equation with memory (Kernel) is proven for the first time in the literature (Section 2.3). The analysis is carried out in an open bounded and simply connected convex domain,  $\Omega \subset \mathbb{R}^d (d = 2, 3)$ , with a Lipschitz boundary  $\partial \Omega$ .
- **Regularity results**: The novel regularity results for the equation (1.1) in a convex domain or a domain with a C<sup>2</sup>-boundary under smoothness assumptions on the initial data and external forcing have been obtained in Section 2.4. These results are significant from a numerical analysis standpoint as well.
- Semi-discrete error estimates: This work provides, for the first time, semi-discrete error estimates for non-linear integro-differential equations with weakly singular kernels under minimal regularity assumptions.
- Fully-discrete FEM error estimates: The major contribution in the numerical aspect lies in the error estimates for the fully-discrete finite element method (FEM). Unlike the previous literature that necessitates  $u_{tt} \in L^2(0,T;L^2(\Omega))$  (Thomée (2007)) to handle the time derivative term, which requires the boundary to be at least C<sup>4</sup>, our results do not impose such higher regularity requirements. Notably, the minimum regularity result has been established for the linear parabolic equation Hou & Zhu (2006), but our work is the first to extend it for a non-linear equation with weakly singular kernels.

We show the existence and uniqueness of a global weak solution to the system (1.1), using a Faedo-Galerkin method. Regularity results of the weak solution are obtained under smoothness assumptions on the initial data and external forcing. Here, we propose the conforming finite elements (CFEM) scheme to obtain the numerical approximation of the exact solution under minimal regularity and the required regularity is also proved for GBHE with weakly singular kernels. We rigorously derive a priori error estimates indicating first-order convergence in both space and time of the CFEM. We also include a set of computational tests that confirm the theoretical error bounds and show some model equation properties. The finite element approximation using non-conforming and discontinuous Galerkin (DG) finite element method has been discussed in Mahajan & Khan (Prep).

4 of 45

#### S. MAHAJAN, A. KHAN, M.T. MOHAN

The paper is organized as follows: In section 2, we will discuss the notational conventions and provide the abstract formulation of our problem with the necessary functional spaces. The global solvability results for the system (1.1) are established in Theorem 2.2; that is, the existence and uniqueness of a global weak solution is proved by taking the initial data  $u_0 \in L^2(\Omega)$  and external forcing  $f \in L^2(0,T; H^{-1}(\Omega))$ . The uniqueness result of weak solutions under different choices on the initial data is discussed in Theorem 2.3. The regularity results are obtained in Theorem 2.4 and Theorem 2.5 under smoothness assumptions on the initial data and the external forcing. In the subsequent section, the semi-discrete and the fully-discrete scheme are introduced, and a priori error estimates are derived for the CFEM in section 3. Finally, numerical results are presented in section 4 for 2D and 3D to support our theoretical results.

### 2. Mathematical Setting and Solvability

This section lists necessary function spaces and notations used throughout the paper. Also, we define the operators used to obtain the global solvability results of the system (1.1), discuss its properties and establish the existence of a unique weak solution for our system and its regularity.

# 2.1 Functional setting

We use the standard notation for function spaces. In particular, the Lebesgue spaces are denoted by  $L^p(\Omega)$  for  $p \in [1,\infty]$  and the corresponding norm is denoted by  $\|\cdot\|_{L^p}$  (omitting the domain specification whenever clear from the context). For p = 2,  $L^2(\Omega)$  is a Hilbert space with the inner product denoted by  $(\cdot, \cdot)$ . The Hilbertian Sobolev spaces are denoted by  $H^k(\Omega)$ . Let  $C_0^{\infty}(\Omega)$  denote the space of all infinitely differentiable functions with compact support in  $\Omega$  and  $H_0^1(\Omega)$  be the closure of  $C_0^{\infty}(\Omega)$  in H<sup>1</sup>-norm. As  $\Omega$  is a bounded domain,  $\|\nabla \cdot\|_{L^2}$  defines an equivalent norm on  $H_0^1(\Omega)$  by using the Poincaré inequality. Moreover, we have the Gelfand triplet  $H_0^1(\Omega) \hookrightarrow L^2(\Omega) \hookrightarrow H^{-1}(\Omega)$ , where  $H^{-1}(\Omega)$  is the dual of  $H_0^1(\Omega)$ , and the embedding  $H_0^1(\Omega) \hookrightarrow L^2(\Omega)$  is compact. The duality paring between  $H_0^1(\Omega)$  and  $H^{-1}(\Omega)$  as well as  $L^p(\Omega)$  and its dual  $L^{p'}(\Omega)$ , where  $\frac{1}{p} + \frac{1}{p'} = 1$  is denoted by  $\langle \cdot, \cdot \rangle$ . The sum space  $H^{-1}(\Omega) + L^{p'}(\Omega)$  is well defined and

$$(\mathrm{H}^{-1}(\varOmega) + \mathrm{L}^{p'}(\varOmega))' = \mathrm{H}^{1}_{0}(\varOmega) \cap \mathrm{L}^{p}(\varOmega) \text{ and } (\mathrm{H}^{1}_{0}(\varOmega) \cap \mathrm{L}^{p}(\varOmega))' = \mathrm{H}^{-1}(\varOmega) + \mathrm{L}^{p'}(\varOmega),$$

where  $||y||_{H_0^1 \cap L^p} = \max\{||y||_{H_0^1}, ||y||_{L^p}\}$ , which is equivalent to the norms  $||y||_{H_0^1} + ||y||_{L^p}$  and  $\sqrt{||y||_{H_0^1}^2 + ||y||_{L^p}^2}$ , and

$$\begin{split} \|y\|_{\mathrm{H}^{-1}+\mathrm{L}^{p'}} &= \inf\{\|y_1\|_{\mathrm{H}^{-1}} + \|y_2\|_{\mathrm{L}^{p'}} : y = y_1 + y_2, y_1 \in \mathrm{H}^{-1}(\Omega) \text{ and } y_2 \in \mathrm{L}^{p'}(\Omega)\}\\ &= \sup\left\{\frac{|\langle y_1 + y_2, f\rangle|}{\|f\|_{\mathrm{H}^1_0\cap\mathrm{L}^p}} : 0 \neq f \in \mathrm{H}^1_0(\Omega) \cap \mathrm{L}^p(\Omega)\right\}. \end{split}$$

Note that  $H_0^1(\Omega) \cap L^p(\Omega)$  and  $H^{-1}(\Omega) + L^{p'}(\Omega)$  are Banach spaces. Moreover, we have the continuous embedding  $H_0^1(\Omega) \cap L^p(\Omega) \hookrightarrow H_0^1(\Omega) \hookrightarrow L^2(\Omega) \hookrightarrow H^{-1}(\Omega) \hookrightarrow H^{-1}(\Omega) + L^{p'}(\Omega)$ . The following operators will be used in further analysis:

2.1.1 *Linear operator.* Let A be the unbounded and self-adjoint linear operator on  $L^2(\Omega)$  defined by:

$$Au := -\Delta u, D(A) = \mathrm{H}^{2}(\Omega) \cap \mathrm{H}^{1}_{0}(\Omega),$$

5 of 45

whenever  $\Omega$  is convex, bounded open subset of  $\mathbb{R}^n$  with a C<sup>2</sup> boundary  $\partial \Omega$  (Theorem 3.1.2.1, Grisvard (1985)). Note that  $D(A) \subset L^p$ , for  $p \in [1, \infty]$  and d = 2, 3, using the Sobolev Embedding Theorem (see Mitrovic & Zubrinic, 1997). As the embedding  $H_0^1(\Omega) \hookrightarrow L^2(\Omega)$  is compact,  $A^{-1}$  exists and is a symmetric compact operator on  $L^2(\Omega)$ . Then, by using spectral theorem (see Dautray & Lions, 1990), there exists a sequence of eigenvalues of A,  $0 < \lambda_1 \leq \lambda_2 \leq \ldots \rightarrow \infty$  and an orthonormal basis  $\{w_k\}_{k=1}^{\infty}$  of  $L^2(\Omega)$  consisting of eigenfunctions of A (see Dautray & Lions, 1990, p. 504).

An integration by parts yields

$$\langle Au, v \rangle = (\nabla u, \nabla v), \text{ for all } u, v \in \mathrm{H}_0^1(\Omega),$$

so that  $A : \mathrm{H}^{1}_{0}(\Omega) \to \mathrm{H}^{-1}(\Omega)$ .

2.1.2 *Non-linear operators.* Let us define the non-linear operator  $b: (H_0^1(\Omega) \cap L^{2(\delta+1)}(\Omega)) \times H_0^1(\Omega) \times (H_0^1(\Omega) \cap L^{2(\delta+1)}(\Omega)) \to \mathbb{R}$  by

$$b(u, v, w) = \int_{\Omega} (u(x))^{\delta} \sum_{i=1}^{d} \frac{\partial v(x)}{\partial x_i} w(x) \, \mathrm{d}x.$$

The map is well-defined since  $|b(u, v, w)| \leq ||u||_{L^{2(\delta+1)}}^{\delta} ||v||_{H_{0}^{1}} ||w||_{L^{2(\delta+1)}}$ , so that we can define a mapping  $B(\cdot, \cdot) : (H_{0}^{1}(\Omega) \cap L^{2(\delta+1)}(\Omega)) \times H_{0}^{1}(\Omega) \to H^{-1}(\Omega) + L^{\frac{2(\delta+1)}{2\delta+1}}(\Omega)$  by  $\langle B(u, v), w \rangle = b(u, v, w)$ . For simplicity, we use the notation B(u) = B(u, u).

For all  $p \ge 2$  and  $u \in H_0^1(\Omega) \cap L^{2(\delta+1)}(\Omega)$ , using an integration by parts and boundary conditions, we have

$$\begin{split} b(u,u,|u|^{p-2}u) &= \left(u^{\delta}\sum_{i=1}^{d}\frac{\partial u}{\partial x_{i}},|u|^{p-2}u\right) = \frac{1}{\delta+2}\sum_{i=1}^{d}\int_{\Omega}\frac{\partial}{\partial x_{i}}(u(x))^{\delta+2}|u(x)|^{p-2} dx \\ &= -\frac{1}{\delta+2}\sum_{i=1}^{d}\int_{\Omega}(u(x))^{\delta+2}\frac{\partial}{\partial x_{i}}|u(x)|^{p-2} dx = -\frac{p-2}{\delta+2}\sum_{i=1}^{d}\int_{\Omega}(u(x))^{\delta}|u(x)|^{p-2}\frac{\partial u(x)}{\partial x_{i}} dx \\ &= -\frac{p-2}{\delta+2}\left(u^{\delta}\sum_{i=1}^{d}\frac{\partial u}{\partial x_{i}},|u|^{p-2}u\right). \end{split}$$

The above expression implies

$$b(u,u,|u|^{p-2}u) = \left(u^{\delta}\sum_{i=1}^{d}\frac{\partial u}{\partial x_{i}},|u|^{p-2}u\right) = 0.$$
(2.1)

Let us now show that the operator  $B(\cdot)$  is locally Lipschitz. For  $u, v, z \in H_0^1(\Omega) \cap L^{2(\delta+1)}(\Omega)$ , and w = u - v, using integration by parts and Taylor's formula for  $0 < \theta_1 < 1$ , we have

$$\langle B(u) - B(v), z \rangle$$

$$= \left( u^{\delta} \sum_{i=1}^{d} \frac{\partial u}{\partial x_{i}} - v^{\delta} \sum_{i=1}^{d} \frac{\partial v}{\partial x_{i}}, z \right) = \left( u^{\delta} \sum_{i=1}^{d} \frac{\partial w}{\partial x_{i}}, z \right) + \left( (u^{\delta} - v^{\delta}) \sum_{i=1}^{d} \frac{\partial v}{\partial x_{i}}, z \right)$$

$$= -\delta \left( u^{\delta-1} \sum_{i=1}^{d} \frac{\partial u}{\partial x_{i}} w, z \right) - \left( u^{\delta} w, \sum_{i=1}^{d} \frac{\partial z}{\partial x_{i}} \right) + \delta \left( (\theta_{1}u + (1 - \theta_{1})v)^{\delta-1} \sum_{i=1}^{d} \frac{\partial v}{\partial x_{i}} w, z \right)$$

$$\leq \left( \delta \|u\|_{L^{2(\delta+1)}}^{\delta-1} \|\nabla u\|_{L^{2}} \|z\|_{L^{2(\delta+1)}} + \|u\|_{L^{2(\delta+1)}}^{\delta} \|\nabla z\|_{L^{2}} + \delta 2^{\delta} (\|u\|_{L^{2(\delta+1)}}^{\delta-1} + \|v\|_{L^{2(\delta+1)}}^{\delta-1}) \|\nabla v\|_{L^{2}} \|z\|_{L^{2(\delta+1)}} \right) \\ \times \|w\|_{L^{2(\delta+1)}} \leq C \left[ \left( \|u\|_{L^{2(\delta+1)}}^{\delta-1} + \|v\|_{L^{2(\delta+1)}}^{\delta-1} \right) \left( \|u\|_{H^{1}_{0}}^{1} + \|v\|_{H^{1}_{0}}^{1} \right) + \|u\|_{L^{2(\delta+1)}}^{\delta} \right] \|w\|_{L^{2(\delta+1)}} \|z\|_{H^{1}_{0} \cap L^{2(\delta+1)}},$$

so that

$$\begin{aligned} \|B(u) - B(v)\|_{\mathbf{H}^{-1} + \mathbf{L}^{\frac{2(\delta+1)}{2\delta+1}}} &\leq C \left[ \left( \|u\|_{\mathbf{L}^{2(\delta+1)}}^{\delta-1} + \|v\|_{\mathbf{L}^{2(\delta+1)}}^{\delta-1} \right) \left( \|u\|_{\mathbf{H}^{1}_{0}}^{1} + \|v\|_{\mathbf{H}^{1}_{0}}^{1} \right) + \|u\|_{\mathbf{L}^{2(\delta+1)}}^{\delta} \right] \|w\|_{\mathbf{L}^{2(\delta+1)}} \\ &\leq Cr^{\delta} \|u - v\|_{\mathbf{L}^{2(\delta+1)}}, \end{aligned}$$

$$(2.2)$$

for all  $\|u\|_{\mathrm{H}^{1}_{0}\cap\mathrm{L}^{2(\delta+1)}}, \|v\|_{\mathrm{H}^{1}_{0}\cap\mathrm{L}^{2(\delta+1)}} \leq r$ . Therefore, the operator  $B:\mathrm{H}^{1}_{0}(\Omega)\cap\mathrm{L}^{2(\delta+1)}(\Omega) \to \mathrm{H}^{-1}(\Omega) + \mathrm{L}^{\frac{2(\delta+1)}{2\delta+1}}(\Omega)$  is locally Lipschitz.

Let us now consider the non-linear operator  $c(u) = u(1-u^{\delta})(u^{\delta}-\gamma)$ . It should be noted that

$$(c(u), u) = (u(1 - u^{\delta})(u^{\delta} - \gamma), u) = ((1 + \gamma)u^{\delta + 1} - \gamma u - u^{2\delta + 1}, u)$$
  
$$= (1 + \gamma)(u^{\delta + 1}, u) - \gamma ||u||_{L^{2}}^{2} - ||u||_{L^{2(\delta + 1)}}^{2(\delta + 1)}$$
  
$$\leq (1 + \gamma) ||u||_{L^{2(\delta + 1)}}^{\delta + 1} ||u||_{L^{2}} - \gamma ||u||_{L^{2}}^{2} - ||u||_{L^{2(\delta + 1)}}^{2(\delta + 1)},$$
(2.3)

for all  $u \in L^{2(\delta+1)}(\Omega)$ . Using Taylor's formula and Hölder's inequality, for  $u, v, z \in L^{2(\delta+1)}(\Omega)$  with w = u - v,  $0 < \theta_2 < 1$  and  $0 < \theta_3 < 1$ , we get

$$\begin{aligned} \langle c(u) - c(v), z \rangle &= \langle (1+\gamma)(u^{\delta+1} - v^{\delta+1}) - \gamma(u-v) - (u^{2\delta+1} - v^{2\delta+1}), z \rangle \\ &= (1+\gamma)(\delta+1)(w(\theta_2 u + (1-\theta_2)v)^{\delta}, z) - \gamma(w, z) \\ &- (2\delta+1)\langle w(\theta_3 u + (1-\theta_3)v)^{2\delta}, z \rangle \\ &\leqslant (1+\gamma)(\delta+1)2^{\delta-1} \left( \|u\|_{L^{2(\delta+1)}}^{\delta} + \|v\|_{L^{2(\delta+1)}}^{\delta} \right) \|w\|_{L^{2(\delta+1)}} \|z\|_{L^2} + \gamma \|w\|_{L^2} \|z\|_{L^2} \\ &+ (2\delta+1)2^{2\delta-1} \left( \|u\|_{L^{2(\delta+1)}}^{2\delta} + \|v\|_{L^{2(\delta+1)}}^{\delta} \right) \|w\|_{L^{2(\delta+1)}} \|z\|_{L^{2(\delta+1)}} \\ &\leqslant \left( (1+\gamma)(\delta+1)2^{\delta} |\Omega|^{\frac{\delta}{2(\delta+1)}} r^{\delta} + \gamma |\Omega|^{\frac{\delta}{\delta+1}} + (2\delta+1)2^{2\delta} r^{2\delta} \right) \|w\|_{L^{2(\delta+1)}} \|z\|_{L^{2(\delta+1)}} \|z\|_{L^{2(\delta+1)}} \|v\|_{L^{2(\delta+1)}} \|z\|_{L^{2(\delta+1)}} \|v\|_{L^{2(\delta+1)}} \|v\|_$$

for  $\|u\|_{L^{2(\delta+1)}}, \|v\|_{L^{2(\delta+1)}} \leq r$ , where  $|\Omega|$  is the Lebesgue measure of  $\Omega$ . Thus, the operator  $c(\cdot)$ :  $L^{2(\delta+1)}(\Omega) \rightarrow L^{\frac{2(\delta+1)}{2\delta+1}}(\Omega)$  is locally Lipschitz.

The following lemmas are useful for the further analysis:

LEMMA 2.1 (Mohan, 2020) Let  $K \in L^1(0,T)$  and  $\phi \in L^2(0,T)$  for some T > 0. Then

$$\left(\int_0^T \left(\int_0^s K(s-\tau)\phi(\tau) \,\mathrm{d}\tau\right)^2 \,\mathrm{d}s\right)^{\frac{1}{2}} \leq \left(\int_0^T |K(s)| \,\mathrm{d}s\right) \left(\int_0^T \phi^2(s) \,\mathrm{d}s\right)^{\frac{1}{2}}.$$

LEMMA 2.2 (Remark A.2, Mohan & Sritharan (2019)) If the kernel  $K(\cdot)$  satisfies the conditions

$$\begin{cases} K \in \mathbf{C}^{2}(0,\infty) \cap \mathbf{C}[0,\infty), \\ (-1)^{k} K^{(k)}(t) \ge 0, \text{ for all } t > 0, k = 0, 1, 2, \end{cases}$$

then for any right continuous function  $f : [0,T] \to [0,\infty)$  with left limits and for  $g : [0,T] \to \mathbb{R}$  such that  $(K * g)(\cdot)$  is well defined, we have

$$\int_0^T f(t)(K * g)(t)g(t)dt \ge 0.$$
(2.5)

REMARK 2.1 By approximating a weakly singular kernel with a smooth kernel having the properties obtained in Lemma 2.2, one can obtain the result (2.5) for weakly singular kernels also. For instance, one can approximate  $K(t) = \frac{1}{\Gamma(\alpha)} \frac{1}{t^{1-\alpha}}$ ,  $0 < \alpha < 1$ , by  $K_{\varepsilon}(t) = \frac{1}{\Gamma(\alpha)} \frac{1}{(t+\varepsilon)^{1-\alpha}}$ , for some  $\varepsilon > 0$ .

# 2.2 Abstract formulation

By using the above functional setting, the abstract formulation of the system (1.1) is given by

$$\begin{cases} \frac{\mathrm{d}u(t)}{\mathrm{d}t} + vAu(t) + \eta(K * Au)(t) = -\alpha B(u(t)) + \beta c(u(t)) + f(t), & \text{in } \mathrm{H}^{-1}(\Omega) + \mathrm{L}^{\frac{2(\delta+1)}{2\delta+1}}(\Omega), \\ u(0) = u_0 \in \mathrm{L}^2(\Omega), \end{cases}$$
(2.6)

for a.e.  $t \in [0,T]$ , where  $(K * Au)(t) = \int_0^t K(t-s)Au(s)ds$ .

Let us now provide the definition of weak solution of the system (2.6).

DEFINITION 2.1 (Weak solution) For the initial data  $u_0 \in L^2(\Omega)$  and external forcing  $f \in L^2(0, T; H^{-1}(\Omega))$ , we say that a function

$$u \in \mathcal{L}^{\infty}(0,T;\mathcal{L}^{2}(\Omega)) \cap \mathcal{L}^{2}(0,T;\mathcal{H}^{1}_{0}(\Omega)) \cap \mathcal{L}^{2(\delta+1)}(0,T;\mathcal{L}^{2(\delta+1)}(\Omega)),$$

with

$$\partial_t u \in \mathrm{L}^{\frac{2(\delta+1)}{2\delta+1}}(0,T;\mathrm{H}^{-1}(\Omega) + \mathrm{L}^{\frac{2(\delta+1)}{2\delta+1}}(\Omega)),$$

is called *a weak solution* to the system (2.6), if, for a.e.  $t \in (0,T)$ ,  $u(\cdot)$  satisfies:

$$\begin{cases} \langle \partial_t u(t), v \rangle + v(\nabla u(t), \nabla v) + \alpha b(u(t), u(t), v) + \eta((K * \nabla u)(s), \nabla v) = \beta \langle c(u(t)), v \rangle + \langle f(t), v \rangle, \\ (u(0), v) = (u_0, v), \end{cases}$$

for any  $v \in H_0^1(\Omega) \cap L^{2(\delta+1)}(\Omega)$ .

# 2.3 Existence and uniqueness of weak solution

Let us now discuss the existence and uniqueness of the weak solution for the system (1.1).

THEOREM 2.2 (Existence of a weak solution) For  $u_0 \in L^2(\Omega)$  and  $f \in L^2(0,T;H^{-1}(\Omega))$ , there exists a weak solution to the system (1.1) in the sense of Definition 2.1.

*Proof.* We establish the existence of a weak solution to the system (1.1) in the following steps:

**Step 1:** *Faedo-Galerkin approximatiiron.* Let the set  $\{w_k(x)\}_{k=1}^{\infty}$  be an orthogonal basis of  $H_0^1(\Omega)$  and orthonormal basis of  $L^2(\Omega)$  (page 504, Dautray & Lions (1990)). We look for a function  $u_m : [0,T] \rightarrow H_0^1(\Omega)$  of the form  $u_m(t) = \sum_{k=1}^m d_m^k(t)w_k$ , where *m* is a fixed positive integer and the coefficients  $d_m^k(t)$ ,

for  $0 \le t \le T$  and k = 1, ..., m are selected so that  $d_m^k(0) = (u_0, w_k)$ , for k = 1, 2, ..., m and

$$(u'_{m}(t), w_{k}) + \nu(\nabla u_{m}(t), \nabla w_{k}) + \alpha \left( u_{m}(t)^{\delta} \sum_{i=1}^{d} \frac{\partial u_{m}(t)}{\partial x_{i}}, w_{k} \right) + \eta((K * \nabla u_{m})(t), \nabla w_{k})$$
$$= \beta(u_{m}(t)(1 - (u_{m}(t))^{\delta})((u_{m}(t))^{\delta} - \gamma), w_{k}) + \langle f(t), w_{k} \rangle,$$
(2.7)

for a.e.  $0 \le t \le T$  and k = 1, ..., m. As the non-linear operators are locally Lipschitz (see (2.2) and (2.4)) using Carathéodory's existence theorem, the finite dimensional problem (2.7) has a local solution  $u_m \in C([0,T]; H_m)$ , where  $H_m = \text{span}\{w_1, ..., w_m\}$ .

**Step 2:** L<sup>2</sup>-energy estimate. Multiplying (2.7) by  $d_m^k(t)$ , and then summing for k = 1, ..., m, we have

$$\begin{aligned} &\frac{1}{2} \frac{d}{dt} \|u_m(t)\|_{L^2}^2 + \nu \|\nabla u_m(t)\|_{L^2}^2 + \alpha \left( u_m(t)^{\delta} \sum_{i=1}^d \frac{\partial u_m(t)}{\partial x_i}, u_m(t) \right) + \eta ((K * \nabla u_m)(t), \nabla u_m(t)) \\ &= \beta (u_m(t)(1 - (u_m(t))^{\delta})((u_m(t))^{\delta} - \gamma), u_m(t)) + \langle f(t), u_m(t) \rangle, \end{aligned}$$

for a.e.  $t \in [0,T]$ . Using the property of the non-linear operator (2.1) and the estimate (2.3), we get

$$\begin{split} &\frac{1}{2} \frac{\mathrm{d}}{\mathrm{d}t} \|u_m(t)\|_{\mathrm{L}^2}^2 + \nu \|\nabla u_m(t)\|_{\mathrm{L}^2}^2 + \beta \gamma \|u_m(t)\|_{\mathrm{L}^2}^2 + \beta \|u_m(t)\|_{\mathrm{L}^{2(\delta+1)}}^{2(\delta+1)} + \eta ((K * \nabla u_m)(t), \nabla u_m(t)) \\ &= \beta (1+\gamma) ((u_m(t))^{\delta+1}, u_m(t)) + \langle f(t), u_m(t) \rangle \\ &\leq \beta (1+\gamma) \|u_m(t)\|_{\mathrm{L}^{2(\delta+1)}}^{\delta+1} \|u_m(t)\|_{\mathrm{L}^2} + \|f(t)\|_{\mathrm{H}^{-1}} \|\nabla u_m(t)\|_{\mathrm{L}^2}^2 \\ &\leq \frac{\beta}{2} \|u_m(t)\|_{\mathrm{L}^{2(\delta+1)}}^{2(\delta+1)} + \frac{\beta (1+\gamma)^2}{2} \|u_m(t)\|_{\mathrm{L}^2}^2 + \frac{\nu}{2} \|\nabla u_m(t)\|_{\mathrm{L}^2}^2 + \frac{1}{2\nu} \|f(t)\|_{\mathrm{H}^{-1}}^2. \end{split}$$

Integrating the above inequality from 0 to t, we have

$$\begin{aligned} \|u_{m}(t)\|_{L^{2}}^{2} + \nu \int_{0}^{t} \|\nabla u_{m}(s)\|_{L^{2}}^{2} \, \mathrm{d}s + \beta \int_{0}^{t} \|u_{m}(s)\|_{L^{2(\delta+1)}}^{2(\delta+1)} \, \mathrm{d}s + 2\eta \int_{0}^{t} ((K * \nabla u_{m})(s), \nabla u_{m}(s)) \, \mathrm{d}s \\ &\leq \|u_{0}\|_{L^{2}}^{2} + \frac{1}{\nu} \int_{0}^{t} \|f(s)\|_{\mathrm{H}^{-1}}^{2} \, \mathrm{d}s + \beta (1 + \gamma^{2}) \int_{0}^{t} \|u_{m}(s)\|_{\mathrm{L}^{2}}^{2} \, \mathrm{d}s, \end{aligned}$$
(2.8)

for all  $t \in [0,T]$ , where we have used the fact that  $||u_m(0)||_{L^2} \leq ||u_0||_{L^2}$ . Using (1.2) and an application of Gronwall's inequality in (2.8) yields

$$\|u_m(t)\|_{L^2}^2 + \nu \int_0^t \|\nabla u_m(s)\|_{L^2}^2 \, \mathrm{d}s + \beta \int_0^t \|u_m(s)\|_{L^{2(\delta+1)}}^{2(\delta+1)} \, \mathrm{d}s \leq \left(\|u_0\|_{L^2}^2 + \frac{1}{\nu} \int_0^T \|f(t)\|_{H^{-1}}^2 \, \mathrm{d}t\right) e^{\beta(1+\gamma^2)T} \, \mathrm{d}s$$

for all  $t \in [0,T]$ . Taking supremum over time  $0 \le t \le T$ , we have

$$\sup_{0 \le t \le T} \|u_m(t)\|_{L^2}^2 + \nu \int_0^T \|\nabla u_m(t)\|_{L^2}^2 dt + \beta \int_0^T \|u_m(t)\|_{L^{2(\delta+1)}}^{2(\delta+1)} dt$$
  
$$\leq \left( \|u_0\|_{L^2}^2 + \frac{1}{\nu} \int_0^T \|f(t)\|_{H^{-1}}^2 dt \right) e^{\beta(1+\gamma^2)T},$$
(2.9)

and the right hand side of the inequality (2.9) is independent of m.

**Step 3:** *Estimate for the time derivative.* For any  $v \in H_0^1(\Omega) \cap L^{2(\delta+1)}(\Omega)$ , with  $||v||_{H_0^1 \cap L^{2(\delta+1)}} \leq 1$ , we have

$$\langle u'_m, v \rangle = (u'_m, v) = -v(\nabla u_m, \nabla v) - \alpha b(u_m, u_m, v) + \eta(K * \nabla u_m, \nabla v) + \beta \langle c(u_m), v \rangle + \langle f, v \rangle.$$

An application of the Cauchy-Schwarz inequality and Hölder's inequality yield

$$\begin{aligned} |\langle u'_{m}, v \rangle| &\leq v |(\nabla u_{m}, \nabla v)| + \alpha |(b(u_{m}, u_{m}, v^{1}))| + \eta |(K * \nabla u_{m}, \nabla v)| + \beta |(c(u_{m}), v)| + |\langle f, v \rangle| \\ &\leq \left( v \|\nabla u_{m}\|_{L^{2}} + \frac{\alpha}{\delta + 1} \|u_{m}\|_{L^{2(\delta+1)}}^{\delta+1} + \|f\|_{H^{-1}} + \eta \|K * \nabla u_{m}\|_{L^{2}} \right) \|\nabla v\|_{L^{2}} \\ &+ \beta \left( (1 + \gamma) \|u_{m}\|_{L^{2(\delta+1)}}^{\delta+1} + \gamma \|u_{m}\|_{L^{2}} \right) \|v\|_{L^{2}} + \beta \|u_{m}\|_{L^{2(\delta+1)}}^{2\delta+1} \|v\|_{L^{2(\delta+1)}}. \end{aligned}$$

Taking supremum over all  $v \in H^1_0(\Omega) \cap L^{2(\delta+1)}(\Omega)$ , we have

$$\begin{aligned} \|u_m'\|_{\mathrm{H}^{-1}+\mathrm{L}^{\frac{2(\delta+1)}{2\delta+1}}} &\leqslant \left( \nu \|\nabla u_m\|_{\mathrm{L}^2} + \frac{\alpha}{\delta+1} \|u_m\|_{\mathrm{L}^{2(\delta+1)}}^{\delta+1} + \|f\|_{\mathrm{H}^{-1}} + \eta \|K * \nabla u_m\|_{\mathrm{L}^2} + C\beta(1+\gamma) \|u_m\|_{\mathrm{L}^{2(\delta+1)}}^{\delta+1} + C\gamma \|u_m\|_{\mathrm{L}^2}^{2} + \beta \|u_m\|_{\mathrm{L}^{2(\delta+1)}}^{2\delta+1} \right). \end{aligned}$$

Integrating from 0 to T and using Young's inequality, we arrive at

$$\begin{split} \int_{0}^{T} \|u_{m}'(t)\|_{H^{-1}+L^{\frac{2(\delta+1)}{2\delta+1}}}^{\frac{2(\delta+1)}{2\delta+1}} dt \leqslant C \bigg\{ v^{\frac{2(\delta+1)}{2\delta+1}} \int_{0}^{T} \|\nabla u_{m}(t)\|_{L^{2}}^{\frac{2(\delta+1)}{2\delta+1}} dt + \bigg(\frac{\alpha}{\delta+1}\bigg)^{\frac{2(\delta+1)}{2\delta+1}} \int_{0}^{T} \|u_{m}(t)\|_{L^{2(\delta+1)}}^{\frac{2(\delta+1)^{2}}{2\delta+1}} dt \\ &+ \int_{0}^{T} \|f(t)\|_{H^{-1}}^{\frac{2(\delta+1)}{2\delta+1}} dt + \eta^{\frac{2(\delta+1)}{2\delta+1}} \int_{0}^{T} \|K * \nabla u_{m}(t)\|_{L^{2}}^{\frac{2(\delta+1)}{2\delta+1}} dt \\ &+ (\beta(1+\gamma))^{\frac{2(\delta+1)}{2\delta+1}} \int_{0}^{T} \|u_{m}(t)\|_{L^{2(\delta+1)}}^{\frac{2(\delta+1)}{2\delta+1}} dt \\ &+ \gamma^{\frac{2\delta}{2\delta+1}} \int_{0}^{T} \|u_{m}(t)\|_{L^{2}}^{\frac{2(\delta+1)}{2\delta+1}} dt + \beta^{\frac{2(\delta+1)}{2\delta+1}} \int_{0}^{T} \|u_{m}(t)\|_{L^{2(\delta+1)}}^{\frac{2(\delta+1)}{2\delta+1}} dt \end{split}$$

$$(2.10)$$

where  $\int_0^T \|K * \nabla u_m(t)\|_{L^2}^{\frac{2(\delta+1)}{2\delta+1}} dt$  is bounded since (cf. Lemma 2.1)

$$\begin{split} \int_{0}^{T} \|K * \nabla u_{m}(t)\|_{L^{2}}^{\frac{2(\delta+1)}{2\delta+1}} \, \mathrm{d}t &\leq T^{\frac{\delta}{2\delta+1}} \left( \int_{0}^{T} \|K * \nabla u_{m}(t)\|_{L^{2}}^{2} \, \mathrm{d}t \right)^{\frac{\delta+1}{2\delta+1}} \\ &\leq T^{\frac{\delta}{2\delta+1}} \left( \int_{0}^{T} |K(t)| \, \mathrm{d}t \right)^{\frac{2(\delta+1)}{2\delta+1}} \left( \int_{0}^{T} \|\nabla u_{m}(t)\|_{L^{2}}^{2} \, \mathrm{d}t \right)^{\frac{\delta+1}{2\delta+1}} < \infty, \end{split}$$

and

$$\int_0^T \|f(t)\|_{\mathrm{H}^{-1}}^{\frac{2(\delta+1)}{2\delta+1}} \, \mathrm{d}t \leqslant T^{\frac{\delta}{2\delta+1}} \left(\int_0^T \|f(t)\|_{\mathrm{H}^{-1}}^2 \, \mathrm{d}t\right)^{\frac{\delta+1}{2\delta+1}}.$$

**Step 4:** *Passing to limit.* As the energy estimates (2.9) and (2.10) are uniformly bounded and independent of *m*, using the Banach-Alaoglu theorem, we can extract a subsequence  $\{u_{m_j}\}_{j=1}^{\infty}$  of  $\{u_m\}_{m=1}^{\infty}$  such that

$$\begin{cases} u_{m_j} \stackrel{w^*}{\longrightarrow} u \text{ in } L^{\infty}(0,T;L^2(\Omega)), \\ u_{m_j} \stackrel{w}{\longrightarrow} u \text{ in } L^2(0,T;H_0^1(\Omega)), \\ u_{m_j} \stackrel{w}{\longrightarrow} u \text{ in } L^{2(\delta+1)}(0,T;L^{2(\delta+1)}(\Omega)), \\ \partial_t u_{m_j} \stackrel{w}{\longrightarrow} \partial_t u \text{ in } \partial_t u \in L^{\frac{2(\delta+1)}{2\delta+1}}(0,T;H^{-1}(\Omega) + L^{\frac{2(\delta+1)}{2\delta+1}}(\Omega)), \end{cases}$$

$$(2.11)$$

as  $j \to \infty$ , for all  $\delta \ge 1$ . Since we have compact embedding of  $H_0^1(\Omega) \subset L^2(\Omega)$  (page 284, Evans (2010)), an application of the Aubin-Lions compactness lemma yields the following strong convergence:

$$u_{m_i} \to u \text{ in } L^2(0,T;L^2(\Omega)),$$

$$(2.12)$$

as  $j \to \infty$ . The above strong convergence also implies  $u_{m_{k_j}} \to u$ , for a.e.  $(t,x) \in [0,T] \times \Omega$  as  $j \to \infty$ (that is, along a subsequence). Let us now take limit in (2.7) along the subsequence. Choose a function  $v \in C^1([0,T]; H^1_0(\Omega))$  having the form  $v(t) = \sum_{k=1}^N d_m^k(t) w_k$ , where  $\{d_m^k(t)\}_{k=1}^N$  are given smooth functions and N is a fixed integer. Let us choose  $m \ge N$ , multiply (2.7) by  $d_m^k(t)$ , sum from k = 1 to N and then integrate it from 0 to T to find

$$\int_{0}^{T} \left[ \langle \partial_{t} u_{m}(t), v(t) \rangle + v(\nabla u_{m}(t), \nabla v(t)) + \eta \langle (K * \nabla u_{m}(t)), \nabla v(t) \rangle + \alpha \left( (u_{m}(t))^{\delta} \sum_{i=1}^{d} \frac{\partial u_{m}}{\partial x_{i}}(t), v(t) \right) \right] dt$$
$$= \int_{0}^{T} \left[ \beta (u_{m}(t)(1 - (u_{m}(t))^{\delta})((u_{m}(t))^{\delta} - \gamma), v(t)) + \langle f(t), v(t) \rangle \right] dt.$$
(2.13)

Setting  $m = m_j$  (more specifically  $m_{k_j}$ ) and using the convergences given in (2.11) and (2.12), we can take weak limit in (2.13) to obtain

$$\int_{0}^{T} \left[ \langle \partial_{t} u(t), v(t) \rangle + v(\nabla u(t), \nabla v(t)) + \eta \langle (K * \nabla u(t)), \nabla v(t) \rangle + \alpha \left( (u(t))^{\delta} \sum_{i=1}^{d} \frac{\partial u_{m}}{\partial x_{i}}(t), v(t) \right) \right] dt$$
$$= \int_{0}^{T} \left[ \beta (u(t)(1 - (u(t))^{\delta})((u(t))^{\delta} - \gamma), v(t)) + \langle f(t), v(t) \rangle \right] dt, \qquad (2.14)$$

provided if we show that

$$B(u_{m_{k_j}}) \xrightarrow{w} B(u)$$
, and  $c(u_{m_{k_j}}) \xrightarrow{w} c(u)$  in  $H^{-1}(\Omega) + L^{\frac{2(\delta+1)}{2\delta+1}}(\Omega)$ , as  $j \to \infty$ .

Using density argument  $C_0^{\infty}(\Omega)$  is dense in  $H_0^1(\Omega)$ , we will show that  $b(u_{m_{k_j}}, u_{m_{k_j}}, v) \to b(u, u, v)$ , for all  $v \in C_0^{\infty}(\Omega)$  to obtain that  $B(u_{m_{k_j}}) \xrightarrow{w} B(u)$  in  $H^{-1}(\Omega) + L^{\frac{2(\delta+1)}{2\delta+1}}(\Omega)$ , as  $j \to \infty$ . By an application of Taylor's formula, integration by parts, Hölder's inequality, and convergence, we obtain

$$|b(u_{m_{k_j}}, u_{m_{k_j}}, v) - b(u, u, v)|$$
  
=  $\left|\frac{1}{\delta + 1} \sum_{i=1}^d \int_{\Omega} (u_{m_{k_j}}^{\delta + 1}(x) - u^{\delta + 1}(x)) \frac{\partial v(x)}{\partial x_i} dx\right|$ 

# GENERALIZED BURGERS-HUXLEY EQUATION WITH WEAKLY SINGULAR KERNEL

$$= \left| \sum_{i=1}^{d} \int_{\Omega} (\theta u_{m_{k_j}}(x) + (1-\theta)u(x))^{\delta} (u_{m_{k_j}}(x) - u(x)) \frac{\partial v(x)}{\partial x_i} dx \right|$$
  
$$\leq \|u_{m_{k_j}} - u\|_{\mathrm{L}^2} \left( \|u_{m_{k_j}}\|_{\mathrm{L}^{2(\delta+1)}}^{\delta} + \|u\|_{\mathrm{L}^{2(\delta+1)}}^{\delta} \right) \|\nabla v\|_{\mathrm{L}^{2(\delta+1)}}$$
  
$$\to 0 \text{ as } j \to \infty, \text{ for all } v \in C_0^{\infty}(\Omega).$$

Again using Taylor's formula, interpolation and Hölder's inequalities, , we find

$$\begin{split} |(c(u_{m_{k_{j}}}) - c(u), v)| \\ &\leqslant (1 + \gamma) \left| \int_{\Omega} (u_{m_{k_{j}}}^{\delta+1}(x) - u^{\delta+1}(x))v(x) \, dx \right| + \left| \int_{\Omega} (u_{m_{k_{j}}}(x) - u(x))v(x) \, dx \right| \\ &+ \left| \int_{\Omega} (u_{m_{k_{j}}}^{2\delta+1}(x) - u^{2\delta+1}(x))v(x) \, dx \right| \\ &\leqslant \left( (1 + \gamma)(\delta + 1) \left( \|u_{m_{k_{j}}}\|_{L^{2}(\delta+1)}^{\delta} + \|u\|_{L^{2}(\delta+1)}^{\delta} \right) \|v\|_{L^{2}(\delta+1)} + \|v\|_{L^{2}} \right) \|u_{m_{k_{j}}} - u\|_{L^{2}} \\ &+ (1 + 2\delta) \|u_{m_{k_{j}}} - u\|_{L^{2}}^{\frac{1}{\delta}} \left( \|u_{m_{k_{j}}}\|_{L^{2}(\delta+1)}^{1-\frac{1}{\delta}} + \|u\|_{L^{2}(\delta+1)}^{1-\frac{1}{\delta}} \right) \left( \|u_{m_{k_{j}}}\|_{L^{2}(\delta+1)}^{2\delta} + \|u\|_{L^{2}(\delta+1)}^{2\delta} \right) \|v\|_{L^{\infty}} \\ &\to 0 \text{ as } j \to \infty, \text{ for all } v \in C_{0}^{\infty}(\Omega). \end{split}$$

which completes the convergence proof.

**Step 5:** *Initial data.* Let us now show that  $u(0) = u_0$ . From (2.14), we note that

$$\int_0^T \left[ -\langle u(t), v'(t) \rangle + v(\nabla u(t), \nabla v(t)) + \eta \langle (K * \nabla u(t)), \nabla v(t) \rangle + \alpha \left( (u(t))^{\delta} \sum_{i=1}^2 \frac{\partial u}{\partial x_i}(t), v(t) \right) \right] dt$$
$$= \int_0^T \left[ \beta (u(t)(1 - (u(t))^{\delta})((u(t))^{\delta} - \gamma), v(t)) + (f(t), v(t)) \right] dt + (u(0), v(0)), \tag{2.15}$$

for each  $v \in C^1([0,T]; H^1_0(\Omega))$  with v(T) = 0. In a similar fashion, we deduce from (2.13) that

$$\begin{split} &\int_0^T \left[ -\langle u_m(t), v'(t) \rangle + v(\nabla u_m(t), \nabla v(t)) + \eta \langle (K * \nabla u_m(t)), \nabla v(t) \rangle + \alpha \left( (u_m(t))^{\delta} \sum_{i=1}^2 \frac{\partial u_m}{\partial x_i}(t), v(t) \right) \right] \, \mathrm{d}t \\ &= \int_0^T \left[ \beta (u_m(t)(1 - (u_m(t))^{\delta})((u_m(t))^{\delta} - \gamma), v(t)) + (f(t), v(t)) \right] \, \mathrm{d}t + (u_m(0), v(0)). \end{split}$$

Taking  $m = m_j$  in the above expression, using the above convergences ((2.11) and (2.12)) and passing then to limit, we find

$$\int_0^T \left[ -\langle u(t), v'(t) \rangle + v(\nabla u(t), \nabla v(t)) + \eta \langle (K * \nabla u(t)), \nabla v(t) \rangle + \alpha \left( (u(t))^{\delta} \sum_{i=1}^2 \frac{\partial u}{\partial x_i}(t), v(t) \right) \right] dt$$
$$= \int_0^T \left[ \beta (u(t)(1 - (u(t))^{\delta})((u(t))^{\delta} - \gamma), v(t)) + (f(t), v(t)) \right] dt + (u_0, v(0)), \tag{2.16}$$

since  $u_m(0) \to u_0$  as  $m \to \infty$  in  $L^2(\Omega)$ . As v(0) is arbitrary, comparing (2.15) and (2.16), one can easily conclude that  $u(0) = u_0$  in  $L^2(\Omega)$ .

**Step 6:** Energy equality. As  $u \in L^2(0,T; H^1_0(\Omega)) \cap L^{2(\delta+1)}(0,T; L^{2(\delta+1)}(\Omega))$  with  $\partial_t u \in L^{\frac{2(\delta+1)}{2\delta+1}}(0,T; H^{-1}(\Omega) + L^{\frac{2(\delta+1)}{2\delta+1}}(\Omega))$  using an argument similar to Theorem 7.2 (Robinson & Pierre (2003) (Exercise 8.2)) we have  $u \in C^0([0,T]; L^2(\Omega))$  and the following energy equality holds

$$\begin{aligned} \|u(t)\|_{L^{2}}^{2} + 2\nu \int_{0}^{t} \|\nabla u(s)\|_{L^{2}}^{2} ds + 2\eta \int_{0}^{t} (K * \nabla u(s), \nabla u(s)) ds + 2\beta \gamma \int_{0}^{t} \|u(s)\|_{L^{2}}^{2} ds \\ + 2\beta \int_{0}^{T} \|u_{m}(t)\|_{L^{2(\delta+1)}}^{2(\delta+1)} ds = \|u_{0}\|_{L^{2}}^{2} + 2\beta \int_{0}^{t} (u^{\delta+1}(s), u(s)) ds + \int_{0}^{t} \langle f(s), u(s) \rangle ds, \end{aligned}$$

for all  $t \in [0, T]$ . This proves the existence of weak solution.

REMARK 2.2 If we take the initial data  $u_0 \in L^{2(\delta+1)}(\Omega)$ , and the external forcing,  $f \in L^2(0,T;L^2(\Omega))$ , then we can obtain the  $L^{2(\delta+1)}$ -energy estimate as follows:

Multiplying (2.7) by  $d_m^k(t)|u_m(t)|^{2\delta}$ , and then summing for k = 1, ..., m, we obtain for a.e.  $t \in [0,T]$ 

$$\begin{aligned} &(\partial_t u_m(t), |u_m(t)|^{2\delta} u_m(t)) + \mathbf{v}(2\delta + 1)(\nabla u_m(t), |u_m(t)|^{2\delta} \nabla u_m(t)) \\ &+ \eta(2\delta + 1)((K * \nabla u_m)(t), |u_m(t)|^{2\delta} \nabla u_m(t)) + \alpha \left( (u_m(t))^{\delta} \sum_{i=1}^d \frac{\partial u_m}{\partial x_i}(t), |u_m(t)|^{2\delta} u_m(t) \right) \\ &= \beta (u_m(t)(1 - (u_m(t))^{\delta})((u_m(t))^{\delta} - \gamma, |u_m(t)|^{2\delta} u_m(t)) + (f, |u_m(t)|^{2\delta} u_m(t)). \end{aligned}$$

Now, using (2.1), the Cauchy-Schwarz, Hölder's and Young's inequalities, we arrive at

$$\begin{split} &\frac{1}{2(\delta+1)} \frac{\mathrm{d}}{\mathrm{d}t} \|u_m(t)\|_{\mathrm{L}^{2(\delta+1)}}^{2(\delta+1)} + \nu(2\delta+1) \||u_m(t)|^{\delta} \nabla u_m(t)\|_{\mathrm{L}^2}^2 \\ &+ \eta(2\delta+1)((K*\nabla u_m)(t), |u_m(t)|^{2\delta} \nabla u_m(t)) + \beta \|u_m(t)\|_{\mathrm{L}^{4\delta+2}}^{4\delta+2} + \beta \gamma \|u_m(t)\|_{\mathrm{L}^{2(\delta+1)}}^{2(\delta+1)} \\ &\leqslant \frac{\beta}{2} \|u_m(t)\|_{\mathrm{L}^{4\delta+2}}^{4\delta+2} + \beta (1+\gamma)^2 \|u_m(t)\|_{\mathrm{L}^{2(\delta+1)}}^{2(\delta+1)} + \frac{1}{\beta} \|f(t)\|_{\mathrm{L}^2}^2, \end{split}$$

for a.e.  $t \in [0, T]$ . Integrating the above inequality from 0 to t and using Lemma 2.2 and Remark 2.1, we have

$$\frac{1}{2(\delta+1)} \|u_m(t)\|_{L^{2(\delta+1)}}^{2(\delta+1)} + \nu(2\delta+1) \int_0^t \||u_m(s)|^{\delta} \nabla u_m(s)\|_{L^2}^2 \, \mathrm{d}s + \frac{\beta}{2} \int_0^t \|u_m(s)\|_{L^{4\delta+2}}^{4\delta+2} \, \mathrm{d}s$$

$$\leq \frac{1}{2(\delta+1)} \|u_0\|_{L^{2(\delta+1)}}^{2(\delta+1)} + \frac{1}{\beta} \int_0^t \|f(s)\|_{L^2}^2 \, \mathrm{d}s + \beta(1+\gamma+\gamma^2) \int_0^t \|u_m(s)\|_{L^{2(\delta+1)}}^{2(\delta+1)} \, \mathrm{d}s,$$

for all  $t \in [0, T]$ . Applying Gronwall's inequality, we get

$$\|u_m(t)\|_{\mathrm{L}^{2(\delta+1)}}^{2(\delta+1)} \leqslant \left(\|u_0\|_{\mathrm{L}^{2(\delta+1)}}^{2(\delta+1)} + \frac{2(\delta+1)}{\beta} \int_0^T \|f(t)\|_{\mathrm{L}^2}^2 \mathrm{d}t\right) e^{\beta(1+\gamma+\gamma^2)T},$$

for all  $t \in [0, T]$ . Thus, it is immediate that

$$\sup_{0 \leqslant t \leqslant T} \|u_m(t)\|_{L^{2(\delta+1)}}^{2(\delta+1)} + \nu(2\delta+1)(2(\delta+1)) \int_0^T \||u_m(t)|^{\delta} \nabla u_m(t)\|_{L^2}^2 dt + \frac{\beta(2(\delta+1))}{2} \int_0^T \|u_m(t)\|_{L^{4\delta+2}}^{4\delta+2} dt$$

$$\leqslant \left( \|u_0\|_{L^{2(\delta+1)}}^{2(\delta+1)} + \frac{2(\delta+1)}{\beta} \int_0^T \|f(t)\|_{L^2}^2 dt \right) e^{2\beta(1+\gamma+\gamma^2)T}, \qquad (2.17)$$

and the right hand side of the above inequality is independent of *m*. Since the right hand side of (2.17) is independent of *m*, an application of the Banach-Alaoglu theorem yields  $u \in L^{\infty}(0,T;L^{2(\delta+1)}(\Omega)) \cap L^{2(2\delta+1)}(0,T;L^{2(2\delta+1)}(\Omega))$ .

We will now discuss a result on the uniqueness of the weak solution that we have obtained in Theorem 2.2.

THEOREM 2.3 (Uniqueness of weak solution) For  $u_0 \in L^{d\delta}(\Omega)$ , for d = 2, 3, the weak solution to the system (1.1) obtained in Theorem 2.2 is unique.

*Proof.* Let  $u_1(\cdot)$  and  $u_2(\cdot)$  be two weak solutions of the system (1.1) with the same external forcing  $f(\cdot, \cdot)$  and initial data  $u_0$  with the regularity given in the statement. Then,  $w = u_1 - u_2$  satisfies:

$$\begin{cases} \frac{\partial w(x,t)}{\partial t} + \alpha \left[ u_1(x,t)^{\delta} \sum_{i=1}^d \frac{\partial u_1(x,t)}{\partial x_i} - u_2(x,t)^{\delta} \sum_{i=1}^d \frac{\partial u_2(x,t)}{\partial x_i} \right] - \nu \Delta w(x,t) - \eta \left( K * \Delta w \right)(x,t) \\ = \beta \left[ u_1(x,t)(1 - u_1(x,t)^{\delta})(u_1(x,t)^{\delta} - \gamma) - u_2(x,t)(1 - u_2(x,t)^{\delta})(u_2(x,t)^{\delta} - \gamma) \right], \\ w(x,t) = 0, \ x \in \partial \Omega, \ t \in (0,T), \\ w(x,0) = 0, \end{cases}$$

for  $(x,t) \in \Omega \times (0,T)$  in the weak sense. One can write the above system in the weak sense as

$$\langle w', v \rangle + \alpha(B(u_1) - B(u_2), v) + v(\nabla w, \nabla v) + \eta((K * \nabla w), \nabla v) = \beta \langle c(u_1) - c(u_2), v \rangle,$$
(2.18)

for all  $v \in H_0^1(\Omega) \cap L^{2(\delta+1)}(\Omega)$  and a.e.  $t \in [0,T]$ . Using v = w in (2.18), we find

$$\frac{1}{2}\frac{d}{dt}\|w\|_{L^{2}}^{2}+\nu\|\nabla w\|_{L^{2}}^{2}+\eta((K*\nabla w),\nabla w)=-\alpha(B(u_{1})-B(u_{2}),w)+\beta\langle c(u_{1})-c(u_{2}),w\rangle.$$
 (2.19)

It can be easily seen that

$$\beta(c(u_1) - c(u_2), w) = \beta \left[ (u_1(1 - u_1^{\delta})(u_1^{\delta} - \gamma) - u_2(1 - u_2^{\delta})(u_2^{\delta} - \gamma), w) \right]$$
  
=  $-\beta \gamma ||w||_{L^2}^2 - \beta (u_1^{2\delta+1} - u_2^{2\delta+1}, w) + \beta (1 + \gamma)(u_1^{\delta+1} - u_2^{\delta+1}, w).$  (2.20)

Estimating the term  $-\beta(u_1^{2\delta+1}-u_2^{2\delta+1},w)$  from (2.20) as

$$-\beta(u_{1}^{2\delta+1} - u_{2}^{2\delta+1}, w) = -\beta(u_{1}^{2\delta}, w^{2}) - \beta(u_{2}^{2\delta}, w^{2}) - \beta(u_{1}^{2\delta}u_{2} - u_{2}^{2\delta}u_{1}, w)$$

$$= -\beta \|u_{1}^{\delta}w\|_{L^{2}}^{2} - \beta\|u_{2}^{\delta}w\|_{L^{2}}^{2} - \beta(u_{1}u_{2}, u_{1}^{2\delta} + u_{2}^{2\delta}) + \beta(u_{1}^{2}, u_{2}^{2\delta}) + \beta(u_{2}^{2}, u_{1}^{2\delta})$$

$$= -\frac{\beta}{2} \|u_{1}^{\delta}w\|_{L^{2}}^{2} - \frac{\beta}{2} \|u_{2}^{\delta}w\|_{L^{2}}^{2} - \frac{\beta}{2} (u_{1}^{2\delta} - u_{2}^{2\delta}, u_{1}^{2} - u_{2}^{2})$$

$$\leqslant -\frac{\beta}{2} \|u_{1}^{\delta}w\|_{L^{2}}^{2} - \frac{\beta}{2} \|u_{2}^{\delta}w\|_{L^{2}}^{2}.$$
(2.21)

We estimate the term  $\beta(1+\gamma)(u_1^{\delta+1}-u_2^{\delta+1},w)$  from (2.20), using Taylor's formula, Hölder's and Young's inequalities as

$$\beta(1+\gamma)(u_1^{\delta+1}-u_2^{\delta+1},w) = \beta(1+\gamma)(\delta+1)((\theta_1u_1+(1-\theta)u_2)^{\delta}w,w)$$

$$\leq \beta (1+\gamma)(\delta+1)2^{\delta-1} (\|u_1^{\delta}w\|_{L^2} + \|u_2^{\delta}w\|_{L^2})\|w\|_{L^2} \leq \frac{\beta}{4} \|u_1^{\delta}w\|_{L^2}^2 + \frac{\beta}{4} \|u_2^{\delta}w\|_{L^2}^2 + \frac{\beta}{2} 2^{2\delta} (1+\gamma)^2 (\delta+1)^2 \|w\|_{L^2}^2.$$
 (2.22)

Combining (2.21)-(2.22) and substituting it in (2.20), we obtain

$$\beta \left[ (u_1(1-u_1^{\delta})(u_1^{\delta}-\gamma)-u_2(1-u_2^{\delta})(u_2^{\delta}-\gamma),w) \right] \\ \leqslant -\beta \gamma \|w\|_{L^2}^2 - \frac{\beta}{4} \|u_1^{\delta}w\|_{L^2}^2 - \frac{\beta}{4} \|u_2^{\delta}w\|_{L^2}^2 + \frac{\beta}{2} 2^{2\delta} (1+\gamma)^2 (\delta+1)^2 \|w\|_{L^2}^2.$$
(2.23)

On the other hand, we derive a bound for  $-\alpha \langle B(u_1) - B(u_2), w \rangle$ , using Taylor's formula, Hölder's, Ladyzhenskaya's and Young's inequalities as

$$-\alpha \langle B(u_{1}) - B(u_{2}), w \rangle = \frac{\alpha}{\delta + 1} \left( (u_{1}^{\delta + 1} - u_{2}^{\delta + 1}) \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix}, \nabla w \right)$$

$$= \alpha \left( (u_{1} - u_{2})(\theta u_{1} + (1 - \theta)u_{2})^{\delta} \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix}, \nabla w \right)$$

$$\leq 2^{\delta - 1} \alpha (\|u_{1}\|_{L^{4\delta}}^{\delta} + \|u_{2}\|_{L^{4\delta}}^{\delta}) \|w\|_{L^{4}} \|\nabla w\|_{L^{2}}$$

$$\leq 2^{\delta - 1} \alpha (\|u_{1}\|_{L^{4\delta}}^{\delta} + \|u_{2}\|_{L^{4\delta}}^{\delta}) \|w\|_{L^{4}}^{\frac{4 - d}{4}} \|\nabla w\|_{L^{2}}$$

$$\leq \frac{\nu}{2} \|\nabla w\|_{L^{2}}^{2} + C(\alpha, \nu) \left( \|u_{1}\|_{L^{4\delta}}^{\frac{8 \delta}{4 - d}} + \|u_{2}\|_{L^{4\delta}}^{\frac{8 \delta}{4 - d}} \right) \|w\|_{L^{2}}^{2}.$$
(2.24)

where  $C(\alpha, \nu) = \left(\frac{4+d}{4\nu}\right)^{\frac{4+d}{4-d}} \left(\frac{4-d}{8}\right) \left(2^{\delta-1}\alpha\right)^{\frac{8}{4-d}}$ . Combining (2.23) and (2.24), we have

$$\frac{\mathrm{d}}{\mathrm{d}t} \|w(t)\|_{\mathrm{L}^{2}}^{2} + \nu \|\nabla w(t)\|_{\mathrm{L}^{2}}^{2} + \frac{\beta}{2} \|u_{1}^{\delta}(t)w(t)\|_{\mathrm{L}^{2}}^{2} + \frac{\beta}{2} \|u_{2}^{\delta}(t)w(t)\|_{\mathrm{L}^{2}}^{2} + \beta\gamma \|w(t)\|_{\mathrm{L}^{2}}^{2} \\
\leqslant \beta 2^{2\delta+1} (1+\gamma)^{2} (\delta+1)^{2} \|w(t)\|_{\mathrm{L}^{2}}^{2} + C(\alpha,\nu) \left( \|u_{1}(t)\|_{\mathrm{L}^{4\delta}}^{\frac{8\delta}{4-d}} + \|u_{2}(t)\|_{\mathrm{L}^{4\delta}}^{\frac{8\delta}{4-d}} \right) \|w(t)\|_{\mathrm{L}^{2}}^{2}, \qquad (2.25)$$

for a.e.  $t \in [0,T]$ . An application of Gronwall's inequality in (2.25) gives for all  $t \in [0,T]$ 

$$\|w(t)\|_{L^{2}}^{2} \leq \|w(0)\|_{L^{2}}^{2} e^{\beta 2^{2\delta}(1+\gamma)^{2}(\delta+1)^{2}T} \exp\left\{C(\alpha, \nu) \int_{0}^{T} \left(\|u_{1}(t)\|_{L^{4\delta}}^{\frac{8\delta}{4-d}} + \|u_{2}(t)\|_{L^{4\delta}}^{\frac{8\delta}{4-d}}\right) dt\right\}.$$
 (2.26)

For d = 2 and  $u_0 \in L^{2\delta}(\Omega)$ , from Remark 2.2, it is clear that  $u_1, u_2 \in L^{4\delta}(0, T; L^{4\delta}(\Omega))$ , and hence the uniqueness follows. For d = 3,  $u_0 \in L^{3\delta}(\Omega)$ , the integral appearing in the exponential is finite by using interpolation inequality as

$$\int_0^T \|u_1(t)\|_{\mathrm{L}^{4\delta}}^{8\delta} dt \leq \sup_{t \in [0,T]} \|u_1(t)\|_{\mathrm{L}^{3\delta}}^{3\delta} \int_0^T \|u_1(t)\|_{\mathrm{L}^{5\delta}}^{5\delta} dt < \infty.$$

Hence, the uniqueness follows if we take the initial data  $u_0 \in L^{d\delta}(\Omega)$ .

REMARK 2.3 For d = 3, using Gagliardo-Nirenberg's inequality in (2.24), one can find a sufficient criterion for the uniqueness of weak solution for  $u_0 \in L^{3\delta}(\Omega)$  as

$$\int_0^T \|u(t)\|_{\mathrm{L}^{r\delta}}^{s\delta} \mathrm{d} t < \infty, \text{ where } \frac{3}{r} + \frac{2}{s} = 1, \ r \in (3,\infty).$$

REMARK 2.4 If we consider our initial data  $u_0 \in L^2(\Omega)$  and  $\beta \nu > (2^{\delta} \alpha)^2$ , then there exists a unique weak solution to the system (1.1). One can estimate  $-\alpha \langle B(u_1) - B(u_2), w \rangle$  in (2.24) using integrating by parts, Taylor's formula, Hölder's and Young's inequalities as:

$$-\alpha \langle B(u_{1}) - B(u_{2}), w \rangle = \frac{\alpha}{\delta + 1} \left( (u_{1}^{\delta + 1} - u_{2}^{\delta + 1}) \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix}, \nabla w \right)$$
$$= \alpha \left( (u_{1} - u_{2})(\theta u_{1} + (1 - \theta)u_{2})^{\delta} \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix}, \nabla w \right)$$
$$\leq 2^{\delta - 1} \alpha \|\nabla w\|_{L^{2}} \left( \|u_{1}^{\delta} w\|_{L^{2}} + \|u_{2}^{\delta} w\|_{L^{2}} \right)$$
$$\leq \frac{\nu}{2} \|\nabla w\|_{L^{2}}^{2} + \frac{2^{2\delta} \alpha^{2}}{4\nu} \|u_{1}^{\delta} w\|_{L^{2}}^{2} + \frac{2^{2\delta} \alpha^{2}}{4\nu} \|u_{2}^{\delta} w\|_{L^{2}}^{2}.$$
(2.27)

Combining (2.23) and (2.27) in (2.19), we find

$$\begin{split} &\frac{1}{2} \frac{\mathrm{d}}{\mathrm{d}t} \|w(t)\|_{\mathrm{L}^{2}}^{2} + \frac{\nu}{2} \|\nabla w(t)\|_{\mathrm{L}^{2}}^{2} + \frac{\beta}{4} \|u_{1}^{\delta}(t)w(t)\|_{\mathrm{L}^{2}}^{2} + \frac{\beta}{4} \|u_{2}(t)^{\delta}w(t)\|_{\mathrm{L}^{2}}^{2} + \beta\gamma \|w(t)\|_{\mathrm{L}^{2}}^{2} + \eta((K*\nabla w)(t),\nabla w(t)) \\ &\leq \frac{\beta}{2} 2^{2\delta} (1+\gamma)^{2} (\delta+1)^{2} \|w(t)\|_{\mathrm{L}^{2}}^{2} + \frac{\alpha^{2}}{4\nu} 2^{2\delta} \|u_{1}^{\delta}w(t)\|_{\mathrm{L}^{2}}^{2} + \frac{\alpha^{2}}{4\nu} 2^{2\delta} \|u_{2}(t)^{\delta}w(t)\|_{\mathrm{L}^{2}}^{2}. \end{split}$$

From the above expression, it is immediate that

$$\begin{split} & \frac{\mathrm{d}}{\mathrm{d}t} \|w(t)\|_{\mathrm{L}^{2}}^{2} + v \|\nabla w(t)\|_{\mathrm{L}^{2}}^{2} + \left(\frac{\beta}{2} - \frac{\alpha^{2}}{2v}2^{2\delta}\right) \|u_{1}(t)^{\delta}w(t)\|_{\mathrm{L}^{2}}^{2} \\ & + \left(\frac{\beta}{2} - \frac{\alpha^{2}}{2v}2^{2\delta}\right) \|u_{2}(t)^{\delta}w(t)\|_{\mathrm{L}^{2}}^{2} + \eta\left((K * \nabla w)(t), \nabla w(t)\right) \\ & \leq \beta 2^{2\delta}(1 + \gamma)^{2}(\delta + 1)^{2} \|w(t)\|_{\mathrm{L}^{2}}^{2}. \end{split}$$

For  $\beta v > (2^{\delta} \alpha)^2$ , integrating the above inequality from 0 to *t*, (1.2) and then applying of Gronwall's inequality results to

$$\|w(t)\|_{\mathrm{L}^{2}}^{2} \leq \|w(0)\|_{\mathrm{L}^{2}}^{2} e^{\beta 2^{2\delta}(1+\gamma)^{2}(\delta+1)^{2}T},$$

for all  $t \in [0, T]$ . Since  $w_0 = 0$  and  $u_1$  and  $u_2$  are weak solutions of the system (1.1), uniqueness follows easily.

# 2.4 Regularity results

If we take our initial data  $u_0 \in H_0^1(\Omega) \cap L^{2(\delta+1)}(\Omega)$ , and the external forcing,  $f \in L^2(0,T; L^2(\Omega))$ , where  $\Omega \subset \mathbb{R}^d$ , d = 2, 3, is either convex, or a domain with C<sup>2</sup>-boundary, then we can obtain the  $H_0^1(\Omega)$ energy estimate. Under smoothness assumptions on the domain, initial data and external forcing, let us now establish the regularity of the weak solution obtained in Theorem 2.2.

THEOREM 2.4 (Regularity) Let *u* be a weak solution to the system (1.1). If  $\Omega \subset \mathbb{R}^d$ , d = 2, 3, is either convex, or a domain with  $C^2$ -boundary and let  $u_0 \in H_0^1(\Omega) \cap L^{2(\delta+1)}(\Omega)$  and  $f \in L^2(0,T; L^2(\Omega))$  be given. Then, *u* is a strong solution with

$$u \in \mathrm{C}([0,T];\mathrm{H}^1_0(\Omega)) \cap \mathrm{L}^2(0,T;\mathrm{H}^2(\Omega)) \cap \mathrm{L}^{2(\delta+1)}(0,T;\mathrm{L}^{6(\delta+1)}(\Omega)),$$

and  $\partial_t u \in L^2(0,T; L^2(\Omega))$  and the following system is satisfied:

$$\begin{cases} \frac{\mathrm{d}u(t)}{\mathrm{d}t} + vAu(t) + \eta(K * Au)(s) = -\alpha B(u(t)) + \beta c(u(t)) + f(t), & \text{in } \mathrm{L}^{2}(\Omega), \\ u(0) = u_{0} \in \mathrm{L}^{2}(\Omega), \end{cases}$$

for a.e.  $t \in [0, T]$ .

Proof.

**Step 1:**  $H_0^1$ -energy estimate. Let  $\lambda_k$  denote the  $k^{th}$  eigenvalue of  $-\Delta$  in  $H_0^1(\Omega)$ . Multiplying the identity (2.7) by  $\lambda_k d_k^m(t)$  and summing it from k = 1, 2, ..., m, we have

$$(\partial_t u_m, Au_m) - \nu(\Delta u_m, Au_m) - \eta(K * \Delta u_m, Au_m) = -\alpha(B(u_m), Au_m) + \beta(c(u_m), Au_m) + (f, Au_m).$$
(2.28)

Now, using integrating by parts and applying Hölder's and Young's inequalities, we find

$$\alpha |(B(u_m), Au_m)| \leq \alpha ||B(u_m)||_{L^2} ||Au_m||_{L^2} \leq \alpha ||u_m^{\delta} \nabla u_m||_{L^2} ||Au_m||_{L^2} \leq \frac{\nu}{4} ||Au_m||_{L^2}^2 + \frac{\alpha^2}{\nu} ||u_m^{\delta} \nabla u_m||_{L^2}^2.$$

Estimating  $\beta(c(u_m), Au_m)$  using Hölder's and Young's inequalities as follows:

$$\begin{split} &\beta(c(u_m), Au_m) \\ &= \beta((1+\gamma)u_m^{\delta+1} - \gamma u_m - u_m^{2\delta+1}, Au_m) \\ &= \beta((1+\gamma)(\delta+1)u_m^{\delta}\nabla u_m - \gamma \nabla u_m - (2\delta+1)u_m^{2\delta}\nabla u_m, \nabla u_m) \\ &= \beta(1+\gamma)(\delta+1)(u_m^{\delta}\nabla u_m, \nabla u_m) - \beta\gamma(\nabla u_m, \nabla u_m) - \beta(2\delta+1)(u_m^{2\delta}\nabla u_m, \nabla u_m) \\ &\leqslant \beta(1+\gamma)(\delta+1) \|u_m^{\delta}\nabla u_m\|_{L^2} \|\nabla u_m\|_{L^2} - \beta\gamma\|\nabla u_m\|_{L^2}^2 - \beta(2\delta+1) \|u_m^{\delta}\nabla u_m\|_{L^2}^2 \\ &\leqslant \frac{\beta(1+\gamma)^2(\delta+1)^2}{2(2\delta+1)} \|\nabla u_m\|_{L^2}^2 + \frac{\beta(2\delta+1)}{2} \|u_m^{\delta}\nabla u_m\|_{L^2}^2 - \beta\gamma\|\nabla u_m\|_{L^2}^2 - \beta(2\delta+1) \|u_m^{\delta}\nabla u_m\|_{L^2}^2 \\ &\leqslant \frac{\beta(1+\gamma)^2(\delta+1)^2}{2(2\delta+1)} \|\nabla u_m\|_{L^2}^2 - \beta\gamma\|\nabla u_m\|_{L^2}^2 - \frac{\beta(2\delta+1)}{2} \|u_m^{\delta}\nabla u_m\|_{L^2}^2. \end{split}$$

Combining the above estimates and substituting in (2.28), we get

$$\frac{1}{2} \frac{\mathrm{d}}{\mathrm{d}t} \|\nabla u_m(t)\|_{\mathrm{L}^2}^2 + \frac{v}{2} \|\Delta u_m(t)\|_{\mathrm{L}^2}^2 + \frac{\beta(2\delta+1)}{2} \|u_m(t)^{\delta} \nabla u_m(t)\|_{\mathrm{L}^2}^2 + \eta((K * \Delta u_m)(t), \Delta u_m(t))$$

GENERALIZED BURGERS-HUXLEY EQUATION WITH WEAKLY SINGULAR KERNEL

$$\leq \frac{\alpha^2}{\nu} \|u_m(t)^{\delta} \nabla u_m(t)\|_{\mathrm{L}^2}^2 + \frac{\beta((1+\gamma^2)(\delta+1)^2+2\gamma\delta^2)}{2(2\delta+1)} \|\nabla u_m(t)\|_{\mathrm{L}^2}^2 + \frac{1}{\nu} \|f(t)\|_{\mathrm{L}^2}^2.$$

Integrating from 0 to t and using (1.2), we find

$$\begin{aligned} \|\nabla u_{m}(t)\|_{L^{2}}^{2} + \nu \int_{0}^{t} \|\Delta u_{m}(s)\|_{L^{2}}^{2} ds + \beta (2\delta + 1) \int_{0}^{t} \|u_{m}(s)^{\delta} \nabla u_{m}(s)\|_{L^{2}}^{2} ds \\ &\leq \|u_{0}\|_{H^{1}_{0}}^{2} + \frac{2\alpha^{2}}{\nu} \int_{0}^{t} \|u_{m}(s)^{\delta} \nabla u_{m}(s)\|_{L^{2}}^{2} ds + \frac{\beta ((1 + \gamma^{2})(\delta + 1)^{2} + 2\gamma\delta^{2})}{(2\delta + 1)} \int_{0}^{t} \|\nabla u_{m}(s)\|_{L^{2}}^{2} ds \\ &+ \frac{2}{\nu} \int_{0}^{t} \|f(s)\|_{L^{2}}^{2} ds \\ &\leq C \left(\|u_{0}\|_{H^{1}_{0}}, \|f\|_{L^{2}(0,T;L^{2}(\Omega))}, \alpha, \beta, \gamma, \delta, \nu, T\right), \end{aligned}$$

$$(2.29)$$

since  $u_m(\cdot)$  satisfies the energy estimates (2.9) and (2.17). Whenever  $u_0 \in H_0^1(\Omega)$  and  $f \in L^2(0,T; L^2(\Omega))$ , from (2.29), we have

$$u_m \in \mathcal{L}^{\infty}(0,T;\mathcal{H}^1_0(\Omega)) \cap \mathcal{L}^2(0,T;\mathcal{H}^2(\Omega))$$

by using the elliptic regularity (Theorem 3.1.2.1, Grisvard (1985)). Using Sobolev's inequality,  $||u||_{L^6} \leq C ||\nabla u||_{L^2}$ , we have  $||u_m^{\delta+1}||_{L^6}^2 \leq C ||\nabla u_m^{\delta+1}||_{L^2}^2 \leq C(\delta+1) ||u_m^{\delta} \nabla u_m||_{L^2}$ , so that

$$\int_0^t \|u_m(s)\|_{\mathrm{L}^{6(\delta+1)}}^{2(\delta+1)} \mathrm{d}s = \int_0^t \|u_m(s)^{\delta+1}\|_{\mathrm{L}^6}^2 \mathrm{d}s \leqslant C(\delta+1)^2 \int_0^t \|u_m(s)^{\delta} \nabla u_m(s)\|_{\mathrm{L}^2}^2 \mathrm{d}s < \infty.$$

Thus, we infer

$$u_m \in \mathcal{L}^{2(\delta+1)}(0,T;\mathcal{L}^{6(\delta+1)}(\Omega)).$$

**Step 2:** Estimate for time derivative. For  $m \ge 1$ , multiplying (2.7) by  $d_m^{k'}(t)$  and sum it over k = 1, 2, ..., m, we have

$$\begin{aligned} \|\partial_{t}u_{m}(t)\|_{L^{2}}^{2} &+ \frac{\nu}{2} \frac{d}{dt} \|\nabla u_{m}(t)\|_{L^{2}}^{2} \\ &= \eta((K * \Delta u_{m})(t), \partial_{t}u_{m}(t)) - \alpha \left( (u_{m}(t))^{\delta} \sum_{i=1}^{d} \frac{\partial u_{m}}{\partial x_{i}}(t), \partial_{t}u_{m}(t) \right) \\ &+ \beta (u_{m}(t)(1 - (u_{m}(t))^{\delta})((u_{m}(t))^{\delta} - \gamma), \partial_{t}u_{m}(t)) + (f(t), \partial_{t}u_{m}(t)) \\ &= \eta((K * \Delta u_{m})(t), \partial_{t}u_{m}(t)) - \alpha \left( (u_{m}(t))^{\delta} \sum_{i=1}^{d} \frac{\partial u_{m}}{\partial x_{i}}(t), \partial_{t}u_{m}(t) \right) + \beta (1 + \gamma)(u_{m}(t)^{\delta+1}, \partial_{t}u_{m}(t)) \\ &- \beta \gamma \frac{d}{dt} \|u_{m}(t)\|_{L^{2}}^{2} - \frac{\beta}{2(\delta+1)} \frac{d}{dt} \|u_{m}(t)\|_{L^{2(\delta+1)}}^{2(\delta+1)} + (f(t), \partial_{t}u_{m}(t)). \end{aligned}$$
(2.30)

Using Hölder's and Young's inequalities, we estimate the first term from the right hand side of the equality (2.30) as

$$\alpha\left((u_m)^{\delta}\sum_{i=1}^d \frac{\partial u_m}{\partial x_i}, \partial_t u_m\right) \leqslant \frac{1}{4} \|\partial_t u_m\|_{L^2}^2 + \alpha^2 \|u_m^{\delta}\nabla u_m\|_{L^2}^2.$$
(2.31)

Similarly, we estimate  $\beta(1+\gamma)(u_m^{\delta+1},\partial_t u_m)$  and  $\eta(K * \Delta u_m,\partial_t u_m)$  as

$$\beta(1+\gamma)|(u_{m}^{\delta+1},\partial_{t}u_{m})| \leq \frac{1}{4} \|\partial_{t}u_{m}\|_{L^{2}}^{2} + \beta^{2}(1+\gamma)^{2}\|u_{m}\|_{L^{2}(\delta+1)}^{2(\delta+1)},$$
  
$$\eta(K*\Delta u_{m},\partial_{t}u_{m}) \leq \frac{\eta^{2}}{2}\|K*\Delta u_{m}\|_{L^{2}}^{2} + \frac{1}{2}\|\partial_{t}u_{m}\|_{L^{2}}^{2},$$
(2.32)

The final term from the right hand side of the equality (2.30) can be estimated as

$$|(f,\partial_t u_m)| \leq ||f||_{\mathbf{L}^2} ||\partial_t u_m||_{\mathbf{L}^2} \leq \frac{1}{4} ||\partial_t u_m||_{\mathbf{L}^2}^2 + ||f||_{\mathbf{L}^2}^2.$$
(2.33)

Combining (2.31)-(2.33), substituting it in (2.30) and then integrating from 0 to t, we get

$$\begin{aligned} &2\mathbf{v} \|\nabla u_m(t)\|_{L^2}^2 + \frac{2\beta}{\delta+1} \|u_m(t)\|_{L^{2(\delta+1)}}^{2(\delta+1)} + 4\beta\gamma \|u_m(t)\|_{L^2}^2 + \frac{1}{2} \int_0^t \|\partial_t u_m(s)\|_{L^2}^2 \,\mathrm{d}s \\ &\leqslant 2\mathbf{v} \|u_0\|_{\mathrm{H}_0^1}^2 + \frac{2\beta}{\delta+1} \|u_0\|_{L^{2(\delta+1)}}^{2(\delta+1)} + 4\beta\gamma \|u_0\|_{L^2}^2 + 4 \int_0^t \|f(s)\|_{L^2}^2 \,\mathrm{d}s + 4\alpha^2 \int_0^t \|u_m^{\delta}(s)\nabla u_m(s)\|_{L^2}^2 \,\mathrm{d}s \\ &+ \frac{\eta^2 C}{2} \int_0^t \|\Delta u_m(s)\|_{L^2}^2 \,\mathrm{d}s + 4\beta^2 (1+\gamma)^2 \int_0^t \|u_m(s)\|_{L^{2(\delta+1)}}^{2(\delta+1)} \,\mathrm{d}s. \end{aligned}$$

Applying Gronwall's inequality, we arrive at

$$2\nu \|\nabla u_{m}(t)\|_{L^{2}}^{2} + \frac{2\beta}{\delta+1} \|u_{m}(t)\|_{L^{2(\delta+1)}}^{2(\delta+1)} + 4\beta\gamma \|u_{m}(t)\|_{L^{2}}^{2} + \frac{1}{2}\int_{0}^{t} \|\partial_{t}u_{m}(s)\|_{L^{2}}^{2} ds$$

$$\leq \left(2\nu \|u_{0}\|_{H_{0}^{1}}^{2} + \frac{2\beta}{\delta+1} \|u_{0}\|_{L^{2(\delta+1)}}^{2(\delta+1)} + 4\beta\gamma \|u_{0}\|_{L^{2}}^{2} + 4\int_{0}^{T} \|f(t)\|_{L^{2}}^{2} dt + 4\alpha^{2}\int_{0}^{T} \|u_{m}^{\delta}(t)\nabla u_{m}(t)\|_{L^{2}}^{2} dt\right)$$

$$\times \exp\left(4\beta(1+\gamma)^{2}(1+\delta)T\right)e^{\frac{\eta^{2}C}{2\nu}T},$$
(2.34)

for all  $t \in [0, T]$ . Using the estimate given in (2.29), we obtain that the right hand side of (2.34) is independent of *m*. Thus, it is immediate that  $\partial_t u_m \in L^2(0, T; L^2(\Omega))$ . Using arguments similar to Steps 4 and 5 in Theorem 2.2, we obtain the existence of a  $\tilde{u}(\cdot)$  such that

$$\begin{split} \tilde{u} \in \mathrm{L}^{\infty}(0,T;\mathrm{H}^{1}_{0}(\Omega) \cap \mathrm{L}^{2(\delta+1)}(\Omega)) \cap \mathrm{L}^{2}(0,T;\mathrm{H}^{2}(\Omega)) \cap \mathrm{L}^{2(\delta+1)}(0,T;\mathrm{L}^{6(\delta+1)}(\Omega)) \quad \text{and} \\ \partial_{t}\tilde{u} \in \mathrm{L}^{2}(0,T;\mathrm{L}^{2}(\Omega)), \end{split}$$

so that we get  $\tilde{u} \in C([0,T]; H_0^1(\Omega))$  and is a strong solution to the system (1.1). The uniqueness follows from the estimate (2.26) and we have  $\tilde{u} = u$ .

THEOREM 2.5 Let  $1 \leq \delta < \infty$  for d = 2 and,  $1 \leq \delta \leq 2$  for d = 3 and assume that  $u_0 \in H^2(\Omega) \cap H^1_0(\Omega)$ 

(i) If 
$$f \in H^1(0,T;L^2(\Omega))$$
, then, we have  $\partial_t u \in L^{\infty}(0,T;L^2(\Omega)) \cap L^2(0,T;H^1_0(\Omega))$ .

(ii) If  $f \in H^1(0,T; L^2(\Omega)) \cap L^2(0,T; H^1(\Omega))$ , then  $u \in L^{\infty}(0,T; H^2(\Omega))$ .

*Proof.* We divide the proof into the following steps:

**Step 1.**  $\partial_t u \in L^{\infty}(0,T;L^2(\Omega))$ , and  $\partial_t \nabla u \in L^2(0,T;L^2(\Omega))$ .

Let  $1 \leq \delta < \infty$  for d = 2 and,  $1 \leq \delta \leq 2$  for d = 3. Differentiating (2.7) with respect to t, we find

$$(\partial_{tt}u_m(t), w_k) + \mathbf{v}(\partial_t \nabla u_m(t), \nabla w_k)$$

# GENERALIZED BURGERS-HUXLEY EQUATION WITH WEAKLY SINGULAR KERNEL 19 of 45

$$+\alpha \sum_{i=1}^{d} \left( (u_m(t))^{\delta} \partial_{tx_i} u_m(t) + \delta u_m(t)^{\delta-1} \partial_t u_m(t) \partial_{x_i} u_m(t), w_k \right) - \eta \left( (K * \partial_t \Delta u_m)(t) + K(t) \Delta u_m(0), w_k \right)$$
$$= \beta \left( (\gamma+1)(\delta+1)u_m(t)^{\delta} \partial_t u_m(t) - \gamma \partial_t u_m(t) - (2\delta+1)u_m(t)^{2\delta} \partial_t u_m(t), w_k \right) + (\partial_t f(t), w_k).$$
(2.35)

Multiplying (2.35) with  $d_m^{k'}(\cdot)$  and summing it over k = 1, 2, ..., m, we obtain

$$\frac{1}{2} \frac{d}{dt} \|\partial_{t} u_{m}(t)\|_{L^{2}}^{2} + v \|\partial_{t} \nabla u_{m}(t)\|_{L^{2}}^{2} + \beta \gamma \|\partial_{t} u_{m}(t)\|_{L^{2}}^{2} + \beta (2\delta + 1) \|u_{m}(t)^{\delta} \partial_{t} u_{m}(t)\|_{L^{2}}^{2} 
+ \eta ((K * \partial_{t} \nabla u_{m})(t), \partial_{t} \nabla u_{m}(t)) 
= -\alpha \sum_{i=1}^{d} \left( (u_{m}(t))^{\delta} \partial_{tx_{i}} u_{m}(t), \partial_{t} u_{m} \right) + \beta (\gamma + 1) (\delta + 1) (u_{m}(t)^{\delta} \partial_{t} u_{m}(t), \partial_{t} u_{m}(t)) 
+ (\partial_{t} f(t), \partial_{t} u_{m}(t)) + \eta (K(t) \Delta u_{m}(0), \partial_{t} u_{m}(t)).$$
(2.36)

We estimate the term  $-\alpha \sum_{i=1}^{d} ((u_m)^{\delta} \partial_{tx_i} u_m, \partial_t u_m)$  using integration by parts, Hölder's, Ladyzhenskaya's inequality and Young's inequalities as

$$\begin{split} &-\sum_{i=1}^{d} \alpha \left( (u_{m})^{\delta} \partial_{tx_{i}} u_{m}, \partial_{t} u_{m} \right) \\ &= -\frac{1}{2} \sum_{i=1}^{d} \alpha \left( u_{m}^{\delta}, \frac{\partial}{\partial x_{i}} \left( \partial_{t} u_{m} \right)^{2} \right) = \frac{\alpha \delta}{2} \sum_{i=1}^{d} \left( u_{m}^{\delta-1} \frac{\partial u_{m}}{\partial x_{i}}, \left( \partial_{t} u_{m} \right)^{2} \right) \\ &= \frac{\alpha \delta}{2} \sum_{i=1}^{d} \int_{\Omega} u_{m}^{\delta-1} \left( \frac{\partial u_{m}}{\partial x_{i}} \right)^{\frac{\delta-1}{\delta}} \left( \frac{\partial u_{m}}{\partial x_{i}} \right)^{\frac{1}{\delta}} \left( \partial_{t} u_{m} \right)^{2} dx \\ &\leq \frac{\alpha \delta}{2} \sum_{i=1}^{d} \left( \int_{\Omega} u_{m}^{2\delta} \left( \frac{\partial u_{m}}{\partial x_{i}} \right)^{2} dx \right)^{\frac{\delta-1}{2\delta}} \left( \int_{\Omega} \left( \frac{\partial u_{m}}{\partial x_{i}} \right)^{\frac{2}{\delta+1}} \left( \partial_{t} u_{m} \right)^{\frac{4\delta}{\delta+1}} dx \right)^{\frac{\delta+1}{2\delta}} \\ &\leq \frac{\alpha \delta}{2} \| u_{m}^{\delta} \nabla u_{m} \|_{L^{2}}^{\frac{\delta-1}{\delta}} \| (\nabla u_{m})^{\frac{1}{\delta+1}} \left( \partial_{t} u_{m} \right)^{\frac{2\delta}{\delta+1}} \|_{L^{2}}^{\frac{\delta+1}{\delta}} \\ &\leq \frac{\alpha \delta}{2} \| u_{m}^{\delta} \nabla u_{m} \|_{L^{2}}^{\frac{\delta-1}{\delta}} \| \nabla u_{m} \|_{L^{2}}^{\frac{1}{\delta}} \| \partial_{t} u_{m} \|_{L^{2}}^{2} \\ &\leq \frac{\alpha \delta}{\sqrt{2}} \| u_{m}^{\delta} \nabla u_{m} \|_{L^{2}}^{\frac{\delta-1}{\delta}} \| \nabla u_{m} \|_{L^{2}}^{\frac{1}{\delta}} \| \partial_{t} u_{m} \|_{L^{2}}^{\frac{4-d}{\delta}} \\ &\leq \frac{\alpha \delta}{\sqrt{2}} \| u_{m}^{\delta} \nabla u_{m} \|_{L^{2}}^{\frac{\delta-1}{\delta}} \| \nabla u_{m} \|_{L^{2}}^{\frac{1}{\delta}} \| \partial_{t} u_{m} \|_{L^{2}}^{\frac{4-d}{\delta}} \\ &\leq \frac{\nu}{2} \| \partial_{t} \nabla u_{m} \|_{L^{2}}^{\frac{2}{\delta}} + \left( \frac{4-d}{4} \right) \left( \frac{d}{2\nu} \right)^{\frac{d}{4-d}} \left( \frac{\alpha \delta}{\sqrt{2}} \right)^{\frac{4-d}{4-d}} \| u_{m}^{\delta} \nabla u_{m} \|_{L^{2}}^{\frac{4(\delta-1)}{\delta}} \| \nabla u_{m} \|_{L^{2}}^{\frac{4}{\delta}} \| \partial_{t} u_{m} \|_{L^{2}}^{\frac{2}{\delta}}. \tag{2.37}$$

We estimate  $\eta |(K(t)\Delta u_m(0), \partial_t u_m)|$  as

$$\eta |(K(t)\Delta u_m(0), \partial_t u_m(t))| \leq \eta |K(t)| ||\Delta u_m(0)||_{L^2} ||\partial_t u_m(t)||_{L^2} \leq \frac{\eta}{2} |K(t)| (||\Delta u_m(0)||_{L^2}^2 + ||\partial_t u_m(t)||_{L^2}^2).$$

Similarly, we estimate the terms  $\beta(\gamma+1)(\delta+1)(u_m^{\delta}\partial_t u_m, \partial_t u_m)$  and  $(\partial_t f, \partial_t u_m)$  as

$$\beta(\gamma+1)(\delta+1)|(u_m^{\delta}\partial_t u_m,\partial_t u_m)| \leq \beta(\gamma+1)(\delta+1)||u_m^{\delta}\partial_t u_m||_{L^2}||\partial_t u_m||_{L^2}$$

$$\leq \frac{\beta(2\delta+1)}{2} \|u_{m}^{\delta}\partial_{t}u_{m}\|_{L^{2}}^{2} + \frac{\beta(\gamma+1)^{2}(\delta+1)^{2}}{2(2\delta+1)} \|\partial_{t}u_{m}\|_{L^{2}}^{2},$$
$$|(\partial_{t}f,\partial_{t}u_{m})| \leq \|\partial_{t}f\|_{L^{2}} \|\partial_{t}u_{m}\|_{L^{2}} \leq \frac{1}{2\beta\gamma} \|\partial_{t}f\|_{L^{2}}^{2} + \frac{\beta\gamma}{2} \|\partial_{t}u_{m}\|_{L^{2}}^{2}.$$
(2.38)

Substituting the estimates (2.37)-(2.38) in (2.36), we have

$$\begin{aligned} \frac{\mathrm{d}}{\mathrm{d}t} \|\partial_{t}u_{m}(t)\|_{\mathrm{L}^{2}}^{2} + \nu \|\partial_{t}\nabla u_{m}(t)\|_{\mathrm{L}^{2}}^{2} + \beta \gamma \|\partial_{t}u_{m}(t)\|_{\mathrm{L}^{2}}^{2} + \beta (2\delta + 1)\|u_{m}(t)^{\delta}\partial_{t}u_{m}(t)\|_{\mathrm{L}^{2}}^{2} \\ + 2\eta ((K * \partial_{t}\nabla u_{m})(t), \partial_{t}\nabla u_{m}(t)) \\ \leqslant \eta |K(t)|\|\Delta u_{m}(0)\|_{\mathrm{L}^{2}}^{2} + \frac{1}{\beta \gamma} \|\partial_{t}f(t)\|_{\mathrm{L}^{2}}^{2} \\ + \left(C(\alpha, \nu, \delta)\|u_{m}(t)^{\delta}\nabla u_{m}(t)\|_{\mathrm{L}^{2}}^{\frac{4(\delta-1)}{(4-d)\delta}} \|\nabla u_{m}(t)\|_{\mathrm{L}^{2}}^{\frac{4}{\delta(4-d)}} + \eta |K(t)| + C(\beta, \gamma, \delta)\right) \|\partial_{t}u_{m}(t)\|_{\mathrm{L}^{2}}^{2}, \end{aligned}$$

where  $C(\alpha, \nu, \delta) = \left(\frac{4-d}{2}\right) \left(\frac{d}{2\nu}\right)^{\frac{d}{4-d}} \left(\frac{\alpha\delta}{\sqrt{2}}\right)^{\frac{4}{4-d}}$  and  $C(\beta, \gamma, \delta) = \frac{\beta(\gamma+1)^2(\delta+1)^2}{(2\delta+1)}$ .

$$\begin{aligned} \|\partial_{t}u_{m}(t)\|_{L^{2}}^{2} + \nu \int_{0}^{t} \|\partial_{t}\nabla u_{m}(s)\|_{L^{2}}^{2} ds + \beta\gamma \int_{0}^{t} \|\partial_{t}u_{m}(s)\|_{L^{2}}^{2} ds + \beta(2\delta+1) \int_{0}^{t} \|u_{m}(s)^{\delta}\partial_{t}u_{m}(s)\|_{L^{2}}^{2} ds \\ &\leq \|\partial_{t}u_{m}(0)\|_{L^{2}}^{2} + \eta \|\Delta u_{m}(0)\|_{L^{2}}^{2} \int_{0}^{t} |K(s)| ds + \frac{1}{\beta\gamma} \int_{0}^{t} \|\partial_{t}f(s)\|_{L^{2}}^{2} ds \\ &+ \int_{0}^{t} \left( C(\alpha,\nu,\delta) \|u_{m}(s)^{\delta}\nabla u_{m}(s)\|_{L^{2}}^{\frac{4(\delta-1)}{(4-d)\delta}} \|\nabla u_{m}(s)\|_{L^{2}}^{\frac{4}{\delta(4-d)}} + \eta |K(s)| + C(\beta,\gamma,\delta) \right) \|\partial_{t}u_{m}(s)\|_{L^{2}}^{2} ds. \end{aligned}$$

As  $K(t) \in L^1(0,T)$  and note that  $||u_m(0)||_{H^2} \leq C ||u_0||_{H^2}$  (see page 385, Evans (2010)) and an application of Gronwall's inequality yields

$$\begin{aligned} \|\partial_{t}u_{m}(t)\|_{L^{2}}^{2} + \nu \int_{0}^{t} \|\partial_{t}\nabla u_{m}(s)\|_{L^{2}}^{2} ds + \beta \gamma \int_{0}^{t} \|\partial_{t}u_{m}(s)\|_{L^{2}}^{2} ds \\ &+ \beta(2\delta+1) \int_{0}^{t} \|u_{m}(s)^{\delta}\partial_{t}u_{m}(s)\|_{L^{2}}^{2} ds \\ \leqslant \left( \|\partial_{t}u_{m}(0)\|_{L^{2}}^{2} + \eta \|u_{0}\|_{H^{2}} \int_{0}^{t} |K(s)| ds + \frac{1}{\beta\gamma} \int_{0}^{t} \|\partial_{t}f(s)\|_{L^{2}}^{2} ds \right) \\ &\times \exp\left( \int_{0}^{t} \left[ C(\alpha,\nu,\delta) \|u_{m}(s)^{\delta}\nabla u_{m}(s)\|_{L^{2}}^{\frac{4(\delta-1)}{4-d)\delta}} \|\nabla u_{m}(s)\|_{L^{2}}^{\frac{4(\delta-1)}{2}} + \eta \|u_{0}\|_{H^{2}} |K(s)| + C(\beta,\gamma,\delta) \right] \right) \\ \leqslant C\left( \|u_{0}\|_{H^{2}}, \|f\|_{H^{1}(0,T;L^{2}(\Omega))}, \alpha, \nu, \beta, \gamma, \delta, T \right), \end{aligned}$$
(2.39)

for all  $t \in [0,T]$ , where the first term in exponential is bounded, provided  $1 \le \delta < \infty$ , for d = 2 and,  $1 \le \delta \le 2$ , for d = 3. Taking supremum over time  $0 \le t \le T$ , we have

$$\sup_{0 \le t \le T} \|\partial_t u_m(t)\|_{L^2}^2 + \nu \int_0^T \|\partial_t \nabla u_m(s)\|_{L^2}^2 \, \mathrm{d}s \le C \left( \|u_0\|_{\mathrm{H}^2}, \|f\|_{\mathrm{H}^1(0,T;\mathrm{L}^2(\Omega))}, \alpha, \beta, \gamma, \delta, \nu, T \right).$$

+

# GENERALIZED BURGERS-HUXLEY EQUATION WITH WEAKLY SINGULAR KERNEL 21 of 45

**Step 2.**  $\nabla(u^{\delta+1}) \in L^{\infty}(0,T;L^2(\Omega))$ . In (2.7), if we take  $w_k = -u_m^{2\delta}\Delta u_m$ , then we have for a.e.  $t \in [0,T]$ 

$$(\partial_{t}u_{m}(t), -u_{m}(t)^{2\delta}\Delta u_{m}(t)) + \nu(\Delta u_{m}(t), u_{m}(t)^{2\delta}\Delta u_{m}(t)) + \eta((K * \Delta u_{m})(t), u_{m}(t)^{2\delta}\Delta u_{m}(t)) = \alpha(B(u_{m}(t)), u_{m}(t)^{2\delta}\Delta u_{m}(t)) - \beta(c(u_{m}(t)), u_{m}(t)^{2\delta}\Delta u_{m}(t)) - (f(t), u_{m}(t)^{2\delta}\Delta u_{m}(t)).$$
(2.40)

Using integration by parts, one can estimate the time derivative term as

$$\begin{aligned} (\partial_t u_m, -u_m^{2\delta} \Delta u_m) &= -\frac{1}{2\delta + 1} \left( \partial_t u_m^{2\delta + 1}, \Delta u_m \right) = \frac{1}{2\delta + 1} \left( \nabla \left( \partial_t u_m^{2\delta + 1} \right), \nabla u_m \right) \\ &= \left( \partial_t \left( u_m^{2\delta} \nabla u_m \right), \nabla u_m \right) = \frac{d}{dt} \left( u_m^{2\delta} \nabla u_m, \nabla u_m \right) - \left( u_m^{2\delta} \nabla u_m, \partial_t \nabla u_m \right) \\ &= \frac{d}{dt} \| u^{\delta} \nabla u_m \|_{L^2}^2 - \left( u_m^{2\delta} \nabla u_m, \partial_t \nabla u_m \right). \end{aligned}$$

Using the above estimate, (2.40) can be rewritten as

$$\frac{d}{dt} \|u_m(t)^{\delta} \nabla u_m(t)\|_{L^2}^2 + \nu \|u_m(t)^{\delta} \Delta u\|_{L^2}^2 + \eta ((K * \Delta) u_m(t), u_m(t)^{2\delta} \Delta u_m(t)) 
= (u_m(t)^{2\delta} \nabla u_m(t), \partial_t \nabla u_m(t)) + \alpha (B(u_m(t)), u^{2\delta} \Delta u_m(t)) 
- \beta (c(u_m(t)), u_m(t)^{2\delta} \Delta u_m(t)) - (f(t), u_m(t)^{2\delta} \Delta u_m(t)),$$
(2.41)

for a.e.  $t \in [0,T]$ . For the term  $-\beta(c(u_m), u_m^{2\delta} \Delta u_m)$ , using integration by parts and Young's inequality, we get

$$\begin{split} -\beta(c(u_m), u_m^{2\delta}\Delta u_m) &= -\beta(u_m(1-u_m^{\delta})(u_m^{\delta}-\gamma), u_m^{2\delta}\Delta u_m) \\ &= -\beta(1+\gamma)(u_m^{\delta+1}, u_m^{2\delta}\Delta u_m) + \beta\gamma(u_m, u_m^{2\delta}\Delta u_m) + \beta(u^{2\delta+1}, u_m^{2\delta}\Delta u_m) \\ &= -\beta(1+\gamma)(u_m^{3\delta+1}, \Delta u_m) + \beta\gamma(u_m^{2\delta+1}, \Delta u_m) + \beta(u_m^{4\delta+1}, \Delta u_m) \\ &= \beta(1+\gamma)(3\delta+1)(u_m^{3\delta}\nabla u_m, \nabla u_m) - \beta\gamma(2\delta+1)(u_m^{2\delta}\nabla u_m, \nabla u_m) - \beta(4\delta+1)(u_m^{4\delta}\nabla u_m, \nabla u_m) \\ &\leqslant \frac{\beta}{4}(4\delta+1) \|u_m^{2\delta}\nabla u_m\|_{L^2}^2 + \frac{\beta(1+\gamma)^2(3\delta+1)^2}{(4\delta+1)} \|u_m^{\delta}\nabla u_m\|_{L^2}^2 \\ &- \beta\gamma(2\delta+1) \|u_m^{\delta}\nabla u_m\|_{L^2}^2 - \beta(4\delta+1) \|u^{2\delta}\nabla u_m\|_{L^2}^2. \end{split}$$

An application of Cauchy-Schwarz and Young's inequalities yields

$$\begin{aligned} \alpha(B(u_m), u_m^{2\delta} \Delta u_m) &= \sum_{i=1}^d \left( u_m^{\delta} \frac{\partial u_m}{\partial x_i}, u_m^{2\delta} \Delta u_m \right) \leqslant \|u_m^{2\delta} \nabla u_m\|_{L^2} \|u_m^{\delta} \Delta u_m\|_{L^2} \\ &\leqslant \frac{1}{v} \|u_m\|_{L^{\infty}}^{2\delta} \|u_m^{\delta} \nabla u_m\|_{L^2}^2 + \frac{v}{4} \|u_m^{\delta} \Delta u_m\|_{L^2}^2, \\ (f, u_m^{2\delta} \Delta u_m) &= (u_m^{\delta} f, u_m^{\delta} \Delta u_m) \leqslant \frac{v}{4} \|u_m^{\delta} \Delta u_m\|_{L^2}^2 + \frac{1}{v} \|u_m^{\delta} f\|_{L^2}^2 \leqslant \frac{v}{4} \|u_m^{\delta} \Delta u_m\|_{L^2}^2 + \frac{1}{v} \|u_m\|_{L^{\infty}}^{2\delta} \|f\|_{L^2}^2, \\ (u_m^{2\delta} \nabla u_m, \partial_t \nabla u_m) \leqslant \|u_m^{2\delta} \nabla u_m\|_{L^2}^2 \|\partial_t \nabla u_m\|_{L^2}^2 \leqslant \frac{\beta}{4} (4\delta+1) \|u_m^{2\delta} \nabla u_m\|_{L^2}^2 + \frac{1}{\beta(4\delta+1)} \|\partial_t \nabla u_m\|_{L^2}^2. \end{aligned}$$

Combining the above estimates and integrating from 0 to t, and using Lemma 2.2 and Remark 2.1, we have

$$\begin{aligned} \|u_{m}(t)^{\delta}\nabla u_{m}(t)\|_{L^{2}}^{2} + \frac{\nu}{2} \int_{0}^{t} \|u_{m}(s)^{\delta}\Delta u_{m}\|_{L^{2}}^{2} ds + \frac{\beta}{2}(4\delta+1) \int_{0}^{t} \|u_{m}(s)^{2\delta}\nabla u_{m}(s)\|_{L^{2}}^{2} ds \\ &+ \beta\gamma(2\delta+1) \int_{0}^{t} \|u_{m}(s)^{\delta}\nabla u_{m}(s)\|_{L^{2}}^{2} ds \\ \leqslant \|u_{m}(0)^{\delta}\nabla u_{m}(0)\|_{L^{2}}^{2} + \frac{1}{\beta(4\delta+1)} \int_{0}^{t} \|\partial_{t}\nabla u_{m}(s)\|_{L^{2}}^{2} ds + \frac{\beta(1+\gamma)^{2}(3\delta+1)^{2}}{(4\delta+1)} \int_{0}^{t} \|u_{m}(s)^{\delta}\nabla u_{m}(s)\|_{L^{2}}^{2} ds \\ &+ \frac{1}{\nu} \int_{0}^{t} \|u_{m}(s)\|_{L^{\infty}}^{2\delta} \|u_{m}(s)^{\delta}\nabla u_{m}(s)\|_{L^{2}}^{2\delta} ds + \frac{1}{\nu} \int_{0}^{t} \|u_{m}(s)\|_{L^{\infty}}^{2\delta} \|f(s)\|_{L^{2}}^{2} ds. \end{aligned}$$
(2.42)

As  $f \in H^1(0,T;L^2(\Omega))$ , we have  $f \in L^2(0,T;L^2(\Omega))$  and  $\partial_t f \in L^2(0,T;L^2(\Omega))$ , which implies  $f \in L^2(0,T;L^2(\Omega))$  $C([0,T],L^2(\Omega))$ , so we have

$$\int_0^t \|u_m(s)\|_{L^{\infty}}^{2\delta} \|f(s)\|_{L^2}^2 \, \mathrm{d} s \leq \sup_{0 \leq t \leq T} \|f(t)\|_{L^2}^2 \int_0^t \|u_m(s)\|_{L^{\infty}}^{2\delta} \, \mathrm{d} s.$$

Using the last estimate and an application of Gronwall's inequality in (2.42) yield

$$\begin{split} \sup_{0 \le t \le T} \|u_m(t)^{\delta} \nabla u_m(t)\|_{L^2}^2 &+ \frac{\nu}{2} \int_0^t \|u_m(s)^{\delta} \Delta u_m\|_{L^2}^2 \, ds + \frac{\beta}{2} (4\delta + 1) \int_0^t \|u_m(s)^{2\delta} \nabla u_m(s)\|_{L^2}^2 \, ds \\ &\le \left( \|u_m(0)^{\delta} \nabla u_m(0)\|_{L^2}^2 + \frac{1}{\beta(4\delta + 1)} \int_0^T \|\partial_t \nabla u_m(s)\|_{L^2}^2 \, ds + \sup_{0 \le t \le T} \|f(t)\|_{L^2}^2 \int_0^T \|u_m(s)\|_{L^{\infty}}^{2\delta} \, ds \right) \\ &\times e^{\left(\frac{\beta(1+\gamma)^2(3\delta+1)^2}{(4\delta+1)}\right)} \exp\left\{ \frac{1}{\nu} \int_0^T \|u_m(s)\|_{L^{\infty}}^{2\delta} \, ds \right\}. \end{split}$$

For the right hand side to be bounded, we need to show that  $\int_0^T \|u_m(t)\|_{L^{\infty}}^{2\delta} ds < \infty$ . As  $u \in L^{\infty}(0,T; H_0^1(\Omega)) \cap L^2(0,T; H^2(\Omega))$ , using Agmon's inequality we have for 3D(d=3)

$$\int_0^T \|u_m(s)\|_{L^{\infty}}^{2\delta} \, \mathrm{d} s \leqslant C \int_0^T \|u_m(s)\|_{\mathrm{H}^1}^{\delta} \|u_m(s)\|_{\mathrm{H}^2}^{\delta} \, \mathrm{d} s \leqslant C \sup_{0 \leqslant t \leqslant T} \|u_m(s)\|_{\mathrm{H}^1}^{\delta} \int_0^T \|u_m(s)\|_{\mathrm{H}^2}^{\delta} \, \mathrm{d} s < \infty,$$

for  $1 \leq \delta \leq 2$ . For the 2D (d = 2), for  $\varepsilon > 0$ , using Agmon's inequality  $||u||_{L^{\infty}} \leq C ||u||_{H^{1-\varepsilon}}^{\frac{1}{1+\varepsilon}} ||u||_{H^{2}}^{\frac{\varepsilon}{1+\varepsilon}}$ , so

$$\int_0^T \|u_m(s)\|_{\mathrm{L}^{\infty}}^{2\delta} \,\mathrm{d} s \leqslant C \int_0^T \|u_m(s)\|_{\mathrm{H}^{1-\varepsilon}}^{\frac{2\delta}{1+\varepsilon}} \|u_m(s)\|_{\mathrm{H}^2}^{\frac{2\varepsilon\delta}{1+\varepsilon}} \,\mathrm{d} s \leqslant C \sup_{0\leqslant t\leqslant T} \|u_m(s)\|_{\mathrm{H}^{1-\varepsilon}}^{\frac{2\delta}{1+\varepsilon}} \int_0^t \|u_m(s)\|_{\mathrm{H}^2}^{\frac{2\varepsilon\delta}{1+\varepsilon}} \,\mathrm{d} s < \infty,$$

provided  $\frac{2\varepsilon\delta}{1+\varepsilon} \leq 2 \implies \delta \leq \frac{1}{\varepsilon} + 1$ . Since  $\varepsilon > 0$  is arbitrary, the above estimate is finite for  $1 \leq \delta < \infty$ . Hence  $\nabla u_m^{\delta+1} \in L^{\infty}(0,T; L^2(\Omega))$ . **Step 3.**  $u \in L^{\infty}(0,T; H^2(\Omega))$ . Let  $\lambda_k$  denote the  $k^{\text{th}}$  eigenvalue of  $\Delta$  in  $H_0^1(\Omega)$ . Multiplying the identity (2.7) by  $-\lambda_k^2 d_k^m(t)$  and summing it from k = 1, 2, ..., m, we have

$$(\partial_t u_m(t), A^2 u_m(t)) - \mathbf{v}(A u_m(t), A^2 u_m(t)) - \eta(K * A u_m(t), A^2 u_m(t))$$

GENERALIZED BURGERS-HUXLEY EQUATION WITH WEAKLY SINGULAR KERNEL 23 of 45

$$= -\alpha(B(u_m(t)), A^2u_m(t)) + \beta(c(u_m(t)), A^2u_m(t)) + (f(t), A^2u_m(t)).$$
(2.43)

An application of integration by parts yields

$$\frac{1}{2} \frac{\mathrm{d}}{\mathrm{d}t} \|Au_m(t)\|_{\mathrm{L}^2}^2 + \mathbf{v} \|A^{\frac{3}{2}}u_m(t)\|_{\mathrm{L}^2}^2 + \eta \left(K * A^{\frac{3}{2}}u_m(t), A^{\frac{3}{2}}u_m(t)\right) 
= -\alpha \left(B(u_m(t)), A^2u_m(t)\right) + \beta \left(c(u_m(t)), A^2u_m(t)\right) + \left(f(t), A^2u_m(t)\right).$$
(2.44)

Note that  $D(A^{\alpha}) = \begin{cases} H^{2\alpha}(\Omega) & \text{if } 0 < \alpha < 1/4, \\ H_0^{2\alpha}(\Omega) & \text{if } 1/4 < \alpha < 1. \end{cases}$  Since  $u_m |_{\partial \Omega} = 0$ , using integration by parts, Cauchy-Schwarz and Young's inequalities, we have

$$\begin{aligned} \alpha |(B(u_m), A^2 u_m)| &\leq \alpha ||A^{\frac{1}{2}} B(u_m)||_{L^2} ||A^{\frac{3}{2}} u_m||_{L^2} &\leq \alpha ||A^{\frac{1}{2}} (u_m^{\delta} \nabla u_m)||_{L^2} ||A^{\frac{3}{2}} u_m||_{L^2} \\ &\leq \frac{\nu}{8} ||A^{\frac{3}{2}} u_m||_{L^2}^2 + \frac{2\alpha^2}{\nu(\delta+1)} ||A^{\frac{1}{2}} \nabla u_m^{\delta+1}||_{L^2}^2 \\ &\leq \frac{\nu}{8} ||A^{\frac{3}{2}} u_m||_{L^2}^2 + \frac{C\alpha^2}{\nu} \left( ||u_m^{\delta} A u_m||_{L^2}^2 + \delta ||u_m^{\delta-1} (\nabla u_m)^2||_{L^2}^2 \right). \end{aligned}$$

Now we estimate  $||u_m^{\delta-1}(\nabla u_m)^2||_{L^2}^2$  using Sobolev embedding (for  $d = 2, 1 \le \delta < \infty$  and for  $d = 3, 1 \le \delta \le 2$ ), Hölder's and Young's inequalities as

$$\begin{split} \|u_{m}^{\delta-1}(\nabla u_{m})^{2}\|_{\mathrm{L}^{2}}^{2} &= \int_{\Omega} |u_{m}|^{2(\delta-1)} |\nabla u_{m}|^{4} \mathrm{d}x = \int_{\Omega} |u_{m}|^{2(\delta-1)} |\nabla u_{m}|^{\frac{\delta-1}{\delta}} |\nabla u_{m}|^{\frac{3\delta+1}{\delta}} \mathrm{d}x \\ &\leq \left(\int_{\Omega} |u_{m}|^{4\delta} |\nabla u_{m}|^{2} \mathrm{d}x\right)^{\frac{\delta-1}{2\delta}} \left(\int_{\Omega} |\nabla u_{m}|^{\frac{2(3\delta+1)}{\delta+1}} \mathrm{d}x\right)^{\frac{\delta+1}{2\delta}} \\ &= \|u_{m}^{2\delta} \nabla u_{m}\|_{\mathrm{L}^{2}}^{\frac{\delta-1}{\delta}} \|\nabla u_{m}\|_{\mathrm{L}^{2}^{\frac{3\delta+1}{\delta}}}^{\frac{3\delta+1}{\delta}} \\ &\leq C \|u_{m}^{2\delta} \nabla u_{m}\|_{\mathrm{L}^{2}}^{\frac{\delta-1}{\delta}} \|Au_{m}\|_{\mathrm{L}^{2}}^{\frac{3\delta+1}{\delta}} = C \||u_{m}|^{2\delta} \nabla u_{m}\|_{\mathrm{L}^{2}}^{\frac{\delta-1}{\delta}} \|Au_{m}\|_{\mathrm{L}^{2}}^{\frac{2(\delta+1)}{\delta}} \\ &\leq \frac{1}{2} \|u_{m}^{2\delta} \nabla u_{m}\|_{\mathrm{L}^{2}}^{2} \|Au_{m}\|_{\mathrm{L}^{2}}^{2} + C \left(\frac{2(\delta-1)}{2\delta}\right)^{\frac{\delta-1}{\delta+1}} \left(\frac{\delta+1}{2\delta}\right) \|Au_{m}\|_{\mathrm{L}^{2}}^{4}. \end{split}$$

Combining, we have

$$\alpha|(B(u_m),A^2u_m)| \leq \frac{\nu}{8} ||A^{\frac{3}{2}}u_m||_{L^2}^2 + \frac{C\alpha^2}{\nu} ||u_m^{\delta}Au_m||_{L^2}^2 + \frac{\delta}{2} ||u_m^{2\delta}\nabla u_m||_{L^2}^2 ||Au_m||_{L^2}^2 + C(\alpha,\nu,\delta) ||Au_m||_{L^2}^4$$

Again using Cauchy-Schwarz and Young's inequalities, we get

$$\begin{aligned} |\beta(c(u_m), A^2 u_m)| &= |\beta((1+\gamma)u_m^{\delta+1} - \gamma u_m - u_m^{2\delta+1}, A^2 u_m)| \\ &= \beta(1+\gamma)(A^{\frac{1}{2}}u_m^{\delta+1}, A^{\frac{3}{2}}u_m) + \beta\gamma(Au_m, Au_m) + \beta(A^{\frac{1}{2}}u_m^{2\delta+1}, A^{\frac{3}{2}}u_m) \\ &\leqslant \beta(1+\gamma)(\delta+1) \|u_m^{\delta}A^{\frac{1}{2}}u_m\|_{L^2} \|A^{\frac{3}{2}}u_m\|_{L^2} + \beta\gamma\|Au_m\|_{L^2}^2 + \beta(2\delta+1) \|u_m^{2\delta}A^{\frac{1}{2}}u_m\|_{L^2} \|A^{\frac{3}{2}}u_m\|_{L^2} \\ &\leqslant \frac{\beta^2(1+\gamma)^2(\delta+1)^2}{\nu} \|u_m^{\delta}A^{\frac{1}{2}}u_m\|_{L^2}^2 + \beta\gamma\|Au_m\|_{L^2}^2 + \frac{\beta^2(2\delta+1)^2}{\nu} \|u_m^{2\delta}A^{\frac{1}{2}}u_m\|_{L^2}^2 + \frac{\gamma}{4} \|A^{\frac{3}{2}}u_m\|_{L^2}^2. \end{aligned}$$

Finally, for the forcing term using Cauchy-Schwarz and Young's inequalities, we obtain

$$|(f,A^{2}u_{m})| \leq |(A^{\frac{1}{2}}f,A^{\frac{3}{2}}u_{m})| \leq \frac{1}{\nu} ||A^{\frac{1}{2}}f||_{L^{2}}^{2} + \frac{\nu}{8} ||A^{\frac{3}{2}}u_{m}||_{L^{2}}^{2}.$$

Combing the above estimates we have

$$\begin{split} &\frac{1}{2} \frac{\mathrm{d}}{\mathrm{d}t} \|Au_m(t)\|_{\mathrm{L}^2}^2 + \frac{\nu}{2} \|A^{\frac{3}{2}} u_m(t)\|_{\mathrm{L}^2}^2 + \eta \left(K * A^{\frac{3}{2}} u_m(t), A^{\frac{3}{2}} u_m(t)\right) \\ &\leqslant \frac{\beta^2 (1+\gamma)^2 (\delta+1)^2}{\nu} \|u_m(t)^{\delta} A^{\frac{1}{2}} u_m(t)\|_{\mathrm{L}^2}^2 + \frac{\beta^2 (2\delta+1)^2}{\nu} \|u_m(t)^{2\delta} A^{\frac{1}{2}} u_m(t)\|_{\mathrm{L}^2}^2 + \frac{1}{\nu} \|A^{\frac{1}{2}} f(t)\|_{\mathrm{L}^2}^2 \\ &+ \frac{C\alpha^2}{\nu} \|u_m(t)^{\delta} A u_m(t)\|_{\mathrm{L}^2}^2 + \left(\frac{\delta}{2} \|u_m(t)^{2\delta} \nabla u_m(t)\|_{\mathrm{L}^2}^2 + \beta\gamma + C(\alpha,\nu,\delta) \|A u_m(t)\|_{\mathrm{L}^2}^2\right) \|A u_m(t)\|_{\mathrm{L}^2}^2. \end{split}$$

Integrating from 0 to t, using positivity of the kernel K and Gronwall's inequality, and taking supremum over time, we have

$$\begin{split} \sup_{0 \leq t \leq T} \|Au_m(t)\|_{L^2}^2 + \nu \int_0^t \|A^{\frac{3}{2}}u_m(s)\|_{L^2}^2 ds \\ &\leq \left(\frac{2\beta^2(1+\gamma)^2(\delta+1)^2}{\nu} \int_0^T \|u_m(s)^{\delta}A^{\frac{1}{2}}u_m(s)\|_{L^2}^2 ds + \frac{2\beta^2(2\delta+1)^2}{\nu} \int_0^T \|u_m(s)^{2\delta}A^{\frac{1}{2}}u_m(s)\|_{L^2}^2 ds \\ &+ \frac{2}{\nu} \int_0^T \|A^{\frac{1}{2}}f(s)\|_{L^2}^2 ds + \|u_0\|_{H^2}\right) \\ &\times \exp\left(\delta \int_0^T \|u_m(s)^{2\delta}\nabla u_m(s)\|_{L^2}^2 ds + 2\beta\gamma T + C(\alpha,\nu,\delta) \int_0^T \|Au_m(s)\|_{L^2}^2 ds\right). \end{split}$$

Using the estimate (2.29) and (2.42) the right hand side is bounded. Hence we have  $u_m \in L^{\infty}(0,T; H^2(\Omega))$ ,  $\partial_t u_m \in L^{\infty}(0,T; L^2(\Omega)) \cap L^2(0,T; H^1_0(\Omega))$ . Passing limit along a subsequence  $m_j \to \infty$ , we obtain the required bound for u.

REMARK 2.5 In Theorem 2.5 (*ii*), we need  $f \in L^2(0,T; H^1(\Omega))$  for the memory term. However, if we take  $\eta = 0$ , that is, GBHE without memory using elliptic regularity we can prove that for  $u_0 \in H^2(\Omega) \cap H^1_0(\Omega)$  and  $f \in H^1(0,T; L^2(\Omega))$ , we have  $u \in L^{\infty}(0,T; H^2(\Omega))$ .

# 3. Finite Element Method

#### 3.1 Semidiscrete Galerkin approximation and error estimates

This section is devoted to the error estimates for the semidiscrete Galerkin approximation. We partition the domain  $\Omega$  into shape-regular meshes (triangular or rectangles) denoted by  $\mathcal{T}_h$  and consider a finite dimensional space  $V_h$ , defined as

$$V_h = \left\{ v_h : v_h \in C^0(\bar{\Omega}) \cap \mathrm{H}^1_0(\Omega), v_h |_T \in \mathbb{P}_1(T) \text{ for all } T \in \mathscr{T}_h \right\},$$
(3.1)

where  $\mathbb{P}_1$  is the space of polynomials which have degree at most 1. Note that,  $V_h$  is a finite dimensional subspace of  $\mathrm{H}_0^1(\Omega)$  and 0 < h < 1 is the associated small mesh parameter. The aim is to prove the estimate of the type (see Thomée, 2007)

$$\inf_{\chi \in V_h} \{ \|u - \chi\|_{L^2} + h \|\nabla(u - \chi)\|_{L^2} \} \leq Ch^2 \|u\|_{H^2},$$

for all  $u \in H^2(\Omega) \cap H^1_0(\Omega)$ . The continuous time Galerkin approximation of the problem (2.6) is defined in the following way: Find  $u_h(\cdot, t) \in V_h$ , such that for  $t \in (0, T)$ ,

$$\langle \partial_t u_h(t), \chi \rangle + \mathbf{v} (\nabla u_h(t), \nabla \chi) + \alpha b(u_h(t), u_h(t), \chi) + \eta ((K * \nabla u_h)(t), \nabla \chi) = \beta (c(u_h(t)), \chi) + \langle f(t), \chi \rangle,$$

$$(u_h(0), \chi) = (u_0^h, \chi), \text{ for } \chi \in V_h,$$

$$(3.2)$$

where  $u_0^h$  approximates  $u_0$  in  $V_h$ . Central to the analysis of finite element methods, we define an elliptic projection associated to our model onto  $V_h$ , the Ritz projection (Thomée (2007)) of *u* defined by

$$(\nabla R_h u, \nabla \chi) = (\nabla u, \nabla \chi), \text{ for all } \chi \in V_h.$$

By setting  $\chi = R_h u$  in the above equality, we obtain that the Ritz projection is stable, that is,  $\|\nabla R_h u\|_{L^2} \leq \|\nabla u\|_{L^2}$ , for all  $u \in H_0^1(\Omega)$ . If we take  $\Lambda = R_h u - u$ , then, we have the following estimate for  $\Lambda$  (Thomée (2007)),

$$\|\Lambda(t)\|_{L^{2}} + h\|\nabla\Lambda(t)\|_{L^{2}} \leqslant Ch^{s}\|u(t)\|_{H^{s}}, \text{ for } s = 1, 2.$$
(3.3)

THEOREM 3.1 (Existence of a discrete solution) The discrete equation (3.2) admits at least one solution  $u_h \in V_h$ .

*Proof.* It follows as a direct consequence of Theorem 2.2.  $\Box$ For all  $t \in [0, T]$ , the error estimates in this semi-discretization are given in the following theorem,

THEOREM 3.2 Let  $V_h$  be the finite dimensional subspace of  $H_0^1(\Omega)$  with parameter *h* defined in (3.1). Assume that  $u_0 \in L^{d\delta}(\Omega) \cap H_0^1(\Omega)$  and  $f \in L^2(0,T;L^2(\Omega))$ , and  $u(\cdot)$ , the solution of (2.1) has the following regularity:

$$u \in \mathcal{L}^{\infty}(0,T;\mathcal{L}^{2(\delta+1)}(\Omega)) \cap \mathcal{L}^{2}(0,T;\mathcal{H}^{1}_{0}(\Omega)) \cap \mathcal{L}^{(d+2)\delta}(0,T;\mathcal{L}^{(d+2)\delta}(\Omega)), \ \partial_{t}u \in \mathcal{L}^{2}(0,T;\mathcal{H}^{1}_{0}(\Omega)).$$

Then the error in the semi-discretization satisfies the following estimate:

$$\|u_{h} - u\|_{\mathrm{L}^{\infty}(0,T;\mathrm{L}^{2}(\Omega))}^{2} + \|u_{h} - u\|_{\mathrm{L}^{2}(0,T;\mathrm{H}_{0}^{1}(\Omega))}^{2}$$
  
$$\leq C \bigg\{ \|u_{0}^{h} - u_{0}\|_{\mathrm{L}^{2}}^{2} + h^{2} \int_{0}^{T} \|\partial_{t}u(t)\|_{\mathrm{H}_{0}^{1}}^{2} \,\mathrm{d}t + h^{2} \int_{0}^{T} \|u(t)\|_{\mathrm{H}_{0}^{1}}^{2} \,\mathrm{d}t + h^{2} \int_{0}^{T} \|u(t)\|_{\mathrm{H}^{2}}^{2} \,\mathrm{d}t \bigg\},$$
(3.4)

where the constant *C* is independent of *h* and depends only on  $\|u_0\|_{H^1_0}$ ,  $\nu$ ,  $\alpha$ ,  $\beta$ ,  $\gamma$ ,  $\eta$ ,  $\delta$ , *T*,  $\|f\|_{L^2(0,T;L^2(\Omega))}$ .

Note: If  $u_0 \in L^{d\delta}(\Omega) \cap H^1_0(\Omega)$  and  $f \in L^2(0,T;L^2(\Omega))$ , then using Theorem 2.4 we have  $u \in L^{\infty}(0,T;L^{2(\delta+1)}(\Omega)) \cap L^2(0,T;H^1_0(\Omega)) \cap L^{(d+2)\delta}(0,T;L^{(d+2)\delta}(\Omega))$ . *Proof.* Using (2.1) and (3.2), we know that  $(u_h - u)(\cdot)$  satisfies

$$\langle \partial_t (u_h(t) - u(t)), \boldsymbol{\chi} \rangle + v(\nabla(u_h(t) - u(t)), \nabla \boldsymbol{\chi}) + \eta((K * \nabla(u_h - u))(t), \nabla \boldsymbol{\chi}) = -\alpha[b(u_h(t), u_h(t), \boldsymbol{\chi}) - b(u(t), u(t), \boldsymbol{\chi})] + \beta[c(u_h(t), \boldsymbol{\chi}) - c(u(t), \boldsymbol{\chi})],$$
(3.5)

for all  $\chi \in V_h$  and a.e.  $t \in [0, T]$ . Let us choose  $\chi = u_h - W \in V_h$  in (3.5) to obtain for a.e.  $t \in [0, T]$ 

$$\langle \partial_t (u_h(t) - u(t)), u_h(t) - W(t) \rangle + v(\nabla (u_h(t) - u(t)), \nabla (u_h(t) - W(t)))$$

$$+ \eta \langle (K * \nabla(u_h - u))(t), \nabla(u_h(t) - W(t)) \rangle$$
  
=  $-\alpha [b(u_h(t), u_h(t), u_h(t) - W(t)) - b(u(t), u(t), u_h(t) - W(t))]$   
+  $\beta [c(u_h(t), u_h(t) - W(t)) - c(u(t), u_h(t) - W(t))].$  (3.6)

Next, we write  $u_h - W$  as  $u_h - u + u - W$  in (3.6) to find

$$\frac{1}{2} \frac{d}{dt} \|u_{h}(t) - u(t)\|_{L^{2}}^{2} + \nu \|\nabla(u_{h}(t) - u(t))\|_{L^{2}}^{2} + \eta \left((K * \nabla(u_{h} - u))(t), \nabla(u_{h}(t) - u(t))\right)\right) 
= -\langle \partial_{t}(u_{h}(t) - u(t)), u(t) - W(t) \rangle - (\nabla(u_{h}(t) - u(t)), \nabla(u(t) - W(t))) 
- \eta \langle (K * \nabla(u_{h} - u))(t), \nabla(u(t) - W(t)) \rangle 
- \alpha \left(u_{h}(t)^{\delta} \sum_{i=1}^{d} \frac{\partial u_{h}(t)}{\partial x_{i}} - u(t)^{\delta} \sum_{i=1}^{d} \frac{\partial u(t)}{\partial x_{i}}, u_{h}(t) - u(t)\right) 
- \alpha \left(u_{h}(t)^{\delta} \sum_{i=1}^{d} \frac{\partial u_{h}(t)}{\partial x_{i}} - u(t)^{\delta} \sum_{i=1}^{d} \frac{\partial u(t)}{\partial x_{i}}, u(t) - W(t)\right) 
+ \beta \left(u_{h}(t)(1 - u_{h}(t)^{\delta})(u_{h}(t)^{\delta} - \gamma) - u(t)(1 - u(t)^{\delta})(u(t)^{\delta} - \gamma), u_{h}(t) - u(t)\right) 
+ \beta \left(u_{h}(t)(1 - u_{h}(t)^{\delta})(u_{h}(t)^{\delta} - \gamma) - u(t)(1 - u(t)^{\delta})(u(t)^{\delta} - \gamma), u(t) - W(t)\right), \qquad (3.7)$$

for a.e.  $t \in [0, T]$ . Using calculations similar to (2.25) yields

$$\begin{aligned} \frac{\mathrm{d}}{\mathrm{d}t} \|u_{h}(t) - u(t)\|_{\mathrm{L}^{2}}^{2} + \nu \|\nabla(u_{h}(t) - u(t))\|_{\mathrm{L}^{2}}^{2} + 2\eta((K * \nabla(u_{h} - u))(t), \nabla(u_{h}(t) - u(t)))) \\ &+ \frac{\beta}{2} \|u_{h}(t)^{\delta}(u_{h}(t) - u(t))\|_{\mathrm{L}^{2}}^{2} + \frac{\beta}{2} \|u(t)^{\delta}(u_{h}(t) - u(t))\|_{\mathrm{L}^{2}}^{2} + \beta\gamma\|u_{h}(t) - u(t)\|_{\mathrm{L}^{2}}^{2} \\ &\leq \left[ C(\beta, \gamma, \delta) + C(\alpha, \nu) \left( \|u_{h}(t)\|_{\mathrm{L}^{4\delta}}^{\frac{8\delta}{4-d}} + \|u(t)\|_{\mathrm{L}^{4\delta}}^{\frac{8\delta}{4-d}} \right) \right] \|u_{h}(t) - u(t)\|_{\mathrm{L}^{2}}^{2} \\ &- 2\partial_{t}(u_{h}(t) - u(t), u(t) - W(t)) + 2(u_{h}(t) - u(t), \partial_{t}(u(t) - W(t)))) \\ &- 2(\nabla(u_{h}(t) - u(t)), \nabla(u(t) - W(t))) - 2\eta\langle(K * \nabla(u_{h} - u))(t), \nabla(u(t) - W(t))\rangle) \\ &- 2\alpha \left( u_{h}(t)^{\delta} \sum_{i=1}^{d} \frac{\partial u_{h}(t)}{\partial x_{i}} - u(t)^{\delta} \sum_{i=1}^{d} \frac{\partial u(t)}{\partial x_{i}}, u(t) - W(t) \right) \\ &+ 2\beta \left( u_{h}(t)(1 - u_{h}(t)^{\delta})(u_{h}(t)^{\delta} - \gamma) - u(t)(1 - u(t)^{\delta})(u(t)^{\delta} - \gamma), u(t) - W(t) \right) \\ &= \left( C(\beta, \gamma, \delta) + C(\alpha, \nu) \left( \|u_{h}(t)\|_{\mathrm{L}^{4\delta}}^{\frac{8\delta}{4-d}} + \|u(t)\|_{\mathrm{L}^{4\delta}}^{\frac{8\delta}{4-d}} \right) \right) \|u_{h}(t) - u(t)\|_{\mathrm{L}^{2}}^{2} \\ &- 2\partial_{t}(u_{h}(t) - u(t), u(t) - W(t)) - 2\eta\langle(K * \nabla(u_{h} - u))(t), \nabla(u(t) - W(t))\rangle + \sum_{i=1}^{4} J_{i}, \end{aligned}$$
(3.8)

where the positive constants  $C(\beta, \gamma, \delta) = \beta 2^{2\delta} (1+\gamma)^2 (\delta+1)^2$  and  $C(\alpha, \nu) = \left(\frac{4+d}{4\nu}\right)^{\frac{4+d}{4-d}} \left(\frac{4-d}{8}\right) (2^{\delta-1}\alpha)^{\frac{4-d}{8}}$ . We estimate  $J_1$  and  $J_2$  using the Cauchy-Schwarz and Young's inequalities as

$$|J_1| \leq 2||u_h - u||_{L^2} ||\partial_t (u - W)||_{L^2} \leq ||u_h - u||_{L^2}^2 + ||\partial_t (u - W)||_{L^2}^2,$$
(3.9)

GENERALIZED BURGERS-HUXLEY EQUATION WITH WEAKLY SINGULAR KERNEL 27 of 45

$$|J_2| \leq 2 \|\nabla(u_h - u)\|_{L^2} \|\nabla(u - W)\|_{L^2} \leq \frac{\nu}{4} \|\nabla(u_h - u)\|_{L^2}^2 + \frac{4}{\nu} \|\nabla(u - W)\|_{L^2}^2.$$
(3.10)

Using an integration by parts, Taylor's formula, Hölder's and Young's inequalities, we rewrite  $J_3$  as

$$J_{3} = -\frac{2\alpha}{\delta+1} \sum_{i=1}^{d} \left( \frac{\partial}{\partial x_{i}} (u_{h}^{\delta+1} - u^{\delta+1}), u - W \right) = \frac{2\alpha}{\delta+1} \sum_{i=1}^{d} \left( \left( u_{h}^{\delta+1} - u^{\delta+1} \right), \frac{\partial}{\partial x_{i}} (u - W) \right)$$
  
$$= 2\alpha \sum_{i=1}^{d} \left( (\theta u_{h} + (1 - \theta) u)^{\delta} (u_{h} - u), \frac{\partial}{\partial x_{i}} (u - W) \right)$$
  
$$\leq 2\alpha \sum_{i=1}^{d} \left\| (\theta u_{h} + (1 - \theta) u)^{\delta} (u_{h} - u) \right\|_{L^{2}} \left\| \frac{\partial}{\partial x_{i}} (u - W) \right\|_{L^{2}}$$
  
$$\leq 2^{\delta} \alpha \left( \left\| u_{h}^{\delta} (u_{h} - u) \right\|_{L^{2}} + \left\| u^{\delta} (u_{h} - u) \right\|_{L^{2}} \right) \left\| \nabla (u - W) \right\|_{L^{2}}$$
  
$$\leq \frac{\beta}{8} \left\| u_{h}^{\delta} (u_{h} - u) \right\|_{L^{2}}^{2} + \frac{\beta}{8} \left\| u^{\delta} (u_{h} - u) \right\|_{L^{2}}^{2} + \frac{2^{2(\delta+1)}\alpha^{2}}{\beta} \left\| \nabla (u - W) \right\|_{L^{2}}^{2}.$$
(3.11)

Let us rewrite  $J_4$  as

$$J_{4} = 2\beta(1+\gamma)(u_{h}^{\delta+1} - u^{\delta+1}, u - W) - 2\beta\gamma(u_{h} - u, u - W)$$
$$-2\beta(u_{h}^{2\delta+1} - u^{2\delta+1}, u - W) := \sum_{i=5}^{7} J_{i}.$$
(3.12)

We estimate  $J_5$  using Taylor's formula, Hölder's and Young's inequalities as

$$J_{5} = 2\beta(1+\gamma)(\delta+1)((\theta u_{h}+(1-\theta)u)^{\delta}(u_{h}-u),u-W)$$

$$\leq 2^{\delta}\beta(1+\gamma)(\delta+1)\left(\|u_{h}^{\delta}(u_{h}-u)\|_{L^{2}}+\|u^{\delta}(u_{h}-u)\|_{L^{2}}\right)\|u-W\|_{L^{2}}$$

$$\leq \frac{\beta}{8}\|u_{h}^{\delta}(u_{h}-u)\|_{L^{2}}^{2}+\frac{\beta}{8}\|u^{\delta}(u_{h}-u)\|_{L^{2}}^{2}+2^{2(\delta+1)}\beta(1+\gamma)^{2}(\delta+1)^{2}\|u-W\|_{L^{2}}^{2}.$$
(3.13)

Using the Cauchy-Schwarz and Young's inequalities, we estimate  $J_6$  as

$$J_{6} \leq 2\beta\gamma \|u_{h} - u\|_{L^{2}} \|u - W\|_{L^{2}} \leq \beta\gamma \|u_{h} - u\|_{L^{2}}^{2} + \beta\gamma \|u - W\|_{L^{2}}^{2}.$$
(3.14)

Making use of Taylor's formula, Hölder's, Young's and Gagliardo-Nirenberg inequalities, we estimate  $J_7$  as

$$\begin{split} J_{7} &= -2(2\delta+1)\beta\left((\theta u_{h} + (1-\theta)u)^{2\delta}(u_{h}-u), u-W\right) \\ &\leq 2^{2\delta}(2\delta+1)\beta\left((|u_{h}|^{2\delta} + |u|^{2\delta})(u_{h}-u), u-W\right) \\ &\leq 2^{2\delta}(2\delta+1)\beta\left(\|u_{h}\|_{L^{4\delta}}^{2\delta} + \|u\|_{L^{4\delta}}^{2\delta}\right)\|u_{h}-u\|_{L^{\frac{2d}{d-1}}}\|u-W\|_{L^{2d}} \\ &\leq 2^{2\delta}(2\delta+1)\beta\left(\|u_{h}\|_{L^{4\delta}}^{2\delta} + \|u\|_{L^{4\delta}}^{2\delta}\right)\|u_{h}-u\|_{L^{2}}^{\frac{1}{2}}\|\nabla(u_{h}-u)\|_{L^{2}}^{\frac{1}{2}}\|u-W\|_{L^{2d}} \\ &\leq 2^{2\delta-1}(2\delta+1)\beta\left(\|\nabla(u-W)\|_{L^{2}}^{2} + 2^{2\delta-1}(2\delta+1)\beta\left(\|u_{h}\|_{L^{4\delta}}^{4\delta} + \|u\|_{L^{4\delta}}^{4\delta}\right)\|u_{h}-u\|_{L^{2}}\|\nabla(u_{h}-u)\|_{L^{2}} \end{split}$$

$$\leq 2^{2\delta-1}(2\delta+1)\beta \|\nabla(u-W)\|_{L^{2}}^{2} + \frac{\nu}{4} \|\nabla(u_{h}-u)\|_{L^{2}}^{2} + \frac{2^{4\delta-2}(2\delta+1)^{2}\beta^{2}}{\nu} \left(\|u_{h}\|_{L^{4\delta}}^{8\delta} + \|u\|_{L^{4\delta}}^{8\delta}\right) \|u_{h}-u\|_{L^{2}}^{2},$$
(3.15)

where we used the fact that  $H^1_0(\boldsymbol{\varOmega}) \subset L^6(\boldsymbol{\varOmega}).$  Note also that

$$\|u_{h} - u\|_{L^{2(\delta+1)}}^{2(\delta+1)} = \int_{\Omega} |u_{h}(x) - u(x)|^{2\delta} |u_{h}(x) - u(x)|^{2} dx$$
  
$$\leq 2^{2\delta-1} \int_{\Omega} \left( |u_{h}(x)|^{2\delta} + |u(x)|^{2\delta} \right) |u_{h}(x) - u(x)|^{2} dx$$
  
$$= 2^{2\delta-1} \|u_{h}^{\delta}(u_{h} - u)\|_{L^{2}}^{2} + 2^{2\delta-1} \|u^{\delta}(u_{h} - u)\|_{L^{2}}^{2}.$$
(3.16)

Combining (3.9)-(3.16) and then substituting it in (3.8), we deduce

$$\begin{split} &\frac{\mathrm{d}}{\mathrm{d}t} \|u_{h}(t) - u(t)\|_{\mathrm{L}^{2}}^{2} + \frac{\nu}{2} \|\nabla(u_{h}(t) - u(t))\|_{\mathrm{L}^{2}}^{2} + \eta \langle (K * \nabla(u_{h} - u))(t), \nabla(u_{h}(t) - u(t)) \rangle \\ &+ \frac{\beta}{2^{2\delta+1}} \|u_{h}(t) - u(t))\|_{\mathrm{L}^{2\delta+2}}^{2\delta+2} \\ &\leqslant -2\partial_{t}(u_{h}(t) - u(t), u(t) - W(t)) + \|\partial_{t}(u(t) - W(t))\|_{\mathrm{L}^{2}}^{2} \\ &- \eta \langle (K * \nabla(u_{h} - u))(t), \nabla(u(t) - W(t)) \rangle + \left(2^{2\delta-1}(2\delta+1)\beta + \frac{4}{\nu} + \frac{2^{2(\delta+1)}\alpha^{2}}{\beta}\right) \|\nabla(u(t) - W(t))\|_{\mathrm{L}^{2}}^{2} \\ &+ \left(2^{2(\delta+1)}\beta(1+\gamma)^{2}(\delta+1)^{2} + \frac{\beta\gamma}{2}\right) \|u(t) - W(t)\|_{\mathrm{L}^{2}}^{2} + \left(1+2\beta\gamma + C(\beta,\gamma,\delta)\right) \|u_{h}(t) - u(t)\|_{\mathrm{L}^{2}}^{2} \\ &+ \left(\frac{2^{4\delta-2}(2\delta+1)^{2}\beta^{2}}{\nu} \left(\|u_{h}(t)\|_{\mathrm{L}^{4\delta}}^{8\delta} + \|u(t)\|_{\mathrm{L}^{4\delta}}^{8\delta}\right) + C(\alpha,\nu) \left(\|u_{h}(t)\|_{\mathrm{L}^{4\delta}}^{\frac{8\delta}{4-d}} + \|u(t)\|_{\mathrm{L}^{4\delta}}^{\frac{8\delta}{4-d}}\right) \left\|u_{h}(t) - u(t)\|_{\mathrm{L}^{2}}^{2} \end{split}$$

Integrating the above inequality from 0 to t and using positivity of kernel  $K(\cdot)$ , we have

$$\begin{split} \|u_{h}(t) - u(t)\|_{L^{2}}^{2} + \frac{v}{2} \int_{0}^{t} \|\nabla(u_{h}(s) - u(s))\|_{L^{2}}^{2} ds + \frac{\beta}{2^{2\delta+1}} \int_{0}^{t} \|u_{h}(s) - u(s))\|_{L^{2\delta+2}}^{2\delta+2} ds \\ &\leq \|u_{h}^{0} - u_{0}\|_{L^{2}}^{2} - 2(u_{h}(t) - u(t), u(t) - W(t)) + 2(u_{h}^{0} - u_{0}, u_{0} - W(0)) \\ &+ \int_{0}^{t} \|\partial_{t}(u(s) - W(s))\|_{L^{2}}^{2} ds + \left(2^{2\delta-1}(2\delta+1)\beta + \frac{4}{v} + \frac{2^{2(\delta+1)}\alpha^{2}}{\beta} + \frac{\eta}{4C_{K}v}\right) \int_{0}^{t} \|\nabla(u(s) - W(s))\|_{L^{2}}^{2} ds \\ &+ \left(2^{2(\delta+1)}\beta(1+\gamma)^{2}(\delta+1)^{2} + \frac{\beta\gamma}{2}\right) \int_{0}^{t} \|u(s) - W(s)\|_{L^{2}}^{2} ds + \left(1+2\beta\gamma+C(\beta,\alpha,\delta)\right) \int_{0}^{t} \|u_{h}(s) - u(s)\|_{L^{2}}^{2} ds \\ &+ \int_{0}^{t} \left(\frac{2^{4\delta-2}(2\delta+1)^{2}\beta^{2}}{v} \left(\|u_{h}(s)\|_{L^{4\delta}}^{8\delta} + \|u(s)\|_{L^{4\delta}}^{8\delta}\right) + C(\alpha,v) \left(\|u_{h}(s)\|_{L^{4\delta}}^{\frac{8\delta}{4-d}} + \|u(s)\|_{L^{4\delta}}^{\frac{8\delta}{4-d}}\right) \right) \|u_{h}(s) - u(s)\|_{L^{2}}^{2} ds \\ &+ \int_{0}^{t} \left(\frac{2^{4\delta-2}(2\delta+1)^{2}\beta^{2}}{v} \left(\|u_{h}(s)\|_{L^{4\delta}}^{8\delta} + \|u(s)\|_{L^{4\delta}}^{8\delta}\right) + C(\alpha,v) \left(\|u_{h}(s)\|_{L^{4\delta}}^{\frac{8\delta}{4-d}} + \|u(s)\|_{L^{4\delta}}^{\frac{8\delta}{4-d}}\right) \right) \|u_{h}(s) - u(s)\|_{L^{2}}^{2} ds \\ &+ \int_{0}^{t} \left(\frac{2^{4\delta-2}(2\delta+1)^{2}\beta^{2}}{v} \left(\|u_{h}(s)\|_{L^{4\delta}}^{8\delta} + \|u(s)\|_{L^{4\delta}}^{8\delta}\right) + C(\alpha,v) \left(\|u_{h}(s)\|_{L^{4\delta}}^{\frac{8\delta}{4-d}} + \|u(s)\|_{L^{4\delta}}^{\frac{8\delta}{4-d}}\right) \right) \|u_{h}(s) - u(s)\|_{L^{2}}^{2} ds \\ &+ \int_{0}^{t} \left(\frac{2^{4\delta-2}(2\delta+1)^{2}\beta^{2}}{v} \left(\|u_{h}(s)\|_{L^{4\delta}}^{8\delta} + \|u(s)\|_{L^{4\delta}}^{8\delta}\right) + C(\alpha,v) \left(\|u_{h}(s)\|_{L^{4\delta}}^{\frac{8\delta}{4-d}} + \|u(s)\|_{L^{4\delta}}^{\frac{8\delta}{4-d}}\right) \right) \|u_{h}(s) - u(s)\|_{L^{2}}^{2} ds \\ &+ \int_{0}^{t} \left(\frac{2^{4\delta-2}(2\delta+1)^{2}\beta^{2}}{v} \left(\|u_{h}(s)\|_{L^{4\delta}}^{8\delta} + \|u(s)\|_{L^{4\delta}}^{8\delta}\right) + C(\alpha,v) \left(\|u_{h}(s)\|_{L^{4\delta}}^{\frac{8\delta}{4-d}} + \|u(s)\|_{L^{4\delta}}^{\frac{8\delta}{4-d}}\right) \right) \|u_{h}(s) - u(s)\|_{L^{2}}^{2} ds \\ &+ \int_{0}^{t} \left(\frac{2^{4\delta-2}(2\delta+1)^{2}\beta^{2}}{v} \left(\|u_{h}(s)\|_{L^{4\delta}}^{\frac{8\delta}{4-d}}\right) + C(\alpha,v) \left(\|u_{h}(s)\|_{L^{4\delta}}^{\frac{8\delta}{4-d}} + \|u(s)\|_{L^{4\delta}}^{\frac{8\delta}{4-d}}\right) \right) \|u_{h}(s) - u(s)\|_{L^{4\delta}}^{\frac{8\delta}{4-d}} + \|u(s)\|_{L^{4\delta}}^{\frac{8\delta}{4-d}} + \|u(s)\|_{L^{4\delta}}^{\frac{8\delta}{4-d}} + \|u(s)\|_{L^{4\delta}}^{\frac{8\delta}{4-d}} + \|u(s)\|_{L^{4\delta}}^{\frac{8\delta}{4-d}} + \|u(s)\|_{L^{4\delta}}^{\frac{8\delta}{4-d}} + \|u(s)\|_{L^{4\delta}}^{\frac{8\delta}{4-d}} + \|u(s)\|_{L^{$$

for all  $t \in [0, T]$ , where we estimate the memory term using Lemma 2.1 as

$$\eta \int_0^t \langle K * \nabla(u_h(s) - u(s)), \nabla(u(s) - W(s)) \rangle \, \mathrm{d}s$$

29 of 45

$$\leq \eta \int_{0}^{t} \|K * \nabla(u_{h}(s) - u(s))\|_{L^{2}} \|\nabla(u(s) - W(s))\|_{L^{2}} ds$$

$$\leq \frac{\nu}{4C_{K}} \int_{0}^{t} \|K * \nabla(u_{h}(s) - u(s))\|_{L^{2}}^{2} ds + \frac{C_{K}\eta^{2}}{\nu} \int_{0}^{t} \|\nabla(u(s) - W(s))\|_{L^{2}}^{2} ds$$

$$\leq \frac{\nu}{4} \int_{0}^{t} \|\nabla(u_{h}(s) - u(s))\|_{L^{2}}^{2} ds + \frac{C_{K}\eta^{2}}{\nu} \int_{0}^{t} \|\nabla(u(s) - W(s))\|_{L^{2}}^{2} ds,$$

for the constant  $C_K = \left(\int_0^T |K(t)| dt\right)^2$ . Using the Cauchy-Schwarz inequality and Young's inequality, we estimate  $-2(u_h - u, u - W)$  and  $2(u_h^0 - u_0, u_0 - W(0))$  as

$$-2(u_h - u, u - W) \leq 2||u_h - u||_{L^2} ||u - W||_{L^2} \leq \frac{1}{2} ||u_h - u||_{L^2}^2 + 2||u - W||_{L^2}^2,$$
  
$$2(u_h^0 - u_0, u_0 - W(0)) \leq 2||u_h^0 - u_0||_{L^2} ||u_0 - W(0)||_{L^2} \leq ||u_h^0 - u_0||_{L^2}^2 + ||u_0 - W(0)||_{L^2}^2.$$

Using the above estimates in (3.17) and then applying Gronwall's inequality, we get

$$\begin{split} \sup_{t\in[0,T]} \|u_{h}(t) - u(t)\|_{L^{2}}^{2} + \nu \int_{0}^{T} \|\nabla(u_{h}(t) - u(t))\|_{L^{2}}^{2} dt + \left(\frac{\beta}{2^{2\delta}}\right) \int_{0}^{T} \|u_{h}(t) - u(t)\|_{L^{2}(\delta+1)}^{2(\delta+1)} dt \\ &\leqslant 2 \bigg\{ 2\|u_{h}^{0} - u_{0}\|_{L^{2}}^{2} + \|u_{0} - W(0)\|_{L^{2}}^{2} + \int_{0}^{T} \|\partial_{t}(u(t) - W(t))\|_{L^{2}}^{2} dt \\ &+ \bigg( 2^{2\delta-1}(2\delta+1)\beta + \frac{4}{\nu} + \frac{2^{2(\delta+1)}\alpha^{2}}{\beta} + \frac{\eta}{4C\nu} \bigg) \int_{0}^{T} \|\nabla(u(t) - W(t))\|_{L^{2}}^{2} dt \\ &+ \bigg( 2^{2(\delta+1)}\beta(1+\gamma)^{2}(\delta+1)^{2} + \frac{\beta\gamma}{2} \bigg) \int_{0}^{T} \|u(t) - W(t)\|_{L^{2}}^{2} dt \bigg\} \times e^{(1+2\beta\gamma+C(\beta,\gamma,\delta))T} \\ &\times \exp\bigg\{ \int_{0}^{T} \bigg( \frac{2^{4\delta-2}(2\delta+1)^{2}\beta^{2}}{\nu} \left( \|u_{h}(t)\|_{L^{4\delta}}^{8\delta} + \|u(t)\|_{L^{4\delta}}^{8\delta} \bigg) + C(\alpha,\nu) \bigg( \|u_{h}(t)\|_{L^{4\delta}}^{\frac{8\delta}{4-d}} + \|u(t)\|_{L^{4\delta}}^{\frac{8\delta}{4-d}} \bigg) \bigg) dt \bigg\} \end{aligned}$$

$$(3.18)$$

The term appearing in the exponential is uniformly bounded and is independent of *h* for  $u_0 \in L^{d\delta}$  (see 2.26). Using (3.3), we get

$$\begin{aligned} \|u_0 - W(0)\|_{L^2}^2 &= \|u_0 - R^h u_0\|_{L^2}^2 \leqslant Ch^2 \|u_0\|_{H_0^1}^1, \\ \int_0^T \|u(t) - W(t)\|_{L^2}^2 \, dt &= \int_0^T \|u(t) - R^h u(t)\|_{L^2}^2 \, dt \leqslant Ch^4 \int_0^T \|u(t)\|_{H^2}^2 \, dt, \\ \int_0^T \|\nabla(u(t) - W(t))\|_{L^2}^2 \, dt &= \int_0^T \|\nabla(u(t) - R^h u(t))\|_{L^2}^2 \, dt \leqslant Ch^2 \int_0^T \|u(t)\|_{H^2}^2 \, dt, \\ \int_0^T \|\partial_t (u(t) - W(t))\|_{L^2}^2 \, dt &= \int_0^T \|\partial_t (u(t) - R^h u(t))\|_{L^2}^2 \, dt \leqslant Ch^2 \int_0^T \|\partial_t u(t)\|_{H_0^1}^2 \, dt, \end{aligned}$$

Thus, (3.18) implies that

$$v \int_0^T \|\nabla(u^h(t) - u(t))\|_{L^2}^2 dt \leq C(\|u_0\|_{H_0^1}, v, \alpha, \beta, \gamma, \eta, \delta, T, \|f\|_{L^2(0,T; L^2(\Omega))})$$

$$\times \left\{ \|u_0^h - u_0\|_{\mathbf{L}^2}^2 + h^2 \int_0^T \|\partial_t u(t)\|_{\mathbf{H}_0^1}^2 \, \mathrm{d}t + h^2 \int_0^T \|u(t)\|_{\mathbf{H}_0^1}^2 \, \mathrm{d}t + h^2 \int_0^T \|u(t)\|_{\mathbf{H}^2}^2 \, \mathrm{d}t \right\},$$

and hence the estimate (3.4) follows.

REMARK 3.1 1. For d = 2 with  $\delta \in [1, \infty)$  and d = 3 with  $\delta \in [1, 2]$ , we infer from Theorem 2.5  $\partial_t u \in L^2(0, T; H_0^1(\Omega))$ . For large values of  $\alpha, \beta, \gamma$  (cf. Remark 2.4), one can get the same regularity for the case d = 3 with  $\delta \in (2, \infty)$ .

2. In the above Theorem we need  $\partial_t u \in L^2(0,T; H^1_0(\Omega))$ . However, if we assume  $u_h \in H^1(0,T; V_h)$  and  $P_h$  denotes the  $L^2(\Omega)$  projection from  $L^2(\Omega)$  onto  $V_h$  as in Hou & Zhu (2006) and Chrysafinos & Hou (2002), namely, for each  $v \in L^2(\Omega)$ 

$$(P_h v - v, w^h) = 0$$
 for all  $w^h \in V_h$ , a.e.  $t$ . (3.19)

Also, there exists a constant C independent of v and h such that

$$\|u - P_h v\|_{\mathbf{H}^r} \leq C \|u - R_h v\|_{\mathbf{H}^r}, \quad \text{for all} \quad u \in \mathbf{H}^1_0(\Omega), \ r \in [0, 1],$$

We can estimate the first term on the right hand side of (3.7) using the L<sup>2</sup> projection (3.19) repeatedly as

$$-\langle \partial_t (u_h(t) - u(t)), u(t) - P_h u(t) \rangle = \langle \partial_t u(t), u(t) - P_h u(t) \rangle$$
$$= \langle \partial_t (u(t) - P_h u(t)), u(t) - P_h u(t) \rangle = \frac{1}{2} \frac{\mathrm{d}}{\mathrm{d}t} \| u(t) - P_h u(t) \|_{\mathrm{L}^2}^2$$

for a.e.  $t \in [0, T]$ , which upon integration can be estimated as

$$\int_0^T \|u(t) - P_h u(t)\|_{L^2}^2 dt \leq \int_0^T \|u(t) - R^h u(t)\|_{L^2}^2 dt \leq Ch^4 \int_0^T \|u(t)\|_{H^2}^2 dt.$$

So, we need only our initial data  $u_0 \in L^{d\delta}(\Omega) \cap H^1_0(\Omega)$  and  $\partial_t u \in L^2(0,T;L^2(\Omega))$ .

### 3.2 Fully-discrete Galerkin approximation and error estimates

In this section, we introduce a fully-discrete conforming finite element scheme. We partition the time interval [0,T] into  $0 = t_0 < t_1 < t_2 \cdots < t_N = T$  with uniform time stepping  $\Delta t$ , that is,  $t_k = k\Delta t$ . We replace the time derivative with a backward Euler discretization and approximating the memory term in the following way: We shall approximate  $\psi$  in  $J(\psi) = \int_0^t K(t-s)\psi(s)ds$  by the piecewise constant function taking value  $\psi^k = \psi(t_k)$  in  $(t_{k-1}, t_k)$  and the quadrature rule as

$$J(\boldsymbol{\psi}) = \int_0^t K(t-s)\boldsymbol{\psi}(s) \mathrm{d}s \approx \frac{1}{(\Delta t)^2} \int_{t_{k-1}}^{t_k} \int_0^t K(t-s)\Delta t \boldsymbol{\psi}(s) \, \mathrm{d}s \mathrm{d}t = \sum_{j=1}^k \omega_{kj} \Delta t \, \boldsymbol{\psi}^j,$$

where  $\omega_{kj} = \frac{1}{(\Delta t)^2} \int_{t_{k-1}}^{t_k} \int_{t_{j-1}}^{\min(t,t_j)} K(t-s) \, ds \, dt$ , for  $1 \le k \le N$ . We denote *C* as a generic constant which is independent of *h* and  $\Delta t$ . We then define a generic finite element approximate solution  $u_{kh}$  by

$$u_{kh}|_{[t_{k-1},t_k]} = u_h^{k-1} + \left(\frac{t - t_{k-1}}{\Delta t}\right) (u_h^k - u_h^{k-1}), \qquad 1 \le k \le N, \qquad \text{for } t \in [t_{k-1},t_k].$$
(3.20)

Note here that  $\partial_t u_{kh} = \frac{u_h^k - u_h^{k-1}}{\Delta t}$ . The continuous time Galerkin approximation of the problem (2.6) is defined in the following way: Find  $u_h^k \in V_h$ , for k = 1, 2, ..., N such that for  $\chi \in V_h$ :

$$\begin{cases} \left(\frac{u_h^k - u_h^{k-1}}{\Delta t}, \chi\right) + \mathbf{v}a(u_h^k, \chi) + \alpha b(u_h^k, u_h^k, \chi) + \left(\sum_{j=1}^k \omega_{kj}(\Delta t) \nabla u_h^j, \nabla \chi\right) \\ &= \beta(u_h^k(1 - (u_h^k)^\delta)((u_h^k)^\delta - \gamma), \chi) + \langle f^k, \chi \rangle, \\ &(u_h^0(x_i), \chi) = (u_0(x_i), \chi), \text{ for } i = 1, 2, \dots, N, \end{cases}$$
(3.21)

 $f^k = (\Delta t)^{-1} \int_{t_{k-1}}^{t_k} f(s) ds$  for  $f \in L^2(0, T; H^{-1}(\Omega))$ ,  $u_h^0$  approximates  $u_0$  in  $V_h$  and  $x_i$ , i = 1, ..., n, represent the nodes of the triangulation  $\mathscr{T}^h$  of  $\Omega$ .

The existence of the discrete solution can be proved using an application of Brouwer's fixed point theorem [Lemma 1.4, Temam (2001)] as done in Theorem 3.1 Khan *et al.* (2021). To show the error estimates, we need a stability result. It is natural to see that for the quadratic form analogous to the double integral in (1.2), we have

$$\sum_{k=1}^{N} \left( \sum_{j=1}^{k} \omega_{kj} \Delta t \nabla u_{h}^{j}, \nabla u_{h}^{k} \right) \ge 0,$$
(3.22)

as discussed in Mustapha & McLean (2009) and Lemma 4.7 McLean & Thomée (1993). The following lemma is useful for the analysis:

LEMMA 3.1 Assume  $f \in L^2(0,T; H^{-1}(\Omega))$ . Then the set  $\{f^k\}_{k=1}^N$  defined by  $f^k = (\Delta t)^{-1} \int_{t_{k-1}}^{t_k} f(s) ds$  satisfies

$$\Delta t \sum_{k=1}^{N} \|f^{k}\|_{\mathbf{H}^{-1}}^{2} \leqslant C \|f\|_{\mathbf{L}^{2}(0,T;\mathbf{H}^{-1}(\Omega))}^{2}$$

and

$$\sum_{k=1}^{N} \int_{t_{k-1}}^{t_{k}} \|f^{k} - f(t)\|_{\mathbf{H}^{-1}}^{2} \mathrm{d}t \to 0 \quad \Delta t \to 0.$$

If  $f \in H^{\gamma}(0,T; \mathrm{H}^{-1}(\Omega))$  for some  $\gamma \in [0,1]$ , then

$$\sum_{k=1}^{N} \int_{t_{k-1}}^{t_{k}} \|f^{k} - f(t)\|_{\mathbf{H}^{-1}}^{2} \mathrm{d}t \leqslant C(\Delta t)^{2\gamma} \|f\|_{\mathbf{H}^{\gamma}(0,T;\mathbf{H}^{-1}(\Omega))}^{2}$$

*Proof.* For a proof, we refer to Lemma 3.2, Hou & Zhu (2006).

Let us now discuss the stability of the fully-discrete approximation (3.21).

LEMMA 3.2 (Stability) Assume  $f \in L^2(0,T; H^{-1}(\Omega))$  and  $u_h^0 \in L^2(\Omega)$ . Let  $\{u_h^k\}_{k=1}^N \subset V_h$  be defined by (3.21). Then

$$\sum_{k=1}^{N} \Delta t \|\nabla u_{h}^{k}\|_{\mathrm{L}^{2}}^{2} \leqslant \left(\frac{1}{\nu} \|f\|_{\mathrm{L}^{2}(0,T;H^{-1}(\Omega))}^{2} + \|u_{h}^{0}\|_{\mathrm{L}^{2}}^{2}\right) \times e^{\beta T(1+\gamma)^{2}}.$$
(3.23)

*Proof.* Suppose  $f \in L^2(0,T; H^{-1}(\Omega))$ , taking  $\chi = u_h^k$  in (3.21), we obtain

$$\frac{1}{2\Delta t} \|u_h^k\|_{L^2}^2 - \frac{1}{2\Delta t} \|u_h^{k-1}\|_{L^2}^2 + \frac{1}{2\Delta t} \|u_h^k - u_h^{k-1}\|_{L^2}^2 + \nu a(u_h^k, u_h^k) + \alpha b(u_h^k, u_h^k, u_h^k) + \eta \left(\sum_{j=1}^k \Delta t \omega_{kj} \nabla u_h^j, \nabla u_h^k\right) = \beta (u_h^k (1 - (u_h^k)^\delta) ((u_h^k)^\delta - \gamma), u_h^k) + \langle f^k, u_h^k \rangle.$$

Using (2.1), (2.3), Cauchy-Schwarz and Young's inequalities, we have

$$\frac{1}{2\Delta t} \|u_h^k\|_{L^2}^2 - \frac{1}{2\Delta t} \|u_h^{k-1}\|_{L^2}^2 + \frac{1}{2\Delta t} \|u_h^k - u_h^{k-1}\|_{L^2}^2 + \nu \|\nabla u_h^k\|_{L^2}^2 + \beta \gamma \|u_h^k\|_{L^2}^2 + \beta \|u_h^k\|_{L^{2(\delta+1)}}^{2(\delta+1)} + \eta \left(\sum_{j=1}^k \Delta t \omega_{kj} \nabla u_h^j, \nabla u_h^k\right) \leqslant \frac{\beta}{2} \|u_h^k\|_{L^{2(\delta+1)}}^{2(\delta+1)} + \frac{\beta (1+\gamma)^2}{2} \|u_h^k\|_{L^2}^2 + \frac{1}{2\nu} \|f^k\|_{H^{-1}}^2 + \frac{\nu}{2} \|\nabla u_h^k\|_{L^2}^2$$

This may be simplified as

$$\frac{1}{2\Delta t} \|u_h^k\|_{L^2}^2 - \frac{1}{2\Delta t} \|u_h^{k-1}\|_{L^2}^2 + \frac{\nu}{2} \|\nabla u_h^k\|_{L^2}^2 + \eta \left(\sum_{j=1}^k \omega_{kj} \Delta t \nabla u_h^j, \nabla u_h^k\right) \leq \frac{\beta (1+\gamma)^2}{2} \|u_h^k\|_{L^2}^2 + \frac{1}{2\nu} \|f^k\|_{H^{-1}}^2$$

Summing over k for k = 1, 2, ..., N and using positivity of the Kernel  $K(\cdot)$  (3.22) and Lemma 3.1, we have

$$\frac{1}{\Delta t} \|u_h^N\|_{\mathrm{L}^2}^2 - \frac{1}{\Delta t} \|u_h^0\|_{\mathrm{L}^2}^2 + v \sum_{k=1}^N \|\nabla u_h^k\|_{\mathrm{L}^2}^2 \leqslant \beta (1+\gamma)^2 \sum_{k=1}^N \|u_h^k\|_{\mathrm{L}^2}^2 + \frac{1}{v\Delta t} \Delta t \sum_{k=1}^N \|f^k\|_{\mathrm{H}^{-1}}^2$$

Finally, an application of the discrete Gronwall inequality (Lemma 9 Pani & Peterson (1996)) yields the required bound (3.23).  $\Box$ 

LEMMA 3.3 Let  $1 \leq \delta < \infty$  for d = 2 and,  $1 \leq \delta \leq 2$  for d = 3 and assume that  $u_0 \in H^2(\Omega) \cap H^1_0(\Omega)$ and  $f \in H^1(0,T; L^2(\Omega))$ . Let  $u_h$  be the semi-discrete solution (3.2) and  $u_{kh}$  be the generic fully-discrete solution defined in (3.20), then

$$\begin{aligned} \|u_{h} - u_{kh}\|_{\mathbf{L}^{\infty}(0,T;\mathbf{L}^{2}(\Omega))}^{2} + \|u_{h} - u_{kh}\|_{\mathbf{L}^{2}(0,T;\mathbf{H}_{0}^{1}(\Omega))}^{2} &\leq C(\Delta t)^{2}(\|f\|_{\mathbf{H}^{1}(0,T;\mathbf{L}^{2}(\Omega))}^{2} + \|u_{0}\|_{\mathbf{H}^{2}}^{2}) \\ &+ \eta^{2}(\Delta t)^{2} \sup_{k,j} \omega_{kj}^{2}(\|f\|_{\mathbf{H}^{1}(0,T;\mathbf{L}^{2}(\Omega))}^{2} + \|u_{0}\|_{\mathbf{H}^{2}}^{2}). \end{aligned}$$

*Proof.* We divide our proof into two steps: In Step 1, we prove the error estimates on the nodal values and in Step 2, we prove the final estimate.

**Step 1:** *Estimates at t<sub>k</sub>* : Integrating the semi-discrete scheme (3.2) from  $t_{k-1}$  to  $t_k$ , we obtain

$$(u_{h}(t_{k}) - u_{h}(t_{k-1}), \boldsymbol{\chi}) + \boldsymbol{\nu} \left( \int_{t_{k-1}}^{t_{k}} \nabla u_{h}(t) \, \mathrm{d}t, \nabla \boldsymbol{\chi} \right) + \boldsymbol{\alpha} \left( \int_{t_{k-1}}^{t_{k}} B(u_{h}(t)) \, \mathrm{d}t, \boldsymbol{\chi} \right) + \boldsymbol{\eta} \left( \int_{t_{k-1}}^{t_{k}} (K * \nabla u_{h})(t) \, \mathrm{d}t, \nabla \boldsymbol{\chi} \right) = \boldsymbol{\beta} \left( \int_{t_{k-1}}^{t_{k}} c(u_{h}(t)) \, \mathrm{d}t, \boldsymbol{\chi} \right) + \left( \int_{t_{k-1}}^{t_{k}} f(t) \, \mathrm{d}t, \boldsymbol{\chi} \right).$$
(3.24)

The fully-discrete scheme (3.21) is given as

$$\left(\frac{u_h^k - u_h^{k-1}}{\Delta t}, \chi\right) + v(\nabla u_h^k, \nabla \chi) + \alpha b(u_h^k, u_h^k, \chi) + \left(\sum_{j=1}^k \omega_{kj} \Delta t \nabla u_h^k, \nabla \chi\right)$$

# GENERALIZED BURGERS-HUXLEY EQUATION WITH WEAKLY SINGULAR KERNEL 33 of 45

$$=\beta(u_{h}^{k}(1-(u_{h}^{k})^{\delta})((u_{h}^{k})^{\delta}-\gamma),\chi)+(f^{k},\chi).$$
(3.25)

Subtracting (3.25) from (3.24), we have

$$(u_{h}(t_{k})-u_{h}^{k},\boldsymbol{\chi})-(u_{h}(t_{k-1})-u_{h}^{k-1},\boldsymbol{\chi})+\boldsymbol{v}\left(\int_{t_{k-1}}^{t_{k}}\nabla u_{h}(t)\,\mathrm{d}t-\Delta t\nabla u_{h}^{k},\nabla\boldsymbol{\chi}\right)+\eta\left(\int_{t_{k-1}}^{t_{k}}(K*\nabla u_{h})(t)\,\mathrm{d}t,\nabla\boldsymbol{\chi}\right)$$
$$-\Delta t\eta\left(\sum_{j=1}^{k}\Delta t\omega_{kj}\nabla u_{h}^{j},\nabla\boldsymbol{\chi}\right)+\alpha\left(\left(\int_{t_{k-1}}^{t_{k}}B(u_{h}(t))\,\mathrm{d}t\right)-\Delta tB(u_{h}^{k}),\boldsymbol{\chi}\right)=\beta\left(\int_{t_{k-1}}^{t_{k}}c(u_{h}(t))\,\mathrm{d}t-\Delta tc(u_{h}^{k}),\boldsymbol{\chi}\right).$$

Rewriting the above equation and taking  $\chi = u_h(t_k) - u_h^k$ , we deduce

$$\begin{split} \|u_{h}(t_{k}) - u_{h}^{k}\|_{L^{2}}^{2} + \nu\Delta t \|\nabla(u_{h}(t_{k}) - u_{h}^{k})\|_{L^{2}}^{2} + \Delta t \eta \left(\sum_{j=1}^{k} \omega_{kj}\Delta t \nabla(u_{h}(t_{j}) - u_{h}^{j}), \nabla(u_{h}(t_{k}) - u_{h}^{k})\right) \\ &= (u_{h}(t_{k-1}) - u_{h}^{k-1}, u_{h}(t_{k}) - u_{h}^{k}) - \nu \left(\int_{t_{k-1}}^{t_{k}} \nabla u_{h}(t) \, dt - \Delta t \nabla u_{h}(t_{k}), \nabla(u_{h}(t_{k}) - u_{h}^{k})\right) \\ &+ \eta \left(\Delta t \sum_{j=1}^{k} \omega_{kj}\Delta t \nabla u_{h}(t_{j}) - \int_{t_{k-1}}^{t_{k}} (K * \nabla u_{h})(t) \, dt, \nabla(u_{h}(t_{k}) - u_{h}^{k})\right) \\ &- \alpha \left(\int_{t_{k-1}}^{t_{k}} B(u_{h}(t)) \, dt - \Delta t B(u_{h}(t_{k})), u_{h}(t_{k}) - u_{h}^{k}\right) - \alpha\Delta t \left(B(u_{h}(t_{k})) - B(u_{h}^{k}), u_{h}(t_{k}) - u_{h}^{k}\right) \\ &+ \beta \left(\int_{t_{k-1}}^{t_{k}} c(u_{h}(t)) \, dt - \Delta t c(u_{h}(t_{k})), u_{h}(t_{k}) - u_{h}^{k}\right) + \beta\Delta t \left(c(u_{h}(t_{k})) - c(u_{h}^{k}), u_{h}(t_{k}) - u_{h}^{k}\right). \end{split}$$

Using Cauchy-Schwarz and Young's inequalities, we can estimate the first term on right hand side as

$$(u_h(t_{k-1}) - u_h^{k-1}, u_h(t_k) - u_h^k) \leqslant \frac{1}{2} \|u_h(t_{k-1}) - u_h^{k-1}\|_{L^2}^2 + \frac{1}{2} \|u_h(t_k) - u_h^k\|_{L^2}^2,$$
(3.26)

Again an application of the Cauchy-Schwarz and Young's inequalities yield

$$\begin{split} \mathbf{v} &\left( \int_{t_{k-1}}^{t_k} \nabla u_h(t) \, \mathrm{d}t - \Delta t \nabla u_h(t_k), \nabla (u_h(t_k) - u_h^k) \right) \\ &\leqslant \mathbf{v} \Big\| \int_{t_{k-1}}^{t_k} \nabla u_h(t) \, \mathrm{d}t - \Delta t \nabla u_h(t_k) \Big\|_{\mathrm{L}^2} \Big\| \nabla (u_h(t_k) - u_h^k) \Big\|_{\mathrm{L}^2} \\ &\leqslant \frac{2}{\mathbf{v} \Delta t} \Big\| \int_{t_{k-1}}^{t_k} \nabla u_h(t) \, \mathrm{d}t - \Delta t \nabla u_h(t_k) \Big\|_{\mathrm{L}^2}^2 + \frac{\mathbf{v} \Delta t}{8} \Big\| \nabla (u_h(t_k) - u_h^k) \Big\|_{\mathrm{L}^2}^2, \end{split}$$

where the first term on the right hand side can be estimated as

$$\frac{1}{\Delta t} \left\| \int_{t_{k-1}}^{t_k} \nabla u_h(t) \, dt - \Delta t \nabla u_h(t_k) \right\|_{L^2}^2 = \frac{1}{\Delta t} \left\| \int_{t_{k-1}}^{t_k} (\nabla u_h(t) - \nabla u_h(t_k)) \, dt \right\|_{L^2}^2 
= \frac{1}{\Delta t} \left\| \int_{t_{k-1}}^{t_k} \int_{t_k}^t \nabla \partial_t u_h(s) \, ds \, dt \right\|_{L^2}^2 \leqslant \frac{1}{\Delta t} \left( \int_{t_{k-1}}^{t_k} \int_{t_{k-1}}^{t_k} \| \partial_t \nabla u_h(s) \|_{L^2} \, ds \, dt \right)^2 
\leqslant \Delta t \left( \int_{t_{k-1}}^{t_k} \| \partial_t \nabla u_h(s) \|_{L^2} \, ds \right)^2 \leqslant (\Delta t)^2 \int_{t_{k-1}}^{t_k} \| \partial_t \nabla u_h(s) \|_{L^2}^2 \, ds.$$
(3.27)

The memory term can be estimated as

$$\eta \left( \Delta t \sum_{j=1}^{k} \Delta t \omega_{kj} \nabla u_h(t_j) - \int_{t_{k-1}}^{t_k} (K * \nabla u_h)(t) \, \mathrm{d}t, \nabla (u_h(t_k) - u_h^k) \right)$$

$$\leq \eta \left\| \Delta t \sum_{j=1}^{k} \Delta t \omega_{kj} \nabla u_h(t_j) - \int_{t_{k-1}}^{t_k} (K * \nabla u_h)(t) \, \mathrm{d}t \right\|_{\mathrm{L}^2} \| \nabla (u_h(t_k) - u_h^k) \|_{\mathrm{L}^2}$$

$$\leq \frac{2\eta^2}{\nu \Delta t} \left\| (\Delta t)^2 \sum_{j=1}^{k} \omega_{kj} \nabla u_h(t_j) - \int_{t_{k-1}}^{t_k} (K * \nabla u_h)(t) \, \mathrm{d}t \right\|_{\mathrm{L}^2}^2 + \frac{\nu \Delta t}{8} \| \nabla (u_h(t_k) - u_h^k) \|_{\mathrm{L}^2}^2$$

where the term  $\frac{2\eta^2}{v\Delta t} \| (\Delta t)^2 \sum_{j=1}^k \omega_{kj} \nabla u_h(t_j) - \int_{t_{k-1}}^{t_k} (K * \nabla u_h)(t) dt \|_{L^2}^2$  can be estimated using (3.27) as

$$\begin{split} &\frac{1}{\Delta t} \left\| (\Delta t)^2 \sum_{j=1}^k \omega_{kj} \nabla u_h(t_j) - \int_{t_{k-1}}^{t_k} (K * \nabla u_h)(t) \, \mathrm{d}t \right\|_{\mathrm{L}^2}^2 \\ &= \frac{1}{\Delta t} \left\| \int_{t_{k-1}}^{t_k} \sum_{j=1}^k \left( \int_{t_{j-1}}^{\min(t,t_j)} K(t-s) (\nabla u_h(t_j) - \nabla u_h(s)) \, \mathrm{d}s \right) \, \mathrm{d}t \right\|_{\mathrm{L}^2}^2 \\ &\leqslant \frac{1}{\Delta t} \left\| \sum_{j=1}^k \int_{t_{k-1}}^{t_k} \left( \int_{t_{j-1}}^{\min(t,t_j)} |K(t-s)| \int_{t_{j-1}}^{t_j} |\partial_t \nabla u_h(\tau)| \, \mathrm{d}\tau \right) \, \mathrm{d}s \, \mathrm{d}t \right\|_{\mathrm{L}^2}^2 \\ &\leqslant \frac{1}{\Delta t} \left\| \sum_{j=1}^k (\Delta t)^2 \left( \frac{1}{(\Delta t)^2} \int_{t_{k-1}}^{t_k} \int_{t_{j-1}}^{\min(t,t_j)} |K(t-s)| \, \mathrm{d}s \, \mathrm{d}t \int_{t_{j-1}}^{t_j} |\partial_t \nabla u_h(\tau)| \, \mathrm{d}\tau \right) \right\|_{\mathrm{L}^2}^2 \\ &\leqslant (\Delta t)^3 \left\| \sum_{j=1}^k \overline{K}_{kj} \int_{t_{j-1}}^{t_j} |\partial_t \nabla u_h(\tau)| \, \mathrm{d}\tau \right\|_{\mathrm{L}^2}^2 \\ &\leqslant K (\Delta t)^4 \sum_{j=1}^k \overline{K}_{kj}^2 \int_{t_{j-1}}^{t_j} \|\partial_t \nabla u_h(\tau)\|_{\mathrm{L}^2}^2 \, \mathrm{d}\tau \\ &\leqslant T (\Delta t)^3 \sum_{j=1}^k \overline{K}_{kj}^2 \int_{t_{j-1}}^{t_j} \|\partial_t \nabla u_h(\tau)\|_{\mathrm{L}^2}^2 \, \mathrm{d}\tau, \end{split}$$

where  $\overline{K}_{kj} = \frac{1}{(\Delta t)^2} \int_{t_{k-1}}^{t_k} \int_{t_{j-1}}^{\min(t,t_j)} |K(t-s)| \, ds \, dt$ . An application of integration by parts and Cauchy-Schwarz inequality gives

$$\alpha \left( \int_{t_{k-1}}^{t_k} B(u_h(t)) \, dt - \Delta t B(u_h(t_k)), u_h(t_k) - u_h^k \right)$$

$$= \frac{\alpha}{\delta + 1} \sum_{i=1}^d \left( \int_{t_{k-1}}^{t_k} u_h^{\delta + 1}(t) \, dt - \Delta t u_h^{\delta + 1}(t_k), \frac{\partial}{\partial x_i} (u_h(t_k) - u_h^k) \right)$$

$$\leqslant \frac{\alpha}{\delta + 1} \left\| \int_{t_{k-1}}^{t_k} u_h^{\delta + 1}(t) \, dt - \Delta t u_h^{\delta + 1}(t_k) \right\|_{L^2} \left\| \nabla (u_h(t_k) - u_h^k) \right\|_{L^2}$$

$$\leqslant \frac{2\alpha^2}{\nu(\delta + 1)^2 \Delta t} \left\| \int_{t_{k-1}}^{t_k} u_h^{\delta + 1}(t) \, dt - \Delta t u_h^{\delta + 1}(t_k) \right\|_{L^2}^2 + \frac{\nu \Delta t}{8} \left\| \nabla (u_h(t_k) - u_h^k) \right\|_{L^2}.$$

$$(3.28)$$

+

# GENERALIZED BURGERS-HUXLEY EQUATION WITH WEAKLY SINGULAR KERNEL

Now, we estimate  $\frac{1}{\Delta t} \left\| \int_{t_{k-1}}^{t_k} u_h^{\delta+1}(t) dt - \Delta t u_h^{\delta+1}(t_k) \right\|_{L^2}^2$  similar to (3.27) as

$$\begin{split} &\frac{1}{\Delta t} \left\| \int_{t_{k-1}}^{t_k} u_h^{\delta+1}(t) \, \mathrm{d}t - \Delta t u_h^{\delta+1}(t_k) \right\|_{\mathrm{L}^2}^2 = \frac{1}{\Delta t} \left\| \int_{t_{k-1}}^{t_k} (u_h^{\delta+1}(t) - u_h^{\delta+1}(t_k)) \, \mathrm{d}t \right\|_{\mathrm{L}^2}^2 \\ &= \frac{1}{\Delta t} \left\| \int_{t_{k-1}}^{t_k} \int_{t_k}^{t} \partial_t u_h^{\delta+1}(s) \, \mathrm{d}s \, \mathrm{d}t \right\|_{\mathrm{L}^2}^2 \leqslant \frac{1}{\Delta t} \left( \int_{t_{k-1}}^{t_k} \int_{t_{k-1}}^{t_k} \| \partial_t u_h^{\delta+1}(s) \|_{\mathrm{L}^2} \, \mathrm{d}s \, \mathrm{d}t \right)^2 \\ &\leqslant \Delta t \left( \int_{t_{k-1}}^{t_k} \| \partial_t u_h^{\delta+1}(s) \|_{\mathrm{L}^2} \, \mathrm{d}s \right)^2 \leqslant (\Delta t)^2 \int_{t_{k-1}}^{t_k} \| \partial_t u_h^{\delta+1}(s) \|_{\mathrm{L}^2}^2 \, \mathrm{d}s \\ &= (\Delta t)^2 (\delta + 1)^2 \int_{t_{k-1}}^{t_k} \| u_h^{\delta}(s) \partial_t u_h(s) \|_{\mathrm{L}^2}^2 \, \mathrm{d}s. \end{split}$$

Finally the reaction term can be simplified as

$$\left(\int_{t_{k-1}}^{t_k} c(u_h(t)) \, \mathrm{d}t\right) - \Delta t c(u_h(t_k)) = \beta (1+\gamma) \left(\int_{t_{k-1}}^{t_k} u_h^{\delta+1}(t) \, \mathrm{d}t - \Delta t u_h^{\delta+1}(t_k)\right) \\ - \beta \gamma \left(\left(\int_{t_{k-1}}^{t_k} u_h(t) \, \mathrm{d}t\right) - \Delta t u_h(t_k)\right) - \beta \left(\int_{t_{k-1}}^{t_k} u_h^{2\delta+1}(t) \, \mathrm{d}t - \Delta t u_h^{2\delta+1}(t_k)\right).$$

Firstly, we estimate the term with coefficient  $\gamma$ , using Cauchy-Schwarz inequality and the estimate similar to (3.27) as

$$\begin{split} \beta \gamma \left( \int_{t_{k-1}}^{t_k} u_h(t) \, \mathrm{d}t - \Delta t u_h(t_k), u_h(t_k) - u_h^k \right) &\leq \frac{(\beta \gamma)^2}{\nu \Delta t} \left\| \int_{t_{k-1}}^{t_k} u_h(t) \, \mathrm{d}t - \Delta t u_h(t_k) \right\|_{\mathrm{L}^2}^2 + \frac{\nu \Delta t}{8} \|u_h(t_k) - u_h^k\|_{\mathrm{L}^2}^2 \\ &\leq \frac{(\beta \gamma)^2 (\Delta t)^2}{\nu} \int_{t_{k-1}}^{t_k} \|\partial_t u_h(s)\|_{\mathrm{L}^2}^2 \, \mathrm{d}s + \frac{\nu \Delta t}{8} \|(u_h(t_k) - u_h^k)\|_{\mathrm{L}^2}^2. \end{split}$$

Using an estimate similar to (3.28), we have

$$\beta(1+\gamma)\left(\int_{t_{k-1}}^{t_k} u_h^{\delta+1}(t) \, \mathrm{d}t - \Delta t u_h^{\delta+1}(t_k), u_h(t_k) - u_h^k\right) \\ \leqslant \frac{(\Delta t)^2 (1+\gamma)^2 \beta^2 (\delta+1)^2}{\nu} \int_{t_{k-1}}^{t_k} \|u_h^{\delta}(s) \partial_t u_h(s)\|_{\mathrm{L}^2}^2 + \frac{\nu \Delta t}{8} \|u_h(t_k) - u_h^k\|_{\mathrm{L}^2}^2.$$

Also for the final term, we obtain

$$\begin{split} \left| \beta \left( \int_{t_{k-1}}^{t_k} u_h^{2\delta+1}(t) \, \mathrm{d}t - \Delta t u_h^{2\delta+1}(t_k), u_h(t_k) - u_h^k \right) \right| &= \left| \beta \int_{t_{k-1}}^{t_k} \left( u_h^{2\delta+1}(t) - u_h^{2\delta+1}(t_k), u_h(t_k) - u_h^k \right) \, \mathrm{d}t \right| \\ &= \left| \beta \int_{t_{k-1}}^{t_k} \left( \int_{t_k}^{t} \partial_t u_h^{2\delta+1}(s) \, \mathrm{d}s, u_h(t_k) - u_h^k \right) \, \mathrm{d}t \right| \leq \beta (2\delta+1) \int_{t_{k-1}}^{t_k} \int_{t_{k-1}}^{t_k} \left| \left( u_h^{2\delta}(s) \partial_t u_h(s), u_h(t_k) - u_h^k \right) \right| \, \mathrm{d}s \, \mathrm{d}t \\ &\leq \beta (2\delta+1) \Delta t \int_{t_{k-1}}^{t_k} \left| \left( u_h^{\delta}(t) (u_h^{\delta}(t) \partial_t u_h(t)), u_h(t_k) - u_h^k \right) \right| \, \mathrm{d}t \\ &\leq \beta (2\delta+1) \Delta t \int_{t_{k-1}}^{t_k} \left\| u_h^{\delta}(t) \right\|_{\mathrm{L}^{\frac{2(\delta+1)}{\delta}}} \left\| u_h^{\delta}(t) \partial_t u_h(t) \right\|_{\mathrm{L}^2} \left\| u_h(t_k) - u_h^k \right\|_{\mathrm{L}^{2(\delta+1)}} \, \mathrm{d}t \\ &\leq \beta (2\delta+1) \Delta t \int_{t_{k-1}}^{t_k} \left\| u_h^{\delta}(t) \right\|_{\mathrm{L}^{\frac{2(\delta+1)}{\delta}}} \left\| u_h^{\delta}(t) \partial_t u_h(t) \right\|_{\mathrm{L}^2} \left\| u_h(t_k) - u_h^k \right\|_{\mathrm{L}^{2(\delta+1)}} \, \mathrm{d}t \\ &\leq \beta (2\delta+1) \Delta t \left\| u_h \right\|_{\mathrm{L}^{\infty}((t_{k-1}, t_k); \mathrm{L}^{2(\delta+1)}(\Omega))} \int_{t_{k-1}}^{t_k} \left\| u_h^{\delta}(t) \partial_t u_h(t) \right\|_{\mathrm{L}^2} \left\| \nabla (u_h(t_k) - u_h^k) \right\|_{\mathrm{L}^2} \, \mathrm{d}t \end{split}$$

$$\leq \frac{2\beta^{2}(2(\delta+1))^{2}(\Delta t)^{2}}{\nu} \|u_{h}\|_{L^{\infty}(0,T;L^{2(\delta+1)}(\Omega))}^{2\delta} \int_{t_{k-1}}^{t_{k}} \|u_{h}^{\delta}(t)\partial_{t}u_{h}(t)\|_{L^{2}}^{2} dt + \frac{\nu\Delta t}{8} \|\nabla(u_{h}(t_{k}) - u_{h}^{k})\|_{L^{2}}^{2} dt + \frac{\nu\Delta t}{8} \|\nabla(u_{h}(t_{k}) - u_{h}^{k})\|_{L^{2$$

for  $1 \le \delta < \infty$  (d = 2) and  $1 \le \delta \le 2$  (d = 3). For,  $w = u_h(t_k) - u_h^k$ , we derive a bound for  $-\alpha \Delta t(B(u_h(t_k)) - B(u_h^k), w)$ , using Taylor's formula, Hölder's, Gagliardo-Nirenberg interpolation and Young's inequalities as

$$\begin{aligned} -\alpha \Delta t \left( B(u_h(t_k)) - B(u_h^k), u_h(t_k) - u_h^k \right) &= \frac{\alpha}{\delta + 1} \left( (u_h(t_k)^{\delta + 1} - (u_h^k)^{\delta + 1}) \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix}, \nabla w \right) \\ &= \alpha \left( (u_h(t_k) - u_h^k) (\theta u_h(t_k) + (1 - \theta) u_h^k)^{\delta} \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix}, \nabla w \right) \\ &\leqslant 2^{\delta - 1} \alpha (\|u_h(t_k)\|_{L^{2(\delta + 1)}}^{\delta} + \|u_h^k\|_{L^{2(\delta + 1)}}^{\delta})) \|w\|_{L^{2(\delta + 1)}} \|\nabla w\|_{L^2} \\ &\leqslant 2^{\delta - 1} C \alpha (\|u_h(t_k)\|_{L^{2(\delta + 1)}}^{\delta} + \|u_h^k\|_{L^{2(\delta + 1)}}^{\delta})) \|\nabla w\|_{L^2}^{\frac{(2 - d)\delta + 2}{2(\delta + 1)}} \\ &\leqslant \frac{\nu}{8} \|\nabla w\|_{L^2}^2 + C(\alpha, \nu) \left( \|u_h(t_k)\|_{L^{2(\delta + 1)}}^{\frac{4\delta(\delta + 1)}{2}} + \|u_h^k\|_{L^{2(\delta + 1)}}^{\frac{4\delta(\delta + 1)}{2}} + \|u_h^k\|_{L^{2(\delta + 1)}}^{\frac{4\delta(\delta + 1)}{2}} \right) \|w\|_{L^2}^2. \end{aligned}$$

where  $C(\alpha, \nu) = \left(\frac{2((2+d)\delta+2)}{\nu(\delta+1)}\right)^{\frac{(2-d)\delta+2}{(2+d)\delta+2}} \times \left(\frac{(2-d)\delta+2}{4(\delta+1)}\right) (2^{\delta-1}\alpha)^{\frac{4(\delta+1)}{(2-d)\delta+2}}$ . Combining the above estimates and using the calculations similar to (2.22), we deduce

$$\begin{split} &\frac{1}{2} \|u_{h}(t_{k}) - u_{h}^{k}\|_{\mathrm{L}^{2}}^{2} - \frac{1}{2} \|u_{h}(t_{k-1}) - u_{h}^{k-1}\|_{\mathrm{L}^{2}}^{2} + \frac{\nu}{2} \Delta t \|\nabla(u_{h}(t_{k}) - u_{h}^{k})\|_{\mathrm{L}^{2}}^{2} \\ &+ \Delta t \eta \left( \sum_{j=1}^{k} \omega_{kj} \Delta t \nabla(u_{h}(t_{j}) - u_{h}^{j}), \nabla(u_{h}(t_{k}) - u_{h}^{k}) \right) + \frac{\beta \Delta t}{2} \|u_{h}(t_{k})^{\delta}(u_{h}(t_{k}) - u_{h}^{k})\|_{\mathrm{L}^{2}}^{2} \\ &+ \frac{\beta \Delta t}{2} \|(u_{h}^{k})^{\delta}(u_{h}(t_{k}) - u_{h}^{k})\|_{\mathrm{L}^{2}}^{2} + \beta \gamma \Delta t \|(u_{h}(t_{k}) - u_{h}^{k})\|_{\mathrm{L}^{2}}^{2} \\ \leqslant \frac{2\alpha^{2} (\Delta t)^{2}}{\nu} \int_{t_{k-1}}^{t_{k}} \|\partial_{t} \nabla u_{h}(s)\|_{\mathrm{L}^{2}}^{2} \, ds + \frac{(\Delta t)^{2} \beta^{2} \gamma^{2}}{\nu} \int_{t_{k-1}}^{t_{k}} \|\partial_{t} u_{h}(s)\|_{\mathrm{L}^{2}}^{2} \, ds + \frac{(\Delta t)^{2} \beta^{2} \gamma^{2}}{\nu} \int_{t_{k-1}}^{t_{k}} \|\partial_{t} u_{h}(s)\|_{\mathrm{L}^{2}}^{2} \, ds \\ &+ \Delta t \left( C(\alpha, \nu) \left( \|u_{h}(t_{k})\|_{\mathrm{L}^{2(\delta+1)}}^{\frac{4\delta(\delta+1)}{2}} + \|u_{h}^{k}\|_{\mathrm{L}^{2(\delta+1)}}^{\frac{4\delta(\delta+1)}{2}} \right) + \frac{\beta}{2} 2^{2\delta} (1+\gamma)^{2} (\delta+1)^{2} + \frac{3\nu}{8} \right) \|u_{h}(t_{k}) - u_{h}^{k}\|_{\mathrm{L}^{2}}^{2} \\ &+ \frac{(\Delta t)^{2}}{\nu} \left( 2\alpha^{2} + \beta^{2} (1+\gamma)^{2} (\delta+1)^{2} \right) \int_{t_{k-1}}^{t_{k}} \|u_{h}^{\delta}(s) \partial_{t} u_{h}(s)\|_{\mathrm{L}^{2}}^{2} \, ds + \sum_{j=1}^{k} \frac{2\eta^{2} T (\Delta t)^{3}}{\nu} \overline{K}_{kj}^{2} \int_{t_{j-1}}^{t_{j}} \|\partial_{t} \nabla u_{h}(\tau)\|_{\mathrm{L}^{2}}^{2} \, d\tau \end{split}$$

Summing over all k = 1, 2, ..., N, and using the positivity of the kernel (cf. (3.22)), we attain

$$\begin{aligned} \|u_{h}(t_{N}) - u_{h}^{N}\|_{L^{2}}^{2} + \sum_{k=1}^{N} \nu \Delta t \|\nabla(u_{h}(t_{k}) - u_{h}^{k})\|_{L^{2}}^{2} \\ \leqslant \frac{4\alpha^{2}(\Delta t)^{2}}{\nu} \int_{0}^{T} \|\partial_{t} \nabla u_{h}(s)\|_{L^{2}}^{2} ds + \frac{2(\Delta t)^{2}\beta^{2}\gamma^{2}}{\nu} \int_{0}^{T} \|\partial_{t} u_{h}(s)\|_{L^{2}}^{2} ds \end{aligned}$$

GENERALIZED BURGERS-HUXLEY EQUATION WITH WEAKLY SINGULAR KERNEL 37 of 45

$$+2\Delta t \sum_{k=1}^{N} \left( C(\alpha, \mathbf{v}) \left( \left\| u_{h}(t_{k}) \right\|_{\mathbf{L}^{2}(\delta+1)}^{\frac{4\delta(\delta+1)}{(2-d)\delta+2}} + \left\| u_{h}^{k} \right\|_{\mathbf{L}^{2}(\delta+1)}^{\frac{4\delta(\delta+1)}{(2-d)\delta+2}} \right) + \frac{\beta}{2} 2^{2\delta} (1+\gamma)^{2} (\delta+1)^{2} + \frac{3\nu}{8} \right) \left\| u_{h}(t_{k}) - u_{h}^{k} \right\|_{\mathbf{L}^{2}}^{2} + \frac{2(\Delta t)^{2}}{\nu} \left( 2\alpha^{2} + C\beta^{2} (1+\gamma)^{2} (\delta+1)^{2} \right) \int_{0}^{T} \left\| u_{h}(s)^{\delta} \partial_{t} u_{h}(s) \right\|_{\mathbf{L}^{2}}^{2} \, \mathrm{d}s + \frac{4\eta^{2} T(\Delta t)^{3}}{\nu} \sum_{k=1}^{N} \sum_{j=1}^{k} \overline{K}_{kj}^{2} \int_{t_{j-1}}^{t_{j}} \left\| \partial_{t} \nabla u_{h}(\tau) \right\|_{\mathbf{L}^{2}}^{2} \, \mathrm{d}\tau.$$

$$(3.29)$$

For the memory term, we get

$$T(\Delta t)^{3} \sum_{k=1}^{N} \sum_{j=1}^{k} \overline{K}_{kj}^{2} \int_{t_{j-1}}^{t_{j}} \|\partial_{t} \nabla u_{h}(\tau)\|_{L^{2}}^{2} d\tau \leq \sup_{k,j} \overline{K}_{kj}^{2} T(\Delta t)^{3} \sum_{k=1}^{N} \sum_{j=1}^{k} \int_{t_{j-1}}^{t_{j}} \|\partial_{t} \nabla u_{h}(\tau)\|_{L^{2}}^{2} d\tau$$
$$\leq \sup_{k,j} \overline{K}_{kj}^{2} T^{2} (\Delta t)^{2} \int_{0}^{T} \|\partial_{t} \nabla u_{h}(\tau)\|_{L^{2}}^{2} d\tau.$$

Using the above estimate and an application of Gronwall inequality in (3.29) yields

$$\begin{aligned} \|u_{h}(t_{N}) - u_{h}^{N}\|_{L^{2}}^{2} + \sum_{k=1}^{N} \nu \Delta t \|\nabla(u_{h}(t_{k}) - u_{h}^{k})\|_{L^{2}}^{2} \\ &\leq C(\Delta t)^{2} \left( \int_{0}^{T} \|\partial_{t} \nabla u_{h}(s)\|_{L^{2}}^{2} \, \mathrm{d}s + \int_{0}^{T} \|u_{h}(s)^{\delta} \partial_{t} u_{h}(s)\|_{L^{2}}^{2} + \frac{4\eta^{2}T^{2}}{\nu} \sup_{k,j} \overline{K}_{kj}^{2} \int_{0}^{T} \|\partial_{t} u_{h}(s)\|_{L^{2}}^{2} \, \mathrm{d}s \right) \\ &\times \exp\left\{ 2\Delta t \left[ C(\alpha, \nu) \left( \|u_{h}(t_{k})\|_{L^{2(\delta+1)}}^{\frac{4\delta(\delta+1)}{(2-d)\delta+2}} + \|u_{h}^{k}\|_{L^{2(\delta+1)}}^{\frac{4\delta(\delta+1)}{(2-d)\delta+2}} \right) + \frac{\beta}{2} 2^{2\delta} (1+\gamma)^{2} (\delta+1)^{2} + \frac{3\nu}{8} \right] \right\}. \end{aligned}$$
(3.30)

One can prove the estimates obtained in (2.39) for the semi-discrete solution  $u_h$  also, so that the right hand side of (3.30) is bounded (see Remark 3.2 below).

**Step 2:** *Estimate for any*  $t \in [t_{k-1}, t_k]$ . Let us now introduce a linear interpolation  $\mathscr{I}u_h$  for the semidiscrete solution  $u_h$  [Yi & Guo (2015) (Section 3.1)] by:

$$\mathscr{I}u_h(t) = u_h(t_{k-1}) + \left(\frac{t - t_{k-1}}{\Delta t}\right) (u_h(t_k) - u_h(t_{k-1})) \quad \text{for } t \in [t_{k-1}, t_k].$$

Let us write  $u_h - u_{kh} = u_h - \mathscr{I}u_h + \mathscr{I}u_h - u_{kh}$ , where  $\mathscr{I}u_h$  is the linear interpolation as defined above. Using the triangle inequality, we have

$$\|u_h - u_{kh}\|_{L^2(0,T;H^1_0(\Omega))}^2 \leq 2\|u_h - \mathscr{I}u_h\|_{L^2(0,T;H^1_0(\Omega))}^2 + 2\|\mathscr{I}u_h - u_{kh}\|_{L^2(0,T;H^1_0(\Omega))}^2$$

The first term can be estimated using (Lemma 3.2 [Yi & Guo (2015)]) and the final term using (3.30) as

$$\|u_{h} - \mathscr{I}u_{h}\|_{L^{2}(0,T;H_{0}^{1}(\Omega))}^{2} = \sum_{i=1}^{N} \int_{t_{i-1}}^{t_{i}} \|u_{h}(t) - \mathscr{I}u_{h}(t)\|_{H_{0}^{1}(\Omega)}^{2} dt \leq C(\Delta t)^{2} \int_{0}^{T} \|\partial_{t}u_{h}(t)\|_{H_{0}^{1}}^{2} dt,$$
  
$$\|\mathscr{I}u_{h} - u_{kh}\|_{L^{2}(0,T;H_{0}^{1}(\Omega))}^{2} = \sum_{i=1}^{N} \int_{t_{i-1}}^{t_{i}} \|\mathscr{I}u_{h}(t) - u_{kh}(t)\|_{H_{0}^{1}(\Omega)}^{2} dt \leq C \sum_{i=1}^{N} \Delta t \|u_{h}(t_{i}) - u_{h}^{i}\|_{H_{0}^{1}}^{2}.$$
(3.31)

Again using triangle inequality, we have

$$|u - u_{kh}||^{2}_{L^{\infty}(0,T;L^{2}(\Omega))} \leq 2||u_{h} - \mathscr{I}u_{h}||^{2}_{L^{\infty}(0,T;L^{2}(\Omega))} + 2||\mathscr{I}u_{h} - u_{kh}||^{2}_{L^{\infty}(0,T;L^{2}(\Omega))}.$$

Similarly as in (3.31), by using (Corollary 3.1 [Yi & Guo (2015)]) and the estimate (3.30), we deduce

$$\begin{split} \|u_{h} - \mathscr{I}u_{h}\|_{L^{\infty}(0,T;L^{2}(\Omega))}^{2} &\leq \sup_{1 \leq i \leq N} \left( \sup_{t_{i-1} \leq t \leq t_{i}} \|u_{h} - \mathscr{I}u_{h}\|_{L^{2}(\Omega)}^{2} \right) \leq C(\Delta t)^{2} \|u_{h}\|_{W^{1,\infty}(0,T;L^{2}(\Omega))}^{2}, \\ \|\mathscr{I}u_{h} - u_{kh}\|_{L^{\infty}(0,T;L^{2}(\Omega))}^{2} &\leq \sup_{1 \leq i \leq N} \left( \sup_{t_{i-1} \leq t \leq t_{i}} \|\mathscr{I}u_{h} - u_{kh}\|_{L^{2}(\Omega)}^{2} \right) \leq C \sup_{1 \leq i \leq N} \left( \|u_{h}(t_{i}) - u_{h}^{i}\|_{L^{2}(\Omega)}^{2} \right), \end{split}$$

and the required result follows.

The main result of this section is

THEOREM 3.3 Assume that  $u_0 \in H^2(\Omega) \cap H_0^1(\Omega)$  and  $f \in H^1(0,T;L^2(\Omega))$  such that  $u \in L^{\infty}(0,T;H^2(\Omega))$ ,  $u_t \in L^{\infty}(0,T;L^2(\Omega)) \cap L^2(0,T;H_0^1(\Omega))$  be the solution of (2.1) on the interval (0,T]. Then, the finite element approximation  $u_{kh}$  converges to u as  $\Delta t, h \to 0$ . In addition, there exists a constant  $C \ge 0$  such that the approximation  $u_{kh}$  satisfies the following error estimate:

$$\begin{aligned} \|u - u_{kh}\|_{L^{\infty}(0,T;L^{2}(\Omega))}^{2} + \|u - u_{kh}\|_{L^{2}(0,T;H_{0}^{1}(\Omega))}^{2} \\ &\leqslant \eta^{2}(\Delta t)^{2} \sup_{k,j} \overline{K}_{kj}^{2}(\|f\|_{H^{1}(0,T;L^{2}(\Omega))}^{2} + \|u_{0}\|_{H^{2}}^{2}) + C((\Delta t)^{2} + h^{2})(\|f\|_{H^{1}(0,T;L^{2}(\Omega))}^{2} + \|u_{0}\|_{H^{2}}^{2}). \end{aligned}$$

*Proof.* The proof is an application of triangle inequality, Theorem 3.2 and Lemma 3.3.

REMARK 3.2 1. In the literature most of the estimates needs higher regularity  $u_{tt} \in L^1(0,T;L^2(\Omega))$  but we have proved estimates under the minimal regularity assumptions.

- 2. For the generalised Burgers- Huxley equation without memory (that is,  $\eta = 0$ ), we have obtained the optimal rate of convergence. Also, if we assume that  $K(\cdot) \in L^{\infty}(0,T)$ , then  $\overline{K}_{kj}^2 \leq C$ , so that we obtain the optimal convergence again.
- 3. As proved in McLean *et al.* (1996) and McLean & Mustapha (2007), if we assume  $|K(t-s)| \le |t-s|^{-\alpha}$ , for  $0 < \alpha < 1$ , then we have  $\overline{K}_{kj}^2 \le O((\Delta t)^{-2\alpha})$  and in that case the error estimates are given as

$$\|u - u_{kh}\|_{L^{\infty}(0,T;L^{2}(\Omega))}^{2} + \|u - u_{kh}\|_{L^{2}(0,T;H_{0}^{1}(\Omega))}^{2} \leq C((\Delta t)^{2(1-\alpha)} + h^{2})(\|f\|_{H^{1}(0,T;L^{2}(\Omega))}^{2} + \|u_{0}\|_{H^{2}}^{2}).$$

For  $\alpha = \frac{1}{2}$ , we have sub-optimal convergence rate and as  $\alpha \to 0$ , the convergence tends to an optimal one.

- 4. If we assume  $K(\cdot) \in L^p(0,T)$ , then we have  $\overline{K}_{kj}^2 \leq (\Delta t)^{\frac{2-2q}{q}}$ , where  $\frac{1}{p} + \frac{1}{q} = 1$ . Therefore, the error estimates are of order  $((\Delta t)^{\frac{2}{q}} + h^2)$ . For p = q = 2, that is,  $K(\cdot) \in L^2(0,T)$ , then we have the error estimates are of order  $(\Delta t + h^2)$ .
- 5. If we use the time discretization with non uniform refinement as done in McLean *et al.* (1996) and McLean & Mustapha (2007), we can get the optimal convergence rate.

#### 4. Numerical studies

In this section, we perform numerical computations to verify the theoretical results obtained in Theorem 3.3. The results are obtained in the open-source finite element library FEniCS (Alnæs *et al.*, 2015; Langtangen & Logg, 2016).

The error tables shown below present the error estimates and order of convergence for the generalized Burgers-Huxley equation (GBHE) with kernel ( $\eta \neq 0$ ) and without Kernel ( $\eta = 0$ ) for the equation

$$\frac{\partial u}{\partial t} + \alpha u^{\delta} \sum_{i=1}^{d} \frac{\partial u}{\partial x_{i}} - \nu \Delta u - \eta \int_{0}^{t} K(t-s) \Delta u(s) \, \mathrm{d}s = \beta u (1-u^{\delta}) (u^{\delta} - \gamma) + f, \tag{4.1}$$

under the following norm

$$|||u^{k} - u_{h}^{k}||| = ||u_{h}(t_{N}) - u_{h}^{N}||_{L^{2}}^{2} + \sum_{k=1}^{N} \nu \Delta t ||\nabla(u_{h}(t_{k}) - u_{h}^{k})||_{L^{2}}^{2}.$$

In all the experiments, we are using a backward Euler discretization in time and FEM in space. Consider the temporal discretization  $t_k = k\Delta t$ , for a given M,  $\Delta t = \frac{T}{M}$ ,  $t_k = k\Delta t$  and h is the spatial discretization parameter. We choose  $\Delta t \propto h$  and the rate of convergence is given by  $r = \log(||u - u_h^1||/||u - u_h^2||)/\log(h^1/h^2)$ .

### 4.1 Accuracy verification for the smooth kernel

Consider the problem (4.1) defined on the domain  $(0, 1)^d$ . Table 1 represents the errors and convergence rates for the numerical solution  $u_h$  for the smooth kernel  $K(t) = e^{-\delta t}$ ,  $t \in [0, T]$  with the exact solution  $u = \prod_{i=1}^d e^{-t} \sin(\pi x_i)$ , where we have used positive quadrature rule for the memory term.

We choose the value of the parameters as follows:  $\alpha = \beta = \delta = v = 1$ , and  $\gamma = 0.5$ , the forcing term has been calculated using the closed-form solution. We are using first-order polynomials to approximate the numerical solution for all the stimulations. Table 1 contains the error history for a sequence of refined uniform meshes for the numerical solution constructed using CFEM. The convergence results for both the GBHE with and without memory are represented in the table for both 2D and 3D. In the table, we can observe that the error in the energy norm decreases with the mesh size at the rate O(h). We require at most three Newton iterations and three linear solver iterations to reach the prescribed tolerance of  $10^{-10}$ .

#### 4.2 Weakly singular kernel

Consider the problem (4.1) defined on the domain  $(0,1)^d$ . We shall approximate  $\psi$  in  $J_k(\psi) = \int_0^{t_n} K(t_n - s)\psi(s)ds$  by the piecewise constant function taking value  $\psi(t_j)$  in  $(t_{j-1}, t_j)$ . We have obtained the errors and convergence rates for the numerical solution  $u_h^k$  for the weakly singular kernel  $K(t) = \frac{1}{\sqrt{t}}$  with two different expressions of the exact solution

Case 1 : 
$$u = (t^3 - t^2 + 1) \prod_{i=1}^{d} \sin(\pi x_i)$$
, Case 2 :  $u = t \sqrt{t} \prod_{i=1}^{d} \sin(\pi x_i)$ .

The parameters are chosen as in Khan *et al.* (2021), that is,  $\alpha = \beta = \delta = v = 1$ , and  $\gamma = 0.5$ . Table 2 represents the convergence results related to the Case 1 and Table 3 corresponds to the Case 2. In both cases, we can observe that the error in the energy norm decreases with the mesh size at the rate O(h). We require at most three Newton iterations and three linear solver iterations to reach the prescribed tolerance of  $10^{-10}$ .

Table 1. Errors and convergence rates for the numerical solutions  $u_h^k$  for the smooth function  $K(t) = e^{-\delta t}$  and the given solution  $u = \prod_{i=1}^d e^{-t} \sin(\pi x_i)$ .

Error history in 2D								
CGFEM	Mesh	$\ \cdot\ $ -error for $\eta = 0$	O(h)	$\ \cdot\ $ -error for $\eta = 1$	O(h)			
	$4 \times 4$	7.21(-01)	_	6.77(-01)	-			
	$8 \times 8$	3.54(-01)	1.03	3.28(-01)	1.04			
	16×16	1.74(-01)	1.02	1.60(-01)	1.04			
	$32 \times 32$	8.62(-02)	1.01	7.90(-02)	1.02			
	$64 \times 64$	4.29(-02)	1.01	3.92(-02)	1.01			
	$128 \times 128$	2.14(-02)	1.00	1.95(-02)	1.00			
Error history in 3D								
CGFEM	Mesh	$\ \cdot\ $ -error for $\eta = 0$	O(h)	$\ \cdot\ $ -error for $\eta = 1$	O(h)			
	$2 \times 2 \times 2$	1.35(00)	_	1.33(00)	-			
	$4 \times 4 \times 4$	7.40(-01)	0.86	7.13(-01)	0.90			
	$8 \times 8 \times 8$	3.71(-01)	1.00	3.53(-01)	1.01			
	$16 \times 16 \times 16$	1.83(-01)	1.02	1.73(-01)	1.03			
	$32 \times 32 \times 32$	9.06(-02)	1.01	8.55(-02)	1.02			
	$64 \times 64 \times 64$	4.50(-02)	1.01	4.24(-02)	1.01			

Table 2. Errors and convergence rates for the numerical solutions  $u_h^k$  for the weakly singular kernel  $K(t) = \frac{1}{\sqrt{t}}$  and the given solution  $u = \prod_{i=1}^d (t^3 - t^2 + 1) \sin(\pi x_i)$ .

Error history in 2D							
CGFEM	Mesh	$\ \cdot\ $ -error for $\eta = 0$	O(h)	$\ \cdot\ $ -error for $\eta = 1$	O(h)		
	$4 \times 4$	8.04(-01)	_	8.05(-01)	_		
	$8 \times 8$	4.14(-01)	0.96	4.14(-01)	0.96		
	16×16	2.09(-01)	0.99	2.09(-01)	0.99		
	$32 \times 32$	1.05(-01)	0.99	1.05(-01)	0.99		
	$64 \times 64$	5.26(-02)	1.00	5.25(-02)	1.00		
	$128 \times 128$	2.63(-02)	1.00	2.62(-02)	1.00		
Error history in 3D							
CGFEM	Mesh	$\ \cdot\ $ -error for $\eta = 0$	O(h)	$\ \cdot\ $ -error for $\eta = 1$	O(h)		
	$2 \times 2 \times 2$	1.48(00)	—	1.50(00)	-		
	$4 \times 4 \times 4$	8.62(-01)	0.78	8.64(-01)	0.80		
	$8 \times 8 \times 8$	4.51(-01)	0.93	4.51(-01)	0.94		
	$16 \times 16 \times 16$	2.23(-01)	1.01	2.28(-01)	0.98		
	$32 \times 32 \times 32$	1.15(-01)	0.96	1.15(-01)	0.99		
	$64 \times 64 \times 64$	5.74(-02)	1.00	5.74(-02)	1.00		

Error history in 2D with weakly singular kernel, $u = t\sqrt{t} \sin(2\pi x) \sin(2\pi y)$							
CGFEM	Mesh	$\ \cdot\ $ -error for $\eta = 0$	<i>O</i> ( <i>h</i> )	$\ \cdot\ $ -error for $\eta = 1$	O(h)		
	$4 \times 4$	1.44(00)	_	1.47(00)	_		
	$8 \times 8$	9.24(-01)	0.64	9.35(-01)	0.65		
	$16 \times 16$	5.02(-01)	0.88	5.06(-01)	0.89		
	$32 \times 32$	2.59(-01)	0.95	2.60(-01)	0.96		
	$64 \times 64$	1.31(-01)	0.98	1.32(-01)	0.99		
	$128 \times 128$	6.59(-02)	0.99	6.61(-02)	1.00		
Error history in 3D with weakly singular kernel, $u = t\sqrt{t} \sin(2\pi x) \sin(2\pi y) \sin(2\pi z)$							
CGFEM	Mesh	$\ \cdot\ $ -error for $\eta = 0$	O(h)	$\ \cdot\ $ -error for $\eta = 1$	O(h)		
	$4 \times 4$	1.31(00)	—	1.32(00)	—		
	8×8	9.20(-01)	0.51	9.26(-01)	0.51		
	16×16	5.15(-01)	0.84	5.18(-01)	0.84		
	$32 \times 32$	2.68(-01)	0.94	2.61(-01)	0.99		
	$64 \times 64$	1.36(-01)	0.98	1.37(-01)	0.93		

Table 3. Errors and convergence rates for the numerical solutions  $u_h^k$  for the weakly singular kernel  $K(t) = \frac{1}{\sqrt{t}}$  and the given solution  $u = t\sqrt{t} \sin(2\pi x) \sin(2\pi y)$ .

# 4.3 Application to Nerve Pulse Propagation

We conclude this section by showing a difference of qualitative illustration between a classical bistable equation with a weakly singular kernel (without advection,  $\alpha = 0$  and with a simplified cubic nonlinearity induced by  $\delta = 1$ ) and the generalized Burgers-Huxley equation with a weakly singular kernel. In this study, we perform a straightforward simulation to a transient problem which incorporates an additional ordinary differential equation (ODE). This inclusion allows for the emergence of self-sustained patterns, as discussed in Bini *et al.* (2010). The system reads

$$\partial_t u + \alpha u^{\delta} \sum_{i=1}^d \partial_i u - v \Delta u - \eta \int_0^t K(t-s) \Delta u(s) \, \mathrm{d}s = \beta u (1-u^{\delta}) (u^{\delta} - \gamma) - v,$$
  
$$\partial_t v = \varepsilon (u - \rho v), \tag{4.2}$$

where  $\varepsilon$  and  $\rho$  are parameters that can change the rest state and dynamics. Setting  $\delta = 1$ ,  $\eta = 0$  and  $\alpha = 0$ , one recovers the well-known FitzHugh-Nagumo equations

 $\partial_t u - v \Delta u - \beta u (1-u)(u-\gamma) + v = 0, \qquad \partial_t v = \varepsilon (u-\rho v).$ 

To obtain the weak formulation similar to (3.21) for the CFEM, we apply a simple backward Euler time discretization with constant time step  $\Delta t = 0.2$  and other parameters similar to Khan *et al.* (2021). For this example, we prescribe Neumann boundary conditions for u on  $\partial \Omega$ . Figure 1 depicts three snapshots of the evolution of u (representing the action potential propagation in a piece of nerve tissue, cardiac muscle, or any excitable media) for the classical FitzHugh-Nagumo system with weakly singular kernel vs. the modified delayed generalized Burgers-Huxley system with weakly singular kernel (4.2). All numerical solutions computed using the CFEM setting. The difference in the spiral dynamics appears to be more responsive to the degree of non-linearity (governed by  $\delta$ ) rather than the additional advection term (influenced by  $\alpha$ ).

FIG. 1. Example 3. Snapshots at t = 200 of  $u_h$  with memory coefficient  $\eta = 0.0001, 0.01, 1$  respectively for the FitzHugh-Nagumo model using  $\delta = 1$ ,  $\alpha = 0$  (top panels) and for the modified generalized Burgers-Huxley system with  $\delta = 1$ ,  $\alpha = 0.1$  (middle row) and with  $\delta = 2$ ,  $\alpha = 0.1$  (bottom).



# 5. Acknowledgement

The first author would like to thank Ministry of Education, Government of India (Prime Minister Research Fellowship, PMRF ID : 2801816), for financial support to carry out his research work.

# References

- ABLOWITZ, M. J., FUCHSSTEINER, B. & KRUSKAL, M. (1987) Topics in soliton theory and exactly solvable nonlinear equations: proceedings of the conference on nonlinear evolution equations, solitons and the inverse scattering transform. World Scientific.
- ADOLFSSON, K., ENELUND, M. & LARSSON, S. (2003) Adaptive discretization of an integrodifferential equation with a weakly convolution kernel. *Comput. Methods Appl. Mech. Engrg.*, **192**, 5285–5304.

- ALNÆS, M., BLECHTA, J., HAKE, J., JOHANSSON, A., KEHLET, B., LOGG, A., RICHARDSON, C., RING, J., ROGNES, M. E. & WELLS, G. N. (2015) The fenics project version 1.5. Archive of Numerical Software, 3.
- BINI, D., CHERUBINI, C., FILIPPI, S., GIZZI, A. & RICCI, P. E. (2010) On spiral waves arising in natural systems. *Commun. Comput. Phys.*, **8**, 610–622.
- BURGERS, J. M. (1948) A mathematical model illustrating the theory of turbulence. Academic Press, Inc., New York, N. Y.
- CANNON, J. & LIN, Y. (1987) A priori  $L^2$  error estimates for galerkin method for nonlinear parabolic integro-differential equations. *Manuscript*.
- CANNON, J. & LIN, Y. (1988) Non-classical  $H^1$  projection and galerkin methods for non-linear parabolic integro-differential equations. *Calcolo*, **25**, 187–201.
- CHEN, C., THOMÉE, V. & WAHLBIN, L. B. (1992) Finite element approximation of a parabolic integro-differential equation with a weakly singular kernel. *Math. Comp.*, **58**, 587–602.
- CHRYSAFINOS, K. & HOU, L. S. (2002) Error estimates for semidiscrete finite element approximations of linear and semilinear parabolic equations under minimal regularity assumptions. *SIAM J. Numer. Anal.*, **40**, 282–306.
- DAUTRAY, R. & LIONS, J.-L. (1990) Mathematical analysis and numerical methods for science and technology. Vol. 4. Springer-Verlag, Berlin.
- EL-DANAF, T. S. (2007) Solitary wave solutions for the generalized Burgers-Huxley equation. *Int. J. Nonlinear Sci. Numer. Simul.*, **8**, 315–318.
- EVANS, L. C. (2010) Partial differential equations. American Mathematical Soc.
- GRISVARD, P. (1985) *Elliptic problems in nonsmooth domains*. Pitman (Advanced Publishing Program), Boston, MA.
- HASHIM, I., NOORANI, M. S. M. & SAID AL-HADIDI, M. R. (2006) Solving the generalized Burgers-Huxley equation using the Adomian decomposition method. *Math. Comput. Modelling*, **43**, 1404–1411.
- HOU, L. S. & ZHU, W. (2006) Error estimates under minimal regularity for single step finite element approximations of parabolic partial differential equations. *Int. J. Numer. Anal. Model.*, **3**, 504–524.
- KHAN, A., MOHAN, M. T. & RUIZ-BAIER, R. (2021) Conforming, nonconforming and DG methods for the stationary generalized Burgers-Huxley equation. J. Sci. Comput., 88, 1–26.
- KHATTAK, A. J. (2009) A computational meshless method for the generalized Burgers'-Huxley equation. *Appl. Math. Model.*, **33**, 3718–3729.
- KUMAR, A. & MOHAN, M. T. (2022) Large deviation principle for occupation measures of stochastic generalized burgers-huxley equation. *Journal of Theoretical Probability*, **36**, 1–49.

- KUMAR, B. V. R., SANGWAN, V., MURTHY, S. V. S. S. N. V. G. K. & NIGAM, M. (2011) A numerical study of singularly perturbed generalized Burgers-Huxley equation using three-step Taylor-Galerkin method. *Comput. Math. Appl.*, 62, 776–786.
- LANGTANGEN, H. P. & LOGG, A. (2016) Solving PDEs in Python. Springer, Cham.
- MAHAJAN, S. & KHAN, A. (Prep.) Finite element approximaton for the delayed generalised Burgers-Huxley equation with weakly singular kernels: Part II Non-conforming and DG approximaton (under prepartion).
- MCLEAN, W., THOMÉE, V. & WAHLBIN, L. B. (1996) Discretization with variable time steps of an evolution equation with a positive-type memory term. *J. Comput. Appl. Math.*, **69**, 49–69.
- MCLEAN, W. & MUSTAPHA, K. (2007) A second-order accurate numerical method for a fractional wave equation. *Numer. Math.*, **105**, 481–510.
- MCLEAN, W. & THOMÉE, V. (1993) Numerical solution of an evolution equation with a positive-type memory term. J. Austral. Math. Soc. Ser. B, 35, 23–70.
- MITROVIC, D. & ZUBRINIC, D. (1997) Fundamentals of applied functional analysis. CRC Press.
- MOHAN, M. T. (2020) On the three dimensional Kelvin-Voigt fluids: global solvability, exponential stability and exact controllability of Galerkin approximations. *Evol. Equ. Control Theory*, **9**, 301–339.
- MOHAN, M. T. & KHAN, A. (2021) On the generalized Burgers-Huxley equation: Existence, uniqueness, regularity, global attractors and numerical studies. *Discrete Contin. Dyn. Syst. Ser. B*, **26**, 3943– 3988.
- MOHAN, M. T. & SRITHARAN, S. S. (2019) Stochastic Navier-Stokes equations perturbed by Lévy noise with hereditary viscosity. *Infin. Dimens. Anal. Quantum Probab. Relat. Top.*, **22**, 1950006.
- MUSTAPHA, K. & MCLEAN, W. (2009) Discontinuous Galerkin method for an evolution equation with a memory term of positive type. *Math. Comp.*, **78**, 1975–1995.
- MUSTAPHA, K. & MUSTAPHA, H. (2010) A second-order accurate numerical method for a semilinear integro-differential equation with a weakly singular kernel. *IMA J. Numer. Anal.*, **30**, 555–578.
- OLDHAM, K. & SPANIER, J. (1974) The fractional calculus theory and applications of differentiation and integration to arbitrary order. Elsevier.
- PANI, A. K. & PETERSON, T. E. (1996) Finite element methods with numerical quadrature for parabolic integrodifferential equations. *SIAM J. Numer. Anal.*, **33**, 1084–1105.
- ROBINSON, J. C. & PIERRE, C. (2003) Infinite-dimensional dynamical systems: An introduction to dissipative parabolic pdes and the theory of global attractors. cambridge texts in applied mathematics. *Appl. Mech. Rev.*, 56, B54–B55.
- SARI, M., GURARSLAN, G. & ZEYTINOUGU, A. (2011) High-order finite difference schemes for numerical solutions of the generalized Burgers-Huxley equation. *Numer. Methods Partial Differential Equations*, 27, 1313–1326.

- SLOAN, I. H. & THOMÉE, V. (1986) Time discretization of an integro-differential equation of parabolic type. SIAM J. Numer. Anal., 23, 1052–1061.
- TEMAM, R. (2001) *Navier-Stokes equations: theory and numerical analysis*, vol. 343. American Mathematical Soc.
- THOMÉE, V. (2007) Galerkin finite element methods for parabolic problems. Springer Science & Business Media.
- THOMÉE, V. & ZHANG, N. Y. (1989) Error estimates for semidiscrete finite element methods for parabolic integro-differential equations. *Math. Comp.*, **53**, 121–139.
- WANG, X., ZHU, Z. & LU, Y. (1990) Solitary wave solutions of the generalised Burgers-Huxley equation. *Journal of Physics A: Mathematical and General*, 23, 271.
- WANG, X.-Y. (1985) Nerve propagation and wall in liquid crystals. Physics Letters A, 112, 402–406.
- WAZWAZ, A.-M. (2005) Travelling wave solutions of generalized forms of Burgers, Burgers–Kdv and Burgers–Huxley equations. *Applied Mathematics and Computation*, **169**, 639–656.
- WAZWAZ, A.-M. (2009) Burgers, fisher and related equations. Partial Differential Equations and Solitary Waves Theory. Springer, pp. 665–681.
- YANIK, E. G. & FAIRWEATHER, G. (1988) Finite element methods for parabolic and hyperbolic partial integro-differential equations. *Nonlinear Anal.*, **12**, 785–809.
- YI, L. & GUO, B. (2015) An *h-p* version of the continuous Petrov-Galerkin finite element method for Volterra integro-differential equations with smooth and nonsmooth kernels. *SIAM J. Numer. Anal.*, 53, 2677–2704.
- ZHANG, N. Y. (1993) On fully discrete Galerkin approximations for partial integro-differential equations of parabolic type. *Math. Comp.*, **60**, 133–166.
- ZHOU, J., XU, D. & DAI, X. (2019) Weak Galerkin finite element method for the parabolic integrodifferential equation with weakly singular kernel. *Comput. Appl. Math.*, **38**, Paper No. 38, 12.