Understanding the first measurement of $\mathcal{B}(B \to K \nu \bar{\nu})$

L. Allwicher,^{1,*} D. Bečirević,^{2,†} G. Piazza,^{2,‡} S. Rosauro-Alcaraz,^{2,§} and O. Sumensari^{2,¶}

¹Physik-Institut, Universität Zürich, CH-8057 Zürich, Switzerland

²IJCLab, Pôle Théorie (Bat. 210), CNRS/IN2P3 et Université, Paris-Saclay, 91405 Orsay, France

Recently, Belle II reported on the first measurement of $\mathcal{B}(B^{\pm} \to K^{\pm}\nu\bar{\nu})$ which appears to be almost 3σ larger than predicted in the Standard Model. We point out the important correlation with $\mathcal{B}(B \to K^*\nu\bar{\nu})$ so that the measurement of that decay mode could help restrain the possible options for building the model of New Physics. We interpret this new experimental result in terms of physics beyond the Standard Model by using SMEFT and find that a scenario with coupling only to τ can accommodate the current experimental constraints but fails in getting a desired $R_{D^{(*)}}^{exp}/R_{D^{(*)}}^{SM}$, unless one turns the other SMEFT operators that are not related to $b \to s\ell\ell$ or/and $b \to s\nu\nu$.

I. INTRODUCTION

Experimental studies of the angular distribution of $B \to K^*(\to K\pi)\mu\mu$ [1, 2] and $B \to K\mu\mu$ [3, 4] offered a number of observables, the study of which could help unveiling the effects of presence of New Physics (NP), i.e. physics beyond the Standard Model (BSM). It appeared, however, that interpreting several apparent deviations with respect to the Standard Model (SM) predictions required a very good control over the hadronic uncertainties, and in particular those related to the lowenergy operators coupling to $\bar{c}\gamma_{\mu}c$. Those latter contributions are particularly problematic in the regions of $q^2 = (p_{\mu^+} + p_{\mu^-})^2$ populated by the $c\bar{c}$ resonances. Clearly, in order to resolve the effects of BSM physics from those related to the SM weak interaction in these regions, one needs to evaluate the relevant hadronic matrix element of a non-local operator, which cannot be done vet in lattice QCD. Instead, one resorts to using various assumptions such as the guark-hadron duality to treat the problem perturbatively (even though the energy window is narrow) [5–7], or a specific hadronic model [8– 11]. That problem is commonly circumvented when measuring $R_{K^{(*)}} = \mathcal{B}'(B \to K^{(*)}\mu\mu)/\mathcal{B}'(B \to K^{(*)}ee)$ [12], where \mathcal{B}' is used to indicate that the partial branching fractions are measured in the interval $q^2 \in [1.1, 6] \text{ GeV}^2$ well below the first $c\bar{c}$ resonance, $m_{J/\psi}^2 = (3.097 \,\text{GeV})^2$. First measurements of R_K and R_{K^*} , as well as of $\mathcal{B}(B_s \to \mu\mu)$ [13–16] indicated an important departure from the SM predictions. Many models used to accommodate these deviations were used to constrain the BSM couplings, most of which also implied a significant deviation of $\mathcal{B}(B \to K^{(*)}\nu\bar{\nu})$ from its SM value. Very recently, however, LHCb reported $R_K = 0.998(90)$, $R_{K^*} = 0.930(97)$ [17], thus fully consistent with $R_{K^{(*)}}^{SM} =$

1.00(1) [18].

Even though it is experimentally much more challenging, $B \to K^{(*)} \nu \bar{\nu}$ is theoretically cleaner than the equivalent mode with charged leptons instead of neutrinos in the final state. This is so because the coupling to the problematic operators involving $c\bar{c}$ resonances is absent. The remaining non-perturbative QCD obstacle, that both decay modes share, is a reliable theoretical estimate of the hadronic matrix elements of the local operators in the entire physical region, $0 \leq q^2 \leq$ $(m_B - m_{K^{(*)}})^2$. That task is also very challenging for lattice QCD because the available q^2 range is too large. This is why the lattice QCD results are used to constrain the parameters entering a model q^2 -dependence of the relevant form factor, necessary for the SM prediction of $\mathcal{B}(B \to K \nu \bar{\nu})$. Interestingly, in the case of $B \to K \nu \bar{\nu}$ the q^2 shape of the hadronic form factor can be checked experimentally by measuring the partial branching fractions $\mathcal{B}'(B \to K \nu \bar{\nu})$, as discussed in Ref. [19].

In the following we will often use the ratio $R_{\nu\nu}^{K^{(*)}} = \mathcal{B}(B \to K^{(*)}\nu\bar{\nu})/\mathcal{B}(B \to K^{(*)}\nu\bar{\nu})^{\text{SM}}$, for which the experimental upper bounds exist [20]:

$$R_{\mu\nu}^K < 3.6 \quad (90\% \text{ C.L.}),$$
 (1)

$$R_{\nu\nu}^{K^*} < 2.7 \quad (90\% \text{ C.L.}).$$
 (2)

Very recently, the first of these bounds was superseded by the first measurement of this decay mode by Belle II, $\mathcal{B}(B^{\pm} \to K^{\pm} \nu \bar{\nu}) = 2.40(67) \times 10^{-5}$ [21], which appears to be 2.9 σ larger than its SM estimate.¹ This then leads to

$$R_{\nu\nu}^{K} = 5.4 \pm 1.5 \,. \tag{3}$$

In this letter we discuss how this departure from the SM prediction can be interpreted in terms of generic BSM

^{*} lukall@physik.uzh.ch

[†] damir.becirevic@ijclab.in2p3.fr

 $^{^{\}ddagger}$ gioacchino.piazza@ijclab.in2p3.fr

 $^{^{\$}}$ salvador.rosauro@ijclab.in2p3.fr

[¶] olcyr.sumensari@ijclab.in2p3.fr

¹ Note that for the SM estimate we take $\mathcal{B}(B^{\pm} \to K^{\pm}\nu\bar{\nu}) = 4.44(14)(27) \times 10^{-5}[19]$ i.e. the value which does not include the tree level contribution [22] which was also subtracted away during the experimental data analysis [21].

scenarios in the SMEFT framework, which then rec one to remain consistent with the stringent bound NP in $b \rightarrow s$ transitions arising from the measured . mentioned above, as well as from the measured $\mathcal{B}($ $\mu\mu$) = 3.35(27) × 10⁻⁸ [16], which is consistent wit SM estimate $\mathcal{B}(B_s \to \mu\mu)^{\text{SM}} = 3.66(4) \times 10^{-8}$ [23].

II. EFFECTIVE THEORY CONSIDERATIC

The low energy effective theory relevant to the b – decay is described by the effective Hamiltonian,

$$\mathcal{L}_{\text{eff}}^{\text{b} \to \text{s}\nu\nu} = \frac{4G_F}{\sqrt{2}} \lambda_t \sum_a C_a \mathcal{O}_a + \text{h.c.} \,,$$

with

$$\mathcal{O}_{L}^{\nu_{i}\nu_{j}} = \frac{e^{2}}{(4\pi)^{2}} (\bar{s}_{L}\gamma_{\mu}b_{L}) (\bar{\nu}_{i}\gamma^{\mu}(1-\gamma_{5})\nu_{j}) ,$$

$$\mathcal{O}_{R}^{\nu_{i}\nu_{j}} = \frac{e^{2}}{(4\pi)^{2}} (\bar{s}_{R}\gamma_{\mu}b_{R}) (\bar{\nu}_{i}\gamma^{\mu}(1-\gamma_{5})\nu_{j}) ,$$

in a standard notation with $\lambda_t = V_{tb}V_{ts}^*$, the suitable combination of the Cabibbo-Kobayashi-Maskawa (CKM) matrix elements. In the SM we know that $C_R^{\text{SM}} = 0$ and $\left[C_L^{\nu_i\nu_j}\right]_{\text{SM}} \equiv \delta_{ij} C_L^{\text{SM}}$, with $C_L^{\text{SM}} = -6.32(7)$ [24–28]. A detailed discussion of hadronic uncertainties and those arising from the choice of λ_t can be found in Ref. [19]. Here we will just mention that in the SM the values of decay rates of the charged *B*-meson read:

$$\mathcal{B}(B^{\pm} \to K^{\pm} \nu \nu) = (4.44 \pm 0.30) \times 10^{-6} ,$$

$$\mathcal{B}(B^{\pm} \to K^{\pm *} \nu \nu) = (9.8 \pm 1.4) \times 10^{-6} , \qquad (6)$$

and they do not comprise the triply Cabibbo-suppressed tree-level contribution, $B^+ \to \tau^+ (\to K^+ \bar{\nu}) \nu$ (also subtracted away in the experimental data analysis [21]). It is convenient to factor out the SM contribution to the branching fractions and write

$$\mathcal{B}(B \to K^{(*)}\nu\nu) = \mathcal{B}(B \to K^{(*)}\nu\nu)\Big|_{\mathrm{SM}} \left(1 + \delta \mathcal{B}_{K^{(*)}}^{\nu\nu}\right),$$
(7)

so that the NP piece, $\delta \mathcal{B}_{K^{(*)}}^{\nu \bar{\nu}}$, can be expressed in terms of $\delta C_{L,R}^{\nu_i \nu_j}$, the BSM contribution to the left- or right-handed operator given in Eq. (5). It is then straightforward to express $R_{\nu\nu}^{K^{(*)}} = 1 + \delta \mathcal{B}_{K^{(*)}}^{\nu\bar{\nu}}$. If we write $C_{L,R}^{\nu_i\nu_j} = \delta_{ij}C_{L,R}^{\mathrm{SM}} +$ $\delta C_{L,R}^{\nu_i\nu_j}$, then we have [19, 24]:

$$\delta \mathcal{B}_{K^{(*)}}^{\nu\bar{\nu}} = \sum_{i} \frac{2 \text{Re}[C_{L}^{\text{SM}} \left(\delta C_{L}^{\nu_{i}\nu_{i}} + \delta C_{R}^{\nu_{i}\nu_{i}}\right)]}{3|C_{L}^{\text{SM}}|^{2}} \\ + \sum_{i,j} \frac{|\delta C_{L}^{\nu_{i}\nu_{j}} + \delta C_{R}^{\nu_{i}\nu_{j}}|^{2}}{3|C_{L}^{\text{SM}}|^{2}} \\ - \eta_{K^{(*)}} \sum_{i,j} \frac{\text{Re}[\delta C_{R}^{\nu_{i}\nu_{j}} (C_{L}^{\text{SM}} \delta_{ij} + \delta C_{L}^{\nu_{i}\nu_{j}})]}{3|C_{L}^{\text{SM}}|^{2}} , \qquad (8)$$



FIG. 1. The correlation between $\mathcal{B}(B \to K^* \nu \nu)$ and $\mathcal{B}(B \to K^* \nu \nu)$ $K\nu\nu$) decays with respect to the variation of δC_L or δC_R . The shaded gray area correspond to 1σ and 2σ of the recent Belle II result for $\mathcal{B}(B \to K\nu\nu)$. The red point corresponds to the SM predictions for these observables. We also show the region of experimentally excluded $\mathcal{B}(B \to K^* \nu \nu)$ values [20] (gray hatched area), as well as the region which is not accessible within the EFT approach (purple hatched area), cf. Eq. (9)

where the sum over neutrino flavor indices is understood, $i, j \in \{1, 2, 3\}$. In the above expression, $\eta_K = 0$, and $\eta_{K^*} = 3.33(7)$. In that way, one can easily check the response of $\mathcal{B}(B \to K^* \nu \nu)$ to the new experimentally established $\mathcal{B}(B \to K\nu\nu)$, which is shown in Fig. 1.

Note, in particular, that we find the following relation between $B \to K^* \nu \bar{\nu}$ and $B \to K \nu \bar{\nu}$,

$$\frac{\mathcal{B}(B \to K^* \nu \bar{\nu})}{\mathcal{B}(B \to K \nu \bar{\nu})} \ge \frac{\mathcal{B}(B \to K^* \nu \bar{\nu})}{\mathcal{B}(B \to K \nu \bar{\nu})} \bigg|_{\rm SM} \left(1 - \frac{\eta_{K^*}}{4}\right), \quad (9)$$

which is depicted by the hatched-dark blue region la-

belled as "EFT" in Fig. 1 (see also [29]). By switching on either $\delta C_L^{\nu_i\nu_j}$ or $\delta C_R^{\nu_i\nu_j}$ and by considering all three diagonal and universal couplings to neutrino flavors, such that $\delta C_{L(R)}^{\nu_i\nu_j} = \delta_{ij}\delta C_{L(R)}$, we obtain that accommodating the measured $\mathcal{B}(B \to K \nu \nu)$ to 2σ results in:

$$\delta C_{L,R} \in [-12.0, -3.5] \cup [16.1, 24.7]. \tag{10}$$

These solutions become strongly restricted after imposing the experimental bound $\mathcal{B}(B \to K^* \nu \nu)^{\exp} < 2.7 \times$ 10^{-5} [20]. In particular, the large positive values of δC_R are discarded, and the above ranges become:

$$\delta C_L \neq 0 : \delta C_L \in [-4.2, -3.5] \cup [16.9, 17.3],$$

$$R_{\nu\nu}^{K^{*}} \in (2.4, 2.7),$$

$$\delta C_{R} \neq 0 : \delta C_{R} \in [-12.0, -3.5],$$

$$R_{\nu\nu}^{K^{*}} \in (0.6, 2.1), \qquad (11)$$

where we also give the resulting $\mathcal{B}(B \to K^* \nu \nu)$ compatible with $\mathcal{B}(B \to K \nu \nu)^{\exp}$ to 2σ and the experimental bound on $\mathcal{B}(B \to K^* \nu \nu)^{\exp}$. These allowed ranges are highlighted in Fig. 1 with darker hues. At this stage we consider the BSM contributions to be neutrino flavor universal, which is a relevant information when interpreting the values for $\delta C_{L,R}$. However, the correlation shown in Fig. 1 and the ranges of $R_{\nu\nu}^{K^*}$ given in Eq. (11) remain as such regardless of whether we assume the NP to be lepton flavor universal or not.

One can go a step further and predict the behavior of a fraction of the $B \to K^* \nu \nu$ corresponding to a specific polarization state of the outgoing K^* . For example, one can check how F_L , the fraction of decay rate corresponding to the longitudinally polarized K^* (cf. Refs. [24, 30]), responds to a non-zero δC_L or δC_R . $\mathcal{R}_{F_L} = F_L/F_L^{\text{SM}}$ is presented in Fig. 2 to better emphasize the fact that F_L remains insensitive to $\delta C_L \neq 0$, whereas it becomes drastically depleted with respect to the SM value, $F_L^{\text{SM}} = 0.49(7)$, when $\delta C_R \neq 0$ is chosen such that it is consistent with both $\mathcal{B}(B \to K \nu \nu)^{\exp}$ and the experimental bound on $\mathcal{B}(B \to K^* \nu \nu)^{\exp}$, corresponding to negative values of δC_R and the darker purple region in Fig. 2. In fact, we obtain that with $\delta C_R \neq 0$, the value of F_L gets more than 50% suppressed with respect to its SM value. From the plot in Fig. 2 we read off:

$$0 \le \mathcal{R}_{F_L} \le 0.44 \quad \Rightarrow \quad F_L \in [0, 0.21] \,. \tag{12}$$

This is a clear prediction that could be tested experimentally. Note, in particular, that F_L has a much smaller theoretical uncertainty in the SM than the $\mathcal{B}(B \to K^* \nu \bar{\nu})$, since most of the form factor uncertainties cancel out in the ratio.

III. SMEFT CONSIDERATIONS

When discussing the BSM scenarios in which the new degrees of freedom enter the stage well above the electroweak scale, it is convenient to work in the SM effective field theory (SMEFT), invariant under the full $SU(3)_c \times SU(2)_L \times U(1)_Y$ gauge symmetry [31]. This allows one to relate the $b \to s\nu\bar{\nu}$ and $b \to s\ell\ell$ via $SU(2)_L$ gauge symmetry [24, 32–36]. Of all the d = 6 operators in

$$\mathcal{L}_{\text{SMEFT}}^{(6)} \supset \sum_{i} \frac{\mathcal{C}_{i}}{\Lambda^{2}} \mathcal{O}_{i} ,$$
 (13)



FIG. 2. Similar to Fig. 1 we show how the fraction of $B \to K^* \nu \nu$ with longitudinally polarized K^* responds to the variation of δC_L or δC_R . Note that the yellow and purple curves correspond to those shown in Fig. 1, each associated with the allowed ranges of the Wilson coefficient given in Eq. 11.

we select those relevant to our study, namely,

$$\begin{split} \left[\mathcal{O}_{Hq}^{(1)}\right]_{kl} &= \left(\overline{Q}_{k}\gamma^{\mu}Q_{l}\right)\left(H^{\dagger}\overleftarrow{D}_{\mu}H\right), \\ \left[\mathcal{O}_{Hq}^{(3)}\right]_{kl} &= \left(\overline{Q}_{k}\tau^{I}\gamma_{\mu}Q_{l}\right)\left(H^{\dagger}\overleftarrow{D}_{\mu}\tau^{I}H\right), \\ \left[\mathcal{O}_{Hd}\right]_{kl} &= \left(\overline{d}_{kR}\gamma_{\mu}d_{lR}\right)\left(H^{\dagger}\overleftarrow{D}_{\mu}H\right), \\ \left[\mathcal{O}_{lq}^{(1)}\right]_{ijkl} &= \left(\overline{L}_{i}\gamma^{\mu}L_{j}\right)\left(\overline{Q}_{k}\gamma_{\mu}Q_{l}\right), \\ \left[\mathcal{O}_{lq}^{(3)}\right]_{ijkl} &= \left(\overline{L}_{i}\gamma^{\mu}\tau^{I}L_{j}\right)\left(\overline{Q}_{k}\tau^{I}\gamma_{\mu}Q_{l}\right), \\ \left[\mathcal{O}_{ld}\right]_{ijkl} &= \left(\overline{L}_{i}\gamma^{\mu}L_{j}\right)\left(\overline{d}_{kR}\gamma_{\mu}d_{lR}\right), \end{split}$$
(15)

where Q and L denote the quark and lepton $SU(2)_L$ doublet, respectively, while u, d, e stand for the quark and lepton weak singlets. In what follows, we will work in the flavor basis defined with diagonal down-type quark Yukawa matrix, i.e. with the CKM matrix element in the upper component of $Q_i = [(V^{\dagger} u)_i, d_i]^T$. Since we are focusing on the $b \to s$ processes we will fix k = 2 and l = 3, and discuss the two subsets of operators (14,15) separately.

A. Quark bilinears and Higgs

Firstly, we consider the operators with quark-bilinears and the Higgs current, as defined in Eq. (14). These contribute to $B \to K \nu \bar{\nu}$ via a tree-level induced Z-coupling to the $(\bar{s}\gamma^{\mu}b)$ current, as depicted in the right panel of Fig. 3. Clearly, these contributions would impact not only the $b \to s\nu_{\ell}\bar{\nu}_{\ell}$ transition, but also $b \to s\ell\ell$, with



FIG. 3. Tree-level contributions of four-fermion (left panel) and Higgs-fermion (right panel) operators to the $b \rightarrow s\nu\bar{\nu}$ transition in the SMEFT.

lepton-flavor universal contributions [24], e.g. the precisely determined $B_s \rightarrow \mu\mu$ branching ratio. Moreover, a double insertion of these coefficients will also have an effect in $B_s - \bar{B}_s$ mixing. Writing the Z-boson interaction Lagrangian with down-type quarks as

$$\mathcal{L}_{\text{eff}}^{Z} = -\frac{g}{\cos\theta_{W}} \sum_{i,j} \bar{d}_{i} \gamma^{\mu} \left(g_{d_{L}}^{Z\,ij} P_{L} + g_{d_{R}}^{Z\,ij} P_{R} \right) d_{j} Z_{\mu} \,, \quad (16)$$

where we find

L

$$\delta g_{d_L}^{Z\,ij} = -\frac{v^2}{2\Lambda^2} \left\{ \left[\mathcal{C}_{Hq}^{(1)} \right]_{ij} + \left[\mathcal{C}_{Hq}^{(3)} \right]_{ij} \right\} \,, \qquad (17)$$

$$\delta g_{d_R}^{Z\,ij} = -\frac{v^2}{2\Lambda^2} \left[\mathcal{C}_{Hd} \right]_{ij},\tag{18}$$

with $g_{\psi}^{Z\,ij} = \delta_{ij} g_{\psi}^{Z} + \delta g_{\psi}^{Z\,ij}$, $g_{d_L}^{Z} = -1/2 + 1/3 \sin^2 \theta_W$ and $g_{d_R}^{Z} = +1/3 \sin^2 \theta_W$, and θ_W the Weinberg angle. These relations can be used to match onto the relevant low-energy effective four-fermion operators, after integration of the SM vector bosons [37]. In particular, one finds, for example, that the contribution from NP to $B_s \to \mu\mu$ depends on the SMEFT operators as $\delta C_{10}^{\ell_i\ell_i} \propto [\mathcal{C}_{Hq}^{(1)}]_{23} + [\mathcal{C}_{Hq}^{(3)}]_{23} - [\mathcal{C}_{Hd}]_{23}$. Using the constraints from $\mathcal{B}(B_s \to \mu\mu)$ [16] and Δm_{B_s} [38, 39], we determine the 2σ confidence intervals for the coefficients $[\mathcal{C}_{Hq}^{(1)}]_{23}, [\mathcal{C}_{Hq}^{(3)}]_{23}$, and $[\mathcal{C}_{Hd}]_{23}$, switching on one at a time at $\Lambda = 1$ TeV and accounting for the renormalization group evolution from Λ down to $\mu = m_b$ [40]. The allowed intervals for the $B \to K\nu\bar{\nu}\bar{\nu}$ and $B \to K^*\nu\bar{\nu}$ branching fractions,

$$\begin{aligned} \mathcal{C}_{Hq}^{(1)} &\neq 0 : \mathcal{C}_{Hq}^{(1)} / \Lambda^2 \in [-0.86, 0.45] \times 10^{-3} \,\mathrm{TeV}^{-2}, \\ R_{\nu\nu}^K &\in (0.85, 1.08), R_{\nu\nu}^{K^*} \in (0.85, 1.08), \\ \mathcal{C}_{Hq}^{(3)} &\neq 0 : \mathcal{C}_{Hq}^{(3)} / \Lambda^2 \in [-0.78, 0.41] \times 10^{-3} \,\mathrm{TeV}^{-2}, \\ R_{\nu\nu}^K &\in (0.85, 1.08), R_{\nu\nu}^{K^*} \in (0.85, 1.08), \\ \mathcal{C}_{Hd} &\neq 0 : \mathcal{C}_{Hd} / \Lambda^2 \in [-0.45, 0.86] \times 10^{-3} \,\mathrm{TeV}^{-2}, \\ R_{\nu\nu}^K &\in (0.92, 1.16), R_{\nu\nu}^{K^*} \in (0.90, 1.05). \end{aligned}$$

We find that the $\mathcal{B}(B^{\pm} \to K^{\pm}\nu\bar{\nu})$ value can only be enhanced by $\approx 20\%$, which is largely insufficient to accommodate the deviation shown in Belle-II data.

B. Four-fermion operators

We turn now our attention to the four-fermion operators defined in Eq. (15). Let us first explicitly rewrite $\mathcal{L}_{\text{SMEFT}}^{(6)}$ using the operators given in Eq. (15). We have:

$$\mathcal{L}_{\text{SMEFT}}^{(6)} \supset \frac{1}{\Lambda^2} \left\{ \left(\mathcal{C}_{lq}^{(1)} + \mathcal{C}_{lq}^{(3)} \right)_{ij} (\bar{s}_L \gamma^\mu b_L) (\bar{e}_{Li} \gamma_\mu e_{Lj}) \right. \\ \left. + \left(\mathcal{C}_{lq}^{(1)} - \mathcal{C}_{lq}^{(3)} \right)_{ij} (\bar{s}_L \gamma^\mu b_L) (\bar{\nu}_{Li} \gamma_\mu \nu_{Lj}) \right. \\ \left. + 2 V_{cs} \left[\mathcal{C}_{lq}^{(3)} \right]_{ij} (\bar{c}_L \gamma^\mu b_L) (\bar{e}_{Li} \gamma_\mu \nu_{Lj}) \right. \\ \left. + \left[\mathcal{C}_{ld} \right]_{ij} (\bar{s}_R \gamma^\mu b_R) \left[(\bar{\nu}_{Li} \gamma_\mu \nu_{Lj}) + (\bar{e}_{Li} \gamma_\mu e_{Lj}) \right] + \text{h.c.} \right\},$$

$$(20)$$

where we have not written the charged-current operators that are CKM-suppressed. By comparing the above Lagrangian with the one given in Eq. (4) it is easy to identify:

$$\delta C_L^{\nu_i \nu_j} = \frac{\pi}{\alpha_{\rm em} \lambda_t} \frac{v^2}{\Lambda^2} \left\{ \left[\mathcal{C}_{lq}^{(1)} \right]_{ij} - \left[\mathcal{C}_{lq}^{(3)} \right]_{ij} \right\},$$

$$\delta C_R^{\nu_i \nu_j} = \frac{\pi}{\alpha_{\rm em} \lambda_t} \frac{v^2}{\Lambda^2} \left[\mathcal{C}_{ld} \right]_{ij}.$$
(21)

As it was discussed in Ref. [19], several simple scenarios can be distinguished if we switch one of the above SMEFT operators at the time. For example, by allowing only $C_{lq}^{(1)} \neq 0$ we get the simplest scenario with a Z' boson coupled only to the left-handed fermions. If, instead, the Z' is considered to be member of the weak triplet then one has only $C_{lq}^{(3)} \neq 0$. Quite peculiar is the case with $C_{lq}^{(3)} = C_{lq}^{(1)} \neq 0$, which corresponds to the BSM scenario often encountered in the literature and involves a singlet vector leptoquark state of hypercharge Y = 2/3. Similarly, a choice $C_{lq}^{(1)} = 3C_{lq}^{(3)}$ corresponds to the BSM model with a triplet of scalar leptoquarks with hypercharge Y = 1/3 [41].

IV. PHENOMENOLOGY

So far we did not touch on the issue of lepton flavor. In this Section we discuss simple BSM scenarios and check whether or not one can build a simple scenario compatible with experimental data, and most importantly with $\mathcal{B}(B \to K\nu\nu)^{\exp}$. In doing so we will separately treat the lepton flavor conserving case in which the couplings are diagonal (i = j), from the lepton flavor violating, i.e. with non-diagonal couplings $(i \neq j)$ being non-zero.

A. New Physics coupling to muons only

We first attempt attributing the discrepancy between the measured $\mathcal{B}(B \to K\nu\nu)$ and $\mathcal{B}(B \to K\nu\nu)^{\text{SM}}$ to



FIG. 4. Turning on the couplings to μ 's only, also labeled as i = 2, i.e. $\left[C_{l_q}^{(1,3)} \right]_{22} \equiv C_{l_q}^{(1,3)} \neq 0$ or $[C_{l_d}]_{22} \equiv C_{l_d} \neq 0$. Note that $C_{l_q}^{(3)}$ and C_{l_d} lead to the same prediction (purple curve in the plot) which cannot be reconciled with $\mathcal{B}(B \to K\nu\nu)^{\exp}$. Instead, by only allowing $C_{l_q}^{(1)} \neq 0$ (orange curve) one can get its small fraction to be consistent with both $\mathcal{B}(B \to K\nu\nu)^{\exp}$ and $\mathcal{B}(B_s \to \mu\mu)^{\exp}$ to 2σ , which is also highlighted in orange in the plot. The red point corresponds to the SM values (i.e. $C_{l_q}^{(1)} = C_{l_q}^{(3)} = C_{l_d} = 0$).

the couplings to muon only, i.e. i = j = 2. The best suited quantity to check the validity of such a solution is $\mathcal{B}(B_s \to \mu\mu)$ for which we need to include the modification of the Wilson coefficient $C_{10} = C_{10}^{\text{SM}} + \delta C_{10}$, that in terms of $\mathcal{C}_{la}^{(1,3)}$ and \mathcal{C}_{ld} writes:

$$\delta C_{10}^{\ell_i \ell_i} = \frac{\pi}{\alpha_{\rm em} \lambda_t} \frac{v^2}{\Lambda^2} \left\{ \left[\mathcal{C}_{ld} \right]_{ii} - \left[\mathcal{C}_{lq}^{(1)} \right]_{ii} - \left[\mathcal{C}_{lq}^{(3)} \right]_{ii} \right\}, \quad (22)$$

In Fig. 4 we show how $\mathcal{B}(B_s \to \mu\mu)$ varies when switching $\mathcal{C}_{lq}^{(1,3)}$ or \mathcal{C}_{ld} one at the time. Clearly, any $\mathcal{C}_{lq}^{(3)}$ or \mathcal{C}_{ld} cannot alone enhance $\mathcal{B}(B \to K\nu\nu)^{\text{SM}}$ in order to be consistent with experiment. The situation improves when $\mathcal{C}_{lq}^{(1)} \neq 0$ in which case one can reach the 2σ edge of $\mathcal{B}(B \to K\nu\nu)^{\exp}$, while remaining consistent with $\mathcal{B}(B_s \to \mu\mu)^{\exp}$ to 2σ as well. We find that the resulting allowed $\mathcal{C}_{lq}^{(1)}/\Lambda^2 \in [0.0129, 0.0131] \text{ TeV}^{-2}$, corresponding to the highlighted darker orange band in Fig. 5, and which translates into a very stringent bound: $2.16 \leq \mathcal{B}(B \to K^*\nu\nu) \times 10^5 \leq 2.19$. This scenario, however, seems quite unlikely even though it would be consistent with the recent LHCb results regarding the lepton flavor universality $R_{K^{(*)}}$. In fact, consistency with $R_{K^{(*)}}$ would require $\mathcal{C}_{lq}^{(1)}/\Lambda^2 \in [0.0129, 0.0134] \text{ TeV}^{-2}$.



FIG. 5. Similar to Fig. 4, except that in this case we turn on all of the lepton species with $[\mathcal{C}]_{11} = [\mathcal{C}]_{22} = [\mathcal{C}]_{33}$ where \mathcal{C} stands for either $\mathcal{C}_{lq}^{(1)}$, $\mathcal{C}_{lq}^{(3)}$ or \mathcal{C}_{ld} .

B. New Physics coupling to two or three lepton species

With respect to the previous case, the situation considerably improves if we turn on the couplings to both electrons and muons, in such a way that $\left[\mathcal{C}_{lq}^{(1)}\right]_{11}$ = $\left[\mathcal{C}_{lq}^{(1)}\right]_{22} \equiv \mathcal{C}.$ Quite obviously, in that case $R_{K^{(*)}}$ remains at its SM value and thus consistent with recent experimental analyses at LHCb. Compatibility with $\mathcal{B}(B \to K \nu \nu)^{\exp}$ is so improved that even a small 1σ overlap with $\mathcal{B}(B \to K\nu\nu)^{\exp}$ can be reached. To go deep into the 1σ region of both $\mathcal{B}(B \to K\nu\nu)^{\exp}$ and $\mathcal{B}(B_s \to \mu\mu)$ one can take the flavor universal situation, and assume $\begin{bmatrix} \mathcal{C}_{lq}^{(1)} \end{bmatrix}_{11} = \begin{bmatrix} \mathcal{C}_{lq}^{(1)} \end{bmatrix}_{22} = \begin{bmatrix} \mathcal{C}_{lq}^{(1)} \end{bmatrix}_{33} \equiv \mathcal{C}, \text{ which is what we plot} \\ \text{ in Fig. 5. Like before, } R_{K^{(*)}} \text{ remains unchanged with re$ spect to its SM value, and overall consistency with the $b \rightarrow s$ data is achieved. However, as a result of the severe constraint arising from $\mathcal{B}(B_s \to \mu\mu)^{\exp}$, in both cases we obtain $\mathcal{C}/\Lambda^2 \in [0.012, 0.013] \text{ TeV}^{-2}$, which then results in either $R_{\nu\nu}^{K^*} \in (3.6, 3.9)_{e,\mu}$ or $R_{\nu\nu}^{K^*} \in (4.8, 5.3)_{e,\mu,\tau}$. Since both these intervals are larger than $R_{\nu\nu}^{K^* (exp)} < 2.7$, all three scenarios discussed so far cannot meet the experimental constraints.

Another important shortcoming of the two scenarios discussed here, is that the contribution to ratios $R_D^{(*)} = \mathcal{B}(B \to D^{(*)}\tau\bar{\nu})/\mathcal{B}(B \to D^{(*)}l\bar{\nu})$, where $l \in (e,\mu)$, is absent, since this needs to go through the triplet operator $C_{lq}^{(3)}$ (cf. Eq. (20)). We remind the reader that the most recent averages of experimental values for $R_{D^{(*)}}$ [42],

$$R_D = 0.257(29), \quad R_{D^*} = 0.284(12)$$
 (23)

are larger than predicted in the SM [43–45], leading to $R_D/R_D^{\text{SM}} = 1.19(10)$ and $R_{D^*}/R_{D^*}^{\text{SM}} = 1.15(5)$, which can

be averaged for our purpose to:

$$R_{D^{(*)}}^{\exp}/R_{D^*}^{SM} = 1.16(5).$$
 (24)

From Figure 4 one can clearly see that the constraints from $B_s \to \mu\mu$ cannot be respected in the case $\left[\mathcal{C}_{lq}^{(3)}\right]_{11} = \left[\mathcal{C}_{lq}^{(3)}\right]_{22} = \left[\mathcal{C}_{lq}^{(3)}\right]_{33}$. Moreover, this lepton-flavor universal scenario predicts $R_{D^{(*)}} = R_{D^{(*)}}^{\mathrm{SM}}$ which disagrees with Eq. (24). Also the case $\left[\mathcal{C}_{lq}^{(3)}\right]_{11} = \left[\mathcal{C}_{lq}^{(3)}\right]_{22}$ is in disagreement with data, since it predicts $R_{D^{(*)}} < R_{D^{(*)}}^{\mathrm{SM}}$ in correlation with an enhanced $\mathcal{B}(B \to K\nu\nu)$.

One concludes that modification of the couplings to τ 's is what is desired in order to be consistent with experimental data. More specifically one needs $\left[\mathcal{C}_{lq}^{(3)}\right]_{33} \neq 0$.

C. New Physics coupling to taus only

Finding $C_{lq}^{(3)} \equiv \left[C_{lq}^{(3)}\right]_{33} \neq 0$ such that $\mathcal{B}(B \to K\nu\nu)$ and $R_{D^{(*)}}$ are simultaneously consistent with their experimental values is possible as we avoid the constraint coming from $\mathcal{B}(B_s \to \mu\mu)^{\exp}$. We note that in this situation

$$\frac{R_{D^{(*)}}}{R_{D^{(*)}}^{\rm SM}} = \left(1 - \frac{v^2}{\Lambda^2} \frac{V_{cs}}{V_{cb}} \mathcal{C}_{lq}^{(3)}\right)^2, \qquad (25)$$

from which we see that $C_{lq}^{(3)} \neq 0$ is significantly enhanced by $V_{cs}/V_{cb} = 24(1)$. The solution is shown in Fig. 6. We find that the acceptable $C_{lq}^{(3)}$ falls in:

$$C_{lq}^{(3)} \in [-0.039, -0.019] \,\mathrm{TeV}^{-2}.$$
 (26)

As can be seen from (25), and in view of Eq. (24), only negative $C_{lq}^{(3)}$ values will lead to $R_{D^{(*)}}^{\exp}/R_{D^*}^{SM} > 1$, and we obtain $R_{D^{(*)}}/R_{D^*}^{SM} \in [1.06, 1.12]$, also highlighted in Fig. 6.

However, the $C_{lq}^{(3)}$ range allowed by $R_{\nu\nu}^{K^*(\exp)} < 2.7$ and $\mathcal{B}(B \to K\nu\nu)^{\exp}$ is $C_{lq}^{(3)}/\Lambda^2 \in [-0.015, -0.013] \cup$ $[0.033, 0.036] \text{ TeV}^{-2}$, incompatible with Eq. (26). Nonetheless, including other operators contributing to $b \to c\tau\nu$, but not to $b \to s\nu\bar{\nu}$, one can find scenarios allowing us to simultaneously explain the deviations in $R_K^{\nu\nu}$ and $R_{D^{(*)}}$. Example of such scenarios are $C_{lq}^{(1)} = C_{lq}^{(3)}$ [59], or combinations of scalar $C_{lequ}^{(1)}$ and tensor $C_{lequ}^{(3)}$, e.g. Ref. [60].

As it is well known, accommodating $R_{D^{(*)}}^{(\exp)}$ results in an increase of decay rates to a pair of τ -leptons in the final state [46]. Indeed we find that the scenarios compatible with $R_{D^{(*)}}^{(\exp)}$ and $\mathcal{B}(B \to K\nu\nu)^{(\exp)}$, and consistent with the $\mathcal{C}_{l_{0}}^{(a)}$ values given in Eq. (26) yield:



FIG. 6. The curve shows $R_{D^{(*)}}/R_{D^*}^{\text{SM}}$ as a function of $\mathcal{B}(B \to K \nu \nu)$ varying $[\mathcal{C}_{lq}^{(3)}]_{33}$ (purple) or $[\mathcal{C}_{ld}]_{33}$ (blue). The highlighted purple region correspond to $[\mathcal{C}_{lq}^{(3)}]_{33}$ allowed by $\mathcal{B}(B \to K \nu \nu)^{\text{exp}}$ which becomes restricted to the hatched region once the condition $R_{\nu\nu}^{K^*}(\text{exp}) < 2.7$ is imposed. That range is smaller than $R_{D^{(*)}}^{\text{exp}}/R_{D^*}^{\text{SM}}$ to 2σ (horizontal gray area). The red point corresponds to the SM.

$$\frac{\mathcal{B}(B \to K\tau\tau)}{\mathcal{B}(B \to K\tau\tau)^{\rm SM}} = \frac{\mathcal{B}(B \to K^*\tau\tau)}{\mathcal{B}(B \to K^*\tau\tau)^{\rm SM}} \in [15, 49]. \quad (27)$$

If we drop the requirement of compatibility with $R_{D^{(*)}}^{(\exp)}$ but instead insist on respecting the bound on $R_{\nu\nu}^{K^*(\exp)}$, we find the following values, ²

$$\frac{\mathcal{B}(B_s \to \tau\tau)}{\mathcal{B}(B_s \to \tau\tau)^{\mathrm{SM}}} \in [9, 10],$$

$$\frac{\mathcal{B}(B \to K\tau\tau)}{\mathcal{B}(B \to K\tau\tau)^{\mathrm{SM}}} = \frac{\mathcal{B}(B \to K^*\tau\tau)}{\mathcal{B}(B \to K^*\tau\tau)^{\mathrm{SM}}} \in [9, 11]. \quad (28)$$

Note, in particular, that we only consider $\mathcal{O}_{lq}^{(3)}$ in the predictions shown in Eq. (27)–(28). However, the connection between the $b \to s\nu_{\tau}\bar{\nu}_{\tau}$ and $b \to s\tau\tau$ transitions is more general, as the operators $\mathcal{O}_{lq}^{(1)}$ and \mathcal{O}_{ld} also contribute to both transitions. Therefore, if $\mathcal{B}(B \to K\nu\nu)$ is confirmed to be considerably larger than the SM value, one should expect a sizable deviation from the SM in $\mathcal{B}(B_s \to \tau\tau)$ and $\mathcal{B}(B \to K^{(*)}\tau\tau)$ as well. The only exception to this conclusion is the scenario where $\mathcal{C}_{lq}^{(1)} = -\mathcal{C}_{lq}^{(3)}$, which only affects neutral currents with neutrinos, cf. Eq. (20).

D. Lepton flavor violating case

We can now assume that the difference between $\mathcal{B}(B \to K\nu\nu)^{\exp}$ and its SM prediction can be described by

6

$$\frac{\mathcal{B}(B_s \to \tau \tau)}{\mathcal{B}(B_s \to \tau \tau)^{\rm SM}} \in [15, 46],$$

² Note that we neglect the range for which $R_{D^{(*)}}^{\exp}/R_{D^{*}}^{SM} < 1$.

turning on the off-diagonal couplings to lepton flavors $(i \neq j)$. This can be done with a number of various assumptions. Here, as an example, we turn either $[\mathcal{C}_{lq}^{(3)}]_{ij}$, or $[\mathcal{C}_{lq}]_{ij}$, or $[\mathcal{C}_{ld}]_{ij}$ at the time and assume the couplings to μ and τ to be non-zero by taking $\mathcal{C}_{32} = \mathcal{C}_{23}$. As a result we get that the 2σ compatibility with $\mathcal{B}(B \to K\nu\nu)^{\exp}$ is achieved for $\mathcal{C}_{32}/\Lambda^2 = \mathcal{C}_{23}/\Lambda^2 \in$ $[-0.034, -0.015] \cup [0.015, 0.034] \text{ TeV}^{-2}$, which, after imposing $R_{\nu\nu}^{K^*}(\exp) < 2.7$, reduces to $\mathcal{C}_{32}/\Lambda^2 = \mathcal{C}_{23}/\Lambda^2 \in$ $[-0.017, -0.015] \cup [0.015, 0.017] \text{ TeV}^{-2}$. As a consequence the decay rates of the corresponding lepton flavor violating modes will be significant. By using the expressions derived in Ref. [47] we obtain:

$$\mathcal{B}(B \to K\mu\tau) \in [2,3] \times 10^{-6},\tag{29}$$

not far from but consistent with the experimental bound [48, 49], $\mathcal{B}(B \to K \mu \tau)^{\exp} < 4.8 \times 10^{-5}$ at 90% C.L.

E. Expectations from concrete models

So far we have been agnostic about the possible origin of these effective operators, not focusing on any concrete model [41, 50, 51]. It is worth mentioning that a BSM model with a weak triplet of scalar leptoquarks S_3 cannot be made consistent with all the data discussed here. The reason is well known, namely it cannot give a significant increase to $R_{D^{(*)}}$ [50, 51]. The U_1 vector leptoquark, in a simplified model approach (involving couplings to the left-handed operators only), cannot be made consistent with data. This is so because at tree-level $C_{lq}^{(1)} = C_{lq}^{(3)}$, or $\delta C_L = 0$, which is excluded by the new Belle II result for $\mathcal{B}(B \to K \nu \nu)^{\text{exp}}$. It is important to stress, however, that the U_1 being a massive vector necessarily calls for a UV completion, which typically requires also the inclusion of additional fermionic degrees of freedom. The full theory of the U_1 has been extensively studied, and in particular it has been shown that the relation $C_{lq}^{(1)} = C_{lq}^{(3)}$ is broken at one-loop, leading to an increase of $\mathcal{B}(B \to K\nu\nu)$ of up to $\approx 50\%$ with respect to the Standard Model, given the current values of $R_{D^{(*)}}$ [52, 53]. Another possibility is to take the couplings to τ 's verifying $C_{lq}^{(1)} = -C_{lq}^{(3)}$, which is valid in a scenario with the so called S_1 lepto-quark. In that case one gets $R_{D^{(*)}}/R_{D^*}^{\text{SM}} \approx 1.02$, thus much smaller than in the case discussed in Sec. IVC. The way out would then be to turn on the right handed couplings [50]. Similarly, in the case of the so called R_2 leptoquarks one needs to turn on the right-handed couplings to quarks (and left-handed to τ), which in the SMEFT language means that $C_{ld} \neq 0$ [54]. It thus can explain the new experimental result $\mathcal{B}(B \to K\nu\nu)^{\exp}$. Moreover, by combining the constraints $\mathcal{B}(B \to K \nu \nu)^{\exp}$ and $[\Delta m_s]^{\text{exp}}/[\Delta m_s]^{\text{SM}}$ we get the bound $m_{\tilde{R}_2} \lesssim 3$ TeV and $m_{S_1} \lesssim 4.5$ TeV. This scenario, however, fails to ex-plain $R_{D^{(*)}}^{\text{exp}}/R_{D^*}^{\text{SM}}$. Another similar model is the one with a Z' boson coupled to left-handed τ and $\bar{s}\gamma_{\mu}P_{R}b$, again giving rise to $C_{ld} \neq 0$ [55]. While it can accommodate all of the $b \rightarrow s$ constraints, it does not contribute to $b \rightarrow c\tau\nu$.

F. Summary

In this letter, we discuss the possible consequences on $\mathcal{B}(B \to K^* \nu \nu)$ due to the recent Belle II measurement of $\mathcal{B}(B \to K \nu \nu)$ which was found to be nearly 3σ larger than predicted. We find that the values $\mathcal{B}(B \to K^* \nu \nu) \lesssim$ $1.8 \times \mathcal{B}(B \to K^* \nu \nu)^{\text{SM}}$ could only occur if the coupling to the right-handed operator $\bar{b}_R \gamma_\mu s_R$ is non-zero, while larger branching fractions would be consistent with couplings to either left-handed or right-handed operators (or a combination of both).

By relying on SMEFT we show that the increase in $\mathcal{B}(B \to K \nu \nu)$ cannot be described by switching on the couplings to one or more lepton flavors if that choice comprises the coupling to muons. The reason is that $\mathcal{B}(B_s \to \mu \mu)^{\text{exp}}$ provides a very stringent constraint on the values of the Wilson coefficient $C_{lq}^{(1)}$, which is the only one that can provide the desired enhancement to $\mathcal{B}(B \to K^{(*)}\nu\nu)$. In the SMEFT Lagrangian the Wilson coefficients relevant to $b \to s\ell\ell$ and $b \to s\nu\nu$ decay modes are also related to the semileptonic $b \to c\ell\nu$, so that one should make sure that the resulting $R_{D^{(*)}} > R_{D^*}^{\text{SM}}$, as suggested by experiment. For that it is necessary that $\mathcal{C}_{lg}^{(3)} \neq 0$ or to have the scalar and tensor operators that are not related to $b \to s\ell\ell$ and $b \to s\nu\nu$ [50]. We find that a significant increase in $R_{D^{(*)}}/R_{D^*}^{\rm SM}$ can be achieved if we allow only the coupling to τ and not to other species. In this way one can find the agreement with all of the abovementioned experimental constraints, including $R_{K^{(*)}}$, except that $R_{D^{(*)}}/R_{D^*}^{SM} \approx 1.05$.

We also considered a possibility with the off-diagonal couplings to lepton species. Accommodating $\mathcal{B}(B \rightarrow K\nu\nu)^{\exp}$ in this setting implies a large $\mathcal{B}(B \rightarrow K\mu\tau)$, not too far from the current experimental bound.

We need to emphasize that all the couplings discussed in this paper are well within the ranges allowed by the experimental data collected in the region of high- p_T tails of the differential distribution of $pp \rightarrow \ell \ell$ (+ soft jets), cf. Ref. [56].

One final comment regards the experimental value used in our analysis, which is the new measurement of $\mathcal{B}(B \to K\nu\nu)$ only. In [21], also an average with the previous bound from Belle-II was provided [20], giving $\mathcal{B}(B \to K\nu\nu)^{\exp} = 1.4(4) \times 10^{-5}$. How considering this average affects our conclusions can be mostly read off the same plots we presented, the main effect being a higher compatibility with the bound on $R_{\nu}^{K^*}$ in the left-handed scenario, and an increased (although still below the SM) range for F_L . Similarly, the τ -only coupling scenario would call for a larger right-handed contribution in order to enhance the contributions to $R_D^{(*)}$.

ACKNOWLEDGMENTS

This project has received funding /support from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 860881-HIDDeN and No 101086085-ASYMMETRY. The work of LA is supported by the Swiss National Science Foundation (SNF) under contract 200020-204428. LA is also grateful to the Mainz Institute for Theoretical Physics (MITP) of the Cluster of Excel-

- R. Aaij et al. [LHCb], JHEP 02, 104 (2016)
 [arXiv:1512.04442 [hep-ex]]; R. Aaij et al. [LHCb], Phys. Rev. Lett. 125, no.1, 011802 (2020) [arXiv:2003.04831
 [hep-ex]]; R. Aaij et al. [LHCb], Phys. Rev. Lett. 126, no.16, 161802 (2021) [arXiv:2012.13241 [hep-ex]].
- [2] A. M. Sirunyan *et al.* [CMS], JHEP **04** (2021), 124 [arXiv:2010.13968 [hep-ex]].
- [3] R. Aaij et al. [LHCb], JHEP 05, 082 (2014)
 [arXiv:1403.8045 [hep-ex]]; R. Aaij et al. [LHCb], Eur.
 Phys. J. C 77, no.3, 161 (2017) [arXiv:1612.06764 [hep-ex]].
- [4] A. M. Sirunyan *et al.* [CMS], Phys. Rev. D **98** (2018) no.11, 112011 [arXiv:1806.00636 [hep-ex]].
- [5] C. Greub, V. Pilipp and C. Schupbach, JHEP 12 (2008), 040 [arXiv:0810.4077 [hep-ph]]; H. H. Asatryan, H. M. Asatrian, C. Greub and M. Walker, Phys. Rev. D 66 (2002), 034009 [arXiv:hep-ph/0204341 [hep-ph]]; H. H. Asatryan, H. M. Asatrian, C. Greub and M. Walker, Phys. Rev. D 66 (2002), 034009 [arXiv:hep-ph/0204341 [hep-ph]].
- [6] H. H. Asatryan, H. M. Asatrian, C. Greub and M. Walker, Phys. Rev. D 65, 074004 (2002) [arXiv:hepph/0109140 [hep-ph]].
- [7] H. M. Asatrian, C. Greub and J. Virto, JHEP 04 (2020), 012 [arXiv:1912.09099 [hep-ph]].
- [8] A. Khodjamirian, T. Mannel, A. A. Pivovarov and Y. M. Wang, JHEP **09**, 089 (2010) [arXiv:1006.4945 [hepph]].
- [9] A. Khodjamirian, T. Mannel and Y. M. Wang, JHEP 02, 010 (2013) [arXiv:1211.0234 [hep-ph]].
- [10] N. Gubernari, D. van Dyk and J. Virto, JHEP 02, 088 (2021) [arXiv:2011.09813 [hep-ph]].
- [11] N. Gubernari, M. Reboud, D. van Dyk and J. Virto, JHEP 09 (2022), 133 [arXiv:2206.03797 [hep-ph]].
- [12] G. Hiller and F. Kruger, Phys. Rev. D 69, 074020 (2004) [arXiv:hep-ph/0310219 [hep-ph]].
- [13] M. Aaboud *et al.* [ATLAS], JHEP **04**, 098 (2019) [arXiv:1812.03017 [hep-ex]].
- [14] "Combination of the ATLAS, CMS and LHCb results on the $B_{(s)}^0 \rightarrow \mu^+ \mu^-$ decays," CMS-PAS-BPH-20-003.
- [15] R. Aaij et al. [LHCb], Phys. Rev. D 105, no.1, 012010 (2022) [arXiv:2108.09283 [hep-ex]].
- [16] A. Tumasyan *et al.* [CMS], Phys. Lett. B **842** (2023), 137955 [arXiv:2212.10311 [hep-ex]];
- [17] R. Aaij et al. [LHCb], [arXiv:2212.09152 [hep-ex]]; [arXiv:2212.09153 [hep-ex]].

lence PRISMA+ (Project ID 39083149), for its hospitality and its partial support during the completion of this work.

NOTE ADDED

While this paper was in writing the results of studies similar to our's were released in Refs. [29, 57]. In particular the authors of Ref. [29] interpret the new Belle II results in terms of the SMEFT operators reaching the conclusions which agree with ours for the most part.

- [18] M. Bordone, G. Isidori and A. Pattori, Eur. Phys. J. C 76 (2016) no.8, 440 [arXiv:1605.07633 [hep-ph]].
- [19] D. Bečirević, G. Piazza and O. Sumensari, Eur. Phys. J. C 83 (2023) no.3, 252 [arXiv:2301.06990 [hep-ph]].
- [20] J. Grygier *et al.* [Belle], Phys. Rev. D **96** (2017) no.9, 091101 [arXiv:1702.03224 [hep-ex]]; F. Abudinén *et al.* [Belle-II], Phys. Rev. Lett. **127** (2021) no.18, 181802 [arXiv:2104.12624 [hep-ex]].
- [21] I. Adachi et al. [Belle-II], [arXiv:2311.14647 [hep-ex]].
- [22] J. F. Kamenik and C. Smith, Phys. Lett. B 680 (2009), 471-475 [arXiv:0908.1174 [hep-ph]].
- [23] M. Beneke, C. Bobeth and R. Szafron, JHEP 10 (2019),
 232 [erratum: JHEP 11 (2022), 099] [arXiv:1908.07011
 [hep-ph]]; C. Bobeth, M. Gorbahn, T. Hermann,
 M. Misiak, E. Stamou and M. Steinhauser, Phys. Rev.
 Lett. 112 (2014), 101801 [arXiv:1311.0903 [hep-ph]].
- [24] A. J. Buras, J. Girrbach-Noe, C. Niehoff and D. M. Straub, JHEP 02, 184 (2015) [arXiv:1409.4557 [hep-ph]]; W. Altmannshofer, A. J. Buras, D. M. Straub and M. Wick, JHEP 04 (2009), 022 [arXiv:0902.0160 [hep-ph]].
- [25] G. Buchalla and A. J. Buras, Nucl. Phys. B 400, 225-239 (1993)
- [26] G. Buchalla and A. J. Buras, Nucl. Phys. B 548, 309-327 (1999) [arXiv:hep-ph/9901288 [hep-ph]].
- [27] M. Misiak and J. Urban, Phys. Lett. B 451, 161-169 (1999) [arXiv:hep-ph/9901278 [hep-ph]].
- [28] J. Brod, M. Gorbahn and E. Stamou, Phys. Rev. D 83, 034030 (2011) [arXiv:1009.0947 [hep-ph]].
- [29] R. Bause, H. Gisbert and G. Hiller, [arXiv:2309.00075 [hep-ph]].
- [30] D. Das, G. Hiller and I. Nisandzic, Phys. Rev. D 95 (2017) no.7, 073001 [arXiv:1702.07599 [hep-ph]].
- [31] W. Buchmuller and D. Wyler, Nucl. Phys. B 268, 621-653 (1986); B. Grzadkowski, M. Iskrzynski, M. Misiak and J. Rosiek, JHEP 10 (2010), 085 [arXiv:1008.4884 [hep-ph]].
- [32] R. Alonso, B. Grinstein and J. Martin Camalich, Phys. Rev. Lett. **113** (2014), 241802 [arXiv:1407.7044 [hepph]].
- [33] J. Aebischer, A. Crivellin, M. Fael and C. Greub, JHEP 05 (2016), 037 [arXiv:1512.02830 [hep-ph]].
- [34] R. Bause, H. Gisbert, M. Golz and G. Hiller, JHEP 12, 061 (2021) [arXiv:2109.01675 [hep-ph]].
- [35] A. de Giorgi and G. Piazza, [arXiv:2211.05595 [hep-ph]].

- [36] N. Rajeev and R. Dutta, Phys. Rev. D 105 (2022) no.11, 115028 [arXiv:2112.11682 [hep-ph]].
- [37] E. E. Jenkins, A. V. Manohar and P. Stoffer, JHEP 03 (2018), 016 [arXiv:1709.04486 [hep-ph]].
- [38] E. E. Jenkins, A. V. Manohar and P. Stoffer, JHEP 01 (2018), 084 [arXiv:1711.05270 [hep-ph]].
- [39] J. Aebischer, C. Bobeth, A. J. Buras and J. Kumar, JHEP **12** (2020), 187 [arXiv:2009.07276 [hep-ph]].
- [40] E. E. Jenkins, A. V. Manohar and M. Trott, JHEP 10 (2013), 087 [arXiv:1308.2627 [hep-ph]]; E. E. Jenkins, A. V. Manohar and M. Trott, JHEP 01 (2014), 035 [arXiv:1310.4838 [hep-ph]]; R. Alonso, E. E. Jenkins, A. V. Manohar and M. Trott, JHEP 04 (2014), 159 [arXiv:1312.2014 [hep-ph]].
- [41] W. Buchmuller, R. Ruckl and D. Wyler, Phys. Lett. B 191 (1987), 442-448 [erratum: Phys. Lett. B 448 (1999), 320-320]; I. Doršner, S. Fajfer, A. Greljo, J. F. Kamenik and N. Košnik, Phys. Rept. 641 (2016), 1-68 [arXiv:1603.04993 [hep-ph]].
- [42] Y. S. Amhis *et al.* [Heavy Flavor Averaging Group and HFLAV], Phys. Rev. D **107** (2023) no.5, 052008 [arXiv:2206.07501 [hep-ex]].
- [43] Y. Aoki *et al.* [Flavour Lattice Averaging Group (FLAG)], Eur. Phys. J. C 82 (2022) no.10, 869 [arXiv:2111.09849 [hep-lat]];
- [44] J. A. Bailey *et al.* [MILC], Phys. Rev. D **92** (2015) no.3, 034506 [arXiv:1503.07237 [hep-lat]]; H. Na *et al.* [HPQCD], Phys. Rev. D **92** (2015) no.5, 054510 [erratum: Phys. Rev. D **93** (2016) no.11, 119906] [arXiv:1505.03925 [hep-lat]].
- [45] A. Bazavov *et al.* [Fermilab Lattice, MILC, Fermilab Lattice and MILC], Eur. Phys. J. C 82 (2022) no.12, 1141 [erratum: Eur. Phys. J. C 83 (2023) no.1, 21] [arXiv:2105.14019 [hep-lat]]; J. Harrison and C. T. H. Davies, [arXiv:2304.03137 [hep-lat]]; Y. Aoki *et al.* [JLQCD], [arXiv:2306.05657 [hep-lat]].
- [46] B. Capdevila, A. Crivellin, S. Descotes-Genon, L. Hofer and J. Matias, Phys. Rev. Lett. **120** (2018) no.18, 181802 [arXiv:1712.01919 [hep-ph]].
- [47] D. Bečirević, O. Sumensari and R. Zukanovich Funchal, Eur. Phys. J. C **76** (2016) no.3, 134 [arXiv:1602.00881 [hep-ph]].
- [48] R. Aaij *et al.* [LHCb], JHEP **06** (2023), 143 [arXiv:2209.09846 [hep-ex]].

- [49] R. L. Workman *et al.* [Particle Data Group], PTEP **2022**, 083C01 (2022)
- [50] A. Angelescu, D. Bečirević, D. A. Faroughy and O. Sumensari, JHEP **10** (2018), 183 [arXiv:1808.08179 [hep-ph]]; A. Angelescu, D. Bečirević, D. A. Faroughy, F. Jaffredo and O. Sumensari, Phys. Rev. D **104** (2021) no.5, 055017 [arXiv:2103.12504 [hep-ph]].
- [51] A. Greljo, G. Isidori and D. Marzocca, JHEP 07, 142 (2015) [arXiv:1506.01705 [hep-ph]]; D. Buttazzo, A. Greljo, G. Isidori and D. Marzocca, JHEP 11 (2017), 044 doi:10.1007/JHEP11(2017)044 [arXiv:1706.07808 [hep-ph]].
- [52] J. Fuentes-Martín, G. Isidori, M. König and N. Selimović, Phys. Rev. D 101 (2020) no.3, 035024 [arXiv:1910.13474 [hep-ph]].
- [53] J. Fuentes-Martín, G. Isidori, M. König and N. Selimović, Phys. Rev. D **102** (2020) no.3, 035021 doi:10.1103/PhysRevD.102.035021 [arXiv:2006.16250 [hep-ph]].
- [54] D. Bečirević, S. Fajfer, N. Košnik and O. Sumensari, Phys. Rev. D 94 (2016) no.11, 115021 [arXiv:1608.08501 [hep-ph]].
- [55] D. Bečirević, S. Fajfer and N. Košnik, Phys. Rev. D 92 (2015) no.1, 014016 [arXiv:1503.09024 [hep-ph]];
 D. Bečirević, S. Fajfer, N. Košnik and O. Sumensari, Phys. Rev. D 94 (2016) no.11, 115021 [arXiv:1608.08501 [hep-ph]].
- [56] L. Allwicher, D. A. Faroughy, F. Jaffredo, O. Sumensari and F. Wilsch, JHEP **03** (2023), 064 [arXiv:2207.10714 [hep-ph]]; L. Allwicher, D. A. Faroughy, F. Jaffredo, O. Sumensari and F. Wilsch, high-p_T Drell-Yan tails beyond the standard model," Comput. Phys. Commun. **289** (2023), 108749 [arXiv:2207.10756 [hep-ph]].
- [57] P. Athron, R. Martinez and C. Sierra, [arXiv:2308.13426 [hep-ph]].
- [58] A. M. Sirunyan *et al.* [CMS], JHEP **04** (2020), 188 doi:10.1007/JHEP04(2020)188 [arXiv:1910.12127 [hepex]].
- [59] D. Buttazzo, A. Greljo, G. Isidori and D. Marzocca, JHEP **11** (2017), 044 doi:10.1007/JHEP11(2017)044 [arXiv:1706.07808 [hep-ph]].
- [60] F. Feruglio, P. Paradisi and O. Sumensari, JHEP
 11 (2018), 191 doi:10.1007/JHEP11(2018)191
 [arXiv:1806.10155 [hep-ph]].