Macroscopic Quantum Response to Gravitational Waves

Asuka Ito^{1, 2, *} and Ryuichiro Kitano^{2, 3, †}

¹International Center for Quantum-field Measurement Systems for Studies

of the Universe and Particles (QUP), KEK, Tsukuba 305-0801, Japan

²KEK Theory Center, Tsukuba 305-0801, Japan

³Graduate University for Advanced Studies (Sokendai), Tsukuba 305-0801, Japan

We study the excitation of a one-electron quantum cyclotron by gravitational waves. The electron in such as a penning trap is prepared to be at the lowest Landau level, which has an infinite degeneracy parameterized by the spread of the wave function in position space. We find that the excitation rate from the ground state to the first excited state is enhanced by the size of the electron wave function: an electron with a larger wave function feels gravitational waves more. As a consequence, we derive a good sensitivity to gravitational waves at a macroscopic one-electron quantum cyclotron.

Gravitational waves are a powerful tool to explore our universe since they bring valuable information as it has been demonstrated by detection of gravitational waves around kHz with interferometers [1] and of gravitational waves around nHz with pulsar timing arrays [2]. Certainly, advancing multi-frequency gravitational wave observations is crucial for exploring our universe since different sources predict different frequencies of gravitational waves [3]. In the lower frequency range of approximately 10^{-18} to 10^{-16} Hz, there is a viable method for observing gravitational waves, namely, through the cosmic microwave background observations [4-6]. On the flip side, the detection of gravitational waves at frequencies higher than kHz is still in the process of being developed, and it even calls for novel ideas [7-20]. This is the case even though high frequency gravitational waves hold theoretical interest for investigating new areas of physics [10].

One organic strategy to detect high frequency gravitational waves involves employing small-scale experimental setups. This is attributed to the fact that gravitational wave detectors tend to attain heightened sensitivity when their dimensions align with the wavelengths of the gravitational waves. For example, a magnon gravitational wave detector has been proposed, which utilizes the resonant excitation of magnons by gravitational waves to detect signals around the GHz range [7, 8, 15]. In the context of this, there have been substantial proposals for pioneering methods of detecting high frequency gravitational waves by leveraging experiments designed for axion detection [9-14]. However, further improvement of sensitivity is required to probe theoretically interesting targets [10]. One promising way to this end is employing quantum sensing [21].

In this letter, we propose the search for high frequency gravitational waves by using a one-electron quantum cyclotron, which can also be utilized for the dark photon dark matter search [22]. By recasting the reported dark count rate in a penning trap [22], we give upper bounds on gravitational waves at frequencies of O(100) GHz, which have not been searched for. Generically, signal rates of gravitational wave detection are suppressed by the ratio of the size of the detector and the wavelength of the gravitational wave, which is O(mm) in the case of 100 GHz wave. In this sense, a single electron does not sound a good detector as it is small. However, our discussion reveals that the state with a macroscopic wave function can compensate this factor. In an ideal one-electron quantum cyclotron, each energy level has an infinite number of degenerate energy states due to the translational (rotational) invariance of the system. In the symmetric gauge where the Hamiltonian has the rotational invariance, the degenerate states can be labeled by the size of the wave function on the transverse plane. We show that the excitation rate from the ground state to the first excited state is enhanced when the ground state has a large wave function. In a realistic penning trap system where the degeneracy is lifted slightly, the size of (the extent of) the wave function corresponds to that of a magnetron orbit. Similarly, one also finds that the large axial motion enhances the transition rate. It is interesting to note that an electron with a larger wave function feels gravitational waves more. The size of the wave function can be macroscopic such as $O(\mu m) \sim O(mm)$ in the actual setup. This fact may provide insights into the search for high frequency gravitational waves using quantum sensing.

I. ONE-ELECTRON QUANTUM CYCLOTRON

In order to observe the magnetic moment of electrons, a one-electron quantum cyclotron has been utilized [23– 25]. Let us briefly review the principle of the penning trap (see [26–28] for more details). In a penning trap, an external magnetic field is applied on an electron. Taking the direction of the magnetic field B along the zdirection, the electron experiences the cyclotron motion in the x-y plane and the spin precession. Furthermore, an external inhomogeneous electric field E is applied to

^{*} asuka.ito@kek.jp

[†] ryuichiro.kitano@kek.jp

bound the electron in the z-direction. The Hamiltonian of the system is given by

$$H = \frac{1}{2m_e} \left(p_x + \frac{eBy}{2} \right)^2 + \frac{1}{2m_e} \left(p_y - \frac{eBx}{2} \right)^2 + \frac{1}{2m_e} p_z^2 + \frac{1}{2} m_e \omega_z^2 z^2 - 2\mu_B S_z B , \qquad (1)$$

where m_e and e are the mass and charge of an electron, p_i represents the momentum, ω_z represents the angular frequency of the harmonic oscillator along z-axis, μ_B is the Bohr magneton, and S_z is the z component of the spin operator. Note that we took the symmetric gauge for the magnetic field, namely the vector potential was taken to be $A_i = \left(-\frac{By}{2}, \frac{Bx}{2}, 0\right)$, since this gauge is the most convenient for the actual experimental setup, which usually has a rotationally symmetric boundary. Also, the electric field is identified as $E = -\frac{m_e \omega_z^2 z}{e}$. We note that although an electrostatic quadrupole potential is applied in the penning trap actually, which affects the dynamics of the electron in the x-y plane, we will temporarily disregard the effect and reassess it later. The g-factor of the electron has been set as exactly 2 for simplicity.

One can solve the time independent Schrödinger equation with the Hamiltonian (1), and the following solution is obtained:

$$\Psi = \phi_k(z)\psi_{n,m}(x,y)\chi_{\pm} , \qquad (2)$$

where

$$\phi_k(z) = \frac{1}{\sqrt{2^k k!}} \left(\frac{m_e \omega_z}{\pi}\right)^{1/4} e^{-\frac{m_e \omega_z}{2} z^2} H_k\left(\sqrt{m_e \omega_z} z\right) ,$$
(3)

is the wave function of the axial motion with $k = 0, 1, \ldots \infty$ (H_k is the Hermite polynomial), and

$$\psi_{n,m}(x,y) = \frac{eB}{\sqrt{2\pi}} e^{im\varphi} \sqrt{\frac{(n-\frac{m+|m|}{2})!}{(n-\frac{m-|m|}{2})!}} e^{-\frac{eB}{4}\rho^2} \times \left(\frac{eB\rho^2}{2}\right)^{|m|/2} L_{n-\frac{1}{2}(m+|m|)}^{|m|} \left(\frac{eB\rho^2}{2}\right) (4)$$

with $n = 0, 1, ..., \infty$, $m = -\infty, ..., n - 1, n$, is the wave function of the cyclotron in the symmetric gauge in the *xy* plane specified by the radial distance ρ and the azimuth φ (L_n^m is the generalized Laguerrel polynomial), and

$$\chi_{+} = \begin{pmatrix} 1\\0 \end{pmatrix}, \quad \chi_{-} = \begin{pmatrix} 0\\1 \end{pmatrix}, \tag{5}$$

are the spinors for the spin up and the down states. The excitation of the quantum cyclotron described by Eq. (4) from the ground state (n = 0) to the excited state (n = 1) can be measured with penning traps [23–25]. In principle, we have infinite degenerate states labeled by $m (\leq n)$ for each energy level specified by n. However in practice, the quantum number m has a minimum value corresponding

to the size of the experimental apparatus because the size of the wave functions become larger for larger |m|. Indeed, using the cyclotron frequency $\omega_c = eB/m_e$, the root mean square of the radial distance of the wave functions is given by $\sqrt{2(1-m)/m_e\omega_c}$ for $(n = 0, m \le 0)$ and $\sqrt{2(3-m)/m_e\omega_c}$ for $(n = 1, m \le 1)$, and the wave functions exponentially decrease over them. Therefore, the value of m for these energy levels in a penning trap with an radius R must satisfies¹

$$|m| \lesssim m_e \omega_c R^2 \sim 10^9 \times \left(\frac{\omega_c/2\pi}{100 \text{GHz}}\right) \left(\frac{R}{0.5 \text{mm}}\right)^2.$$
 (6)

As we will see in the next section, this huge degeneracy can give rise to enhancement of transition probability from the ground state to the excited state by gravitational waves.

In the actual experimental setup of such as a penning trap, the degeneracy for the m quantum number is lifted by the electric field such that each energy level has the interval of the magnetron frequency $\sim \omega_z/2\omega_c^2$. In such a case, for a state with a large expectation value of |m|, it may be more appropriate to take the coherent state. The coherent state corresponds to the classical magnetron motion with the radius $r_m \sim \sqrt{2|m|/(m_e\omega_c)}$. When |m| is large, one can understand |m| as $m_e\omega_c r_m^2/2$ by ignoring O(1/|m|) quantities. Similarly, we usually have large k for the axial motion in experiments, and then the axial motion would also take the coherent state, i.e., $k \sim m_e\omega_z A^2/2$ with A representing the amplitude of the oscillation in the axial direction.

In experiments [23–25], the measurement of the excitation has been operated with the technique of the quantum jump spectroscopy; one applies an additional weak magnetic bottle field to couple the axial motion to the quantum cyclotron, so that the energy transition of the quantum cyclotron can be measured indirectly by monitoring the the axial frequency, which can be easily measured by detecting the current induced by the axial motion. Importantly, the interaction Hamiltonian commutes with the cyclotron Hamiltonian, so that a quantum nondemolition measurement of the cyclotron states is allowed. We note that since the spin precession also couples to the axial motion due to the magnetic bottle field, one can measure the spin excitation in the same way. However, we will see that the spin excitation by gravitational waves is subdominant compared with the excitation of the quantum cyclotron. In the following section, we will review how to formulate the interaction between an electron in a penning trap and gravitational waves.

¹ Since the wave functions exponentially decrease with a decay constant $(m_e\omega_c)^{-1/2}$ ($\ll R$) outside the root mean square of the radial distance of the wave functions, the definition of m_{\min} is actually equivalent to require that the wave functions sufficiently small at R.

II. A DIRAC PARTICLE IN CURVED SPACETIME

To investigate the effect of gravitational waves on a one-electron quantum cyclotron, we consider the Dirac equation in curved spacetime and take the nonrelativistic limit assuming that the velocity of the electron is much smaller than the speed of light as is satisfied in real experiments. Furthermore, we need to use an appropriate coordinate which comoves with the system. Such a procedure has been carried out to study excitation of spins by gravitational waves [8] and/or to evaluate the inertial and gravitational effects due to the Earth on the magnetic moment of electrons/muons [28]. More explicitly, the non-relativistic Hamiltonian for a Dirac particle such as an electron with mass m_e in curved spacetime up to the order of m_e^0 is given by [28]

$$H_{\text{int}} = \left(1 + \frac{1}{2}R_{0k0l}x^{k}x^{l}\right)m_{e} - eA_{0} + \frac{1}{3}R_{0kil}\left(\Pi_{i}x^{k}x^{l} + x^{k}x^{l}\Pi_{i}\right) + \frac{1}{2}\epsilon_{0ijl}S^{l}R_{ijk0}x^{k},$$
(7)

where S^i stands for the spin operator, and A_{μ} represents the vector potential of the electromagnetic field. The Riemann tensor $R_{\mu\nu\rho\sigma}$ is evaluated at the origin of the coordinate $x^i = 0$, and $\Pi_j = -i\partial_j - eA_j$ is the canonical momentum. The first term can be regarded as the rest mass and its correction due to gravity. Compared with it, the third and the fourth terms are higher order effects of $v/c \ll 1$ (v: speed of the electron, c: speed of light) or $1/m_e |x^i| \ll 1$. Thus, we will neglect them and focus on only the first term, which is at the order of m_e . In particular, we see that there is no interaction between the spin of the fermion and gravitational waves at the order. We now consider gravitational waves on the Minkowski spacetime, the metric is given by $g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}$, where $\eta_{\mu\nu}$ is the Minkowski metric and $h_{\mu\nu}$ is the traceless transverse metric, namely, it represents gravitational waves. Then, the Riemann tensor is

$$R^{\alpha}{}_{\mu\beta\nu} = \frac{1}{2} (h^{\alpha}{}_{\nu,\mu\beta} - h^{,\alpha}{}_{\mu\nu}{}_{,\beta} - h^{\alpha}{}_{\beta,\mu\nu} + h^{,\alpha}{}_{\mu\beta}{}_{,\nu}) .$$
 (8)

It should be noted that although we employs the transverse-traceless coordinate to calculate the Riemann tensor, the result is the same as that obtained in a Fermi normal coordinate since the Riemann tensor is gauge invariant at linear order. As a gravitational wave, which drives the excitation of the one-electron quantum cyclotron, we consider a planer gravitational wave:

$$h_{ij}(\boldsymbol{x},t) = h^{(+)}\cos\left(\omega t - \boldsymbol{k}\cdot\boldsymbol{x}\right)e_{ij}^{(+)} +h^{(\times)}\cos\left(\omega t - \boldsymbol{k}\cdot\boldsymbol{x}+\alpha\right)e_{ij}^{(\times)}, \quad (9)$$

where the polarization tensors satisfy $e_{ij}^{(\sigma)} e_{ij}^{(\sigma')} = \delta_{\sigma\sigma'}$. α represents the difference of the phases of the polarization. Note that the polarization tensors can be explicitly constructed as

$$e_{ij}^{(+)} = \frac{1}{\sqrt{2}} \begin{pmatrix} \cos\theta^2 & 0 & -\cos\theta\sin\theta \\ 0 & -1 & 0 \\ -\cos\theta\sin\theta & 0 & \sin\theta^2 \end{pmatrix}, (10)$$
$$e_{ij}^{(\times)} = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & \cos\theta & 0 \\ \cos\theta & 0 & -\sin\theta \\ 0 & -\sin\theta & 0 \end{pmatrix}.$$
(11)

In the above Eqs. (10) and (11), we defined the + mode as a deformation in the *y*-direction. Using Eqs. (8)-(11) in Eq. (7), we obtain the interaction Hamiltonian between a one-electron quantum cyclotron and a planner gravitational wave:

$$H_{\rm int} \simeq \frac{m_e \omega^2 e^{i\omega t}}{8\sqrt{2}} \left[\left(\cos^2 \theta \, x^2 + \sin^2 \theta \, z^2 - y^2 - 2\cos \theta \sin \theta z x \right) h^{(+)} + e^{i\alpha} \left(2\cos \theta x y - \sin \theta y z \right) h^{(\times)} \right] + (\text{h.c.}). \quad (12)$$

In the next section, using the above interaction Hamiltonian, we will calculate the excitation rate of a oneelectron quantum cyclotron by a planner gravitational wave and evaluate the ability of the penning trap as a gravitational wave detector.

III. GRAVITATIONAL WAVE INDUCED EXCITATION

We are now ready to calculate the excitation rate of a one-electron quantum cyclotron by gravitational waves. One can calculate the excitation probability P during

an observation time τ at the resonance point, i.e., the frequency of a gravitational wave coincides with the cyclotron frequency, by using Eq. (12) [29]:

$$P \simeq \sin^2 \left(\frac{\left| \langle \Psi_{\text{out}} | \tilde{H}_{\text{int}} | \Psi_{\text{in}} \rangle \right|}{2} \tau \right), \qquad (13)$$

where \tilde{H}_{int} is defined by $H_{int} = \tilde{H}_{int}e^{i\omega t} + (h.c.)$. Note that we implicitly assumed that the gravitational wave has a coherence during the observation time. More explicitly, from Eqs. (4) and (12), one can calculate the matrix element in Eq. (13) for the transition from a ground state to an excited state as

$$\left| \langle 1, m+2, k | \tilde{H}_{\text{int}} | 0, m, k \rangle \right|^2 = \frac{|m|\omega^2}{128} \left[\left((1 + \cos^2 \theta) h^{(+)} - 2\cos\theta\sin\alpha h^{(\times)} \right)^2 + 4\cos^2\theta\cos^2\alpha (h^{(\times)})^2 \right], \quad (14a)$$

$$\left| \langle 1, m, k | \tilde{H}_{\text{int}} | 0, m, k \rangle \right|^2 = \frac{(|m|+1)\omega^2}{128} \sin^4 \theta (h^{(+)})^2, \tag{14b}$$

$$\left| \langle 1, m+1, k+1 | \tilde{H}_{\text{int}} | 0, m, k \rangle \right|^2 = \frac{(k+1)\omega^3}{32\omega_z} \sin^2 \theta \left[\left(\cos \theta h^{(+)} - \sin \alpha h^{(\times)} \right)^2 + \cos^2 \alpha (h^{(\times)})^2 \right],$$
(14c)

$$\left| \langle 1, m+1, k-1 | \tilde{H}_{\text{int}} | 0, m, k \rangle \right|^2 = \frac{k\omega^3}{32\omega_z} \sin^2 \theta \left[\left(\cos \theta h^{(+)} - \sin \alpha h^{(\times)} \right)^2 + \cos^2 \alpha (h^{(\times)})^2 \right].$$
(14d)

Eqs. (14a) and (14b) come from $\tilde{H}_{\rm int}$ including only xand/or y components. It is worth noting that the transition probability increases for larger |m|. This is a sort of particular feature of the excitation by gravitational waves in contrast to the excitation caused by electromagnetic fields; we have the same transition probability from $|0, m\rangle$ to $|1, m + 1\rangle$ for arbitrary m in the case of dipole excitation by electric fields. Interestingly, it enables us to increase the sensitivity of a one-electron quantum cyclotron to gravitational waves by enlarging the wave function, whose size is proportional to $\sqrt{|m|}$. On the other hand, Eqs. (14c) and (14d) are calculated from $\tilde{H}_{\rm int}$ including zcomponent. It also gets increased for larger k. Thus, we can also increase the transition rate by preparing a large wave function in the z-direction.

Now let us evaluate the sensitivity of a one-electron quantum cyclotron to a planer gravitational wave by using Eqs. (13)-(14d). Supposing that we have an initial state $|\Psi_{\rm in}\rangle = |0, m, k\rangle$, it can be excited to $|1, m+2, k\rangle$ through Eq. (14a), $|1, m, k\rangle$ through Eq. (14b), or $|1, m+1, k \pm 1\rangle$ through Eqs. (14c) and (14d). Since the transition rate depends on m and k, we adopt their values used in experiments [23-25] as a reference. We then need to look back on Eq. (1), in which we just considered an electric field along the axial direction. However, in real measurements, a quadruple potential is used to combine an electron in the axial axis, so that the cyclotron motion in the x-y plane is also modified by the potential. Then the degeneracy of the cyclotron energy levels are lifted due to the appearance of the magnetron motion [26-28]. A specific value of *m* would be chosen in such a measurement. Indeed, $|m| \sim k \sim 10^3$ is obtained in the cooling limit with a sideband cooling [26]. As we already mentioned, when the quantum numbers |m| and k are much larger than unity, coherent states are realized for the magnetron motion and the axial motion, respectively. However, it does not change the evaluation of the excitation of a one-electron quantum cyclotron by gravitational waves at least approximately, since the coherent state has a Gaussian distribution with a peak at $|m| \sim k \sim 10^3$. Using the upper limit on the dark count rate per unit time, 4.33×10^{-6} s⁻¹, obtained in [22], one can evaluate the sensitivity of a one-electron quantum cyclotron to a planner gravitational wave through the

transition of (14a) and (14b),

$$h_0 \sim 1.2 \times 10^{-18} \left(\frac{\omega/2\pi}{200 \text{GHz}}\right)^{-1} \left(\frac{|m|}{1000}\right)^{-1/2} \left(\frac{\tau}{1 \text{day}}\right)^{-1/2},$$
(15)

and of (14c) and (14d),

$$h_0 \sim 3.8 \times 10^{-20} \quad \left(\frac{\omega/2\pi}{200 \text{GHz}}\right)^{-3/2} \left(\frac{\omega_z/2\pi}{100 \text{MHz}}\right)^{1/2} \times \left(\frac{k}{1000}\right)^{-1/2} \left(\frac{\tau}{1 \text{day}}\right)^{-1/2}.$$
 (16)

In the estimation, we took average of θ because we do not know the value in advance and assumed no polarization by setting $\alpha = 0$, and $h^{(+)} = h^{(\times)} = h_0$. Eqs. (15) and (16) would be valid for a frequency range from 20 GHz to 200 GHz because the cyclotron frequency can be tuned in the range, as discussed in [22]. We took an observation time of 1 day just as a reference value. Note that it may significantly deviate from the coherence time of expected gravitational wave sources [10].

Finally, let us consider a strategy to improve the sensitivity to gravitational waves by preparing a macroscopic one-electron quantum cyclotron. As Eqs. (14a)-(14d) tell us, the transition rate by gravitational waves is proportional to the size of (the extent of) the wave function, which is parameterized by |m| and k. One can increase |m| and k arbitrarily in principle. However, in practice, there would be a maximum value for them like Eq. (6) specified by the size of the apparatus of a penning trap. Supposing that we can tune the value of |m|to be the maximum value, $m_e \omega_c R^2$, and k is also taken to be $k \sim |m|$ by a sideband cooling, the sensitivity of a one-electron quantum cyclotron to a gravitational wave would be

$$h_0 \sim 2.2 \times 10^{-21} \left(\frac{\omega/2\pi}{200 \text{GHz}}\right)^{-3/2} \left(\frac{R}{0.2 \text{mm}}\right)^{-1} \left(\frac{\tau}{1 \text{day}}\right)^{-1/2}$$
(17)

$$h_0 \sim 6.9 \times 10^{-23} \quad \left(\frac{\omega/2\pi}{200 \text{GHz}}\right)^{-2} \left(\frac{\omega_z/2\pi}{100 \text{MHz}}\right)^{1/2} \\ \times \left(\frac{R}{0.2 \text{mm}}\right)^{-1} \left(\frac{\tau}{1 \text{day}}\right)^{-1/2}, (18)$$

through the transition (14c) and (14d). As expected, we see that the sensitivity is increased for larger R because then the wave function becomes more macroscopic. Thus, we can improve the sensitivity by preparing a large apparatus. However, we note that one can not take arbitrary large value of R in our formalism since $\omega R \lesssim 1$ should be satisfied to validate the interaction Hamiltonian (7). It is because that Eq. (7) has been derived by truncating the infinite sum of terms in the metric of Fermi normal coordinates by assuming $\omega R \lesssim 1$ [8, 28]. If the assumption does not hold, one needs to take into account infinite terms in the metric as was done in [15]. As a consequence, it is natural to expect that the sensitivity would have a peak around $\omega R \sim 1$ at which the wavelength of the gravitational wave is comparable to the size of the wave function [15]. In the evaluation (15), we took R = 0.2mm to satisfy $\omega R \lesssim 1$ at $\omega/2\pi = 200$ GHz. Therefore, for example, we can take R = 2mm when $\omega/2\pi = 20$ GHz is considered. Notably, such a setup can be realized in the existing experiments [23-25].

IV. CONCLUSION

We studied the excitation of a one-electron quantum cyclotron by gravitational waves. We calculated the transition rate from degenerate ground states, in which each wave function has a different size, to first excited states. It turned out that the transition rate from the ground state to the first excited state is enhanced when the ground state has a larger wave function. As a consequence, good sensitivities to high frequency gravitational waves are derived for a macroscopic one-electron quantum cyclotron. Therefore, in principle, one can increase the sensitivity to gravitational waves just simply preparing a ground state with a larger wave function. This fact may shed light on searches for high frequency gravitational waves by quantum sensing. There is a related study in Ref. [30] where it is found that gravitational wave signals are enhanced by using an atom in a highly excited state such as a Rydberg atom. In contrast to atoms, the penning trap experiment does not require preparing such a special state, and the enhancement is more or less automatic. Another virtue of the experiment is the possibility to scan the frequencies of probed gravitational waves by changing the strength of the magnetic field and the frequencies of driving fields. Indeed, one can search for gravitational waves within a frequency range of 20 to 200GHz with a penning trap, maintaining the sensitivity [22].

ACKNOWLEDGMENTS

A. I. was supported by World Premier International Research Center Initiative (WPI), MEXT, Japan. This work is supported by JSPS KAKENHI Grant Numbers JP22K14034 (A.I.), JP22K21350 (R.K.), JP21H01086 (R.K.), and JP19H00689 (R.K.).

- B. P. Abbott <u>et al.</u> (LIGO Scientific, Virgo), Phys. Rev. Lett. **116**, 061102 (2016), arXiv:1602.03837 [gr-qc].
- [2] G. Agazie <u>et al.</u> (NANOGrav), Astrophys. J. Lett. **951**, L8 (2023), arXiv:2306.16213 [astro-ph.HE].
- [3] K. Kuroda, W.-T. Ni, and W.-P. Pan, Int. J. Mod. Phys. D 24, 1530031 (2015), arXiv:1511.00231 [gr-qc].
- [4] Y. Akrami <u>et al.</u> (Planck), Astron. Astrophys. **641**, A10 (2020), arXiv:1807.06211 [astro-ph.CO].
- [5] D. Paoletti, F. Finelli, J. Valiviita, and M. Hazumi, Phys. Rev. D 106, 083528 (2022), arXiv:2208.10482 [astro-ph.CO].
- [6] E. Allys <u>et al.</u> (LiteBIRD), PTEP **2023**, 042F01 (2023), arXiv:2202.02773 [astro-ph.IM].
- [7] A. Ito, T. Ikeda, K. Miuchi, and J. Soda, Eur. Phys. J. C 80, 179 (2020), arXiv:1903.04843 [gr-qc].

- [8] A. Ito and J. Soda, Eur. Phys. J. C 80, 545 (2020), arXiv:2004.04646 [gr-qc].
- [9] A. Ejlli, D. Ejlli, A. M. Cruise, G. Pisano, and H. Grote, Eur. Phys. J. C 79, 1032 (2019), arXiv:1908.00232 [gr-qc].
- [10] N. Aggarwal et al., Living Rev. Rel. 24, 4 (2021), arXiv:2011.12414 [gr-qc].
- [11] A. Berlin, D. Blas, R. Tito D'Agnolo, S. A. R. Ellis, R. Harnik, Y. Kahn, and J. Schütte-Engel, Phys. Rev. D 105, 116011 (2022), arXiv:2112.11465 [hep-ph].
- [12] V. Domcke, C. Garcia-Cely, and N. L. Rodd, Phys. Rev. Lett. **129**, 041101 (2022), arXiv:2202.00695 [hep-ph].
- [13] M. E. Tobar, C. A. Thomson, W. M. Campbell, A. Quiskamp, J. F. Bourhill, B. T. McAllister, E. N. Ivanov, and M. Goryachev, Symmetry 14, 2165 (2022), arXiv:2209.03004 [physics.ins-det].
- [14] A. Berlin, D. Blas, R. Tito D'Agnolo, S. A. R. Ellis, R. Harnik, Y. Kahn, J. Schütte-Engel, and M. Wentzel, (2023), arXiv:2303.01518 [hep-ph].

- [15] A. Ito and J. Soda, Eur. Phys. J. C 83, 766 (2023), arXiv:2212.04094 [gr-qc].
- [16] M. S. Pshirkov and D. Baskaran, Phys. Rev. D 80, 042002 (2009), arXiv:0903.4160 [gr-qc].
- [17] A. D. Dolgov and D. Ejlli, JCAP 12, 003 (2012), arXiv:1211.0500 [gr-qc].
- [18] V. Domcke and C. Garcia-Cely, Phys. Rev. Lett. **126**, 021104 (2021), arXiv:2006.01161 [astro-ph.CO].
- [19] S. Ramazanov, R. Samanta, G. Trenkler, and F. R. Urban, (2023), arXiv:2304.11222 [astro-ph.HE].
- [20] A. Ito, K. Kohri, and K. Nakayama, (2023), arXiv:2305.13984 [gr-qc].
- [21] C. L. Degen, F. Reinhard, and P. Cappellaro, Rev. Mod. Phys. 89, 035002 (2017).
- [22] X. Fan, G. Gabrielse, P. W. Graham, R. Harnik, T. G. Myers, H. Ramani, B. A. D. Sukra, S. S. Y. Wong,

and Y. Xiao, Phys. Rev. Lett. **129**, 261801 (2022), arXiv:2208.06519 [hep-ex].

- [23] B. C. Odom, D. Hanneke, B. D'Urso, and G. Gabrielse, Phys. Rev. Lett. 97, 030801 (2006).
- [24] D. Hanneke, S. Fogwell, and G. Gabrielse, Phys. Rev. Lett. **100**, 120801 (2008), arXiv:0801.1134 [physics.atom-ph].
- [25] X. Fan, T. G. Myers, B. A. D. Sukra, and G. Gabrielse, (2022), arXiv:2209.13084 [physics.atom-ph].
- [26] L. S. Brown and G. Gabrielse, Rev. Mod. Phys. 58, 233 (1986).
- [27] D. Hanneke, S. F. Hoogerheide, and G. Gabrielse, Phys. Rev. A 83, 052122 (2011), arXiv:1009.4831 [physics.atom-ph].
- [28] A. Ito, Class. Quant. Grav. 38, 195015 (2021), arXiv:2011.11217 [gr-qc].
- [29] D. J. Griffiths, <u>Introduction of Quantum Mechanics</u> (Prentice Hall, Inc., 1995).
- [30] U. Fischer, Class. Quant. Grav. **11**, 463 (1994).