All-charm tetraquark mass and possible quantum numbers of X(6900)

Morgan Kuchta*

University of Wrocław, Institute of Theoretical Physics, pl. Maxa Borna 9 Wrocław, 50-204, Poland

Abstract

In this work we propose possible quantum numbers of X(6900) and suggest a model for it internal structure that explains its unusually high mass. We solve the Schrödinger Equation with Mathematica 12, first for charmonium spectrum, then for all-charm tetraquark spectrum which is understood as a pair of two-particle states, mesons or diquark-antidiquark states. The obtained candidates for all-charm te-traquark will be separated into contributors to various resonances and structures that are visible in the experiments then an explanation for the prominence of X(6900) will be proposed.

1 Introduction

In June 2020 in LHCb experiment found in proton-proton collisions exotic meson originally named X(6900) along with it a broad structure between 6200 - 6800 MeV and a smaller peak around 7200 MeV were found [1]. Until now, the parity and the charge symmetry, as well as the total angular momentum of any of the states mentioned here were not identified. The mass of X(6900) was surprisingly high, as before its discovery the expected mass of the ground state was placed around 6200 MeV, which means that it should have been found within the region of the broad structure. In 2022, the ATLAS collaboration searched for potential all-charm tetraquark ($cc\bar{c}\bar{c}$) states in two different decay channels [2]. Results of the search also feature a peak around 6900 MeV and the broad structure visible in results archived in the previous experiment. The experiment differs from the one conducted in 2020 by

investigating the $J/\psi + \Psi(2S)$ invariant mass spectrum, where the two significant peaks were found, one supposedly corresponding to the peak around 6900 MeV found in LHCb, another around 7200 MeV, both of them have statistical significance lesser than 5σ . Other significant results were presented by the CMS collaboration [3] that suggests three peaks in the $di - J/\psi$ spectrum. A summary of the results can be found in the Table 1.

In this work we will suggest possible quantum numbers for the X(6900) together with its quark structure that explains seemingly lacking prominent ground state and create tables of possible masses of other all-charm and allbottom tetraquarks applying a compact tetraquark framework. The discussion will not involve in-medium effects. Numerical calculations have been performed using Wolfram Language and Mathematica 13 and the base code by F. Schöberl and W. Lucha [4] which had been modified and adjusted to its application in the present work. The code can be found in the supplemental file 1. In this work we will use new notation for exotic hadrons suggested in reference [5].

2 Charmonia and bottomonia

The charmonium spectrum can be obtained through solving the time-independent Schrödinger equation with Fermi-Breit Hamiltonian [6] with reduced mass $\mu_{12} = \frac{m_1 m_2}{m_1 + m_2}$,

$$\begin{bmatrix} m_1 + m_2 + \frac{1}{2\mu_{12}} \left(-\frac{d^2}{dr^2} + \frac{l(l+1)}{r^2} \right) + \\ + V_{12}^S + V_{12}^{SS} + V_{12}^{LS} + V_{12}^T \right] \Psi = E_{12}^{n,l} \Psi.$$
(1a)

^{*309312@}uwr.edu.pl

Table 1: Summary of estimated masses (in MeV) and widths of four resonances from three different sources. If the source includes several estimations they are included with brief descriptions. More detailed explanations can be found in the sources. Marker "NA" suggests that resonance was either not identified or its mass was not estimated by the model.

		X(6200)	X(6500)	X(6900)	X(7200)
LHCb [1]	No interference	Observed	Observed	m = 6905 MeV $\Gamma = 80 \text{ MeV}$	Observed
	NRSPS interference	Observed	Observed	m = 6886 MeV $\Gamma = 168 \text{ MeV}$	Observed
ATLAS [2]	Model A	$m = 6.22 \text{ GeV}$ $\Gamma = 0.31 \text{ GeV}$	$m = 6.62 \text{ GeV}$ $\Gamma = 0.31 \text{ GeV}$	m = 6.87 GeV $\Gamma = 0.12 \text{ GeV}$	$m = 7.22 \text{ GeV}$ $\Gamma = 0.10 \text{ GeV}$
	Model B	NA NA	NA NA	NA NA	$m = 6.78 \text{ GeV}$ $\Gamma = 0.39 \text{ GeV}$
CMS [3]		NA NA	m = 6552 MeV $\Gamma = 124 \text{ MeV}$	m = 6927 MeV $\Gamma = 122 \text{ MeV}$	$m = 7287 \text{ MeV}$ $\Gamma = 95 \text{ MeV}$

The Hamiltonian includes kinetic energy and the contribution of strong interaction between the quark-antiquark pair. The pair interaction includes the one gluon exchange (OGE). For the charmonium and the bottomonium spectrum Cornell potential applies,

$$V_{ij}^G(r_{ij}) = \kappa_s \frac{\alpha_s}{r_{12}} + \sigma r_{12}, \qquad (1b)$$

along with three spin dependent terms; spin-spin term (V_{ij}^{SS}) , spin-orbit term (V_{ij}^{SL}) and tensor term (V_{ij}^{T}) ;

$$V_{ij}^{SS}(r_{ij}) = -\frac{8\kappa_s \alpha_s \pi}{3m^2} (\frac{\sigma_{ss}}{\sqrt{\pi}})^3 e^{-\sigma_{ss}^2 r_{ij}^2} S_i S_j, \qquad (1c)$$

$$V_{ij}^{LS}(r_{ij}) = \left[-\frac{3\kappa_s \alpha_s}{2m^2} \frac{1}{r_{ij}^3} - \frac{b}{2m^2} \frac{1}{r_{ij}} \right] LS, \quad (1d)$$

$$V_{ij}^{T}(r_{ij}) = -\frac{12\kappa_{s}\alpha_{s}}{4m^{2}}\frac{1}{r_{ij}^{3}}\left(\frac{(S_{i}r_{ij})(S_{j}r_{ij})}{r_{ij}^{2}} - \frac{S_{i}S_{j}}{3}\right).$$
 (1e)

In the equation (1b) there are three parameters to be set. The first one is the Casimir coefficient κ_s that can be obtained by calculating scattering amplitude for the quark-

antiquark pair that form a colour-singlet:

$$\kappa_s^1 = -f_1 = \frac{1}{4} \Sigma_\alpha (c_3^{\dagger} \lambda^\alpha c_1) (c_2^{\dagger} \lambda^\alpha c_4) =$$

= $\frac{1}{4} (\frac{1}{\sqrt{3}})^2 Tr(\lambda^\alpha \lambda^\alpha) = \frac{-4}{3}.$ (2)

The other two parameters, strong coupling constant α_s and string tension σ are obtained from a fit to the charmonium and bottomonium spectra and radiative decays [7]. Spin dependent terms include one additional parameter σ_{ss} which is also impossible to determine empirically. Mesons are formed from two quarks with spin $\frac{1}{2}$ that can either form spin singlets (S = 0) or spin triplets (S = 3):

$$2 \otimes 2 = 1 \oplus 3. \tag{3}$$

The contribution of the spin-spin internation in the V^{SS} is evaluated using:

$$\langle S_i S_j \rangle = \frac{1}{2} \langle S^2 - S_1^2 - S_2^2 \rangle = \begin{cases} \frac{-3}{4}, & S = |0\rangle \\ \frac{1}{4}, & S = |1\rangle, \end{cases}$$
(4)

where S_1, S_2 are the spins of each quark and S is the total

spin. In V^{SL} for $L \neq 0$:

$$\langle LS \rangle_{S=|1\rangle} = \begin{cases} -(l-1), & J = L-1 \\ -1, & J = L \\ l, & J = L+1. \end{cases}$$
(5)

The tensor dependent part of the potential (1e) ought to be transformed with the usage of the identity (6) [8] to remove r_{ij} dependency and include J and L instead (7).

$$(a \cdot b)r^2 - 3(a \cdot r)(b \cdot r) = \frac{r^2}{(2l+3)(2l+1)} \times (6) \times (k^2(a \cdot b) - 3(a \cdot k)(b \cdot k) - 3(b \cdot k)(a \cdot k))$$

$$\langle T_{ij} \rangle = 12 \left\langle \frac{(S_i r_{ij})(S_j r_{ij})}{r_{ij}^2} - \frac{S_i S_j}{3} \right\rangle =$$

$$= \frac{4}{(2l+3)(2l-1)} \left\langle S^2 L^2 + \frac{3}{2}(LS) + 3(LS)^2 \right\rangle$$
(7)

Therefore $V^T \neq 0$ for $S \neq 0$ and $L \neq 0$:

$$\langle T_{ij} \rangle = \begin{cases} 2, \quad J = L \\ \frac{-2(l+1)}{2l-1}, \quad J = L-1 \\ \frac{-2l}{2l+3}, \quad J = L+1. \end{cases}$$
(8)

We made appropriate choice of the parameters for charmonia (9a) and bottomonia (9b):

$$\alpha_0 = 0.198, \quad \sigma = 0.177 \text{ GeV}^2,$$

 $\sigma_{ee} = 1.08, \quad m_e = 1.263 \text{ GeV}.$
(9a)

$$\alpha_0 = 0.164, \quad \sigma = 0.177 \text{ GeV}^2,$$

 $\sigma_{ss} = 1.08, \quad m_b = 4.581 \text{ GeV}.$
(9b)

Masses and widths of the resonances were sourced from PDG [9].

3 Tetraquarks

The tools that were first used to properly describe nonexotic hadrons [10][11] may be applied to the tetraquarks and other exotic hadrons [12][13]. The discussion of potential tetraquark states predates the discovery of the first exotic hadron candidate $\chi_{c1}(3872)$, previously known as X(3872), in 2003 [14]. In terms of the fundamental representation, a quark and an antiquark could form a coloursinglet meson that can be observed in the resonances or a colour octet (10a). Non-singlet colour states have not been observed [9] due to the colour confinement, however we can treat newly build colour octets as our new "building blocks" that may be combined into a colour singlet and multiple non-singlets (10b).

$$3 \otimes \bar{3} = 1 \oplus 8 \tag{10a}$$

$$8 \otimes 8 = 1 \oplus 8 \oplus 8 \oplus 10 \oplus \overline{10} \oplus 27 \tag{10b}$$

The concept of a diquark has been frequently used to describe exotic hadrons since the conception of the quark model [15]. Compact tetraquarks are usually described as bound states of a diquark and an antidiquark forming an exotic meson [16]. In this work we will call this configuration "diquark-antidiquark configuration" or "DA". In addition to the diquark picture we will also consider combination of two colour octets, referred to as "mesonmeson configuration" or "MM". To obtain masses of the tetraquarks we use the parameters calculated in the previous section to calculate masses of colour non-singlet states.

$$3 \otimes 3 = \overline{3} \oplus 6, \quad \overline{3} \otimes \overline{3} = 3 \oplus \overline{6}$$
 (10c)

$$6 \otimes \overline{6} = 1 \oplus 8 \oplus 27 \tag{10d}$$

We can see that there are several possible methods of "composing" tetraquaks. Using the same method as in Eq. (2) to obtain Casimir Coefficient for colour singlet, we archive the results displayed in the Table 4 and Table 5. The parameters used in the Cornell Potential Eq. (1b) for interactions between the pairs are identical for those used to obtain charmonium and bottomonium spectra, modified by the Casimir scaling (11) [17][18]. A transition from the DA to the MM configuration could be performed unless the distance between the diquark and the antidiquark is sufficiently larger than the distance between their constituents [19], therefore the parameters α_s and σ need to be scaled accordingly.

$$\frac{V_1(r)}{V_r(2)} = \frac{\kappa_{s1}}{\kappa_{s2}} \tag{11}$$

The compact tetraquark picture is a description in which interaction between a diquark and an antidiquark is treated similarly to interaction between a quark and an

Name	$N^{2S+1}l_J$	J^{PC}	$M_{exp}(MeV)$	$\Gamma({\rm MeV})$	$M_{cal}({ m MeV})$	$rac{M_{cal}-M_{exp}}{M_{exp}}(\%)$
$\eta_c(1S)$	$1^{1}S_{0}$	1^{-+}	2983.9 ± 0.5	32.0 ± 0.7	2983.8	< 0.1
J/ψ	$1^{3}S_{1}$	1	3096.900 ± 0.006	92.9 ± 2.8	3094.7	< 0.1
$h_c(1P)$	$1^{1}P_{1}$	1^{+-}	3525.38 ± 0.11	0.7 ± 0.4	3576.1	1.4
$\chi_{c0}(1P)$	$1^{3}P_{0}$	0^{++}	3414.71 ± 0.30	10.8 ± 0.6	3286.5	3.8
$\chi_{c1}(1P)$	$1^{3}P_{1}$	1^{++}	3510.67 ± 0.05	0.84 ± 0.04	3580.9	1.9
$\chi_{c2}(1P)$	$1^{3}P_{2}$	2^{++}	3556.17 ± 0.07	1.97 ± 0.09	3365.6	5.4
$\eta_c(2S)$	$2^{1}S_{0}$	1^{-+}	3637.5 ± 1.1	11.3 ± 3.2	3576.4	1.6
$\psi(2S)$	$2^{3}S_{1}$	$1^{}$	3686.10 ± 0.06	294 ± 8	3641.9	1.2
$\psi(3770)$	$1^{3}D_{1}$	$1^{}$	3773.7 ± 0.4	27.2 ± 1.0	3758.5	0.4
$\psi_2(3823)*$	$1^{3}D_{2}$	$2^{}$	3823.7 ± 0.5	< 5.2	3816.4	0.4
$\psi_3(3842)*$	$1^{3}D_{3}$	3	3842.71 ± 0.20	2.8 ± 0.6	3602.8	6.2
$Z_c(3900)$	$2^{1}P_{1}$	1^{+-}	3887.1 ± 2.6	28.4 ± 2.6	4108.3	5.7
$\chi_{c0}(3915)$	$2^{3}P_{0}$	0^{++}	3921.7 ± 1.9	18.8 ± 3.5	3561.7	9.2
$\chi_{c1}(3872)$	$2^{3}P_{1}$	1^{++}	3871.65 ± 0.06	1.19 ± 0.21	4120.9	6.4

Table 2: The $c\bar{c}$ mesons and their respective masses (M_{exp}) compared to the masses calculated in this work (M_{cal}) using parameters (9a). The last column incudes the comparison of M_{exp} with M_{calc} in percentages.

Name	$N^{2S+1}l_J$	J^{PC}	$M_{exp}(\text{MeV})$	$\Gamma({\rm MeV})$	$M_{cal}({ m MeV})$	$rac{M_{cal}-M_{exp}}{M_{exp}} (\%)$
$\eta_b(1S)$	$1^{1}S_{0}$	1^{-+}	9398.7 ± 2.0	10 ± 5	9417.1	0.2
$\Upsilon(1S)$	$1^{3}S_{1}$	1	9460.30 ± 0.26	$\approx 0.054 \pm 0.001$	9397.4	0.7
$\chi_{b0}(1P)$	$1^{3}P_{0}$	0^{++}	9859.44 ± 0.26	NA	9417.1	4.4
$\chi_{b1}(1P)$	$1^{3}P_{1}$	1^{++}	9892.78 ± 0.05	NA	9708.2	1.9
$\chi_{b2}(1P)$	$1^{3}P_{2}$	2^{++}	9912.21 ± 0.26	NA	9674.6	2.3
$\Upsilon(2S)$	$2^{3}S_{1}$	1	10023.26 ± 0.31	$pprox 0.031 \pm 0.003$	9814.6	2.1
$\Upsilon(3S)$	$2^{3}S_{1}$	$1^{}$	10355.2 ± 0.56	$\approx 0.020 \pm 0.002$	10102.4	2.4
$\Upsilon(4S)$	$2^{3}S_{1}$	1	10579.4 ± 1.2	20.5 ± 2.5	10358.4	1.3
$\Upsilon_2(1D)$	$1^{3}D_{2}$	2	10163.7 ± 1.4	NA	10062.4	1.0
$h_b(1P)$	$1^{1}P_{1}$	1^{+-}	9899.3 ± 0.8	NA	9713.0	1.9

Table 3: The $b\bar{b}$ mesons and their respective masses (M_{exp}) compared to the masses calculated in this paper (M_{cal}) using parameters (9b). The last column incudes the comparison of M_{exp} with M_{cal} in percentages.

Table 4: The Casimir coefficients for the various colour structures of two partons.

	qar q		qq		
Colour state	1	8	3	6	
κ_s	-4/3	1/6	-2/3	1/3	

Table 5: Casimir coefficients for the two cluster interaction.

	ММ		DD	
Colour state	MM_{11}	MM ₈₈	$DA_{3\bar{3}}$	DA ₆₆
κ_s	-4/3	-3	-4/3	-10/3

antiquark in a non-exotic meson [20]. The method of using diquarks in the description of hadronic bound states is not unique to tetraquarks as it can be used for baryons [21][22] and pentaquarks [23][24]. Using the numerical method described in previous section we will calculate masses of the diquarks and the non-singlet quarkantiquark states and use the obtained result to calculate masses of the tetraquarks composed of those pairs, or in other words, we treat the problem like a set of three different two-body problems. In this work we will use Columblike potential to describe boson exchange between clusters.

The Hamiltonian used to calculate masses of the tetraquarks is presented in Eq. (12a) with H_T representing contribution of kinetic energy and (V_{ij}) representing all the potential contributions including the spin contributions.

$$H = H_T + V_{(12)(34)} + + (m_1 + m_2 + V_{12}) + (m_3 + m_4 + V_{34})$$
(12a)

$$H_T = \frac{\hat{p}_{12}^2}{2\mu_{12}} + \frac{\hat{p}_{34}^2}{2\mu_{34}} + \frac{\hat{p}_{1234}^2}{2\mu_{1234}}$$
(12b)

$$\mu_{12} = \frac{m_1 m_2}{m_1 + m_2}, \quad \mu_{34} = \frac{m_3 m_4}{m_3 + m_4}$$

$$\mu_{1234} = \frac{\mu_{12} \mu_{34}}{\mu_{12} + \mu_{34}}$$
(12c)

$$\vec{r}_{12} = \vec{r}_1 - \vec{r}_2, \quad \vec{r}_{34} = \vec{r}_3 - \vec{r}_4$$

 $\vec{r}_{1234} = \vec{r}_{34} - \vec{r}_{12}$ (12d)

1	S=0		S=1			
$\kappa_s = \frac{1}{3}$	L=0	L=1	L=0		L=1	
	J= 0	J=1	J=1	J=0	J=1	J=2
N=1	2819.4	2929.4	2815.1	2997.3	2929.2	2912.0
N=2	3016.0	3100.7	3011.9	3188.6	3100.3	3084.1

Table 6: Masses of colour sextet diquarks for possible combinations of their quantum numbers.

Table 7: Masses of colour triplet diquarks for possible combinations of their quantum numbers.

$\kappa_s = -\frac{2}{3}$	S=0		S=1			
	L=0	L=1	L=0	L=1		
	J= 0	J=1	J=1	J=0	J=1	J=2
N=1 N=2	2882.3 3226.2	3172.6 3503.8	2908.8 3245.6	3072.3 3263.1	3173.8 3503.2	3083.6 3363.5

$$\vec{R}_{12} = \frac{m_1 r_1}{m_1 + m_2} + \frac{m_2 r_2}{m_1 + m_2},$$

$$\vec{R}_{34} = \frac{m_3 r_3}{m_2 + m_4} + \frac{m_4 r_4}{m_2 + m_4}$$
(12e)

$$\vec{R}_{1234} = \frac{\mu_{12}r_{12}}{\mu_{12} + \mu_{34}} + \frac{\mu_{34}r_{34}}{\mu_{12} + \mu_{34}}$$
(12f)

One of the significant differences between classic meson and compact tetraquark is the fact that pairs are bosons, therefore we cannot use identity found in Eq. (6). Complete representation can be written as:

$$(2\otimes 2)\otimes(2\otimes 2) = (1\oplus 3)\otimes(1\oplus 3) = 1\oplus 3\oplus 3\oplus 1\oplus 3\oplus 5,$$
(13)

therefore for the pair-pair interaction in (1c):

$$\langle S_{12}S_{34} \rangle = \frac{1}{2} \langle S^2 - S_1^2 - S_2^2 \rangle =$$

$$= \begin{cases} 0, \quad S = |0\rangle, S_{12} = S_{34} = |0\rangle \\ 0, \quad S = |1\rangle, S_{12} \neq S_{34} \\ -2, \quad S = |0\rangle, S_{12} = S_{34} = |1\rangle \\ -1, \quad S = |1\rangle, S_{12} = S_{34} = |1\rangle \\ 1, \quad S = |2\rangle, S_{12} = S_{34} = |1\rangle \end{cases}$$

$$(14)$$

Because all of the resonances were identified in $di - J/\psi$ channel (except for X(7200) which was additionally identified in $J/\psi + \Psi(2S)$ [2]) we will assume that total angular momentum is L = 0, so V^{SL} (1d) and V^T (1e) can be neglected. Additional adjustment should be made to the α_s to include its scale dependence. Relation (15a) [25][26] with parameters (15b) and μ as the reduced mass of the system.

$$\alpha_s(\mu) = \frac{\alpha_0}{\ln(\frac{\mu^2 + \mu_0^2}{\Lambda_0^2})}$$
(15a)

$$\mu_0 \approx 0 \text{ MeV}, \quad \Lambda_0 = 0.112 \text{ MeV}, \quad \alpha_0 = 3.2524 \text{ (15b)}$$

For the purposes of this work we will use following naming scheme for potential tetraquark candidates that includes two-letter code, "MM" or "DA", with subscript denting possible colour configuration and three sets of quantum terms; one for total system, two for the pairs. An example of the usage this naming scheme can be found in Eq. (16.) The naming scheme can potentially be used to label wave functions for those states.

$$DA_{3\bar{3}}(2^{3}S_{1} \leftarrow 1^{1}L_{1} + 1^{1}L_{0}).$$
 (16)

	S=0		S=1			
$\kappa_s = \frac{1}{6}$	L=0	L=1	L=0		L=1	
	J= 0	J=1	J= 1	J=0	J=1	J=2
N=1 N=2	2705.2 2830.4	2777.3 2886.9	2704.0 2829.3	2808.8 2922.4	2777.2 2886.8	2768.6 2878.3

Table 8: Masses of colour octet quark-antiquark states for possible combinations of their quantum numbers.

Results with masses of the diquarks have been calculated with use of the code and placed in the Tables 6-8. Masses of the tetraquarks were calculated by choosing quantum numbers of the components, quantum numbers of their product and possible colour combination. The code allows to consider other types of interaction than the Cornell potential or to consider massive force carriers, however, we want to stay on safe ground concerning the heavy quarkonia spectroscopy. The exact results for the ground state (N = 0) and the first orbitral excitation (N = 1) can be for in the tables and energy level diagrams can be found in Figs 1-3.

4 Discussion of the results

We can clearly see that regardless of the composition of the tetraquark there are multiple candidates for for X(6900). All of the possible candidates are N = 2 states. Only sextet-antisextet states have suitable X(7200) candidates and all the suitable X(6900) candidates in that configuration have components with L = 0. Due to the use of a simple model one definite configuration cannot be suggested with certainty, however the closest lying candidate is $DA_{6\overline{6}}(2^3S_1 \leftarrow 1^3S_1 + 1^3S_1)$. In the previous section the subject of the wave function was omitted, as it held no significant impact on the calculation. However, we have to consider influence of the Pauli exclusion principle on the possibility of the existence of several resonances. Using the example shown in the Eq. (16) we will show the use of the notation we introduced to write down the wave function of the tetraquark:

$$|DA_{3\bar{3}}(2^{3}S_{1} \leftarrow 1^{1}L_{1} + 1^{1}L_{0})\rangle = |D_{\bar{3}}(1^{1}L_{1})\rangle \langle A_{3}(1^{1}L_{1})|,$$

$$|X_{c}(N^{2S+1}l_{J})\rangle = |Y_{l}^{m} \times X^{\sigma} \times X^{c} \times X^{f}\rangle,$$

(17)

The four parts of the wave wave function are the spherical harmonics Y_l^m , the spin wave function X^σ , the colour wave function X^c and the flavour wave function X^f . The flavour wave functions X^f of a diquark $|D\rangle$ or antiquark $|A\rangle$ is symmetric, unlike the the wave function for a meson or a meson-like state $|M\rangle$;

$$|X_D^f\rangle = |cc\rangle, \quad |X_A^f\rangle = |\bar{c}\bar{c}\rangle, \quad |X_M^f\rangle = |c\bar{c}\rangle.$$
 (18a)

The spin wave function is identical for any pair and the total spin wave function X_{S,S_z}^{σ} is dependent on the spin wave functions of the pairs:

$$\begin{split} |X_{0,0}^{\sigma}\rangle &= \frac{1}{\sqrt{2}}(|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle), \\ |X_{1,0}^{\sigma}\rangle &= \frac{1}{\sqrt{2}}(|\uparrow\downarrow\rangle + |\downarrow\uparrow\rangle), \\ |X_{1,1}^{\sigma}\rangle &= |\uparrow\uparrow\rangle, \qquad |X_{1,-1}^{\sigma}\rangle = |\downarrow\downarrow\rangle. \end{split}$$
(18b)

For the pairs only the $|X_{0,0}\rangle$ is antisymmetric. The colour wave functions for the pairs will not be shown explicitly for the sake of brevity, however it needs to be explicitly stated that $|X_3^c\rangle$ is fully antisymmetric while $|X_6^c\rangle$ is fully symmetric.

The wave function for the colour antisextet $1^{3}S_{1}$ diquark would have a wave function that would violate the Pauli exclusion principle, therefore any state with that subsystem is marked as forbidden. We have only two possible flavour wave functions for the tetraquark:

$$\begin{aligned} |X_{MM}^{f}\rangle &= |X_{M}^{f}\rangle\langle X_{M}^{f}| = |c\bar{c}c\bar{c}\rangle, \\ |X_{DA}^{f}\rangle &= |X_{D}^{f}\rangle\langle X_{A}^{f}| = |cc\bar{c}\bar{c}\rangle. \end{aligned}$$
(19a)

Assuming that the total spin of the tetraquark is known as well as the spin of the both pairs we can obtain the total of six possible total spin wave functions:

$$\begin{split} |X_{0,0}^{\sigma_{1}}\rangle &= |X_{0,0}^{\sigma}\rangle \langle X_{0,0}^{\sigma}| \\ |X_{0,0}^{\sigma_{2}}\rangle &= \frac{1}{\sqrt{3}} (|X_{1,1}^{\sigma}\rangle \langle X_{1,-1}^{\sigma}| + |X_{1-,1}^{\sigma}\rangle \langle X_{1,1}^{\sigma}| \\ &- |X_{1,0}^{\sigma}\rangle \langle X_{1,0}^{\sigma}|) \\ |X_{1,1}^{\sigma_{3}}\rangle &= |X_{1,1}^{\sigma}\rangle \langle X_{0,0}^{\sigma}| \qquad (19b) \\ |X_{1,1}^{\sigma_{4}}\rangle &= |X_{0,0}^{\sigma}\rangle \langle X_{1,1}^{\sigma}| \\ |X_{1,1}^{\sigma_{5}}\rangle &= \frac{1}{\sqrt{2}} (|X_{1,1}^{\sigma}\rangle \langle X_{1,0}^{\sigma}| + |X_{1,0}^{\sigma}\rangle \langle X_{1,1}^{\sigma}|) \\ &|X_{2,2}^{\sigma_{6}}\rangle &= |X_{1,1}^{\sigma}\rangle \langle X_{1,1}^{\sigma}|. \end{split}$$

There are only three total colour wave functions that we need to consider:

$$|X_{88}^c\rangle = |X_8^c\rangle \langle X_8^c|,$$

$$|X_{3\bar{3}}^c\rangle = |X_{\bar{3}}^c\rangle \langle X_3^c|, \quad |X_{6\bar{6}}^c\rangle = |X_6^c\rangle \langle X_{\bar{6}}^c|.$$
(19c)

For the L = 0 the spherical harmonic Y_0^0 is symmetric therefore $X^{\sigma} \times X^c \times X^f$ must be antisymmetric, therefore we can mark $DA_{3\bar{3}}(1^1S_0 \leftarrow 1^1S_0 + 1^1S_0)$ as forbidden.

One of the problems that we are attempting to solve in this work is finding the reason of the high mass of the X(6900) and the lack of more prominent lighter ground state. If we assume that the resonance has $DA_{6\bar{6}}$ structure we could easily explain lack of a prominent resonances closer to $di - J/\psi$ and $di - \eta_c$ mass threshold could be explained by them violating Pauli exclusion principle, while states with N = 2 could possibly be observed. Further inspection of the data suggest that X(6900) most likely is either 0^{-+} or 1^{--} state that have substructures with orbital angular momentum $L_{12} = L_{34} = 0$. No suggestion for X(7200) can be made, however if we assume that it has the same colour composition as X(6900) we could assume that it state with N = 2 and $L_{12} = L_{34} = 1$. The possibility of X(7200) being a N = 3 state was not evaluated due to a lack of precision during the calculation.



Figure 1: Energy level diagram for $DA_{3\bar{3}}$ all-charm tetraquark structure. Numerical data can be found in Table 9 and Table 10. Masses and widths of the resonances X(6900) and X(7200) were obtained by ATLAS with model A [2].

The other possible compositions of all-charm tetraquarks might be contributing to the broad structure. Since the background consists of multiple components and is possibly composed of several all-charm states with different quantum numbers, the confirmation of quantum numbers of X(6900) and the other structures would require extraordinary precision and the ability to separate data from multiple hadrons with identical content.

Assuming that the $bb\bar{b}\bar{b}$ would behave in the same manner as $cc\bar{c}\bar{c}$, we can calculate potential mass of allbottom tetraquark. Assuming that the composition of $DA_{6\bar{6}}(2^3S_1 \leftarrow 1^3S_1 + 1^3S_1)$ we archive:

$$m(bb\bar{b}\bar{b}) \approx 19.23 \text{ GeV}.$$
 (20)

Due to the high mass, it might be unlikely to confirm the existence of the resonance in the next few years.

Investigation of the additional decay channels may be prolific, in particular investigation of channel involving another exotic hadron and/or D mesons. Despite the lack



M(Mev) 7300 X(7200) 7200 7100 7000 $J/\psi + \chi_{c1}(3872)$ 6900 X(6900) 6800 6700 6600 6500 2S 6400 6300 1S 6200 1/w + 1/w6100 6000 5900 5800 5700 5600 5500 5400 5300 0 1 2-+ 5200

Figure 2: Energy level diagram for $DA_{6\bar{6}}$ all-charm tetraquark structure. Numerical data can be found in Table 9 and Table 10. Masses and widths of the resonances X(6900) and X(7200) were obtained by ATLAS with model A [2].

of significant data about some of the XYZ states, we have hypotheses regarding their internal dynamics, therefore gaining more information about possibility of the investigation of intermediate decays may possibly bring more information about tetraquark structure. Possible candidates for decay channels are $X(6900) \rightarrow \chi_{c1}(3872) + J/\psi$ or $X(6900) \rightarrow \chi_{c1}(3872) + \eta_c$. State $\chi_{c1}(3872)$ is one of the lowest-lying exotic states and potential identification of it in the decays may help with obtaining the threshold from sharp all-charm resonances, since mass of X(6900)is approximately equal to the sum of masses of the ground state charmonium and $\chi_{c1}(3872)$.

5 Conclusions

Assuming L = 0, the X(6900) might have sextetantisextet colour structure with N = 1 and possible terms 0^{-+} or 1^{--} , possibly a mixed state of both. The lack of

Figure 3: Energy level diagram for MM_{88} all-charm tetraquark structure. Numerical data can be found in Table 9 and Table 10. Masses and widths of the resonances X(6900) and X(7200) were obtained by ATLAS with model A [2].

a more prominent resonance with lower energy could be explained by Pauli exclusion principle. Some of the low-lying $DA_{6\bar{6}}$ resonances do not appear as sharp peaks due to the impact of the non-zero angular momenta and due to the large number of the states with similar energies, instead they contribute to the broad structures around 6200 MeV and 6500 MeV.

The octet-octet and triplet-antitriplet structures contribute to the background and to the low-lying broad structures. One could consider the possibility that the $DA_{3\bar{3}}$ structure is less prominent than $DA_{6\bar{6}}$ due to a higher likelihood of the formation of a doubly-charmed baryon and and its antiparticle instead of tightly bound tetraquark. More experimental data is needed to determine the quantum numbers of X(6900) and investigating the $J/\psi + \Psi(2S)$ invariant mass spectrum to more precisely determine the mass of X(7200) and the quantum numbers of the resonances. Additional decay channels involving the XYZ states, in particular $\chi_{c1}(3872)$ should



Figure 4: Energy level diagram for $DA_{6\bar{6}}$ and N = 2separated by orbital angular momentum of the substructures. Masses and widths of the resonances X(6900) and X(7200) were obtained by ATLAS with model A [2].

be considered.

5.1 Acknowledgement

I would like to express my gratitude to my supervisor, prof. dr hab. David Blaschke for the support during my thesis research and writing of this paper.

References

- [1] R. Aaij, et al., Sci. Bull. 65(23), 1983 (2020). DOI 10.1016/j.scib.2020.08.032
- [2] G. Aad, et al. Observation of an excess of dicharmonium events in the four-muon final state with the ATLAS detector (2022)
- 10.22323/1.414.0775

- [4] W. Lucha, F.F. Schoberl, Int. J. Mod. Phys. C 10, 607 (1999). DOI 10.1142/S0129183199000450
- [5] T. Gershon. Exotic hadron naming convention (2022)
- [6] W. Lucha, F.F. Schoberl, D. Gromes, Phys. Rept. **200**, 127 (1991). DOI 10.1016/0370-1573(91) 90001-3
- [7] K.G. Wilson, Phys. Rev. D 10, 2445 (1974). DOI 10.1103/PhysRevD.10.2445
- [8] H.A. Bethe, E.E. Salpeter, *Quantum Mechanics* of One- and Two-Electron Atoms (Springer Berlin, 1957). DOI 10.1007/978-3-662-12869-5
- [9] R.L. Workman, et al., PTEP 2022, 083C01 (2022). DOI 10.1093/ptep/ptac097
- [10] M. Gell-Mann, Phys. Lett. 8, 214 (1964). DOI 10. 1016/S0031-9163(64)92001-3
- [11] M. Gell-Mann. The Eightfold Way: A Theory of strong interaction symmetry (1961). DOI 10.2172/ 4008239
- [12] R.L. Jaffe, Phys. Rev. D 15, 267 (1977). DOI 10. 1103/PhysRevD.15.267
- [13] R.L. Jaffe, Phys. Rev. D 15, 281 (1977). DOI 10. 1103/PhysRevD.15.281
- [14] S.K. Choi, et al., Phys. Rev. Lett. 91, 262001 (2003). DOI 10.1103/PhysRevLett.91.262001
- [15] M. Anselmino, E. Predazzi, S. Ekelin, S. Fredriksson, D.B. Lichtenberg, Rev. Mod. Phys. 65, 1199 (1993). DOI 10.1103/RevModPhys.65.1199
- [16] A. Ali, L. Maiani, A.D. Polosa, Multiquark Hadrons, 1st edn. (Cambridge University Press, 2019). DOI 10.1017/9781316761465
- [17] G.S. Bali, Phys. Rev. D 62, 114503 (2000). DOI 10.1103/PhysRevD.62.114503
- [3] J. Zhang, K. Yi, PoS ICHEP2022, 775 (2022). DOI [18] S. Deldar, Phys. Rev. D 62, 034509 (2000). DOI 10.1103/PhysRevD.62.034509

- 1007/BF01585627
- [20] Progress in Particle and Nuclear Physics 116, 103835 (2021). DOI https://doi.org/10.1016/j.ppnp. 2020.103835.
- [21] D. Blaschke, A.S. Dubinin, D. Zablocki, PoS BaldinISHEPPXXII, 083 (2015). DOI 10.22323/1. 225.0083
- [22] J.L. Basdevant, S. Boukraa, Z. Phys. C 30, 103 (1986). DOI 10.1007/BF01560683

- [19] D. Gromes, Z. Phys. C 41, 427 (1988). DOI 10. [23] R.L. Jaffe, F. Wilczek, Phys. Rev. Lett. 91, 232003 (2003). DOI 10.1103/PhysRevLett.91.232003
 - [24] H.X. Chen, W. Chen, X. Liu, S.L. Zhu, Phys. Rept. 639, 1 (2016). DOI 10.1016/j.physrep.2016.05.004
 - [25] A. Deur, S.J. Brodsky, G.F. de Téramond, Progress in Particle and Nuclear Physics 90, 1 (2016). DOI https://doi.org/10.1016/j.ppnp.2016.04.003
 - [26] W. Buchmuller, S.H.H. Tye, Phys. Rev. D 24, 132 (1981). DOI 10.1103/PhysRevD.24.132

			$M_{12,34}$ (MeV)		
$(N^{2S+1}l_J)_{12}$	$(N^{2S+1}l_J)_{34}$	$(N^{2S+1}l_J)_{12,34}$	MM ₈₈	DA ₆₆	$DA_{3\bar{3}}$
$1^{1}S_{0}$	$1^{1}S_{0}$	$1^{1}S_{0}$	5794.3	5980.8	6114.0*
$1^{1}S_{0}$	$1^{3}S_{1}$	$1^{3}S_{1}$	5793.2	5977.1*	6139.3
$1^{3}S_{1}$	$1^{3}S_{1}$	$1^{1}S_{0}$	5395.0	5530.2*	6085.0
$1^{3}S_{1}$	$1^{3}S_{1}$	$1^{3}S_{1}$	5607.5	5765.4*	6127.4
$1^{3}S_{1}$	$1^{3}S_{1}$	1^5S_2	5946.2	6150.1*	6197.0
$1^{1}L_{1}$	$1^{1}L_{1}$	$1^{1}S_{0}$	5924.2	6182.0	6673.8
$1^{1}L_{1}$	$1^{3}L_{0}$	$1^{3}S_{1}$	5956.4	6243.0	6577.4
$1^{1}L_{1}$	$1^{3}L_{1}$	$1^{3}S_{1}$	5924.2	6177.4	6674.9
$1^{1}L_{1}$	$1^{3}L_{2}$	$1^{3}S_{1}$	5916.3	6162.0	6617.8
$1^{3}L_{0}$	$1^{3}L_{0}$	$1^{1}S_{0}$	5613.4	5904.9	6406.3
$1^{3}L_{0}$	$1^{3}L_{0}$	$1^{3}S_{1}$	5811.3	6115.2	6445.2
$1^{3}L_{0}$	$1^{3}L_{0}$	1^5S_2	6131.8	6467.8	6510.1
$1^{3}L_{1}$	$1^{3}L_{1}$	$1^{1}S_{0}$	5542.9	5760.2	6606.4
$1^{3}L_{1}$	$1^{3}L_{1}$	$1^{3}S_{1}$	5746.2	5980.6	6643.2
$1^{3}L_{1}$	$1^{3}L_{1}$	1^5S_2	6074.1	6346.6	6705.6
$1^{3}L_{2}$	$1^{3}L_{2}$	$1^{1}S_{0}$	5525.4	5725.7	6428.4
$1^{3}L_{2}$	$1^{3}L_{2}$	$1^{3}S_{1}$	5729.8	5948.2	6467.1
$1^{3}L_{2}$	$1^{3}L_{2}$	1^5S_2	6058.9	6316.9	6531.7
$1^{3}L_{0}$	$1^{3}L_{1}$	$1^{3}S_{1}$	5781.3	6051.0	6507.3
$1^{3}L_{0}$	$1^{3}L_{2}$	1^5S_2	6096.6	6393.9	6456.2
$1^{3}L_{1}$	$1^{3}L_{2}$	$1^{3}S_{1}$	5738.0	5964.4	6555.9
$1^{3}L_{1}$	$1^{3}L_{2}$	1^5S_2	6066.5	6331.7	6618.9

Table 9: Possible masses of the all-charm tetraquark for N = 1. States marked with * are forbidden by the Pauli exclusion principle.

			<i>M</i> _{12,34} (MeV)		
$(N^{2S+1}l_J)_{12}$	$(N^{2S+1}l_J)_{34}$	$(N^{2S+1}l_J)_{12,34}$	MM ₈₈	$DA_{6\bar{6}}$	$DA_{3\bar{3}}$
$1^{1}S_{0}$	$1^{1}S_{0}$	$2^{1}S_{0}$	6666.5	6931.5	6558.9
$1^{1}S_{0}$	$1^{3}S_{1}$	$2^{3}S_{1}$	6665.5	6927.8	6583.7
$1^{3}S_{1}$	$1^{3}S_{1}$	$2^{1}S_{0}$	6550.2	6801.5	6573.7
$1^{3}S_{1}$	$1^{3}S_{1}$	$2^{3}S_{1}$	6605.0	6860.6	6591.0
$1^{3}S_{1}$	$1^{3}S_{1}$	$2^{5}S_{2}$	6728.6	6992.9	6626.1
$1^{1}L_{1}$	$1^{1}L_{1}$	$2^{1}S_{0}$	6793.5	7127.0	7108.4
$1^{1}L_{1}$	$1^{3}L_{0}$	$2^{3}S_{1}$	6823.4	7187.0	7013.5
$1^{1}L_{1}$	$1^{3}L_{1}$	$2^{3}S_{1}$	6793.4	7124.4	7109.5
$1^{1}L_{1}$	$1^{3}L_{2}$	$2^{3}S_{1}$	6785.8	7109.2	7110.0
$1^{3}L_{0}$	$1^{3}L_{0}$	$2^{1}S_{0}$	6744.3	7136.0	6885.9
$1^{3}L_{0}$	$1^{3}L_{0}$	$2^{3}S_{1}$	6795.8	7189.8	6901.8
$1^{3}L_{0}$	$1^{3}L_{0}$	$2^{5}S_{2}$	6911.2	7308.1	6933.9
$1^{3}L_{1}$	$1^{3}L_{1}$	$2^{1}S_{0}$	6683.8	7008.6	7080.5
$1^{3}L_{1}$	$1^{3}L_{1}$	$2^{3}S_{1}$	6736.5	7064.6	7095.4
$1^{3}L_{1}$	$1^{3}L_{1}$	$2^{5}S_{2}$	6854.8	7188.4	7125.8
$1^{3}L_{2}$	$1^{3}L_{2}$	$2^{1}S_{0}$	6668.0	6977.4	6907.5
$1^{3}L_{2}$	$1^{3}L_{2}$	$2^{3}S_{1}$	6720.9	7033.8	6923.2
$1^{3}L_{2}$	$1^{3}L_{2}$	$2^{5}S_{2}$	6839.7	7158.8	6955.2
$1^{3}L_{0}$	$1^{3}L_{1}$	$2^{3}S_{1}$	6767.3	7128.6	6983.7
$1^{3}L_{0}$	$1^{3}L_{2}$	$2^{5}S_{2}$	6876.3	7234.5	6912.5
$1^{3}L_{1}$	$1^{3}L_{2}$	$2^{3}S_{1}$	6728.7	7049.2	7009.8
$1^{3}L_{1}$	$1^{3}L_{2}$	2^5S_2	6847.3	7173.6	7040.8

Table 10: Possible masses of the all-charm tetraquark for N = 2.