Induced Lorentz Violation on a Moving Braneworld

Daniel Kabat^{1,3[∗](#page-0-0)}, Marcelo Nomura^{2,3[†](#page-0-1)}

¹Department of Physics and Astronomy Lehman College, City University of New York 250 Bedford Park Blvd. W, Bronx, NY 10468, USA

²Physics Department City College of the City University of New York 160 Convent Avenue, New York, NY 10031, USA

³Graduate School and University Center, City University of New York 365 Fifth Avenue, New York, NY 10016, USA

We consider a braneworld scenario in which a flat 4-D brane, embedded in $M^{3,1} \times S^1$, is moving on or spiraling around the $S¹$. Although the induced metric on the brane is 4-D Minkowski, the would-be Lorentz symmetry of the brane is broken globally by the compactification. As recently pointed out this means causal bulk signals can propagate superluminally and even backwards in time according to brane observers. Here we consider the effective action on the brane induced by loops of bulk fields. We consider a variety of self-energy and vertex corrections due to bulk scalars and gravitons and show that bulk loops with non-zero winding generate UV-finite Lorentz-violating terms in the 4-D effective action. The results can be accommodated by the Standard Model Extension, a general framework for Lorentz-violating effective field theory.

[∗]daniel.kabat@lehman.cuny.edu

[†]mail@marcelonomura.com

Contents

1 Introduction

The simplest braneworld scenario posits a spacetime of the form $M^{3,1} \times S^1$, with a single extra dimension compactified on a circle of radius R . The brane, assumed to be at a fixed position on the S^1 , has a Minkowski metric induced on its worldvolume. In this scenario worldvolume Lorentz invariance is an exact symmetry, inherited from a symmetry of the underlying 5-D spacetime.

A straightforward generalization of this scenario allows the brane to either move on or spiral around the $S¹$. This generalization might seem quite innocuous. The induced metric on the brane is still 4-D Minkowski, so it would seem that brane observers might be hard-pressed to find any evidence that their brane has been boosted or rotated into the compact direction.

From a different perspective, however, the effects of this generalization are quite dramatic. Compactification on S^1 preserves an $SO(3,1)$ symmetry that acts on the directions orthogonal to the $S¹$. Once the brane is moving on or spiraling around the $S¹$ this exact $SO(3, 1)$ symmetry no longer aligns with the would-be Lorentz symmetry

of the brane worldvolume. Although there is no local indication of the violation, worldvolume Lorentz invariance is broken globally by the compactification. Without Lorentz invariance all bets are off, and indeed $\left[1, 2, 3\right]$ $\left[1, 2, 3\right]$ $\left[1, 2, 3\right]$ showed that bulk signals can propagate faster than light and even backwards in time according to brane observers. Fortunately causality – a more robust feature – remains intact, inherited from the causality of the underlying 5-D spacetime.

Here we consider the effects of virtual bulk particles in this generalized braneworld scenario. Such particles can leave a moving or rotated brane, propagate around the compact dimension, and return. Bulk loops have no reason to respect worldvolume Lorentz symmetry and might be expected to induce Lorentz-violating terms in the brane effective action. We will see that this is indeed the case. We focus on Lorentz-violating dimension-4 terms in the effective action, especially the electron self-energy and the electron – photon vertex, and show that bulk loops induce specific Lorentz-violating terms with finite, calculable coefficients. These terms are part of the Standard Model Extension, a general framework for Lorentz-violating effective field theories developed in [\[4,](#page-29-0) [5\]](#page-29-1). There are stringent experimental bounds on the Lorentz-violating coefficients which have been tabulated in [\[6\]](#page-29-2).

An outline of this paper is as follows. In section [2](#page-2-0) we describe the braneworld scenario we will consider. Section [3](#page-6-0) discusses the propagator for a bulk scalar field. In sections [4](#page-9-0) and [5](#page-13-0) we evaluate corrections to the electron self-energy and the electron – photon vertex due to a bulk scalar loop. Section [6](#page-15-0) considers the self-energy for a scalar field on the brane induced by a bulk scalar. Sections [7](#page-17-0) and [8](#page-20-0) evaluate the electron and scalar self-energy induced by a bulk graviton loop. We conclude in section [9](#page-23-0) with a discussion of experimental bounds and future directions.

Cosmological implications of this scenario have been studied in [\[7\]](#page-29-3) and a different approach to braneworld Lorentz violation has been developed in [\[8\]](#page-29-4).

2 Boosted and rotated branes

Consider a 5-D spacetime $\mathcal{M}^{3,1} \times S^1$. To describe this we begin from 5-D Minkowski space $\mathcal{M}^{4,1}$ with coordinates and metric

$$
X^{M} = (X^{\mu}, Z) \qquad M = 0, ..., 4 \qquad \mu = 0, ..., 3
$$

\n
$$
\eta_{MN} = \text{diag}(+ - \cdots -)
$$
\n(1)

We obtain an S^1 by periodically identifying the Z coordinate, $Z \approx Z + 2\pi R$. It's convenient to describe this identification as

$$
X^{M} \approx X^{M} + A^{M} \qquad A^{M} = (0, 0, 0, 0, 2\pi R) \tag{2}
$$

These upper-case coordinates define the preferred frame for the compactification, with an exact $SO(3,1)$ symmetry that acts on the coordinates X^{μ} .

The standard braneworld scenario would be to place a 4-D braneworld at rest at $Z = 0$. We are instead interested in braneworld which is moving in the Z direction and $/$ or has been rotated into the Z direction. To describe this we transform to a new frame with lower-case coordinates x^M via

$$
x^M = L^M{}_N X^N \tag{3}
$$

Here $L^M{}_N$ is an $SO(4,1)$ transformation that acts non-trivially on the Z coordinate. In the x^M coordinates there is a boosted and $/$ or rotated identification

$$
x^M \approx x^M + a^M \qquad a^M = L^M{}_N A^N \tag{4}
$$

We set

$$
x^M = (x^\mu, z) \tag{5}
$$

and imagine a braneworld at $z = 0$. The coordinates x^M can be thought of as co-moving and / or co-rotated with the brane. Since all we've done is a 5-D Lorentz transformation, in the co-moving coordinates the metric still has the form

$$
ds^2 = \eta_{\mu\nu} dx^{\mu} dx^{\nu} - dz^2
$$
\n(6)

The compactification is hidden in the identification [\(4\)](#page-3-0). Thus the induced metric on the brane is 4-D Minkowski, however the $SO(3,1)$ symmetry of the brane metric does not align with the $SO(3, 1)$ symmetry that is preserved by the compactification [\(2\)](#page-2-1). Instead worldvolume Lorentz symmetry is broken globally by the compactification, which leads to the curious possibilities of superluminal and even backwards-in-time signaling explored in $[1, 2, 3]$ $[1, 2, 3]$ $[1, 2, 3]$.

From the brane point of view it's natural to decompose a^M into components tangent and normal to the brane, so we set

$$
a^M = (a^\mu, 2\pi r) \tag{7}
$$

 a^{μ} becomes a preferred 4-vector on the brane, which shows that 4-D Lorentz symmetry on the brane is spontaneously broken. The fifth component $2\pi r$ is a scalar on the brane. In the calculations below we will find it useful to work with the combination

$$
b^{\mu} = \frac{a^{\mu}}{2\pi r} \tag{8}
$$

Up to Lorentz transformations on the brane there are three cases to consider.

1. Timelike b^{μ}

In this case we can go to a reference frame on the brane in which b^{μ} only has a time component. This can be obtained directly from [\(2\)](#page-2-1) by boosting with velocity β in the Z direction.

$$
\begin{pmatrix} T \\ Z \end{pmatrix} = \begin{pmatrix} \gamma & \gamma \beta \\ \gamma \beta & \gamma \end{pmatrix} \begin{pmatrix} t \\ z \end{pmatrix}
$$
 (9)

This leads to

$$
a^{\mu} = (-\gamma \beta 2\pi R, 0, 0, 0) \qquad r = \gamma R \tag{10}
$$

Note that

$$
b^{\mu} = (-\beta, 0, 0, 0) \tag{11}
$$

with

$$
b^2 = \beta^2 \in (0, 1) \tag{12}
$$

or alternatively

$$
1 - b^2 = \frac{1}{\gamma^2} \in (0, 1)
$$
\n(13)

This corresponds to the "boost-like isotropic" case discussed in [\[3\]](#page-28-2). As seen on the brane, bulk signals propagate isotropically in all directions at superluminal speeds.

2. Spacelike b^{μ}

In this case we can go to a reference frame on the brane in which b^{μ} only has an x component. This can be obtained directly from [\(2\)](#page-2-1) by rotating through an angle θ in the XZ plane.

$$
\begin{pmatrix}\nX \\
Z\n\end{pmatrix} = \begin{pmatrix}\n\cos \theta & -\sin \theta \\
\sin \theta & \cos \theta\n\end{pmatrix} \begin{pmatrix}\nx \\
z\n\end{pmatrix}
$$
\n(14)

This leads to

$$
a^{\mu} = (0, \sin \theta \, 2\pi R, 0, 0) \qquad r = \cos \theta R \tag{15}
$$

Note that

$$
b^{\mu} = (0, \tan \theta, 0, 0) \tag{16}
$$

with

$$
b^2 = -\tan^2 \theta \in (-\infty, 0)
$$
 (17)

or alternatively

$$
1 - b^2 = \frac{1}{\cos^2 \theta} \in (1, \infty)
$$
\n⁽¹⁸⁾

This corresponds to the "tilt-like anisotropic" case discussed in [\[3\]](#page-28-2). As seen on the brane, bulk signals propagate superluminally in the x direction and at the speed of light in perpendicular directions.

3. Lightlike b^{μ}

Finally we consider the case of lightlike b^{μ} .^{[1](#page-5-0)} This can be obtained starting from [\(2\)](#page-2-1) by making a Lorentz rotation in the $X-Z$ plane.^{[2](#page-5-1)} Here we've introduced light-front coordinates $X^{\pm} = T \pm X$ with

$$
ds^{2} = dX^{+}dX^{-} - |d\mathbf{Y}|^{2} - dZ^{2}
$$
\n(19)

The form of the Lorentz transformation is a little unfamiliar. Introducing a parameter $\lambda \in \mathbb{R}$ it takes the form

$$
\begin{pmatrix}\nT \\
X \\
Z\n\end{pmatrix} = \begin{pmatrix}\n1 + \frac{1}{2}\lambda^2 & \frac{1}{2}\lambda^2 & \lambda \\
-\frac{1}{2}\lambda^2 & 1 - \frac{1}{2}\lambda^2 & -\lambda \\
\lambda & \lambda & 1\n\end{pmatrix} \begin{pmatrix}\nt \\
x \\
z\n\end{pmatrix}
$$
\n(20)

or equivalently

$$
\begin{pmatrix} X^+ \\ X^- \\ Z \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ \lambda^2 & 1 & 2\lambda \\ \lambda & 0 & 1 \end{pmatrix} \begin{pmatrix} x^+ \\ x^- \\ z \end{pmatrix}
$$
 (21)

To see that this is the appropriate Lorentz transformation note that it leaves X^+ invariant, $X^+ = x^+$, so it acts on $X^- Z$ planes. Also it preserves the metric [\(19\)](#page-5-2), with $ds^2 = dx^+dx^- - |dy|^2 - dz^2$. Applying the (inverse of) the transformation [\(20\)](#page-5-3) gives

$$
a^{\mu} = (-\lambda 2\pi R, \lambda 2\pi R, 0, 0) \qquad r = R \tag{22}
$$

So the radius is unchanged, while

$$
b^{\mu} = (-\lambda, \lambda, 0, 0) \tag{23}
$$

is indeed a null vector on the brane.

A null vector has no invariant length, so one can go to an infinitely-boosted frame in which $b^{\mu} = 0$. This restores conventional Lorentz invariance on the brane. However if any matter (e.g. CMB photons) is present on the brane one may not wish to perform an infinite boost. In section [3](#page-6-0) we show that when λ is non-zero bulk signals can have a negative light-front velocity in the x^- direction. With respect to Minkowski time this means that as seen on the brane a bulk signal can travel faster than light and even backwards in time in the x direction. For further discussion of the geometry of this case see appendix [A.](#page-26-0)

Note that in all three cases we have $b^2 < 1$. The range $-\infty < b^2 < 0$ is tilt-like, $b^2 = 0$ is null, and $0 < b^2 < 1$ is boost-like. Alternatively we can say that we always have $1 - b^2 > 0$.^{[3](#page-5-4)} The range $0 < 1 - b^2 < 1$ is boost-like, $1 - b^2 = 1$ is null, and $1 < 1 - b^2 < \infty$ is tilt-like.

 1 We are grateful to Alexios Polychronakos and Massimo Porrati for bringing up this possibility.

²The case of a rotation in the $X^+ Z$ plane proceeds similarly.

³This is important for convergence of the loop integrals we encounter below.

3 Bulk scalar propagator

We expect that bulk loops should induce Lorentz-violating terms on the brane. Before turning to explicit calculations we start with a discussion of the bulk propagator. We focus on bulk scalar fields for simplicity.

The retarded propagator for a bulk field was discussed in [\[1\]](#page-28-0) while the static Green's function was studied in [\[9\]](#page-29-5). Here we consider the Feynman propagator. It's straightforward to impose the appropriate periodicity $(x^{\mu}, z) \approx (x^{\mu} + a^{\mu}, z + 2\pi r)$ using a winding sum (equivalently, a sum over image charges). In position space this gives the propagator for a bulk scalar of mass μ as

$$
\Delta = \sum_{w=-\infty}^{\infty} \int \frac{d^4k}{(2\pi)^4} \int \frac{dq}{2\pi} \frac{i}{k^2 - q^2 - \mu^2 + i\epsilon} e^{-ik \cdot (x - wa)} e^{iq(z - 2\pi rw)} \tag{24}
$$

It's convenient to shift

$$
q \to q + \frac{k \cdot a}{2\pi r} = q + k \cdot b \tag{25}
$$

so that

$$
\Delta = \sum_{w=-\infty}^{\infty} \int \frac{d^4k}{(2\pi)^4} \int \frac{dq}{2\pi} \frac{i}{k^2 - (q + k \cdot b)^2 - \mu^2 + i\epsilon} e^{-ik \cdot x} e^{i(q + k \cdot b)z} e^{-iq2\pi rw}
$$
(26)

We set $z = 0$ since we will only be interested in bulk propagation that starts and ends on the brane. Also we work in momentum space along the brane, which amounts to dropping $\int \frac{d^4k}{(2\pi)}$ $\frac{d^4k}{(2\pi)^4}e^{-ik\cdot x}$. Then we are left with the winding-sum form for the bulk propagator,

$$
\Delta = \sum_{w=-\infty}^{\infty} \int \frac{dq}{2\pi} \frac{i}{k^2 - (q + k \cdot b)^2 - \mu^2 + i\epsilon} e^{-iq2\pi rw}
$$
\n(27)

We can switch from a sum over windings to a sum over Kaluza-Klein momentum using the Poisson resummation identity

$$
\sum_{w} \int \frac{dq}{2\pi} f(q) e^{-iq2\pi rw} = \frac{1}{2\pi r} \sum_{n=-\infty}^{\infty} f\left(\frac{n}{r}\right)
$$
 (28)

This puts the bulk propagator in the form

$$
\Delta = \frac{1}{2\pi r} \sum_{n=-\infty}^{\infty} \frac{i}{k^2 - (k \cdot b + \frac{n}{r})^2 - \mu^2 + i\epsilon}
$$
\n(29)

It's clear that $b^{\mu} \neq 0$ breaks 4-D Lorentz invariance. We can look for poles in the propagator and read off the dispersion relation for the Kaluza-Klein tower, to see how it is modified from the perspective of a moving or rotated brane [\[9,](#page-29-5) [3\]](#page-28-2). There are three cases to consider.

1. Timelike b^{μ}

In this case we set $b^{\mu} = (-\beta, 0, 0, 0)$ and $k^{\mu} = (\omega, \mathbf{k})$. The propagator has poles at

$$
\omega = \gamma \left[\frac{\beta \gamma n}{r} \pm \sqrt{\frac{\gamma^2 n^2}{r^2} + |\mathbf{k}|^2 + \mu^2} \right] \mp i\epsilon \qquad n \in \mathbb{Z}
$$
 (30)

One branch of solutions has $\omega > 0$ and a pole that is displaced slightly below the real axis. The other branch has $\omega < 0$ and a pole that is displaced slightly above the real axis. Although we don't have 4-D Lorentz invariance, the poles are displaced in the standard way that allows for a Wick rotation to Euclidean signature. One can check that there are no tachyons from a 4-D perspective, $\omega^2 - |\mathbf{k}|^2 \geq 0$. Finally we can evaluate the group velocity

$$
v_g = \frac{d\omega}{dk} = \frac{\gamma |\mathbf{k}|}{\sqrt{\left(\frac{\gamma n}{r}\right)^2 + |\mathbf{k}|^2 + \mu^2}}\tag{31}
$$

This makes it clear that wave propagation is isotropic, with a velocity $0 \le v_g < \gamma$ that exceeds the speed of light if $|\mathbf{k}|$ is sufficiently large.

2. Spacelike b^{μ}

In this case we set $b^{\mu} = (0, \tan \theta, 0, 0)$ and $k^{\mu} = (\omega, k_x, \mathbf{k}_{\perp})$. The propagator has poles at

$$
\omega = \pm \sqrt{\left(\frac{k_x}{\cos \theta} - \frac{n \sin \theta}{r}\right)^2 + |\mathbf{k}_\perp|^2 + \left(\frac{n \cos \theta}{r}\right)^2 + \mu^2} \mp i\epsilon \qquad n \in \mathbb{Z} \tag{32}
$$

Again one branch of solutions has $\omega > 0$ and a pole that is displaced slightly below the real axis, while the other branch has $\omega < 0$ and a pole that is displaced slightly above the real axis, so we can perform Wick rotation in the standard way. One can check that there are no tachyons from a 4-D perspective, $\omega^2 - k_x^2 - |\mathbf{k}_\perp|^2 \ge 0$. Finally the group velocity is anisotropic. For a wave propagating in the x direction

$$
v_{gx} = \frac{d\omega}{dk_x}\bigg|_{\mathbf{k}_{\perp}=0} = \frac{\left(\frac{k_x}{\cos\theta} - \frac{n\sin\theta}{r}\right)}{\cos\theta\sqrt{\left(\frac{k_x}{\cos\theta} - \frac{n\sin\theta}{r}\right)^2 + \left(\frac{n\cos\theta}{r}\right)^2 + \mu^2}}
$$
(33)

while for a wave propagating in one of the perpendicular directions

$$
v_{g\perp} = \frac{d\omega}{dk_{\perp}}\bigg|_{k_x=0} = \frac{|k_{\perp}|}{\sqrt{|k_{\perp}|^2 + \left(\frac{n}{r}\right)^2 + \mu^2}}
$$
(34)

In the perpendicular directions we have the familiar group velocity for a Kaluza-Klein tower of particles with masses $\mu_n^2 = \left(\frac{n}{r}\right)^2$ $(\frac{n}{r})^2 + \mu^2$. In the x direction we have a group velocity $0 \leq |v_{gx}| < \frac{1}{\cos \theta}$ $\frac{1}{\cos \theta}$ that exceeds the speed of light if $|k_x|$ is sufficiently large.

3. Lightlike b^{μ}

For the lightlike case we set $b^{\mu} = (-\lambda, \lambda, 0, 0)$. It's convenient to introduce lightfront coordinates on the brane.

$$
x^{\pm} = t \pm x \qquad k^{\pm} = \omega \pm k_x \tag{35}
$$

We'll interpret $\tau = x^+$ as light-front time and the conjugate momentum $k_+ = \frac{1}{2}$ $\frac{1}{2}k^{-}$ as light-front energy. The propagator has poles at

$$
k^{-} = \frac{1}{k^{+}} \left[\left(\lambda k^{+} - \frac{n}{r} \right)^{2} + |\mathbf{k}_{\perp}|^{2} + \mu^{2} \right] \tag{36}
$$

This fixes the dispersion relation. As usual there are two branches of solutions. Positive frequency modes have $k^+ > 0$ and $k^- > 0$, while negative frequency modes have $k^+ < 0$ and $k^- < 0$. Given a positive-frequency plane-wave solution

$$
e^{-i\left(\frac{1}{2}k^{-}x^{+}+\frac{1}{2}k^{+}x^{-}-\mathbf{k}_{\perp}\cdot\mathbf{x}_{\perp}\right)}\tag{37}
$$

a stationary-phase approximation lets us read off the group velocities with respect to light-front time.

$$
v^{-} = \frac{dx^{-}}{d\tau} = -\frac{\partial k^{-}}{\partial k^{+}} = \frac{|\mathbf{k}_{\perp}|^{2} + \left(\frac{n}{r}\right)^{2} + \mu^{2}}{\left(k^{+}\right)^{2}} - \lambda^{2}
$$
(38)

$$
\mathbf{v}_{\perp} = \frac{d\mathbf{x}_{\perp}}{d\tau} = \frac{1}{2} \frac{\partial k^{-}}{\partial \mathbf{k}_{\perp}} = \frac{\mathbf{k}_{\perp}}{k^{+}}
$$
(39)

In the transverse directions we have conventional light-front kinematics.^{[4](#page-8-0)} In the longitudinal direction there is a shift which allows the longitudinal velocity to be negative, $-\lambda^2 < v^- < \infty$. This means that in Minkowski coordinates bulk signals can travel faster than light and even backwards in time in the x direction. To see this note that in Minkowski coordinates a trajectory $x^- = -\lambda^2 x^+$ corresponds to

$$
x = \frac{1 + \lambda^2}{1 - \lambda^2} t \tag{40}
$$

The Minkowski velocity is superluminal for $0 < \lambda^2 < 1$. The velocity diverges at $\lambda^2 = 1$ and becomes negative for $\lambda^2 > 1$, which as in [\[2\]](#page-28-1) indicates that the signal is traveling backwards in time. For further discussion of this case see appendix [A.](#page-26-0)

⁴Although the transverse kinematics look conventional, with respect to Minkowski time the limiting transverse velocity is $\sqrt{1 + \lambda^2}$. See appendix [A.](#page-26-0)

 $\frac{1}{2}$ $\frac{1}{2}$ Figure 1: One-loop electron self-energy arising from a Yukawa coupling to a bulk scalar.

4 Electron self-energy

I he world-volume metric matted on the brane is 4-D Minkowski, even if the brane is
boosted or rotated in the Z direction. Particles that solely propagate on the brane are The world-volume metric induced on the brane is 4-D Minkowski, even if the brane is describe these "standard model" particles using an effective action with 4-D Lorentz symmetry. However particles that propagate in the bulk can leave the brane, travel not sensitive to the breaking of 4-D Lorentz invariance and it would be reasonable to around the compactification manifold, and return. Such particles notice the global breaking of 4-D Lorentz invariance by the compactification and loops of such particles should induce Lorentz-violating terms in the 4-D effective action.

Yukawa coupling to the electron. We describe the coupled system with the action to the electron self-energy. We imagine a real bulk scalar field χ of mass μ that has a Here we study this effect, beginning with the simple example of radiative corrections

$$
S = \int d^5x \left[\frac{1}{2} \partial_M \chi \partial^M \chi - \frac{1}{2} \mu^2 \chi^2 \right] + \int d^4x \left[\bar{\psi} \left(i \gamma^\mu \partial_\mu - m \right) \psi - \lambda \bar{\psi} \psi \chi \big|_{z=0} \right] \tag{41}
$$

Note that the coupling λ has units (mass)^{-1/2}. The diagram we wish to consider is shown in Fig. [1.](#page-9-1)

Our goal is to evaluate the diagram and expand in powers of the external momentum p. In this way we will make contact with the Standard Model Extension, a general effective field theory framework for Lorentz-violating effects developed in [\[4,](#page-29-0) [5\]](#page-29-1). The basic diagram is easy to write down. Writing the brane-to-brane bulk propagator with a sum over Kaluza-Klein momentum as in [\(29\)](#page-6-1) we have

$$
i\Sigma = \frac{\lambda^2}{2\pi r} \sum_{n=-\infty}^{\infty} \int \frac{d^4k}{(2\pi)^4} \frac{k+m}{k^2 - m^2 + i\epsilon} \frac{1}{(k-p)^2 - ((k-p)\cdot b + \frac{n}{r})^2 - \mu^2 + i\epsilon} \tag{42}
$$

As pointed out in section [3,](#page-6-0) even though the bulk propagator is not Lorentz invariant, it still has poles that allow for a standard Wick rotation. So we Wick rotate in the usual way, setting

$$
k_E = (-ik^0, \mathbf{k}) \qquad d^4k = id^4k_E \qquad k^2 = -k_E^2 \tag{43}
$$

with a similar rotation for all other 4-vectors. We introduce a pair of Schwinger parameters s, t to represent the propagators via the identity

$$
\int_0^\infty ds \, e^{-As} = \frac{1}{A} \tag{44}
$$

It's convenient to use a frame in which only the first component of b_E is non-zero.

$$
b_E = (b_E, \mathbf{0})
$$

\n
$$
p_E = (p_E, \mathbf{p}_E)
$$

\n
$$
k_E = (k_E, \mathbf{k}_E)
$$

\n
$$
\gamma_E = (\gamma_E, \gamma_E)
$$
\n(45)

This leads to

$$
i\Sigma = \frac{i\lambda^2}{2\pi r} \sum_{n=-\infty}^{\infty} \int_0^{\infty} ds \int_0^{\infty} dt \int \frac{d^4k_E}{(2\pi)^4} \left(-\gamma_E k_E - \gamma_E \cdot \mathbf{k}_E + m \right) \exp\left[-s \left(k_E^2 + |\mathbf{k}_E|^2 + m^2 \right) \right]
$$

\n
$$
\exp\left\{ -t \left[(1 + b_E^2)(k_E - p_E)^2 - \frac{2n}{r} b_E (k_E - p_E) + |\mathbf{k}_E - \mathbf{p}_E|^2 + \left(\frac{n}{r} \right)^2 + \mu^2 \right] \right\}
$$
\n(46)

The momentum integrals are Gaussian and lead to the rather tedious expression

$$
i\Sigma = \frac{i\lambda^2}{32\pi^3 r} \sum_{n=-\infty}^{\infty} \int_0^{\infty} ds \int_0^{\infty} dt \frac{1}{(s+t)^{3/2} (s+t(1+b_E^2))^{1/2}}
$$

\n
$$
\left(-\frac{t(1+b_E^2)}{s+t(1+b_E^2)}\gamma_E p_E - \frac{t}{s+t} \gamma_E \cdot \mathbf{p}_E - \frac{t}{s+t(1+b_E^2)} \gamma_E b_E \frac{n}{r} + m\right)
$$

\n
$$
\exp\left\{-sm^2 - t\left(\left(\frac{n}{r}\right)^2 + \mu^2\right) - \frac{st}{s+t(1+b_E^2)} \left(p_E^2(1+b_E^2) + 2b_E p_E \frac{n}{r}\right) - \frac{st}{s+t} |\mathbf{p}_E|^2 + \frac{t^2}{s+t(1+b_E^2)} b_E^2 \left(\frac{n}{r}\right)^2\right\}
$$
(47)

Now we expand in powers of the external momentum p . At zeroth order, after continuing back to Lorentzian signature and restoring Lorentz covariance, we find

$$
i\Sigma^{(0)} = \frac{i\lambda^2}{32\pi^3 r} \sum_{n=-\infty}^{\infty} \int_0^{\infty} ds \int_0^{\infty} dt \frac{1}{(s+t)^{3/2} (s+t(1-b^2))^{1/2}} \left(\frac{t}{s+t(1-b^2)} \frac{n}{r} \not{b} + m\right)
$$

\n
$$
\exp\left\{-sm^2 - t\left(\left(\frac{n}{r}\right)^2 + \mu^2\right) - \frac{t^2 b^2}{s+t(1-b^2)} \left(\frac{n}{r}\right)^2\right\}
$$
(48)

The term proportional to m is Lorentz invariant and therefore not interesting to us. The term proportional to β has the potential to violate Lorentz invariance, but it vanishes once the sum over n is performed. This follows from a symmetry: the underlying expression [\(47\)](#page-10-0) is invariant under $b_E \rightarrow -b_E$, $n \rightarrow -n$ which implies that only even powers of b^{μ} can appear.

At first order in p^{μ} , after continuing back to Lorentzian signature and restoring Lorentz covariance, we find

$$
i\Sigma^{(1)} = \frac{i\lambda^2}{32\pi^{5/2}} \sum_{w=-\infty}^{\infty} \int_0^{\infty} ds \int_0^{\infty} dt \left(\frac{t^{1/2}}{(s+t)^3} \not{p} - \frac{2(\pi rw)^2 s}{t^{1/2}(s+t)^4} b_\mu b_\nu \gamma^\mu p^\nu \right)
$$

\n
$$
\exp\left\{-sm^2 - t\mu^2 - \frac{s + t(1-b^2)}{t(s+t)}(\pi rw)^2\right\} \tag{49}
$$

Here we've used the identity [\(28\)](#page-6-2) in reverse to replace the momentum sum with a winding sum and an integral over q. The integral over q is Gaussian and leads to (49) .

Working with a winding sum is advantageous for the following reasons.

- Lorentz symmetry is broken globally by the compactification. Particle trajectories with $w = 0$ are not sensitive to the breaking and are guaranteed to respect Lorentz invariance. Indeed in [\(49\)](#page-11-0) we see that the term with $w = 0$ is proportional to $\rlap{/}p$.
- Ultraviolet divergences can only arise from the $w = 0$ term in the sum, since nonzero winding means the loop can never shrink to a point. Indeed in [\(49\)](#page-11-0) we see that for $w \neq 0$ the exponential in the second line serves to cut off the short-distance regime $s, t \rightarrow 0$.

Since we are only interested in Lorentz-violating terms, we could simply discard the $w = 0$ term to obtain a finite result. However we might as well discard all terms proportional to $\rlap{/}p$. This means discarding the first term in parenthesis in [\(49\)](#page-11-0) as well as the trace part of $b_{\mu}b_{\nu}$. In this way we obtain the Lorentz-violating contribution

$$
i\Sigma_{LV}^{(1)} = -\frac{i\lambda^2 r^2}{16\pi^{1/2}} \left(b_\mu b_\nu - \frac{1}{4} \eta_{\mu\nu} b^2 \right) \gamma^\mu p^\nu \sum_{w=-\infty}^{\infty} \int_0^\infty ds \int_0^\infty dt \, \frac{sw^2}{t^{1/2} (s+t)^4}
$$

$$
\exp\left\{-sm^2 - t\mu^2 - \frac{s + t(1-b^2)}{t(s+t)} (\pi rw)^2 \right\} \tag{50}
$$

This corresponds to a Lorentz-violating term in the 4-D effective Lagrangian for the electron. In the notation of [\[5\]](#page-29-1) the relevant term is $\mathcal{L} = i c_{\mu\nu} \bar{\psi} \gamma^{\mu} \partial^{\nu} \psi$ which makes a contribution $ic_{\mu\nu}\gamma^{\mu}p^{\nu}$ to i Σ . Comparing to [\(50\)](#page-11-1) we can read off the Lorentz violating coefficient $c_{\mu\nu}$, which can be conveniently presented as

$$
c_{\mu\nu} = -\frac{1}{16\pi^2} \frac{\lambda^2}{\pi r} \left(b_{\mu} b_{\nu} - \frac{1}{4} \eta_{\mu\nu} b^2 \right) I_1 \tag{51}
$$

Figure 2: The quantity I_1 appearing in the electron self-energy as a function of b^2 .

where we've defined^{[5](#page-12-0)}

$$
I_n = \frac{1}{\sqrt{\pi}} \sum_{w=-\infty}^{\infty} \int_0^{\infty} ds \int_0^{\infty} dt \, \frac{s^n w^2}{t^{1/2} (s+t)^4} \exp\left\{-s(\pi mr)^2 - t(\pi \mu r)^2 - \frac{s + t(1-b^2)}{t(s+t)} w^2\right\}
$$
(52)

The induced coefficients $c_{\mu\nu}$ are real, dimensionless, traceless and symmetric. They make a Lorentz-violating but CPT-even contribution to the effective action.

We can think of [\(51\)](#page-11-2) as a product of a loop factor $\frac{1}{16\pi^2}$, a dimensionless coupling $\frac{\lambda^2}{\pi r}$ $\frac{\lambda^2}{\pi r},$ a tensor structure $b_{\mu}b_{\nu} - \frac{1}{4}$ $\frac{1}{4}\eta_{\mu\nu}b^2$, and a function I_1 of the dimensionless parameters b^2 , πmr , $\pi \mu r$. As can be seen in Fig. [2,](#page-12-1) I_1 is an increasing function of b^2 . It vanishes as $b^2 \to -\infty$ and (perhaps despite appearances) approaches a finite limit as $b^2 \to 1$.

The expression for I_1 simplifies if we set $m = 0$ (a massless fermion on the brane) and $b^2 = 0$ (a small boost and / or rotation). Then the sum and integrals can be performed

⁵We rescaled the Schwinger parameters, $s \to \pi^2 r^2 s$ and $t \to \pi^2 r^2 t$.

Figure 3: One-loop vertex correction due to a bulk scalar.

and the behavior for small and large μr can be extracted.^{[6](#page-13-1)} This leads to

$$
b^{2} \approx 0, m \approx 0: \quad I_{1} \approx \begin{cases} \frac{1}{6} \log \frac{1}{\mu r} & \text{as } \mu r \to 0\\ \frac{\pi}{3} \mu r e^{-2\pi \mu r} & \text{as } \mu r \to \infty \end{cases} \tag{53}
$$

5 Electron – photon vertex

Next we consider the one-loop correction to the electron – photon vertex due to a bulk scalar. The diagram is shown in Fig. [3.](#page-13-2)

Suppressing the external polarizations and writing the bulk propagator with a momentum sum as in [\(29\)](#page-6-1), the diagram is

$$
i\Gamma^{\mu} = -\frac{e\lambda^{2}}{2\pi r} \sum_{n=-\infty}^{\infty} \int \frac{d^{4}k}{(2\pi)^{4}} \frac{\rlap{\,/}{p_{2} + k} + m}{(p_{2} + k)^{2} - m^{2} + i\epsilon} \gamma^{\mu} \frac{\rlap{\,/}{p_{1} + k} + m}{(p_{1} + k)^{2} - m^{2} + i\epsilon} \frac{1}{k^{2} - (k \cdot b + \frac{n}{r})^{2} - \mu^{2} + i\epsilon} \tag{54}
$$

We can evaluate this similarly to the electron self-energy. We Wick rotate, introduce a

 $^6\rm{For}$ a massless fermion on the brane we must keep the bulk scalar mass μ non-zero to avoid an IR divergence.

series of Schwinger parameters

$$
\frac{1}{(p_{1E} + k_E)^2 + m^2} = \int_0^\infty ds_1 e^{-s_1[(p_{1E} + k_E)^2 + m^2]}
$$
\n
$$
\frac{1}{(p_{2E} + k_E)^2 + m^2} = \int_0^\infty ds_2 e^{-s_2[(p_{2E} + k_E)^2 + m^2]}
$$
\n
$$
\frac{1}{k_E^2 + (k_E \cdot b_E - \frac{n}{r})^2 + \mu^2} = \int_0^\infty dt \, e^{-t \left[k_E^2 + (k_E \cdot b_E - \frac{n}{r})^2 + \mu^2\right]},
$$
\n(55)

and evaluate the Gaussian integral over k_E^{μ} E ^{μ}. This gives

$$
i\Gamma^{\mu} = \frac{ie\lambda^2}{2\pi r} \sum_{n=-\infty}^{\infty} \int_0^{\infty} ds_1 \int_0^{\infty} ds_2 \int_0^{\infty} dt \frac{1}{16\pi^2} \sqrt{\det h} \, e^{-s_1(p_{1E}^2 + m^2) - s_2(p_{2E}^2 + m^2) - t(\frac{n^2}{r^2} + \mu^2)}
$$
\n
$$
e^{h_{\mu\nu}v^{\mu}v^{\nu}} \left[\frac{1}{2} h_{\alpha\beta} \gamma_E^{\alpha} \gamma_E^{\mu} \gamma_E^{\beta} + \left(p_{2E} \cdot \gamma_E - h_{\alpha\beta} v^{\alpha} \gamma_E^{\beta} - m \right) \gamma_E^{\mu} \left(p_{1E} \cdot \gamma_E - h_{\gamma\delta} v^{\gamma} \gamma_E^{\delta} - m \right) \right]
$$
\n(56)

where we've introduced the convenient notation

$$
h_{\mu\nu} = \text{diag.}\left(\frac{1}{s_1 + s_2 + t(1 + b_E^2)}, \frac{1}{s_1 + s_2 + t}, \frac{1}{s_1 + s_2 + t}, \frac{1}{s_1 + s_2 + t}\right)
$$

$$
v^{\mu} = s_1 p_{1E}^{\mu} + s_2 p_{2E}^{\mu} - t b_E^{\mu} \frac{n}{r}
$$
 (57)

Now we expand in powers of the external momenta. At leading (zeroth) order, after continuing back to Lorentzian signature, switching to a winding sum for the bulk propagator and doing a bit of Dirac algebra, we find

$$
i\Gamma^{(0)\mu} = ie\lambda^2 \sum_{w=-\infty}^{\infty} \int_0^{\infty} ds_1 \int_0^{\infty} ds_2 \int_0^{\infty} dt \frac{1}{(4\pi t)^{1/2} (4\pi (s_1 + s_2 + t))} e^{-s_1 m^2 - s_2 m^2 - t\mu^2}
$$

\n
$$
\exp\left\{-\frac{s_1 + s_2 + t(1 - b^2)}{t(s_1 + s_2 + t)} \pi^2 r^2 w^2\right\} \left[\frac{1}{s_1 + s_2 + t} \gamma^{\mu} + m^2 \gamma^{\mu} - \frac{(\pi r w)^2}{(s_1 + s_2 + t)^2} (2\beta b^{\mu} - b^2 \gamma^{\mu})\right]
$$
\n(58)

We drop all Lorentz-invariant terms, which includes the UV-divergent terms with $w = 0$. Setting $s = s_1 + s_2$ we're left with the Lorentz-violating contribution

$$
i\Gamma_{LV}^{(0)\mu} = -\frac{ie\lambda^2 r^2}{16\pi^{1/2}} \left(b_{\alpha}b_{\beta} - \frac{1}{4}\eta_{\alpha\beta}b^2\right)\gamma^{\alpha}\eta^{\beta\mu} \sum_{w=-\infty}^{\infty} \int_0^{\infty} ds \int_0^{\infty} dt \frac{sw^2}{t^{1/2}(s+t)^4}
$$

$$
\exp\left\{-sm^2 - t\mu^2 - \frac{s + t(1-b^2)}{t(s+t)}\pi^2r^2w^2\right\} \tag{59}
$$

Figure 4: One-loop scalar self-energy arising from a cubic coupling to a bulk scalar.

This pairs nicely with [\(50\)](#page-11-1) to produce a gauge-invariant but Lorentz-violating dimension-4 term in the effective action, namely

$$
\mathcal{L} = i c_{\mu\nu} \bar{\psi} \gamma^{\mu} D^{\nu} \psi \qquad D_{\mu} = \partial_{\mu} - ie A_{\mu} \tag{60}
$$

consistency check on our results. Expanding [\(56\)](#page-14-0) beyond zeroth order in the external The coefficient $c_{\mu\nu}$ is given in [\(51\)](#page-11-2). Since we stopped at zeroth order in the momentum this outcome, required by gauge invariance and Ward identities, can be thought of as a momenta would give higher-derivative corrections to the electron – photon vertex.

6 Scalar self-energy

Having calculated the one-loop Lorentz-violating correction to the self-energy of an electron, we now perform a similar calculation for a real scalar field ϕ on the brane with a cubic coupling to a bulk scalar χ . We start from the action

$$
S = \int d^5x \left[\frac{1}{2} \partial_M \chi \partial^M \chi - \frac{1}{2} \mu^2 \chi^2 \right] + \int d^4x \left[\frac{1}{2} \partial_\mu \phi \partial^\mu \phi - \frac{1}{2} m^2 \phi^2 - \frac{1}{2} g \phi^2 \chi \Big|_{z=0} \right] \tag{61}
$$

Note that the coupling g has units $(mass)^{+1/2}$. The diagram we wish to consider is shown in Fig. [4.](#page-15-1)

Writing the brane-to-brane bulk propagator with a sum over Kaluza-Klein momentum as in [\(29\)](#page-6-1), the diagram is

$$
\frac{g^2}{2\pi r} \sum_{n=-\infty}^{\infty} \int \frac{d^4k}{(2\pi)^4} \frac{1}{k^2 - m^2 + i\epsilon} \frac{1}{(k-p)^2 - ((k-p)\cdot b + \frac{n}{r})^2 - \mu^2 + i\epsilon} \tag{62}
$$

We Wick rotate to Euclidean signature as in [\(43\)](#page-9-2) and introduce a pair of Schwinger parameters as in [\(44\)](#page-10-1). Parametrizing the Euclidean momenta as in [\(45\)](#page-10-2) and performing the Gaussian integral over k_E we find

$$
\frac{ig^2}{32\pi^3 r} \sum_{n=-\infty}^{\infty} \int_0^{\infty} ds \int_0^{\infty} dt \frac{1}{(s+t)^{3/2} (s+t(1+b_E^2))^{1/2}}
$$

\n
$$
\exp\left\{-sm^2 - t\left(\left(\frac{n}{r}\right)^2 + \mu^2\right) - \frac{st}{s+t(1+b_E^2)} \left(p_E^2(1+b_E^2) + 2b_E p_E \frac{n}{r}\right) - \frac{st}{s+t} |\mathbf{p}_E|^2 + \frac{t^2}{s+t(1+b_E^2)} b_E^2 \left(\frac{n}{r}\right)^2\right\}
$$
(63)

Now we expand in powers of the external momentum p. At zeroth order the result is Lorentz invariant and can be ignored. At first order the sum over Kaluza-Klein momentum vanishes because it is odd under $n \to -n$. At second order, after continuing back to Lorentzian signature and restoring Lorentz covariance, we find

$$
\frac{ig^2}{32\pi^{5/2}}\sum_{w=-\infty}^{\infty}\int_0^{\infty}ds\int_0^{\infty}dt\left(\frac{st^{1/2}}{(s+t)^3}p^2-\frac{2(\pi rw)^2s^2}{t^{1/2}(s+t)^4}b_\mu b_\nu p^\mu p^\nu\right)
$$

\n
$$
\exp\left\{-sm^2-t\mu^2-\frac{s+t(1-b^2)}{t(s+t)}(\pi rw)^2\right\}
$$
\n(64)

Again we've used the identity [\(28\)](#page-6-2) in reverse to replace the momentum sum with a winding sum and an integral over q. The integral over q is Gaussian and leads to (64) . The first term in parenthesis is Lorentz invariant and can be dropped. The second term can be matched to a Lorentz-violating term in the effective action [\[5\]](#page-29-1)

$$
\mathcal{L} = \frac{1}{2} k_{\mu\nu} \partial^{\mu} \phi \partial^{\nu} \phi \tag{65}
$$

with a traceless coefficient $k_{\mu\nu}$. Removing the Lorentz-invariant trace from the second term in [\(64\)](#page-16-0) we identify

$$
k_{\mu\nu} = -\frac{1}{16\pi^2} g^2 \pi r \left(b_{\mu} b_{\nu} - \frac{1}{4} \eta_{\mu\nu} b^2 \right) I_2 \tag{66}
$$

where I_n is defined in [\(52\)](#page-12-2). The induced coefficients $k_{\mu\nu}$ are real, dimensionless, traceless and symmetric. They make a Lorentz-violating but CPT-even contribution to the effective action.

We can think of [\(66\)](#page-16-1) as a product of a loop factor $\frac{1}{16\pi^2}$, a dimensionless coupling $g^2 \pi r$, a tensor structure $b_\mu b_\nu - \frac{1}{4}$ $\frac{1}{4}\eta_{\mu\nu}b^2$, and a function I_2 of the dimensionless parameters b^2 , πmr , $\pi \mu r$. As can be seen in Fig. [5,](#page-17-1) I_2 is an increasing function of b^2 . It vanishes as $b^2 \to -\infty$ and approaches a finite limit as $b^2 \to 1$.

Figure 5: The quantity I_2 appearing in the scalar self-energy as a function of b^2 .

The expression for I_2 simplifies if we set $m = 0$ (a massless scalar on the brane) and $b^2 = 0$ (a small boost and / or rotation). Then the sum and integrals can be performed and the behavior for small and large μr can be extracted.^{[7](#page-17-2)} This leads to

$$
b^{2} \approx 0, m \approx 0: I_{2} \approx \begin{cases} \frac{1}{6\pi^{2}\mu^{2}r^{2}} & \text{as } \mu r \to 0\\ \frac{2}{3}e^{-2\pi\mu r} & \text{as } \mu r \to \infty \end{cases}
$$
(67)

7 Electron self-energy from a bulk graviton

The graviton is the most likely candidate for a bulk field. It also provides an interesting contrast to the bulk scalars we have considered so far, since it carries spin and has a non-renormalizeable coupling to the stress tensor on the brane. For these reasons we consider corrections to the electron self-energy induced by a bulk graviton loop. The diagram is shown in Fig. [6.](#page-18-0)

⁷For a massless scalar on the brane we must keep the bulk scalar mass μ non-zero to avoid an IR divergence.

Figure 6: Electron self-energy due to a bulk graviton loop. \mathcal{C} Figure self energy set \mathcal{C}

Bulk gravitons in the large extra dimension scenario $[10, 11, 12]$ $[10, 11, 12]$ $[10, 11, 12]$ have been considered in [\[13\]](#page-29-9) and we borrow several of their expressions. We expand the 5-D metric about flat space,

$$
g_{AB} = \eta_{AB} + \frac{2}{\overline{M}_5^{3/2}} h_{AB} \tag{68}
$$

3 Franch mass. The state to state graviton propagator w where M_5 is the 5-D reduced Planck mass. The brane-to-brane graviton propagator is

AB source- CD
$$
\frac{1}{2\pi r} \sum_{n=-\infty}^{\infty} \frac{i}{2} \frac{\eta_{AC} \eta_{BD} + \eta_{AD} \eta_{BC} - \frac{2}{3} \eta_{AB} \eta_{CD}}{k^2 - (k \cdot b + \frac{n}{r})^2 - \mu^2 + i\epsilon}
$$
(69)

 μ as an infrared regulator. The propagator is written in de Donder gauge, $\xi = 1$ in the
notation of [\[13\]](#page-29-9). We assume the graviton couples to the 4-D stress tensor on the brane, where κ is the 4-D momentum, κ is the Natuza-Niem momentum and we ve introduced μ as an infrared regulator. The propagator is written in de Donder gauge, $\xi = 1$ in the where k is the 4-D momentum, n is the Kaluza-Klein momentum and we've introduced Ĭ

$$
\mathcal{L} = \bar{\psi} \left(i \gamma^{\mu} \partial_{\mu} - m \right) \psi - \frac{1}{\overline{M}_{5}^{3/2}} T^{\mu \nu} h_{\mu \nu} \big|_{z=0} \tag{70}
$$
\n
$$
T_{\mu \nu} = \frac{i}{4} \bar{\psi} \left(\gamma_{\mu} \partial_{\nu} + \gamma_{\nu} \partial_{\mu} \right) \psi - \frac{i}{4} \left(\partial_{\mu} \bar{\psi} \gamma_{\nu} + \partial_{\nu} \bar{\psi} \gamma_{\mu} \right) \psi
$$

which leads to the vertex $\frac{1}{2}$

$$
\sum_{k_1}^{k_2} \text{where } \mu\nu = \frac{i}{4\overline{M}_5^{3/2}} \left[(k_1 + k_2)_{\mu} \gamma_{\nu} + (k_1 + k_2)_{\nu} \gamma_{\mu} \right] \tag{71}
$$

So far the motion of the brane has only entered in the graviton propagator [\(69\)](#page-18-1), in a manner exactly analogous to the scalar propagator [\(29\)](#page-6-1). However the motion of the brane also enters in the effective 4-D coupling. The Newtonian potential on a moving brane was studied by Greene et al. in [\[9\]](#page-29-5), who found that the relation between the 4-D and 5-D reduced Planck masses becomes 8

$$
\overline{M}_4 = (\gamma 2\pi R)^{1/2} \, \overline{M}_5^{3/2} \tag{72}
$$

For a moving brane $r = \gamma R$ so the reduced 4-D Planck mass is

$$
\overline{M}_4 = (2\pi r)^{1/2} \,\overline{M}_5^{3/2} = 2.4 \times 10^{18} \,\text{GeV} \tag{73}
$$

We take this relation to hold in general, i.e. even for a brane that is tilt-like rather than boost-like.

After all these preliminaries we are ready to evaluate the diagram in Fig. [6.](#page-18-0)[9](#page-19-1) The basic diagram is straightforward to write down.

$$
i\Sigma = \frac{1}{16\pi r \overline{M}_5^3} \sum_{n=-\infty}^{\infty} \int \frac{d^4k}{(2\pi)^4} \frac{(k+p)^2 \gamma_\mu (k+m)\gamma^\mu + \frac{1}{3}(k+p)(k+m)(k+p)}{(k^2 - m^2 + i\epsilon)\left((k-p)^2 - \left((k-p)\cdot b + \frac{n}{r}\right)^2 - \mu^2 + i\epsilon\right)}
$$
(74)

With some Dirac algebra the numerator can be simplified so it is at most linear in Dirac matrices. We Wick rotate, introduce Schwinger parameters, and perform the Gaussian integral over k_E . It is convenient to do this in a frame in which the Euclidean vectors have components

$$
b_E = (b_1, 0, 0, 0)
$$

\n
$$
p_E = (p_1, p_2, 0, 0)
$$

\n
$$
k_E = (k_1, k_2, k_3, k_4)
$$
\n(75)

Then we continue back to Lorentzian signature, restore Lorentz covariance, and expand in powers of p. At zeroth order all terms are either Lorentz invariant or vanish because they are odd under $n \to -n$. At first order in p, after switching from a momentum sum to a winding sum, we find that many terms are either Lorentz invariant or vanish because they are odd under $w \rightarrow -w$. Discarding all such terms we are left with a

⁹There is another self-energy diagram at one loop $\hat{\xi}$ $\hat{\xi}$ but it does not induce Lorentz violation on the brane.

⁸In [\[9\]](#page-29-5) this was expressed in terms of Newton's constant, $G_4 = G_5/\gamma 2\pi R$ with $G_4 = 1/8\pi \overline{M}_4^2$ and $G_5=1/8\pi \overline{M}_5^3$ $\frac{6}{5}$.

Lorentz-violating contribution to the effective Lagrangian, $\mathcal{L} = ic_{\mu\nu} \bar{\psi} \gamma^{\mu} \partial^{\nu} \psi$ where^{[10](#page-20-1)}

$$
c_{\mu\nu} = -\frac{1}{16\pi^2} \frac{1}{(\pi r \overline{M}_4)^2} \left(b_{\mu} b_{\nu} - \frac{1}{4} \eta_{\mu\nu} b^2 \right) I_{\text{gravity}}
$$
(76)

$$
I_{\text{gravity}} = \frac{1}{6\sqrt{\pi}} \sum_{w=1}^{\infty} \int_0^{\infty} ds \int_0^{\infty} dt \frac{9s^2 - 2st - 11t^2 + 5sb^2}{w^3 t^{1/2} (s+t)^6}
$$

$$
\exp \left\{ -s(\pi m r w)^2 - t(\pi \mu r w)^2 - \frac{s + t(1-b^2)}{t(s+t)} \right\}
$$

We have written [\(76\)](#page-20-2) as a product of

- a loop factor $\frac{1}{16\pi^2}$,
- a dimensionless coupling $\frac{1}{(\pi r \overline{M}_4)^2}$ built from the effective radius r and the reduced 4-D Planck mass M_4 ,
- a symmetric traceless tensor structure $b_{\mu}b_{\nu} \frac{1}{4}$ $\frac{1}{4}\eta_{\mu\nu}b^2,$
- a function I_{gravity} of the dimensionless parameters b^2 , πmr , $\pi \mu r$.

The function I_{gravity} is shown in Fig. [7.](#page-21-0) It simplifies if we set $m = 0$ (a massless fermion on the brane) and $b^2 = 0$ (a small boost and / or rotation). Then the sum and integrals can be performed and the behavior for large and small μ can be extracted. For graviton loops there is no IR divergence, even for a massless fermion on the brane, and we find

$$
b^2 \approx 0, m \approx 0: \quad I_{\text{gravity}} \approx \begin{cases} -\frac{1}{4}\zeta(3) & \text{as } \mu r \to 0\\ -\frac{1}{3}(\pi\mu r)^2 e^{-2\pi\mu r} & \text{as } \mu r \to \infty \end{cases} \tag{77}
$$

8 Scalar self-energy from a bulk graviton

Finally we consider corrections to the self-energy of a minimally-coupled scalar field due to a bulk graviton loop. We assume the graviton couples to the 4-D stress tensor on the brane,

$$
\mathcal{L} = \frac{1}{2} \partial_{\mu} \phi \partial^{\mu} \phi - \frac{1}{2} m^2 \phi^2 - \frac{1}{\overline{M}_5^{3/2}} T^{\mu \nu} h_{\mu \nu} \big|_{z=0} \tag{78}
$$
\n
$$
T_{\mu \nu} = \partial_{\mu} \phi \partial_{\nu} \phi - \frac{1}{2} \eta_{\mu \nu} \big(\partial_{\lambda} \phi \partial^{\lambda} \phi - m^2 \phi^2 \big)
$$

¹⁰We rescaled the Schwinger parameters, $s \to (\pi r w)^2 s$ and $t \to (\pi r w)^2 t$.

Figure 7: The quantity I_{gravity} appearing in the electron self-energy due to a bulk graviton loop. The function decreases rapidly but has a finite limit as $b^2 \to 1$.

Figure 8: Scalar self-energy due to a bulk graviton loop.

which leads to the scalar $-$ graviton vertex

$$
\frac{k_2}{\sqrt{\frac{1}{k_1}}} \left[k_1 \right]_{\mu\nu} = \frac{i}{\overline{M}_5^{3/2}} \left[k_1 \right]_{\mu\nu} k_2 \left[k_1 \right]_{\mu\nu} + k_1 \left[k_2 \right]_{\mu\nu} + k_2 \left[k_1 \right]_{\mu\nu} (k_1 \cdot k_2 - m^2) \right] \tag{79}
$$

graviton propagator [\(69\)](#page-18-1) the basic diagram is straightforward to write down. The diagram we wish to evaluate is shown in Fig. 8^{11} 8^{11} 8^{11} With the brane-to-brane

$$
i\Sigma = \frac{1}{4\pi r \overline{M}_5^3} \sum_{n=-\infty}^{\infty} \int \frac{d^4k}{(2\pi)^4} \frac{4k^2p^2 + \frac{4}{3}(k \cdot p)^2 + \frac{8}{3}m^2k \cdot p - \frac{8}{3}m^4}{(k^2 - m^2 + i\epsilon)\left((k - p)^2 - \left((k - p) \cdot b + \frac{n}{r}\right)^2 - \mu^2 + i\epsilon\right)} (80)
$$

the contractions of the stress tensors in the numerator. As is by now familiar we Wick Compared to the electron self-energy considered in section [7](#page-17-0) the main difference is in rotate, introduce Schwinger parameters, and perform the Gaussian integral over k_E . It is convenient to do this in a frame in which the Euclidean vectors have components

$$
b_E = (b_1, 0, 0, 0)
$$

\n
$$
p_E = (p_1, p_2, 0, 0)
$$

\n
$$
k_E = (k_1, k_2, k_3, k_4)
$$
\n(81)

the expression is Lorentz invariant. At first order in p the result vanishes because all Then we continue back to Lorentzian signature, restore Lorentz covariance, switch from a momentum sum to a winding sum, and expand in powers of p . At zeroth order in p terms are odd under $w \to -w$. At second order in p many of the terms are Lorentz

¹¹There is another self-energy diagram at one loop $\mathcal{E}_{\epsilon_{\alpha}}\mathcal{F}$ but it does not induce Lorentz violation on the brane.

invariant. Discarding all Lorentz-invariant terms we are left with a Lorentz-violating contribution to the effective Lagrangian, $\mathcal{L} = \frac{1}{2}$ $\frac{1}{2}k_{\mu\nu}\partial^{\mu}\phi\partial^{\nu}\phi$ where^{[12](#page-23-1)}

$$
k_{\mu\nu} = -\frac{1}{16\pi^2} \frac{1}{(\pi r \overline{M}_4)^2} \left(b_{\mu} b_{\nu} - \frac{1}{4} \eta_{\mu\nu} b^2 \right) I_{\text{gravity}}^{\text{scalar}} \tag{82}
$$

$$
I_{\text{gravity}}^{\text{scalar}} = \frac{4}{3\sqrt{\pi}} \sum_{w=1}^{\infty} \int_0^{\infty} ds \int_0^{\infty} dt \, \frac{1 + 4s(\pi m r w)^2 - 4s^2(\pi m r w)^4}{w^3 t^{1/2} (s + t)^4}
$$

$$
\exp \left\{ -s(\pi m r w)^2 - t(\pi \mu r w)^2 - \frac{s + t(1 - b^2)}{t(s + t)} \right\}
$$

We have written [\(82\)](#page-23-2) as a product of

- a loop factor $\frac{1}{16\pi^2}$,
- a dimensionless coupling $\frac{1}{(\pi r\overline{M}_4)^2}$ built from the effective radius r and the reduced 4-D Planck mass M_4 ,
- a symmetric traceless tensor structure $b_{\mu}b_{\nu} \frac{1}{4}$ $\frac{1}{4}\eta_{\mu\nu}b^2,$
- a function $I_{\text{gravity}}^{\text{scalar}}$ of the dimensionless parameters b^2 , πmr , $\pi \mu r$.

The function $I_{\text{gravity}}^{\text{scalar}}$ is shown in Fig. [9.](#page-24-0) It simplifies if we set $m = 0$ (a massless scalar on the brane) and $b^2 = 0$ (a small boost and / or rotation). Then the sum and integrals can be performed and the behavior for large and small μ can be extracted. There is no IR divergence in $I_{\text{gravity}}^{\text{scalar}}$, even for a massless scalar on the brane, and we find

$$
b^2 \approx 0, m \approx 0: \quad I_{\text{gravity}}^{\text{scalar}} \approx \begin{cases} \frac{1}{3}\zeta(3) & \text{as } \mu r \to 0\\ \frac{4}{9}(\pi\mu r)^2 e^{-2\pi\mu r} & \text{as } \mu r \to \infty \end{cases} \tag{83}
$$

9 Conclusions

In this work we considered a braneworld which is moving or spiraling around a compact extra dimension which we take to be a circle of radius R . The configuration is described by an effective radius r for the compactification and a 4-vector b^{μ} that spontaneously

¹²We rescaled the Schwinger parameters, $s \to (\pi r w)^2 s$ and $t \to (\pi r w)^2 t$.

Figure 9: The quantity $I_{\text{gravity}}^{\text{scalar}}$ appearing in the scalar self-energy due to a bulk graviton loop. The function increases rapidly but has a finite limit as $b^2 \to 1$.

breaks the Lorentz symmetry of the brane worldvolume.

$$
r = \begin{cases} \gamma R & \text{boost-like} \\ R & \text{null} \\ \cos \theta R & \text{tilt-like} \end{cases}
$$

$$
b^{\mu} = \begin{cases} (-\beta, 0, 0, 0) & \text{boost-like} \\ (-\lambda, \lambda, 0, 0) & \text{null} \\ (0, \tan \theta, 0, 0) & \text{tilt-like} \end{cases}
$$
(84)

Loops of bulk fields are sensitive to the parameter b^{μ} and can induce Lorentz-violating terms in the 4-D effective action. We explored this, emphasizing the dimension-4 terms which correct the electron self-energy and the electron – photon vertex.

$$
\mathcal{L} = i c_{\mu\nu} \bar{\psi} \gamma^{\mu} D^{\nu} \psi \qquad D_{\mu} = \partial_{\mu} - ie A_{\mu} \qquad (85)
$$

The one-loop coefficients $c_{\mu\nu} \sim b_{\mu}b_{\nu} - \frac{1}{4}$ $\frac{1}{4}\eta_{\mu\nu}b^2$ due to bulk scalars and gravitons are given in [\(51\)](#page-11-2), [\(76\)](#page-20-2). There are stringent experimental bounds on Lorentz violation, reviewed in [\[6\]](#page-29-2). For the electron, for example, laboratory bounds on the dimensionless coefficients $c_{\mu\nu}$ have reached the level of $\sim 10^{-21}$ [\[14\]](#page-29-10).

The Standard Model Extension is a general framework for incorporating Lorentz violation and provides many effects to explore. In addition to the QED effects mentioned above, we considered Lorentz-violating corrections to the self-energy of a scalar field, $\mathcal{L}\,=\,\frac{1}{2}$ $\frac{1}{2}k_{\mu\nu}\partial^{\mu}\phi\partial^{\nu}\phi$ with coefficients $k_{\mu\mu}$ given in [\(66\)](#page-16-1) and [\(82\)](#page-23-2). Taking the scalar field as a proxy for the Higgs field the experimental bounds on $k_{\mu\nu}$ are surprisingly good [\[6\]](#page-29-2), having reached the level of $10^{-12} - 10^{-20}$ [\[15\]](#page-29-11) or $10^{-13} - 10^{-27}$ [\[16\]](#page-29-12).

While many similar calculations could be done, there are also theoretical issues worth exploring. In particular it would be interesting to understand soft emission from a moving braneworld. This should be related to the infrared behavior of the diagrams we have considered. For example for $\mu^2 = 0$ the vertex correction [\(54\)](#page-13-3) has an IR divergence when $p_1^2 = p_2^2 = m^2$ that should cancel against soft emission in suitable inclusive observables.

Any signal for Lorentz violation in the present epoch would be of the utmost significance. One can also entertain the idea that, although Lorentz-violating effects are extremely small today, they may have been larger in the early universe. Perhaps a braneworld was highly boosted in the early universe and only slowed and stabilized with time. Could the attendant violation of Lorentz symmetry in the early universe leave an observable imprint on cosmology?

Figure 10: The case of light-like b^{μ} . The angle between the brane and the X axis is $\theta = \tan^{-1} \lambda$. The brane moves along the Z axis with velocity λ ; the perpendicular component of the velocity is denoted β .

Acknowledgements

We are grateful to Brian Greene, Janna Levin, Alexios Polychronakos and Massimo Porrati for valuable discussions. DK is supported by U.S. National Science Foundation grant PHY-2112548.

A More on lightlike b^{μ}

Since the geometry of the null case may be a little unfamiliar we give some further explanation. According to [\(21\)](#page-5-5) a brane at $z = 0$ has $Z = \lambda X^{+}$, so the brane spirals around the S^1 as one moves in the X^+ direction. At $T = 0$ the brane is located at $Z = \lambda X$, which means it has been rotated in the XZ plane by an angle $\theta = \tan^{-1} \lambda$. Setting $X = 0$ we have $Z = \lambda T$, which means the brane is moving along the Z axis with velocity λ . (As in the "closing scissors" effect this velocity can be arbitrarily large.) However it's the component of the velocity perpendicular to the brane which is physically relevant, and as can be seen in Fig. [10](#page-26-1) this is given by

$$
\beta = \lambda \cos \theta = \frac{\lambda}{\sqrt{1 + \lambda^2}} \qquad \gamma = \sqrt{1 + \lambda^2} \tag{86}
$$

Thus we can summarize the light-like case as a combination of a boost and a rotation with 13

$$
\gamma \beta = \tan \theta = \lambda \tag{87}
$$

We'd like to understand what a causal signal in the bulk looks like on such a brane. Following the analysis in [\[2\]](#page-28-1) we consider a bulk signal sent out from the origin $T = X =$

 $13\,\text{We're grateful to Massimo Porrati for pointing out this relation.}$

Figure 11: At fixed t the future lightcones of the image charges form circles in the xz plane. Their envelope forms a cone which moves in the direction indicated by the arrow. The lower part of the envelope is parallel to the z axis and moves downward at the speed of light. The upper part of the envelope intersects the x axis at $x = \frac{1+\lambda^2}{1-\lambda^2}t$.

 $Y = Z = 0$ and ask where its future lightcone intersects the brane. It's simplest to work in the covering space where the origin corresponds to an infinite series of image charges located along the Z axis at

$$
Z_w = 2\pi R w \qquad w \in \mathbb{Z} \tag{88}
$$

In the bulk the future lightcones of the image charges are given by

$$
X^{+}X^{-} - |\mathbf{Y}|^{2} - (Z - 2\pi R w)^{2} = 0
$$
\n(89)

Using [\(21\)](#page-5-5) to switch to co-moving coordinates and recalling that for this case $r = R$, the future lightcones are given by

$$
(x - 2\pi r w \lambda)^2 + |\mathbf{y}|^2 + (z - 2\pi r w)^2 = (t + 2\pi r w \lambda)^2
$$
\n(90)

At fixed t we see that the future lightcones are spheres of radius $t + 2\pi r w\lambda$ centered at

$$
x = 2\pi r w \lambda \qquad \mathbf{y} = 0 \qquad z = 2\pi r w \tag{91}
$$

The situation in the xz plane is shown in Fig. [11.](#page-27-0) The centers of the spheres lie along the line $x = \lambda z$. Their envelope defines a cone with opening angle $\theta = \tan^{-1} \lambda$. The tip of the cone is located at^{14} at^{14} at^{14}

$$
x = -t \t z = -t/\lambda \t(92)
$$

The bottom part of the envelope is horizontal and intersects the x axis at

$$
x = -t \tag{93}
$$

¹⁴At time t the tip, where the radius shrinks to zero, corresponds to a fictitious image charge with $w = -t/2\pi r\lambda.$

This means bulk signals propagate in the $-x$ direction at the speed of light. The top part of the envelope, on the other hand, intersects the x axis at

$$
x = -t + \frac{t}{\lambda} \tan(2\theta)
$$

=
$$
\frac{1 + \lambda^2}{1 - \lambda^2} t
$$
 (94)

Thus bulk signals propagate in the $+x$ direction with speed $\frac{1+\lambda^2}{1-\lambda^2}$ $\frac{1+\lambda^2}{1-\lambda^2}$ in agreement with [\(40\)](#page-8-1). For $0 < \lambda < 1$ there is superluminal propagation in the $+x$ direction. At $\lambda = 1$ the velocity diverges and propagation is instantaneous. For $\lambda > 1$ the velocity is negative, which can be thought of as a signal from the origin that is traveling in the $+x$ direction but backwards in time. Alternatively it can be thought of as a signal going forward in time that was emitted in the far past at $x = +\infty$, destined to reach the origin at $t = 0$. This can be seen geometrically in Fig. [11](#page-27-0) from the fact that the range $1 < \lambda < \infty$ corresponds to $\pi/2 < 2\theta < \pi$.

If we include the transverse directions, then at time t and position $x = z = 0$ the envelope extends into the transverse directions a distance

$$
|\mathbf{y}| = \sqrt{t^2 + \left(\frac{t}{\lambda}\right)^2} \tan \theta
$$

= $\sqrt{1 + \lambda^2} t$ (95)

Thus bulk signals propagate in the transverse directions at a superluminal speed $\sqrt{1 + \lambda^2}$. To relate this to [\(39\)](#page-8-2), note that a particle moving in the transverse directions has $x = 0$, which means $x^+ = x^- = t$ and hence $v^- = 1$. Then [\(38\)](#page-8-2) becomes $|\mathbf{v}_\perp|^2 + \frac{(n/r)^2 + \mu^2}{(k^+)^2}$ $\frac{(r)^2 + \mu^2}{(k+1)^2} =$ $1 + \lambda^2$, so the transverse velocity is bounded above by $\sqrt{1 + \lambda^2}$.

References

- [1] B. Greene, D. Kabat, J. Levin, and A. S. Menon, "Superluminal propagation on a moving braneworld," Phys. Rev. D 106 [no. 8, \(2022\) 085001,](http://dx.doi.org/10.1103/PhysRevD.106.085001) [arXiv:2206.13590](http://arxiv.org/abs/2206.13590) [\[hep-th\]](http://arxiv.org/abs/2206.13590).
- [2] B. Greene, D. Kabat, J. Levin, and M. Porrati, "Back to the future: Causality on a moving braneworld," Phys. Rev. D 107 [no. 2, \(2023\) 025016,](http://dx.doi.org/10.1103/PhysRevD.107.025016) [arXiv:2208.09014 \[gr-qc\]](http://arxiv.org/abs/2208.09014).
- [3] A. P. Polychronakos, "Superluminal anisotropic propagation and wavefront splitting on tilted and boosted braneworlds," Phys. Lett. B 841 [\(2023\) 137917,](http://dx.doi.org/10.1016/j.physletb.2023.137917) [arXiv:2210.11497 \[hep-th\]](http://arxiv.org/abs/2210.11497).
- [4] D. Colladay and V. A. Kostelecky, "CPT violation and the standard model," Phys. Rev. D 55 (1997) 6760-6774, [arXiv:hep-ph/9703464](http://arxiv.org/abs/hep-ph/9703464).
- [5] D. Colladay and V. A. Kostelecky, "Lorentz violating extension of the standard model," Phys. Rev. D 58 [\(1998\) 116002,](http://dx.doi.org/10.1103/PhysRevD.58.116002) [arXiv:hep-ph/9809521](http://arxiv.org/abs/hep-ph/9809521).
- [6] V. A. Kostelecky and N. Russell, "Data tables for Lorentz and CPT violation," *[Rev. Mod. Phys.](http://dx.doi.org/10.1103/RevModPhys.83.11)* 83 (2011) 11-31, $arXiv:0801.0287$ [hep-ph].
- [7] H. Bernardo and F. Guo, "Brane motion in a compact space: Adiabatic perturbations of brane-bulk coupled fluids," [arXiv:2308.16821 \[hep-th\]](http://arxiv.org/abs/2308.16821).
- [8] D.-C. Dai and D. Stojkovic, "Superluminal propagation along the brane in space with extra dimensions," [arXiv:2306.04069 \[gr-qc\]](http://arxiv.org/abs/2306.04069).
- [9] B. Greene, J. Levin, and M. Parikh, "Brane-world motion in compact dimensions," [Class. Quant. Grav.](http://dx.doi.org/10.1088/0264-9381/28/15/155013) 28 (2011) 155013, [arXiv:1103.2174 \[hep-th\]](http://arxiv.org/abs/1103.2174).
- [10] I. Antoniadis, "A possible new dimension at a few TeV," [Phys. Lett. B](http://dx.doi.org/10.1016/0370-2693(90)90617-F) 246 (1990) [377–384.](http://dx.doi.org/10.1016/0370-2693(90)90617-F)
- [11] N. Arkani-Hamed, S. Dimopoulos, and G. R. Dvali, "The hierarchy problem and new dimensions at a millimeter," *Phys. Lett. B* 429 [\(1998\) 263–272,](http://dx.doi.org/10.1016/S0370-2693(98)00466-3) [arXiv:hep-ph/9803315](http://arxiv.org/abs/hep-ph/9803315).
- [12] I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos, and G. R. Dvali, "New dimensions at a millimeter to a Fermi and superstrings at a TeV," [Phys. Lett. B](http://dx.doi.org/10.1016/S0370-2693(98)00860-0) 436 [\(1998\) 257–263,](http://dx.doi.org/10.1016/S0370-2693(98)00860-0) [arXiv:hep-ph/9804398](http://arxiv.org/abs/hep-ph/9804398).
- [13] G. F. Giudice, R. Rattazzi, and J. D. Wells, "Quantum gravity and extra dimensions at high-energy colliders," *[Nucl. Phys. B](http://dx.doi.org/10.1016/S0550-3213(99)00044-9)* 544 (1999) 3–38, [arXiv:hep-ph/9811291](http://arxiv.org/abs/hep-ph/9811291).
- [14] L. S. Dreissen, C.-H. Yeh, H. A. Fürst, K. C. Grensemann, and T. E. Mehlstäubler, "Improved bounds on Lorentz violation from composite-pulse Ramsey spectroscopy in a trapped ion," $arXiv:2206.00570$ [physics.atom-ph].
- [15] A. I. Hernández-Juárez, J. Montaño, H. Novales-Sánchez, M. Salinas, J. J. Toscano, and O. Vázquez-Hernández, "One-loop structure of the photon propagator in the Standard Model Extension," [Phys. Rev. D](http://dx.doi.org/10.1103/PhysRevD.99.013002) 99 no. 1, (2019) [013002,](http://dx.doi.org/10.1103/PhysRevD.99.013002) [arXiv:1812.05051 \[hep-ph\]](http://arxiv.org/abs/1812.05051).
- [16] D. L. Anderson, M. Sher, and I. Turan, "Lorentz and CPT violation in the Higgs sector," Phys. Rev. D 70 [\(2004\) 016001,](http://dx.doi.org/10.1103/PhysRevD.70.016001) [arXiv:hep-ph/0403116](http://arxiv.org/abs/hep-ph/0403116).