Testing Bell inequality through $h \to \tau \tau$ at CEPC

Kai Ma^{1, *} and Tong Li $^{a2, \dagger}$

¹Faculty of Science, Xi'an University of Architecture and Technology, Xi'an, 710055, China ²School of Physics, Nankai University, Tianjin 300071, China

(Dated: July 10, 2024)

Abstract

The decay of Higgs boson into two spin-1/2 particles provides an ideal system to reveal quantum entanglement and Bell-nonlocality. Future e^+e^- colliders can improve the measurement accuracy of the spin correlation of tau lepton pairs from Higgs boson decay. We show the testability of Bell inequality through $h \to \tau \tau$ at Circular Electron Positron Collider (CEPC). Two realistic methods of testing Bell inequality are investigated, i.e., Törnqvist's method and Clauser-Home-Shimony-Holt (CHSH) inequality. In the simulation, we take into account the detector effects of CEPC including uncertainties for tracks and jets from Z boson in the production of $e^+e^- \to Zh$. Necessary reconstruction approaches are described to measure quantum entanglement between τ^+ and τ^- . Finally, we show the sensitivity of CEPC to the Bell inequality violation for the two methods.

^a corresponding author

^{*} makai@ucas.ac.cn

 $^{^\}dagger$ litong@nankai.edu.cn

CONTENTS

I. Introduction	2
II. Local Quantum Model and Bell Inequality	4
A. Törnqvist's method	5
B. Clauser-Home-Shimony-Holt Inequality	7
III. Measurements at Future Lepton Colliders	8
A. Simulation and Detector Effects	8
B. Reconstruction Method	10
C. Reconstruction by using Impact Parameters	13
IV. Sensitivity of CEPC to the Bell Inequality Violation	17
V. Discussions and Conclusions	20
Acknowledgements	21
A. The spin correlation coefficients in Törnqvist's method	21
References	22

I. INTRODUCTION

It is well known that the most important debate on whether the Quantum Mechanics (QM) is a complete local theory is the challenge raised by Einstein, Podolsky, and Rosen (EPR), named EPR paradox [1]. The interpretation of the EPR paradox in local hidden variable theory (LHVT) shows the contradiction of LHVT with QM and presents the non-local nature of QM. Later on, Bohm et al. proposed a realistic experiment with a system of two spin-1/2 particles to illustrate the EPR paradox [2]. Based on this consideration, Bell established a theorem that the two particles' spin correlation satisfies a Bell inequality (BI) in realistic LHVT [3]. By contrast, the QM predictions may violate this inequality in some certain parameter space. Clauser, Horne, Shimony and Holt (CHSH) also generalized the original Bell inequality and established a more practical inequality [4]. The test of the Bell

inequality delivers a direct justification if the QM is a complete local theory [5].

In the past years, the violation of this Bell inequality has been observed in many lowenergy experiments (such as optical experiments) [6–13] as the foundation of the modern quantum information theory. The predictions of QM are proved to be consistent with the results of these experiments. However, for testing the completeness of QM beyond the electromagnetic interaction regime, it is still a challenge of the test of Bell inequality in high-energy physics (see review Ref. [14] and references therein). At e^+e^- colliders, the testability of BI was first suggested by using the polarization correlation in the process of $e^+e^- \rightarrow \Lambda \bar{\Lambda} \rightarrow \pi^- p \pi^+ \bar{p}$ [15, 16] or $e^+e^- \rightarrow Z \rightarrow \tau^+ \tau^-$ [17–19]. Based on the CHSH method, dedicated proposals were also raised to test the BI in the final states of $t\bar{t}$ pair [20–29] or two weak gauge bosons [30–38] at the Large Hadron Collider (LHC). Nevertheless, the spin-0 state formed by a pair of spin-1/2 particles in the process such as $e^+e^- \rightarrow \Lambda \bar{\Lambda} \rightarrow \pi^- p \pi^+ \bar{p}$ has the largest entanglement. It is because the two spin-1/2 particles emerging from the decay of a spin-0 particle always have the same helicity. In contrast, for the production in a mixture of spin singlet and triplet channels, the magnitude of the correlation is smaller because of the cancellation of different contributions.

The Higgs boson is the only spin-0 elementary particle in the Standard Model (SM) and can play as a natural spin singlet state to test LHVT through the Bell inequality at high energies. The properties of SM Higgs boson will be measured to high precision at future e^+e^- colliders such as the Circular Electron Positron Collider (CEPC) [39]. We thus propose to test the Bell inequality at CEPC through the Higgsstrahlung process with subsequent decay $h \to \tau^+\tau^-$ [27, 40]

$$e^+e^- \to Zh \to Z\tau^+\tau^-$$
 (1)

The tau lepton pair is correlated in the decay process, and Bell inequality (or quantum entanglement) can be tested by measuring their spin correlation. However, spin information of the tau leptons can only be partially inferred from its decay particle. Here we consider only the tau leptons followed by the 1-prong decay mode $\tau^{\pm} \rightarrow \pi^{\pm}\nu_{\tau}$ which is the best spin analyzer for the tau lepton polarization. In principle, the other decay modes can also be employed. However, it is more challenging in practice because of the kinematic reconstruction of the tau lepton as well as limited spin analyzing power. In order to have higher statistics, the associated Z boson will be reconstructed by both its leptonic and hadronic decay modes. Furthermore, both Törnqvist's method [15] and the CHSH method [4] are explored to evaluate the violation of Bell inequality. To have a more realistic estimation on the experimental sensitivities, we investigate the kinematic reconstruction and simulate the detector effects to reveal the quantum entangled spin correlations in the decay of tau pairs.

Note that QM predicts a larger bound of the joint expectation values in CHSH method, that is $2\sqrt{2}$ but not 2 [41]. One may suspect a certain QM setup would not violate the Bell inequality. New physics beyond the SM may modify the angular distributions of the tau-lepton decay products. However, such kind of new physics has no solid evidence up to now and is restricted to be very small compared to the SM interactions. Therefore, our work is conducted based on the assumption that both the $h \to \tau \tau$ decay and tau decay processes are described by the SM.

This paper is organized as follows. In Sec. II, we first outline the LHVT and Bell inequality. Then we show Törnqvist's method and the CHSH method in terms of the polarization correlation in decay $h \to \tau^+ \tau^- \to \pi^+ \bar{\nu}_\tau \pi^- \nu_\tau$. In Sec. III, we describe the simulation of process $e^+e^- \to Zh \to Z\tau^+\tau^-$ and discuss the detector effects as well as reconstruction methods. The results of projected sensitivity to the Bell inequality violation are given in Sec. IV. Finally, in Sec. V we summarize our conclusions.

II. LOCAL QUANTUM MODEL AND BELL INEQUALITY

In this section, we describe the original and generalized expressions of Bell inequality and the realistic methods of testing it in high-energy physics.

In the LHVT with the hidden variable being λ , the Bell inequality can be phased in terms of the polarization correlation

$$P(\vec{a}, \vec{b}) = \int d\lambda \, q(\lambda) \cdot \mathcal{P}_A(\vec{a}, \lambda) \cdot \mathcal{P}_B(\vec{b}, \lambda) \,, \tag{2}$$

where $\mathcal{P}_{A(B)}(\vec{x}, \lambda)$ is the probability of the fermion A (or B) with spin along the direction $\vec{x} = \vec{a} (\vec{b})$ for given hidden variable λ , and $q(\lambda)$ is the corresponding probability distribution of the hidden variable λ . The original expression of Bell inequality refers to three independent spatial directions \vec{a}, \vec{b} and \vec{c} as

$$\left| P(\vec{a}, \vec{b}) - P(\vec{a}, \vec{c}) \right| \le 1 + P(\vec{b}, \vec{c}) .$$
(3)

On the other hand, in QM the quantum average of the correlation operator $\mathcal{O}(\vec{a}, \vec{b}) \equiv [\vec{\sigma}^A \cdot \vec{a}] [\vec{\sigma}^B \cdot \vec{b}]$ is given by

$$P(\vec{a},\vec{b}) = \langle 00 | \left[\vec{\sigma}^A \cdot \vec{a} \right] \left[\vec{\sigma}^B \cdot \vec{b} \right] | 00 \rangle = -\vec{a} \cdot \vec{b} \,, \tag{4}$$

where $\langle 00 | \text{ or } | 00 \rangle$ refers to a singlet state of the total spin. After inserting the QM prediction Eq. (4) in Eq. (3), the Bell inequality Eq. (3) may be violated in some region of phase space. However, in realistic investigations, the spin correlation of the two fermions A and B can only be transferred to the kinematics of their decay products. In terms of $h \to \tau^+ \tau^-$ and tau leptons' hadronic decay mode $\tau^{\pm} \to \pi^{\pm} \nu_{\tau}$, we will describe two existing methods to perform the test of Bell inequality at high energy colliders.

A. Törnqvist's method

In Ref. [15], Törnqvist suggested to test the BI by using the polarization correlation in the process

$$e^+e^- \to \Lambda\bar{\Lambda} \to \pi^- p \pi^+ \bar{p}$$
 (5)

The parent particle of $\Lambda\bar{\Lambda}$ could be either spin-0 η_c or spin-1 J/ψ . We instead establish the polarization correlation of decay $h \to \tau^+ \tau^- \to \pi^+ \bar{\nu}_\tau \pi^- \nu_\tau$. Since the Higgs boson is a scalar and the τ -lepton is a spin-1/2 particle, the decay process $h \to \tau^+ \tau^-$ provides an ideal system for testing the Bell inequality. The joint spin density matrix for the $\tau^+ \tau^-$ system is given by

$$\rho_{\tau\bar{\tau}} = \frac{1}{4} \left(1 - \vec{\sigma}_{\tau} \cdot \vec{\sigma}_{\bar{\tau}} \right), \tag{6}$$

which means the state with parallel $\vec{\sigma}_{\tau}$ and $\vec{\sigma}_{\bar{\tau}}$ vanishes because of spin-zero condition. For the correlation operator $\mathcal{O}(\vec{a}, \vec{b})$, one can easily find the probability is given as

$$P(\vec{a}, \vec{b}) = \left\langle 00 \middle| \rho_{\tau\bar{\tau}} \mathcal{O}(\vec{a}, \vec{b}) \middle| 00 \right\rangle = -\vec{a} \cdot \vec{b} \,. \tag{7}$$

We show the calculation of the spin correlation coefficients in Appendix A.

However, the spin states of the τ -leptons can not be measured directly at collider, and can only be accessed by the angular distributions of their decay products. Here we only investigate the 1-prong decay mode $\tau^- \to \pi^- \nu_{\tau}$, in which the momentum direction of the charged pion (or equivlently the neutrino) is correlated to the spin direction of the tau lepton. Thus, this decay mode has the largest spin analyzing power compared to the cases of the other decay modes. The decay amplitude of the process $\tau^- \to \pi^- \nu_{\tau}$ in the rest frame of the mother particle can be written as

$$\mathcal{M}_{\tau} = \frac{1}{\sqrt{4\pi}} \left(S + P \vec{\sigma}_{\tau} \cdot \vec{a} \right), \tag{8}$$

where \vec{a} is the unit vector along the π^- momentum direction in the rest frame of τ^- , S and P are the S- and P-wave amplitudes respectively. Similar expression is valid for decay process $\tau^+ \to \pi^+ \bar{\nu}_{\tau}$ as well. Then, the probability of having π^- flying along \vec{a} and π^+ flying along \vec{b} (\vec{b} is the unit vector along the π^+ momentum direction in the rest frame of τ^+) becomes

$$\widetilde{P}(\vec{a},\vec{b}) = \langle 00 \big| \rho_{\tau\bar{\tau}} \big[\mathcal{M}_{\tau} \mathcal{M}_{\bar{\tau}} \big]^{\dagger} \big[\mathcal{M}_{\tau} \mathcal{M}_{\bar{\tau}} \big] \big| 00 \rangle = \left[\frac{1}{4\pi} \Big(\big| S \big|^2 + \big| P \big|^2 \Big) \right]^2 \Big(1 + \alpha^2 \vec{a} \cdot \vec{b} \Big), \quad (9)$$

where

$$\alpha = -\frac{2\Re SP^*}{|S|^2 + |P|^2} \approx 0.573.$$
⁽¹⁰⁾

The above value is obtained by fitting in our numerical simulation. One can see that $\tilde{P}(\vec{a}, \vec{b})$ is a partial measurement of the spin states of the τ -lepton pair. Its normalized value $\tilde{P}^{N}(\vec{a}, \vec{b})$ is related to $P(\vec{a}, \vec{b})$ by the following relation

$$P(\vec{a}, \vec{b}) = \frac{1}{\alpha^2} \left[1 - \widetilde{P}^N(\vec{a}, \vec{b}) \right].$$
(11)

The normalized differential cross section is given as

$$\frac{1}{\sigma} \frac{d\sigma}{d\cos\theta_{ab}} = \frac{1}{2} \widetilde{P}^N(\vec{a}, \vec{b}) = \frac{1}{2} \left[1 - \alpha^2 P(\vec{a}, \vec{b}) \right], \qquad (12)$$

where $\cos \theta_{ab} \equiv \vec{a} \cdot \vec{b} = -P(\vec{a}, \vec{b})$. On the other hand, hidden variable theory predicts [15]

$$\left| P(\vec{a}, \vec{b}) \right| \le 1 - \frac{2}{\pi} \theta_{ab} \,, \quad \theta_{ab} \in [0, \pi] \,. \tag{13}$$

Then we have the following classical region satisfying the Bell inequality

$$\begin{cases} \frac{1}{2} - \alpha^2 \left(\frac{1}{2} - \frac{\theta_{ab}}{\pi}\right) \le \frac{1}{\sigma} \frac{d\sigma}{d\cos\theta_{ab}} \le \frac{1}{2} + \alpha^2 \left(\frac{1}{2} - \frac{\theta_{ab}}{\pi}\right), & \theta_{ab} \in [0, \pi/2] \\ \frac{1}{2} + \alpha^2 \left(\frac{1}{2} - \frac{\theta_{ab}}{\pi}\right) \le \frac{1}{\sigma} \frac{d\sigma}{d\cos\theta_{ab}} \le \frac{1}{2} - \alpha^2 \left(\frac{1}{2} - \frac{\theta_{ab}}{\pi}\right), & \theta_{ab} \in (\pi/2, \pi] \end{cases}$$
(14)

B. Clauser-Home-Shimony-Holt Inequality

Clauser, Horne, Shimony and Holt (CHSH) generalized the original Bell inequality Eq. (3) by considering general properties of the quantum density matrix of a spin-1/2 particles system [4]. Density matrix of the quantum state having two spin-1/2 particles can be expressed in general as

$$\rho = \frac{1}{4} \left[\mathbb{I}_A \otimes \mathbb{I}_B + A_i \cdot \left(\sigma_{A,i} \otimes \mathbb{I}_B \right) + B_j \cdot \left(\mathbb{I}_A \otimes \sigma_{B,j} \right) + C_{ij} \left(\sigma_{A,i} \otimes \sigma_{B,j} \right) \right]$$
(15)

where $\sigma_{A(B),i}$ and $\mathbb{I}_{A(B)}$ are Pauli matrices and the unit 2 × 2 matrix for the particle A(B), respectively. The Bell operator associated with the quantum CHSH inequality is defined as

$$\mathcal{B}_{\text{CHSH}} = \vec{a} \cdot \vec{\sigma}_A \otimes \left(\vec{b} + \vec{b}'\right) \cdot \vec{\sigma}_B + \vec{a}' \cdot \vec{\sigma}_A \otimes \left(\vec{b} - \vec{b}'\right) \cdot \vec{\sigma}_B, \qquad (16)$$

where $\vec{a}, \vec{a}', \vec{b}, \vec{b}'$ are unit vectors. Then the CHSH inequality is given by [42]

$$\left| \operatorname{Tr}(\rho \mathcal{B}_{\mathrm{CHSH}}) \right| \le 2$$
 (17)

Again, in practice it is hard to test the above inequality directly because of the challenge in measuring of the spin directions \vec{a} , $\vec{a'}$, \vec{b} , $\vec{b'}$. Alternatively, the matrix with following coefficients

$$C_{ij} = \operatorname{Tr}\left[\rho\sigma_i \otimes \sigma_j\right],\tag{18}$$

can provide an indirect inequality. It was shown that if sum of the two largest eigenvalues of the matrix $U = C^T C$ is larger than 1, then the CHSH inequality is violated [42].

At colliders, the density matrix can be estimated by angular distributions of the two spin-1/2 particles' decay products. The normalized differential cross section can be generally parameterized as [43]

$$\frac{\sigma^{-1}d\sigma}{d\cos\theta_{A,i}d\cos\theta_{B,j}} = \frac{1}{4} \left[1 + A_i\cos\theta_{A,i} + B_j\cos\theta_{B,j} + C_{ij}\cos\theta_{A,i}\cos\theta_{B,j} \right], \quad (19)$$

where $\theta_{A(B),i(j)}$ are the polar angle of charged particles A(B) from the decays of their mother particles, and measured from the i(j)-th axis. The helicity basis is always chosen for the spins of the two taus. In our case of $h \to \tau^+ \tau^- \to \pi^+ \bar{\nu}_\tau \pi^- \nu_\tau$, the cosine quantities of the above polar angles are defined as

$$\cos \theta_{\pi^+,i} = \hat{p}_{\pi^+} \cdot \hat{i} , \quad \cos \theta_{\pi^-,j} = \hat{p}_{\pi^-} \cdot \hat{j} ,$$
 (20)

where the unit vectors \hat{i} and \hat{j} are defined in the rest frames of the τ^+ and τ^- , respectively. They belong to a chosen orthonormal basis $\hat{j} \in \{\hat{k}, \hat{r}, \hat{n}\}$ and satisfy the relation $\hat{i} = -\hat{j}$. More precisely, we define an unit vector \hat{k} as the direction of τ^- momentum in the rest frame of the Higgs boson. In the rest frame of the τ^- lepton, we define an unit vector \hat{r} in the decay plane of the τ^- lepton and perpendicular to \hat{k} , and an unit vector $\hat{n} = \hat{k} \times \hat{r}$. It was shown that the matrix C can be calculated as [21, 43].

$$C_{ij} = -9 \int d\cos\theta_{A,i} d\cos\theta_{B,j} \frac{\sigma^{-1} d\sigma}{d\cos\theta_{A,i} d\cos\theta_{B,j}} \cos\theta_{A,i} \cos\theta_{B,j} .$$
(21)

Then, one can diagonalize the spin correlation matrix $C^T C$ and find the two largest eigenvalues to test the CHSH inequality. Since the Higgs boson is considered to be on-shell, and its spin is zero, there is no any invariant mass and orientation dependencies, compared to the $t\bar{t}$ final states in Ref. [21].

III. MEASUREMENTS AT FUTURE LEPTON COLLIDERS

At e^+e^- colliders, the dominant production mode of the Higgs boson is the so-called Higgsstrahlung channel, $e^+e^- \to Zh$. For our interested mode $h \to \tau^+\tau^-$ with subsequent decay channels $\tau^{\pm} \to \pi^{\pm}\nu$, two neutrinos appear in the final state. Hence kinematic reconstruction is necessary in order to measure quantum entanglement between τ^+ and τ^- . Next, we describe our numerical simulation and implementation of the detector effects, and then the reconstruction approaches in both the leptonic and hadronic decay modes of Z boson.

A. Simulation and Detector Effects

Our numerical simulations are conducted using the MadGraph5_aMC@NLO [44] package, and the quantum entangled spin correlations in the tau-lepton decay are preserved by the TauDecay [45] package. For a realistic simulation, detector resolutions have to be included for the objects. Charged tracks can be precisely measured by the CEPC detector for the decay products in $Z \to \ell^+ \ell^-$ ($\ell = e, \mu$) or $\tau^{\pm} \to \pi^{\pm} \nu$. Table I lists typical uncertainties of the azimuthal angle (ϕ), rapidity (η) and magnitude of the transverse momentum ($|\vec{p}_T|$) at CEPC [39]. One can see that the CEPC uncertainties for tracks are quite small. We smear tracks (both leptons and pions) by randomly sampling the azimuthal angle, pseudo-rapidity and transverse momentum according to Gaussian distribution with standard deviations given in Table I [46, 47].

Observables	Uncertainties	
ϕ	$0.0002 \eta + 0.000022$	
η	$0.000016 \eta + 0.00000022$	
$ ec{p_T} $	$0.036 ec{p_T} $	

TABLE I. CEPC uncertainties for tracks

However, uncertainties of jet from Z boson's hadronic decay are relatively large. Measurement of jet is not only smeared by fragmentation of partons and the corresponding jet clustering processes, but also by the jet clustering of reconstructed objects after detector response and its matching to jet at the generator level [48]. The results in Ref. [49, 50] indicate that the uncertainty induced by the jet clustering and matching can be as significant as those from the detector response, and becomes the dominant uncertainty especially for final state with more than two jets. Hence, sophisticated jet clustering algorithm has to be used [48]. The energy resolution of the jet from light quarks can be described as [39]

$$\sigma_{\rm jet}(E) = \frac{25.7\%}{\sqrt{E}} \oplus 2.4\%.$$
(22)

Jets from charm and bottom quarks have slightly larger uncertainties because of neutrinos in their decays [39]. In consideration of this, in this paper we also use a smearing algorithm to account for detector resolutions. Detector responses to the partons (for the channel $Z \rightarrow q\bar{q}$) are included by smearing energy of the partons according to Gaussian distribution with standard deviations given in Eq. (22).

To see the impact of the above uncertainties, in Fig. 1, we show the distributions of the difference between the real value and the smeared value of transverse momentum p_T , azimuthal angle ϕ and rapidity η (defined as Δp_T , $\Delta \phi$ and $\Delta \eta$) for the objects in different decay modes of Z boson. One can see that, due to the jet energy smearing, the p_T uncertainties of jets in Z boson's hadronic decay are quite large compared to those in the leptonic mode. As a result, the Z boson decaying to dijet is not well reconstructed as shown in Fig. 2.

Furthermore, detection efficiency is also affected by particle identification. For CEPC, the detector is designed to identify prompt leptons with high efficiency and high purity [39].

For leptons with energies above 5 GeV, the identification efficiency is higher than 99% and misidentification rate is smaller than 2%. For the τ -jet with visible energy between 20 and 80 GeV, the identification efficiency is above 80% with a purity closing to 90% [39], further improvement can be expected by optimizations. In our simulation, we ignore the momentum dependence and use an universal identification efficiency 80% estimate experimental significance for τ -jet. For jets from hadronic decay of the Z boson, b-jets can be tagged with an efficiency of 80% and a purity of 90%. Similarly, an efficiency of 60% and a purity of 60% can be achieved for the c-jet tagging [39]. In our case, since the Z boson is treated inclusively, jet-tagging is irrelevant to our analysis. Therefore, we will use an factor of 0.8 to account for possible efficiency loss in reconstruction at the detector level.



FIG. 1. Normalized number of events as a function of Δp_T (left), $\Delta \phi$ (middle) and $\Delta \eta$ (right) for the objects (blue: e^{\pm} or jet j, green: π^{\pm}) in leptonic (top) and hadronic (bottom) decay modes of Z boson.

B. Reconstruction Method

For decay of the Higgs boson, the degree of freedom of the corresponding phase space is 8, and 6 of them can be measured thanks to the two charged pions. As a result, only 2 of them are undetermined. Considering that the decay width of τ -lepton is very small compared to its mass, it is an excellent approximation to assume that both τ s are on-shell. With help of the on-shell conditions, the 8 kinematic degree of freedom can be determined. In the



FIG. 2. Normalized number of events as a function of Δp_T (left), $\Delta \phi$ (middle) and $\Delta \eta$ (right) for the Z boson in leptonic (top) and hadronic (bottom) decay modes.

following studies, we always assume that the Z-boson is reconstructed by its visible decay products, and momentum of the Higgs boson is obtained by energy-momentum conservation condition, *i.e.*, $p_h = p_{e^+} + p_{e^-} - p_Z$. In the approximation of that $P^0 = \sqrt{s}$, and $\vec{P} = 0$ with $P = p_{e^+} + p_{e^-}$, invariant mass of the Higgs momentum is given by $p_h^2 = s + p_Z^2 - 2E_Z\sqrt{s}$. Since the decay width of Higgs boson is also expected to be very small, $p_h^2 \approx m_h^2$ is again an excellent approximation. In practice, $p_h^2 \equiv \hat{m}_h^2$ may deviate from m_h^2 significantly due to experimental uncertainties in measurement of the Z boson momentum.

Since the Higgs boson decay isotropically, the reconstruction is done in the rest frame of h. Assuming that both τ -leptons are on-shell, then energy and magnitude of the τ -lepton momentum in this reference frame can be obtained directly,

$$E_{\tau}^{\star} = \frac{1}{2}\hat{m}_h, \quad p_{\tau}^{\star} = \frac{1}{2}\hat{m}_h \sqrt{1 - \frac{4m_{\tau}^2}{\hat{m}_h^2}}.$$
 (23)

Intersection angles between momentum of the τ^{\pm} and π^{\pm} in this frame are given as,

$$\cos\theta_{\pm}^{\star} = \frac{2E_{\tau}^{\star}E_{\pi^{\pm}}^{\star} - m_{\tau}^{2} - m_{\pi^{\pm}}^{2}}{2p_{\tau}^{\star}p_{\pi^{\pm}}^{\star}}$$
(24)

Without loss of generality, we define the z-axis as the momentum direction of the negatively charged decay product, and the positively charged decay product lies in the x - z plane. In this reference frame, momenta of the of the charged decay products can be written as,

$$p_{\pi^{-}}^{\star\mu} = E_{\pi^{-}}^{\star} \left(1, \, 0, \, 0, \, \beta_{\pi^{-}}^{\star} \right), \tag{25}$$

$$p_{\pi^+}^{\star\mu} = E_{\pi^+}^{\star} \left(1, \, \beta_{\pi^+}^{\star} \sin \theta_{\pi^+}^{\star}, \, 0, \, \beta_{\pi^+}^{\star} \cos \theta_{\pi^+}^{\star} \right).$$
(26)

Furthermore, defining the azimuthal angle of the τ^- -lepton as ϕ_-^* , its momentum can be parameterized as

$$p_{\tau^-}^{\star\mu} = E_{\tau}^{\star} \left(1, \ \beta_{\tau}^{\star} \sin \theta_{-}^{\star} \cos \phi_{-}^{\star}, \ \beta_{\tau}^{\star} \sin \theta_{-}^{\star} \sin \phi_{-}^{\star}, \ \beta_{\tau}^{\star} \cos \theta_{-}^{\star} \right), \tag{27}$$

where $\beta_{\tau}^{\star} = \sqrt{1 - 4m_{\tau}^2/\hat{m}_h^2}$. It turns out that momentum of the τ^+ -lepton is given as

$$p_{\tau^+}^{\star\mu} = E_{\tau}^{\star} \left(1, -\beta_{\tau}^{\star} \sin \theta_{-}^{\star} \cos \phi_{-}^{\star}, -\beta_{\tau}^{\star} \sin \theta_{-}^{\star} \sin \phi_{-}^{\star}, -\beta_{\tau}^{\star} \cos \theta_{-}^{\star} \right).$$
(28)

Using the equation, $\vec{p}_{\tau^+}^{\star} \cdot \vec{p}_{\pi^+}^{\star} = \cos \theta_+^{\star} |\vec{p}_{\tau^+}^{\star}| |\vec{p}_{\pi^+}^{\star}|$, one can immediately have,

$$-\sin\theta_{\pi^+}^{\star}\sin\theta_{-}^{\star}\cos\left(\phi_{-}^{\star}\right) - \cos\theta_{\pi^+}^{\star}\cos\theta_{-}^{\star} = \cos\theta_{+}^{\star}.$$
(29)

Then we get following solutions,

$$\phi_{-}^{\star} = \pm \arccos\left[-\frac{\cos\theta_{\pi^{+}}^{\star}\cos\theta_{-}^{\star} + \cos\theta_{+}^{\star}}{\sin\theta_{\pi^{+}}^{\star}\sin\theta_{-}^{\star}}\right].$$
(30)

Both solutions satisfy all the kinematic constraints, hence there is a two-fold ambiguity.

We then test the above analytical reconstruction method at parton level. Fig. 3 shows the longitudinal and transverse correlations as a result of the above analytical solutions for the leptonic decay mode of Z boson. In the left panel of Fig. 3, the colored densities indicate the true values of $\cos \theta_{\pi^+}$ versus $\cos \theta_{\pi^-}$ and the black contours show the reconstructed values. The above two-fold ambiguity induces a reduction of the transverse spin correction which is described by the azimuthal angle difference of the two decay planes $\delta \phi$, as shown in the right panel of Fig. 3. In the leptonic decay mode of Z, one can see that the above analytical reconstruction method works well.

However, there are some drawbacks in this analytical reconstruction method. First of all, the two-fold ambiguity of kinematic solutions exists as mentioned above. One cannot determine the complete ϕ_{-}^{\star} distribution at a time. More importantly, due to the energy uncertainty of jets, the Z boson cannot be well reconstructed in its hadronic decay mode. As a result, the uncertainties of the Higgs momentum given by analytical reconstruction are very large in the hadronic decay mode of the Z boson. Quantum correlation effects are completely washed out, and hence it is nearly impossible to observe violation of the Bell identity in the hadronic mode. Therefore, we adopt the other reconstruction method by using impact parameters in next section.



FIG. 3. Longitudinal (left) and transverse (right) correlations as a result of analytical reconstruction method for leptonic decay mode of Z boson at parton level.

C. Reconstruction by using Impact Parameters

It was shown that impact parameters of the charged π s are very useful to reconstruct the full decay kinematics [51–53]. The τ -leptons emerging from Higgs decay are strongly boosted. As a result, its typical decay length ~ 3000 μ m is long enough to induce sizable impact parameter for the charged decay product. The CMS group has used the impact parameter to study CP property of the interaction between Higgs and the tau pair [54]. Here we adopt a similar method proposed in Ref. [51]. Furthermore, excellent impact parameter resolution can be achieved by the CEPC vertex detector. The main performance goals for spatial resolution near the IP can be better than 3 μ m [55]. Here, the real impact parameters of the pions are smeared according to Gaussian distribution with standard deviations $\sigma_{IP} = 3 \ \mu$ m.

We use magnitudes of the τ^{\pm} momenta, $|\vec{p}_{\tau^{\pm}}|$, as the free parameters for finding the best fit. For single tau decay, for instance $\tau^- \to \pi^- \nu_{\tau}$, the opening angle between τ^- and π^- is

$$\cos\theta_{\tau^-\pi^-} = \frac{2E_{\tau^-}E_{\pi^-} - m_{\tau}^2 - m_{\pi^-}^2}{2|\vec{p}_{\tau^-}||\vec{p}_{\pi^-}|},\tag{31}$$

where $E_{\tau^-} = \sqrt{m_{\tau}^2 + |\vec{p}_{\tau^-}|^2}$ given by the on-shell condition. Then, the momentum of τ^- is given as

$$\vec{p}_{\tau^{-}} = |\vec{p}_{\tau^{-}}| \cdot \frac{\vec{b}_{\pi^{-}} + \frac{|\vec{b}_{\pi^{-}}|}{\tan \theta_{\tau^{-}\pi^{-}}} \frac{\vec{p}_{\pi^{-}}}{|\vec{p}_{\pi^{-}}|}}{\left| \vec{b}_{\pi^{-}} + \frac{|\vec{b}_{\pi^{-}}|}{\tan \theta_{\tau^{-}\pi^{-}}} \frac{\vec{p}_{\pi^{-}}}{|\vec{p}_{\pi^{-}}|} \right|},$$
(32)

where \vec{b}_{π^-} is the impact parameter of the π^- . Momentum of the neutrino can be obtained by the momentum conservation condition, $p^{\mu}_{\nu_{\tau}} = p^{\mu}_{\tau^-} - p^{\mu}_{\pi^-}$. Similarly, one can obtain momenta of the τ^+ and anti-neutrino as functions of the parameter $|\vec{p}_{\tau^+}|$ and the impact parameter \vec{b}_{π^+} . The best values of the parameters $|\vec{p}_{\tau^\pm}|$ are obtained by minimizing following likelihood function

$$L = L_{BW}(\hat{m}_{\tau\tau}, m_Z, \Gamma_Z) \cdot L_G(\hat{m}_{\tau\tau} - m_Z, \Gamma_Z) \cdot \prod_{\mu=0,1,2,3} L_G(\hat{p}_Z^{\mu} - p_Z^{\mu}, \sigma_Z^{\mu}),$$
(33)

where L_{BW} is the usual Breit-Wigner distribution for the resonant production of the Z boson, and $\hat{m}_{\tau\tau}$ is the reconstructed invariant mass of the tau-lepton pair; \hat{p}_Z^{μ} is the reconstructed momentum of the Z boson, and p_Z^{μ} is the momentum obtained by summing momenta of its decay product; $L_G(x, y)$ is the Gaussian function with mean value x and variance y. Here σ_Z^{μ} is estimated by our numerical simulation.

In Figs. 4 and 5, after using the method of impact parameters, we show the observable uncertainties for the reconstructed τ lepton and the reconstructed Z mass in different decay modes of Z boson, respectively. It turns out that the reconstruction in hadronic mode is as good as that in leptonic mode for azimuthal angle, rapidity and the reconstructed Z boson mass. The reconstructed transverse momentum p_T in hadronic mode is still worse than that in leptonic mode. We also display the reconstructed longitudinal and transverse correlations for hadronic and leptonic decay modes of Z in Fig. 6. As seen in the bottom panel, the reconstructed transverse correlation agrees well with the true result at parton level. It turns out that the method of impact parameters has no the drawback of two-fold ambiguity.



FIG. 4. Normalized number of events as a function of Δp_T (left), $\Delta \phi$ (middle) and $\Delta \eta$ (right) for the reconstructed τ lepton in leptonic (blue) and hadronic (red) decay modes of Z boson.

Finally, in Fig. 7, we show the distribution of $\cos \theta_{\pi\pi}$ for hadronic (red) and leptonic (blue) decay modes of Z and compare with the Bell inequality. The LHVT holds between the two dashed lines (gray fitted region) given by the inequality in Eq. (14). The black solid line shows the normalized differential cross section in Eq. (12) and the result at parton level



FIG. 5. Normalized number of events as a function of reconstructed Z mass $m_{f\bar{f}}^{\text{Rec}}$ in leptonic (red) and hadronic (blue) decay modes of Z boson.



FIG. 6. Reconstructed longitudinal (top) and transverse (bottom) correlations for hadronic (top left) and leptonic (top right) decay modes of Z boson.

is shown in green dots. The simulation results after using the method of impact parameters are represented by the red and blue histograms for hadronic and leptonic decay modes of Z boson, respectively. Naively seen from the distributions, they stay outside the region satisfying the Bell inequality and agree with the QM/QFT prediction in black solid line.

Fig. 8 shows the reconstructed angular distributions of the charged pions. In general, quantum entanglement disappears when we observe any single angular observable of the six



FIG. 7. Reconstructed distributions of $\cos \theta_{\pi\pi}$ for Törnqvist's test of Bell inequality. The grayfitted region is the phase space consistent with classical prediction.

angles. This is because quantum correlations among π^+ and π^- are integrated out, as shown by the flat distributions in the middle panel for the π^+ and the right panel for both π^+ and π^- . Since the x - z plane is by definition spanned by the momentum direction of the τ^- and π^- , the observable $\cos \theta_{\pi^-,n}$ can only be zero as shown in the middle panel. The nontrivial distributions shown in the left-panel of the Fig. 8 are purely kinematic. Since $d\sigma/d \cos \theta_{\pi^-,k}$ is proportional to a constant and $\theta_{\pi^-,r} = \frac{\pi}{2} - \theta_{\pi^-,k}$, we have

$$\frac{\sigma^{-1}d\sigma}{d\cos\theta_{\pi^-,r}} = \frac{\sigma^{-1}d\sigma}{d\sin\theta_{\pi^-,k}} \propto \text{const.} \times \cot\theta_{\pi^-,r} , \qquad (34)$$

which is essentially reconstructed in our approach as shown by the red-solid and green-dashed lines in the left-panel of the Fig. 8. Similarly, the asymmetric distribution for the π^+ is due to the fact that we have defined $\cos \theta_{\pi-,r} > 0$, which can give a nontrivial integration on the differential cross section $d\sigma/d \cos \theta_{\pi-,r} d \cos \theta_{\pi+,r} \propto 1 + C_{rr} \cos \theta_{\pi-,r} \cos \theta_{\pi+,r}$ (the integration region is limited to the range $\cos \theta_{\pi-,r} \in [0, 1]$).

One should note that the calculated coefficients C_{ij} are not really the spin correlation coefficients of the τ -lepton pair. The polar angles in Eq. (19) are defined for the π s rather than τ s. This is a general property of testing Bell-type inequality at colliders, because the spin (or helicity state) of the particle under consideration can not be measured directly. The helicity state can only be inferred partially via the angular distribution of its decay products. For instance, the decay process $\tau \to \pi \nu$ is a good channel to infer the polarization of the

mother τ -lepton.



FIG. 8. Reconstructed angular distributions of π^{\pm} for $\cos \theta_{\pi,r}$ (left), $\cos \theta_{\pi,n}$ (middle) and $\cos \theta_{\pi,k}$ (right).

IV. SENSITIVITY OF CEPC TO THE BELL INEQUALITY VIOLATION

In this section, we show the sensitivity of CEPC to the Bell inequality violation. As stated in Sec. III B, the analytical reconstruction method suffers from the ambiguous two-fold problem etc. We instead adopt the method of impact parameters for the reconstruction of tau leptons as described in Sec. III C.

At the CEPC with $\sqrt{s} = 240 \text{ GeV}$, the total cross section of the Higgsstrahlung process is

$$\sigma_{Zh} = 196.2 \,\text{fb} \,.$$
 (35)

A huge number of the Higgs boson events will be produced with an expected integrated luminosity of $\mathcal{L} = 5.6 \text{ ab}^{-1}$ [39]. However, since both the branching ratios $\mathcal{B}(h \to \tau \tau) =$ 6.32% and $\mathcal{B}(\tau \to \pi \nu_{\tau}) = 10.82\%$ are small, only hundreds of the events are available to test the Bell inequality. The following kinematic cuts are used to select well-reconstructed events, and match to the real detector configuration

$$p_T(\ell/j) > 10 \,\text{GeV}, \ |\eta(\ell/j)| < 3, \ \left|m_{ff}^{\text{Rec.}} - m_Z\right| < 10 \,\text{GeV}.$$
 (36)

The efficiency for the above kinematic cuts is 0.645 for the decay mode $Z \rightarrow \ell \ell$, and 0.648 for the hadronic decay mode $Z \rightarrow jj$. Furthermore, as we have mentioned, a universal jet reconstruction efficiency 0.8 will be used in our following estimation. The number of events can be further reduced by τ -jet identification which is assumed to be 0.8 for a purity closing to 90% at the CEPC [39]. Table II gives the expected number of events at the CEPC.

CEPC (240 GeV, 5.6 ab^{-1})	$Z \to \ell \ell$	$Z \to jj$
No. of Events	55	568
Kin. Cuts and jet reconstruction	22	151
τ -identification	14	97

TABLE II. Number of events used to test the Bell inequality at the CEPC.

The experimental sensitivity for the Törnqvist's approach is studied by defining the following asymmetric observable

$$\mathcal{A} = \frac{N(\cos\theta_{\pi\pi} < 0) - N(\cos\theta_{\pi\pi} > 0)}{N(\cos\theta_{\pi\pi} < 0) + N(\cos\theta_{\pi\pi} > 0)}.$$
(37)

The analytical prediction of the observable gives an upper bound $\mathcal{A} = 0.119$ in LHVT. The experimental sensitivity at the CEPC is estimated by performing 10000 pseudo-experiments and in each experiment we generate as many as possible events to predict the central values and the corresponding uncertainties. Then, we scale the uncertainties to the corresponding luminosities. We obtain $\mathcal{A} = 0.133 \pm 0.269$ for $Z \to \ell \ell$ channel and $\mathcal{A} = 0.137 \pm 0.1$ for $Z \rightarrow jj$ channel, respectively, as listed in Table III. Smaller uncertainties can be obtained with $\mathcal{A} = 0.133 \pm 0.142$ or $\mathcal{A} = 0.137 \pm 0.053$ for updated luminosity $\mathcal{L} = 20$ ab⁻¹. Both central values are above the bound in LHVT. The channel from the hadronic decay mode of Zboson produces more events and gives more reasonable result with small error. In the CHSH approach, the LHVT supports the fact that the sum of the two largest eigenvalues of the matrix $U = C^T C$ (denoted by $m_1 + m_2$) is not larger than 1. It turns out that both channels lead to $m_1 + m_2 > 1$, as listed in Table III. As we can see, for both the Törnqvist's and CHSH approachs, the Bell inequality can only be tested below 1σ level at the CEPC. It is expected that the sensitivity can be further improved by using sophisticated jet reconstruction method and enhanced τ -jet identification efficiency. Note that the results in the fourth and fifth columns of Table III are from our detector-level simulation. The simulation results of SM expectation change from parton-level to detector-level. The parton-level and detector-level predictions are different because the experimental uncertainties can diminish the magnitudes of the spin correlation effects as a result of imperfect reconstruction (and also reconstruction efficiency). In this work, we compare the detector-level QM simulation results with the

Channels	Obs.	Clas.	Exp. $@5.6 ab^{-1}$	Exp. $@20 ab^{-1}$
$Z \to \ell \ell$	${\cal A}$	≤ 0.119	0.133 ± 0.269	0.133 ± 0.142
	$m_1 + m_2$	≤ 1	1.04 ± 0.921	1.04 ± 0.481
$Z \rightarrow jj$	${\cal A}$	≤ 0.119	0.137 ± 0.1	0.137 ± 0.053
	$m_1 + m_2$	≤ 1	1.05 ± 0.355	1.05 ± 0.188

bound in LHVT directly. We will leave the study of a more reasonable comparison in a future work.

TABLE III. The results of observables testing the Bell inequality in Törnqvist's method and the CHSH approach. The experimental predictions are given for the CEPC with colliding energy $\sqrt{s} = 240 \text{ GeV}$ and total luminosities 5.6 ab⁻¹ and 20 ab⁻¹.

It is worthy to point out that, compared to the Törnquivst's method, the calculation of the sum of two largest eigenvalues of $C^T C$ as an estimator in the CHSH approach needs the estimation of the spin projections along all the 3 possible independent directions. Hence, it is relatively more difficult for small data sample. For the Törnquivst's method, the bounds given by the Bell inequality on the LHVT are shown by dashed lines in the distribution of $\cos \theta_{\pi\pi}$ in Fig. 7. In order to give a quantitative determination of the Bell inequality violation, we define the above asymmetric observable \mathcal{A} . In this way, the LHVT has a quantitative upper bound as shown in Table III. Different (quantum) models lead to different values of \mathcal{A} and $m_1 + m_2$, and are certainly affected by new physics. As stated in Introduction, it is reasonable to assume that both the $h \to \tau \tau$ decay and tau decay processes are described by the SM. The QM prediction in Table III is obtained in the SM. We thus claim that the Bell inequality can be tested under the assumption of SM below 1σ level at the CEPC.

In summary, the QM prediction in Table III is given by the parton-level events. The experimental values are based on our Monte-Carlo simulation. There are two factors affecting the disagreement. One is the experimental uncertainty due to the detector effects which can reduce the quantum correlation such that the result tends to deviate from the QM prediction. The second factor is the reconstruction method. The correlation can not be reconstructed precisely. Even if the number of events is increased, the deviation would still exist.

V. DISCUSSIONS AND CONCLUSIONS

Since spin state can not be directly measured at collider, it is a challenge of the test of quantum entanglement and Bell-nonlocality in high-energy collider physics. However, testing Bell-nonlocality in high energy scattering process is essentially important because it provides a unique way to address the quantum entanglement at high energy scale. We investigate the testability of Bell inequality through $h \to \tau^+ \tau^-$, which is an ideal system to observe the LHVT violation, at future e^+e^- collider CEPC. We demonstrated how to use angular distributions of decay products of the spin-correlated τ -pair to address the Bell-nonlocality. Future e^+e^- colliders can improve the measurement accuracy of the spin correlation of tau lepton pairs from Higgs boson decay. Two realistic methods of testing Bell inequality, i.e., Törnqvist's method and the CHSH inequality are studied in terms of the polarization correlation in decay $h \to \tau^+\tau^- \to \pi^+\bar{\nu}_\tau\pi^-\nu_\tau$.

We simulate the production of $e^+e^- \rightarrow Zh \rightarrow Z\tau^+\tau^-$ as well as the Z boson's leptonic and hadronic decay modes. The detector effects of CEPC including uncertainties for tracks and jets from Z boson are taken into account. We also describe necessary reconstruction approaches to measure quantum entanglement between τ^+ and τ^- . Finally, we find that for both the Törnqvist's and CHSH approachs, the Bell inequality can be tested at the CEPC below 1σ level. Further improvements are expected by employing sophisticated jet reconstruction method and enhanced τ -jet identification efficiency.

We also noticed that in Ref. [40] the authors studied the same topic at ILC and FCC-ee. The expected total number of events is 385 at ILC (250 GeV, $\mathcal{L} = 3 \text{ ab}^{-1}$) or 663 at FCC-ee (240 GeV, $\mathcal{L} = 5 \text{ ab}^{-1}$). In contrast, our estimation for CEPC (240 GeV, $\mathcal{L} = 5.6 \text{ ab}^{-1}$) is 111. Even if the total cross section at ILC or FCC-ee is slightly larger (240.1 fb or 240.3 fb) than the one (196.2 fb) at the CEPC, it is too small to have more than 2 times or 5 times larger number of events. Such a difference can only come from the detector effects. The reconstruction of the mode $e^+e^- \rightarrow Z(\rightarrow jj)h(\rightarrow \tau_{\pi}\tau_{\pi})$, which has the largest production rate, is rather difficult and thus influences the detector effects on the reconstruction. Since the jets (from Z or h) are relatively soft at the CEPC (and ILC-250), the usual jet clustering does not give proper output. Hence, the reconstruction efficiency is rather low. Some sophisticated clustering method has to be employed. In our work, we chose the smearing method for CEPC to solve this problem. In addition, we also proposed to use the Törnqvist's method to test the Bell inequality at CEPC. The reason is twofold. Firstly, it is much simple and straightforward. Secondly, in case of the (very) small number of events, the estimation of the correlation efficiency C_{ij} via the integration of the phase space is unstable/unsafe, and hence induces a very large fluctuation of the central/expected value of the observable. This can be seen in Table III, particularly for the leptonic decay mode of the Z boson.

ACKNOWLEDGEMENTS

T.L. would like to thank Xue-Qian Li for helpful discussion. T.L. is supported by the National Natural Science Foundation of China (Grants No. 12375096, 12035008, 11975129) and "the Fundamental Research Funds for the Central Universities", Nankai University (Grant No. 63196013). K. M. was supported by the Natural Science Basic Research Program of Shaanxi (Program No. 2023-JC-YB-041), and the Innovation Capability Support Program of Shaanxi (Program No. 2021KJXX-47).

Appendix A: The spin correlation coefficients in Törnqvist's method

For the Törnqvist's method, we calculate the correlation coefficients in the following way. Assuming the τ and $\bar{\tau}$ are polarized along \vec{a} and \vec{b} , respectively, their helicity basis can be chosen to be the eigenstates of the helicity opeators $\vec{a} \cdot \vec{\sigma}_{\tau}$ and $\vec{b} \cdot \vec{\sigma}_{\bar{\tau}}$, respectively. Then, the expected value of the correlation matrix $\mathcal{O}(\vec{a}, \vec{b}) = (\vec{a} \cdot \vec{\sigma}_{\tau})(\vec{b} \cdot \vec{\sigma}_{\bar{\tau}})$ is given as

$$\langle h_{\tau}h_{\bar{\tau}}|\rho_{\tau\bar{\tau}}\mathcal{O}(\vec{a},\vec{b})|h_{\tau}h_{\bar{\tau}}\rangle = \langle h_{\tau}h_{\bar{\tau}}|\frac{1}{4}(1-\vec{\sigma}_{\tau}\cdot\vec{\sigma}_{\bar{\tau}})(\vec{a}\cdot\vec{\sigma}_{\tau})(\vec{b}\cdot\vec{\sigma}_{\bar{\tau}})|h_{\tau}h_{\bar{\tau}}\rangle, \qquad (A1)$$

where $h_{\tau(\bar{\tau})}$ denotes the helicity of $\tau(\bar{\tau})$. By using the following relation

$$\vec{\sigma}_{\tau} \cdot \vec{\sigma}_{\bar{\tau}} = \frac{1}{2} \left(\vec{\Sigma}^2 - 6 \right) \,, \tag{A2}$$

where $\vec{\Sigma} = \vec{\sigma}_{\tau} + \vec{\sigma}_{\bar{\tau}}$ and $\vec{\Sigma}^2 = 0$ for spin singlet, we have

$$\langle h_{\tau}h_{\bar{\tau}}|\rho_{\tau\bar{\tau}} \mathcal{O}(\vec{a},\vec{b})|h_{\tau}h_{\bar{\tau}}\rangle = \langle h_{\tau}h_{\bar{\tau}}|(\vec{a}\cdot\vec{\sigma}_{\tau})(\vec{b}\cdot\vec{\sigma}_{\bar{\tau}})|h_{\tau}h_{\bar{\tau}}\rangle = h_{\tau}h_{\bar{\tau}}.$$
 (A3)

For $\vec{a} = \hat{\vec{e}}_z$ and $\vec{b} = \hat{\vec{e}}_z$, the spin singlet requires the helicities of τ and $\bar{\tau}$ satisfy

$$h = h_\tau + h_{\bar{\tau}} = 0. \tag{A4}$$

Hence, we obtain the correlation coefficients as

$$C_{zz} = \langle h_{\tau} h_{\bar{\tau}} | \rho_{\tau\bar{\tau}} \mathcal{O}(\hat{\vec{e}}_z, -\hat{\vec{e}}_z) | h_{\tau} h_{\bar{\tau}} \rangle = h_{\tau} h_{\bar{\tau}} = -1, \qquad (A5)$$

and similarly

$$C_{xx} = \langle h_{\tau} h_{\bar{\tau}} | \rho_{\tau\bar{\tau}} \mathcal{O}(\hat{\vec{e}}_x, -\hat{\vec{e}}_x) | h_{\tau} h_{\bar{\tau}} \rangle = -1,$$

$$C_{yy} = \langle h_{\tau} h_{\bar{\tau}} | \rho_{\tau\bar{\tau}} \mathcal{O}(\hat{\vec{e}}_y, -\hat{\vec{e}}_y) | h_{\tau} h_{\bar{\tau}} \rangle = -1.$$
(A6)

The above calculations show that the correlation coefficients in our helicity basis are (-1, -1, -1). In the above calculations, the condition $\vec{\Sigma}^2 = 0$, which means the $\tau \bar{\tau}$ system is a spin-singlet, is crucial. One can have different results if the spin projection axis is defined in a different way.

- Albert Einstein, Boris Podolsky, and Nathan Rosen, "Can quantum mechanical description of physical reality be considered complete?" Phys. Rev. 47, 777–780 (1935).
- [2] D. Bohm and Y. Aharonov, "Discussion of Experimental Proof for the Paradox of Einstein, Rosen, and Podolsky," Phys. Rev. 108, 1070–1076 (1957).
- [3] J. S. Bell, "On the Einstein-Podolsky-Rosen paradox," Physics Physique Fizika 1, 195–200 (1964).
- [4] John F. Clauser, Michael A. Horne, Abner Shimony, and Richard A. Holt, "Proposed experiment to test local hidden variable theories," Phys. Rev. Lett. 23, 880–884 (1969).
- R. A. Bertlmann, "Entanglement, Bell inequalities and decoherence in particle physics," Lect. Notes Phys. 689, 1–45 (2006), arXiv:quant-ph/0410028.
- [6] Alain Aspect, Philippe Grangier, and Gerard Roger, "Experimental realization of Einstein-Podolsky-Rosen-Bohm Gedankenexperiment: A New violation of Bell's inequalities," Phys. Rev. Lett. 49, 91–97 (1982).
- [7] Alain Aspect, Jean Dalibard, and Gerard Roger, "Experimental test of Bell's inequalities using time varying analyzers," Phys. Rev. Lett. 49, 1804–1807 (1982).
- [8] Z. Y. Ou and L. Mandel, "Violation of Bell's Inequality and Classical Probability in a Two-Photon Correlation Experiment," Phys. Rev. Lett. 61, 50–53 (1988).

- [9] Gregor Weihs, Thomas Jennewein, Christoph Simon, Harald Weinfurter, and Anton Zeilinger,
 "Violation of Bell's inequality under strict Einstein locality conditions," Phys. Rev. Lett. 81, 5039–5043 (1998), arXiv:quant-ph/9810080.
- [10] Dik Bouwmeester, Jian-Wei Pan, Matthew Daniell, Harald Weinfurter, and Anton Zeilinger,
 "Observation of three photon Greenberger-Horne-Zeilinger entanglement," Phys. Rev. Lett.
 82, 1345–1349 (1999), arXiv:quant-ph/9810035.
- [11] J. W. Pan, D. Bouwmeester, M. Daniell, H. Weinfurter, and A. Zeilinger, "Experimental test of quantum nonlocality in three-photon Greenberger-Horne-Zeilinger entanglement," Nature 403, 515–519 (2000).
- [12] Andrew G. White, Daniel F. V. James, Philippe H. Eberhard, and Paul G. Kwiat, "Nonmaximally entangled states: Production, characterization, and utilization," Phys. Rev. Lett. 83, 3103–3107 (1999).
- [13] Tao Yang, Qiang Zhang, Jun Zhang, Juan Yin, Zhi Zhao, Marek Żukowski, Zeng-Bing Chen, and Jian-Wei Pan, "All-versus-nothing violation of local realism by two-photon, fourdimensional entanglement," Phys. Rev. Lett. 95, 240406 (2005).
- [14] Yi-Bing Ding, Jun-li Li, and Cong-Feng Qiao, "Bell Inequalities in High Energy Physics," (2007), arXiv:hep-ph/0702271.
- [15] Nils A. Tornqvist, "Suggestion for Einstein-podolsky-rosen Experiments Using Reactions Like $e^+e^- \rightarrow \Lambda \bar{\Lambda} \rightarrow \pi^- p \pi^+ \bar{p}$," Found. Phys. **11**, 171–177 (1981).
- [16] Xi-Qing Hao, Hong-Wei Ke, Yi-Bing Ding, Peng-Nian Shen, and Xue-Qian Li, "Testing the Bell Inequality at Experiments of High Energy Physics," Chin. Phys. C 34, 311–318 (2010), arXiv:0904.1000 [hep-ph].
- [17] Paolo Privitera, "Decay correlations in e+ e- —> tau+ tau- as a test of quantum mechanics," Phys. Lett. B 275, 172–180 (1992).
- [18] S. A. Abel, M. Dittmar, and Herbert K. Dreiner, "Testing locality at colliders via Bell's inequality?" Phys. Lett. B 280, 304–312 (1992).
- [19] Herbert K. Dreiner, "Bell's inequality and tau physics at LEP," in 2nd Workshop on Tau Lepton Physics (1992) arXiv:hep-ph/9211203.
- [20] Yoav Afik and Juan Ramón Muñoz de Nova, "Entanglement and quantum tomography with top quarks at the LHC," Eur. Phys. J. Plus 136, 907 (2021), arXiv:2003.02280 [quant-ph].

- [21] M. Fabbrichesi, R. Floreanini, and G. Panizzo, "Testing Bell Inequalities at the LHC with Top-Quark Pairs," Phys. Rev. Lett. 127, 161801 (2021), arXiv:2102.11883 [hep-ph].
- [22] Claudio Severi, Cristian Degli Esposti Boschi, Fabio Maltoni, and Maximiliano Sioli, "Quantum tops at the LHC: from entanglement to Bell inequalities," Eur. Phys. J. C 82, 285 (2022), arXiv:2110.10112 [hep-ph].
- [23] Yoav Afik and Juan Ramón Muñoz de Nova, "Quantum information with top quarks in QCD,"
 Quantum 6, 820 (2022), arXiv:2203.05582 [quant-ph].
- [24] J. A. Aguilar-Saavedra and J. A. Casas, "Improved tests of entanglement and Bell inequalities with LHC tops," Eur. Phys. J. C 82, 666 (2022), arXiv:2205.00542 [hep-ph].
- [25] Yoav Afik and Juan Ramón Muñoz de Nova, "Quantum Discord and Steering in Top Quarks at the LHC," Phys. Rev. Lett. 130, 221801 (2023), arXiv:2209.03969 [quant-ph].
- [26] Rafael Aoude, Eric Madge, Fabio Maltoni, and Luca Mantani, "Quantum SMEFT tomography: Top quark pair production at the LHC," Phys. Rev. D 106, 055007 (2022), arXiv:2203.05619 [hep-ph].
- [27] Marco Fabbrichesi, Roberto Floreanini, and Emidio Gabrielli, "Constraining new physics in entangled two-qubit systems: top-quark, tau-lepton and photon pairs," Eur. Phys. J. C 83, 162 (2023), arXiv:2208.11723 [hep-ph].
- [28] Mira Varma and O. K. Baker, "Quantum Entanglement in Top Quark Pair Production," (2023), arXiv:2306.07788 [hep-ph].
- [29] Zhongtian Dong, Dorival Gonçalves, Kyoungchul Kong, and Alberto Navarro, "When the Machine Chimes the Bell: Entanglement and Bell Inequalities with Boosted tt," (2023), arXiv:2305.07075 [hep-ph].
- [30] Alan J. Barr, "Testing Bell inequalities in Higgs boson decays," Phys. Lett. B 825, 136866 (2022), arXiv:2106.01377 [hep-ph].
- [31] Alan J. Barr, Pawel Caban, and Jakub Rembieliński, "Bell-type inequalities for systems of relativistic vector bosons," (2022), arXiv:2204.11063 [quant-ph].
- [32] J. A. Aguilar-Saavedra, "Laboratory-frame tests of quantum entanglement in $H \to WW$," Phys. Rev. D 107, 076016 (2023), arXiv:2209.14033 [hep-ph].
- [33] J. A. Aguilar-Saavedra, A. Bernal, J. A. Casas, and J. M. Moreno, "Testing entanglement and Bell inequalities in $H \rightarrow ZZ$," Phys. Rev. D 107, 016012 (2023), arXiv:2209.13441 [hep-ph].

- [34] M. Fabbrichesi, R. Floreanini, E. Gabrielli, and L. Marzola, "Bell inequalities and quantum entanglement in weak gauge bosons production at the LHC and future colliders," (2023), arXiv:2302.00683 [hep-ph].
- [35] Rafael Aoude, Eric Madge, Fabio Maltoni, and Luca Mantani, "Probing new physics through entanglement in diboson production," (2023), arXiv:2307.09675 [hep-ph].
- [36] Alexander Bernal, Paweł Caban, and Jakub Rembieliński, "Entanglement and Bell inequalities violation in $H \rightarrow ZZ$ with anomalous coupling," (2023), arXiv:2307.13496 [hep-ph].
- [37] Federica Fabbri, James Howarth, and Theo Maurin, "Isolating semi-leptonic $H \to WW^*$ decays for Bell inequality tests," (2023), arXiv:2307.13783 [hep-ph].
- [38] Qi Bi, Qing-Hong Cao, Kun Cheng, and Hao Zhang, "New observables for testing Bell inequalities in W boson pair production," (2023), arXiv:2307.14895 [hep-ph].
- [39] Mingyi Dong et al. (CEPC Study Group), "CEPC Conceptual Design Report: Volume 2 -Physics & Detector," (2018), arXiv:1811.10545 [hep-ex].
- [40] Mohammad Mahdi Altakach, Priyanka Lamba, Fabio Maltoni, Kentarou Mawatari, and Kazuki Sakurai, "Quantum information and CP measurement in H → τ⁺τ⁻ at future lepton colliders," Phys. Rev. D 107, 093002 (2023), arXiv:2211.10513 [hep-ph].
- [41] B. S. Cirelson, "QUANTUM GENERALIZATIONS OF BELL'S INEQUALITY," Lett. Math. Phys. 4, 93–100 (1980).
- [42] R. Horodecki, P. Horodecki, and M. Horodecki, "Violating bell inequality by mixed spin-12 states: necessary and sufficient condition," Physics Letters A 200, 340–344 (1995).
- [43] Werner Bernreuther, Dennis Heisler, and Zong-Guo Si, "A set of top quark spin correlation and polarization observables for the LHC: Standard Model predictions and new physics contributions," JHEP 12, 026 (2015), arXiv:1508.05271 [hep-ph].
- [44] J. Alwall, R. Frederix, S. Frixione, V. Hirschi, F. Maltoni, O. Mattelaer, H. S. Shao, T. Stelzer,
 P. Torrielli, and M. Zaro, "The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations," JHEP 07, 079 (2014), arXiv:1405.0301 [hep-ph].
- [45] Kaoru Hagiwara, Tong Li, Kentarou Mawatari, and Junya Nakamura, "TauDecay: a library to simulate polarized tau decays via FeynRules and MadGraph5," Eur. Phys. J. C 73, 2489 (2013), arXiv:1212.6247 [hep-ph].

- [46] J. de Favereau, C. Delaere, P. Demin, A. Giammanco, V. Lemaître, A. Mertens, and M. Selvaggi (DELPHES 3), "DELPHES 3, A modular framework for fast simulation of a generic collider experiment," JHEP 02, 057 (2014), arXiv:1307.6346 [hep-ex].
- [47] Toshinori Abe et al. (Linear Collider ILD Concept Group -), "The International Large Detector: Letter of Intent," (2010), 10.2172/975166, arXiv:1006.3396 [hep-ex].
- [48] Manqi Ruan et al., "Reconstruction of physics objects at the Circular Electron Positron Collider with Arbor," Eur. Phys. J. C 78, 426 (2018), arXiv:1806.04879 [hep-ex].
- [49] Pei-Zhu Lai, "Jet Reconstruction at the CEPC (2017)," CEPC-REC-2017-002.
- [50] Pei-Zhu Lai, Manqi Ruan, and Chia-Ming Kuo, "Jet performance at the circular electronpositron collider," JINST 16, P07037 (2021), arXiv:2104.05029 [hep-ex].
- [51] Kaoru Hagiwara, Kai Ma, and Shingo Mori, "Probing CP violation in $h \to \tau^- \tau^+$ at the LHC," Phys. Rev. Lett. **118**, 171802 (2017), arXiv:1609.00943 [hep-ph].
- [52] D. Jeans and G. W. Wilson, "Measuring the CP state of tau lepton pairs from Higgs decay at the ILC," Phys. Rev. D 98, 013007 (2018), arXiv:1804.01241 [hep-ex].
- [53] Xin Chen and Yongcheng Wu, "Probing the CP-Violation effects in the $h\tau\tau$ coupling at the LHC," Phys. Lett. B **790**, 332–338 (2019), arXiv:1708.02882 [hep-ph].
- [54] Armen Tumasyan *et al.* (CMS), "Analysis of the *CP* structure of the Yukawa coupling between the Higgs boson and τ leptons in proton-proton collisions at $\sqrt{s} = 13$ TeV," JHEP **06**, 012 (2022), arXiv:2110.04836 [hep-ex].
- [55] Muhammd Ahmad *et al.*, "CEPC-SPPC Preliminary Conceptual Design Report. 1. Physics and Detector," (2015).