

Vertex Operators of the KP hierarchy and Singular Algebraic Curves

Atsushi Nakayashiki *

Abstract

Quasi-periodic solutions of the KP hierarchy acted by vertex operators are studied. We show, with the aid of the Sato Grassmannian, that solutions thus constructed correspond to torsion free rank one sheaves on some singular algebraic curves whose normalizations are the non-singular curves corresponding to the seed quasi-periodic solutions. It means that the action of the vertex operator has an effect of creating singular points on an algebraic curve. We further check, by examples, that solutions obtained here can be considered as solitons on quasi-periodic backgrounds, where the soliton matrices are determined by parameters in the vertex operators.

1 Introduction

We revisit the vertex operators of the KP-hierarchy [5]. We apply them to quasi-periodic solutions, that is, solutions which are expressed by Riemann's theta functions of non-singular algebraic curves [22]. The problem we consider here is what kind of solutions we get in this way. To study this problem we use the Sato Grassmannian [36]. We show that solutions obtained here correspond to certain singular algebraic curves whose normalizations are the non-singular curves of the seed quasi-periodic solutions. It implies two things. First is that the action of the vertex operator of the KP-hierarchy has the effect of creating certain singularities on a curve. Second is that the solutions created by vertex operators describe certain limits of quasi-periodic solutions, since singular curves may be considered as limits of non-singular curves. We check, by computer simulations, that the solutions here represent solitons on the quasi-periodic backgrounds, where the soliton matrices can be extracted from the parameters of the vertex operators. It implies that wave patterns of quasi-periodic solutions of the KP equation contain various shapes of soliton solutions [16, 17] as a part. Recently interactions of solitons and quasi-periodic solutions attracted much attention in relation with soliton gases [10, 18, 7].

*Department of Mathematics, Tsuda University, Kodaira, Tokyo 187-8577, Japan, atsushi@tsuda.ac.jp

It is interesting to study whether the results in this paper can have some application to this subject.

Now let us explain the results in more detail along the history of researches. During the last two decades it is revealed that the shapes of soliton solutions of the KP equation form various wave patterns like web diagrams [4, 16, 17]. Those wave patterns are related with combinatorics of non-negative Grassmannians and cluster algebras [20, 21]. Mathematically soliton solutions, in terms of tau function, are those described by linear combinations of exponential functions and are known to be constructed from singular algebraic curves of genus 0 [23, 38]. Quasi-periodic solutions of the KP equation are those written by Riemann's theta function of non-singular algebraic curves of positive genus [22, 37]. It is expected that quasi-periodic solutions tend to soliton solutions in certain genus zero limits [23, 38, 1]. Therefore it is quite interesting to study the wave patterns of quasi-periodic solutions incorporating the recent development on soliton solutions.

One strategy to study this problem is to take limits of quasi-periodic solutions and make correspondence between quasi-periodic solutions and soliton solutions. However, to carry out this program was not very easy because it is difficult to compute limits of period matrices. We have avoided this difficulty by using the Sato Grassmannian and have calculated limits of quasi-periodic solutions for several examples [31, 32, 33, 34]. Recently there are important progress in computing the limits of quasi-periodic solutions [2, 3, 12, 11]. In [12], in particular, some kind of limits have been computed for any Riemann surface. However it seems difficult to understand how solitonic structure is incorporated in the wave patterns of quasi-periodic solutions from those results.

In this paper we change the direction of study. Instead of studying the limits of quasi-periodic solutions we construct solutions corresponding to degenerate algebraic curves which are more convenient to see the relation with soliton solutions. In course of studying the degeneration of quasi-periodic solutions we found the following formula (Theorem 4.5 of [33]):

$$\begin{aligned} \lim \tau_{g,0}(t) &= C e^{-2 \sum_{i=1}^{\infty} \alpha^i t_{2i}} \\ &\times \left(e^{\eta(t, \alpha^{1/2})} \tau_{g-1,0}(t - [\alpha^{-1/2}]) + (-1)^n e^{\eta(t, -\alpha^{1/2})} \tau_{g-1,0}(t - [-\alpha^{-1/2}]) \right), \end{aligned}$$

where C is a certain constant, $t = (t_1, t_2, t_3, \dots)$, $[\kappa] = (\kappa, \kappa^2/2, \kappa^3/3, \dots)$, $\eta(t, p) = \sum_{j=1}^{\infty} t_j p^j$, $\tau_{n,0}$ is a quasi-periodic solution of the KdV hierarchy, expressed in some standard form, corresponding to a hyperelliptic curve of genus n , \lim means pinching a pair of branching points and α is the parameter corresponding to the pinched point. The term inside the bracket of the right hand side has a very special form. It is rewritten using the vertex operator introduced in [5]:

$$\begin{aligned} X(p, q) &= e^{\eta(t,p) - \eta(t,q)} e^{-\eta(\tilde{\partial}, p^{-1}) + \eta(\tilde{\partial}, q^{-1})}, \\ \tilde{\partial} &= (\partial_1, \partial_2/2, \partial_3/3, \dots). \end{aligned}$$

In fact, for any function $\tau(t)$, we have

$$e^{\eta(t,q)} e^{aX(p,q)} \tau(t - [q^{-1}]) = e^{\eta(t,q)} \tau(t - [q^{-1}]) + ae^{\eta(t,p)} \tau(t - [p^{-1}]).$$

If we take $(p, q) = (-\alpha^{1/2}, \alpha^{1/2})$, $a = (-1)^n$ and $\tau(t) = \tau_{g-1,0}$, we recover the above formula. The vertex operator transforms a solution of the KP-hierarchy to another one, that is, if $\tau(t)$ is a solution then $e^{aX(p,q)} \tau(t)$ is again a solution for any constant a [5]. The above fact suggests that the limits of quasi-periodic solutions may be described by the action of vertex operators on quasi-periodic solutions. The aim of this paper is, in a sense, to show that this is actually the case.

Let $\tau_0(t)$ be the quasi-periodic solution of the KP-hierarchy constructed from the data $(C, L_{\Delta,e}, p_\infty, z)$, where C is a compact Riemann surface of genus $g > 0$, $L_{\Delta,e}$ is a certain holomorphic line bundle on C of degree $g - 1$, p_∞ a point of C and z is a local coordinate around p_∞ [22, 37, 15]. We apply vertex operators with various parameters successively to $\tau_0(t)$. More precisely, let $M, N \geq 1$, p_i, q_j , $1 \leq i \leq N$, $1 \leq j \leq M$ distinct complex parameters, $A = (a_{i,j})$ an $M \times N$ matrix. We consider the vertex operator of the form

$$G = e^{\sum_{i=1}^M \sum_{j=1}^N a_{i,j} X(q_i^{-1}, p_j^{-1})}.$$

We make a certain shift on $\tau_0(t)$, apply G and multiply it by a constant times exponential function and we get a new solution $\tau(t)$ (cf. (3.3)). We show that $\tau(t)$ is a solution constructed from the data $(C', \mathcal{W}_e, p'_\infty, z)$ where C' is a certain singular algebraic curve whose normalization is C , \mathcal{W}_e is a certain torsion free sheaf of rank one on C' , p'_∞ is a point of C' and z a local coordinate around p'_∞ . To prove these properties we use the Sato Grassmannian, which we denote by UGM. It is the set of certain subspaces of the vector space $V = \mathbb{C}((z))$ and parametrizes all formal power solutions of the KP-hierarchy. We recall here that if $\tau(t)$ is a solution of the KP-hierarchy so is $\tau(-t)$. We remark that the descriptions of the points of UGM corresponding to $\tau(t)$ and $\tau(-t)$ are not very symmetric. The point corresponding to $\tau(-t)$ is more suitable to describe the geometry. By this reason we determine the point W_e of UGM corresponding to $\tau(-t)$. This is done by examining the properties of the wave function associated with $\tau(t)$ (cf. (2.4)).

The subspace of V corresponding to $\tau_0(-t)$ is a module over the affine coordinate ring R of $C' \setminus \{p_\infty\}$. Since W_e is a subspace of this space, we consider the stabilizer R_e of W_e in R . Following Mumford [27] and Mulase [25, 26] we define C' and \mathcal{W}_e as a scheme and a sheaf on it respectively using R_e and W_e .

Finally it should be mentioned that the relation of the vertex operator and the degeneration of Riemann's theta function has also been observed by Yuji Kodama from a different point of view [18, 19]. It is interesting to investigate the relation of the results of this paper and those of Kodama.

The paper is organized as follows. In section 2 the relation of the KP-hierarchy and the Sato Grassmannian is reviewed. The main results are formulated and stated in section 3. The formula for a quasi-periodic solution, the definition and the properties of the vertex operator and the result of the action of the vertex operator on

a quasi-periodic solution are given here. In section 4 the singular algebraic curve and the sheaf on it associated with the solution considered in the main theorem are defined and their properties are studied. The example of genus one is studied in detail in section 5. Figures of computer simulation by Mathematica are presented here. The proof of the main theorem is given in section 4 and 5. Regarding the readability of the paper proofs of assertions in section 4 are given in appendix A to E.

2 KP-hierarchy and Sato Grassmannian

The KP-hierarchy is the equation for a function $\tau(t)$, $t = (t_1, t_2, t_3, \dots)$ of the form

$$\oint \tau(t - s - [\lambda^{-1}])(t + s + [\lambda^{-1}])e^{-2\eta(s,\lambda)} \frac{d\lambda}{2\pi i} = 0, \quad (2.1)$$

where $s = (s_1, s_2, s_3, \dots)$,

$$[\mu] = \left(\mu, \frac{\mu^2}{2}, \frac{\mu^3}{3}, \dots\right), \quad \eta(t, \lambda) = \sum_{n=1}^{\infty} t_n \lambda^n,$$

and $\oint \cdot \frac{d\lambda}{2\pi i}$ means taking the residue at $\lambda = \infty$. By expanding in $\{s_j\}$ the KP-hierarchy is equivalent to the infinite set of differential equations which include the KP equation in the bilinear form:

$$(D_1^4 - 3D_2^2 - 4D_1D_3)\tau \cdot \tau = 0, \quad (2.2)$$

where the Hirota derivatives D_i is defined, in general, for a function $f(t)$, as the Taylor coefficients of the expansion of $f(t+s)f(t-s)$ in s :

$$f(t+s)f(t-s) = \sum \frac{D^\alpha f \cdot f}{\alpha!} s^\alpha,$$

where $\alpha = (\alpha_1, \alpha_2, \dots)$, $D^\alpha = D_1^{\alpha_1} D_2^{\alpha_2} \dots$, $\alpha! = \alpha_1! \alpha_2! \dots$, $s^\alpha = s_1^{\alpha_1} s_2^{\alpha_2} \dots$. If $x = t_1$, $y = t_2$, $t = t_3$ and $u = 2\partial_x^2 \log \tau(t)$, (2.2) implies the KP equation

$$3u_{yy} + (-4u_t + 6uu_x + u_{xxx})_x = 0. \quad (2.3)$$

The Sato Grassmannian, which we denote by UGM after Sato, parametrizes all formal power series solutions of the KP-hierarchy. It is defined as follows.

Let $V = \mathbb{C}((z))$ be the space of formal Laurent series in one variable z , $V_\phi = \mathbb{C}[z^{-1}]$ the subspace of polynomials in z^{-1} and $V_0 = z\mathbb{C}[[z]]$ the subspace of formal power series vanishing at $z = 0$. Then $V = V_\phi \oplus V_0$. The Sato Grassmannian is the set of subspaces of V with the same size as V_ϕ . More precisely let $\pi : V \rightarrow V_\phi$ be the projection. Then UGM is the set of a subspace U such that $\text{Ker}\pi|_U$ and $\text{Coker}\pi|_U$ is both finite dimensional and their dimensions are same.

Here we shall give a criterion for a subspace of V to be a point of UGM. To this end, for $f(z) \in V$, define the order of f by,

$$\text{ord } f = -N \text{ if } f(z) = cz^N + O(z^{N+1}) \text{ with } c \neq 0.$$

It describes the order of a pole at $z = 0$ if N is negative. We set

$$V(n) = \{f \in V \mid \text{ord } f \leq n\},$$

and, for a subspace U of V , set $U(n) = U \cap V(n)$. Then

Proposition 2.1. A subspace U of V belongs to UGM if and only if $\dim U(n) = n+1$ for all sufficiently large n .

Proof. Suppose that $\dim U(n) = n + 1$ for $n \geq n_0$ with $n_0 \geq 0$. Then $\dim U(n+1)/U(n) = 1$ for $n \geq n_0$. It means that there exist $f_n \in U$, $n \geq n_0 + 1$, such that

$$f_n = z^{-n} + O(z^{-n+1}), \quad n \geq n_0 + 1.$$

We can take f_n , $n \geq n_0 + 1$, as a part of a basis of U . For the remaining part, since $\dim U(n_0) = n_0 + 1$, there exist $f_{m_i} \in U$, $0 \leq i \leq n_0$ such that $\text{ord } f_{m_i} = m_i$ with $m_0 < \dots < m_{n_0}$. By the definition

$$r := \dim \text{Ker } \pi|_U = \#\{j \mid m_j < 0\}.$$

Then

$$\dim \text{Coker } \pi|_U = n_0 + 1 - (n_0 + 1 - r) = r.$$

Thus $U \in \text{UGM}$. □

Corollary 2.2. Let W be a subspace of V . If there exists an integer N such that $\dim W(n) = n - N$ for all sufficiently large n , then $U = z^{N+1}W$ is a point of UGM.

Proof. Since $U(n) = z^{N+1}W(n + N + 1)$ and therefore $\dim U(n) = \dim W(n + N + 1) = n + 1$ for sufficiently large n . Thus the assertion follows from Proposition 2.1. □

Example 2.3. $W = \mathbb{C}z^3 + \mathbb{C}z^2 + \sum_{i=4}^{\infty} \mathbb{C}z^{-i}$. For $n \geq 4$ $\dim W(n) = n - 1$. Then $U = z^2W = \mathbb{C}z^5 + \mathbb{C}z^4 + \sum_{i=2}^{\infty} \mathbb{C}z^{-i}$ is a point of UGM since $\text{Ker } \pi|_U = \mathbb{C}z^5 + \mathbb{C}z^4$ and $\text{Coker } \pi|_U = \mathbb{C}1 + \mathbb{C}z^{-1}$.

Next we explain the correspondence between solutions of the KP-hierarchy and points of UGM. For a point of UGM the corresponding solution of the KP-hierarchy is constructed as a series using Schur functions and Plücker coordinates. For this see [32].

We need, in this paper, the converse construction of the point of UGM from a solution of the KP-hierarchy. Let $\tau(t)$ be a solution of the KP-hierarchy. Define the wave function $\Psi(t; z)$ and the adjoint wave function $\Psi^*(t; z)$ by

$$\Psi(t; z) = \frac{\tau(t - [z])}{\tau(t)} e^{\eta(t, z^{-1})}, \quad \Psi^*(t; z) = \frac{\tau(t + [z])}{\tau(t)} e^{-\eta(t, z^{-1})}. \quad (2.4)$$

Theorem 2.4. [36, 15] Let U be the vector space spanned by the expansion coefficients of $\tau(t)\Psi^*(t; z)$ in t . Then U is the point of UGM corresponding to $\tau(t)$.

It is easy to verify that if $\tau(t)$ is a solution of the KP-hierarchy so is $\tau(-t)$.

Corollary 2.5. Let U' be the vector space spanned by the expansion coefficients of $\tau(t)\Psi(t; z)$ in t . Then U' is the point of UGM corresponding to $\tau(-t)$.

Proof. Let $\Psi_-^*(t; z)$ be the adjoint wave function of $\tau(-t)$. Then

$$\tau(-t)\Psi_-^*(t; z) = \tau(-t - [z])e^{-\eta(t, z^{-1})}. \quad (2.5)$$

If we set $s = -t$, it is equal to

$$\tau(s - [z])e^{\eta(s, z^{-1})} = \tau(s)\Psi(s; z). \quad (2.6)$$

Since the vector space generated by the expansion coefficients of (2.6) in s is the same as that generated by expansion coefficients of (2.5) in t , the assertion of the lemma follows from Theorem 2.4. \square

3 Main results

Let C be a compact Riemann surface of genus $g > 0$, $\{\alpha_i, \beta_i\}_{i=1}^g$ a canonical basis of $H^1(C, \mathbb{Z})$, $\{dv_i\}_{i=1}^g$ the normalized basis of holomorphic one forms, $\Omega = (\Omega_{i,j})_{1 \leq i, j \leq g}$ with $\Omega_{i,j} = \int_{\beta_j} dv_i$ the period matrix, $J(C) = \mathbb{C}^g / L_\Omega$ with $L_\Omega = \mathbb{Z}^g + \Omega\mathbb{Z}^g$ the Jacobian variety of C , p_∞ a point of C , $I(p) = \int_{p_\infty}^p dv$ with $dv = {}^t(dv_1, \dots, dv_g)$ the Abel map, K Riemann's constant, $\theta(z|\Omega)$ Riemann's theta function

$$\theta(z|\Omega) = \sum_{n \in \mathbb{Z}^g} \exp(\pi i {}^t n \Omega n + 2\pi i {}^t n z), \quad z = {}^t(z_1, \dots, z_g),$$

where $n \in \mathbb{Z}^g$ is considered as a column vector.

We extend the the definition of the Abel map to divisors of any degree by, for

$$D = \sum_{j=1}^m p_j - \sum_{j=1}^n q_j,$$

$$I(D) = \sum_{j=1}^m I(p_j) - \sum_{j=1}^n I(q_j).$$

We denote by Δ the Riemann divisor. It is the divisor of degree $g - 1$ which satisfies $2\Delta \equiv \Omega_C^1$ and $I(\Delta) = K$, where \equiv signifies the linear equivalence of divisors and Ω_C^1 is the linear equivalence class of divisors of holomorphic one forms. The Riemann divisor is uniquely determined from the canonical homology basis by the condition [28]

$$\{I(p_1 + \cdots + p_{g-1} - \Delta) \mid p_1, \dots, p_{g-1} \in C\} = \{z \in J(C) \mid \theta(z|\Omega) = 0\}.$$

Notice that the left hand side does not depend on the choice of the base point p_∞ of the Abel map.

Let $E(p_1, p_2)$ be the prime form [8, 29] (see also [30]):

$$E(p_1, p_2) = \frac{\theta[\delta](\int_{p_1}^{p_2} dv)}{h_\delta(p_1)h_\delta(p_2)},$$

where $\delta = \begin{pmatrix} \delta' \\ \delta'' \end{pmatrix}$, $\delta', \delta'' \in \frac{1}{2}\mathbb{Z}^g$ is a non-singular odd half characteristic and $h_\delta(p)$ is the half differential satisfying

$$h_\delta^2(p) = \sum_{j=1}^g \frac{\partial \theta[\delta]}{\partial z_j}(0) dv_j(p).$$

Take a local coordinate z around p_∞ and write

$$\begin{aligned} E(P_1, P_2) &= \frac{E(z_1, z_2)}{\sqrt{dz_1}\sqrt{dz_2}}, \quad P_i \in C, \quad z_i = z(P_i), \\ d_{z_1} d_{z_2} \log E(z_1, z_2) &= \left(\frac{1}{(z_1 - z_2)^2} + \sum_{i,j=1}^{\infty} q_{i,j} z_1^{i-1} z_2^{j-1} \right) dz_1 dz_2, \quad (3.1) \\ dv_i &= \sum_{j=1}^{\infty} v_{i,j} z^{j-1} dz. \end{aligned}$$

We set

$$\mathcal{V} = (v_{i,j})_{1 \leq i \leq g, 1 \leq j}, \quad q(t) = \sum_{i,j=1}^{\infty} q_{i,j} t_i t_j.$$

Then

$$\tau_0(t) = e^{\frac{1}{2}q(t)}\theta(\mathcal{V}t + e|\Omega)$$

is a solution of the KP-hierarchy for arbitrary $e \in \mathbb{C}^g$ [37](see also [35, 15, 30]).

Let

$$\begin{aligned} X(p, q) &= e^{\eta(t,p)-\eta(t,q)} e^{-\eta(\tilde{\partial}, p^{-1})+\eta(\tilde{\partial}, q^{-1})}, \\ \eta(t, p) &= \sum_{j=1}^{\infty} t_j p^j, \quad \tilde{\partial} = (\partial_1, \partial_2/2, \partial_3/3, \dots), \end{aligned}$$

be the vertex operator.

The following theorem is known.

Theorem 3.1. [5] If $\tau(t)$ is a solution of the KP-hierarchy, so is $e^{aX(p,q)}\tau(t)$ for any $a \in \mathbb{C}$.

Vertex operators satisfy

$$X(p_1, q_1)X(p_2, q_2) = \frac{(p_1 - p_2)(q_1 - q_2)}{(p_1 - q_2)(q_1 - p_2)} : X(p_1, q_1)X(p_2, q_2) : \quad (3.2)$$

where $: \ :$ denotes the normal ordering taking all differential operators to the right of all multiplication operators, that is,

$$: X(p_1, q_1)X(p_2, q_2) : := e^{\sum_{j=1}^2 (\eta(t,p_j)-\eta(t,q_j))} e^{\sum_{j=1}^2 (-\eta(\tilde{\partial}, p_j^{-1})+\eta(\tilde{\partial}, q_j^{-1}))}.$$

The following properties follow from this.

$$\begin{aligned} X(p_1, q_1)X(p_2, q_2) &= X(p_2, q_2)X(p_1, q_1) \quad \text{if } p_1 \neq q_2 \text{ and } q_1 \neq p_2, \\ X(p_1, q_1)X(p_2, q_2) &= 0 \quad \text{if } p_1 = p_2 \text{ or } q_1 = q_2. \end{aligned}$$

Let M, N be positive integers, $q_i, 1 \leq i \leq M, p_j, 1 \leq j \leq N$ non-zero complex numbers and $(a_{i,j})$ an $M \times N$ complex matrix. We set $p_{N+j} = q_j$ and use both notation p_{N+j} and q_j .

Set

$$G = e^{\sum_{i=1}^M \sum_{j=1}^N a_{i,j} X(q_i^{-1}, p_j^{-1})}.$$

Define

$$\tau(t) = \Delta(p_1^{-1}, \dots, p_N^{-1}) e^{\sum_{j=1}^N \eta(t, p_j^{-1})} G \tau_0(t - \sum_{j=1}^N [p_j]), \quad (3.3)$$

where $\Delta(p_1, \dots, p_n) = \prod_{1 \leq i < j \leq n} (p_j - p_i)$. It can be computed explicitly as follows.

Set $L = M + N$ and define the $L \times N$ matrix $B = (b_{i,j})$ by

$$\begin{aligned} b_{i,j} &= \delta_{i,j} \quad \text{for } 1 \leq i, j \leq N, \\ b_{N+i,j} &= a_{i,j} \prod_{m \neq j}^N \frac{p_j^{-1} - p_m^{-1}}{q_i^{-1} - p_m^{-1}} \quad \text{for } 1 \leq i \leq M, 1 \leq j \leq N, \end{aligned} \quad (3.4)$$

that is,

$$B = \begin{pmatrix} 1 & & & & & \\ & \ddots & & & & \\ & & 1 & & & \\ b_{N+1,1} & \cdots & b_{N+1,N} & & & \\ \vdots & & \vdots & & & \\ b_{N+M,1} & \cdots & b_{N+M,N} & & & \end{pmatrix}.$$

We set $[L] = \{1, \dots, L\}$ and denote by $\binom{[L]}{N}$ the set of (i_1, \dots, i_N) , $1 \leq i_1 < \dots < i_N \leq L$ [16, 17]. For $I = (i_1, \dots, i_N) \in \binom{[L]}{N}$ set

$$\Delta_I^- = \Delta(p_{i_1}^{-1}, \dots, p_{i_N}^{-1}), \quad \eta_I = \sum_{i \in I} \eta(t, p_i^{-1}), \quad [p_I] = \sum_{i \in I} [p_i],$$

$$B_I = \det(b_{i_r, s})_{1 \leq r, s \leq N}.$$

By a direct calculation using the commutation relation (3.2) we have

Proposition 3.2. [34] The function $\tau(t)$ of (3.3) has the following expression

$$\tau(t) = \sum_{I \in \binom{[L]}{N}} B_I \Delta_I^- e^{\eta_I} \tau_0(t - [p_I]). \quad (3.5)$$

Remark 3.3. The matrices $(a_{i,j})$ and B correspond to each other. When we study the positivity of $\tau(t)$, it is convenient to begin with the matrix B and define the matrix $(a_{i,j})$ from it. For example, in the case where $\tau_0(t) > 0$ for any real t , $\{p_i\}$ are real and $p_1^{-1} < \dots < p_L^{-1}$, then $\tau(t) > 0$ if $B_I \geq 0$ for any I . Later in section 5 this view point is used.

The part $\tau_0(t - [p_I])$ can further be expressed by theta function. To write it let $d\tilde{r}_k$, $k \geq 1$, be the normalized differential of the second kind with a pole only at p_∞ of order $k + 1$, that is, it satisfies

$$\int_{\alpha_j} d\tilde{r}_k = 0 \quad \text{for any } j, \quad d\tilde{r}_k = d(z^{-k} - O(z)) \quad \text{near } p_\infty.$$

The expansion of $d\tilde{r}_k$ near p_∞ can be written more explicitly using $\{q_{i,j}\}$. In integral form it is given by

$$\int^p d\tilde{r}_k = z^{-k} - \sum_{j=1}^{\infty} q_{k,j} \frac{z^j}{j}, \quad p \in C, z = z(p). \quad (3.6)$$

Then

Proposition 3.4. In terms of the theta function $\tau(t)$ defined by (3.3) is written as

$$\tau(t) = e^{\frac{1}{2}q(t)} \sum_{J \in \binom{[L]}{N}} B_J C_J e^{\sum_{j \in J} \sum_{k=1}^{\infty} t_k \int^{P_j} d\tilde{\tau}_k} \theta \left(\mathcal{V}t - \sum_{j \in J} I(Q_j) + e \right), \quad (3.7)$$

where

$$C_J = \prod_{i < j, i, j \in J} E(p_j, p_i) \prod_{j \in J} \frac{p_j}{E(0, p_j)^N},$$

and $P_j, Q_j \in C$ such that $z(P_j) = p_j$, $z(Q_j) = q_j$.

This proposition is proved by a direct calculation using the following lemma which can be derived from (3.1) .

Lemma 3.5. Let $Q(t|s) = \sum_{i,j=1}^{\infty} q_{i,j} t_i s_j$. Then

$$e^{Q(|z||[w])} = \frac{E(z, w)}{w - z} \frac{zw}{E(0, z)E(0, w)}, \quad q^{\frac{1}{2}Q(|z||[z])} = \frac{z}{E(0, z)}.$$

For $e = {}^t(e_1, \dots, e_g) \in \mathbb{C}^g$ L_e denotes the holomorphic line bundle of degree 0 on C whose characteristic homomorphism is specified by

$$\chi(\alpha_j) = 1, \quad \chi(\beta_j) = e^{2\pi i e_j}.$$

If $c \in \mathbb{C}^g$ is taken such that $\theta(c)\theta(e + c) \neq 0$ then $\theta(I(p) + e + c)/\theta(I(p) + c)$ is a meromorphic section of L_{-e} .

We denote L_{Δ} the holomorphic line bundle of degree $g - 1$ corresponding to Δ . Then we consider $L_{\Delta, -e} := L_{\Delta} \otimes L_{-e}$ which has $\theta(I(p) + e)/E(p, p_{\infty})$ as a meromorphic section if $\theta(e) \neq 0$.

Let $H^0(C, L_{\Delta, -e}(*p_{\infty}))$ be the vector space of meromorphic sections of $L_{\Delta, -e}$ which are holomorphic on $C \setminus \{p_{\infty}\}$. Using the local coordinate z we embed this space into $V = \mathbb{C}((z))$ as follows. We consider a section of $L_{\Delta, -e}$ as $E(p, p_{\infty})^{-1}$ times a multi-valued meromorphic function on C whose transformation rule is the same as that of $\theta(I(p) + e)$, that is,

$$f(p + \alpha_j) = f(p), \quad f(p + \beta_j) = e^{-\pi i \Omega_{j,j} - 2\pi i (\int_{p_{\infty}}^p dv_j + e_j)} f(p).$$

We realize a section of $L_{\Delta, -e}$ using a function on C in this way. Then we expand elements of $H^0(C, L_{\Delta, -e}(*p_{\infty}))$ around p_{∞} in z as $\sum a_n z^n \sqrt{dz}$ and get elements $\sum a_n z^n$ of V . In the following we always consider $H^0(C, L_{\Delta, -e}(*p_{\infty}))$ as a subspace of V in this way.

Theorem 3.6. [15] The point of UGM corresponding to $\tau_0(t)$ is $zH^0(C, L_{\Delta, -e}(*p_{\infty}))$ and that corresponding to $\tau_0(-t)$ is $zH^0(C, L_{\Delta, e}(*p_{\infty}))$.

For the solution $\tau(t)$ of (3.3) the descriptions of the points of UGM corresponding to $\tau(t)$ and $\tau(-t)$ are not very symmetric as opposed to $\tau_0(t)$. We consider $\tau(-t)$ here, since it is more conveniently related with the geometry of C as in the case of soliton solutions [38, 23].

Set

$$\begin{aligned} b'_{N+j,i} &= b_{N+j,i}(p_i^{-1}q_j), \\ W_e &= \{f \in H^0(C, L_{\Delta,e}(*p_\infty)) \mid f(p_i) = -\sum_{j=1}^M b'_{N+j,i}f(q_j), \quad 1 \leq i \leq N\}, \\ U_e &= z^{N+1}W_e. \end{aligned} \tag{3.8}$$

Our main theorem is

Theorem 3.7. (i) The subspace U_e is a point of UGM.

(ii) The point of UGM corresponding to $\tau(-t)$ is U_e .

4 Singular curve created by vertex operators

In this section we study the geometry of U_e .

By Theorem 3.6 the point of UGM corresponding to the solution $\tau_0(-t)$ is associated with $(C, L_{\Delta,e}, p_\infty, z)$, where, as in the previous section, C is a non-singular algebraic curve of genus $g > 0$, $L_{\Delta,e}$ is the holomorphic line bundle on C , p_∞ a point of C and z a local coordinate around p_∞ [37, 15].

We show that the point U_e of UGM corresponding to the solution $\tau(-t)$ is associated with $(C', \mathcal{W}_e, p'_\infty, z)$, where C' is a singular algebraic curve whose normalization is C , \mathcal{W}_e is a rank one torsion free sheaf on C' , p'_∞ is a point of C' and z a local coordinate at p'_∞ . It suggests that, geometrically, the action of the vertex operator has an effect of creating some kind of singularities on a curve. Moreover singular curves may be considered as degenerate limits of non-singular curves. Therefore $\tau(-t)$ and consequently $\tau(t)$ can be considered as a certain limit of a quasi-periodic solution.

In order to define the curve C' from U_e the most appropriate way in the present case is the abstract algebraic method of Mumford and Mulase [27, 25, 26]. Namely we define C' as a complete integral scheme and \mathcal{W}_e as a sheaf on it.

We referred to [9, 13, 14, 24] as references on algebraic geometry and commutative algebras.

Let $R := H^0(C, \mathcal{O}(*p_\infty))$ be the vector space of meromorphic functions on C which have a pole only at p_∞ . By expanding functions in the local coordinate z around p_∞ we consider R as a subspace of $V = \mathbb{C}((z))$. It is the affine coordinate ring of $C \setminus \{p_\infty\}$. The vector space $H^0(C, L_{\Delta,e}(*p_\infty))$ is an R -module and W_e is a vector subspace of it. Let

$$R_e = \{f \in R \mid fW_e \subset W_e\}$$

be the stabilizer of W_e in R . Then

Proposition 4.1. We have

$$R_e = \{f \in R \mid f(p_i) = f(q_j) \text{ if } b_{N+j,i} \neq 0\}. \quad (4.1)$$

The proof of this proposition is given in Appendix A.

To study the structure of R_e we introduce a directed graph G_B associated with the matrix B . The vertices of G_B consists of $\{p_1, \dots, p_L\}$. The vertices p_i and p_{N+j} are connected by an edge with the weight $b_{N+j,i}$. The direction of the edge is from p_{N+j} to p_i . We understand the edge with the weight 0 is the same as that there is no edge. Other edges are not connected. Notice that one can recover the matrix B from G_B .

Let s be the number of connected components of G_B . We divide the set of vertices $\{p_i \mid 1 \leq i \leq L\}$ according as connected components and rename them as $\{p_{i,j} \mid 1 \leq j \leq n_i, 1 \leq i \leq s\}$. We denote $P_{i,j}$ the point on C such that $z(P_{i,j}) = p_{i,j}$.

Example 4.2. Consider

$$B = \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ 0 & a \\ -b & 0 \end{pmatrix}, \quad a, b \neq 0.$$

In this case G_B is


$$\begin{array}{ccc} p_1 & \xleftarrow{-b} & p_4 \\ p_2 & \xleftarrow{a} & p_3 \end{array}$$

and $s = 2$. Then $(p_{1,1}, p_{1,2}) = (p_1, p_4)$, $(p_{2,1}, p_{2,2}) = (p_2, p_3)$ for example.

Example 4.3. Let

$$B = \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ -c & a \\ -d & b \end{pmatrix}, \quad a, b, c, d \neq 0.$$

Then G_B is



In this case $s = 1$ and $(p_{1,1}, p_{1,2}, p_{1,3}, p_{1,4}) = (p_1, p_2, p_3, p_4)$ for example.

In the notation introduced above R_e is described as

$$R_e = \{f \in R \mid f(p_{i,j}) = f(p_{i,j'}) \text{ for any } j, j', 1 \leq i \leq s\}. \quad (4.2)$$

Let $H^0(C, \mathcal{O}(np_\infty))$ be the space of meromorphic functions on C with a pole only at p_∞ of order at most n and

$$\begin{aligned} R(n) &= H^0(C, \mathcal{O}(np_\infty)), \\ R_e(n) &= R(n) \cap R_e. \end{aligned} \quad (4.3)$$

Notice that $R(n) = R_e(n) = \{0\}$ for $n < 0$ and $R(0) = R_e(0) = \mathbb{C}$. The set of subspaces $\{R_e(n)\}$ satisfies $R_e(n) \subset R_e(n+1)$ for any n and $R_e = \cup_{n=0}^{\infty} R_e(n)$.

Set

$$\begin{aligned} A' &= \bigoplus_{n=0}^{\infty} R_e(n), \\ C' &= \text{Proj } A', \end{aligned}$$

We call $R_e(n)$ the homogeneous component of A' with degree n . There is a natural injective morphism $\varphi : \text{Spec } R_e \rightarrow C'$ given by

$$\varphi(\mathcal{P}) = \bigoplus_{n=0}^{\infty} \mathcal{P}^{(n)}, \quad \mathcal{P}^{(n)} = \mathcal{P} \cap R_e(n). \quad (4.4)$$

Next define

$$p'_\infty = \bigoplus_{n=0}^{\infty} R_e(n-1),$$

where $R_e(n-1)$ is located at the homogeneous component of A' with degree n . It can be easily checked that $p'_\infty \in C'$.

By the Riemann-Roch theorem there exists N_0 such that

$$\dim R_e(n)/R_e(n-1) = 1 \quad n \geq N_0.$$

Take an arbitrary $m \geq N_0$ and $a \in R_e(m)$ such that

$$a = z^{-m} + O(z^{-m+1}). \quad (4.5)$$

We consider a as a homogeneous element of A' with degree m . Set

$$\begin{aligned} D'_+(a) &= \{\mathcal{P} \in \text{Proj } A' \mid a \notin \mathcal{P}\}, \\ A'_{(a)} &= \{ua^{-n} \mid u \in R_e(mn), n \geq 0\} \end{aligned}$$

= the set of elements of degree zero in $A'[a^{-1}]$.

Then $D'_+(a)$ is an affine open subscheme of C' isomorphic to $\text{Spec} A'_{(a)}$ (c.f. [9]). This isomorphism is given by

$$\mathcal{P} \mapsto \bigoplus_{n=0}^{\infty} a^{-n} \mathcal{P}^{(nm)}.$$

Then

Theorem 4.4. (i) $p'_\infty \notin \varphi(\text{Spec } R_e)$.

(ii) $C' = \varphi(\text{Spec } R_e) \cup \{p'_\infty\}$.

(iii) $H^0(C' \setminus \{p'_\infty\}, \mathcal{O}_{C'}) = R_e$.

(iv) $p'_\infty \in D'_+(a)$ and it corresponds to a maximal ideal of $A'_{(a)}$.

The proof of this theorem is given in Appendix B.

By this theorem

$$C' = \varphi(\text{Spec } R_e) \cup D'_+(a)$$

is an affine open cover of C' . The rings R_e and $A'_{(a)}$ are integral domains, since they are subrings of $\mathbb{C}((z))$. Moreover $\dim C' = 1$ because, as we shall show in Lemma C.1, the quotient field of R_e is isomorphic to the quotient field of R which is the field of meromorphic functions on C . Using Proposition D.3 and the Riemann-Roch theorem we can easily prove that A' is generated over \mathbb{C} by a finite number of homogeneous elements. Then A' can be written as a quotient of polynomial ring by a homogeneous ideal. Therefore C' becomes a closed subscheme of a weighted projective space and consequently of a projective space. Thus C' is a projective integral scheme of dimension one, that is, C' is a projective integral curve.

Next we define a sheaf on C' .

Let $H^0(C, L_{\Delta,e}(np_\infty))$ be the space of meromorphic sections of $L_{\Delta,e}$ on C with a pole only at p_∞ of order at most n and

$$W_e(n) = W_e \cap H^0(C, L_{\Delta,e}(np_\infty)).$$

Define

$$W_e^{gr} = \bigoplus_{n=0}^{\infty} W_e(n).$$

Using Lemma 6.2 and the Riemann-Roch theorem it can easily be proved that W_e^{gr} is a finitely generated A' -module. Therefore W_e^{gr} defines a coherent $\mathcal{O}_{C'}$ module \mathcal{W}_e on C' such that

$$H^0(C' \setminus \{p'_\infty\}, \mathcal{W}_e) = W_e.$$

Moreover we can prove

Proposition 4.5. The $\mathcal{O}_{C'}$ -module \mathcal{W}_e is torsion free and of rank one.

The proof of this proposition is given in Appendix C.

Next we study the relation between C and C' .

A compact Riemann surface can be embedded into a projective space. Therefore there is a projective scheme corresponding to C which we denote by the same symbol C . In terms of $R = \cup_{n=0}^{\infty} R(n)$ C is described as

$$C = \text{Proj} A, \quad A = \bigoplus_{n=0}^{\infty} R(n).$$

There is an injective morphism, $\varphi: \text{Spec} R \rightarrow C$ given by a similar formula to (4.4) which we denote by the same symbol. The affine scheme $\text{Spec} A$ corresponds to $C \setminus \{p_{\infty}\}$. Similarly to p'_{∞} define

$$\tilde{p}_{\infty} = \bigoplus_{n=0}^{\infty} R(n-1),$$

where $R(n-1)$ is situated at the degree n component of A as in the previous case. Set

$$\begin{aligned} D_+(a) &= \{\mathcal{P} \in \text{Proj} A \mid a \notin \mathcal{P}\}, \\ A_{(a)} &= \{ua^{-n} \mid u \in R(mn), n \geq 0\}. \end{aligned}$$

As in the case of C' the following proposition holds.

Proposition 4.6. (i) $\tilde{p}_{\infty} \notin \varphi(\text{Spec} R)$.

(ii) $C = \varphi(\text{Spec} R) \cup \{\tilde{p}_{\infty}\}$.

(iii) $H^0(C \setminus \{\tilde{p}_{\infty}\}, \mathcal{O}_C) = R$.

(iv) $\tilde{p}_{\infty} \in D_+(a)$ and it corresponds to a maximal ideal of $A_{(a)}$.

All elements of the maximal ideal \tilde{p}_{∞} of $A_{(a)}$ vanish at p_{∞} . So \tilde{p}_{∞} can be identified with p_{∞} .

The inclusion map $A' \subset A$ induces a morphism $\psi: C \rightarrow C'$. Then

Proposition 4.7. The morphism $\psi: C \rightarrow C'$ gives the normalization of C' .

The proof of this proposition is given in Appendix D.

Finally we study the singularities of C' .

Let P be a point of the compact Riemann surface C such that $P \neq p_{\infty}$, $z(P) = p$ and $m_P \in \text{Spec} R$ the maximal ideal corresponding to P , that is,

$$m_P = \{f \in R \mid f(p) = 0\}.$$

Then $m' = \psi(m_P) = m_P \cap R_e$ is a maximal ideal of R_e since R is integral over R_e by Corollary D.4. We denote by R_{m_P} the localization of R at m_P etc. Then

Proposition 4.8. (i) If $P \neq P_i$ for any i , then $(R_e)_{m'} \simeq R_{m_P}$. In particular $(R_e)_{m'}$ is a normal ring and the closed point $m' \in \text{Spec}R_e$ is a non-singular point.

(ii) If $P = P_{i,j}$ and $n_i \geq 2$, then $(R_e)_{m'}$ is not a normal ring. In particular $m' \in \text{Spec}R_e$ is a singular point. Moreover $\psi^{-1}(m') = \{P_{i,1}, \dots, P_{i,n_i}\}$ in this case.

The proof of this proposition is given in Appendix E.

This proposition shows that C' is obtained from C by identifying the points $P_{i,1}, \dots, P_{i,n_i}$ for each i such that $n_i > 1$.

5 Solitons on elliptic backgrounds

As remarked in remark 3.3 if all t_i, p_j are real, $\tau_0(t) > 0$ for any t [6], $B_I \geq 0$ for any I and $p_1^{-1} < \dots < p_L^{-1}$, then $\tau(t)$ given by (3.5) is positive. Notice that $\tau(t)$ is a linear combination of $\{e^{\eta t} \tau_0(t - [p_I])\}$. Then, in the region of the xy -plane such that $e^{\eta t} \tau_0(t - [p_I])$ is dominant, $u(t) = 2\partial_x^2 \log \tau(t) \approx 2\partial_x^2 \log \tau_0(t)$ is the quasi-periodic wave corresponding to the shift of $\tau_0(t)$. On the boundary of two such domains, soliton like waves will appear as in the case of soliton solutions[16, 17]. Thus it is expected that $u(t)$ represents a soliton on a quasi-periodic background. In this section we verify it by a computer simulation in the case of genus one .

Let a, b be positive real numbers. Set $2\omega_1 = -ib$, $2\omega_2 = ab$, $\Omega = \omega_2/\omega_1 = ia$ and $\mathbb{L} = 2\omega_1\mathbb{Z} + 2\omega_2\mathbb{Z}$. Define

$$g_2 = 60 \sum_{\omega \in \mathbb{L}, \omega \neq 0} \frac{1}{\omega^4}, \quad g_3 = 140 \sum_{\omega \in \mathbb{L}, \omega \neq 0} \frac{1}{\omega^6}.$$

Then g_2, g_3 are real. Consider the algebraic curve C defined by the corresponding Weierstrass cubic

$$y^2 = 4x^3 - g_2x - g_3.$$

Let $\wp(u)$ be the Weierstrass elliptic function. Then $u \mapsto (\wp(u), \wp'(u))$ gives an isomorphism between the complex torus \mathbb{C}/\mathbb{L} and C , where $u = 0$ corresponds to $\infty \in C$. A basis of holomorphic one forms is $du = dx/y$. A canonical homology basis can be taken such that

$$\int_{\alpha} du = 2\omega_1, \quad \int_{\beta} du = 2\omega_2.$$

Therefore the normalized holomorphic one form is given by

$$dv = (2\omega_1)^{-1} du.$$

We take u as a local coordinate around ∞ . Then the corresponding solution of the KP hierarchy is given by

$$\tau_0(t) = e^{\frac{1}{2}q(t)} \theta \left(\frac{x}{2\omega_1} + e|\Omega \right), \quad (5.1)$$

where e is an arbitrary complex constant. If we take $e \in i\mathbb{R}$ and $x, t_j, j \geq 2$ to be real then $\tau_0(t)$ is real and positive.

In the present case $q_{i,j}$ defining $q(t)$ is described in the following way. Let

$$\theta_{11}(z|\Omega) = \sum_{n \in \mathbb{Z}} e^{\pi i \Omega (n + \frac{1}{2})^2 + 2\pi i (n + \frac{1}{2})(z + \frac{1}{2})}.$$

Sometimes $\theta_{11}(z|\Omega)$ is simply denoted by $\theta_{11}(z)$. It satisfies

$$\theta_{11}(-z) = -\theta_{11}(z), \quad \theta_{11}(z) = \theta'_{11}(0)z + O(z^3), \quad \theta'_{11}(0) \neq 0.$$

The prime form is written as

$$E(z_1, z_2) = \frac{2\omega_1}{\theta'_{11}(0)} \theta_{11}\left(\frac{z_2 - z_1}{2\omega_1}\right).$$

Therefore

$$\frac{\partial^2}{\partial z_1 \partial z_2} \log \frac{\theta_{11}\left(\frac{z_2 - z_1}{2\omega_1}\right)}{z_2 - z_1} = \sum_{i,j=1}^{\infty} q_{i,j} z_1^{i-1} z_2^{j-1}. \quad (5.2)$$

Let us take the elliptic solution (5.1) as $\tau_0(t)$ in the general formula (3.5). To neatly write $\tau(t)$ let us introduce

$$F(z) = \log \theta_{11}\left(\frac{z}{2\omega_1}\right), \quad F_j(z) = \frac{(-1)^{j-1}}{(j-1)!} F^{(j)}(z).$$

By Proposition 3.4 we have

Proposition 5.1. For $\tau_0(t)$ given by (5.1) the $\tau(t)$ defined by (3.3) is written as

$$\begin{aligned} \tau(t) &= c e^{\sum_{j=1}^{\infty} c_j t_j + \frac{1}{2} q(t)} \\ &\times \sum_I B_I C_I e^{\sum_{j \in I} \sum_{k=1}^{\infty} t_k F_k(p_j)} \theta\left(\frac{x - \sum_{j \in I} p_j}{2\omega_1} + e|\Omega\right). \end{aligned} \quad (5.3)$$

Here c, c_j are certain constants and

$$C_I = \prod_{j \in I} \frac{p_j}{\theta_{11}\left(\frac{p_j}{2\omega_1}\right)^N} \prod_{j,k \in I, j < k} \theta_{11}\left(\frac{p_k - p_j}{2\omega_1}\right).$$

We take $\{t_j\}, \{p_j\}$ are real and $e \in i\mathbb{R}$. If $B_I C_I \geq 0$ for any I and some of them is positive then the \sum_I part in the right hand side of (5.3) is positive and $u = 2\partial_x^2 \log \tau(t)$ is non-singular.

For the positivity of C_I we have

Proposition 5.2. If $p_1 < \dots < p_L$, $p_L - p_1 < ab$ and $|p_j| < ab$ for $1 \leq j \leq L$ then $C_I > 0$ for any I .

This proposition follows from the following lemma which can easily be proved.

Lemma 5.3. If $a > 0$, then $i\theta_{11}(ix|ia) > 0$ for $0 < x < a$.

Example 5.4. Take $M = N = 2$, $a = 1$, $b = 6$, $(p_1, p_2, p_3, p_4) = (0.31, 0.46, 0.81, 4.89)$, $t_j = 0$ for $j \geq 4$ and

$$B = \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ -2 & 1 \\ -3 & 0 \end{pmatrix}. \quad (5.4)$$

This B corresponds to (3) of §4.6.4 [17]. Notice that the matrix in [17] is the transpose of our matrix B . In this case $B_{12} = 1$, $B_{13} = 1$, $B_{14} = 0$, $B_{23} = 2$, $B_{24} = 3$, $B_{34} = 3$ and


$$(F^{(1)}(p_1), F^{(1)}(p_2), F^{(1)}(p_3), F^{(1)}(p_4)) = (3.25, 2.21, 1.30, 0.05). \quad (5.5)$$

The results of a computer simulation of $u(t)$ is given in figures 1,2,3. We can see the soliton corresponding to the matrix B on the periodic waves.


Example 5.5. Take M, N, a, b, p_j, t_j the same as those in Example 5.4. Consider B of the form

$$B = \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ -1 & 2 \\ -1 & 1 \end{pmatrix}. \quad (5.6)$$

For this matrix $B_{12} = 1$, $B_{13} = 2$, $B_{14} = 1$, $B_{23} = 1$, $B_{24} = 1$, $B_{34} = 1$. This B corresponds to (1) of §4.6.4 [17]. See figures 4,5,6.




(a) from above




(b) from an angle

Figure 1: Example 5.4 $t = -3$, $-20 \leq x \leq 20$, $0 \leq y \leq 15$




(a) from above




(b) from an angle

Figure 2: Example 5.4 $t = 0$, $-20 \leq x \leq 20$, $-10 \leq y \leq 10$




(a) from above




(b) from an angle

Figure 3: Example 5.4 $t = 3$, $-20 \leq x \leq 35$, $-25 \leq y \leq -5$




(a) from above




(b) from an angle

Figure 4: Example 5.5 $t = -2$, $-35 \leq x \leq 15$, $-5 \leq y \leq 15$




(a) from above




(b) from an angle

Figure 5: Example 5.5 $t = 0$, $-20 \leq x \leq 20$, $-10 \leq y \leq 10$



(a) from above



(b) from an angle

Figure 6: Example 5.5 $t = 2$, $-20 \leq x \leq 32$, $-25 \leq y \leq 0$

6 Proof of Theorem 3.7 (i)

In this section we freely use notation on sheaf cohomologies. Namely, for a holomorphic line bundle \mathcal{L} , distinct points S_i , $1 \leq i \leq m$, S'_j , $1 \leq j \leq m'$ on C , P and positive integers $\{m_i\}$, $\{m'_j\}$ we denote

$$H^0(C, \mathcal{L}(-\sum m_i S_i + \sum m'_j S'_j + *P))$$

the space of meromorphic sections of \mathcal{L} which have a zero at S_i of order at least m_i , a pole at S'_j of order at most m'_j and a pole at P of any order.

To prove Theorem 3.7 (i) it is sufficient to show

$$\dim W_e(n) = n - N \text{ for } n \gg 0, \quad (6.1)$$

by Corollary 2.2.

Lemma 6.1. There exists F_i in $H^0(C, L_{\Delta, e}(*p_\infty))$, $1 \leq i \leq L$ such that

$$F_i(p_j) = \delta_{i,j}.$$

The proof of this lemma is given in the end of this section.

For $1 \leq m \leq M$ set

$$\varphi_m = F_{N+m} - \sum_{i=1}^N b'_{N+m,i} F_i. \quad (6.2)$$

Then

$$\varphi_m \in W_e,$$

since

$$\varphi_m(q_l) = \delta_{m,l}, \quad \varphi_m(p_i) = -b'_{N+m,i}, \quad (6.3)$$

and

$$-\sum_{j=1}^M b'_{N+j,i} \varphi_m(q_j) = -b'_{N+m,i} = \varphi_m(p_i).$$

Notice that the vector space

$$H^0(C, L_{\Delta, e}(-\sum_{j=1}^L P_j + *p_\infty))$$

is a subspace of W_e , since elements of it vanish at all p_j and the linear equations imposed in W_e are trivially satisfied.

Lemma 6.2. The following equation holds,

$$W_e = H^0(C, L_{\Delta, \epsilon}(-\sum_{j=1}^L P_j + *p_\infty)) \oplus \oplus_{m=1}^M \mathbb{C}\varphi_m. \quad (6.4)$$

Proof. It is obvious that the right hand side is included in the left hand side. Let us prove the converse inclusion.

Take any $f \in W_e$ and set $f(q_i) = c_i$. Set

$$F = f - \sum_{m=1}^M c_m \varphi_m.$$

Then

$$F(q_i) = f(q_i) - \sum_{m=1}^M c_m \varphi_m(q_i) = c_i - c_i = 0.$$

Since f and φ_l are both in W_e , $F \in W_e$ and $F(p_i) = 0$ for $1 \leq i \leq N$. Therefore

$$F \in H^0(C, L_{\Delta, \epsilon}(-\sum_{j=1}^L P_j + *p_\infty))$$

and therefore f is in the right hand side of (6.4). \square

Take n larger than the order of a pole of φ_j at p_∞ for any j . Then

$$W_e(n) = H^0(C, L_{\Delta, \epsilon}(-\sum_{j=1}^L P_j + np_\infty)) \oplus \oplus_{m=1}^M \mathbb{C}\varphi_m,$$

and

$$\dim W_e(n) = \dim H^0(C, L_{\Delta, \epsilon}(-\sum_{j=1}^L P_j + np_\infty)) + M.$$

If we further take n larger than $g - 1 + L$,

$$\deg L_{\Delta, \epsilon}(-\sum_{j=1}^L P_j + np_\infty) = g - 1 - L + n > 2g - 2 = \deg \Omega^1,$$

then

$$H^1(C, L_{\Delta, \epsilon}(-\sum_{j=1}^L P_j + np_\infty)) = 0,$$

and, by the Riemann-Roch theorem,

$$\dim H^0(C, L_{\Delta, e}(-\sum_{j=1}^L P_j + np_{\infty})) = n - L.$$

Therefore, for $n \gg 0$,

$$\dim W_e(n) = n - L + M = n - N.$$

□

Proof of Lemma 6.1.

We first show that, for each $1 \leq i \leq L$, there exists an element G_i of $H^0(C, L_{\Delta, e}(*p_{\infty}))$ such that it has only a simple pole at P_i on $C \setminus \{p_{\infty}\}$.

By the Riemann-Roch theorem, for a sufficiently large n , we have

$$\begin{aligned} \dim H^0(C, L_{\Delta, e}(P_i + np_{\infty})) &= n + 1, \\ \dim H^0(C, L_{\Delta, e}(np_{\infty})) &= n. \end{aligned}$$

It means that there exists a meromorphic section of $L_{\Delta, e}$ which has a simple pole at P_i , a pole at p_{∞} of order at most n and has no other poles. Thus G_i exists.

Next we prove that there exists a meromorphic function on C such that it has a simple zero at every P_i , $1 \leq i \leq L$, and it is holomorphic on $C \setminus \{p_{\infty}\}$.

Again, if we take n sufficiently large, we have, by the Riemann-Roch theorem,

$$\begin{aligned} \dim H^0(C, \mathcal{O}(-\sum_{j=1}^L P_j + np_{\infty})) &= 1 - g - L + n, \\ \dim H^0(C, \mathcal{O}(-P_i - \sum_{j=1}^L P_j + np_{\infty})) &= -g - L + n. \end{aligned}$$

It follows that, for each $i \leq L$, there exists a meromorphic function h_i on C which has a simple zero at P_i , a zero at P_j , $j \neq i$, and is holomorphic on $C \setminus \{p_{\infty}\}$. We shall show that a desired h can be constructed as a linear combination of $\{h_i\}$.

Set

$$h = \sum_{i=1}^L \lambda_i h_i.$$

At each P_i take a local coordinate w and write

$$h_i = c_i w + O(w^2), \quad h_j = c_j w^{m_j} + O(w^{m_j+1}), \quad j \neq i,$$

where $c_j \neq 0$ for any j . Let $\{j \mid m_j = 1\} = \{j_1, \dots, j_r\}$, where we set $m_i = 1$. Then

$$h = (c_{j_1} \lambda_{j_1} + \dots + c_{j_r} \lambda_{j_r}) w + O(w^2),$$

and the condition that h has a simple zero at P_i is

$$c_{j_1}\lambda_{j_1} + \cdots + c_{j_r}\lambda_{j_r} \neq 0. \quad (6.5)$$

This is an open condition for $\{\lambda_j\}$. Thus there exists $\{\lambda_j\}$ such that h has a simple zero at every P_i .

Choosing one set of $\{h, G_i | 1 \leq i \leq L\}$ define the element of $H^0(C, L_{\Delta, e}(*p_\infty))$ by

$$F_i = a_i h G_i, \quad a_i = (h G_i)(p_i)^{-1}.$$

Obviously they satisfy $F_i(p_j) = \delta_{i,j}$. □

7 Proof of Theorem 3.7 (ii)

Let $\Psi(t, z)$ be the wave function of $\tau(t)$,

$$\Psi(t, z) = \frac{\tau(t - [z])}{\tau(t)} e^{\eta(t, z^{-1})}. \quad (7.1)$$

We first show

Lemma 7.1. For $1 \leq i \leq N$

$$\Psi(t, p_i) = - \sum_{j=1}^M \tilde{b}_{N+j,i} \Psi(t, p_{N+j}), \quad \tilde{b}_{N+j,i} = b_{N+j,i} (p_i p_{N+j}^{-1})^N \quad (7.2)$$

Proof. Substitute (7.1) into (7.2) we have the equation for $\tau(t)$,

$$\tau(t - [p_i]) e^{\eta_i} = - \sum_{j=1}^M \tilde{b}_{N+j,i} \tau(t - [p_{N+j}]) e^{\eta_{N+j}}. \quad (7.3)$$

Let us prove this equation.

By (3.5) we have, using $e^{-\eta([z], p^{-1})} = 1 - p^{-1}z$,

$$\tau(t - [z]) = \sum_I B_I \Delta_I^- \prod_{j \in I} (1 - p_j^{-1}z) e^{\eta_I} \tau_0(t - [z] - [p_I]). \quad (7.4)$$

Substituting $z = p_i$, $1 \leq i \leq N$ and multiplying by e^{η_i} we get

$$\begin{aligned} \tau(t - [p_i]) e^{\eta_i} &= p_i^N \sum_I B_I \Delta_I^- \prod_{j \in I} (p_i^{-1} - p_j^{-1}) e^{\eta_I + \eta_i} \tau_0(t - [p_i] - [p_I]) \\ &= p_i^N \sum_{i \notin I} B_I \Delta_{(I,i)}^- e^{\eta(I,i)} \tau_0(t - [p_{(I,i)}]). \end{aligned} \quad (7.5)$$

$$\begin{aligned}
&= \frac{1}{r!} \sum_{k=1}^r \sum_{j_1, \dots, j_r=1}^M (-1)^{N-r+k+i} b_{N+j_k, i} B_{(I', N+j_1, \dots, \widehat{N+j_k}, \dots, N+j_r)}^{(i)} \Delta_{(I', N+j_1, \dots, N+j_r, i)}^- \\
&\quad \times T(I', N+j_1, \dots, N+j_r, i). \tag{7.10}
\end{aligned}$$

Use

$$\Delta_{(I', N+j_1, \dots, N+j_r, i)}^- = (-1)^{r-k} \Delta_{(I', N+j_1, \dots, \widehat{N+j_k}, \dots, N+j_r, N+j_k, i)}^-$$

and change the names of indices as $j_k \rightarrow j$, $j_1, \dots, \widehat{j_k}, \dots, j_r \rightarrow j_1, \dots, j_{r-1}$. We get

$$\begin{aligned}
&\text{RHS of (7.10)} \\
&= \frac{1}{r!} \sum_{k=1}^r (-1)^{N+i} \sum_{j=1}^M b_{N+j, i} \sum_{j_1, \dots, j_{r-1}=1}^M B_{(I', N+j_1, \dots, N+j_{r-1})}^{(i)} \Delta_{(I', N+j_1, \dots, N+j_{r-1}, N+j, i)}^- \\
&\quad \times T_{(I', N+j_1, \dots, N+j_{r-1}, N+j, i)}.
\end{aligned}$$

Since the summand does not depend on k , the sum in k gives r times of the summand. Rewrite the summation in $1 \leq j_1, \dots, j_{r-1} \leq M$ to $(r-1)!$ times the summation in (j_1, \dots, j_{r-1}) with $1 < j_1 < \dots < j_{r-1} \leq M$. Substitute it into (7.7) and get

$$\begin{aligned}
&\text{RHS of (7.7)} \\
&= (-1)^{N+i} p_i^N \sum_{j=1}^N b_{N+j, i} \sum_{r=1}^M \sum_{I' \in \binom{[N]}{N-r}, i \notin I'} \sum_{1 \leq j_1 < \dots < j_{r-1} \leq M} B_{(I', N+j_1, \dots, N+j_{r-1})}^{(i)} \\
&\quad \times \Delta_{(I', N+j_1, \dots, N+j_{r-1}, N+j, i)}^- T_{(I', N+j_1, \dots, N+j_{r-1}, N+j, i)}.
\end{aligned}$$

Notice that taking the summation over $r, I', \{j_k\}$ is equivalent to taking the summation over $I' \in \binom{[L]}{N-1}$ with $i, N+j \notin I'$. Thus

$$\begin{aligned}
&\tau(t - [p_i]) e^{\eta_i} \\
&= \text{RHS of (7.7)} \\
&= (-1)^{N+i} p_i^N \sum_{j=1}^N b_{N+j, i} \sum_{I' \in \binom{[L]}{N-1}, i, N+j \notin I'} B_{I'}^{(i)} \Delta_{(I', N+j, i)}^- T(I', N+j, i). \tag{7.11}
\end{aligned}$$

Next, by replacing i by $N+j$ in (7.5) and recalling the definition (7.6) of $T(I, i)$, we have

$$\tau(t - [p_{N+j}]) e^{\eta_{N+j}} = p_{N+j}^N \sum_{N+j \notin I} B_I \Delta_{(I, N+j)}^- T(I, N+j). \tag{7.12}$$

It follows that

$$\sum_{j=1}^M \tilde{b}_{N+j, i} \tau(t - [p_{N+j}]) e^{\eta_{N+j}}$$

$$= p_i^N \sum_{j=1}^M b_{N+j,i} \sum_{N+j \notin I} B_I \Delta_{(I,N+j)}^- T(I, N+j) \quad (7.13)$$

$$= I_+ + I_-, \quad (7.14)$$

where I_+ is the part of the RHS of (7.13) such that I includes i in the summation over I and I_- the part where I does not include i .

We show that I_+ is equal to $-\tau(t - [p_i])e^{\eta_i}$ and $I_- = 0$.

Let us first consider I_+ . Separating i from I we have

$$I_+ = p_i^N \sum_{j=1}^M b_{N+j,i} \sum_{I' \in \binom{[L]}{N-1}, i, N+j \notin I'} B_{(I',i)} \Delta_{(I',i,N+j)}^- T(I', i, N+j).$$

Since the N -th row vector of $B_{(I',i)}$ is the i -th unit vector, we have

$$B_{(I',i)} = (-1)^{N+i} B_{I'}^{(i)}.$$

Therefore

$$\begin{aligned} I_+ &= (-1)^{N+i} p_i^N \sum_{j=1}^M b_{N+j,i} \sum_{I' \in \binom{[L]}{N-1}, i, N+j \notin I'} B_{I'}^{(i)} \Delta_{(I',i,N+j)}^- T(I', i, N+j) \\ &= \text{-RHS of (7.11)} \\ &= -\tau(t - [p_i])e^{\eta_i}, \end{aligned} \quad (7.15)$$

where $\Delta_{(I',i,N+j)}^- = -\Delta_{(I',N+j,i)}^-$ is used.

Next let us consider I_- . In a similar computation to deriving (7.11) we have

$$\begin{aligned} I_- &= (-1)^{N+i} \sum_{j=1}^M b_{N+j,i} \sum_{j'=1}^M b_{N+j',i} \sum_{I' \in \binom{[L]}{N-1}, i, N+j', N+j \notin I'} B_{I'}^{(i)} \Delta_{(I',N+j',N+j)}^- \\ &\quad \times T(I', N+j', N+j) \\ &= (-1)^{N+i} \sum_{j,j'=1}^M b_{N+j,i} b_{N+j',i} \sum_{I' \in \binom{[L]}{N-1}, i, N+j', N+j \notin I'} B_{I'}^{(i)} \Delta_{(I',N+j',N+j)}^- \\ &\quad \times T(I', N+j', N+j). \end{aligned}$$

Since $b_{N+j,i} b_{N+j',i}$ is symmetric in j, j' and the remaining part is skew symmetric in j, j' , the last summation in j, j' becomes zero. Therefore $I_- = 0$. We, then, have (7.3) by (7.14), (7.15). \square

By Lemma 3.5 we have

Lemma 7.2. Let $N \geq 1$, $Q_j \in C$, $1 \leq j \leq N$ and $z_j = z(Q_j)$. Then

$$\begin{aligned}
& \tau_0(t - [z] - \sum_{j=1}^N [z_j]) e^{\eta(t, z^{-1})} \\
&= \left(\frac{z}{E(0, z)} \right)^{N+1} \prod_{j=1}^N \frac{E(z, z_j)}{z_j - z} \prod_{j=1}^N \frac{z_j}{E(0, z_j)} e^{q(\sum_{j=1}^N [z_j]) - \sum_{j=1}^N Q(t|[z_j]) + \frac{1}{2}q(t)} \\
& \quad \times \theta(\mathcal{V}t - I(p) - \sum_{j=1}^N I(Q_j) + e) e^{\sum_{j=1}^{\infty} t_j \int^p d\tilde{r}_j}. \tag{7.16}
\end{aligned}$$

We use (7.16) to compute (7.4) and get

$$\begin{aligned}
& \tau(t - [z]) e^{\eta(t, z^{-1})} \\
&= z^{N+1} \sum_{J \in \binom{[L]}{N}} B_J \Delta_J^- e^{\eta_J} \prod_{j \in J} \frac{E(z, p_j)}{E(0, z) E(0, p_j)} e^{q(\sum_{j \in J} [p_j]) - \sum_{j \in J} Q(t|[p_j]) + \frac{1}{2}q(t)} \\
& \quad \times \frac{1}{E(0, z)} \theta(\mathcal{V}t - I(p) - \sum_{j \in J} I(Q_j) + e) e^{\sum_{j=1}^{\infty} t_j \int^p d\tilde{r}_j}. \tag{7.17}
\end{aligned}$$

Set

$$\Psi'(t, z) = z^{-N-1} \Psi(t, z).$$

Then (7.17) shows that the expansion coefficients of $\tau(t) \Psi'(t, z) = z^{-N-1} \tau(t - [z]) e^{\eta(t, z^{-1})}$ belong to $H^0(C, L_{\Delta, e}(*p_\infty))$. Rewriting the equation (7.2) in terms of $\Psi'(t, z)$ we have

$$\Psi'(t, p_i) = - \sum_{j=1}^M b'_{N+j, i} \Psi'(t, q_j).$$

It means that the expansion coefficients of $\tau(t) \Psi'(t, z)$ are in W_e . Thus the expansion coefficients of $\tau(t) \Psi(t, z)$ are in $z^{N+1} W_e = U_e$. Since $U_e \in UGM$ by (1) of Theorem 3.7 and the strict inclusion relation is impossible for points of UGM, U_e is the point of UGM corresponding to $\tau(-t)$. \square

A Proof of Proposition 4.1

We first show that R_e is contained in the RHS of (4.1). Let $f \in R_e$ and $\varphi_m \in W_e$ be defined in (6.2). Then $f\varphi_m \in W_e$. By (6.3) we see that

$$(f\varphi_m)(p_i) = - \sum_{j=1}^M b'_{N+j, i} (f\varphi_m)(q_j)$$

is equivalent to

$$f(p_i)b'_{N+m,i} = b'_{N+m,i}f(q_m). \quad (\text{A.1})$$

Therefore f is contained in the RHS of (4.1).

Let us prove the converse inclusion. Let f be an element of the RHS of (4.1) and $F \in W_e$. Notice that the equation (A.1) holds for any i, m . Then

$$\begin{aligned} -\sum_{j=1}^M b'_{N+j,i}(fF)(q_j) &= -\sum_{j=1}^M b'_{N+j,i}f(q_j)F(q_j) \\ &= -\sum_{j=1}^M b'_{N+j,i}f(p_i)F(q_j) \\ &= f(p_i)\left(-\sum_{j=1}^M b'_{N+j,i}F(q_j)\right) \\ &= f(p_i)F(p_i) = (fF)(p_i) \end{aligned}$$

which means $fF \in W_e$. Thus $f \in R_e$. □

B Proof of Theorem 4.4

(iii) follows from (ii). Let us prove (i), (ii), (iv).

(i) It can be easily proved that if $\varphi(\mathcal{P}) = p'_\infty$ for some $\mathcal{P} \in \text{Spec}R_e$ then $\mathcal{P} = R_e$. It is absurd.

(ii) Let $\mathcal{P} = \bigoplus_{n=0}^{\infty} \mathcal{P}^{(n)} \in C'$ such that $\mathcal{P}' \neq p'_\infty$. Set

$$q = \bigcup_{n=0}^{\infty} \mathcal{P}^{(n)} \subset R_e,$$

which obviously becomes an ideal of R_e . It is sufficient to prove the following lemma.

Lemma B.1. (i) If $x, y \in R_e$ satisfies $xy \in q$, either $x \in q$ or $y \in q$.

(ii) $q \neq R_e$.

(iii) $\varphi(q) = \mathcal{P}$.

Proof. (i) We can assume that

$$x = z^{-m} + O(z^{-m+1}), \quad y = z^{-n} + O(z^{-n+1})$$

with $m, n \geq 1$. Then

$$xy \in R_e(m+n) \setminus R_e(m+n-1). \quad (\text{B.1})$$

Since $xy \in q$, there exists $N \geq 0$ such that

$$xy \in \mathcal{P}^{(N)} \subset R_e(N).$$

By (B.1) we have $N \geq m + n$. Set

$$N - (m + n) = k \geq 0.$$

Then

$$x \in R_e(m) \subset R_e(m + k), \quad y \in R_e(n).$$

So we consider x and y as homogeneous elements of A' with degree $m + k$ and n respectively. Then

$$xy \in \mathcal{P}^{(N)} \subset A'.$$

Since \mathcal{P} is a prime ideal of A' , $x \in \mathcal{P}$ or $y \in \mathcal{P}$. Therefore $x \in \mathcal{P} \cap R_e(m + k) = \mathcal{P}^{(m+k)}$ or $y \in \mathcal{P} \cap R_e(n) = \mathcal{P}^{(n)}$. It means that $x \in \mathcal{P}$ or $y \in \mathcal{P}$.

(ii) Notice that $1 \in R_e(n)$ for any $n \geq 0$. If we consider 1 as a homogeneous element of A' with degree n we denote it by $1^{(n)}$. We shall show

$$1^{(n)} \notin \mathcal{P}^{(n)}, \quad n \geq 0. \quad (\text{B.2})$$

Since $\mathcal{P} \neq A'$, $1^{(0)} \notin \mathcal{P}^{(0)}$.

Let us consider the case $n = 1$. Suppose that $1^{(1)} \in \mathcal{P}^{(1)}$. Notice that $A'1^{(1)} = p'_\infty$. Therefore

$$p'_\infty \subsetneq \mathcal{P},$$

since $\mathcal{P} \neq p'_\infty$. Then there exists $k \geq 1$ and $f_k \in \mathcal{P}^{(k)}$ such that $f_k \notin R_e(k - 1)$, that is,

$$f_k = z^{-k} + O(z^{-k+1}).$$

By the Riemann-Roch theorem we have

$$\dim R_e(n)/R_e(n - 1) = 1. \quad n \gg 0, \quad (\text{B.3})$$

It follows that, for all sufficiently large n , there exists $h_n \in R_e(n)$ such that

$$h_n = z^{-n} + O(z^{-n+1}).$$

Then $h_n f_k \in \mathcal{P}^{(n+k)}$ and $h_n f_k \in R_e(n + k) \setminus R_e(n + k - 1)$. Taking into account that $\mathcal{P}^{(n+k)} \supset R_e(n + k - 1)$ we see that

$$\mathcal{P}^{(n)} = R_e(n), \quad n \gg 0. \quad (\text{B.4})$$

On the other hand

$$\mathcal{P} \not\subseteq \bigoplus_{n=1}^{\infty} R_e(n)$$

since $\mathcal{P} \in C'$. It follows that there exists $N \geq 1$ and $F_N \in R_e(N)$ such that $F_N \notin \mathcal{P}^{(N)}$. Since $\mathcal{P}^{(N)} \supset R_e(N-1)$, $F_N \in R_e(N) \setminus R_e(N-1)$, that is,

$$F_N = z^{-N} + O(z^{-N+1}).$$

Since \mathcal{P} is a prime ideal,

$$F_N^m \notin \mathcal{P}^{(Nm)},$$

for any $m \geq 1$ which contradicts (B.4). Thus $1^{(1)} \notin \mathcal{P}^{(1)}$.

Notice that $(1^{(1)})^n = 1^{(n)}$ for $n \geq 2$. Thus $1^{(n)} \notin \mathcal{P}^{(n)}$ and (B.2) has been proved.

Then $1 \notin q$ and $q \neq R_e$.

(iii) We have to prove that $q \cap R_e(n) = \mathcal{P}^{(n)}$ for any $n \geq 0$. Suppose that this does not hold. Then there exists $N \geq 1$ such that

$$q \cap R_e(N) \neq \mathcal{P}^{(N)}.$$

Since the right hand side is contained in the left hand side, it means

$$q \cap R_e(N) \supsetneq \mathcal{P}^{(N)}.$$

Therefore there exists $x \in q \cap R_e(N)$ such that $x \notin \mathcal{P}^{(N)}$. Since $x \in q$ there exists M such that $x \in \mathcal{P}^{(M)}$. Then $N < M$. In fact if $N \geq M$, $x \in \mathcal{P}^{(M)} \subset \mathcal{P}^{(N)}$. Since $x \in q \cap R_e(N)$, we can consider x as a homogeneous element of A' with degree N . Then $x \notin \mathcal{P}^{(N)}$ and $1^{(M-N)}x \notin \mathcal{P}^{(M-N)}$ but

$$1^{(M-N)}x = x \in \mathcal{P}^{(M)},$$

which is absurd. Thus $q \cap R_e(n) = \mathcal{P}^{(n)}$ for any n and (iii) of the lemma is proved. \square

Proof of Theorem 4.4 (iv).

It is obvious that $a \notin p'_\infty$ and $p'_\infty \in D_+(a)$. Let

$$(p'_\infty)_{(a)} = \sum_{n=0}^{\infty} \frac{R_e(nm-1)}{a^n}$$

be the image of p'_∞ in $A'_{(a)}$. Any element of $f \in R_e(nm)$ satisfies $f - ca^n \in R_e(nm-1)$ for some constant $c \in \mathbb{C}$. It means that $f/a^n = c$ modulo $(p_\infty)_{(a)}$. Since $(p_\infty)_{(a)} \neq A_{(a)}$, this means that $A_{(a)}/(p'_\infty)_{(a)} \simeq \mathbb{C}$. Thus $(p'_\infty)_{(a)}$ is a maximal ideal of $A_{(a)}$ and (iv) is proved, \square

C Proof of Proposition 4.5

Both W_e and R_e is a subspace of $\mathbb{C}((z))$ and the R_e module structure of W_e is given by the ring structure of $\mathbb{C}((z))$. Therefore W_e is a torsion free R_e module. That W_e is a torsion free $\mathcal{O}_{C'}$ module follows from this.

Let K' be the quotient field of R_e . In order to prove that the rank of W_e is one, it is sufficient to show, by the definition, that $\dim_{K'} K' \otimes_{R_e} W_e = 1$.

Lemma C.1. Let K be the quotient field of R . Then $K' = K$.

Proof. Since $R_e \subset R$, $K' \subset K$. Let us prove the converse inclusion. Take any $f \in K$. Let the pole divisor of f be

$$m_1 R_1 + \cdots + m_s R_s + m_\infty p_\infty, \quad m_i > 0, R_i \in C, R_i \neq p_\infty.$$

Take n sufficiently large such that there exists non zero $F \in H^0(C, \mathcal{O}(-\sum_{i=1}^s m_i R_i - \sum_{i,j} P_{i,j} + np_\infty))$. Then $F \in R_e$ since $F(p_{i,j}) = 0$ for all i, j . Set $h = fF$. Then

$$h \in H^0(C, \mathcal{O}(-\sum_{i,j} P_{i,j} + *p_\infty)) \subset R_e.$$

Thus $f = h/F \in K'$. Thus $K \subset K'$. □

Let us continue the proof of Proposition 4.5. Take any nonzero $f_1 \in W_e$. Notice that K is the field of meromorphic functions on C . Therefore, for any $f_2 \in W_e$, $f_2/f_1 \in K$ and $f_2 \in Kf_1$. By Lemma C.1 we have $W_e \subset Kf_1 = K'f_1$. Thus $K' \otimes_{R_e} W_e = K'f_1$ and $\dim_{K'} K' \otimes_{R_e} W_e = 1$. □

D Proof of Proposition 4.7

We begin by determining the structure of R and R_e .

Lemma D.1. For any (i, j) , $1 \leq i \leq s$, $1 \leq j \leq n_i$, there exists $H_{i,j} \in R$ such that

$$H_{i,j}(p_{k,l}) = \delta_{i,k} \delta_{j,l}.$$

Proof. By the Riemann-Roch formula, for all sufficiently large n , we have

$$\begin{aligned} \dim H^0\left(C, \mathcal{O}\left(-\sum P_{k,l} + np_\infty\right)\right) &= n - L + 1 - g, \\ \dim H^0\left(C, \mathcal{O}\left(-\sum_{(k,l) \neq (i,j)} P_{k,l} + np_\infty\right)\right) &= n - L + 2 - g. \end{aligned}$$

A non-zero element of the latter space which does not belong to the former space satisfies the required property if it is adjusted by a constant multiple. □

Let

$$H_i = H_{i,1} + \cdots + H_{i,n_i}.$$

Then

Lemma D.2. (i) $H_i(p_{i,j}) = 1$ for $1 \leq j \leq n_i$ and $H_i(p_{k,l}) = 0$ if $k \neq i$.

(ii) $H_i \in R_e$.

(iii) $\{H_i \mid 1 \leq i \leq s\}$ is linearly independent.

The lemma can be easily proved from the definition of H_i . So we leave the proof to the reader.

Proposition D.3. (i) $R = H^0(C, \mathcal{O}(-\sum P_{i,j} + *p_\infty)) \oplus \oplus_{i,j} \mathbb{C}H_{i,j}$.

(ii) $R_e = H^0(C, \mathcal{O}(-\sum P_{i,j} + *p_\infty)) \oplus \oplus_{i=1}^s \mathbb{C}H_i$.

Proof. (i) It is sufficient to prove that the left hand side is contained in the right hand side. Take any $f \in R$. Set $f(p_{i,j}) = c_{i,j}$ and $f' = f - \sum c_{i,j}H_{i,j}$. Then $f'(p_{i,j}) = 0$ for any i, j . Thus

$$f' \in H^0\left(C, \mathcal{O}(-\sum P_{i,j} + *p_\infty)\right),$$

which shows that f is contained in the RHS of (i).

(ii) is similarly proved. □

Since R_e is a subring of R , R is considered as an R_e -module. Then

Corollary D.4. The ring R is a finitely generated R_e module.

Proof. By Proposition D.3 we have

$$R = R_e 1 + \sum R_e H_{i,j},$$

which shows the assertion of the corollary. □

Since C is non-singular R is integrally closed in K .

Then we have

Corollary D.5. The ring R is the integral closure of R_e in K .

Proof. By Corollary D.4 R is integral over R_e . An integral element of K over R_e is integral over R and therefore it belongs to R since R is integrally closed. Thus R is the integral closure in K . □

Take $a \in R_e(m) \subset R(m)$ as in (4.5). Consider a as an element of A' and A with the degree m .

Similarly to the above corollary we can prove the following.

Proposition D.6. The ring $A_{(a)}$ is a finitely generated $A'_{(a)}$ module and it is the integral closure of $A'_{(a)}$ in K .

By Corollary D.5 and Proposition D.6 we have Proposition 4.7.

E Proof of Proposition 4.8

(i) Let

$$S = R - m = \{f \in R \mid f(p) \neq 0\}, \quad S' = R_e - m' = \{f \in R_e \mid f(p) \neq 0\}.$$

Then, $S' \subset S$ and, by definition, $R_m = S^{-1}R$, $(R_e)_{m'} = S'^{-1}R_e$. Therefore $(R_e)_{m'} \subset R_m$. Let us prove the converse inclusion. Take any $f \in R_m$ and write it as

$$f = \frac{F}{G}, \quad F \in R, G \in S.$$

Notice that, by the Riemann-Roch theorem, there exists $H \in R$ such that

$$H \in H^0\left(C, \mathcal{O}\left(-\sum P_{i,j} + np_\infty\right)\right), \quad H(p) \neq 0, \quad (\text{E.1})$$

if n is sufficiently large. Then $FH, GH \in R_e$, $(GH)(p) \neq 0$ and $f = FH/GH \in S'^{-1}R_e$. Thus $R_m \subset (R_e)_{m'}$.

(ii) Since R is the integral closure of R_e , the integral closure of $(R_e)_{m'} = S'^{-1}R_e$ is $S'^{-1}R$ (c.f. Proposition 2.1 of [13]). Consider $H_{i,j} \in R \subset S'^{-1}R$ of Lemma D.1. It is not in $S'^{-1}R_e$. In fact if $H_{i,j} \in S'^{-1}R_e$ then $fH_{i,j} \in R_e$ for some $f \in S'$. Then $(fH_{i,j})(p_{i,j}) = f(p_{i,j}) \neq 0$ and $(fH_{i,j})(p_{i,j'}) = f(p_{i,j'})H(p_{i,j'}) = 0$ for $j' \neq j$ which contradicts $fH_{i,j} \in R_e$. Thus $S'^{-1}R_e \neq S'^{-1}R$ and $S'^{-1}R_e$ is not a normal ring.

Let us prove the last statement of (ii). Obviously $m_{P_{i,j'}} \in \psi^{-1}(m')$ for any j' . Since R is integral over R_e , each element of $\psi^{-1}(m')$ is a maximal ideal.

Let Q be a point of the Riemann surface C such that $Q \neq P_{i,j'}$ for any j' and $z(Q) = q$. Suppose that $\psi(m_Q) = m'$. Similarly to (E.1) there exists $H \in R_e$ such that $H(p_{i',j'}) = 0$ for any i', j' and $H(q) \neq 0$. Then $H \in m'$ but $H \notin \psi(m_Q)$. It contradicts the assumption. Thus the assertion is proved. \square

Acknowledgements

Parts of the results of this paper were presented in a series of lectures at University of Tokyo in July, 2022 and at Nagoya University in July, 2023. I would like to thank Junichi Shiraishi and Masashi Hamanaka for invitations and hospitality. I would also like to thank Hiroaki Kanno, Yasuhiko Yamada, Shintaro Yanagida for their interests. I would especially like to thank Yuji Kodama for explaining the contents of the paper [19] as well as related results [18] and for his valuable comments when I gave lectures at Nagoya University. I am also grateful to Yuji Kodama for letting me know his theory on soliton solutions of the KP equation some years ago. That was the starting point of this study. This work was supported by JSPS KAKENHI Grant Number JP19K03528.

References

- [1] S. Abenda and P. Grinevich, Rational degenerations of M-curves, totally positive Grassmannians and KP-solitons, arXiv:1506.00563.
- [2] S. Abenda and P. Grinevich, Real soliton lattices of KP-II and desingularization of spectral curves: the $Gr^{TP}(2, 4)$ case, *Proc. Steklov Inst. Math.* **302;1** (2018) 1-15.
- [3] D. Agostini, C. Fevola, Y. Mandelshtam and B. Sturmfels, KP solitons from tropical Limits, arXiv:2101.10392.
- [4] S. Cakravarty and Y. Kodama, Classification of the line-soliton solutions of KP-II, *J. Phys. A* **41** (2008), 275209.
- [5] E. Date, M. Kashiwara, M. Jimbo, and T. Miwa, Transformation groups for soliton equations, in “Nonlinear Integrable Systems — Classical Theory and Quantum Theory”, M. Jimbo and T. Miwa (eds.), World Sci., Singapore, 1983, pp.39–119.
- [6] B. Dubrovin and S. Natanzon, Real theta function solutions of the Kadomtsev-Petviashvili equation, *Math. USSR Izvestiya* **32-2** (1989) 269-288.
- [7] G.A. El, Soliton gas in integrable dispersive hydrodynamics, *J. Stat. Mechanics: Theory and Experiment* **2021** (2021) 114001.
- [8] J. Fay, Theta-Functions on Riemann Surfaces, Springer Lecture Notes in Mathematics **352**, 1973.
- [9] R. Hartshorne, Algebraic Geometry, Springer, 1977.
- [10] M. Hofer, A. Mukalica and D. Pelinovsky, KdV breathers on a cnoidal wave background, arXiv:2301.08154.
- [11] Takashi Ichikawa, Periods of tropical curves and associated KP solutions, *Comm. Math. Phys.*, **Vol. 402** (2023), 1707-1723.
- [12] Takashi Ichikawa, Tau functions and KP solutions on families of algebraic curves, arXiv:2208.07013.
- [13] S. Iitaka, Algebraic Geometry, Graduate texts in mathematics 76, Springer, 1982.
- [14] Y. Kawamata, Theory of algebraic varieties, Mathematics of the 21st century, 3rd edition, Kyoritsu, 2001 (in Japanese).

- [15] N. Kawamoto, Y. Namikawa, A. Tsuchiya and Y. Yamada, Geometric realization of conformal field theory on Riemann surfaces, *Comm. Math. Phys.* **116** (1988) 247-308.
- [16] Y. Kodama, KP solitons and the Grassmannians, Springer, 2017.
- [17] Y. Kodama, Solitons in two-dimensional shallow water, CBMS-NSF regional conference series in applied mathematics 92, SIAM, 2018.
- [18] Y. Kodama, talk at Nagoya University, July 20, 2023.
- [19] Y. Kodama, KP solitons and the Riemann theta functions, arXiv:2308.06902.
- [20] Y. Kodama and L. Williams, KP solitons, total positivity and cluster algebras, *Proc. Nat. Acad. Sci. USA* **108**(22) (2011), 8984-8989.
- [21] Y. Kodama and L. Williams, KP solitons and total positivity for the Grassmannian, *Invent Math.* **198** (2014), 637-699.
- [22] I.M. Krichever, Methods of algebraic geometry in the theory of nonlinear equations, *Russ. Math. Surv.*, **Vol. 32** (1977), 185-213
- [23] Yu. Manin, Algebraic aspects of nonlinear differential equations, English translation of *Itogi Nauki i Tekhniki, Sovremennye Problemy Matematiki*, **Vol. 11** (1978), 5-152.
- [24] H. Matsumura, Commutative rings, Kyoritsu, 1980 (in Japanese).
- [25] M. Mulase, Cohomological structure in soliton equations and Jacobian varieties, *J. Diff. Geom.* **19** (1984) 403-430.
- [26] M. Mulase, Category of vector bundles on algebraic curves and infinite dimensional Grassmannians, *Int. J. Math.* **01-3** (1990) 293-342.
- [27] D. Mumford, An algebro-geometric construction of commuting operators and of solutions to the Toda lattice equation, Korteweg deVries equation and related nonlinear equation, Intl. Symp. on Algebraic Geometry, Kyoto, 1977, 115-153.
- [28] D. Mumford, Tata lectures on theta I, Birkhauser, 1983.
- [29] D. Mumford, Tata lectures on theta II, Birkhauser, 1983.
- [30] A. Nakayashiki, Tau function approach to theta functions, *Int. Math. Res. Not. IMRN* **2016-17** (2016), 5202–5248.
- [31] A. Nakayashiki, Degeneration of trigonal curves and solutions of the KP-hierarchy, *Nonlinearity* **31** (2018), 3567-3590.

- [32] A. Nakayashiki, On reducible degeneration of hyperelliptic curves and soliton solutions, *SIGMA* **15** (2019), 009, 18 pages, arXiv:1808.06748.
- [33] A. Nakayashiki, One step degeneration of trigonal curves and mixing solitons and quasi-periodic solutions of the KP equation, Geometric methods in physics XXXVIII, P. Kielanowski, A. Odziejewicz and E. Previato eds. Springer, 2020. arXiv:1911.06524
- [34] A. Nakayashiki, Tau functions of $(n, 1)$ curves and soliton solutions on non-zero constant backgrounds, *Lett. Math. Phys.* **111** (2021) Article number: 85.
- [35] T. Shiota, Characterization of jacobian varieties in terms of soliton equations, *Inv. Math.* **83** (1986) 333-382.
- [36] M. Sato and Y. Sato, Soliton equations as dynamical systems on infinite dimensional Grassmann manifold, in “Nolinear Partial Differential Equations in Applied Sciences”, P. D. Lax, H. Fujita and G. Strang (eds.), North-Holland, Amsterdam, and Kinokuniya, Tokyo, 1982, pp.259–271.
- [37] G. Segal and G. Wilson, Loop groups and equations of KdV type, *Publ. Math. IHES* **61** (1985) 5-65.
- [38] S. Tanaka and E. Date, KdV equation, Kinokuniya,1979 (in Japanese).