

# About optimization of methods for mixed derivatives of bivariate functions

Y.V. SEMENOVA<sup>†</sup>, S.G. SOLODKY<sup>†,‡</sup>

<sup>†</sup>Institute of Mathematics, National Academy of Sciences of Ukraine, Kyiv

<sup>‡</sup>University of Giessen, Department of Mathematics, Giessen, Germany

ABSTRACT.

The problem of optimal recovering high-order mixed derivatives of bivariate functions with finite smoothness is studied. On the basis of the truncation method, an algorithm for numerical differentiation is constructed, which is order-optimal both in the sense of accuracy and in terms of the amount of involved Galerkin information.

## 1. Description of the problem

Currently, many research activities on the problem of stable numerical differentiation have been taking place due to the importance of this tool in such areas of science and technology as finance, mathematical physics, image processing, analytical chemistry, viscous elastic mechanics, reliability analysis, pattern recognition, and many others. Among these investigations we highlight [4], which is the first publication on numerical differentiation in terms of the theory of ill-posed problems. Further research [4] has been continued in numerous publications on numerical differentiation (see, for example, [21], [30], [7], [8], [1], [9], [12], [31], [32], [10], [24]), covering different classes of the functions and the types of proposed methods. Despite the abundance of works on this topic, the problem for recovery of high order derivatives was considered only in a few publications, among which we note [5], [22], [11], [19], [12] and [24]. In particular, the results of [24] have opened perspective for further investigation of numerical methods for recovery of high order derivatives. Namely as the main criteria of method's efficiency have been taken its ability to achieve the optimal order of accuracy by using the minimal amount of discrete information. Note that particular these aspects of numerical differentiation remains still insufficiently studied. The present paper continues the research of [24], [23] and proposes a numerical method for recovering the high order mixed derivatives of smooth bivariate functions. The method is not only stable to small perturbations of the input data, but is also optimal in terms of accuracy and quantity of involved Fourier-Legendre coefficients, and also has a simple numerical implementation.

Let  $\{\varphi_k(t)\}_{k=0}^{\infty}$  be the system of Legendre polynomials orthonormal on  $[-1, 1]$  as

$$\varphi_k(t) = \sqrt{k+1/2} (2^k k!)^{-1} \frac{d^k}{dt^k} [(t^2-1)^k], \quad k = 0, 1, 2, \dots$$

By  $L_2 = L_2(Q)$  we mean space of square-summable on  $Q = [-1, 1]^2$  functions  $f(t, \tau)$  with inner product

$$\langle f, g \rangle = \int_{-1}^1 \int_{-1}^1 f(t, \tau) g(t, \tau) d\tau dt$$

and standard norm

$$\|f\|_{L_2}^2 = \sum_{k,j=0}^{\infty} |\langle f, \varphi_{k,j} \rangle|^2 < \infty,$$

---

<sup>†</sup> *Key words.* Numerical differentiation, Legendre polynomials, truncation method, minimal radius of Galerkin information.

where

$$\langle f, \varphi_{k,j} \rangle = \int_{-1}^1 \int_{-1}^1 f(t, \tau) \varphi_k(t) \varphi_j(\tau) dt d\tau, \quad k, j = 0, 1, 2, \dots,$$

are Fourier-Legendre coefficients of  $f$ . Let  $\ell_p$ ,  $1 \leq p \leq \infty$ , be the space of numerical sequences  $\bar{x} = \{x_{k,j}\}_{k,j \in \mathbb{N}_0}$ ,  $\mathbb{N}_0 = \{0\} \cup \mathbb{N}$ , such that the corresponding relation

$$\|\bar{x}\|_{\ell_p} := \begin{cases} \left( \sum_{k,j \in \mathbb{N}_0} |x_{k,j}|^p \right)^{\frac{1}{p}} < \infty, & 1 \leq p < \infty, \\ \sup_{k,j \in \mathbb{N}_0} |x_{k,j}| < \infty, & p = \infty, \end{cases}$$

is fulfilled.

We introduce the space of functions

$$L_{s,2}^\mu(Q) = \{f \in L_2(Q) : \|f\|_{s,\mu}^s := \sum_{k,j=0}^{\infty} (\underline{k} \cdot \underline{j})^{s\mu} |\langle f, \varphi_{k,j} \rangle|^s < \infty\},$$

where  $\mu > 0$ ,  $1 \leq s < \infty$ ,  $\underline{k} = \max\{1, k\}$ ,  $k = 0, 1, 2, \dots$ . Note that in the future we will use the same notations both for space and for a unite ball from this space:  $L_{s,2}^\mu = L_{s,2}^\mu(Q) = \{f \in L_{s,2}^\mu : \|f\|_{s,\mu} \leq 1\}$ , what we call a class of functions. What exactly is meant by  $L_{s,2}^\mu$ , space or class, will be clear depending on the context in each case. It should be noted that  $L_{s,2}^\mu$  is generalization of the class of bivariate functions with dominating mixed partial derivatives. Moreover, let  $C = C(Q)$  be the space of continuous on  $Q$  bivariate functions.

We represent a function  $f(t, \tau)$  from  $L_{s,2}^\mu$  as

$$f(t, \tau) = \sum_{k,j=0}^{\infty} \langle f, \varphi_{k,j} \rangle \varphi_k(t) \varphi_j(\tau),$$

and by its mixed derivative we mean the following series

$$f^{(r_1, r_2)}(t, \tau) = \sum_{k,j=r}^{\infty} \langle f, \varphi_{k,j} \rangle \varphi_k^{(r_1)}(t) \varphi_j^{(r_2)}(\tau), \quad r_1, r_2 = 1, 2, \dots \quad (1.1)$$

Assume that instead of the exact values of the Fourier-Legendre coefficients  $\langle f, \varphi_{k,j} \rangle$  only some their perturbations are known with the error level  $\delta$  in the metrics of  $\ell_p$ ,  $1 \leq p \leq \infty$ . More accurately, we assume that there is a sequence of numbers  $\bar{f}^\delta = \{\langle f^\delta, \varphi_{k,j} \rangle\}_{k,j \in \mathbb{N}_0}$  such that for  $\bar{\xi} = \{\xi_{k,j}\}_{k,j \in \mathbb{N}_0}$ , where  $\xi_{k,j} = \langle f - f^\delta, \varphi_{k,j} \rangle$ , and for some  $1 \leq p \leq \infty$  the relation

$$\|\bar{\xi}\|_{\ell_p} \leq \delta, \quad 0 < \delta < 1, \quad (1.2)$$

is true.

The research of this work is devoted to the optimization of methods for recovering the derivative (1.1) of functions from classes  $L_{s,2}^\mu$ . Further, we give a strict statement of the problem to be studied. In the coordinate space  $[r_1, \infty) \times [r_2, \infty)$  we take an arbitrary bounded domain  $\Omega$ . By  $\text{card}(\Omega)$  we mean the number of points that make up  $\Omega$  and by the information vector

$G(\Omega, \bar{f}^\delta) \in \mathbb{R}^N$ ,  $\text{card}(\Omega) = N$ , we take the set of perturbed values of Fourier-Legendre coefficients  $\{\langle f^\delta, \varphi_{k,j} \rangle\}_{(k,j) \in \Omega}$ .

Let  $X = L_2(Q)$  or  $X = C(Q)$ . By numerical differentiation algorithm we mean any mapping  $\psi^{(r_1, r_2)} = \psi^{(r_1, r_2)}(\Omega)$  that corresponds to the information vector  $G(\Omega, \bar{f}^\delta)$  an element

$\psi^{(r_1, r_2)}(G(\Omega, \bar{f}^\delta)) \in X$ , which is taken as an approximation to the derivative (1.1) of function  $f$  from the class  $L_{s,2}^\mu$ . We denote by  $\Psi(\Omega)$  the set of all algorithms  $\psi^{(r_1, r_2)}(\Omega) : \mathbb{R}^N \rightarrow X$ , that use the same information vector  $G(\Omega, \bar{f}^\delta)$ .

We do not require from algorithms which belong to  $\Psi(\Omega)$ , in generally speaking, either linearity or even stability. The only condition for algorithms from  $\Psi(\Omega)$  is to use input information in the form of perturbed values of the coefficients Fourier-Legendre with indices from the domain  $\Omega$  of the coordinate plane. Such the general understanding of the algorithm is explained by the desire to compare the widest possible range of possible methods of numerical differentiation.

The error of the algorithm  $\psi^{(r_1, r_2)}$  on the class  $L_{s,2}^\mu$  is determined by the quantity

$$\varepsilon_\delta(L_{s,2}^\mu, \psi^{(r_1, r_2)}(\Omega), X, \ell_p) = \sup_{f \in L_{s,2}^\mu, \|f\|_{s,\mu} \leq 1} \sup_{\bar{f}^\delta: (1.2)} \|f^{(r_1, r_2)} - \psi^{(r_1, r_2)}(G(\Omega, \bar{f}^\delta))\|_X. \quad (1.2)$$

The minimal radius of the Galerkin information for the problem of numerical differentiation on the class  $L_{s,2}^\mu$  is given by

$$R_{N,\delta}^{(r_1, r_2)}(L_{s,2}^\mu, X, \ell_p) = \inf_{\Omega: \text{card}(\Omega) \leq N} \inf_{\psi^{(r_1, r_2)} \in \Psi(\Omega)} \varepsilon_\delta(L_{s,2}^\mu, \psi^{(r_1, r_2)}(\Omega), X, \ell_p).$$

This value  $R_{N,\delta}^{(r_1, r_2)}(L_{s,2}^\mu, X, \ell_p)$  describes the minimal possible accuracy in the metric of  $X$ , which can be achieved by numerical differentiation of arbitrary function  $f \in L_{s,2}^\mu$ , while using not more than  $N$  values of its Fourier-Legendre coefficients that are  $\delta$ -perturbed in the  $\ell_p$  metric. Note that the minimal radius of Galerkin information in the problem of recovering the first partial derivative was studied in [27], and for other types of ill-posed problems, similar studies were previously carried out in [18], [13]. It should be added that the minimal radius characterizes the information complexity of the considered problem and is traditionally studied within the framework of the IBC-theory (Information- Based Complexity Theory), the foundations of which are laid in monographs [29] and [28].

The aim of our research is to find sharp estimates (in the power scale) for the quantities  $R_{N,\delta}^{(r_1, r_2)}(L_{s,2}^\mu, C, \ell_p)$  and  $R_{N,\delta}^{(r_1, r_2)}(L_{s,2}^\mu, L_2, \ell_p)$ .

## 2. Truncation method. Error estimate in $L_2$ - metric

It should be noted that at the moment a number of approaches were developed for numerical differentiation (see, for example, [3], [2], [20], [22], and also see [25] and its bibliography). All these methods are accepted to divide into three groups (see [22]): difference methods, interpolation methods and regularization methods. As is known, the first two types of methods have their advantage in the simplicity of implementation, but they guarantee satisfactory accuracy only in the case of exactly given input data about the differentiable function. In the same time regularization methods give stable approximations to the desired derivatives in the case of perturbed input data but most of them (for example, the Tikhonov method and its various variations) are quite complicated for numerical realization in view of their integral form and require hard-to-implement rules for determination of regularization parameters (see [22]). Recently in [24] a concise numerical method, called the truncation method, has been proposed as stable and simple approach to numerical differentiation of multivariable functions. The essence of this method is to replace the Fourier series (1.1) with a finite Fourier sum using perturbed data  $\langle f_\delta, \varphi_{k,j} \rangle$ . In the truncation method to ensure the stability of the approximation and achieve the required order accuracy, it is necessary to choose properly the discretization parameter, which here serves as a regularization parameter. So, the process of regularization in method under consideration consists in matching the discretization parameter with the perturbation level  $\delta$  of the input data. Simplicity of implementation is the main advantage of this method.

In the case of an arbitrary bounded domain  $\Omega$  of the coordinate plane  $[r_1, \infty) \times [r_2, \infty)$ , the truncation method for differentiating functions of two variables has the form

$$\mathcal{D}_{\Omega}^{(r_1, r_2)} f_{\delta}(t, \tau) = \sum_{(k, j) \in \Omega} \langle f_{\delta}, \varphi_{k, j} \rangle \varphi_k^{(r_1)}(t) \varphi_j^{(r_2)}(\tau).$$

In order to increase the efficiency of the approach under study, we take a hyperbolic cross as the domain  $\Omega$  of the following form

$$\Omega = \Gamma_{n, \gamma} := \{(k, j) : k \cdot j^{\gamma} \leq n, \quad k = r_1, \dots, n/r_2^{\gamma}, \quad j = r_2, \dots, (n/r_1)^{1/\gamma}\}, \quad \gamma \geq 1.$$

Then the proposed version of the truncation method can be written as

$$\mathcal{D}_{n, \gamma}^{(r_1, r_2)} f_{\delta}(t, \tau) = \sum_{k=r_1}^{n/r_2^{\gamma}} \sum_{j=r_2}^{(n/k)^{1/\gamma}} \langle f_{\delta}, \varphi_{k, j} \rangle \varphi_k^{(r_1)}(t) \varphi_j^{(r_2)}(\tau). \quad (2.1)$$

We note that earlier the idea of a hyperbolic cross in the problem of numerical differentiation was used in the papers [26], [27], [24] (for more details about usage of hyperbolic cross in solving the other types of ill-posed problems see [18], [13], [15]).

The approximation properties of the method (2.1) will be investigated in Sections 2 and 3 while in Section 4 it will be established that the method (2.1) is order-optimal in the sense of the minimal radius of the Galerkin information.

The parameters  $n$  and  $\gamma$  in (2.1) should be chosen depending on  $\delta$ ,  $p$ ,  $s$ ,  $r_1$ ,  $r_2$  and  $\mu$  so as to minimize the error of the method  $\mathcal{D}_{n, \gamma}^{(r_1, r_2)}$ :

$$\begin{aligned} & f^{(r_1, r_2)}(t, \tau) - \mathcal{D}_{n, \gamma}^{(r_1, r_2)} f_{\delta}(t, \tau) \\ &= \left( f^{(r_1, r_2)}(t, \tau) - \mathcal{D}_{n, \gamma}^{(r_1, r_2)} f(t, \tau) \right) + \left( \mathcal{D}_{n, \gamma}^{(r_1, r_2)} f(t, \tau) - \mathcal{D}_{n, \gamma}^{(r_1, r_2)} f_{\delta}(t, \tau) \right). \end{aligned} \quad (2.2)$$

For the first difference on the right-hand side of (2.2), the representation

$$f^{(r_1, r_2)}(t, \tau) - \mathcal{D}_{n, \gamma}^{(r_1, r_2)} f(t, \tau) = \Delta_1(t, \tau) + \Delta_2(t, \tau) \quad (2.3)$$

holds true, where

$$\Delta_1(t, \tau) = \sum_{k=n/r_2^{\gamma}+1}^{\infty} \sum_{j=r_2}^{\infty} \langle f, \varphi_{k, j} \rangle \varphi_k^{(r_1)}(t) \varphi_j^{(r_2)}(\tau), \quad (2.4)$$

$$\Delta_2(t, \tau) = \sum_{k=r_1}^{n/r_2^{\gamma}} \sum_{j=(n/k)^{1/\gamma}+1}^{\infty} \langle f, \varphi_{k, j} \rangle \varphi_k^{(r_1)}(t) \varphi_j^{(r_2)}(\tau). \quad (2.5)$$

For our calculations, we need the following formula (see Lemma 18 [16])

$$\varphi_k'(t) = 2 \sqrt{k+1/2} \sum_{l=0}^{k-1} \sqrt{l+1/2} \varphi_l(t), \quad k \in \mathbb{N}, \quad (2.6)$$

where in aggregate  $\sum_{l=0}^{k-1} \sqrt{l+1/2} \varphi_l(t)$  the summation is extended over only those terms for which  $k+l$  is odd.

Let us estimate the error of the method (2.1) in the metric of  $L_2$ . Upper bounds for difference (2.3) are contained in the following statement.

**Lemma 2.1** Let  $f \in L_{s,2}^\mu$ ,  $1 \leq s < \infty$ ,  $r_1 \geq r_2$ ,  $\mu > 2r_1 - 1/s + 1/2$ .

I) If  $r_1 = r_2$ , then for  $\gamma = 1$  it holds true

$$\|f^{(r_1, r_1)} - \mathcal{D}_{n,1}^{(r_1, r_1)} f\|_{L_2} \leq c \|f\|_{s,\mu} n^{-\mu+2r_1-1/s+1/2} \ln^{3/2-1/s} n.$$

II) If  $r_1 > r_2$ , then

a) for any  $\gamma \in \left[1, \frac{\mu-2r_2+1/s-1/2}{\mu-2r_1+1/s+1/2}\right) \cup \left(\frac{\mu-2r_2+1/s-1/2}{\mu-2r_1+1/s+1/2}, \frac{\mu-2r_2+1/s+1/2}{\mu-2r_1+1/s+1/2}\right) \cup \left(\frac{\mu-2r_2+1/s+1/2}{\mu-2r_1+1/s+1/2}, \frac{\mu-2r_2+1/s-1/2}{\mu-2r_1+1/s-1/2}\right)$  it holds true

$$\|f^{(r_1, r_2)} - \mathcal{D}_{n,\gamma}^{(r_1, r_2)} f\|_{L_2} \leq c \|f\|_{s,\mu} n^{-\mu+2r_1-1/s+1/2},$$

b) for  $\gamma = \frac{\mu-2r_2+1/s-1/2}{\mu-2r_1+1/s+1/2}$  and  $\gamma = \frac{\mu-2r_2+1/s-1/2}{\mu-2r_1+1/s-1/2}$  it holds true

$$\|f^{(r_1, r_2)} - \mathcal{D}_{n,\gamma}^{(r_1, r_2)} f\|_{L_2} \leq c \|f\|_{s,\mu} n^{-\mu+2r_1-1/s+1/2} \ln^{1/2} n,$$

c) for  $\gamma = \frac{\mu-2r_2+1/s+1/2}{\mu-2r_1+1/s+1/2}$  it holds true

$$\|f^{(r_1, r_2)} - \mathcal{D}_{n,\gamma}^{(r_1, r_2)} f\|_{L_2} \leq c \|f\|_{s,\mu} n^{-\mu+2r_1-1/s+1/2} \ln^{1-1/s} n.$$

*Proof.* Using the formula (2.6), from (2.4) we have

$$\begin{aligned} \Delta_1(t, \tau) &= 2^{r_1+r_2} \sum_{k=n/r_2^\gamma+1}^{\infty} \sum_{j=r_2}^{\infty} \sqrt{k+1/2} \sqrt{j+1/2} \langle f, \varphi_{kj} \rangle \\ &\times \sum_{l_1=r_1-1}^{k-1} (l_1+1/2) \sum_{l_2=r_1-2}^{l_1-1} (l_2+1/2) \dots \sum_{l_{r_1-1}=1}^{l_{r_1-2}-1} (l_{r_1-1}+1/2) \sum_{l_{r_1}=0}^{l_{r_1-1}-1} \sqrt{l_{r_1}+1/2} \varphi_{l_{r_1}}(t) \\ &\times \sum_{m_1=r_2-1}^{j-1} (m_1+1/2) \sum_{m_2=r_2-2}^{m_1-1} (m_2+1/2) \dots \sum_{m_{r_2-1}=1}^{m_{r_2-2}-1} (m_{r_2-1}+1/2) \sum_{m_{r_2}=0}^{m_{r_2-1}-1} \sqrt{m_{r_2}+1/2} \varphi_{m_{r_2}}(\tau). \end{aligned}$$

Further, we change the order of summation and get

$$\Delta_1(t, \tau) = \Delta_{11}(t, \tau) + \Delta_{12}(t, \tau),$$

where

$$\begin{aligned} \Delta_{11}(t, \tau) &= 2^{r_1+r_2} \sum_{l_{r_1}=0}^{n/r_2^\gamma-r_1+1} \sqrt{l_{r_1}+1/2} \varphi_{l_{r_1}}(t) \sum_{m_{r_2}=0}^{\infty} \sqrt{m_{r_2}+1/2} \varphi_{m_{r_2}}(\tau) \\ &\times \sum_{k=n/r_2^\gamma+1}^{\infty} \sum_{j=m_{r_2}+r_2}^{\infty} \sqrt{k+1/2} \sqrt{j+1/2} \langle f, \varphi_{kj} \rangle B_{k,j}^{r_1, r_2}, \end{aligned} \quad (2.7)$$

$$\begin{aligned} \Delta_{12}(t, \tau) &= 2^{r_1+r_2} \sum_{l_{r_1}=n/r_2^\gamma-r_1+2}^{\infty} \sqrt{l_{r_1}+1/2} \varphi_{l_{r_1}}(t) \sum_{m_{r_2}=0}^{\infty} \sqrt{m_{r_2}+1/2} \varphi_{m_{r_2}}(\tau) \\ &\times \sum_{k=l_{r_1}+r_1}^{\infty} \sum_{j=m_{r_2}+r_2}^{\infty} \sqrt{k+1/2} \sqrt{j+1/2} \langle f, \varphi_{kj} \rangle B_{k,j}^{r_1, r_2} \end{aligned} \quad (2.8)$$

and

$$B_{k,j}^{r_1, r_2} := \sum_{l_1=l_{r_1}+r_1-1}^{k-1} (l_1+1/2) \sum_{l_2=l_{r_1}+r_1-2}^{l_1-1} (l_2+1/2) \dots \sum_{l_{r_1-1}=l_{r_1}+1}^{l_{r_1-2}-1} (l_{r_1-1}+1/2)$$

$$\begin{aligned}
& \times \sum_{m_1=m_{r_2}+r_2-1}^{j-1*} (m_1+1/2) \sum_{m_2=m_{r_2}+r_2-2}^{m_1-1*} (m_2+1/2) \dots \sum_{m_{r_2-1}=m_{r_2}+1}^{m_{r_2}-2-1*} (m_{r_2-1}+1/2) \\
& \leq ck^{2(r_1-1)}j^{2(r_2-1)}. \tag{2.9}
\end{aligned}$$

Let's  $1 < s < \infty$ . At first we bound  $\Delta_1(t, \tau)$ . Using Hölder inequality and (2.9), for  $\mu > 2r_1 + \frac{s-1}{s} - 1/2$  we have

$$\begin{aligned}
& \|\Delta_{11}\|_{L_2}^2 \leq 4^{r_1+r_2} \sum_{l_{r_1}=0}^{n/r_2^\gamma-r_1+1} (l_{r_1}+1/2) \sum_{m_{r_2}=0}^{\infty} (m_{r_2}+1/2) \\
& \times \left( \sum_{k=n/r_2^\gamma+1}^{\infty} \sum_{j=m_{r_2}+r_2}^{\infty} (kj)^\mu |\langle f, \varphi_{kj} \rangle| \frac{B_{k,j}^{r_1,r_2}}{(kj)^{\mu-1/2}} \right)^2 \leq c\|f\|_{s,\mu}^2 \sum_{l_{r_1}=0}^{n/r_2^\gamma-r_1+1} (l_{r_1}+1/2) \sum_{m_{r_2}=0}^{\infty} (m_{r_2}+1/2) \\
& \times \left( \sum_{k=n/r_2^\gamma+1}^{\infty} k^{-(\mu-2r_1+3/2)s/(s-1)} \sum_{j=m_{r_2}+r_2}^{\infty} j^{-(\mu-2r_2+3/2)s/(s-1)} \right)^{2(s-1)/s} \\
& \leq c\|f\|_{s,\mu}^2 n^{-2(\mu-2r_1+3/2)+\frac{2(s-1)}{s}} \sum_{l_{r_1}=0}^{n/r_2^\gamma-r_1+1} (l_{r_1}+1/2) \sum_{m_{r_2}=0}^{\infty} (m_{r_2}+1/2)^{-2(\mu-2r_2+3/2)+\frac{2(s-1)}{s}+1} \\
& \leq c\|f\|_{s,\mu}^2 n^{-2(\mu-2r_1+1/s-1/2)}, \\
& \|\Delta_{12}\|_{L_2}^2 \leq c\|f\|_{s,\mu}^2 \sum_{l_{r_1}=n/r_2^\gamma-r_1+2}^{\infty} (l_{r_1}+1/2) \sum_{m_{r_2}=0}^{\infty} (m_{r_2}+1/2) \\
& \times \left( \sum_{k=l_{r_1}+r_1}^{\infty} k^{-(\mu-2r_1+3/2)s/(s-1)} \sum_{j=m_{r_2}+r_2}^{\infty} j^{-(\mu-2r_2+3/2)s/(s-1)} \right)^{2(s-1)/s} \\
& \leq c\|f\|_{s,\mu}^2 \sum_{l_{r_1}=n/r_2^\gamma-r_1+2}^{\infty} (l_{r_1}+1/2)^{-2(\mu-2r_1+3/2)+\frac{2(s-1)}{s}+1} \sum_{m_{r_2}=0}^{\infty} (m_{r_2}+1/2)^{-2(\mu-2r_2+3/2)+\frac{2(s-1)}{s}+1} \\
& \leq c\|f\|_{s,\mu}^2 n^{-2(\mu-2r_1+1/s-1/2)}.
\end{aligned}$$

Combining the estimates for  $\Delta_{11}(t, \tau)$  and  $\Delta_{12}(t, \tau)$  we obtain

$$\|\Delta_1\|_{L_2} \leq c\|f\|_{s,\mu} n^{-\mu+2r_1-1/s+1/2}.$$

Next, using the formula (2.6), from (2.5) we have

$$\begin{aligned}
\Delta_2(t, \tau) &= 2^{r_1+r_2} \sum_{k=r_1}^{n/r_2^\gamma} \sum_{j=(n/k)^{1/\gamma}+1}^{\infty} \sqrt{k+1/2} \sqrt{j+1/2} \langle f, \varphi_{kj} \rangle \\
& \times \sum_{l_1=r_1-1}^{k-1*} (l_1+1/2) \sum_{l_2=r_1-2}^{l_1-1*} (l_2+1/2) \dots \sum_{l_{r_1-1}=1}^{l_{r_1-2}-1*} (l_{r_1-1}+1/2) \sum_{l_{r_1}=0}^{l_{r_1-1}-1*} \sqrt{l_{r_1}+1/2} \varphi_{l_{r_1}}(t) \\
& \times \sum_{m_1=r_2-1}^{j-1*} (m_1+1/2) \sum_{m_2=r_2-2}^{m_1-1*} (m_2+1/2) \dots \sum_{m_{r_2-1}=1}^{m_{r_2-2}-1*} (m_{r_2-1}+1/2) \sum_{m_{r_2}=0}^{m_{r_2-1}-1*} \sqrt{m_{r_2}+1/2} \varphi_{m_{r_2}}(\tau).
\end{aligned}$$

Further, we change the order of summation and get

$$\Delta_2(t, \tau) = \Delta_{21}(t, \tau) + \Delta_{22}(t, \tau) + \Delta_{23}(t, \tau),$$

where

$$\begin{aligned} \Delta_{21}(t, \tau) &= 2^{r_1+r_2} \sum_{l_{r_1}=0}^{n/r_2^\gamma-r_1} \sqrt{l_{r_1}+1/2} \varphi_{l_{r_1}}(t) \sum_{m_{r_2}=0}^{(\frac{n}{l_{r_1}+r_1})^{1/\gamma}-r_2+1} \sqrt{m_{r_2}+1/2} \varphi_{m_{r_2}}(\tau) \\ &\quad \times \sum_{k=l_{r_1}+r_1}^{\frac{n}{(m_{r_2}+r_2-1)^\gamma}} \sum_{j=(n/k)^{1/\gamma}+1}^{\infty} \sqrt{k+1/2} \sqrt{j+1/2} \langle f, \varphi_{kj} \rangle B_{k,j}^{r_1, r_2}, \\ \Delta_{22}(t, \tau) &= 2^{r_1+r_2} \sum_{l_{r_1}=0}^{n/r_2^\gamma-r_1} \sqrt{l_{r_1}+1/2} \varphi_{l_{r_1}}(t) \sum_{m_{r_2}=2}^{(\frac{n}{l_{r_1}+r_1})^{1/\gamma}-r_2+2} \sqrt{m_{r_2}+1/2} \varphi_{m_{r_2}}(\tau) \\ &\quad \times \sum_{k=\frac{n}{(m_{r_2}+r_2-2)^\gamma}}^{n/r_2^\gamma} \sum_{j=m_{r_2}+r_2}^{\infty} \sqrt{k+1/2} \sqrt{j+1/2} \langle f, \varphi_{kj} \rangle B_{k,j}^{r_1, r_2}, \\ \Delta_{23}(t, \tau) &= 2^{r_1+r_2} \sum_{l_{r_1}=0}^{n/r_2^\gamma-r_1} \sqrt{l_{r_1}+1/2} \varphi_{l_{r_1}}(t) \sum_{m_{r_2}=(\frac{n}{l_{r_1}+r_1})^{1/\gamma}-r_2+3}^{\infty} \sqrt{m_{r_2}+1/2} \varphi_{m_{r_2}}(\tau) \\ &\quad \times \sum_{k=l_{r_1}+r_1}^{n/r_2^\gamma} \sum_{j=m_{r_2}+r_2}^{\infty} \sqrt{k+1/2} \sqrt{j+1/2} \langle f, \varphi_{kj} \rangle B_{k,j}^{r_1, r_2}. \end{aligned}$$

Let as before  $1 < s < \infty$ . Now we estimate the norm of  $\Delta_2$  in the case of  $r_1 > r_2$  and  $\gamma = 1$ :

$$\begin{aligned} \|\Delta_{21}\|_{L_2}^2 &\leq c \|f\|_{s, \mu}^2 n^{-2(\mu-2r_1+3/2)+\frac{2(s-1)}{s}} \sum_{l_{r_1}=0}^{n/r_2-r_1} (l_{r_1}+1/2) \sum_{m_{r_2}=0}^{\frac{n}{l_{r_1}+r_1}-r_2+1} (m_{r_2}+1/2) \\ &\quad \times \left( \sum_{k=l_{r_1}+r_1}^{\frac{n}{m_{r_2}+r_2-1}} k^{2(r_1-r_2)s/(s-1)-1} \right)^{2(s-1)/s} \leq c \|f\|_{s, \mu}^2 n^{-2(\mu-2r_1+3/2)+\frac{2(s-1)}{s}} \\ &\quad \times \sum_{l_{r_1}=0}^{n/r_2-r_1} (l_{r_1}+1/2) \sum_{m_{r_2}=0}^{\frac{n}{l_{r_1}+r_1}-r_2+1} (m_{r_2}+1/2)^{1-4(r_1-r_2)} \\ &\leq c \|f\|_{s, \mu}^2 n^{-2(\mu-2r_1+3/2)+\frac{2(s-1)}{s}} \sum_{l_{r_1}=0}^{n/r_2-r_1} (l_{r_1}+1/2) = c \|f\|_{s, \mu}^2 n^{-2(\mu-2r_1+1/s-1/2)}, \\ \|\Delta_{22}\|_{L_2}^2 &\leq c \|f\|_{s, \mu}^2 \sum_{l_{r_1}=0}^{n/r_2-r_1} (l_{r_1}+1/2) \sum_{m_{r_2}=2}^{\frac{n}{l_{r_1}+r_1}-r_2+2} (m_{r_2}+1/2) \\ &\quad \times \left( \sum_{k=\frac{n}{m_{r_2}+r_2-2}}^{n/r_2} k^{-(\mu-2r_1+3/2)s/(s-1)} \sum_{j=m_{r_2}+r_2}^{\infty} j^{-(\mu-2r_2+3/2)s/(s-1)} \right)^{2(s-1)/s} \end{aligned}$$

$$\begin{aligned}
&\leq c \|f\|_{s,\mu}^2 n^{-2(\mu-2r_1+3/2)+\frac{2(s-1)}{s}} \sum_{l_{r_1}=0}^{n/r_2-r_1} (l_{r_1}+1/2) \sum_{m_{r_2}=2}^{\frac{n}{l_{r_1+r_1}}-r_2+2} (m_{r_2}+1/2)^{1-4(r_1-r_2)} \\
&\leq c \|f\|_{s,\mu}^2 n^{-2(\mu-2r_1+1/s-1/2)}, \\
&\|\Delta_{23}\|_{L_2}^2 \leq c \|f\|_{s,\mu}^2 \sum_{l_{r_1}=0}^{n/r_2-r_1} (l_{r_1}+1/2) \sum_{m_{r_2}=\frac{n}{l_{r_1+r_1}}-r_2+3}^{\infty} (m_{r_2}+1/2) \\
&\quad \times \left( \sum_{k=l_{r_1}+r_1}^{n/r_2} k^{-(\mu-2r_1+3/2)s/(s-1)} \sum_{j=m_{r_2}+r_2}^{\infty} j^{-(\mu-2r_2+3/2)s/(s-1)} \right)^{2(s-1)/s} \\
&\leq c \|f\|_{s,\mu}^2 \sum_{l_{r_1}=0}^{n/r_2-r_1} (l_{r_1}+1/2)^{1-2(\mu-2r_1+3/2)+\frac{2(s-1)}{s}} \sum_{m_{r_2}=\frac{n}{l_{r_1+r_1}}-r_2+3}^{\infty} (m_{r_2}+1/2)^{1-2(\mu-2r_2+3/2)+\frac{2(s-1)}{s}} \\
&\leq c \|f\|_{s,\mu}^2 n^{-2(\mu-2r_2+3/2)+\frac{2(s-1)}{s}+2} \sum_{l_{r_1}=0}^{n/r_2-r_1} (l_{r_1}+1/2)^{4(r_1-r_2)-1} \\
&\leq c \|f\|_{s,\mu}^2 n^{-2(\mu-2r_1+1/s-1/2)}.
\end{aligned}$$

Combining the estimates for  $\Delta_{21}(t, \tau)$ ,  $\Delta_{22}(t, \tau)$  and  $\Delta_{23}(t, \tau)$  we obtain

$$\|\Delta_2\|_{L_2} \leq c \|f\|_{s,\mu} n^{-\mu+2r_1-1/s+1/2}.$$

To estimate  $\|\Delta_2\|_{L_2}$  for  $r_1 = r_2$  or  $\gamma > 1$ , it is enough to repeat an analogous line of reasoning. The combination of (2.3) and bounds for the norms of  $\Delta_1$ ,  $\Delta_2$  makes it possible to establish the desired inequality in the case  $1 < s < \infty$ .

For  $s = 1$  the assertion of Lemma is proved similarly.  $\square$

The following statement contains estimates for the second difference from the right-hand side of (2.2) in the metric of  $L_2$ .

**Lemma 2.2** *Let  $f \in L_2(Q)$  and let the condition (1.2) be satisfied.*

*I) If  $r_1 = r_2$ , then*

*a) for  $\gamma = 1$  it holds true*

$$\|\mathcal{D}_{n,1}^{(r_1,r_1)} f - \mathcal{D}_{n,1}^{(r_1,r_1)} f_\delta\|_{L_2} \leq c\delta n^{2r_1-1/p+1/2} \ln^{3/2-1/p} n,$$

*b) for any  $1 < \gamma < \frac{2r_1-1/2+(p-1)/p}{2r_1-3/2+(p-1)/p}$  it holds true*

$$\|\mathcal{D}_{n,\gamma}^{(r_1,r_1)} f - \mathcal{D}_{n,\gamma}^{(r_1,r_1)} f_\delta\|_{L_2} \leq c\delta n^{2r_1-1/p+1/2},$$

*c) for  $\gamma = \frac{2r_1-1/2+(p-1)/p}{2r_1-3/2+(p-1)/p}$  it holds true*

$$\|\mathcal{D}_{n,\gamma}^{(r_1,r_1)} f - \mathcal{D}_{n,\gamma}^{(r_1,r_1)} f_\delta\|_{L_2} \leq c\delta n^{2r_1-1/p+1/2} \ln^{1/2} n.$$

*II) If  $r_1 > r_2$ , then for any  $\gamma \geq 1$  it holds true*

$$\|\mathcal{D}_{n,\gamma}^{(r_1,r_2)} f - \mathcal{D}_{n,\gamma}^{(r_1,r_2)} f_\delta\|_{L_2} \leq c\delta n^{2r_1-1/p+1/2}.$$



*Proof.* Let's write down the representation

$$\mathcal{D}_{n,\gamma}^{(r_1,r_2)} f(t,\tau) - \mathcal{D}_{n,\gamma}^{(r_1,r_2)} f_\delta(t,\tau) = \sum_{k=r_1}^{n/r_2^\gamma} \sum_{j=r_2}^{(n/k)^{1/\gamma}} \langle f - f_\delta, \varphi_{k,j} \rangle \varphi_k^{(r_1)}(t) \varphi_j^{(r_2)}(\tau).$$

Using the formula (2.6), we get

$$\begin{aligned} \mathcal{D}_{n,\gamma}^{(r_1,r_2)} f(t,\tau) - \mathcal{D}_{n,\gamma}^{(r_1,r_2)} f_\delta(t,\tau) &= 2^{r_1+r_2} \sum_{k=r_1}^{n/r_2^\gamma} \sum_{j=r_2}^{(n/k)^{1/\gamma}} \sqrt{k+1/2} \sqrt{j+1/2} \langle f - f_\delta, \varphi_{k,j} \rangle \\ &\times \sum_{l_1=r_1-1}^{k-1}^* (l_1+1/2) \sum_{l_2=r_1-2}^{l_1-1}^* (l_2+1/2) \dots \sum_{l_{r_1-1}=1}^{l_{r_1-2}-1}^* (l_{r_1-1}+1/2) \sum_{l_{r_1}=0}^{l_{r_1-1}-1}^* \sqrt{l_{r_1}+1/2} \varphi_{l_{r_1}}(t) \\ &\times \sum_{m_1=r_2-1}^{j-1}^* (m_1+1/2) \sum_{m_2=r_2-2}^{m_1-1}^* (m_2+1/2) \dots \sum_{m_{r_2-1}=1}^{m_{r_2-2}-1}^* (m_{r_2-1}+1/2) \sum_{m_{r_2}=0}^{m_{r_2-1}-1}^* \sqrt{m_{r_2}+1/2} \varphi_{m_{r_2}}(\tau). \end{aligned}$$

Further, we change the order of summation and get

$$\begin{aligned} \mathcal{D}_{n,\gamma}^{(r_1,r_2)} f(t,\tau) - \mathcal{D}_{n,\gamma}^{(r_1,r_2)} f_\delta(t,\tau) &= 2^{r_1+r_2} \sum_{l_{r_1}=0}^{n/r_2^\gamma-r_1} \sqrt{l_{r_1}+1/2} \varphi_{l_{r_1}}(t) \sum_{m_{r_2}=0}^{(\frac{n}{l_{r_1}+r_1})^{1/\gamma}-r_2} \sqrt{m_{r_2}+1/2} \varphi_{m_{r_2}}(\tau) \\ &\times \sum_{k=l_{r_1}+r_1}^{\frac{n}{(m_{r_2}+r_2)^\gamma}} \sum_{j=m_{r_2}+r_2}^{(\frac{n}{k})^{1/\gamma}} \sqrt{k+1/2} \sqrt{j+1/2} \langle f - f_\delta, \varphi_{k,j} \rangle B_{k,j}^{r_1,r_2}. \end{aligned}$$

Let  $1 < p < \infty$  first. Then, using the Hölder inequality and the estimate (2.9), we find

$$\begin{aligned} \|\mathcal{D}_{n,\gamma}^{(r_1,r_2)} f - \mathcal{D}_{n,\gamma}^{(r_1,r_2)} f_\delta\|_{L_2}^2 &\leq 4^{r_1+r_2} \sum_{l_{r_1}=0}^{n/r_2^\gamma-r_1} (l_{r_1}+1/2) \sum_{m_{r_2}=0}^{(\frac{n}{l_{r_1}+r_1})^{1/\gamma}-r_2} (m_{r_2}+1/2) \\ &\times \left( \sum_{k=l_{r_1}+r_1}^{\frac{n}{(m_{r_2}+r_2)^\gamma}} \sum_{j=m_{r_2}+r_2}^{(\frac{n}{k})^{1/\gamma}} \sqrt{k+1/2} \sqrt{j+1/2} |\langle f - f_\delta, \varphi_{k,j} \rangle| B_{k,j}^{r_1,r_2} \right)^2 \\ &\leq c\delta^2 \sum_{l_{r_1}=0}^{n/r_2^\gamma-r_1} (l_{r_1}+1/2) \sum_{m_{r_2}=0}^{(\frac{n}{l_{r_1}+r_1})^{1/\gamma}-r_2} (m_{r_2}+1/2) \\ &\times \left( \sum_{k=l_{r_1}+r_1}^{\frac{n}{(m_{r_2}+r_2)^\gamma}} k^{(2r_1-3/2)p/(p-1)} \sum_{j=m_{r_2}+r_2}^{(\frac{n}{k})^{1/\gamma}} j^{(2r_2-3/2)p/(p-1)} \right)^{2(p-1)/p} \\ &\leq c\delta^2 \sum_{l_{r_1}=0}^{n/r_2^\gamma-r_1} (l_{r_1}+1/2) \sum_{m_{r_2}=0}^{(\frac{n}{l_{r_1}+r_1})^{1/\gamma}-r_2} (m_{r_2}+1/2) \\ &\times \left( n^{\frac{(2r_2-3/2)p}{\gamma(p-1)}+1/\gamma} \sum_{k=l_{r_1}+r_1}^{\frac{n}{(m_{r_2}+r_2)^\gamma}} k^{\frac{(2r_1-3/2)p}{p-1}-\frac{(2r_2-3/2)p}{\gamma(p-1)}-1/\gamma} \right)^{2(p-1)/p}. \end{aligned} \quad (2.10)$$

Whence it follows for  $r_1 > r_2$  and any  $\gamma \geq 1$  that

$$\|\mathcal{D}_{n,\gamma}^{(r_1,r_2)} f - \mathcal{D}_{n,\gamma}^{(r_1,r_2)} f_\delta\|_{L_2}^2 \leq c\delta^2 n^{4r_1-3+\frac{2(p-1)}{p}} \sum_{l_{r_1}=0}^{n/r_2^\gamma-r_1} (l_{r_1} + 1/2) = c\delta^2 n^{4r_1-2/p+1}.$$

which was required to prove.

Let's now  $r_1 = r_2$  and any  $\gamma = 1$ . Then from (2.10) we obtain

$$\begin{aligned} \|\mathcal{D}_{n,1}^{(r_1,r_1)} f - \mathcal{D}_{n,1}^{(r_1,r_1)} f_\delta\|_{L_2}^2 &\leq c\delta^2 n^{4r_1-3+\frac{2(p-1)}{p}} \ln^{\frac{2(p-1)}{p}} n \sum_{l_{r_1}=0}^{n/r_1-r_1} (l_{r_1} + 1/2) \sum_{m_{r_1}=0}^{\frac{n}{l_{r_1}+r_1}-r_1} (m_{r_1} + 1/2) \\ &\leq c\delta^2 n^{4r_1-1+\frac{2(p-1)}{p}} \ln^{\frac{2(p-1)}{p}} n \sum_{l_{r_1}=0}^{n/r_2-r_1} \frac{1}{l_{r_1} + 1/2} = c\delta^2 n^{4r_1-2/p+1} \ln^{3-2/p} n. \end{aligned}$$

To estimate  $\|\mathcal{D}_{n,\gamma}^{(r_1,r_2)} f - \mathcal{D}_{n,\gamma}^{(r_1,r_2)} f_\delta\|_{L_2}$  for  $\gamma > 1$  and also for  $p = 1, p = \infty$ , it is enough to repeat an analogous line of reasoning.  $\square$

The combination of Lemmas 2.1 and 2.2 gives

**Theorem 2.3** *Let  $f \in L_{s,2}^\mu$ ,  $1 \leq s < \infty$ ,  $r_1 \geq r_2$ ,  $\mu > 2r_1 + 1/2 - 1/s$ , and let the condition (1.2) be satisfied.*

I) *If  $r_1 = r_2$ , then for  $n \asymp \left(\delta / \ln^{1/p-1/s} \frac{1}{\delta}\right)^{-\frac{1}{\mu-1/p+1/s}}$  and  $\gamma = 1$  it holds true*

$$\|f^{(r_1,r_1)} - \mathcal{D}_{n,1}^{(r_1,r_1)} f_\delta\|_{L_2} \leq c \left(\delta \ln^{1/s-1/p} \frac{1}{\delta}\right)^{\frac{\mu-2r_1+1/s-1/2}{\mu-1/p+1/s}} \ln^{3/2-1/s} \frac{1}{\delta}.$$

II) *If  $r_1 > r_2$ , then*

a) *for  $n \asymp \delta^{-\frac{1}{\mu-1/p+1/s}}$  and any  $\gamma \in \left[1, \frac{\mu-2r_2+1/s-1/2}{\mu-2r_1+1/s+1/2}\right) \cup \left(\frac{\mu-2r_2+1/s-1/2}{\mu-2r_1+1/s+1/2}, \frac{\mu-2r_2+1/s+1/2}{\mu-2r_1+1/s+1/2}\right)$*   
 $\cup \left(\frac{\mu-2r_2+1/s+1/2}{\mu-2r_1+1/s+1/2}, \frac{\mu-2r_2+1/s-1/2}{\mu-2r_1+1/s-1/2}\right)$  *it holds true*

$$\|f^{(r_1,r_2)} - \mathcal{D}_{n,\gamma}^{(r_1,r_2)} f_\delta\|_{L_2} \leq c\delta^{\frac{\mu-2r_1+1/s-1/2}{\mu-1/p+1/s}},$$

b) *for  $n \asymp \delta^{-\frac{1}{\mu-1/p+1/s}} \ln^{\frac{1}{2(\mu-1/p+1/s)}} \frac{1}{\delta}$  and  $\gamma = \frac{\mu-2r_2+1/s-1/2}{\mu-2r_1+1/s+1/2}$ ,  $\gamma = \frac{\mu-2r_2+1/s-1/2}{\mu-2r_1+1/s-1/2}$*   
*it holds true*

$$\|f^{(r_1,r_2)} - \mathcal{D}_{n,\gamma}^{(r_1,r_2)} f_\delta\|_{L_2} \leq c \left(\delta / \ln^{1/2} \frac{1}{\delta}\right)^{\frac{\mu-2r_1+1/s-1/2}{\mu-1/p+1/s}} \ln^{1/2} \frac{1}{\delta},$$

c) *for  $n \asymp \left(\delta / \ln^{1-1/s} \frac{1}{\delta}\right)^{-\frac{1}{\mu-1/p+1/s}}$  and  $\gamma = \frac{\mu-2r_2+1/s+1/2}{\mu-2r_1+1/s+1/2}$  it holds true*

$$\|f^{(r_1,r_2)} - \mathcal{D}_{n,\gamma}^{(r_1,r_2)} f_\delta\|_{L_2} \leq c \left(\delta / \ln^{1-1/s} \frac{1}{\delta}\right)^{\frac{\mu-2r_1+1/s-1/2}{\mu-1/p+1/s}} \ln^{1-1/s} \frac{1}{\delta}.$$

### 3. Truncation method. Error estimate in the metric of $C$

Now we have to bound the error of (2.1) in the metric of  $C$ . Upper estimates for the norm of difference (2.3) are contained in the following statement.

**Lemma 3.1** *Let  $f \in L_{s,2}^\mu$ ,  $1 \leq s < \infty$ ,  $r_1 \geq r_2$ ,  $\mu > 2r_1 + 3/2 - 1/s$ .*

*I) If  $r_1 = r_2$ , then for  $\gamma = 1$  it holds true*

$$\|f^{(r_1, r_1)} - \mathcal{D}_{n,1}^{(r_1, r_1)} f\|_C \leq c \|f\|_{s,\mu} n^{-\mu+2r_1-1/s+3/2} \ln^{2-1/s} n.$$

*II) If  $r_1 = r_2 + 1$ , then*

a) for any  $\gamma \in \left(1, \frac{\mu-2r_1+1/s+5/2}{\mu-2r_1+1/s+1/2}\right) \cup \left(\frac{\mu-2r_1+1/s+5/2}{\mu-2r_1+1/s+1/2}, \frac{\mu-2r_1+1/s+1/2}{\mu-2r_1+1/s-3/2}\right)$

*it holds true*

$$\|f^{(r_1, r_1-1)} - \mathcal{D}_{n,\gamma}^{(r_1, r_1-1)} f\|_C \leq c \|f\|_{s,\mu} n^{-\mu+2r_1-1/s+3/2},$$

b) for  $\gamma = 1$  and  $\gamma = \frac{\mu-2r_1+1/s+1/2}{\mu-2r_1+1/s-3/2}$  *it holds true*

$$\|f^{(r_1, r_1-1)} - \mathcal{D}_{n,\gamma}^{(r_1, r_1-1)} f\|_C \leq c \|f\|_{s,\mu} n^{-\mu+2r_1-1/s+3/2} \ln n,$$

c) for  $\gamma = \frac{\mu-2r_1+1/s+5/2}{\mu-2r_1+1/s+1/2}$  *it holds true*

$$\|f^{(r_1, r_1-1)} - \mathcal{D}_{n,\gamma}^{(r_1, r_1-1)} f\|_C \leq c \|f\|_{s,\mu} n^{-\mu+2r_1-1/s+3/2} \ln^{1-1/s} n.$$

*III) If  $r_1 > r_2 + 1$ , then*

a) for any  $\gamma \in \left[1, \frac{\mu-2r_2+1/s-3/2}{\mu-2r_1+1/s+1/2}\right) \cup \left(\frac{\mu-2r_2+1/s-3/2}{\mu-2r_1+1/s+1/2}, \frac{\mu-2r_2+1/s+1/2}{\mu-2r_1+1/s+1/2}\right)$

$\cup \left(\frac{\mu-2r_2+1/s+1/2}{\mu-2r_1+1/s+1/2}, \frac{\mu-2r_2+1/s-3/2}{\mu-2r_1+1/s-3/2}\right)$  *it holds true*

$$\|f^{(r_1, r_2)} - \mathcal{D}_{n,\gamma}^{(r_1, r_2)} f\|_C \leq c \|f\|_{s,\mu} n^{-\mu+2r_1-1/s+3/2},$$

b) for  $\gamma = \frac{\mu-2r_2+1/s-3/2}{\mu-2r_1+1/s+1/2}$  and  $\gamma = \frac{\mu-2r_2+1/s-3/2}{\mu-2r_1+1/s-3/2}$  *it holds true*

$$\|f^{(r_1, r_2)} - \mathcal{D}_{n,\gamma}^{(r_1, r_2)} f\|_C \leq c \|f\|_{s,\mu} n^{-\mu+2r_1-1/s+3/2} \ln n,$$

c) for  $\gamma = \frac{\mu-2r_2+1/s+1/2}{\mu-2r_1+1/s+1/2}$  *it holds true*

$$\|f^{(r_1, r_2)} - \mathcal{D}_{n,\gamma}^{(r_1, r_2)} f\|_C \leq c \|f\|_{s,\mu} n^{-\mu+2r_1-1/s+3/2} \ln^{1-1/s} n.$$

*Proof.* Let's start with the case  $1 < s < \infty$ . Using (2.9), from (2.7) and (2.8) we get

$$\begin{aligned} \|\Delta_{11}\|_C &\leq c \sum_{l_{r_1}=0}^{n/r_2^\gamma - r_1 + 1} (l_{r_1} + 1/2) \sum_{m_{r_2}=0}^{\infty} (m_{r_2} + 1/2) \\ &\times \sum_{k=n/r_2^\gamma + 1}^{\infty} \sum_{j=m_{r_2} + r_2}^{\infty} (kj)^\mu |\langle f, \varphi_{kj} \rangle| \frac{B_{k,j}^{r_1, r_2}}{(kj)^{\mu-1/2}} \leq c \|f\|_{s,\mu} \sum_{l_{r_1}=0}^{n/r_2^\gamma - r_1 + 1} (l_{r_1} + 1/2) \sum_{m_{r_2}=0}^{\infty} (m_{r_2} + 1/2) \\ &\times \left( \sum_{k=n/r_2^\gamma + 1}^{\infty} k^{-(\mu-2r_1+3/2)s/(s-1)} \sum_{j=m_{r_2} + r_2}^{\infty} j^{-(\mu-2r_2+3/2)s/(s-1)} \right)^{(s-1)/s} \end{aligned}$$

$$\begin{aligned}
&\leq c\|f\|_{s,\mu} n^{-(\mu-2r_1+3/2)+\frac{s-1}{s}} \sum_{l_{r_1}=0}^{n/r_2^\gamma-r_1+1} (l_{r_1}+1/2) \sum_{m_{r_2}=0}^{\infty} (m_{r_2}+1/2)^{1-(\mu-2r_2+3/2)+\frac{s-1}{s}} \\
&\leq c\|f\|_{s,\mu} n^{-\mu+2r_1-1/s+3/2}, \\
\|\Delta_{12}\|_C &\leq c\|f\|_{s,\mu} \sum_{l_{r_1}=n/r_2^\gamma-r_1+2}^{\infty} (l_{r_1}+1/2) \sum_{m_{r_2}=0}^{\infty} (m_{r_2}+1/2) \\
&\quad \times \left( \sum_{k=l_{r_1}+r_1}^{\infty} k^{-(\mu-2r_1+3/2)s/(s-1)} \sum_{j=m_{r_2}+r_2}^{\infty} j^{-(\mu-2r_2+3/2)s/(s-1)} \right)^{(s-1)/s} \\
&\leq c\|f\|_{s,\mu} \sum_{l_{r_1}=n/r_2^\gamma-r_1+2}^{\infty} (l_{r_1}+1/2)^{1-(\mu-2r_1+3/2)+\frac{s-1}{s}} \sum_{m_{r_2}=0}^{\infty} (m_{r_2}+1/2)^{1-(\mu-2r_2+3/2)+\frac{s-1}{s}} \\
&\leq c\|f\|_{s,\mu} n^{-\mu+2r_1-1/s+3/2}.
\end{aligned}$$

Thus, we get

$$\|\Delta_1\|_C \leq c\|f\|_{s,\mu} n^{-\mu+2r_1-1/s+3/2}.$$

Further, we estimate the norm of  $\Delta_2$  in the case of  $r_1 > r_2$  and  $\gamma = 1$ :

$$\begin{aligned}
\|\Delta_{21}\|_C &\leq c\|f\|_{s,\mu} n^{-\mu+2r_2-3/2+\frac{s-1}{s}} \sum_{l_{r_1}=0}^{n/r_2-r_1} (l_{r_1}+1/2) \sum_{m_{r_2}=0}^{\frac{n}{l_{r_1}+r_1}-r_2+1} (m_{r_2}+1/2) \\
&\quad \times \left( \sum_{k=l_{r_1}+r_1}^{\frac{n}{m_{r_2}+r_2}-1} k^{2(r_1-r_2)s/(s-1)-1} \right)^{(s-1)/s} \leq c\|f\|_{s,\mu} n^{-\mu+2r_1-3/2+\frac{s-1}{s}} \\
&\quad \times \sum_{l_{r_1}=0}^{n/r_2-r_1} (l_{r_1}+1/2) \sum_{m_{r_2}=0}^{\frac{n}{l_{r_1}+r_1}-r_2+1} (m_{r_2}+1/2)^{1-2(r_1-r_2)} \\
&\leq c\|f\|_{s,\mu} \begin{cases} n^{-\mu+2r_1-1/s+3/2} \ln n, & r_1 = r_2 + 1, \\ n^{-\mu+2r_1-1/s+3/2}, & r_1 > r_2 + 1, \end{cases} \\
\|\Delta_{22}\|_C &\leq c\|f\|_{s,\mu} \sum_{l_{r_1}=0}^{n/r_2-r_1} (l_{r_1}+1/2) \sum_{m_{r_2}=2}^{\frac{n}{l_{r_1}+r_1}-r_2+2} (m_{r_2}+1/2) \\
&\quad \times \left( \sum_{k=\frac{n}{m_{r_2}+r_2-2}}^{n/r_2} k^{-(\mu-2r_1+3/2)s/(s-1)} \sum_{j=m_{r_2}+r_2}^{\infty} j^{-(\mu-2r_2+3/2)s/(s-1)} \right)^{(s-1)/s} \\
&\leq c\|f\|_{s,\mu} n^{-\mu+2r_1-3/2+\frac{s-1}{s}} \sum_{l_{r_1}=0}^{n/r_2-r_1} (l_{r_1}+1/2) \sum_{m_{r_2}=2}^{\frac{n}{l_{r_1}+r_1}-r_2+2} (m_{r_2}+1/2)^{1-2(r_1-r_2)} \\
&\leq c\|f\|_{s,\mu} \begin{cases} n^{-\mu+2r_1-1/s+3/2} \ln n, & r_1 = r_2 + 1, \\ n^{-\mu+2r_1-1/s+3/2}, & r_1 > r_2 + 1, \end{cases}
\end{aligned}$$

$$\begin{aligned}
\|\Delta_{23}\|_C &\leq c\|f\|_{s,\mu} \sum_{l_{r_1}=0}^{n/r_2-r_1} (l_{r_1} + 1/2) \sum_{m_{r_2}=\frac{n}{l_{r_1}+r_1}-r_2+3}^{\infty} (m_{r_2} + 1/2) \\
&\times \left( \sum_{k=l_{r_1}+r_1}^{n/r_2} k^{-(\mu-2r_1+3/2)s/(s-1)} \sum_{j=m_{r_2}+r_2}^{\infty} j^{-(\mu-2r_2+3/2)s/(s-1)} \right)^{(s-1)/s} \\
&\leq c\|f\|_{s,\mu} \sum_{l_{r_1}=0}^{n/r_2-r_1} (l_{r_1} + 1/2)^{1-(\mu-2r_1+3/2)+\frac{s-1}{s}} \sum_{m_{r_2}=\frac{n}{l_{r_1}+r_1}-r_2+3}^{\infty} (m_{r_2} + 1/2)^{1-(\mu-2r_2+3/2)+\frac{s-1}{s}} \\
&\leq c\|f\|_{s,\mu} n^{-(\mu-2r_2+3/2)+\frac{s-1}{s}+2} \sum_{l_{r_1}=0}^{n/r_2-r_1} (l_{r_1} + 1/2)^{2(r_1-r_2)-1} \\
&\leq c\|f\|_{s,\mu} n^{-\mu+2r_1-1/s+3/2}.
\end{aligned}$$

Combining the estimates for  $\Delta_{21}(t, \tau)$ ,  $\Delta_{22}(t, \tau)$  and  $\Delta_{23}(t, \tau)$  we obtain

$$\|\Delta_2\|_C \leq c\|f\|_{s,\mu} \begin{cases} n^{-\mu+2r_1-1/s+3/2} \ln n, & r_1 = r_2 + 1, \\ n^{-\mu+2r_1-1/s+3/2}, & r_1 > r_2 + 1, \end{cases}.$$

To estimate  $\|\Delta_2\|_C$  for  $r_1 = r_2$  or  $\gamma > 1$ , it is enough to repeat an analogous line of reasoning. The combination of (2.3) and bounds for the norms of  $\Delta_1$ ,  $\Delta_2$  makes it possible to establish the desired inequality in the case of  $1 < s < \infty$ .

For  $s = 1$  the assertion of Lemma is proved similarly.  $\square$

The following statement contains upper estimates for the second difference from the right-hand side of (2.2) in the metric of  $C$ .

**Lemma 3.2** *Let  $f \in L_2(Q)$  and let the condition (1.2) be satisfied.*

*I) If  $r_1 = r_2$ , then*

a) *for  $\gamma = 1$  it holds true*

$$\|\mathcal{D}_{n,1}^{(r_1,r_1)} f - \mathcal{D}_{n,1}^{(r_1,r_1)} f_\delta\|_C \leq c\delta n^{2r_1-1/p+3/2} \ln^{2-1/p} n,$$

b) *for any  $1 < \gamma < \frac{2r_1+1/2+(p-1)/p}{2r_1-3/2+(p-1)/p}$  it holds true*

$$\|\mathcal{D}_{n,\gamma}^{(r_1,r_1)} f - \mathcal{D}_{n,\gamma}^{(r_1,r_1)} f_\delta\|_C \leq c\delta n^{2r_1-1/p+3/2},$$

c) *for  $\gamma = \frac{2r_1+1/2+(p-1)/p}{2r_1-3/2+(p-1)/p}$  it holds true*

$$\|\mathcal{D}_{n,\gamma}^{(r_1,r_1)} f - \mathcal{D}_{n,\gamma}^{(r_1,r_1)} f_\delta\|_C \leq c\delta n^{2r_1-1/p+3/2} \ln n.$$

*II) If  $r_1 = r_2 + 1$ , then*

a) *for  $\gamma = 1$  it holds true*

$$\|\mathcal{D}_{n,\gamma}^{(r_1,r_1-1)} f - \mathcal{D}_{n,\gamma}^{(r_1,r_1-1)} f_\delta\|_C \leq c\delta n^{2r_1-1/p+3/2} \ln n,$$

b) *for any  $\gamma > 1$  it holds true*

$$\|\mathcal{D}_{n,\gamma}^{(r_1,r_1-1)} f - \mathcal{D}_{n,\gamma}^{(r_1,r_1-1)} f_\delta\|_C \leq c\delta n^{2r_1-1/p+3/2}.$$

*III) If  $r_1 > r_2 + 1$ , then for any  $\gamma \geq 1$  it holds true*

$$\|\mathcal{D}_{n,\gamma}^{(r_1,r_2)} f - \mathcal{D}_{n,\gamma}^{(r_1,r_2)} f_\delta\|_C \leq c\delta n^{2r_1-1/p+3/2}.$$

*Proof.* Let  $1 < p < \infty$  first. Then, using the Hölder inequality and the estimate (2.9), we find

$$\begin{aligned}
\|\mathcal{D}_{n,\gamma}^{(r_1,r_2)} f - \mathcal{D}_{n,\gamma}^{(r_1,r_2)} f_\delta\|_C &\leq 2^{r_1+r_2} \sum_{l_{r_1}=0}^{n/r_2^\gamma-r_1} (l_{r_1} + 1/2) \sum_{m_{r_2}=0}^{\left(\frac{n}{l_{r_1}+r_1}\right)^{1/\gamma-r_2}} (m_{r_2} + 1/2) \\
&\times \sum_{k=l_{r_1}+r_1}^{\frac{n}{(m_{r_2}+r_2)^\gamma}} \sum_{j=m_{r_2}+r_2}^{\left(\frac{n}{k}\right)^{1/\gamma}} \sqrt{k+1/2} \sqrt{j+1/2} |\langle f - f_\delta, \varphi_{kj} \rangle| B_{k,j}^{r_1,r_2} \\
&\leq c\delta \sum_{l_{r_1}=0}^{n/r_2^\gamma-r_1} (l_{r_1} + 1/2) \sum_{m_{r_2}=0}^{\left(\frac{n}{l_{r_1}+r_1}\right)^{1/\gamma-r_2}} (m_{r_2} + 1/2) \\
&\times \left( \sum_{k=l_{r_1}+r_1}^{\frac{n}{(m_{r_2}+r_2)^\gamma}} k^{(2r_1-3/2)p/(p-1)} \sum_{j=m_{r_2}+r_2}^{\left(\frac{n}{k}\right)^{1/\gamma}} j^{(2r_2-3/2)p/(p-1)} \right)^{(p-1)/p} \\
&\leq c\delta n^{(2r_2-3/2)/\gamma + \frac{p-1}{\gamma p}} \sum_{l_{r_1}=0}^{n/r_2^\gamma-r_1} (l_{r_1} + 1/2) \sum_{m_{r_2}=0}^{\left(\frac{n}{l_{r_1}+r_1}\right)^{1/\gamma-r_2}} (m_{r_2} + 1/2) \\
&\times \left( \sum_{k=l_{r_1}+r_1}^{\frac{n}{(m_{r_2}+r_2)^\gamma}} k^{\frac{(2r_1-3/2)p}{p-1} - \frac{(2r_2-3/2)p}{\gamma(p-1)} - 1/\gamma} \right)^{(p-1)/p} \\
&\leq c\delta n^{2r_1-3/2 + \frac{p-1}{p}} \sum_{l_{r_1}=0}^{n/r_2^\gamma-r_1} (l_{r_1} + 1/2) \\
&\times \sum_{m_{r_2}=0}^{\left(\frac{n}{l_{r_1}+r_1}\right)^{1/\gamma-r_2}} (m_{r_2} + 1/2)^{1-\gamma(2r_1-3/2)+2r_2-3/2 + \frac{p-1}{p} - \frac{\gamma(p-1)}{p}}. \tag{3.1}
\end{aligned}$$

Whence it follows for  $r_1 = r_2 + 1$  and  $\gamma = 1$  that

$$\|\mathcal{D}_{n,1}^{(r_1,r_1-1)} f - \mathcal{D}_{n,1}^{(r_1,r_1-1)} f_\delta\|_C \leq c\delta n^{2r_1-3/2 + \frac{p-1}{p}} \ln n \sum_{l_{r_1}=0}^{n/r_2^\gamma-r_1} (l_{r_1} + 1/2) = c\delta n^{2r_1-1/p+3/2} \ln n,$$

which was required to prove.

Let's now  $r_1 = r_2 + 1$ ,  $\gamma > 1$  or  $r_1 > r_2 + 1$ ,  $\gamma \geq 1$ . Then from (3.1) we obtain the desired estimate

$$\|\mathcal{D}_{n,\gamma}^{(r_1,r_2)} f - \mathcal{D}_{n,\gamma}^{(r_1,r_2)} f_\delta\|_C \leq c\delta n^{2r_1-3/2 + \frac{p-1}{p}} \sum_{l_{r_1}=0}^{n/r_1-r_1} (l_{r_1} + 1/2) \leq c\delta n^{2r_1-1/p+3/2}.$$

To estimate  $\|\mathcal{D}_{n,\gamma}^{(r_1,r_2)} f - \mathcal{D}_{n,\gamma}^{(r_1,r_2)} f_\delta\|_C$  for  $r_1 = r_2$ , it is enough to repeat an analogous line of reasoning.

In the case of  $p = 1$  and  $p = \infty$  the assertion of Lemma is proved similarly.  $\square$

The combination of Lemmas 3.1 and 3.2 gives

**Theorem 3.3** Let  $f \in L_{s,2}^\mu$ ,  $1 \leq s < \infty$ ,  $r_1 \geq r_2$ ,  $\mu > 2r_1 - 1/s + 3/2$ , and let the condition (1.2) be satisfied.

I) If  $r_1 = r_2$ , then for  $n \asymp \left(\delta/\ln^{1/p-1/s} \frac{1}{\delta}\right)^{-\frac{1}{\mu-1/p+1/s}}$  and  $\gamma = 1$  it holds true

$$\|f^{(r_1, r_1)} - \mathcal{D}_{n,1}^{(r_1, r_1)} f_\delta\|_C \leq c \left(\delta \ln^{1/s-1/p} \frac{1}{\delta}\right)^{\frac{\mu-2r_1+1/s-3/2}{\mu-1/p+1/s}} \ln^{2-1/s} \frac{1}{\delta}.$$

II) If  $r_1 = r_2 + 1$ , then

a) for  $n \asymp \delta^{-\frac{1}{\mu-1/p+1/s}}$  and  $\gamma = 1$  it holds true

$$\|f^{(r_1, r_1-1)} - \mathcal{D}_{n,1}^{(r_1, r_1-1)} f_\delta\|_C \leq c \delta^{\frac{\mu-2r_1+1/s-3/2}{\mu-1/p+1/s}} \ln \frac{1}{\delta},$$

b) for  $n \asymp \delta^{-\frac{1}{\mu-1/p+1/s}}$  and any  $\gamma \in \left(1, \frac{\mu-2r_1+1/s+5/2}{\mu-2r_1+1/s+1/2}\right) \cup \left(\frac{\mu-2r_1+1/s+5/2}{\mu-2r_1+1/s+1/2}, \frac{\mu-2r_1+1/s+1/2}{\mu-2r_1+1/s-3/2}\right)$  it holds true

$$\|f^{(r_1, r_1-1)} - \mathcal{D}_{n,\gamma}^{(r_1, r_1-1)} f_\delta\|_C \leq c \delta^{\frac{\mu-2r_1+1/s-3/2}{\mu-1/p+1/s}},$$

c) for  $n \asymp \left(\delta/\ln \frac{1}{\delta}\right)^{-\frac{1}{\mu-1/p+1/s}}$  and  $\gamma = \frac{\mu-2r_1+1/s+1/2}{\mu-2r_1+1/s-3/2}$  it holds true

$$\|f^{(r_1, r_1-1)} - \mathcal{D}_{n,\gamma}^{(r_1, r_1-1)} f_\delta\|_C \leq c \left(\delta/\ln \frac{1}{\delta}\right)^{\frac{\mu-2r_1+1/s-3/2}{\mu-1/p+1/s}} \ln \frac{1}{\delta},$$

d) for  $n \asymp \left(\delta/\ln^{1-1/s} \frac{1}{\delta}\right)^{-\frac{1}{\mu-1/p+1/s}}$  and  $\gamma = \frac{\mu-2r_1+1/s+5/2}{\mu-2r_1+1/s+1/2}$  it holds true

$$\|f^{(r_1, r_1-1)} - \mathcal{D}_{n,\gamma}^{(r_1, r_1-1)} f_\delta\|_C \leq c \left(\delta/\ln^{1-1/s} \frac{1}{\delta}\right)^{\frac{\mu-2r_1+1/s-3/2}{\mu-1/p+1/s}} \ln^{1-1/s} \frac{1}{\delta}.$$

III) If  $r_1 > r_2 + 1$ , then

a) for  $n \asymp \delta^{-\frac{1}{\mu-1/p+1/s}}$  and any  $\gamma \in \left[1, \frac{\mu-2r_2+1/s-3/2}{\mu-2r_1+1/s+1/2}\right) \cup \left(\frac{\mu-2r_2+1/s-3/2}{\mu-2r_1+1/s+1/2}, \frac{\mu-2r_2+1/s+1/2}{\mu-2r_1+1/s+1/2}\right) \cup \left(\frac{\mu-2r_2+1/s+1/2}{\mu-2r_1+1/s+1/2}, \frac{\mu-2r_2+1/s-3/2}{\mu-2r_1+1/s-3/2}\right)$  it holds true

$$\|f^{(r_1, r_2)} - \mathcal{D}_{n,\gamma}^{(r_1, r_2)} f_\delta\|_C \leq c \delta^{\frac{\mu-2r_1+1/s-3/2}{\mu-1/p+1/s}},$$

b) for  $n \asymp \left(\delta/\ln \frac{1}{\delta}\right)^{-\frac{1}{\mu-1/p+1/s}}$  and  $\gamma = \frac{\mu-2r_2+1/s-3/2}{\mu-2r_1+1/s+1/2}$ ,  $\gamma = \frac{\mu-2r_2+1/s-3/2}{\mu-2r_1+1/s-3/2}$  it holds true

$$\|f^{(r_1, r_2)} - \mathcal{D}_{n,\gamma}^{(r_1, r_2)} f_\delta\|_C \leq c \left(\delta/\ln \frac{1}{\delta}\right)^{\frac{\mu-2r_1+1/s-3/2}{\mu-1/p+1/s}} \ln \frac{1}{\delta},$$

c) for  $n \asymp \left(\delta/\ln^{1-1/s} \frac{1}{\delta}\right)^{-\frac{1}{\mu-1/p+1/s}}$  and  $\gamma = \frac{\mu-2r_2+1/s+1/2}{\mu-2r_1+1/s+1/2}$  it holds true

$$\|f^{(r_1, r_2)} - \mathcal{D}_{n,\gamma}^{(r_1, r_2)} f_\delta\|_C \leq c \left(\delta/\ln^{1-1/s} \frac{1}{\delta}\right)^{\frac{\mu-2r_1+1/s-3/2}{\mu-1/p+1/s}} \ln^{1-1/s} \frac{1}{\delta}.$$

**Remark 3.4** The  $\mathcal{D}_n^{(r,r)}$  (2.1) method was studied earlier (see [23]) for the problem of numerical differentiation of functions from  $L_{s,2}^\mu$  in the case of  $r = 1$  and  $p = s = 2$ . Thus, the results of theorems 2.3 and 3.3 generalize studies [23] for the case of arbitrary  $r, p, s$ .

## 4. Minimal radius of Galerkin information

Let us turn to finding sharp estimates (in the power scale) for the minimal radius. First, we establish a lower estimate for the quantity  $R_{N,\delta}^{(r_1,r_2)}(L_{s,2}^\mu, C, \ell_p)$ . We fix an arbitrarily chosen domain  $\hat{\Omega}$ ,  $\text{card}(\hat{\Omega}) \leq N$ , of the coordinate plane  $[r_1, \infty) \times [r_2, \infty)$  and build an auxiliary function

$$f_1(t, \tau) = \tilde{c} \left( \varphi_0(t)\varphi_0(\tau) + N^{-\mu-1/s} r_2^{-\mu} \varphi_{r_2}(\tau) \sum_{k=N+r_1}^{3N+r_1} \varphi_k(t) \right),$$

where the sum  $\sum_{k=N+r_1}^{3N+r_1}$  is taken over any  $N$  pairwise distinct functions  $\varphi_k(t)$  such that  $N + r_1 \leq k \leq 3N + r_1$  and  $(k, r_2) \notin \hat{\Omega}$ . Obviously, there is at least one set of such functions.

Now we estimate the norm of  $f_1$  in the space metric  $L_{s,2}^\mu$ :

$$\|f_1\|_{s,\mu}^s = \tilde{c}^s \left( 1 + N^{-s\mu-1} \sum_{k=N+r_1}^{3N+r_1} k^{s\mu} \right) \leq \tilde{c}^s \left( 1 + 4^{s\mu} \right).$$

Whence it follows that to satisfy the condition  $\|f_1\|_{s,\mu} \leq 1$  it suffices to take

$$\tilde{c} = \left( 1 + 4^{s\mu} \right)^{-1/s}.$$

Next, we take another function from the class  $L_{s,2}^\mu$ :

$$f_2(t, \tau) = \tilde{c} \varphi_0(t)\varphi_0(\tau).$$

Let us find a lower bound for the quantity  $\|f_1^{(r_1,r_2)} - f_2^{(r_1,r_2)}\|_C$ . For this we need formulas

$$\begin{aligned} \varphi_r^{(r)}(t) &= \frac{\sqrt{r+1/2}}{2^{r-1/2}} \frac{(2r)!}{r!} \varphi_0(t), \quad f_2^{(r_1,r_2)}(t, \tau) \equiv 0, \\ f_1^{(r_1,r_2)}(t, \tau) &= \frac{\tilde{c}}{r_2^\mu} N^{-\mu-1/s} \varphi_{r_2}^{(r_2)}(\tau) \sum_{k=N+r_1}^{3N+r_1} \varphi_k^{(r_1)}(t) \\ &= 2^{r_1} \frac{\tilde{c}}{r_2^\mu} N^{-\mu-1/s} \varphi_{r_2}^{(r_2)}(\tau) \sum_{k=N+r_1}^{3N+r_1} \sqrt{k+1/2} \sum_{l_1=r_1-1}^{k-1} (l_1+1/2) \sum_{l_2=r_1-2}^{l_1-1} (l_2+1/2) \\ &\quad \dots \sum_{l_{r_1-1}=1}^{l_{r_1-2}-1} (l_{r_1-1}+1/2) \sum_{l_{r_1}=0}^{l_{r_1-1}-1} \sqrt{l_{r_1}+1/2} \varphi_{l_{r_1}}(t). \end{aligned} \quad (4.1)$$

It is easy to see that

$$\|f_1^{(r_1,r_2)} - f_2^{(r_1,r_2)}\|_C \geq |f_1^{(r_1,r_2)}(1, 1)| \geq \bar{c} N^{-\mu+2r_1-1/s+3/2},$$

where

$$\bar{c} = \frac{\tilde{c} \sqrt{r_2+1/2} (2r_2)!}{2^{r_1+r_2} r_2^\mu r_1! r_2!}.$$

Since for any  $1 \leq p \leq \infty$  it holds true

$$\|\bar{f}_1 - \bar{f}_2\|_{\ell_p} = \frac{\tilde{c}}{r_2^\mu} N^{-\mu-1/s+1/p},$$



then in the case of  $N^{-\mu-1/s+1/p} \leq r_2^\mu \delta / \tilde{c}$  under  $\delta$ -perturbations of the functions  $f_1$  and  $f_2$  can be considered

$$f_1^\delta(t, \tau) = f_2(t, \tau), \quad f_2^\delta(t, \tau) = f_1(t, \tau).$$

Let us find the upper bound for  $\|f_1^{(r_1, r_2)} - f_2^{(r_1, r_2)}\|_C$ . Taking into account the relation  $G(\hat{\Omega}, \bar{f}_1^\delta) = G(\hat{\Omega}, \bar{f}_2^\delta)$ , for any  $\psi^{(r_1, r_2)}(\hat{\Omega}) \in \Psi(\hat{\Omega})$  we find

$$\begin{aligned} \|f_1^{(r_1, r_2)} - f_2^{(r_1, r_2)}\|_C &\leq \|f_1^{(r_1, r_2)} - \psi^{(r_1, r_2)}(G(\hat{\Omega}, \bar{f}_1^\delta))\|_C + \|f_2^{(r_1, r_2)} - \psi^{(r_1, r_2)}(G(\hat{\Omega}, \bar{f}_2^\delta))\|_C \leq \\ &\leq 2 \sup_{f \in L_{s,2}^\mu, \|f\|_{s,\mu} \leq 1} \sup_{\bar{f}^\delta: (1.2)} \|f^{(r_1, r_2)} - \psi^{(r_1, r_2)}(G(\hat{\Omega}, \bar{f}^\delta))\|_C =: 2\varepsilon_\delta(L_{s,2}^\mu, \psi^{(r_1, r_2)}(\hat{\Omega}), C, \ell_p). \end{aligned}$$

That is

$$\varepsilon_\delta(L_{s,2}^\mu, \psi^{(r_1, r_2)}(\hat{\Omega}), C, \ell_p) \geq \frac{\bar{c}}{2} N^{-\mu+2r_1-1/s+3/2}.$$

From the fact that the domain  $\hat{\Omega}$  and the algorithm  $\psi^{(r_1, r_2)}(\hat{\Omega}) \in \Psi(\hat{\Omega})$  are arbitrary, follows that

$$R_{N,\delta}^{(r_1, r_2)}(L_{s,2}^\mu, C, \ell_p) \geq \frac{\bar{c}}{2} N^{-\mu+2r_1-1/s+3/2}.$$

Thus, the following assertion is proved.

**Theorem 4.1** *Let  $1 \leq s < \infty$ ,  $r_1 \geq r_2$ ,  $\mu > 2r_1 - 1/s + 3/2$ ,  $1 \leq p \leq \infty$ . Then for any  $N \geq \left(r_2^\mu \delta / \tilde{c}\right)^{-1/(\mu+1/s-1/p)}$  it holds true*

$$R_{N,\delta}^{(r_1, r_2)}(L_{s,2}^\mu, C, \ell_p) \geq \frac{\bar{c}}{2} N^{-\mu+2r_1-1/s+3/2}.$$

The following assertion contains sharp estimates (in the power scale) for the minimal radius in the uniform metric.

**Theorem 4.2** *Let  $f \in L_{s,2}^\mu$ ,  $1 \leq s < \infty$ ,  $r_1 \geq r_2$ ,  $\mu > 2r_1 + 1/2 - 3/s$ .*

I) *If  $r_1 = r_2$ , then for  $N \asymp \left(\delta / \ln^\mu \frac{1}{\delta}\right)^{-\frac{1}{\mu-1/p+1/s}}$  it holds true*

$$\begin{aligned} N^{-\mu+2r_1-1/s+3/2} &\preceq R_{N,\delta}^{(r_1, r_1)}(L_{s,2}^\mu, C, \ell_p) \preceq N^{-\mu+2r_1-1/s+3/2} \ln^{\mu-2r_1+1/2} N, \\ \left(\delta / \ln^\mu \frac{1}{\delta}\right)^{\frac{\mu-2r_1+1/s-3/2}{\mu-1/p+1/s}} &\preceq R_{N,\delta}^{(r_1, r_1)}(L_{s,2}^\mu, C, \ell_p) \preceq \left(\delta \ln^{1/s-1/p} \frac{1}{\delta}\right)^{\frac{\mu-2r_1+1/s-3/2}{\mu-1/p+1/s}} \ln^{2-1/s} \frac{1}{\delta}. \end{aligned}$$

The upper bounds are implemented by the method  $\mathcal{D}_{n,\gamma}^{(r_1, r_1)}$  (2.1) with  $n \asymp \left(\delta / \ln^{1/p-1/s} \frac{1}{\delta}\right)^{-\frac{1}{\mu-1/p+1/s}}$  and  $\gamma = 1$ .

II) *If  $r_1 = r_2 + 1$ , then for  $N \asymp \delta^{-\frac{1}{\mu-1/p+1/s}}$  it holds true*

$$R_{N,\delta}^{(r_1, r_1-1)}(L_{s,2}^\mu, C, \ell_p) \asymp N^{-\mu+2r_1-1/s+3/2} \asymp \delta^{\frac{\mu-2r_1+1/s-3/2}{\mu-1/p+1/s}}.$$

The order-optimal bounds are implemented by the method  $\mathcal{D}_{n,\gamma}^{(r_1, r_1-1)}$  (2.1) with  $n \asymp \delta^{-\frac{1}{\mu-1/p+1/s}}$  and any  $\gamma \in \left(1, \frac{\mu-2r_1+1/s+5/2}{\mu-2r_1+1/s+1/2}\right) \cup \left(\frac{\mu-2r_1+1/s+5/2}{\mu-2r_1+1/s+1/2}, \frac{\mu-2r_1+1/s+1/2}{\mu-2r_1+1/s-3/2}\right)$ .

III) *If  $r_1 > r_2 + 1$ , then for  $N \asymp \delta^{-\frac{1}{\mu-1/p+1/s}}$  it holds true*

$$R_{N,\delta}^{(r_1, r_2)}(L_{s,2}^\mu, C, \ell_p) \asymp N^{-\mu+2r_1-1/s+3/2} \asymp \delta^{\frac{\mu-2r_1+1/s-3/2}{\mu-1/p+1/s}}.$$

The order-optimal bounds are implemented by the method  $\mathcal{D}_{n,\gamma}^{(r_1, r_2)}$  (2.1) for  $n \asymp \delta^{-\frac{1}{\mu-1/p+1/s}}$  and any  $\gamma \in \left(1, \frac{\mu-2r_2+1/s-3/2}{\mu-2r_1+1/s+1/2}\right) \cup \left(\frac{\mu-2r_2+1/s-3/2}{\mu-2r_1+1/s+1/2}, \frac{\mu-2r_2+1/s+1/2}{\mu-2r_1+1/s+1/2}\right) \cup \left(\frac{\mu-2r_2+1/s+1/2}{\mu-2r_1+1/s+1/2}, \frac{\mu-2r_2+1/s-3/2}{\mu-2r_1+1/s-3/2}\right)$ .

**Proof.** The upper bounds for  $R_{N,\delta}^{(r_1,r_2)}(L_{s,2}^\mu, C, \ell_p)$  follow from Theorem 3.3. The lower bound is found in Theorem 4.1.  $\square$

Let's turn to estimating the minimal radius in the integral metric.

**Theorem 4.3** *Let  $1 \leq s < \infty$ ,  $r_1 \geq r_2$ ,  $\mu > 2r_1 - 1/s + 1/2$ ,  $1 \leq p \leq \infty$ . Then for any  $N \geq \left(r_2^\mu \delta / \tilde{c}\right)^{-1/(\mu+1/s-1/p)}$  it holds true*

$$R_{N,\delta}^{(r_1,r_2)}(L_{s,2}^\mu, L_2, \ell_p) \geq \bar{c} N^{-\mu+2r_1-1/s+1/2},$$

$$\text{where } \bar{c} = \frac{\tilde{c} \sqrt{r_2+1/2}}{2^{3r_1+r_2-3/2} r_2^\mu (r_1-1)! r_2!}.$$

**Proof** The proof of Theorem 4.3 almost completely coincides with the proof of Theorem 4.1, including the form of the auxiliary functions  $f_1, f_1^\delta, f_2, f_2^\delta$ . The only difference is in the lower estimate of the norm of  $f_1^{(r_1,r_2)} - f_2^{(r_1,r_2)}$ . Changing the order of summation in (4.1) yields to the representation

$$\begin{aligned} f_1^{(r_1,r_2)}(t, \tau) &= 2^{r_1} \frac{\tilde{c}}{r_2^\mu} N^{-\mu-1/s} \varphi_{r_2}^{(r_2)}(\tau) \\ &\times \left( \sum_{l_1=0}^N \sqrt{l_{r_1}+1/2} \varphi_{l_{r_1}}(t) \sum_{k=N+r_1}^{3N+r_1} \sqrt{k+1/2} + \sum_{l_1=N+1}^{3N} \sqrt{l_{r_1}+1/2} \varphi_{l_{r_1}}(t) \sum_{k=l_{r_1}+r_1}^{3N+r_1} \sqrt{k+1/2} \right) \\ &\times \sum_{l_1=l_{r_1}+r_1-1}^{k-1} (l_1+1/2) \sum_{l_2=l_{r_1}+r_1-2}^{l_1-1} (l_2+1/2) \dots \sum_{l_{r_1-1}=l_{r_1}+1}^{l_{r_1-2}-1} (l_{r_1-1}+1/2). \end{aligned}$$

Let us introduce the following notation

$$c' = \frac{2^{2(r_1-r_2)+1}}{r_2^{2\mu}} \tilde{c}^2 (r_2+1/2) \frac{((2r_2)!)^2}{(r_2!)^2}.$$

Then it follows that

$$\begin{aligned} \|f_1^{(r_1,r_2)} - f_2^{(r_1,r_2)}\|_{L_2}^2 &= \|f_1^{(r_1,r_2)}\|_{L_2}^2 \geq c' N^{-2\mu-2/s} \sum_{l_{r_1}=0}^N (l_{r_1}+1/2) \\ &\times \left( \sum_{k=N+r_1}^{3N+r_1} \sqrt{k+1/2} \sum_{l_1=l_{r_1}+r_1-1}^{k-1} (l_1+1/2) \sum_{l_2=l_{r_1}+r_1-2}^{l_1-1} (l_2+1/2) \dots \sum_{l_{r_1-1}=l_{r_1}+1}^{l_{r_1-2}-1} (l_{r_1-1}+1/2) \right)^2 \\ &\geq c' \frac{N^{-2\mu-2/s} \cdot N^{4r_1-1}}{4^{4(r_1-1)} ((r_1-1)!)^2} \sum_{l_{r_1}=0}^{N/2} (l_{r_1}+1/2). \end{aligned}$$

That is, the estimate holds

$$\|f_1^{(r_1,r_2)} - f_2^{(r_1,r_2)}\|_{L_2} \geq \frac{\sqrt{c'}}{4^{2r_1-1} (r_1-1)!} N^{-\mu+2r_1-1/s+1/2}.$$

Whence we obtain that the relation

$$\varepsilon_\delta(L_{s,2}^\mu, \psi^{(r_1,r_2)}(\hat{\Omega}), L_2, \ell_p) \geq \bar{c} N^{-\mu+2r_1-1/s+1/2}$$

is true for any  $N \geq \left(r_2^\mu \delta / \bar{c}\right)^{-1/(\mu+1/s-1/p)}$ . From the fact that the domain  $\hat{\Omega}$  and the algorithm  $\psi^{(r_1, r_2)}(\hat{\Omega}) \in \Psi(\hat{\Omega})$  are arbitrary, it follows that

$$R_{N, \delta}^{(r_1, r_2)}(L_{s,2}^\mu, L_2, \ell_p) \geq \bar{c} N^{-\mu+2r_1-1/s+1/2}.$$

Thus, the proof of Theorem 4.3 has been complied.  $\square$

The following assertion contains sharp estimates (in the power scale) for the minimal radius in the integral metric.

**Theorem 4.4** *Let  $f \in L_{s,2}^\mu$ ,  $1 \leq s < \infty$ ,  $r_1 \geq r_2$ ,  $\mu > 2r_1 + 1/2 - 1/s$ .*

I) *If  $r_1 = r_2$ , then for  $N \asymp \left(\delta / \ln^\mu \frac{1}{\delta}\right)^{-\frac{1}{\mu-1/p+1/s}}$  it holds true*

$$N^{-\mu+2r_1-1/s+1/2} \preceq R_{N, \delta}^{(r_1, r_1)}(L_{s,2}^\mu, L_2, \ell_p) \preceq N^{-\mu+2r_1-1/s+1/2} \ln^{\mu-2r_1+1} N,$$

$$\left(\delta / \ln^\mu \frac{1}{\delta}\right)^{\frac{\mu-2r_1+1/s-1/2}{\mu-1/p+1/s}} \preceq R_{N, \delta}^{(r_1, r_1)}(L_{s,2}^\mu, L_2, \ell_p) \preceq \left(\delta \ln^{1/s-1/p} \frac{1}{\delta}\right)^{\frac{\mu-2r_1+1/s-1/2}{\mu-1/p+1/s}} \ln^{3/2-1/s} \frac{1}{\delta}.$$

The upper bounds are implemented by the method  $\mathcal{D}_{n, \gamma}^{(r_1, r_1)}$  (2.1) with  $n \asymp \left(\delta^{-1} \ln^{1/p-1/s} \frac{1}{\delta}\right)^{\frac{1}{\mu-1/p+1/s}}$  and  $\gamma = 1$ .

II) *If  $r_1 > r_2$ , then for  $N \asymp \delta^{-\frac{1}{\mu-1/p+1/s}}$  it holds true*

$$R_{N, \delta}^{(r_1, r_2)}(L_{s,2}^\mu, L_2, \ell_p) \asymp N^{-\mu+2r_1-1/s+1/2} \asymp \delta^{\frac{\mu-2r_1+1/s-1/2}{\mu-1/p+1/s}}.$$

The order-optimal bounds are implemented by the method  $\mathcal{D}_{n, \gamma}^{(r_1, r_2)}$  (2.1) with  $n \asymp \delta^{-\frac{1}{\mu-1/p+1/s}}$  and any  $\gamma \in \left(1, \frac{\mu-2r_2+1/s-1/2}{\mu-2r_1+1/s+1/2}\right) \cup \left(\frac{\mu-2r_2+1/s-1/2}{\mu-2r_1+1/s+1/2}, \frac{\mu-2r_2+1/s+1/2}{\mu-2r_1+1/s+1/2}\right) \cup \left(\frac{\mu-2r_2+1/s+1/2}{\mu-2r_1+1/s+1/2}, \frac{\mu-2r_2+1/s-1/2}{\mu-2r_1+1/s-1/2}\right)$ .

**Proof.** The upper bounds for  $R_{N, \delta}^{(r_1, r_2)}(L_{s,2}^\mu, L_2, \ell_p)$  follow from Theorem 2.3. The lower bound is found in Theorem 4.3.  $\square$

## 5. Acknowledgements

This project has received funding through the MSCA4Ukraine project, which is funded by the European Union. In addition, the first named author is supported by the Volkswagen Foundation project "From Modeling and Analysis to Approximation". Also, the authors acknowledge partial financial support due to the project "Mathematical modelling of complex dynamical systems and processes caused by the state security" (Reg. No. 0123U100853).

## BIBLIOGRAPHY

1. S. Ahn, U.J. Choi, A.G. Ramm, A scheme for stable numerical differentiation // J. Comput. Appl. Math. – 2006.– Vol. 186(2).– P. 325–334
2. R. S. Anderssen, F. R. de Hoog, Finite difference methods for the numerical differentiation of non-exact data, Computing, 33(1984), 259–267.
3. Cullum, J., Numerical Differentiation and Regularization. SIAM Journal on Numerical Analysis.– 1971. – 8(2), 254–265. <http://www.jstor.org/stable/2949474>
4. T.F. Dolgoplova, V.K. Ivanov, On numerical differentiation // Zh. Vychisl. Mat. and Mat. Ph.– 1966. – Vol. 6(3).– P. 223–232.

5. Yu.V.Egorov, V.A.Kondrat'ev, On a problem of numerical differentiation // Vestnik Moskov. Univ. Ser. I Mat. Mekh. 3 (1989), 80-81.
6. Wolfgang Erb, Evgeniya V. Semenova, On adaptive discretization schemes for the solution of ill-posed problems with semiiterative methods// *Applicable Analysis*, Vol.94, 2015(10)
7. Groetsch C.W., Optimal order of accuracy in Vasin's method for differentiation of noisy functions // *J. Optim.Theory Appl.*– 1992.– Vol. 74(2).– P. 373–378.
8. M. Hanke, O. Scherzer, Inverse problems light: numerical differentiation // *Amer. Math. Monthly.*– 2001.– Vol. 108(6).– P. 512–521.
9. S. Lu, V. Naumova, S. V. Pereverzev, Legendre polynomials as a recommended basis for numerical differentiation in the presence of stochastic white noise // *J. Inverse Ill-Posed Probl.*– 2013.– Vol. 21(2).– P. 193–216.
10. Z. Meng, Z. Zhaoa, D. Mei, Y. Zhou, Numerical differentiation for two-dimensional functions by a Fourier extension method // *Inverse Problems in Science and Engineering.*– 2020.– Vol. 28(1).– P. 1–18.
11. Y.B. Wang, Y.C. Hon, J. Cheng Reconstruction of high order derivatives from input data *J. Inverse Ill-posed Probl.*, 14 (2006), pp. 205-218
12. G. Nakamura, S.Z. Wang, Y.B. Wang, Numerical differentiation for the second order derivatives of functions of two variables // *J. Comput. Appl. Math.*– 2008.– Vol. 212(2).– P. 341–358.
13. G.L. Myleiko, S.G. Solodky, The minimal radius of Galerkin information for severely ill-posed problems // *Journal of Inverse and Ill-Posed Problems.*– 2014.– Vol. 22(5).– P. 739–757.
14. G.L. Mileyko, S.G. Solodky, On optimization of projection methods for solving some classes of severely ill-posed problems // *Applicable Analysis.* – 2016. Vol.95(4), P.826-841.
15. G.L. Mileyko, S.G. Solodky, Hyperbolic cross and complexity of different classes of linear ill-posed problems // *Ukr. Mat. J.*– 2017.– Vol. 69(7).– P. 951–963.
16. Müller C., *Foundations of the Mathematical Theory of Electromagnetic Waves* /C. Müller – Springer-Verlag, Berlin, Heidelberg, New York, 1969.
17. S. V. Pereverzev, Optimization of projection methods for solving ill-posed problems, // *Computing.* – 1995 – Vol.55(2), P.113–124.
18. S. V. Pereverzev, S.G. Solodky, The minimal radius of Galerkin information for the Fredholm problem of the first kind, *Journal of Complexity* **12** (4) (1996), 401–415.
19. Qian Z., Fourier truncation method for high order numerical derivatives /Z. Qian, C.L. Fu, X.T. Xiong, T. Wei, // *Appl. Math. Comput.*– 2006.– Vol. 181(2).– P. 940–948.
20. Qu, Ruibin, A new approach to numerical differentiation and integration. *Mathematical and Computer Modelling* 24 (1996): 55-68.
21. Ramm A.G., On numerical differentiation // *Izv. Vuzov. Matem.*– 1968.– Vol. 11.– P. 131–134.
22. A.G. Ramm, A.B. Smirnova, On stable numerical differentiation *Math. Comput.*, 70 (2001), pp. 1131-1153
23. Semenova Y.V., Solodky S.G., Error bounds for Fourier-Legendre truncation method in numerical differentiation, // *Journal of Numerical and Applied Mathematics.*– 2021.– Vol. 137(3).–P.113–130.
24. Semenova, Evgeniya V., Solodky, Sergiy G. and Stasyuk, Serhii A., Application of Fourier Truncation Method to Numerical Differentiation for Bivariate Functions// *Computational Methods in Applied Mathematics*, vol. 22, no. 2, 2022, pp. 477-491. <https://doi.org/10.1515/cmam-2020-0138>
25. Semenova Y., Solodky S.G., Stasyuk S., Truncation method for numerical differentiation problem. *Proceedings of the Institute of Mathematics of the National Academy of Sciences of Ukraine*, Vol. 18 No. 1 (2021): Modern problems of mathematics and its applications, II
26. S. G. Solodky, S. A. Stasyuk, Estimates of efficiency for two methods of stable numerical summation of smooth functions, *Journal of Complexity* **56** (2020) <https://doi.org/10.1016/j.jco.2019.101422>
27. Solodky S.G., Stasyuk S., On optimization of methods of numerical differentiation for bivariate functions // *Ukr. Mat. J.*– 2022.– Vol. 74(2).– P. 253–273.
28. J. F. Traub, G. W. Wasilkowski, H. Wozniakowski, *Information-Based Complexity*, Academic Press, New York, 1988.

29. J. F. Traub, H. Wozniakowski, A General Theory of Optimal Algorithms, Academic Press, New York, 1980.
30. Vasin V.V., Regularization of the numerical differentiation problem /V.V. Vasin // Mat. app. Ural un-t.– 1969.– Vol. 7(2).– P. 29–33.
31. Zhao Z., A truncated Legendre spectral method for solving numerical differentiation // International Journal of Computer Mathematics.– 2010.– Vol. 87.– P. 3209–3217.
32. Z. Zhao, Z. Meng, L. Zhao, L. You, O. Xie, A stabilized algorithm for multi-dimensional numerical differentiation // Journal of Algorithms and Computational Technology.– 2016.– Vol. 10(2).– P. 73–81.

INSTITUTE OF MATHEMATICS, NATIONAL ACADEMY OF SCIENCES OF UKRAINE, 3, TERESCHENKIVSKA STR., 01024, KIEV, UKRAINE  
*E-mail address:* semenovaevgen@gmail.com, solodky@imath.kiev.ua

Received 11.02.2009; revised 23.03.2009