Balanced norm estimates for *rp*-Finite Element Methods applied to singularly perturbed fourth order boundary value problems

Torsten Linß* Christos Xer

Christos Xenophontos[†]

September 20, 2023

Abstract

We establish robust exponential convergence for rp-Finite Element Methods (FEMs) applied to fourth order singularly perturbed boundary value problems, in a *balanced norm* which is stronger than the usual energy norm associated with the problem. As a corollary, we get robust exponential convergence in the maximum norm. rp FEMs are simply p FEMs with possible repositioning of the (fixed number of) nodes. This is done for a C^1 Galerkin FEM in 1-D, and a C^0 mixed FEM in 2-D over domains with smooth boundary. In both cases we utilize the *Spectral Boundary Layer* mesh from [15].

Keywords: fourth order singularly perturbed problem, boundary layers, *rp*-finite element method, uniform, exponential convergence, balanced norm

AMS subject classification (2000): 65N30

1 Introduction

The quest for *balanced norm* estimates for singularly perturbed problems (SPPs) has been going on for a while now (see [7]–[10] and the references therein). When using the Finite Element Method (FEM), the error estimates are usually given in the *energy norm* defined by the bilinear form in the variational formulation of the problem. In certain cases, this norm is too weak and cannot see the layers present in the solution (see, e.g., [10]). While it is straight forward to define a balanced norm, the bilinear form lacks coercivity with respect to it, hence the usual methods of proof cannot be applied. Various approaches have been proposed [7]–[10], dealing mainly with reaction-diffusion type problems. The majority of these works pertains to fixed order FEMs; notable exceptions are [4, 13].

In this article we consider fourth order SPPs and the approximation of their solution by the rp version of the FEM on the Spectral Boundary Layer mesh from [15]. In this version the position (but not the number) of the nodes could possibly change, in addition to the polynomial degree p of the approximating polynomials, being increased. In the literature, one finds results on robust exponential convergence in the energy norm [2, 3, 17]. We prove robust exponential convergence in a balanced norm and as a corollary we get robust exponential convergence in the maximum norm. In one-dimension this is achieved for a C^1 Galerkin FEM, used in [2, 17], and in two-dimensions for the C^0 mixed formulation from [3]. The novelty in our approach lies with the choice of a certain projection of the solution onto the FE space. Specifically, we use an interpolant in the layer region and an appropriate projection in the rest of the domain. While this general approach has been used before (e.g., [9]), our contribution differs twofold: first we consider the rp version (hence exponential rates of convergence are obtained), and second, the projection is used only for the smooth parts of the solution, as opposed to the entire solution as was done in [9]. Finally, our contribution proves the conjectures made in [2] and [3] based on numerical evidence.

^{*}FernUniversität in Hagen, Fakultät für Mathematik und Informatik, Universitätsstraße 1, 58084, Hagen, email: torsten.linss@fernuni-hagen.de

[†]University of Cyprus, Department of Mathematics & Statistics, PO BOX 20537, 1678, Nicosia, email: xenophontos.christos@ucy.ac.cy

The rest of the paper is organized as follows: in Section 2 we consider a one-dimensional problem and its discretization by a C^1 Galerkin FEM. We present the available estimates from the literature and in Section 2.2 we improve them by establishing robust exponential convergence in a balanced, as well as the maximum norm. Section 3 contains the two-dimensional case, over domains with smooth boundary, and the C^0 mixed method from [3]. We, again, describe the estimates from the literature and then in Section 3.2 we improve them, just like in one-dimension.

Notation With $D \subset \mathbb{R}^d$, d = 1, 2, a domain with boundary ∂D and measure |D|, we will denote by $C^k(D)$ the space of continuous functions on D with continuous derivatives up to order k. We will use the usual Sobolev spaces $W^{k,q}(D)$, $q \in [0, \infty]$, $k \in \mathbb{N}_0$ of functions on D generalized derivatives of order $0, 1, \ldots, k$ in $L^q(D)$, equipped with the norm and seminorm $\|\cdot\|_{k,q,D}$ and $|\cdot|_{k,m,D}$, respectively. When q = 2, we will write $H^k(D)$ instead of $W^{k,2}(D)$, and for the norm and seminorm, we will write $\|\cdot\|_{k,D}$ and $|\cdot|_{k,D}$, respectively. The usual $L^2(D)$ inner product will be denoted by $\langle \cdot, \cdot \rangle_D$, with the subscript omitted when $D = \Omega$ and there is no confusion. We will also use the space

$$H_0^2(D) = \left\{ u \in H^2(D) \colon \left. u \right|_{\partial D} = \left. \frac{\partial u}{\partial n} \right|_{\partial D} = 0 \right\},\$$

where $\frac{\partial}{\partial n}$ denotes the normal derivative. The norm of the space $L^{\infty}(D) = W^{0,\infty}(D)$ of essentially bounded functions on *D* is denoted by $\|\cdot\|_{L^{\infty}(D)} = \|\cdot\|_{0,\infty,D} = |\cdot|_{0,\infty,D}$.

Finally, the notation " $a \leq b$ " means " $a \leq Cb$ " with C being a generic positive constant, independent of any discretization or singular perturbation parameters.

2 One-dimensional problems

Since the boundary layer effect is one-dimensional (in the direction normal to the boundary), studying one-dimensional problems is a necessary first step towards two (and higher) dimensions. In fact, most of the ideas are present in the one-dimensional case, and their extention to two-dimensions is (in most cases) straight forward.

We consider the following problem (cf. [2, 17]): find *u* such that

$$\varepsilon^2 u^{(4)} - (bu')' + cu = f \quad \text{in } \Omega := (0, 1),$$
 (1a)

$$u(0) = u(1) = u'(0) = u'(1) = 0,$$
(1b)

where $0 < \varepsilon \le 1$ is a given parameter that can approach zero and the functions *b*, *c* and *f* are given and sufficiently smooth. In particular, we assume that they are (real) *analytic* functions satisfying, for some positive constant γ_d , independent of ε ,

$$|f|_{n,\infty,\Omega} + |c|_{n,\infty,\Omega} + |b|_{n,\infty,\Omega} \leq n! \gamma_d^n \quad \forall \ n \in \mathbb{N}_0.$$
⁽²⁾

In addition, we assume that there exist positive constants b_{\min} and c_{\min} , independent of ε , such that

$$b \ge b_{\min}^2, \quad c \ge c_{\min}^2 \quad \text{on } \bar{\Omega},$$
(3)

It is well known (see, e.g. [16]) that the solution u to (1) can be decomposed into a smooth part, two boundary layer parts and a remainder. We have the following result from [17].

Proposition 1. Let u be the solution to (1), and assume (2) holds. Then, there exist a positive constants K, γ and δ such that for all $n \in \mathbb{N}_0$,

$$|u|_{n,\infty,\Omega} \lesssim K^n \max\left\{n^n, \varepsilon^{1-n}\right\}.$$
(4)

Moreover, u may be decomposed as

$$u = u_S + \tilde{u} + \bar{u} + u_R,\tag{5a}$$

with

$$|u_S|_{n,\infty,\Omega} \lesssim K^n n^n,\tag{5b}$$

$$|u_{S}|_{n,\infty,\Omega} \lesssim K^{n} n^{n},$$

$$\tilde{u}^{(n)}(x) \Big| + \Big| \bar{u}^{(n)}(1-x) \Big| \lesssim K^{n} \varepsilon^{1-n} \mathrm{e}^{-\gamma x/\varepsilon} \quad \forall \ x \in \bar{\Omega}$$
(5b)
(5c)

$$\|u_R\|_{2,\Omega} \lesssim e^{-\delta/\varepsilon}.$$
 (5d)

In this decomposition, u_S denotes the smooth part, \tilde{u} the boundary layer at the left endpoint, \bar{u} denotes the boundary layer at the right endpoint, and u_R is the remainder.

We mention that (4) corresponds to classical differentiability, while (5) correspond to regularity through asymptotic expansions, see [2, 12].

2.1 Discretization by a C^1 rp-FEM

The variational formulation of (1) reads: Find $u \in H^2_0(\Omega)$ such that

$$\mathcal{B}_{\varepsilon}(u,v) = \mathcal{F}(v) \quad \forall v \in H_0^2(\Omega), \tag{6}$$

where

$$\mathcal{B}_{\varepsilon}(w,v) \coloneqq \varepsilon^{2} \langle w'', v'' \rangle_{\Omega} + \langle bw', v' \rangle_{\Omega} + \langle cw, v \rangle_{\Omega}, \quad \mathcal{F}(v) \coloneqq \langle f, v \rangle_{\Omega}.$$
(7)

Because of (3), the bilinear form $\mathcal{B}_{\varepsilon}(\cdot, \cdot)$ induces a norm

$$\|w\|_E \coloneqq \mathcal{B}_{\varepsilon}(w, w), \tag{8}$$

the so called *energy norm*. Clearly, $\mathcal{B}_{\varepsilon}(\cdot, \cdot)$ is coercive (with constant 1) with respect to this norm.

We shall seek an approximation to u in a finite dimensional subspace of $H_0^2(\Omega)$. To this end, let \mathcal{P}_p , $p \in \mathbb{N}_0$, denote the space of polynomials of maximum degree p, and let $\Delta := \{x_i\}_{i=0}^N$ be an arbitrary subdivision of Ω with mesh intervals $\Omega_j := (x_{i-1}, x_i)$ and mesh sizes $h_i := x_i - x_{i-1} > 0$, i = 1, ..., N. We define the finite dimensional spaces

$$\mathcal{S}^p_{\Delta} \coloneqq \left\{ w \in C^1(\bar{\Omega}) \colon w|_{\Omega_i} \in \mathcal{P}_p, \ i = 1, \dots, N \right\}, \quad \mathcal{S}^p_{0,\Delta} \coloneqq \mathcal{S}^p_{\Delta} \cap H^2_0(\Omega).$$

Our discretization of (1) reads: Find $u_{p,\Delta} \in S_{0,\Delta}^p$ such that

$$\mathcal{B}_{\varepsilon}(u_{p,\Delta}, v) = \mathcal{F}(v) \quad \forall \ v \in \mathcal{S}_{0,\Delta}^{p}.$$

This discretization possesses the Galerkin orthogonality property

$$\mathcal{B}_{\varepsilon}\left(u-u_{p,\Delta},v\right)=0 \quad \forall \ v \in \mathcal{S}_{0,\Delta}^{p}$$

as well as best approximation property

$$\left\| u - u_{p,\Delta} \right\|_{E} = \inf_{v \in \mathcal{S}_{0,\Delta}^{p}} \| u - v \|_{E} .$$
⁽⁹⁾

Given $w \in H^2(\Omega)$, an interpolant $I^p_{\Delta} w \in S^p_{\Delta}$, $p \ge 3$, is uniquely defined by

$$(I_{\Delta}^{p}w - w)^{(k)}(x_{i}) = 0, \quad i = 0, \dots, N, \quad k = 0, 1$$

and

$$\int_{\Omega_j} \left(I_{\Delta}^p w - w \right)^{\prime\prime}(x) q(x) \, \mathrm{d}x = 0, \ \forall q \in \mathcal{P}_{p-2}, \ i = 0, \dots, N.$$

Lemma 2. Let $w \in H^{\ell}(\Omega_j)$, $\ell \ge 2$. Then for all $s, 0 \le s \le \min(p-1, \ell-2)$, the interpolation error satisfies the local bounds

$$(p-1)^{2-k} \left| w - I_{\Delta}^{p} w \right|_{k,\Omega_{j}}^{2} \le \left(\frac{\left| \Omega_{j} \right|}{2} \right)^{2(s+2-k)} \frac{(p-1-s)!}{(p+1+s)!} \left| w \right|_{s+2,\Omega_{j}}^{2}, \quad k = 0, 1, 2.$$

Proof. See, [18] and [1].

We next give the definition of the Spectral Boundary Layer Mesh we will utilize throughout the article.

Definition 3. For $\kappa > 0$, $p \in \mathbb{N}$ and $0 < \varepsilon \le 1$, define the Spectral Boundary Layer mesh Δ_{BL} as

 $\Delta_{BL} := \{0, \tau, 1 - \tau, 1\}, \quad \tau := \min\{\kappa p \varepsilon, 1/3\},$

with the layer and coarse-mesh regions

$$\Omega_{\ell} \coloneqq (0,\tau) \cup (1-\tau,1), \quad \Omega_{c} \coloneqq (\tau,1-\tau).$$

We shall consider the sequence $u_p \in S^p_{0,\Delta_{Rl}}$, $p = 3, 4, \dots$, of approximations defined by

$$\mathcal{B}_{\varepsilon}\left(u_{p},v\right) = \mathcal{F}(v) \quad \forall \ v \in \mathcal{S}_{0,\Delta_{BL}}^{p}.$$
(10)

Given $w \in H^2(\Omega)$, we define interpolants $I_p w := I_{\Delta_{BL}}^p w, p = 3, 4, \dots$

The best approximation property (9) implies

$$||u - u_p||_E \le ||u - I_p u||_E$$
, $p = 3, 4, ...$

Proposition 4. Let u be the solution to (1) and assume that (2) holds. Then there exist positive constants κ_0 , κ_1 and $\beta > 0$ independent of ε and p, such that the following is true: For every $p \in \mathbb{N}$ and every $\kappa \in (0, \kappa_0]$ with $\kappa p \ge \kappa_1$, the interpolation error satisfies

$$\left\| u - I_p u \right\|_{1,\infty,\Omega} + \varepsilon^{1/2} \left| u - I_p u \right|_{2,\Omega} \lesssim e^{-\beta p}.$$
 (11)

Moreover,

$$\left\| u_{S} - I_{p} u_{S} \right\|_{2,\Omega} \lesssim e^{-\beta p},\tag{12}$$

and

$$u - I_p u \Big|_{k,\Omega_{BL}} \lesssim \varepsilon^{3/2-k} \mathrm{e}^{-\beta p}.$$
⁽¹³⁾

Proof. Inequality (11) was established in [17] (see also [2]), and inequalities (12), (13) are shown in the Appendix. \Box

Using (9) and (11) the following may be established, cf. [2].

Proposition 5. Let u solve (1) and let u_p , p = 3, 4, ..., be its finite element approximations obtained by (10). Then there exists a constant $\sigma > 0$, independent of ε but depending on the data, such that

$$\left\| u - u_p \right\|_E \lesssim e^{-\sigma p}.$$

2.2 Balanced norm estimates

The energy norm used in Proposition 5 is not the appropriate norm for this problem, since with

$$u_{BL} := \tilde{u} + \bar{u}$$

we calculate

$$\|u_{BL}\|_{E} = O\left(\varepsilon^{1/2}\right)$$
 while $\|u_{S}\|_{E} = O(1)$, $\varepsilon \to 0$

Thus, the energy norm does not see the layers present in the solution of (1), as $\varepsilon \to 0$. The correct weight for $|u|_{2,\Omega}$ is $\varepsilon^{1/2}$ (as opposed to ε , cf. (8)). In which case, the resulting norm would be *balanced*, like the one below:

$$\|u\|_{B}^{2} \coloneqq \varepsilon \|u\|_{2,\Omega}^{2} + \|u\|_{1,\Omega}^{2}.$$
(14)

In this norm we have

$$\|u_{BL}\|_{\mathcal{B}} = O(1) = \|u_{S}\|_{\mathcal{B}}, \quad \varepsilon \to 0.$$

Unfortunately, the bilinear form $\mathcal{B}_{\varepsilon}$ given by (7), is not coercive with respect to this norm. Nevertheless, numerical experiments reported in [2], suggest the following:

$$\left\| u - u_p \right\|_{1,\infty,\Omega} + \left\| u - u_p \right\|_B \lesssim e^{-\sigma p}$$
⁽¹⁵⁾

Our goal in this section is to establish (15).

We begin by studying the auxiliary problem of finding $u_{S,p} \in \mathcal{P}_p$, $p \ge 3$, such that

$$\mathcal{B}_0(u_{S,p} - u_S, v) = 0 \quad \forall \ v \in \mathcal{P}_p \cap H^1_0(\Omega), \quad u_{S,p} - u_S = 0 \text{ on } \partial\Omega$$
(16)

where

$$\mathcal{B}_0(w,v) \coloneqq \langle bw',v' \rangle + \langle cw,v \rangle.$$

Note that \mathcal{B}_0 induces a norm $|||w||| := \mathcal{B}_0(w, w)^{1/2}$ that is equivalent to the standard $H^1(\Omega_{REG})$ norm because of (3). Consequently, $u_{S,p}$ is uniquely defined and we have the following.

Lemma 6. Let $u_{S,p} \in \mathcal{P}_p$, $p \ge 3$, be defined by (16). Then,

$$\begin{aligned} \left| u_{S,p} - u_S \right|_k &\leq e^{-\beta p}, \quad k = 0, 1, 2\\ \left\| u_{S,p} - u_S \right\|_{C^1(\bar{\Omega})} &\leq e^{-\beta p} \end{aligned}$$

and

$$\left|\left(u_{S,p}-u_{S}\right)(x)\right|\lesssim\min(x,1-x)\mathrm{e}^{-\beta p},\ x\in\bar{\Omega}.$$

Proof. Note, that the bilinear form in (16) is coercive and bounded. So standard techniques for spectral methods apply to give the bounds of the first inequality for k = 0 and k = 1.

For k = 2 we proceed as follows. We have

$$||u_{S,p} - u_S||| \le ||v - u_S||| \quad \forall v \in \mathcal{P}_p \text{ with } v = u_S \text{ on } \partial\Omega,$$

which implies

$$|||u_{S,p} - I_p u_S||| \le 2 |||I_p u_S - u_S|||,$$

by the triangle inequality. Next

$$\begin{aligned} \left| u_{S,p} - u_{S} \right|_{2} &\leq \left| u_{S,p} - I_{p} u_{S} \right|_{2} + \left| I_{p} u_{S} - u \right|_{2} \leq (p-1)^{2} \left| u_{S,p} - I_{p} u_{S} \right|_{1} + \left| I_{p} u_{S} - u \right|_{2} \\ &\leq 2 b_{0}^{-1} (p-1)^{2} \left| \left| \left| I_{p} u_{S} - u_{S} \right| \right|_{1} + \left| I_{p} u_{S} - u \right|_{2} \right|_{2} \end{aligned}$$

where we have used an inverse inequality, see Lemma 19. Now the desired result follows from (12).

A Sobolev embedding gives the bound in the C^1 norm, while the final bound follows upon noting that $(u_{S,p} - u_S)(x) = 0$ for $x \in \{0, 1\}$ and the maximum-norm bound for the derivative.

Next, we construct a special FE-function $\hat{u}_p \in S^p_{0,\Delta_{BL}}$ that is particularly close to u. To this end, define auxiliary functions $\chi_0, \chi_1 \in \mathcal{P}_3$ by

$$\chi_0(0) = \chi'_0(0) = \chi_0(\tau) = 0, \quad \chi'_0(\tau) = 1,$$

and

$$\chi_1(0) = \chi'_1(0) = \chi'_1(\tau) = 0, \quad \chi_1(\tau) = 1.$$

A direct calculation establishes the following bounds on the χ_i :

$$|\chi_i|_{k,(0,\tau)} \lesssim \tau^{3/2-k-i}, \ k \in \{0,1,2\}, \ i \in \{0,1\}.$$
 (17)

Now, set

$$\hat{u}_{p}(x) \coloneqq \begin{cases} \left(I_{p}u\right)(x) + \chi_{0}(x)\left(u_{S,p} - u\right)'(\tau) + \chi_{1}(x)\left(u_{S,p} - u\right)(\tau), & x \in [0,\tau], \\ u_{S,p}(x), & x \in [\tau, 1 - \tau], \\ (I_{p}u)(x) - \chi_{0}(1 - x)\left(u_{S,p} - u\right)'(1 - \tau), + \chi_{1}(1 - x)\left(u_{S,p} - u\right)(1 - \tau), & x \in [1 - \tau, 1]. \end{cases}$$

$$(18)$$

We re-iterate that in the layer region, \hat{u}_p is given as the interpolant $I_p u$ of u plus corrections that ensure $\hat{u}_p \in C^1(\bar{\Omega})$. In the regular region, we *only* apply the projection on u_s . We have the following.

Lemma 7. Assume that (5) holds and that \hat{u}_p is defined by (18). Then the following bounds hold true:

$$\left| u - \hat{u}_p \right|_{k,\Omega_{BL}} \lesssim \varepsilon^{3/2-k} \mathrm{e}^{-\beta p}, \qquad k \in \{0, 1, 2\},$$
(19a)

$$\left|u - \hat{u}_p\right|_{k,\Omega_{REG}} \lesssim \max\left(1,\varepsilon^{3/2-k}\right) e^{-\beta p}, k \in \{0,1,2\}.$$
(19b)

Proof. We shall study the two regions Ω_{BL} and Ω_{REG} separately.

 Ω_{BL} : We present the argument for $(0, \tau)$, because identical bounds hold for the interval $(1 - \tau, 1)$. We have

$$\left| u - \hat{u}_p \right|_{k,(0,\tau)} \le \left| u - I_p u \right|_{k,(0,\tau)} + \left| \chi_0 \right|_{k,(0,\tau)} \cdot \left| \left(u_{S,p} - u \right)'(\tau) \right| + \left| \chi_1 \right|_{k,(0,\tau)} \cdot \left| \left(u_{S,p} - u \right)(\tau) \right|, \ k \in \{0, 1, 2\}.$$

Proposition 4 takes care of the first term above. For the other two terms, a triangle inequality, Lemma 6, (5c) and (5d) yield

$$\left| \left(u_{S,p} - u \right)^{(\ell)} (\tau) \right| = \left| \left(u_{S,p} - u_S - u_{BL} - u_R \right)^{(\ell)} (\tau) \right| \le \tau^{1-\ell} \mathrm{e}^{-\beta p} \,, \ \ell \in 0, 1.$$

Recalling (17), we get

$$\left|\chi_{1-\ell}\right|_{k,(0,\tau)} \left| \left(u_{S,p} - u \right)^{(\ell)}(\tau) \right| \lesssim \tau^{3/2-k} \mathrm{e}^{-\beta p}, \ \ell \in 0, 1,$$

and (19a) follows. We mention here that we will routinely "hide" polynomial powers into exponentials, e.g. $p^s e^{-\beta p} \leq e^{-\beta p} \forall s \in \mathbb{R}$.

 Ω_{REG} : We have from the definition of \hat{u}_p ,

$$\left| u - \hat{u}_p \right|_{k,\Omega_{REG}} \le \left| u_S - u_{S,p} \right|_{k,\Omega_{REG}} + |u_{BL}|_{k,\Omega_{REG}} + |u_R|_{k,\Omega_{REG}}, \ k = 0, 1, 2.$$

The first term has been bounded in Lemma 6. For u_{BL} we have by (5c), and because $x \in [\tau, 1 - \tau]$,

$$|u_{BL}|_{k,\Omega_{REG}} \lesssim \varepsilon^{3/2-k} \mathrm{e}^{-\gamma\kappa p}, \ k = 0, 1, 2,$$

while for u_R , ineq. (5d) gives

$$|u_R|_{k,\Omega_{REG}} \lesssim e^{-3\delta\kappa p}, \ k = 0, 1, 2.$$

Combining the last two inequalities with Lemma 6, we obtain (19b).

We now have the necessary tools for proving the following, main result of this section.

Theorem 8. Let u be the solution of (6) and let u_p be the approximation obtained by (10). Then, there exists a positive constant β , independent of ε , such that

$$\left\|u-u_p\right\|_B \lesssim \mathrm{e}^{-\beta p}$$

Proof. By (14)

$$\|u - u_p\|_B^2 = \varepsilon \|u - u_p\|_{2,\Omega}^2 + \|u - u_p\|_{1,\Omega}^2$$

The only "troublesome" term above is $\varepsilon |u - u_p|_{2,\Omega}$. (The other one is handled by Proposition 5). To deal with it, we use the triangle inequality (with \hat{u}_p defined by (18)) to get

$$\varepsilon^{1/2} \left| u - u_p \right|_{2,\Omega} \le \varepsilon^{1/2} \left| u - \hat{u}_p \right|_{2,\Omega} + \varepsilon^{1/2} \left| \hat{u}_p - u_p \right|_{2,\Omega}.$$

Only the term $\varepsilon^{1/2} |\hat{u}_p - u_p|_{2,\Omega}$ needs to be considered, since the rest can be bounded by Lemma 7. Let $\eta := \hat{u}_p - u_p \in S^p_{0,\Delta_{R'}}$. Galerkin orthogonality gives

$$\|\eta\|_{E}^{2} = \varepsilon^{2} \left\langle \left(\hat{u}_{p} - u\right)^{\prime\prime}, \eta^{\prime\prime} \right\rangle + \left\langle b(\hat{u}_{p} - u)^{\prime}, \eta^{\prime} \right\rangle + \left\langle c(\hat{u}_{p} - u), \eta \right\rangle$$

On the coarse mesh region Ω_{REG} , we have

$$\begin{split} \left\langle b\left(\hat{u}_{p}-u\right)',\eta'\right\rangle_{\Omega_{REG}} + \left\langle c\left(\hat{u}_{p}-u\right),\eta\right\rangle_{\Omega_{REG}} \\ &= \left\langle b\left(u_{S,p}-u_{S}\right)',\eta'\right\rangle_{\Omega_{REG}} + \left\langle c\left(u_{S,p}-u_{S}\right),\eta\right\rangle_{\Omega_{REG}} \\ &- \left\langle b\left(u_{BL}+u_{R}\right)',\eta'\right\rangle_{\Omega_{REG}} - \left\langle c\left(u_{BL}+u_{R}\right),\eta\right\rangle_{\Omega_{REG}} \\ &= -\left\langle b\left(u_{S,p}-u_{S}\right)',\eta'\right\rangle_{\Omega_{BL}} - \left\langle c\left(u_{S,p}-u_{S}\right),\eta\right\rangle_{\Omega_{BL}} \\ &- \left\langle b\left(u_{BL}+u_{R}\right)',\eta'\right\rangle_{\Omega_{REG}} - \left\langle c\left(u_{BL}+u_{R}\right),\eta\right\rangle_{\Omega_{REG}} \end{split}$$

where the definition of \hat{u}_p was used. Now, the C¹-norm bounds of Lemma 6, (5c) and (5d) yield

$$\begin{split} \left| \left\langle b\left(\hat{u}_p - u\right)', \eta' \right\rangle_{\Omega_{REG}} + \left\langle c\left(\hat{u}_p - u\right), \eta \right\rangle_{\Omega_{REG}} \right| \\ \lesssim \tau^{1/2} \left\| u_{S,p} - u_S \right\|_{C^1(\bar{\Omega}_{BL})} \|\eta\|_{1,\Omega_{BL}} + \|u_{BL} + u_R\|_{1,\Omega_{REG}} \|\eta\|_{1,\Omega_{REG}} \lesssim \varepsilon^{1/2} \mathrm{e}^{-\beta p} \|\eta\|_{1,\Omega} \,. \end{split}$$

Similarly, using the H^k -norm bounds of Lemma 6, (5c) and (5d), we get

$$\varepsilon^{2} \left| \left\langle (\hat{u}_{p} - u)^{\prime\prime}, \eta^{\prime\prime} \right\rangle_{\Omega_{REG}} \right|$$

$$\leq \varepsilon^{2} \left| \left\langle (u_{S,p} - u_{S})^{\prime\prime}, \eta^{\prime\prime} \right\rangle_{\Omega_{REG}} \right| + \varepsilon^{2} \left| \left\langle (u_{BL} + u_{R})^{\prime\prime}, \eta^{\prime\prime} \right\rangle_{\Omega_{REG}} \right| \lesssim \varepsilon^{3/2} \mathrm{e}^{-\beta p} \left| \eta \right|_{2,\Omega_{REG}} .$$

In the layer region Ω_{BL} , we use the Cauchy-Schwarz inequality and (19a) to obtain

$$\left|\left\langle b\left(\hat{u}_{p}-u\right)',\eta'\right\rangle_{\Omega_{BL}}+\left\langle c\left(\hat{u}_{p}-u\right),\eta\right\rangle_{\Omega_{BL}}\right|\lesssim\varepsilon^{1/2}\mathrm{e}^{\beta p}\left\|\eta\right\|_{1,\Omega_{BL}}$$

and

$$\varepsilon^{2} \left| \left\langle (\hat{u}_{p} - u)^{\prime\prime}, \eta^{\prime\prime} \right\rangle_{\Omega_{BL}} \right| \lesssim \varepsilon^{1/2} \mathrm{e}^{\beta p} \varepsilon \left| \eta \right|_{2, \Omega_{BL}}.$$

Combining the above we have

$$\left\|\eta\right\|_{E}^{2} \lesssim \varepsilon^{1/2} \mathrm{e}^{-\beta p} \left\|\eta\right\|_{E}.$$

Dividing by $\|\eta\|_E$ yields

$$\varepsilon^{1/2} \left| u_p - \hat{u}_p \right|_{2,\Omega} \lesssim \left\| u_p - \hat{u}_p \right\|_E \lesssim \mathrm{e}^{-\beta p},$$

and from the triangle inequality we have

$$\varepsilon^{1/2} \left| u_p - u \right|_{2,\Omega} \lesssim \mathrm{e}^{-\beta p}$$

This completes the proof.

Once we have the balanced norm estimate, we may obtain a maximum norm estimate as follows.

Corollary 9. Let u be the solution of (6) and let u_p be the solution of (10), based on the Spectral Boundary Layer mesh of Definition 3. Then, there exists a positive constant β , independent of ε , such that

$$\left\|u-u_p\right\|_{C^1(\bar{\Omega})} \lesssim \mathrm{e}^{-\beta p}$$

Proof. Let $x \in [0, \tau]$ be arbitrary. Then, using the boundary conditions (1b), we have

$$(u - u_p)^{(k)}(x) = \int_0^x \left(u - u_p\right)^{(k+1)}(t) \, \mathrm{d}t, \ k \in \{0, 1\},$$

and by the Cauchy-Schwarz inequality

$$\left| (u-u_p)^{(k)}(x) \right| \lesssim \tau^{1/2} \left| u-u_p \right|_{k+1,\Omega_{BL}} \lesssim (p\varepsilon)^{1/2} \varepsilon^{-1/2} \mathrm{e}^{-\beta p} \lesssim \mathrm{e}^{-\beta p},$$

where Theorem 8 was used.

The same technique works for the other layer region $[1 - \tau, 1]$, so let us consider the coarse region Ω_{REG} . We use the triangle inequality, with I_p the interpolant of Proposition 4, to get

$$\left|u-u_{p}\right|_{k,\infty,\Omega_{REG}} \leq \left|u-I_{p}u\right|_{k,\infty,\Omega_{REG}} + \left|I_{p}u-u_{p}\right|_{k,\infty,\Omega_{REG}}.$$

The term $|u - I_p u|_{k,\infty,\Omega_{RFG}}$ is handled by Proposition 4, while for the second term an inverse inequality gives

$$\left|I_{p}u-u_{p}\right|_{k,\infty,\Omega_{REG}} \lesssim \left|I_{p}u-u_{p}\right|_{k,2,\Omega_{REG}} \leq \left|I_{p}u-u\right|_{k,2,\Omega_{REG}} + \left|u-u_{p}\right|_{k,2,\Omega_{REG}}.$$

Application of Propositions 4 and 5 completes the proof.

Remark 10. Theorem 8 and Corollary 9 establish the conjecture (15) made in [2].

3 Two-dimensional problems

In two dimensional smooth domains $\Omega \subset \mathbb{R}^2$, we would have great difficulty constructing a C^1 conforming FEM; instead, a C^0 mixed formulation is usually preferred. Before we present the problem under consideration along with existing results, we comment on the differences between the C^1 method and the C^0 mixed method, beyond the obvious ones. Specifically, we want to point out that in order to use the previous methodology, we should define two special representatives, since in the mixed formulation we will be seeking two unknown functions – the second one will be auxiliary. Moreover, for the extra function we only need to use the L^2 projection, while for u we (again) utilize the H^1 projection.

We consider the following problem¹ from [3]: Find *u* such that

$$\varepsilon^2 \Delta^2 u - b \Delta u + c u = f \quad \text{in } \Omega \subset \mathbb{R}^2, \tag{20a}$$

$$u = \frac{\partial u}{\partial n} = 0 \quad \text{on } \partial\Omega, \tag{20b}$$

where $0 < \varepsilon \le 1$ is a given parameter, Δ denotes the Laplacian (and Δ^2 the biharmonic) operator, b, c > 0 are given constants, Ω is an open bounded domain with analytic boundary $\partial \Omega$ and f is a given analytic function, which satisfies

$$\|\nabla^n f\|_{\infty,\Omega} \lesssim n! \gamma_f^n \text{ for } n = 0, 1, 2, \dots,$$

$$(21)$$

for some positive constant γ_f independent of ε . Here we have used the shorthand notation

$$|\nabla^n f|^2 := \sum_{|\alpha|=n} \frac{|\alpha|!}{\alpha!} |D^{\alpha} f|^2,$$

. 1		
. 1		

¹Henceforth, the symbols that appeared in the previous section will represent two-dimensional analogs.

with $D^{\alpha}f$ denoting the generalized derivative of f with respect to the multiindex $\alpha \in \mathbb{N}_{0}^{2}$.

The regularity of the solution to (20) was studied in [2, 13], and will be described in Proposition 1 below. To this end, define *boundary fitted coordinates* (ρ , θ) in a neighborhood of the boundary as follows: Let ($X(\theta), Y(\theta)$), $\theta \in [0, L]$ be a parametrization of $\partial\Omega$ by arclength and let Ω_0 be a tubular neighborhood of $\partial\Omega$ in Ω . For each point $z = (x, y) \in \Omega_0$ there is a unique nearest point $z_0 \in \partial\Omega$, so with θ the arclength parameter (with counterclockwise orientation), we set $\rho = |z - z_0|$ which measures the distance from the point z to $\partial\Omega$. Explicitly, let $\rho_0 > 0$ be less than the minimum radius of curvature of $\partial\Omega$ and set

$$\Omega_0 \coloneqq \{ z - \rho \overrightarrow{n}_z \colon z \in \partial \Omega, \ \rho \in (0, \rho_0) \}, \tag{22}$$

where \overrightarrow{n}_z is the outward unit normal at $z \in \partial \Omega$, and

$$x = X(\theta) - \rho Y'(\theta), \ y = Y(\theta) + \rho X'(\theta),$$

with $\rho \in (0, \rho_0), \theta \in (0, L)$; see Figure 1.



Figure 1: Boundary fitted coordinates.

Proposition 11. The solution u of (20) can be decomposed into a smooth part u^S , a boundary layer part u^{BL} and a remainder u^R , viz.

$$u = u^S + \chi u^{BL} + u^R, \tag{23a}$$

where χ is a smooth cut-off function, satisfying

$$\chi = \begin{cases} 1 & \text{for } 0 < \rho < \rho_0/3, \\ 0 & \text{for } \rho > 2\rho_0/3. \end{cases}$$

Moreover, there exist constants $K_1, K_2, \omega, \delta > 0$, independent of ε , such that

$$\left\| D^{n} u^{S} \right\|_{0,\Omega} \lesssim |n|! K_{1}^{|n|} \quad \forall \ n \in \mathbb{N}_{0}^{2},$$
(23b)

$$\left|\frac{\partial^{m+n} u^{BL}(\rho,\theta)}{\partial \rho^m \partial \theta^n}\right| \leq n! K_2^{m+n} \varepsilon^{1-m} \mathrm{e}^{-\omega \rho/\varepsilon} \quad \forall \ m, n \in \mathbb{N}, \ (\rho,\theta) \in \overline{\Omega}_0,$$
(23c)

and

$$\varepsilon \left| u^{R} \right|_{2,\Omega} + \left\| u^{R} \right\|_{1,\Omega} \lesssim e^{-\delta/\varepsilon}.$$
(23d)

Finally, there exist constants C, K > 0, depending only on the data, such that

$$\|D^{n}u\|_{0,\Omega} \le CK^{|n|} \max\left\{|n|^{|n|}, \varepsilon^{1-|n|}\right\} \quad \forall \ n \in \mathbb{N}_{0}^{2}.$$
(23e)

Proof. All estimates were shown in [2], except for (23b) which appears in [14]. \Box

Letting² $w := \varepsilon \Delta u \in H^2(\Omega)$, we shall study the *mixed* formulation of finding $(u, w) \in H^1_0(\Omega) \times H^1(\Omega)$ such that

$$\varepsilon \langle \nabla u, \nabla \phi \rangle + \langle w, \phi \rangle = 0 \qquad \forall \ \phi \in H^1(\Omega), \tag{24a}$$

$$b\langle u, \nabla \rangle + c\langle u, \psi \rangle - \varepsilon \langle \nabla w, \nabla \rangle = \langle f, \psi \rangle \ \forall \ \psi \in H^1_0(\Omega).$$
(24b)

We define the bilinear form

$$\mathcal{A}_{\varepsilon}((u,w),(\psi,\phi)) \coloneqq \varepsilon \left\langle \nabla u, \nabla \phi \right\rangle + \left\langle w, \phi \right\rangle + b \left\langle \nabla u, \nabla \psi \right\rangle + c \left\langle u, \psi \right\rangle - \varepsilon \left\langle \nabla w, \nabla \psi \right\rangle.$$
(25)

Then the *energy norm* of *u* is given by

$$||u||_{E}^{2} = |||(u, \varepsilon \Delta u)|||^{2} = \varepsilon^{2} ||\Delta u||_{0,\Omega}^{2} + b ||\nabla u||_{0,\Omega}^{2} + c ||u||_{0,\Omega}^{2} = |||(u, w)|||^{2}.$$

A regularity result for *w*, analogous to Proposition 11, is given below.

Corollary 12. Let $w = \varepsilon \Delta u$ and assume Proposition 11 holds. Then

$$w = w^S + \chi w^{BL} + w^R, \tag{26a}$$

where χ is the same cut-off function as in Proposition 11. Moreover, there exist constants K_3 , K_4 , β , $\gamma > 0$, independent of ε , such that

$$\left\| D^n w^{\mathcal{S}} \right\|_{0,\Omega} \lesssim |n|! K_3^{|n|} \quad \forall \ n \in \mathbb{N}_0^2,$$
(26b)

$$\left|\frac{\partial^{m+n} w^{BL}(\rho, \theta)}{\partial \rho^m \partial \theta^n}\right| \lesssim n! K_4^{m+n} \varepsilon^{-m} \mathrm{e}^{-\beta \rho/\varepsilon} \quad \forall \, m, n \in \mathbb{N}, \ (\rho, \theta) \in \overline{\Omega}_0,$$
(26c)

and

$$\left\|w^{R}\right\|_{1,\Omega} \lesssim e^{-\gamma/\varepsilon}.$$
(26d)

Finally, there exists a constant K > 0, depending only on the data, such that

$$\|D^{n}w\|_{0,\Omega} \lesssim K^{[n]} \max\left\{|n|^{[n]}, \varepsilon^{-[n]}\right\} \quad \forall \ n \in \mathbb{N}_{0}^{2}.$$
(26e)

Proof. Follows from Proposition 11 and the definition of $w = \varepsilon \Delta u$.

We close this section with the following result which will be used in the sequel.

Lemma 13. Let u^{BL} and w^{BL} satisfy (23c) and (26c), respectively. With Ω_0 given by (22), we have

$$\varepsilon^{-1} \left\| u^{BL} \right\|_{0,\Omega\setminus\Omega_0} + \left\| w^{BL} \right\|_{0,\Omega\setminus\Omega_0} \lesssim \varepsilon^{1/2} \mathrm{e}^{-\sigma/\varepsilon},$$

for some positive constant σ , independent of ε .

Proof. This follows from direct calculations.

3.1 Discretization by a mixed *rp*-FEM

In order to define our finite dimensional discretization spaces $V_1^N \subset H_0^1(\Omega)$ and $V_2^N \subset H^1(\Omega)$, we let $\Delta := {\Omega_i}_{i=1}^N$ be a mesh consisting of curvilinear quadrilaterals, subject to standard conditions (see, e.g. [12]) and associate with each Ω_i a bijective mapping $M_i: S_{ST} \to \overline{\Omega}_i$, where $S_{ST} = [0, 1]^2$ denotes the reference square. With $Q_p(S_{ST})$ the space of polynomials of degree p (in each variable) on S_{ST} , we define

$$S^{p}(\Delta) \coloneqq \left\{ u \in H^{1}(\Omega) : u|_{\Omega_{i}} \circ M_{i} \in Q_{p}(S_{ST}), \quad i = 1, \dots, N \right\},$$
$$S^{p}_{0}(\Delta) \coloneqq S^{p}(\Delta) \cap H^{1}_{0}(\Omega).$$

²The fact that $w \in H^2$ follows from the smoothness of $\partial \Omega$.

The mesh Δ is chosen following the construction in [3, 13]. We denote by Δ_A a *fixed* (asymptotic) mesh consisting of curvilinear quadrilateral elements Ω_i , $i = 1, ..., N_1$. These elements Ω_i are the images of the reference square S_{ST} under the element mappings $M_{A,i}$, $i = 1, ..., N_1 \in \mathbb{N}$. Moreover, the element mappings $M_{A,i}$ are assumed to be analytic (with analytic inverses). We also assume that the elements do not have a single vertex on the boundary $\partial \Omega$, but only complete, single edges. For convenience, we number the elements along the boundary first, i.e., Ω_i , $i = 1, ..., N_2 < N_1$ for some $N_2 \in \mathbb{N}$.

We now give the definition of the layer-adapted mesh from [13], almost verbatim.

Definition 14. Given parameters $\kappa > 0$, $p \in \mathbb{N}$, $\varepsilon \in (0, 1]$ and the (asymptotic) mesh Δ_A , the Spectral Boundary Layer Mesh Δ_{BL} is defined as follows:

- 1. If $\kappa p \varepsilon \ge 1/2$ then we are in the asymptotic range of p and we use the mesh Δ_A .
- 2. If $\kappa \rho \varepsilon < 1/2$, we need to define so-called needle elements. We do so by splitting the elements Ω_i , $i = 1, ..., N_2$ into two elements Ω_i^{need} and Ω_i^{reg} . To this end, split the reference square S_{ST} into two rectangles

$$S^{need} = [0, \kappa p \varepsilon] \times [0, 1], \quad S^{reg} = [\kappa p \varepsilon, 1] \times [0, 1],$$

and define the elements Ω_i^{need} and Ω_i^{reg} as the images of these two rectangles under the element map $M_{A,i}$ and the corresponding element maps as the concatination of the affine maps

$$\begin{aligned} A^{need} &: S_{ST} \to S^{need}, \quad (\xi, \eta) \to (\kappa \rho \varepsilon \xi, \eta), \\ A^{reg} &: S_{ST} \to S^{reg}, \quad (\xi, \eta) \to (\kappa \rho \varepsilon + (1 - \kappa \rho \varepsilon)\xi, \eta). \end{aligned}$$

with the element map $M_{A,i}$, i.e., $M_i^{need} = M_{A,i} \circ A^{need}$ and $M_i^{reg} = M_{A,i} \circ A^{reg}$.

In total, the mesh Δ_{BL} consists of $N = N_1 + N_2$ elements if $\kappa p \varepsilon < 1/2$. By construction, the resulting mesh

$$\Delta_{BL} = \{\Omega_1^{need}, \ldots, \Omega_{N_1}^{need}, \Omega_1^{reg}, \ldots, \Omega_{N_1}^{reg}, \Omega_{N_1+1}, \ldots, \Omega_N\},\$$

is a regular admissible mesh. In Figure 2, we show an example of such a mesh from [3].



Figure 2: Example of the SBL mesh.

Next, we take $V_1^N = S_0^p(\Delta_{BL})$ and $V_2^N = S^p(\Delta_{BL})$, and seek an approximation $(u_{p,\Delta}, w_{p,\Delta}) \in V_1^N \times V_2^N$ of (24) such that

$$\mathcal{A}_{\varepsilon}\left(\left(u_{p,\Delta}, w_{p,\Delta}\right)\right), (\psi, \phi)\right) = \langle f, \psi \rangle \quad \forall \ (\phi, \psi) \in V_1^N \times V_2^N.$$

$$(27)$$

Subtracting (27) from (24), we get

$$\mathcal{A}_{\varepsilon}\left((u-u_{p,\Delta}, w-w_{p,\Delta}), (\psi, \phi)\right) = 0 \quad \forall \ (\psi, \phi) \in V_1^N \times V_2^N$$

We have the following interpolation estimates, which were based on the Gauss-Lobbato interpolant from [12].

Lemma 15 ([3]). Let (u, w) be the solution to (24) and assume that (21) holds. Then there exist constants $\kappa_0, \kappa_1, \beta > 0$ independent of both $\varepsilon \in (0, 1]$ and of $p \in \mathbb{N}$, such that the following is true: For every p and every $\kappa \in (0, \kappa_0]$ with $\kappa p \ge \kappa_1$, there exist $I_p u \in S_0^p(\Delta_{BL})$ and $J_p w \in S^p(\Delta_{BL})$ such that

$$\left\|u-I_{p}u\right\|_{L^{\infty}(\Omega)}+\left\|\nabla(u-I_{p}u)\right\|_{L^{\infty}(\Omega)}+\left\|w-J_{p}w\right\|_{L^{\infty}(\Omega)}+\varepsilon^{1/2}\left\|\nabla(w-J_{p}w)\right\|_{0,\Omega}\lesssim e^{-\beta p}.$$

With

$$\Omega_{BL} = \bigcup_{i} \Omega_{i}^{need}, \quad \Omega_{REG} = \bigcup_{i} \Omega_{i}^{reg}$$

and $|\Omega_{BL}| = O(\kappa p \varepsilon)$, $|\Omega_{REG}| = O(1)$, we see that additionally there holds

$$\left\|\nabla(u - I_p u)\right\|_{0,\Omega_{BL}} + \left\|w - J_p w\right\|_{0,\Omega_{BL}} \lesssim \varepsilon^{1/2} \mathrm{e}^{-\beta p}$$
⁽²⁸⁾

The following was the main result of [3].

Theorem 16 ([3]). Let (u, w) and $(u_{p,\Delta}, w_{p,\Delta})$ be the solutions to (24) and (27), respectively. Assume Proposition 11 holds. Then there exists a positive constant β , independent of ε , such that

$$\left\| \left\| \left(u - u_{p,\Delta}, w - w_{p,\Delta} \right) \right\| \right\| \lesssim e^{-\beta p}.$$

Note that the norm used in Theorem 16 is not correctly balanced, because

$$\left|\left|\left|(u^{BL},\varepsilon\Delta u^{BL})\right|\right|\right| = O\left(\varepsilon^{1/2}\right), \text{ but } \left|\left|\left|(u^{S},\varepsilon\Delta u^{S})\right|\right|\right| = O(1), \ (\varepsilon \to 0).$$

Just like in the one-dimensional case, as $\varepsilon \to 0$, "the energy norm does not see the layers" [7, 9]. This is due to the fact that the weight on $||w||_{0,\Omega}$ is not the appropriate one. A more suitable, correctly balanced norm is

$$\||(u,w)||_{B}^{2} \coloneqq \varepsilon^{-1} \|w\|_{0,\Omega}^{2} + b \|\nabla u\|_{0,\Omega}^{2} + c \|u\|_{0,\Omega}^{2}, \qquad (29)$$

since then

$$\left\| \left(u^{BL}, \varepsilon \Delta u^{BL} \right) \right\|_{B} = O(1) = \left\| \left(u^{S}, \varepsilon \Delta u^{S} \right) \right\|_{B}, \quad (\varepsilon \to 0)$$

Numerical experiments using the above norm have been reported in [3], suggesting that indeed the method yields uniform exponential convergence in the balanced norm $\|\|\cdot\|\|_B$. Unfortunately, the bilinear form $\mathcal{R}_{\varepsilon}$ given by (25), is not uniformly coercive with respect to the norm (29). To circumvent this obstacle, we will use the methodology derived in Section 2.

3.2 Balanced norm estimates

Recall the decomposition (26), set $w^{SR} := w^S + w^R$ and define the L^2 projection of w^{SR} on the regular region Ω_{REG} , denoted by $\pi_p^1 w^{SR} \in V_2^N|_{\Omega_{REG}}$, via

$$\left\langle \pi_p^1 w^{SR}, v \right\rangle_{\Omega_{REG}} = 0 \quad \forall \ v \in V_2^N |_{\Omega_{REG}}$$

The projection is uniquely defined and consequently, we have the following:

$$\left\| \pi_{p}^{1} w^{SR} - w^{S} \right\|_{0,\Omega_{REG}} + \left\| \pi_{p}^{1} w^{SR} - w^{SR} \right\|_{L^{\infty}(\Omega_{REG})} \lesssim e^{-\kappa p}$$
(30)

where the definition of the projection π_p^1 and Lemma 13 were used, along with $\kappa p\varepsilon < 1/2$, which is when we use the SBL mesh.

Let analogously $u^{SR} := u^S + u^R$ and define $\pi_p^2 u^{SR} \in V_2^N|_{\Omega_{REG}}$ as the weighted H^1 projection of u^{SR} on Ω_{REG} :

$$b\left\langle \nabla(\pi_p^2 u^{SR} - u^{SR}), v\right\rangle_{\Omega_{REG}} + c\left\langle \pi_p^2 u^{SR} - u^{SR}, v\right\rangle_{\Omega_{REG}} = 0 \quad \forall \ v \in V_2^N|_{\Omega_{REG}},$$

and we have

$$\left\|\pi_p^2 u^{SR} - u^{SR}\right\|_{1,\Omega_{REG}} + \left\|\pi_p^2 \nabla (u^{SR} - u^{SR})\right\|_{L^{\infty}(\Omega_{REG})} \lesssim e^{-\beta p}.$$

We next define an auxiliary function $\chi_2 \in \mathcal{P}_1$ by the conditions

$$\chi_2(0) = 0, \quad \chi_2(\kappa p \varepsilon) = 1.$$

A direct calculation establishes the following bounds on χ_2 :

$$\|\chi_2\|_{0,\Omega_{BL}} \lesssim (\kappa p\varepsilon)^{1/2}, \quad \|\chi_2'\|_{0,\Omega_{BL}} \lesssim (\kappa p\varepsilon)^{-1/2}. \tag{31}$$

We are now in a position to define *special representatives* $\tilde{w} \in S^p$ of w and $\tilde{u} \in S_0^p$ of u. Recalling the decomposition (26a), define

$$\tilde{w}(x) \coloneqq \begin{cases} J_p w - \chi_2(\rho) \left(w^{SR} + w^{BL} - \pi_p^1 w^{SR} \right) |_{\partial \Omega_{REG}} & \text{in } \Omega_{BL}, \\ \pi_p^1 w^{SR} & \text{in } \Omega_{REG}. \end{cases}$$
(32a)

$$\tilde{u}(x) \coloneqq \begin{cases} I_p u - \chi_2(\rho) \left(u^{SR} + u^{BL} - \pi_p^2 u^{SR} \right) |_{\partial \Omega_{REG}} & \text{ in } \Omega_{BL}, \\ \pi_p^2 u^{SR} & \text{ in } \Omega_{REG}. \end{cases}$$
(32b)

We have the following.

Lemma 17. There exist a positive contant β such that the following bounds holds true:

$$\|w - \tilde{w}\|_{0,\Omega_{BL}} \lesssim \varepsilon^{1/2} \mathrm{e}^{-\beta p},\tag{33a}$$

$$\|w - \tilde{w}\|_{0,\Omega_{REG}} \lesssim e^{-\beta p},\tag{33b}$$

$$\|u - \tilde{u}\|_{0,\Omega_{BL}} \lesssim \varepsilon^{1/2} \mathrm{e}^{-\beta p},\tag{33c}$$

$$\|u - \tilde{u}\|_{0,\Omega_{REG}} \lesssim e^{-\beta p}.$$
(33d)

Proof. We study the two regions Ω_{BL} and Ω_{REG} separately.

 Ω_{BL} : We have for $w - \tilde{w}$,

$$\|w - \tilde{w}\|_{0,\Omega_{BL}} \le \|w - J_p w\|_{0,\Omega_{BL}} + \|\chi_2\|_{0,\Omega_{BL}} \left\{ \|w^{BL}\|_{L^{\infty}(\partial\Omega_{REG})} + \|\pi_p^1 w^{SR} - w^{SR}\|_{L^{\infty}(\Omega_{BL})} \right\}$$

Equation (28) takes care of the first term above, and equations (23c), (30) and (31), allow us to bound the remaining terms, as desired. This shows (33a). Similarly, for $u - \tilde{u}$ we have

$$\|u-\tilde{u}\|_{0,\Omega_{BL}} \leq \left\|u-I_pu\right\|_{0,\Omega_{BL}} + \|\chi_2\|_{0,\Omega_{BL}} \left\{ \|u^{BL}\|_{L^\infty(\partial\Omega_{REG})} + \left\|\pi_p^2 u^{SR} - u^{SR}\right\|_{L^\infty(\Omega_{BL})} \right\}.$$

Eq. (33c) follows from the same equations.

 Ω_{REG} : We have $w = w^{BL} + w^{SR}$. Therefore,

$$\|\boldsymbol{w}-\tilde{\boldsymbol{w}}\|_{0,\Omega_{REG}} \leq \left\|\boldsymbol{w}^{BL}\right\|_{0,\Omega_{REG}} + \left\|\boldsymbol{w}^{SR}-\pi_p^1\boldsymbol{w}^{SR}\right\|_{0,\Omega_{REG}}.$$

Lemma 13 and (30) then give (33b). Eq. (33d) follows in the same fashion.

We now have the necessary tools for proving the following.

Theorem 18. Let (u, w) be the solution of (24) and let $(u_{p,\Delta}, w_{p,\Delta})$ be the solution of (27), based on the Spectral Boundary Layer mesh of Definition 14. Then, there exists a positive constant β , independent of ε , such that

$$\left|\left|\left|(u-u_{p,\Delta},w-w_{p,\Delta})\right|\right|\right|_{B} \lesssim e^{-\beta p}.$$

Proof. Let \tilde{w} , \tilde{u} be defined by (32a) and (32b), respectively. We have from (29) and the triangle inequality,

$$\begin{split} \left| \left| \left| \left(u - u_{p,\Delta}, w - w_{p,\Delta} \right) \right| \right|_{B} &\leq \left| \left| \left(u - \tilde{u}, w - \tilde{w} \right) \right| \right|_{B} + \left| \left| \left| \left(\tilde{u} - u_{p,\Delta}, \tilde{w} - w_{p,\Delta} \right) \right| \right|_{B} \right|_{B} \\ &\lesssim \varepsilon^{-1/2} \left\| w - \tilde{w} \right\|_{0} + \left\| u - \tilde{u} \right\|_{1} + \varepsilon^{-1/2} \left\| \tilde{w} - w_{p,\Delta} \right\|_{0} + \left\| \tilde{u} - u_{p,\Delta} \right\|_{1} . \end{split}$$

We only need to treat the terms $\varepsilon^{-1/2} \| \tilde{w} - w_{p,\Delta} \|_0$ and $\| \tilde{u} - u_{p,\Delta} \|_1$ since the rest can be handled by Theorem 16 and Lemma 17.

Let $\eta \coloneqq w_{p,\Delta} - \tilde{w}$ and $\xi \coloneqq u_{p,\Delta} - \tilde{u}$. We have by Galerkin orthogonality,

$$|||(\eta,\xi)||| = \langle w - \tilde{w}, \eta \rangle + b \langle \nabla(u - \tilde{u}), \nabla\xi \rangle + c \langle u - \tilde{u}, \xi \rangle.$$
(34)

We will consider the two regions separately.

 Ω_{BL} : We first note that

$$\|\chi_2\|_{0,\Omega_{\mathrm{BL}}}\left\{\left\|w^{BL}\right\|_{L_{\infty}(\partial\Omega_{REG})}+\left\|\pi_p^1w^{SR}-w^{SR}\right\|_{L_{\infty}(\partial\Omega_{REG})}\right\}\lesssim\varepsilon^{1/2}\mathrm{e}^{-\kappa p}.$$

Then, we combine the above with (28), and use the Cauchy-Schwarz inequality, along with the defintion of \hat{w} , to obtain

$$\langle w - \hat{w}, \eta \rangle_{\Omega_{BL}} \lesssim \varepsilon^{1/2} \mathrm{e}^{-\beta p} \|\eta\|_{\Omega_{BL}}.$$

Similarly, we first note that

$$\left|\chi_{2}'\right|_{0,\Omega_{\mathrm{BL}}}\left\{\left\|u^{BL}\right\|_{L_{\infty}(\partial\Omega_{REG}}+\left\|\pi_{p}^{2}u^{SR}-u^{SR}\right\|_{L_{\infty}(\partial\Omega_{REG}}\right\}\lesssim\varepsilon^{1/2}\mathrm{e}^{-\kappa p},$$

hence, as above, we have

$$b \langle \nabla(u-\tilde{u}), \nabla \xi \rangle_{\Omega_{BL}} + c \langle u-\tilde{u}, \xi \rangle_{\Omega_{BL}} \lesssim \varepsilon^{1/2} \mathrm{e}^{-\beta p} \left\| \xi \right\|_{1,\Omega_{BL}}.$$

 Ω_{REG} : We have by the definition of \hat{w} ,

$$\langle w - \hat{w}, \eta \rangle_{\Omega_{REG}} = \left\langle w_{BL} + w^{SR} - \pi_p w^{SR}, \eta \right\rangle_{\Omega_{REG}} = \left\langle w^{BL}, \eta \right\rangle_{\Omega_{REG}} \lesssim \varepsilon^{1/2} \mathrm{e}^{-\kappa p} \|\eta\|_{0,\Omega_{REG}}$$

Similarly,

$$\begin{split} b \left\langle \nabla(u - \tilde{u}), \nabla \xi \right\rangle_{\Omega_{REG}} &+ c \left\langle u - \tilde{u}, \xi \right\rangle_{\Omega_{REG}} \\ &= b \left\langle \nabla u^{BL}, \nabla \xi \right\rangle_{\Omega_{REG}} + c \left\langle u^{BL}, \xi \right\rangle_{\Omega_{REG}} \lesssim \left\| u^{BL} \right\|_{1,\Omega_{BL}} \|\xi\|_{1,\Omega_{BL}} \lesssim \varepsilon^{1/2} \mathrm{e}^{-\beta p} \|\xi\|_{1,\Omega_{BL}} \,. \end{split}$$

Combining the above with (34), we see that

$$\|\|(\eta,\xi)\|\|^2 \leq \varepsilon^{1/2} e^{-\beta p} \|\|(\eta,\xi)\|\|$$

from which we have

$$\varepsilon^{-1/2} \left\| w_{p,\Delta} - \tilde{w} \right\|_{0,\Omega} + \left\| u_{p,\Delta} - \tilde{u} \right\|_{1,\Omega} \lesssim \mathrm{e}^{-\beta p}.$$

This completes the proof.

4 Conclusions

We presented balanced norm estimates for singularly perturbed fourth order problems in one-dimension as well as in smooth (analytic) domains in \mathbb{R}^2 . As a corollary, we also got maximum norm estimates in the one-dimensional case. The methodology is not restricted to conforming methods, and in fact we are in the process of studying balanced norm estimates for non-conforming *rp*-version methods.

Finally, we mention that this contribution proves the conjectures made in [2], [3] based on numerical evidence.

References

- L. Beirão da Veiga, A. Buffa, J. Rivas and G. Sangalli, Some estimates for h-p-k-refinement in isogeometric analysis, Numer. Math. 118 (2011), 271–305.
- [2] P. Constantinou, *The hp Finite Element Method for fourth order singularly perturbed problems*, Doctoral Dissertation, Department of Mathematics and Statistics, University of Cyprus, 2019.
- [3] P. Constantinou, S. Franz, L. Ludwig and C. Xenophontos, *A mixed hp finite element method for a 4th order singularly perturbed problem*, Num. Meth. PDEs, **35** (2019) 114–127.
- [4] P. Constantinou, C. Varnava and C. Xenophontos, *An hp finite element method for fourth order sin*gularly perturbed problems, Numer. Alg., **73** (2016) 567–590.
- [5] P. Constantinou and C. Xenophontos, *An hp finite element method for a 4th order singularly perturbed boundary value problem in two dimensions*, Comp. Math. Appl., **72** (2017) ,1565–1575.
- [6] D. A. Di Pietro and A. Ern, *Mathematical Aspects of Discontinuous Galerkin Methods*, Mathematique et Applications 69, Springer-Verlag, 2012.
- [7] S. Franz and H.-G. Roos, Robust error estimation in energy and balanced norms for singularly perturbed fourth order problems, Comp. Math. Appl., 72 (2016) 233–247.
- [8] S. Franz and H.-G. Roos, On robust error estimation for singularly perturbed fourth-order problems, In Boundary and Interior Layers, Computational and Asymptotic Methods BAIL 2016, Lecture Notes in Computational Science and Engineering 120 (2017), 77–85, Springer, Berlin Heidelberg,
- [9] S. Franz and H.-G. Roos, *Robust error estimation in energy and balanced norms for singularly perturbed fourth order problems*, Model. Anal. Inform. Sist., **23** (2016), 364–369.
- [10] R. Lin and M. Stynes, A balanced finite element method for singularly perturbed reaction-diffusion problems, SIAM J. Numer. Anal., 50 (2012) 2729–2743.
- [11] T. Linß, Layer-adapted meshes for reaction-convection-diffusion problems, Lecture Notes in Mathematics 1985, Springer-Verlag, 2010.
- [12] J. M. Melenk, *hp-Finite Element Methods for Singular Perturbations*, Vol. 1796 of Springer Lecture Notes in Mathematics, Springer Verlag, 2002.
- [13] J. M. Melenk and C. Xenophontos, Robust exponential convergence of hp-FEM in balanced norms for singularly perturbed reaction-diffusion equations, Calcolo, 53 (2016) 105–132 (2016).
- [14] J. M. Melenk and C. Xenophontos, *Analytic regularity results for a fourth singularly perturbed problem in smooth domains*, in preparation (2023).
- [15] J. M. Melenk, C. Xenophontos and L. Oberbroeckling, Robust exponential convergence of hp-FEM for singularly perturbed systems of reaction-diffusion equations with multiple scales, IMA J. Num. Anal., 33 (2013) 609–628.
- [16] R. E. O'Malley, Introduction to Singular Perturbations, Academic Press, New York, 1974.

- [17] P. Panaseti, A. Zouvani, N. Madden and C. Xenophontos, A C¹-conforming hp finite element method for fourth order singularly perturbed boundary value problems, Appl. Num. Math., **104** (2016) 81– 97.
- [18] C. Schwab, p- and hp-Finite Element Methods, Oxford University Press, 1998.

A Auxiliary results

Lemma 19 (Inverse inequality). For any polynomial $q \in \mathcal{P}_p$, $p \in \mathbb{N}$, and any domain D there holds

$$|q|_{k,D} \le \left(\frac{p!}{(p-k)!}\right)^2 |D|^{-k} ||q||_{0,D}, k = 0, 1, \dots, p.$$

Proof. Theorem 4.76 in [18] gives the cases k = 0 and 1. Iterating on that result, we obtain the desired inequality.

B Proof of (12) **and** (13)

We begin with the following results:

Lemma 20. Let $v \in C^{\infty}(I_{ST})$, $I_{ST} = (-1, 1)$ satisfy

$$\left\|v^{(n)}\right\|_{\infty,I_{ST}} \leq C_{v}(\gamma_{v}h)^{n}K^{n-1}, \ \forall \ n \in \mathbb{N},$$

for some constants $C_v, \gamma_v > 0, K \ge 1$, and for $h \in (0, 1]$. Then there exists an approximation $I_p v \in \mathcal{P}_p$ with $I_p^{(k)}v(\pm 1) = v^{(k)}(\pm 1), k = 0, 1$ such that under the condition

$$\frac{hK}{p} \le \eta$$

for some $\eta > 0$, there holds

$$\left\|v-\mathcal{I}_p v\right\|_{2,I_{ST}} \lesssim C_v K^{1/2} h \mathrm{e}^{-\beta p}.$$

Proof. There exists (see, e.g. [17]) $I_p v \in \mathcal{P}_p$ with $(I_p v)^{(k)}(\pm 1) = v^{(k)}(\pm 1), k = 0, 1$ and

$$\left\|v - \mathcal{I}_p v\right\|_{2, I_{ST}}^2 \leq \frac{(p-s)!}{(p+s)!} \left\|v^{(s+1)}\right\|_{0, I_{ST}}^2, \ \forall \ s \in (0,p).$$

Choose $s = \lambda p$, with $\lambda \in (0, 1]$ to be selected shortly. Then we calculate

$$\frac{(p-\lambda p)!}{(p+\lambda p)!} \left\| v^{(\lambda p+1)} \right\|_{0,I_{ST}}^2 \leq \left[\frac{(1-\lambda)^{1-\lambda}}{(1+\lambda)^{1+\lambda}} \right]^p p^{-2\lambda p} \mathrm{e}^{2\lambda p+1} C_{\nu}^2 (h\gamma_{\nu})^{2(\lambda p+1)} K^{2\lambda p+1} \leq \mathrm{e} C_{\nu}^2 h^2 \gamma_{\nu}^2 K \left(\frac{\mathrm{e} \gamma_{\nu} h K}{p} \right)^{2\lambda p} q^p,$$

with $q = (1 - \lambda)^{1-\lambda}/(1 + \lambda)^{1+\lambda} \in (0, 1)$. Using the hypothesis $hK/p \le \eta$ and selecting $\lambda \in (0, 1]$, $\eta > 0$ such that $(\gamma_v e \eta)^{2\lambda p} \le 1$, gives

$$\frac{(p-\lambda p)!}{(p+\lambda p)!} \left\| v^{(\lambda p+1)} \right\|_{0,I_{ST}}^2 \lesssim C_v^2 h^2 K \gamma_v q^p,$$

from which the desired result follows.

By means of affine transformations the interpolation operator I_p of Lemma 20 can be applied to all subintervals of our mesh Δ to give a global piecewise polynomial interpolant. To this end introduce the affine transformations

$$Q_j: I_{ST} \to I_j: t \mapsto Q_j(t) := \frac{1-t}{2} x_{j-1} + \frac{1+t}{2} x_j, \ j = 1, \dots, N,$$

and define

$$\left(\mathcal{I}_{p,\Delta}v\right)\Big|_{I_j} \coloneqq \mathcal{I}_p\left(v \circ Q_j^{-1}\right), \quad j = 1, \dots, N,$$
(35)

As there is no confusion, we drop the Δ from the notation.

Corollary 21. Let $j \in \{1, ..., N\}$ and assume $v \in C^{\infty}(\overline{I}_i)$ satisfies

$$\left\|v^{(n)}\right\|_{\infty,I} \leq C_{v}(h_{j}\gamma_{v})^{n}K^{n-1}, \quad \forall \ n \in \mathbb{N},$$

for some constants $C_{\nu}, \gamma_{\nu} > 0$, $K \ge 1$. Then there exist constants $\eta, \beta > 0$ depending only on γ_{ν} such that under the condition

$$\frac{h_j K}{p} \le \eta$$

the polynomial approximation I_{pv} defined in (35) satisfies

$$\left\| \left(v - I_p v \right)^{(k)} \right\|_{0, I_j} \lesssim C_v K^{1/2} h_j^{3/2 - k} \mathrm{e}^{-\beta p}, \ k = 0, 1, 2.$$

Proof. Set $\hat{v} = v \circ Q_j$. Then

$$\left\|\hat{v}^{(n)}\right\|_{0,I_{ST}} \leq C_{v}(h_{j}/2)^{n}\gamma_{v}^{n}K^{n-1},$$

and by Lemma 20 we have

$$\left\|\hat{v}-\mathcal{I}_p\hat{v}\right\|_{2,I_{ST}} \lesssim C_v K^{1/2} h_j \mathrm{e}^{-\beta p}.$$

Transforming back to I_j gives the result.

For our spectral boundary layer mesh, the previous results give

$$\left\| \left(u - I_p u \right)^{(k)} \right\|_{0,\Omega_{BL}} \lesssim C_u \varepsilon^{-1/2} h_j^{3/2-k} \mathrm{e}^{-\beta p}, \ k = 0, 1, 2,$$

or equivalently, since $h_j = \kappa p \varepsilon$,

$$(\kappa p\varepsilon)^{-1} \left\| u - \mathcal{I}_p u \right\|_{0,\Omega_{BL}} + \left| u - \mathcal{I}_p u \right|_{1,\Omega_{BL}} + \kappa p\varepsilon \left| u - \mathcal{I}_p u \right|_{2,\Omega_{BL}} \lesssim C_u (\kappa p)^{1/2} e^{-\beta p}.$$
(36)

Now, since $(u - \mathcal{I}_p u)^{(k)}(0) = 0, k = 0, 1$, we may write

$$\left|\left(u-\mathcal{I}_{p}u\right)^{(k)}(x)\right|=\left|\int_{0}^{x}\left(u-\mathcal{I}_{p}u\right)^{(k+1)}(t)\,\mathrm{d}t\right|,\ x\in\Omega_{BL}.$$

Then, by the Cauchy-Schwarz inequality and (36) we get for k = 0, 1,

$$\left| \left(u - \mathcal{I}_p u \right)^{(k)} (x) \right| \le (\kappa p \varepsilon)^{1/2} \left| \left(u - \mathcal{I}_p u \right)^{(k)} \right|_{1,\Omega_{BL}} \lesssim \begin{cases} C_u \varepsilon^{-1/2} (\kappa p \varepsilon)^2 e^{-\beta p} , & k = 0, \\ C_u \varepsilon^{-1/2} \kappa p \varepsilon e^{-\beta p} , & k = 1. \end{cases}$$

Thus

$$(\kappa p\varepsilon)^{-3/2} \left\| u - \mathcal{I}_p u \right\|_{\infty,\Omega_{BL}} + (\kappa p\varepsilon)^{-1/2} \left\| \left(u - \mathcal{I}_p u \right)' \right\|_{\infty,\Omega_{BL}} \lesssim C_u(\kappa p)^{1/2} \mathrm{e}^{-\beta p}.$$