

# A Golub-Welsch version for simultaneous Gaussian quadrature

Walter Van Assche\*  
KU Leuven, Belgium

## Abstract

The zeros of type II multiple orthogonal polynomials can be used for quadrature formulas that approximate  $r$  integrals of the same function  $f$  with respect to  $r$  measures  $\mu_1, \dots, \mu_r$  in the spirit of Gaussian quadrature. This was first suggested by Borges in 1994, even though he does not mention multiple orthogonality. We give a method to compute the quadrature nodes and the quadrature weights which extends the Golub-Welsch approach using the eigenvalues and left and right eigenvectors of a banded Hessenberg matrix. This method was already described by Coussement and Van Assche in 2005 but it seems to have gone unnoticed. We describe the result in detail for  $r = 2$  and give some examples.

**Keywords:** Simultaneous Gauss quadrature, multiple orthogonal polynomials, banded Hessenberg matrix.

**MSC Classification:** primary 41A55, 65D32; secondary 15A18, 33C45, 41A21, 42C05

## 1 Introduction

Suppose a real function  $f$  is given and one wants to compute the integrals

$$\int_{\mathbb{R}} f(x) d\mu_j(x), \quad 1 \leq j \leq r$$

for  $r$  positive measures  $\mu_1, \dots, \mu_r$  with only  $N$  function evaluations. In 1994 Borges [4] suggested Gauss-like quadrature rules and showed that one gets an optimal set of quadrature rules if one takes the  $N$  quadrature nodes at the zeros of a polynomial of degree  $N$  which satisfies orthogonality conditions with respect to the measures  $\mu_1, \dots, \mu_r$ . Borges uses the terminology quasi-orthogonality but such a polynomial is now known as a type II multiple orthogonal polynomial. The quadrature rules are of the form

$$\sum_{k=1}^N \lambda_{k,N}^{(j)} f(x_{k,N}) \approx \int_{\mathbb{R}} f(x) d\mu_j(x),$$

so that the integral for the measure  $\mu_j$  corresponds to quadrature weights  $\lambda_{k,N}^{(j)}$ . An important numerical aspect is then how to compute the quadrature nodes  $\{x_{k,N}, 1 \leq k \leq N\}$ .

---

\*Supported by research project G0C9819N of FWO (Research Foundation – Flanders).

$N\}$  and the quadrature weights  $\{\lambda_{k,n}^{(j)}, 1 \leq k \leq N\}$  for  $1 \leq j \leq r$  in an efficient and stable way. This problem was suggested in the survey on classical multiple orthogonal polynomials [25, §4.3]. The computation of the quadrature nodes and weights was first done by Milovanović and Stanić for almost diagonal multi-indices [20] and later this was extended for general multi-indices for the case of two or three quadrature rules by Tomović and Stanić [23] and for arbitrary  $r$  by Jovanović, Stanić and Tomović [15]. The common nodes were calculated as eigenvalues of certain matrices, while the quadrature weights were calculated by solving certain systems of linear equations. Simultaneous Gaussian quadrature for multiple orthogonal polynomials for two measures  $\mu_1$  and  $\mu_2$  supported on two intervals  $\Delta_1$  and  $\Delta_2$  which are disjoint (a so-called Angelesco system) was investigated in [18], and for multiple Hermite polynomials (with  $r = 3$ ) in [27] but mainly from an analytical point of view, leaving out the numerical aspects.

In 1969 Golub and Welsch [12] showed that for Gaussian quadrature (the case when  $r = 1$ ) one can use the Jacobi matrix for the corresponding orthogonal polynomials. The eigenvalues of the  $N \times N$  Jacobi matrix are the quadrature nodes and the first components of the normalized eigenvectors are related to the quadrature weights. This approach was extended to simultaneous Gaussian quadrature (in the sense of Borges) by Coussement and Van Assche [6] but this extension seems to have gone unnoticed. The role of the Jacobi matrix is now played by a banded Hessenberg matrix for which the eigenvalues are the required quadrature nodes, and one needs the right and the left eigenvectors to find the quadrature weights. The goal of this paper is to give a simplified proof of this result for two measures  $\mu_1$  and  $\mu_2$ . We will give some examples involving measures  $d\mu_1 = w_1(x) dx$  and  $d\mu_2(x) = w_2(x) dx$  for which the weight functions are related to modified Bessel functions.

The structure of the paper is as follows. In Section 2 we will give the necessary background about multiple orthogonal polynomials and their relation with Hermite-Padé approximation. In Section 3 we explain what we mean by simultaneous Gaussian quadrature. Multiple orthogonal polynomials satisfy various recurrence relations which we give in Section 4, and in particular the multiple orthogonal polynomials on the stepline (near the diagonal) satisfy a linear recurrence relation of order  $r + 1$ , which gives rise to a banded Hessenberg matrix with one diagonal above the main diagonal and  $r$  diagonals below the main diagonal. The eigenvalues of this Hessenberg matrix correspond to the zeros of the type II multiple orthogonal polynomial. The main theorem (Theorem 2) will be proved in Section 5. We will use the fact that the quadrature weights can be expressed in terms of the Christoffel-Darboux kernel for multiple orthogonal polynomials. One important difference is that the Christoffel-Darboux kernel for multiple orthogonal polynomials contains both the type I and the type II multiple orthogonal polynomials, and hence is not symmetric in the two variables. Two numerical examples involving modified Bessel function weights are worked out in Section 6.

## 2 Multiple orthogonal polynomials

Multiple orthogonal polynomials are polynomials in one variable with orthogonality relations with respect to  $r$  measures  $(\mu_1, \mu_2, \dots, \mu_r)$  (on the real line in this paper). We use a multi-index  $\vec{n} = (n_1, n_2, \dots, n_r) \in \mathbb{N}^r$  of size  $|\vec{n}| = n_1 + n_2 + \dots + n_r$ .

**Definition 1** (type II MOP). *A type II multiple orthogonal polynomial  $P_{\vec{n}}$  is a polynomial*

of degree  $\leq |\vec{n}|$  for which

$$\begin{aligned} \int_{\mathbb{R}} P_{\vec{n}}(x)x^k d\mu_1(x) &= 0, & 0 \leq k \leq n_1 - 1, \\ &\vdots \\ \int_{\mathbb{R}} P_{\vec{n}}(x)x^k d\mu_r(x) &= 0, & 0 \leq k \leq n_r - 1. \end{aligned}$$

**Definition 2** (type I MOP). *Type I multiple orthogonal polynomials are in a vector  $(A_{\vec{n},1}, \dots, A_{\vec{n},r})$ , with  $\deg A_{\vec{n},j} \leq n_j - 1$ , and they satisfy*

$$\sum_{j=1}^r \int_{\mathbb{R}} A_{\vec{n},j}(x)x^k d\mu_j(x) = 0, \quad 0 \leq k \leq |\vec{n}| - 2.$$

The orthogonality conditions for the type II multiple orthogonal polynomial give a homogeneous system of  $|\vec{n}|$  linear equations for the  $|\vec{n}| + 1$  unknown coefficients of the polynomial  $P_{\vec{n}}$ . In a similar way, the orthogonality conditions for the type I multiple orthogonal polynomials give a homogeneous linear system of  $|\vec{n}| - 1$  equations for the  $|\vec{n}|$  unknown coefficients of the polynomials  $A_{\vec{n},j}$  ( $1 \leq j \leq r$ ). When the type I and type II multiple orthogonal polynomials are unique up to a multiplicative factor, then we say that the multi-index  $\vec{n}$  is normal. When this happens, the type II polynomial  $P_{\vec{n}}$  has degree  $|\vec{n}|$  and we will take the leading coefficient to be one. For a normal multi-index we normalize the type I polynomials by setting

$$\sum_{j=1}^r \int_{\mathbb{R}} x^{|\vec{n}|-1} A_{\vec{n},j}(x) d\mu_j(x) = 1. \quad (2.1)$$

In this paper we assume this is the case for all multi-indices  $\vec{n}$ , in which case we deal with a perfect system. Hence our type II multiple orthogonal polynomials are monic and the type I multiple orthogonal polynomials satisfy (2.1).

Multiple orthogonal polynomials appear in simultaneous rational approximation of  $r$  functions  $(f_1, f_2, \dots, f_r)$ , where

$$f_j(z) = \int_{\mathbb{R}} \frac{d\mu_j(x)}{z-x}, \quad z \in \mathbb{C} \setminus \mathbb{R}.$$

**Problem 1** (Type II Hermite-Padé). *Find a polynomial  $P_{\vec{n}}$  of degree  $\leq |\vec{n}|$  and polynomials  $Q_{\vec{n},j}$  such that for  $1 \leq j \leq r$*

$$P_{\vec{n}}(z)f_j(z) - Q_{\vec{n},j}(z) = \mathcal{O}\left(\frac{1}{z^{n_j+1}}\right), \quad z \rightarrow \infty.$$

The common denominator  $P_{\vec{n}}$  is the type II multiple orthogonal polynomial and the numerator polynomials are given by

$$Q_{\vec{n},j}(z) = \int_{\mathbb{R}} \frac{P_{\vec{n}}(z) - P_{\vec{n}}(x)}{z-x} d\mu_j(x).$$

The error can also be expressed in terms of the type II multiple orthogonal polynomial:

$$P_{\vec{n}}(z)f_j(z) - Q_{\vec{n},j}(z) = \int_{\mathbb{R}} \frac{P_{\vec{n}}(x)}{z-x} d\mu_j(x).$$

**Problem 2** (Type I Hermite-Padé). *Find polynomials  $A_{\vec{n},j}$  of degree  $\leq n_j - 1$  and a polynomial  $B_{\vec{n}}$  such that*

$$\sum_{j=1}^r A_{\vec{n},j}(z) f_j(z) - B_{\vec{n}}(z) = \mathcal{O}\left(\frac{1}{z^{|\vec{n}|}}\right), \quad z \rightarrow \infty.$$

The polynomials  $A_{\vec{n},1}, \dots, A_{\vec{n},r}$  are the type I multiple orthogonal polynomials and the polynomial  $B_{\vec{n}}$  is

$$B_{\vec{n}}(z) = \sum_{j=1}^r \int_{\mathbb{R}} \frac{A_{\vec{n},j}(z) - A_{\vec{n},j}(x)}{z - x} d\mu_j(x).$$

The error is given by

$$\sum_{j=1}^r A_{\vec{n},j}(z) f_j(z) - B_{\vec{n}}(z) = \sum_{j=1}^r \int_{\mathbb{R}} \frac{A_{\vec{n},j}(x)}{z - x} d\mu_j(x).$$

Here is some history indicating important progress (but very incomplete). Hermite-Padé approximation was introduced by Charles Hermite in 1873, who used them in his proof that  $e$  is transcendental [14]. The case  $r = 1$  was then investigated in more detail by Hermite's student Henri Padé (Padé approximation, Padé table) [22]. One important system for which all the multi-indices are normal (a perfect system) was described by Aurel Angelescu [2] who used  $r$  measures (weights) supported on  $r$  disjoint intervals. This is nowadays known as an Angelesco system. In 1934 Kurt Mahler [19] investigated perfect systems but his work was published much later 1968. Another important system of measures was introduced by Evgeni Nikishin in 1979 [21]. For  $r = 2$  the measures  $(\mu_1, \mu_2)$  are absolutely continuous on an interval  $\Delta_0$  with weight functions  $(w_1, w_2)$  for which

$$\frac{w_2(x)}{w_1(x)} = \int_{\Delta_1} \frac{d\sigma(t)}{x - t},$$

where  $\sigma$  is a positive measure on an interval  $\Delta_1$  which is disjoint from  $\Delta_0$ . Such a system (and its recursive extensions to  $r > 2$ ) are now known as Nikishin systems and it has been shown that these are perfect systems [9]. Gonchar and Rakhmanov studied the convergence of Hermite-Padé approximants in [13] for which they used an extremal problem in logarithmic potential theory for a vector of measures. The idea of simultaneous Gaussian quadrature was introduced by Carlos F. Borges in [4]. Van Assche, Geronimo and Kuijlaars formulated a Riemann-Hilbert problem for multiple orthogonal polynomials in 2001 [26]. This allows to obtain the asymptotic behavior of multiple orthogonal polynomials.

Nowadays multiple orthogonal polynomials are used in various applications, such as eigenvalues of products of random matrices, non-intersecting Brownian motions, combinatorial problems related to random tilings, rational approximations in number theory, etc. They are also used in numerical analysis for matrix function evaluation (Alqahtani and Reichel [1]).

## 2.1 Examples extending classical orthogonal polynomials

Many examples of multiple orthogonal polynomials are known.

- Multiple Hermite polynomials: the weights are normal densities with non-zero mean

$$w_j(x) = e^{-x^2+c_jx}, \quad c_i \neq c_j, \quad x \in (-\infty, \infty).$$

- Multiple Laguerre polynomials: the weights

$$w_j(x) = x^{\alpha_j} e^{-x}, \quad \alpha_i - \alpha_j \notin \mathbb{Z}, \quad \alpha_j > -1, \quad x \in [0, \infty)$$

correspond to multiple Laguerre polynomials of the first kind. The weights

$$w_j(x) = x^\alpha e^{-c_jx}, \quad c_i \neq c_j, \quad c_j > 0, \quad \alpha > -1, \quad x \in [0, \infty),$$

correspond to multiple Laguerre polynomials of the second kind.

- Jacobi-Piñero polynomials: the weights are

$$w_j(x) = x^{\alpha_j}(1-x)^\beta, \quad \alpha_i - \alpha_j \notin \mathbb{Z}, \quad \alpha_j, \beta > -1, \quad x \in [0, 1].$$

For these extensions of the classical orthogonal polynomials one knows a differential equation for the multiple orthogonal polynomials, a system of nearest neighbor recurrence relations, the asymptotic distribution of the zeros, etc. Other examples include weights in terms of special functions satisfying a second order differential equation.

## 2.2 Modified Bessel functions $(K_\nu, K_{\nu+1})$

The modified Bessel function  $K_\nu$  is given as

$$K_\nu(x) = \frac{1}{2} \left(\frac{x}{2}\right)^\nu \int_0^\infty \exp\left(-t - \frac{x^2}{4t}\right) t^{-\nu-1} dt.$$

It satisfies the modified Bessel equation

$$x^2 y'' + xy' - (x^2 + \nu^2)y = 0.$$

Introduce the functions

$$\rho_\nu(x) = 2x^{\nu/2} K_\nu(2\sqrt{x}), \quad x > 0,$$

which have the moments

$$\int_0^\infty x^n \rho_\nu(x) dx = \Gamma(n + \nu + 1) \Gamma(n + 1).$$

Van Assche and Yakubovich [28] and Ben Cheikh and Douak [3] obtained the multiple orthogonal polynomials for the weights or  $(w_1, w_2) = x^\alpha(\rho_\nu, \rho_{\nu+1})$ , where  $\alpha > -1$  and  $\nu \geq 0$ . The vector Pearson equation is

$$\left[ x \begin{pmatrix} w_1 \\ w_2 \end{pmatrix} \right]' = \begin{pmatrix} \alpha + \nu + 1 & -1 \\ -x & \alpha + 1 \end{pmatrix} \begin{pmatrix} w_1 \\ w_2 \end{pmatrix}.$$

One has

$$\frac{\rho_\nu(x)}{\rho_{\nu+1}(x)} = \frac{1}{\pi^2} \int_0^\infty \frac{1}{(x+s)[J_{\nu+1}^2(2\sqrt{s}) + Y_{\nu+1}^2(2\sqrt{s})]} \frac{ds}{s},$$

so that this is a Nikishin system. Denote the type I function by

$$q_{n,m}^\alpha(x) = A_{n,m}(x)\rho_\nu(x) + B_{n,m}(x)\rho_{\nu+1}(x),$$

and the type II multiple orthogonal polynomials by  $p_{n,m}^\alpha$ .

**Property 1** (Rodrigues formula). *One has*

$$x^\alpha q_{n,n}^\alpha = \frac{d^{2n-1}}{dx^{2n-1}} [x^{2n-1+\alpha} \rho_\nu(x)],$$

$$x^\alpha q_{n,n-1}^\alpha = \frac{d^{2n-2}}{dx^{2n-2}} [x^{2n+\alpha-2} \rho_\nu(x)].$$

**Property 2** (recurrence relation). *Let  $P_{2n} = p_{n,n}^\alpha$  and  $P_{2n+1} = p_{n+1,n}^\alpha$ , then*

$$xP_n(x) = P_{n+1}(x) + b_n P_n(x) + c_n P_{n-1}(x) + d_n P_{n-2}(x)$$

with

$$\begin{aligned} b_n &= (n + \alpha + 1)(3n + \alpha + 2\nu) - (\alpha + 1)(\nu - 1), \\ c_n &= n(n + \alpha)(n + \alpha + \nu)(3n + 2\alpha + \nu), \\ d_n &= n(n - 1)(n + \alpha)(n + \alpha - 1)(n + \alpha + \nu)(n + \alpha + \nu - 1). \end{aligned}$$

Let  $Q_{2n} = q_{n,n}^\alpha$  and  $Q_{2n-1} = q_{n,n-1}^\alpha$ , then

$$xQ_n(x) = Q_{n-1}(x) + b_{n-1}Q_n(x) + c_n Q_{n+1}(x) + d_{n+1}Q_{n+2}(x).$$

### 2.3 Modified Bessel functions ( $I_\nu, I_{\nu+1}$ )

Another solution of the modified Bessel equation is given by the modified Bessel function  $I_\nu(x)$ . In [5] we considered

$$w_1(x) = x^{\nu/2} I_\nu(2\sqrt{x}) e^{-cx}, \quad w_2(x) = x^{(\nu+1)/2} I_{\nu+1}(2\sqrt{x}) e^{-cx}, \quad x > 0,$$

with  $\nu > -1$  and  $c > 0$ . The case  $c = 1$  was first studied in [8]. These weights are related to the noncentral  $\chi^2$ -distribution:

$$p_{\chi^2}^{(\nu,\lambda)}(x) = \frac{1}{2} \left(\frac{x}{\lambda}\right)^{(\nu-2)/4} I_{(\nu-2)/2}(\sqrt{\lambda x}) e^{-(\lambda+x)/2}, \quad x > 0,$$

with  $\lambda > 0$  and  $\nu \in \mathbb{N}$ . The Pearson equation is

$$\left[ x \begin{pmatrix} w_1 \\ w_2 \end{pmatrix} \right]' = \begin{pmatrix} \nu - cx & 1 \\ x & -cx \end{pmatrix} \begin{pmatrix} w_1 \\ w_2 \end{pmatrix}.$$

One has

$$\frac{w_1(x)}{w_2(x)} = \frac{\nu + 1}{x} + \sum_{n=1}^{\infty} \frac{1}{x + j_{\nu+1,n}^2/4} \frac{J_{\nu+2}(j_{\nu+1,n})}{J'_{\nu+1}(j_{\nu+1,n})},$$

so that we again have a Nikishin system. Let

$$Q_{2n}^\nu(x) = q_{n,n}^\nu(x), \quad Q_{2n+1}^\nu(x) = q_{n+1,n}^\nu(x),$$

and similarly

$$P_{2n}^\nu(x) = p_{n,n}^\nu(x), \quad P_{2n+1}^\nu(x) = p_{n+1,n}^\nu(x).$$

**Property 3** (raising-lowering). *For every  $\nu > -1$  and  $c > 0$  one has*

$$\frac{d}{dx}Q_n^{\nu+1}(x) = Q_{n+1}^\nu(x), \quad \text{and} \quad \frac{d}{dx}P_n^\nu(x) = nP_{n-1}^{\nu+1}(x).$$

**Property 4** (recurrence relation). *One has*

$$xP_n^\nu(x) = P_{n+1}^\nu(x) + b_nP_n^\nu(x) + c_nP_{n-1}^\nu(x) + d_nP_{n-2}^\nu(x),$$

with

$$\begin{aligned} b_n &= \frac{1}{c^2}[1 + c(\nu + 2n + 1)], \\ c_n &= \frac{n}{c^3}[2 + c(\nu + n)], \\ d_n &= \frac{n(n-1)}{c^4}. \end{aligned}$$

Furthermore

$$xQ_n^\nu(x) = Q_{n-1}^\nu(x) + b_{n-1}Q_n^\nu(x) + c_nQ_{n+1}^\nu(x) + d_{n+1}Q_{n+2}^\nu(x).$$

### 3 Simultaneous Gaussian quadrature

Suppose we have two measures  $\mu_1, \mu_2$  and one function  $f$ . We want to approximate integrals by sums

$$\begin{aligned} \int_{\mathbb{R}} f(x) d\mu_1(x) &\approx \sum_{k=1}^{2n} \lambda_{k,2n}^{(1)} f(x_{k,2n}), \\ \int_{\mathbb{R}} f(x) d\mu_2(x) &\approx \sum_{k=1}^{2n} \lambda_{k,2n}^{(2)} f(x_{k,2n}). \end{aligned}$$

**Theorem 1** (Borges [4]). *If we take for  $\{x_{k,2n}, 1 \leq k \leq 2n\}$  the zeros of the type II multiple orthogonal  $P_{n,n}$  for the two measures  $(\mu_1, \mu_2)$  and use interpolatory quadrature, then the quadrature is exact for polynomials  $f$  of degree  $\leq 3n - 1$ .*

There are  $2n$  function evaluations. If we use ordinary Gaussian quadrature with  $n$  nodes for each integral, then we also use  $2n$  function evaluations, but the quadrature then is exact for polynomials  $f$  of degree  $\leq 2n - 1$ . Important numerical aspects are:

- How to compute the quadrature nodes  $x_{k,2n}$ ,  $1 \leq k \leq 2n$  (the zeros of  $P_{n,n}$ )?
- How to compute the quadrature weights

$$\lambda_{k,2n}^{(1)} = \int_{\mathbb{R}} \frac{P_{n,n}(x)}{(x - x_{k,2n})P'_{n,n}(x_{k,2n})} d\mu_1(x), \quad (3.1)$$

$$\lambda_{k,2n}^{(2)} = \int_{\mathbb{R}} \frac{P_{n,n}(x)}{(x - x_{k,2n})P'_{n,n}(x_{k,2n})} d\mu_2(x). \quad (3.2)$$

Some relevant analytical problems are

- The convergence of the quadrature rules as  $n \rightarrow \infty$ .
- Are the weights  $\lambda_{k,2n}^{(1)}$  and  $\lambda_{k,2n}^{(2)}$  positive? If not, are they small?

We have investigated some of the analytical problems of simultaneous Gaussian quadrature for an Angelesco system in [18] and for multiple Hermite polynomials in [27].

We formulated the simultaneous Gaussian quadrature for  $2n$  nodes. If the number of nodes is odd ( $2n + 1$ ), then one uses the zeros of the type II multiple orthogonal  $P_{n+1,n}$  and

$$\int_{\mathbb{R}} f(x) d\mu_1(x) = \sum_{k=1}^{2n+1} \lambda_{k,2n+1}^{(1)} f(x_{k,2n+1})$$

holds for polynomials  $f$  of degree  $\leq 3n + 1$ , and

$$\int_{\mathbb{R}} f(x) d\mu_2(x) = \sum_{k=1}^{2n+1} \lambda_{k,2n+1}^{(2)} f(x_{k,2n+1})$$

holds for polynomials  $f$  of degree  $\leq 3n$ . For multi-indices  $(n, m)$  one takes the nodes at the zeros of  $P_{n,m}$  and

$$\int_{\mathbb{R}} f(x) d\mu_1(x) = \sum_{k=1}^{n+m} \lambda_{k,n+m}^{(1)} f(x_{k,n+m}), \quad \deg f \leq 2n + m - 1,$$

$$\int_{\mathbb{R}} f(x) d\mu_2(x) = \sum_{k=1}^{n+m} \lambda_{k,n+m}^{(2)} f(x_{k,n+m}), \quad \deg f \leq n + 2m - 1.$$

There is an important relation with Hermite-Padé approximation. Recall that for type II Hermite-Padé approximation one has

$$P_{\vec{n}}(z)f_j(z) - Q_{\vec{n},j}(z) = \mathcal{O}\left(\frac{1}{z^{n_j+1}}\right), \quad z \rightarrow \infty.$$

For the rational approximants one then has

$$f_j(z) - \frac{Q_{\vec{n},j}(z)}{P_{\vec{n}}(z)} = \mathcal{O}\left(\frac{1}{z^{|\vec{n}|+n_j+1}}\right),$$

and if one decomposes the rational approximant into partial fractions, then

$$\frac{Q_{\vec{n},j}(z)}{P_{\vec{n}}(z)} = \sum_{k=1}^{|\vec{n}|} \frac{\lambda_{k,\vec{n}}^{(j)}}{z - x_{k,\vec{n}}}.$$

Hence the poles of the type II Hermite-Padé approximants are the quadrature nodes and the residues are the quadrature weights. This extends the well-known connection between Gaussian quadrature and Padé approximation. If  $\mu_j$  is supported on  $[a, b]$  and  $g$  is analytic in a neighborhood  $\Omega$  of  $[a, b]$ , then the error is given by

$$\int_a^b g(x) d\mu_j(x) - \sum_{k=1}^{|\vec{n}|} \lambda_{k,\vec{n}}^{(j)} g(x_{k,\vec{n}}) = \frac{1}{2\pi i} \int_{\Gamma} g(z) \left( f_j(z) - \frac{Q_{\vec{n},j}(z)}{P_{\vec{n}}(z)} \right) dz,$$

where  $\Gamma$  is a closed contour in  $\Omega$  around  $[a, b]$ . Hence results about the asymptotic behavior of Hermite-Padé approximation imply results for the convergence of the quadrature formula when the integrand  $g$  is sufficiently smooth.



## 4 Recurrence relations

Denote by  $\vec{e}_j = (0, \dots, 0, 1, 0, \dots, 0)$  the standard unit vectors. Multiple orthogonal polynomials always satisfy a system of linear recurrence relations connecting  $P_{\vec{n}}$  to its nearest neighbors  $P_{\vec{n}+\vec{e}_k}$  from above and  $P_{\vec{n}-\vec{e}_j}$  from below [24]:

$$xP_{\vec{n}}(x) = P_{\vec{n}+\vec{e}_k}(x) + b_{\vec{n},k}P_{\vec{n}}(x) + \sum_{j=1}^r a_{\vec{n},j}P_{\vec{n}-\vec{e}_j}(x), \quad 1 \leq k \leq r. \quad (4.1)$$

For the type I functions

$$Q_{\vec{n}}(x) = \sum_{j=1}^r A_{\vec{n},j}(x)w_j(x), \quad w_j(x) = \frac{d\mu_j(x)}{d\mu(x)},$$

where  $\mu = \mu_1 + \mu_2 + \dots + \mu_r$ , the nearest neighbor recurrence relations are

$$xQ_{\vec{n}}(x) = Q_{\vec{n}-\vec{e}_k}(x) + b_{\vec{n}-\vec{e}_k,k}Q_{\vec{n}}(x) + \sum_{j=1}^r a_{\vec{n},j}Q_{\vec{n}+\vec{e}_j}(x), \quad 1 \leq k \leq r. \quad (4.2)$$

If we combine these  $r$  recurrence relations, then one can find a  $(r+2)$ -term recurrence relation on the stepline: for  $n = kr + \ell$  we take  $\vec{n} = \underbrace{(k+1, \dots, k+1, k, \dots, k)}_{\ell \text{ times}}$  and we

define the type II polynomials on the stepline by  $P_n(x) = P_{\vec{n}}(x)$ . This  $(r+2)$ -term recurrence relation is

$$xP_n(x) = P_{n+1}(x) + \sum_{j=0}^r c_{n,j}P_{n-j}(x). \quad (4.3)$$

There is also a  $(r+2)$ -term recurrence relation for the type I functions on the stepline:

$$xQ_n(x) = Q_{n-1}(x) + \sum_{j=0}^r c_{n+j-1,j}Q_{n+j}(x). \quad (4.4)$$

Filipuk, Haneczok and Van Assche [10] worked out an algorithm to go from the nearest neighbor recurrence coefficients  $(a_{\vec{n},j}, b_{\vec{n},j}; 1 \leq j \leq r)_{\vec{n} \in \mathbb{N}^r}$  to the stepline coefficients  $(c_{n,0}, c_{n,1}, \dots, c_{n,r})_{n \in \mathbb{N}}$ . They also gave algorithms to go from the recurrence coefficients of the orthogonal polynomials of the measures  $\mu_j$  to the nearest neighbor recurrence coefficients. We will illustrate this for the case  $r = 2$ . The nearest neighbor recurrence relations are

$$xP_{n,m}(x) = P_{n+1,m}(x) + c_{n,m}P_{n,m}(x) + a_{n,m}P_{n-1,m}(x) + b_{n,m}P_{n,m-1}(x), \quad (4.5)$$

$$xP_{n,m}(x) = P_{n,m+1}(x) + d_{n,m}P_{n,m}(x) + a_{n,m}P_{n-1,m}(x) + b_{n,m}P_{n,m-1}(x). \quad (4.6)$$

Subtracting both equations gives

$$P_{n+1,m}(x) - P_{n,m+1}(x) = (d_{n,m} - c_{n,m})P_{n,m}(x). \quad (4.7)$$

Denote the type II polynomials on the stepline by  $P_{2k}(x) = P_{k,k}(x)$ ,  $P_{2k+1}(x) = P_{k+1,k}(x)$ . Take  $m = n$  then (4.5) is

$$xP_{n,n}(x) = P_{n+1,n}(x) + c_{n,n}P_{n,n}(x) + a_{n,n}P_{n-1,n}(x) + b_{n,n}P_{n,n-1}(x).$$

Use (4.7) to replace  $P_{n-1,n}(x)$  by  $P_{n,n-1}(x) + (c_{n-1,n-1} - d_{n-1,n-1})P_{n-1,n-1}(x)$  to find

$$xP_{2n}(x) = P_{2n+1}(x) + c_{n,n}P_{2n}(x) + (a_{n,n} + b_{n,n})P_{2n-1}(x) + a_{n,n}(c_{n-1,n-1} - d_{n-1,n-1})P_{2n-2}(x).$$

In a similar way, by taking  $m = n - 1$ , (4.6) is

$$xP_{n,n-1}(x) = P_{n,n}(x) + d_{n,n-1}P_{n,n-1}(x) + a_{n,n-1}P_{n-1,n-1}(x) + b_{n,n-1}P_{n,n-2}(x).$$

Use (4.7) to replace  $P_{n,n-2}(x)$  by  $P_{n-1,n-1}(x) + (d_{n-1,n-2} - c_{n-1,n-2})P_{n-1,n-2}(x)$  to find

$$xP_{2n-1}(x) = P_{2n}(x) + d_{n,n-1}P_{2n-1}(x) + (a_{n,n-1} + b_{n,n-1})P_{2n-2}(x) + b_{n,n-1}(d_{n-1,n-2} - c_{n-1,n-2})P_{2n-3}(x).$$

This means that the four-term recurrence relation for the type II polynomials on the stepline is

$$xP_n(x) = P_{n+1}(x) + b_nP_n(x) + c_nP_{n-1}(x) + d_nP_{n-2}(x), \quad (4.8)$$

with

$$\begin{aligned} b_{2k} &= c_{k,k}, & b_{2k+1} &= d_{k+1,k} \\ c_{2k} &= a_{k,k} + b_{k,k}, & c_{2k+1} &= a_{k+1,k} + b_{k+1,k}, \\ d_{2k} &= a_{k,k}(c_{k-1,k-1} - d_{k-1,k-1}), & d_{2k+1} &= b_{k+1,k}(d_{k,k-1} - c_{k,k-1}). \end{aligned}$$

For the type I functions, the nearest neighbor recurrence relations are

$$\begin{aligned} xQ_{n,m}(x) &= Q_{n-1,m}(x) + c_{n-1,m}Q_{n,m}(x) + a_{n,m}Q_{n+1,m}(x) + b_{n,m}(x)Q_{n,m+1}(x), \\ xQ_{n,m}(x) &= Q_{n,m-1}(x) + d_{n,m-1}Q_{n,m}(x) + a_{n,m}Q_{n+1,m}(x) + b_{n,m}(x)Q_{n,m+1}(x). \end{aligned}$$

Subtracting both equations gives

$$Q_{n-1,m}(x) - Q_{n,m-1}(x) = (d_{n,m-1} - c_{n-1,m})Q_{n,m}(x). \quad (4.9)$$

In a similar way as before, one then finds for the type I functions on the stepline  $Q_{2k}(x) = Q_{k,k}(x)$ ,  $Q_{2k+1}(x) = Q_{k+1,k}(x)$

$$xQ_n(x) = Q_{n-1}(x) + b_{n-1}Q_n(x) + c_nQ_{n+1}(x) + d_{n+1}Q_{n+2}(x). \quad (4.10)$$

Using the coefficients in this four-term recurrence relation, we can create a banded Hessenberg matrix

$$H_N = \begin{pmatrix} b_0 & 1 & 0 & 0 & 0 & 0 & \cdots & 0 \\ c_1 & b_1 & 1 & 0 & 0 & 0 & \cdots & 0 \\ d_2 & c_2 & b_2 & 1 & 0 & 0 & \cdots & 0 \\ 0 & d_3 & c_3 & b_3 & 1 & 0 & \cdots & 0 \\ 0 & 0 & d_4 & c_4 & b_4 & 1 & \cdots & 0 \\ \vdots & & & \ddots & \ddots & \ddots & \ddots & \vdots \\ 0 & \cdots & 0 & 0 & d_{N-2} & c_{N-2} & b_{N-2} & 1 \\ 0 & \cdots & 0 & 0 & 0 & d_{N-1} & c_{N-1} & b_{N-1} \end{pmatrix}. \quad (4.11)$$

From now on we will use  $N$  for the number of quadrature nodes. If  $N = 2n$  then this corresponds to the multi-index  $(n, n)$  and for  $N = 2n + 1$  to the multi-index  $(n + 1, n)$ . The recurrence relation (4.8) for the type II polynomials can then be written as

$$H_N \begin{pmatrix} P_0(x) \\ P_1(x) \\ \vdots \\ P_{N-1}(x) \end{pmatrix} + \begin{pmatrix} 0 \\ \vdots \\ 0 \\ P_N(x) \end{pmatrix} = x \begin{pmatrix} P_0(x) \\ P_1(x) \\ \vdots \\ P_{N-1}(x) \end{pmatrix}.$$

If we evaluate this at a zero  $x_{k,N}$  of  $P_N$ , then we get the following result:

**Property 5.** *If  $P_N(x_{k,N}) = 0$  then  $x_{k,N}$  is an eigenvalue of  $H_N$  with **right** eigenvector  $(P_0(x_{k,N}), P_1(x_{k,N}), \dots, P_{N-1}(x_{k,N}))$ .*

There is a similar property for the type I multiple orthogonal polynomials  $(A_N, B_N)$  for which

$$\begin{aligned} A_{2k}(x) &= A_{k,k}(x), & A_{2k+1}(x) &= A_{k+1,k}(x), \\ B_{2k}(x) &= B_{k,k}(x), & B_{2k+1}(x) &= B_{k+1,k}(x). \end{aligned}$$

**Property 6.** *If  $P_N(x_{k,N}) = 0$  then  $x_{k,N}$  is an eigenvalue of  $H_N$  with **left** eigenvector  $(C_1 A_1(x_{k,N}) + C_2 B_1(x_{k,N}), C_1 A_2(x_{k,N}) + C_2 B_2(x_{k,N}), \dots, C_1 A_n(x_{k,N}) + C_2 B_n(x_{k,N}))$ , with constants  $C_1$  and  $C_2$  satisfying*

$$\begin{cases} C_1 A_{N+1}(x_{k,N}) + C_2 B_{N+1}(x_{k,N}) = 0, \\ C_1 A_{N+2}(x_{k,N}) + C_2 B_{N+2}(x_{k,N}) = 0. \end{cases} \quad (4.12)$$

*Proof.* For the type I function  $Q_n(x) = A_n(x)w_1(x) + B_n(x)w_2(x)$ , the four-term recurrence relation (4.10) is equivalent to

$$\begin{aligned} & (Q_1(x) \quad Q_2(x) \quad \cdots \quad Q_N(x)) H_N \\ & + \begin{pmatrix} 0 & \cdots & 0 & d_N Q_{N+1}(x) & 0 \end{pmatrix} \\ & + \begin{pmatrix} 0 & \cdots & 0 & c_N Q_{N+1}(x) & d_{N+1} Q_{N+2}(x) \end{pmatrix} \\ & = x (Q_1(x) \quad Q_2(x) \quad \cdots \quad Q_N(x)). \end{aligned}$$

This relation also holds for the polynomials  $A_n(x)$  and  $B_n(x)$ , hence

$$\begin{aligned} & (A_1(x) \quad A_2(x) \quad \cdots \quad A_N(x)) H_N \\ & + \begin{pmatrix} 0 & \cdots & 0 & d_N A_{N+1}(x) & 0 \end{pmatrix} \\ & + \begin{pmatrix} 0 & \cdots & 0 & c_N A_{N+1}(x) & d_{N+1} A_{N+2}(x) \end{pmatrix} \\ & = x (A_1(x) \quad A_2(x) \quad \cdots \quad A_N(x)), \end{aligned}$$

and

$$\begin{aligned} & (B_1(x) \quad B_2(x) \quad \cdots \quad B_N(x)) H_N \\ & + \begin{pmatrix} 0 & \cdots & 0 & d_N B_{N+1}(x) & 0 \end{pmatrix} \\ & + \begin{pmatrix} 0 & \cdots & 0 & c_N B_{N+1}(x) & d_{N+1} B_{N+2}(x) \end{pmatrix} \\ & = x (B_1(x) \quad B_2(x) \quad \cdots \quad B_N(x)), \end{aligned}$$

and therefore also for a linear combination of  $A_n$  and  $B_n$

$$\begin{aligned} & (C_1A_1(x) + C_2B_1(x) \quad C_1A_2(x) + C_2B_2(x) \quad \cdots \quad C_1A_N(x) + C_2B_N(x)) H_N \\ & + \begin{pmatrix} 0 & \cdots & 0 & d_N[C_1A_{N+1}(x) + C_2B_{N+1}(x)] & 0 \\ 0 & \cdots & 0 & c_N[C_1A_{N+1}(x) + C_2B_{N+1}(x)] & d_{N+1}[C_1A_{N+2}(x) + C_2B_{N+2}(x)] \end{pmatrix} \\ & = x (C_1A_1(x) + C_2B_1(x) \quad C_1A_2(x) + C_2B_2(x) \quad \cdots \quad C_1A_N(x) + C_2B_N(x)). \end{aligned}$$

Choose  $C_1$  and  $C_2$  such that

$$\begin{cases} C_1A_{N+1}(x) + C_2B_{N+1}(x) = 0, \\ C_1A_{N+2}(x) + C_2B_{N+2}(x) = 0. \end{cases}$$

This is a homogeneous system of equations for  $C_1$  and  $C_2$ , which only has a non-trivial solution if the matrix is singular. Mahler's relation between type I and type II polynomials [26, Eq. (4.2)] gives

$$\det \begin{pmatrix} A_{N+1}(x) & B_{N+1}(x) \\ A_{N+2}(x) & B_{N+2}(x) \end{pmatrix} = C_3P_N(x)$$

for some constant  $C_3$ . Hence if  $x_{k,N}$  is a zero of  $P_N$ , then we can find  $(C_1, C_2) \neq (0, 0)$  so that (4.12) holds.  $\square$

Observe that  $C_1$  and  $C_2$  may depend on the eigenvalue  $x_{k,N}$ .

## 5 Golub-Welsch type result

We are now ready to prove the main result.

**Theorem 2.** *The quadrature nodes  $\{x_{j,N}, 1 \leq j \leq N\}$  for simultaneous Gaussian quadrature are given by the eigenvalues of the banded Hessenberg matrix  $H_N$ . Let  $\vec{v}_{N,j}$  be the right eigenvector of  $H_N$  for the eigenvalue  $x_{j,N}$  with first component 1 and  $\vec{u}_{N,j}$  be a left eigenvector for  $x_{j,N}$ . Then*

$$\begin{aligned} \lambda_{j,N}^{(1)} &= \frac{D_{1,1}\vec{u}_{N,j}(1)}{\langle \vec{u}_{N,j}, \vec{v}_{N,j} \rangle}, \\ \lambda_{j,N}^{(2)} &= \frac{D_{2,1}\vec{u}_{N,j}(1) + D_{2,2}\vec{u}_{N,j}(2)}{\langle \vec{u}_{N,j}, \vec{v}_{N,j} \rangle}, \end{aligned}$$

with constants  $D_{i,j}$  given by

$$\begin{pmatrix} D_{1,1} & 0 \\ D_{2,1} & D_{2,2} \end{pmatrix} = \begin{pmatrix} A_1 & 0 \\ A_2 & B_2 \end{pmatrix}^{-1}.$$

This result was given for general  $r$  in Coussement-Van Assche [6, Thm.3.2].

*Proof.* We first need the Christoffel-Darboux formula for multiple orthogonal polynomials (Daems-Kuijlaars [7]). Let  $N = |\vec{n}|$  and  $(\vec{n}_k)_{0 \leq k \leq N}$  be a path in  $\mathbb{N}^r$  from  $\vec{n}_0 = \vec{0}$  to  $\vec{n}_N = \vec{n}$  such that  $\vec{n}_{k+1} - \vec{n}_k = \vec{e}_j$  for some  $1 \leq j \leq r$ . Then

$$(x - y) \sum_{k=0}^{N-1} P_{\vec{n}_k}(x) Q_{\vec{n}_{k+1}}(y) = P_{\vec{n}}(x) Q_{\vec{n}}(y) - \sum_{j=1}^r a_{\vec{n},j} P_{\vec{n} - \vec{e}_j}(x) Q_{\vec{n} + \vec{e}_j}(y).$$

Here we used the version as it is given in terms of the nearest neighbor recurrence coefficients [24]. For  $r = 2$  and on the stepline, it becomes

$$(x - y) \sum_{k=0}^{N-1} P_k(x) Q_{k+1}(y) = P_N(x) Q_N(y) - c_N P_{N-1}(x) Q_{N+1}(y) \\ - d_N P_{N-2}(x) Q_{N+1}(y) - d_{N+1} P_{N-1}(x) Q_{N+2}(y).$$

The Christoffel-Darboux formula also holds if we replace  $Q_k(y)$  by  $C_1 A_k(y) + C_2 B_k(y)$ :

$$(x - y) \sum_{k=0}^{N-1} P_k(x) [C_1 A_{k+1}(y) + C_2 B_{k+1}(y)] \\ = P_N(x) [C_1 A_N(y) + C_2 B_N(y)] - c_N P_{N-1}(x) [C_1 A_{N+1}(y) + C_2 B_{N+1}(y)] \\ - d_N P_{N-2}(x) [C_1 A_{N+1}(y) + C_2 B_{N+1}(y)] - d_{N+1} P_{N-1}(x) [C_1 A_{N+2}(y) + C_2 B_{N+2}(y)].$$

Take  $y = x_{j,N}$  a zero of  $P_N$ , then (4.12) gives

$$\sum_{k=0}^{N-1} P_k(x) [C_1 A_{k+1}(x_{j,N}) + C_2 B_{k+1}(x_{j,N})] = \frac{P_N(x)}{x - x_{j,N}} [C_1 A_N(x_{j,N}) + C_2 B_N(x_{j,N})]. \quad (5.1)$$

Recall that the quadrature nodes are given by (3.1) and (3.2). Integrating (5.1) with measure  $\mu_1$  gives

$$\sum_{k=0}^{N-1} [C_1 A_{k+1}(x_{j,N}) + C_2 B_{k+1}(x_{j,N})] \int_{\mathbb{R}} P_k(x) d\mu_1(x) \\ = \int_{\mathbb{R}} \frac{P_N(x)}{x - x_{j,N}} d\mu_1(x) [C_1 A_N(x_{j,N}) + C_2 B_N(x_{j,N})],$$

hence if we retain only the non-vanishing integrals

$$\lambda_{j,N}^{(1)} [C_1 A_N(x_{j,N}) + C_2 B_N(x_{j,N})] = \frac{1}{P'_N(x_{j,N})} \int_{\mathbb{R}} P_0(x) d\mu_1(x) [C_1 A_1 + C_2 B_1]. \quad (5.2)$$

In a similar way, integrating (5.1) with measure  $\mu_2$  gives

$$\sum_{k=0}^{N-1} [C_1 A_{k+1}(x_{j,N}) + C_2 B_{k+1}(x_{j,N})] \int_{\mathbb{R}} P_k(x) d\mu_2(x) \\ = \int_{\mathbb{R}} \frac{P_N(x)}{x - x_{j,N}} d\mu_2(x) [C_1 A_N(x_{j,N}) + C_2 B_N(x_{j,N})],$$

hence if we retain only the non-vanishing integrals

$$\lambda_{j,N}^{(2)} [C_1 A_N(x_{j,N}) + C_2 B_N(x_{j,N})] \\ = \frac{1}{P'_N(x_{j,N})} \left( \int_{\mathbb{R}} P_0(x) d\mu_2(x) [C_1 A_1 + C_2 B_1] \right. \\ \left. + \int_{\mathbb{R}} P_1(x) d\mu_2(x) [C_1 A_2 + C_2 B_2] \right). \quad (5.3)$$

If we denote

$$\begin{aligned} D_{1,1} &= \int_{\mathbb{R}} P_0(x) d\mu_1(x), & D_{1,2} &= \int_{\mathbb{R}} P_1(x) d\mu_1(x), \\ D_{2,1} &= \int_{\mathbb{R}} P_0(x) d\mu_2(x), & D_{2,2} &= \int_{\mathbb{R}} P_1(x) d\mu_2(x), \end{aligned}$$

and use the biorthogonality

$$\begin{aligned} \int_{\mathbb{R}} P_0(x)[A_1 w_1(x) + B_1 w_2(x)] d\mu(x) &= 1, & \int_{\mathbb{R}} P_0(x)[A_2 w_1(x) + B_2 w_2(x)] d\mu(x) &= 0, \\ \int_{\mathbb{R}} P_1(x)[A_1 w_1(x) + B_1 w_2(x)] d\mu(x) &= 0, & \int_{\mathbb{R}} P_1(x)[A_2 w_1(x) + B_2 w_2(x)] d\mu(x) &= 1, \end{aligned}$$

then we find

$$\begin{pmatrix} A_1 & 0 \\ A_2 & B_2 \end{pmatrix} \begin{pmatrix} D_{1,1} & D_{1,2} \\ D_{2,1} & D_{2,2} \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

Observe that  $A_1 = A_{1,0}(x)$ ,  $A_2 = A_{1,1}(x)$  and  $B_2 = B_{1,1}(x)$  are polynomials of degree 0 and hence these are constants, whereas  $B_1 = B_{1,0} = 0$  and by the orthogonality of type II multiple orthogonal polynomials we also have  $D_{1,2} = 0$ .

In order to find a formula for  $P'_N(x_{j,N})$ , we take the limit  $x \rightarrow x_{j,N}$  in (5.1) to find

$$\sum_{k=0}^{N-1} P_k(x_{j,N})[C_1 A_{k+1}(x_{j,N}) + C_2 B_{k+1}(x_{j,N})] = P'_N(x_{j,N})[C_1 A_N(x_{j,N}) + C_2 B_N(x_{j,N})]. \quad (5.4)$$

The left hand side is the inner product of the left eigenvector and the right eigenvector of  $H_N$  for the eigenvalue  $x_{j,N}$ . Combining (5.2), (5.3) and (5.4) and taking into account Property 5 and 6 then gives the main result.  $\square$

## 6 Examples

### 6.1 Bessel functions $K_\nu$

We will consider the multiple orthogonal polynomials related to the modified Bessel functions  $K_\nu$  and  $K_{\nu+1}$ . Let  $(w_1, w_2) = x^\alpha(\rho_\nu, \rho_{\nu+1})$  with

$$\rho_\nu(x) = 2x^{\nu/2} K_\nu(2\sqrt{x}), \quad x > 0.$$

We take  $\alpha = 1$  and  $\nu = 0$  and we compute the eigenvalues of  $H_N$  for  $N = 10$ , together with the weights  $\lambda_{j,10}^{(1)}$  and  $\lambda_{j,10}^{(2)}$  using Theorem 2. The type I polynomials  $(A_{1,0}, B_{1,0})$  and  $(A_{1,1}, B_{1,1})$  are the constants

$$A_{1,0} = 1, \quad B_{1,0} = 0, \quad A_{1,1} = \alpha + \nu + 1, \quad B_{1,1} = -1,$$

and if we use the normalization (2.1)

$$\int_0^\infty x^\alpha [\widehat{A}_{n,m}(x)\rho_\nu(x) + \widehat{B}_{n,m}(x)\rho_{\nu+1}(x)] x^{n+m-1} dx = 1,$$

then we get

$$\begin{pmatrix} A_1 & 0 \\ A_2 & B_2 \end{pmatrix} = \begin{pmatrix} \frac{1}{\Gamma(\alpha+\nu+1)\Gamma(\alpha+1)} & 0 \\ -\frac{\alpha+\nu+1}{\Gamma(\alpha+\nu+2)\Gamma(\alpha+2)} & \frac{1}{\Gamma(\alpha+\nu+2)\Gamma(\alpha+2)} \end{pmatrix},$$

and hence for  $\alpha = 1$  and  $\nu = 0$  we have

$$\begin{pmatrix} D_{1,1} & 0 \\ D_{2,1} & D_{2,2} \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ -1/2 & 1/4 \end{pmatrix}^{-1} = \begin{pmatrix} 1 & 0 \\ 2 & 4 \end{pmatrix}.$$

We used Maple with `Digits:=100` for our calculations and we show the 20 first digits after the decimal point in Fig. 6.1.

$j$	$x_{j,10}$	$\lambda_{j,10}^{(1)}$	$\lambda_{j,10}^{(2)}$
1	0.52720348133440875760	0.27736269648616286974	0.26086734230400106004
2	2.74106066716069179819	0.46938499819336417730	0.88799214753397210390
3	8.13937609771412899056	0.21135584109286564463	0.65379039925659229785
4	18.66164146312871349710	0.03854365644852726770	0.17589229666877292663
5	36.89653691488348638176	0.00322544756122977083	0.02038307627872880093
6	66.43703332978391524587	0.00012523808693942895	0.00105166051829272396
7	112.55686514754090244347	0.00000210903533490802	0.00002289663649071884
8	183.67841427499791701294	0.00000001307455465436	0.00000018043669350953
9	295.27746298319776238423	0.00000000002101777610	0.00000000036637784733
10	485.08440564025807348828	0.00000000000000350239	0.00000000000007801100

Figure 6.1: Quadrature nodes and quadrature weights for  $(w_1, w_2) = x(\rho_0, \rho_1)$ .

### Algorithm 1: Maple

```

with(LinearAlgebra):
b:=n->(n+alpha+1)*(3*n+alpha+2*nu)-(alpha+1)*(nu-1);
c:=n->n*(n+alpha)*(n+alpha+nu)*(3*n+2*alpha+nu);
d:=n->n*(n-1)*(n+alpha)*(n+alpha-1)*(n+alpha+nu)*(n+alpha+nu-1);
H:=n->BandMatrix([[seq(d(k),k=2..n-1)],[seq(c(k),k=1..n-1)],
  [seq(b(k),k=0..n-1)],[seq(1,k=1..n-1)]]);
alpha:=1.;nu:=0.;
Digits:=100;
N:=10;
E,RE:=Eigenvectors(H(N)): % right eigenvalues/vectors (descending)
eig:=Eigenvectors(Transpose(H(N)), output = list):
eig := sort(eig, (a, b) -> is(Re(b[1]) < Re(a[1]))):
EE:=Vector[column]([seq(eig[k][1],k=1..N)]): % eigenvalues (descending)
LE:=Matrix([seq(op(eig[k][3]), k = 1 .. N)]): % left eigenvectors
A:=Matrix([[1/(GAMMA(alpha+nu+1)*GAMMA(alpha+1)),0],
  [-(alpha+nu+1)/(GAMMA(alpha+nu+2)*GAMMA(alpha+2)),
  1/(GAMMA(alpha+nu+2)*GAMMA(alpha+2))]]);
D:=MatrixInverse(A);
u:=j->LE(1:N,j); % left eigenvectors
v:=j->RE(1:N,N-j+1)/RE(1,N-j+1); % normalized right eigenvectors

```

```

lambda1:=j->D(1,1)*u(j)[1]/DotProduct(u(j),v(j));
lambda2:=j->(D(2,1)*u(j)[1]+D(2,2)*u(j)[2])/DotProduct(u(j),v(j));
seq(EE(k),k=1..N); % quadrature nodes
seq(lambda1(k),k=1..N); % quadrature weights (first weight)
seq(lambda2(k),k=1..N); % quadrature weights (second weight)

```

To illustrate the quality of the quadratures, we will simultaneously compute

$$I_1 = \int_0^\infty e^{-x} x \rho_0(x) dx = 0.1926947246\dots,$$

$$I_2 = \int_0^\infty e^{-x} x \rho_1(x) dx = 0.2109579130\dots$$

using  $N = 10, 20, 30, 40, 50$  nodes in Maple. We show the first 10 decimals in Fig. 6.2.

$N$	$I_1$	$I_2$
10	0.1940521520	0.2114457811
20	0.1926653563	0.2109395236
30	0.1926958911	0.2109610461
40	0.1926947184	0.2109576142
50	0.1926947165	0.2109579157

Figure 6.2: Simultaneous quadrature to  $I_1$  and  $I_2$

The results show that the quadrature formulas converge rather slowly to the correct value. A possible explanation is that the quadrature nodes become very large and the corresponding quadrature weights are very small, so only a small proportion of terms in the quadrature sum contribute to the result.

## 6.2 Bessel functions $I_\nu$

For the multiple orthogonal polynomials with weights  $(w_1, w_2) = (\omega_{\nu,c}, \omega_{\nu+1,c})$ , where

$$\omega_{\nu,c}(x) = x^{\nu/2} I_\nu(2\sqrt{x}) e^{-cx}, \quad x > 0,$$

we take  $c = 1$  and  $\nu = 0$  and compute the eigenvalues of  $H_N$  for  $N = 10$ , together with the weights  $\lambda_{j,10}^{(1)}$  and  $\lambda_{j,10}^{(2)}$  using Theorem 2. After normalization we get

$$\begin{pmatrix} A_1 & 0 \\ A_2 & B_2 \end{pmatrix} = e^{-1/c} \begin{pmatrix} c^{\nu+1} & 0 \\ -c^{\nu+2} & c^{\nu+3} \end{pmatrix},$$

so that for  $c = 1$  and  $\nu = 0$  we have

$$\begin{pmatrix} D_{1,1} & 0 \\ D_{2,1} & D_{2,2} \end{pmatrix} = \begin{pmatrix} 1/e & 0 \\ -1/e & 1/e \end{pmatrix}^{-1} = \begin{pmatrix} e & 0 \\ e & e \end{pmatrix}.$$

We use Matlab and we show the first 10 digits after the decimal point in Fig. 6.3.



$j$	$x_{j,10}$	$\lambda_{j,10}^{(1)}$	$\lambda_{j,10}^{(2)}$
1	0.1531952228	0.3913749988	0.0557885974
2	0.8105837014	0.8175616919	0.4874004644
3	2.0077223654	0.8459198767	0.9551942639
4	3.7719525634	0.4850707607	0.8091738873
5	6.1482336073	0.1517396396	0.3357737316
6	9.2079873838	0.0246520172	0.0683288497
7	13.0663024491	0.0019027391	0.0063827530
8	17.9203555594	0.0000595495	0.0002366956
9	24.1543375116	0.0000005543	0.0000025816
10	32.7593296369	0.0000000007	0.0000000038

Figure 6.3: Quadrature nodes and quadrature weights for  $(w_1, w_2) = (\omega_{0,1}, \omega_{1,1})$ .

### Algorithm 2: Matlab

```

1  C=1
2  nu=0
3  n=10
4  b=(1+C*(nu+1+2*[0:1:n-1]))/C^2
5  c=[1:1:n-1].*(2+C*(nu+[1:1:n-1]))/C^3
6  d=[2:1:n-1].*([2:1:n-1]-1)/C^4
7  H=diag(ones(1,n-1),1)+diag(b)+diag(c,-1)+diag(d,-2)
8  [R,E,L]=eig(H)
9  U=L
10 V=R/diag(R(1,1:n))
11 D=exp(1/C)*inv([[C^(nu+1),0];[-C^(nu+2),C^(nu+3)]])
12 lambda1=D(1,1)*U(1,1:n)/dot(U,V)
13 lambda2=(D(2,1)*U(1,1:n)+D(2,2)*U(2,1:n))./dot(U,V)
14 [diag(E),lambda1',lambda2']

```

We will simultaneously compute the integrals

$$J_1 = \int_0^\infty \cos(x)\omega_{0,1}(x) dx = 0.328224976685277123104160354501976758\dots$$

$$J_2 = \int_0^\infty \cos(x)\omega_{1,1}(x) dx = -0.39521954160680745592163128352397786234\dots$$

with  $N = 10, 20, 30, 40, 50$  quadrature nodes in Maple (with `Digits:=100`). The results are in Fig. 6.4. The results in Matlab were comparable for  $N = 10$  but for  $N = 20$  and higher some of the eigenvalues and eigenvectors became complex, due to machine precision. This was avoided in Maple by choosing `Digits:=100`. Alternatively one could use variable precision arithmetic (`vpa`) in Matlab but then Algorithm 2 needs to be modified because the command `eig` does not give the left eigenvalues.

Clearly the results of the quadrature are much better than for the previous example.

$N$	$J_1$	$J_2$
10	0.328340082411357	-0.395132567462746
20	0.32822497721656944454	-0.39521953865314722695
30	0.32822497668527696693	-0.395219541606806392096
40	0.328224976685277123103734621725	-0.39521954160680745592554825999940
50	0.32822497668527712310416035472	-0.3952195416068074559216312825809

Figure 6.4: Simultaneous quadrature to  $J_1$  and  $J_2$

## 7 Concluding remarks

We have shown that the nodes and the quadrature weights for simultaneous Gaussian quadrature can be computed numerically using the eigenvalues and the left and right eigenvectors of a banded Hessenberg matrix, thus extending the well-known approach of Golub and Welsch for Gaussian quadrature and symmetric tridiagonal matrices. In the present paper we investigated the case of two measures, but the approach works for any number of measures [6], however the number of diagonals of the Hessenberg matrix increases. Algorithms for the case of three or more measures using this Golub-Welsch approach have not been implemented yet. For two and three measures Tomović and Stanić [23] published an algorithm in which the quadrature weights are computed by solving some system of equations, and this approach was extended to more measures by Jovanović et al. [15]. Their matrices are not of Hessenberg type.

One feature that we noticed is that the computation of the right eigenvectors seems to be easier and faster than that of the left eigenvectors. The software that we used (Matlab and Maple) sometimes gives the left eigenvectors in a different order than the right eigenvectors so that one needs to be careful in pairing up the left and right eigenvectors. For nonsymmetric tridiagonal matrices Van Dooren et al. [29] gave some improvements for computing the left and right eigenvectors, and a similar idea might also be valuable for Hessenberg matrices. These authors already worked on the Hessenberg case in [16] and noticed that the total positivity of the matrices allows a stable and efficient way to compute the eigenvalues. The Hessenberg matrices in our examples §6.1 and §6.2 are indeed totally positive as they can be factored using bidiagonal matrices with positive entries, see Lima [17].

The Bessel weights in our example have the remarkable property that one knows the recurrence coefficients of the multiple orthogonal polynomials on the stepline exactly with simple formulas. This is in sharp contrast with the recurrence coefficients of the corresponding orthogonal polynomials, for which no expression is known, see [11] for weight functions related to the  $K_\nu$  Bessel function. This means that we don't need an additional step to compute the recurrence coefficients from the moments or modified moments of the measures, which is usually a badly conditioned problem. We can start right away from the known recurrence coefficients to build the Hessenberg matrix. In case one knows the nearest neighbor recurrence coefficients of the multiple orthogonal polynomials, one can compute the recurrence coefficients for the polynomials on the stepline using an algorithm of Filipuk et al. [10].

## 8 Declarations

### 8.1 Ethical approval

Not applicable.

### 8.2 Availability of supporting data

All the data are numerical results published in the paper. The algorithms are also in the paper.

### 8.3 Competing interests

Not applicable.

### 8.4 Funding

Supported by research project G0C9819N of FWO (Research Foundation – Flanders).

### 8.5 Author's contribution

This is a single author paper. The paper was written by the author and the numerical results were obtained by the author.

### 8.6 Acknowledgments

The author is grateful to the referees for useful comments and additional references.

## References

- [1] H. Alqahtani, L. Reichel, *Multiple orthogonal polynomials applied to matrix function evaluation*, BIT Numer. Math. **58** (2018), no. 4, 835–849.
- [2] A. Angelesco, *Sur deux extensions des fractions continues algébriques*, C. R. Acad. Sci. Paris **168** (1919), 262–265.
- [3] Y. Ben Cheikh, K. Douak, *On two-orthogonal polynomials related to the Bateman  $j_n^{u,v}$ -function*, Methods Appl. Anal. **7** (2000), 641–662.
- [4] C.F. Borges, *On a class of Gauss-like quadrature rules*, Numer. Math. **67** (1994), 271–288.
- [5] E. Coussement, W. Van Assche, *Multiple orthogonal polynomials associated with the modified Bessel functions of the first kind*, Constr. Approx. **19** (2003), 237–263.
- [6] J. Coussement, W. Van Assche, *Gaussian quadrature for multiple orthogonal polynomials*, J. Comput. Appl. Math. **178** (2005), 131–145.

- [7] E. Daems, A.B.J. Kuijlaars, *A Christoffel-Darboux formula for multiple orthogonal polynomials*, J. Approx. Theory **130** (2004), no. 2, 190–202.
- [8] K. Douak, *On 2-orthogonal polynomials of Laguerre type*, Internat. J. Math. & Math. Sci. **22** (1999), no. 1, 29–48.
- [9] U. Fidalgo Prieto, G. López Lagomasino, *Nikishin systems are perfect*, Constr. Approx. **34** (2011), no. 3, 297–356.
- [10] G. Filipuk, M. Haneczok, W. Van Assche, *Computing recurrence coefficients of multiple orthogonal polynomials* Numer. Algorithms **70** (2015), 519–543.
- [11] W. Gautschi, G.V. Milovanović, *Orthogonal polynomials relative to weight functions of Prudnikov type*, Numer. Algorithms **90** (2022), no. 1, 263–270.
- [12] G.H. Golub, J.H. Welsch, *Calculation of Gauss quadrature rules*, Math. Comp. **23** (1969), 221–230.
- [13] A.A. Gonchar, E. Rakhmanov, *On the convergence of simultaneous Padé approximants for systems of functions of Markov type*, (Russian), Trudy Mat. Inst. Steklov. **157** (1981), 31–48.
- [14] C. Hermite, *Sur la fonction exponentielle*, C.R. Acad. Sci. Paris **77** (1873), 18–24; 74–79; 226–233; 285–293.
- [15] A.N. Jovanović, M.P. Stanić, T.V. Tomović, *Construction of the optimal set of quadrature rules in the sense of Borges*, Electron. Trans. Numer. Anal. (ETNA) **50** (2018), 164–181.
- [16] T. Laudadio, N. Mastronardi, P. Van Dooren, *Computational aspects of simultaneous Gaussian quadrature*, manuscript
- [17] H. Lima, *Bidiagonal matrix factorizations associated with symmetric multiple orthogonal polynomials and lattice paths*, arXiv:2308.03561 [math.CA].
- [18] D.S. Lubinsky, W. Van Assche, *Simultaneous Gaussian quadrature for Angelesco systems*, Jaén Journal of Approximation **8** (2016), nr. 2, 113–149.
- [19] K. Mahler, *Perfect systems*, Compositio Math. **19** (1968), 95–166.
- [20] G.V. Milovanović, M. Stanić, *Construction of multiple orthogonal polynomials by discretized Stieltjes-Gautschi procedure and corresponding Gaussian quadratures*, Facta Univ. Ser. Math. Inform. **18** (2003), 9–29.
- [21] E.M. Nikishin, *A system of Markov functions* (Russian), Vestn. Mosk. Univ., Ser. I (1979) no. 4, 60–63; translation in Mosc. Univ. Math. Bull. **34** (1979), no. 4, 63–66.
- [22] H. Padé, *Sur la représentation approchée d’une fonction par des fractions rationnelles*, Thesis, Ann. École Nor. (3) **9** (1892), pp. 1–93.
- [23] T.V. Tomović, M.P. Stanić, *Construction of the optimal set of two or three quadrature rules in the sense of Borges*, Numer. Algorithms **78** (2018), no. 4, 1087–1109.

- [24] W. Van Assche, *Nearest neighbor recurrence relations for multiple orthogonal polynomials*, J. Approx. Theory **163** (2011), no. 10, 1427–1448.
- [25] W. Van Assche, E. Coussement, *Some classical multiple orthogonal polynomials*, Numerical Analysis 2000, Vol. V, Quadrature and Orthogonal Polynomials, J. Comput. Appl. Math. **127** (2001), no. 1–2, 317–347.
- [26] W. Van Assche, J.S. Geronimo, A.B.J. Kuijlaars, *Riemann-Hilbert problems for multiple orthogonal polynomials*, in ‘Special Functions 2000: Current Perspectives and Future Directions’ (J. Bustoz et al., eds.), NATO Science Series II. Mathematics, Physics and Chemistry **30**, Kluwer, Dordrecht, 2001, pp. 23–59.
- [27] W. Van Assche, A. Vuerinckx, *Multiple Hermite polynomials and simultaneous Gaussian quadrature*, Electron. Trans. Numer. Anal. **50** (2018), 182–198.
- [28] W. Van Assche, S.B. Yakubovich, *Multiple orthogonal polynomials associated with Macdonald functions*, Integral Transforms Spec. Funct. **9** (2000), no. 3, 229–244.
- [29] P. Van Dooren, T. Laudadio, N. Mastronardi, *Computing the eigenvectors of nonsymmetric tridiagonal matrices*, Comput. Math. Math. Phys. **61** (2001), no. 5, 733–749.

Walter Van Assche  
 Department of Mathematics  
 KU Leuven  
 Celestijnenlaan 200B, box 2400  
 BE-3001 Leuven, BELGIUM  
 walter.vanassche@kuleuven.be  
 orcid 0000-0003-3446-6936