

Chimeras in the two-community Kuramoto model with an external drive

Jens Grønberg

Abstract

We study the bifurcations of a special case of the Kuramoto model with two communities of oscillators and an external drive. We use Ott-Antonsen's ansatz to derive the low-dimensional system of differential equations that governs the macroscopic dynamics of the high-dimensional problem. The choice of parameters of the system is motivated by the search for so-called Chimera states; stable phase configurations with partial synchronization [10, 1]. Our main result is the derivation of the low-dimensional system following Ott-Antonsen's Ansatz and findings of periodic and chaotic Chimeras.

Introduction

The Kuramoto model has since its introduction in 1984 [9] found application in scientific fields with interacting oscillators dynamics. Oscillators appear in a wide range of areas from modeling Josephson Junctions [6, 16] used in constructing superconducting integrated circuit chips for quantum computing, to describing modern power grid configurations [5]. In ecology the Kuramoto model is used to model the interaction between pests found on coffee plants [15] and in neuroscience where neuron dynamics are characterized by chemical oscillators [4, 12, 2].

Ott and Antonsen's 2008 paper [13] marked a breakthrough in the study of Kuramoto-like models with a large number of oscillators. The paper discusses the original model as well as variations on it. They all have global dynamics governed by a low-dimensional system under a series of constraints on the initial configuration of the system. This paper is motivated by combining two of these variations [13, p.9-11]; the community generalization with an external drive

$$d\theta_{\sigma k}/dt = \omega_{\sigma k} + \sum_{\sigma'=1}^2 \frac{K_{\sigma\sigma'}}{N_{\sigma'}} \sum_{j=1}^{N_{\sigma'}} \sin(\theta_{\sigma'j} - \theta_{\sigma k} - \beta) + \Lambda \sin(\Omega t - \theta_{\sigma k}). \quad (1)$$

$\theta_{\sigma k}$ is the phase of an oscillator k belonging to the community σ . $\omega_{\sigma k}$ is its initial frequency, $K_{\sigma\sigma'}$ the coupling constants between communities σ and σ' and β the phase distortion factor. The external drive is characterized by a weight Λ and a frequency Ω . The phase distortion parameter β acts as a destabilizer by moving the fixed points of the coupling function away from the synchronized state enabling more complex dynamics [14].

Ott-Antonsen Ansatz

The continuum case of Kuramoto-like models where $N \rightarrow \infty$ has received much attention in parts due to its complexity and in part its applicability. Ott and Antonsen's paper [13] sets off by considering the oscillators from the point of view of statistical mechanics. The starting point is the

Vlasov equation which is a conservation model from Plasma Physics. It models the conservation of oscillators and discounts collision effects, the same conditions that underpin the Kuramoto model

$$\partial f_\sigma / \partial t + \partial / \partial \theta (f_\sigma \cdot v_\sigma) = 0. \quad (2)$$

The phase and frequency of each oscillator are modeled as a continuous random variable with density function $f_\sigma(\omega_\sigma, \theta_\sigma, t)$. The velocity v_σ term is given by the Kuramoto model. Ott and Antonsen proceeds by defining a so-called order parameter $r = 1/N \sum e^{i\theta}$ is introduced which is used to rewrite 1 as a single equation

$$d\theta_\sigma / dt = \omega_\sigma - \Omega + \frac{1}{2i} \sum_{\sigma'=1}^2 (K_{\sigma\sigma'} r_{\sigma'} e^{-i\beta} + \Lambda) e^{-i\theta_\sigma} - c.c., \quad (3)$$

where c.c. is the complex conjugate of the prior term. r encodes the synchronicity of a community of oscillators in its absolute value and is in the continuum case defined in terms of density function f

$$r_\sigma = \int_{-\infty}^{\infty} \int_0^{2\pi} f_\sigma e^{i\theta_\sigma} d\theta_\sigma d\omega_\sigma. \quad (4)$$

Following the steps from Ott and Antonsen [13] we expand the density f_σ as a Fourier series in the phase variable

$$f_\sigma = \frac{g_\sigma(\omega)}{2\pi} \left(1 + \sum_{n=1}^{\infty} f_{\sigma n}(\omega_\sigma, t) e^{in\theta_\sigma} + c.c. \right), \quad (5)$$

Ott-Antonsen chose the Fourier coefficients as a power series of a real function α_σ

$$f_{\sigma n}(\omega_\sigma, t) = (\alpha_\sigma(\omega_\sigma, t))^n. \quad (6)$$

With this special choice of coefficients, we insert 3 and 5 in 2

$$\begin{aligned} \partial f_\sigma / \partial t + \partial / \partial \theta (f_\sigma \cdot v_\sigma) &= \partial \alpha_\sigma / \partial t \sum_{n=1}^{\infty} n \alpha_\sigma^{n-1} e^{in\theta_\sigma} + c.c. \\ &\quad + (\omega_\sigma - \Omega) \sum_{n=1}^{\infty} i n \alpha_\sigma^n e^{in\theta_\sigma} + c.c. \\ &\quad + \left(\frac{1}{2i} \sum_{\sigma'=1}^2 (K_{\sigma\sigma'} r_{\sigma'} e^{-i\beta} + \Lambda) e^{-i\theta_\sigma} - c.c. \right) \sum_{n=1}^{\infty} i n \alpha_\sigma^n e^{in\theta_\sigma} + c.c. \\ &\quad - \left(\frac{1}{2} \sum_{\sigma'=1}^2 (K_{\sigma\sigma'} r_{\sigma'} e^{-i\beta} + \Lambda) e^{-i\theta_\sigma} + c.c. \right) \left(1 + \sum_{n=1}^{\infty} \alpha_\sigma^n e^{in\theta_\sigma} + c.c. \right) \\ &= \partial \alpha_\sigma / \partial t \sum_{n=1}^{\infty} n \alpha_\sigma^{n-1} e^{in\theta_\sigma} + c.c. \\ &\quad + (\omega_\sigma - \Omega) \sum_{n=1}^{\infty} i n \alpha_\sigma^n e^{in\theta_\sigma} + c.c. \\ &\quad + \frac{1}{2} \sum_{n=1}^{\infty} \sum_{\sigma'=1}^2 (K_{\sigma\sigma'} r_{\sigma'} e^{-i\beta} + \Lambda) (n-1) \alpha_\sigma^n e^{i(n-1)\theta_\sigma} - \sum_{\sigma'=1}^2 (K_{\sigma\sigma'} r_{\sigma'} e^{-i\beta} + \Lambda) (n+1) \alpha_\sigma^{n*} e^{-i(n+1)\theta_\sigma} \end{aligned}$$

$$\begin{aligned}
& - \sum_{\sigma'=1}^2 (K_{\sigma\sigma'} r_{\sigma'} e^{-i\beta} + \Lambda)^* (n+1) \alpha_{\sigma}^n e^{i(n+1)\theta_{\sigma}} + \sum_{\sigma'=1}^2 (K_{\sigma\sigma'} r_{\sigma'} e^{-i\beta} + \Lambda)^* (n-1) \alpha_{\sigma}^{n*} e^{-i(n-1)\theta_{\sigma}} \\
& \quad - \left(\frac{1}{2} \sum_{\sigma'=1}^2 (K_{\sigma\sigma'} r_{\sigma'} e^{-i\beta} + \Lambda) e^{-i\theta_{\sigma}} + c.c. \right) \\
& \quad = \partial\alpha_{\sigma}/\partial t \sum_{n=1}^{\infty} n \alpha_{\sigma}^{n-1} e^{in\theta_{\sigma}} + c.c. + (\omega_{\sigma} - \Omega) i \alpha_{\sigma} \sum_{n=1}^{\infty} n \alpha_{\sigma}^{n-1} e^{in\theta_{\sigma}} + c.c. \\
& \quad + \left(\frac{1}{2} \sum_{\sigma'=1}^2 \alpha_{\sigma}^2 (K_{\sigma\sigma'} r_{\sigma'} e^{-i\beta} + \Lambda) - (K_{\sigma\sigma'} r_{\sigma'} e^{-i\beta} + \Lambda)^* \right) \sum_{n=1}^{\infty} n \alpha_{\sigma}^{n-1} e^{in\theta_{\sigma}} + c.c. = 0 \quad (7)
\end{aligned}$$

We notice that surprisingly the θ -dependency can be factored out leaving

$$\partial\alpha_{\sigma}/\partial t + i(\omega_{\sigma} - \Omega)\alpha_{\sigma} + \frac{1}{2} \sum_{\sigma'=1}^2 (K_{\sigma\sigma'} r_{\sigma'} e^{-i\beta} + \Lambda) \alpha_{\sigma}^2 - (K_{\sigma\sigma'} r_{\sigma'} e^{-i\beta} + \Lambda)^* = 0. \quad (8)$$

Despite the simplification, eq. 4 and 8 which determines the dynamics of the global order parameter r is still an infinite-dimensional problem because α is a function. To address this choose the frequency density g to be a Lorentzian, $g_{\sigma}(\omega) = \Delta_{\sigma}/\pi((\omega_{\sigma} - \omega_{\sigma 0}^2) + \Delta_{\sigma}^2)^{-1}$, and the integrals in 4 have a closed analytic form

$$r_{\sigma} = \int_{-\infty}^{\infty} \int_0^{2\pi} f_{\sigma} e^{i\theta} d\theta d\omega_{\sigma} = \int_{-\infty}^{\infty} \alpha(\omega, t)^* g(\omega) d\omega_{\sigma} = \alpha_{\sigma}(\omega_{\sigma 0} - i\Delta_{\sigma}, t)^*. \quad (9)$$

With this result 8 simplifies to a set of complex ODEs

$$\partial r_{\sigma}/\partial t = \Delta_{\sigma} + i(\omega_{\sigma 0} - \Omega) r_{\sigma}^* + \frac{1}{2} \sum_{\sigma'=1}^2 (K_{\sigma\sigma'} r_{\sigma'}^* e^{-i\beta} + \Lambda) r_{\sigma}^2 - (K_{\sigma\sigma'} r_{\sigma'}^* e^{-i\beta} + \Lambda)^*. \quad (10)$$

Numerical analysis

Any numerical analysis of 10 will be local in nature and non-exhaustive. Our focus is to show that 1 has stable Chimera configurations in the continuum limit. We make parameter reductions in line with what can be found in related works [1, 3].

We simplify the search by choosing an initial value that is a Chimera. This is achieved by letting the first community of oscillators be synchronized, $|r_1| = 1$, and let the oscillators be identical by pinching the scale parameter $\Delta_{\sigma} \rightarrow 0$.

We only consider the case when the internal and external coupling constants are symmetric $K_{12} = K_{21}, K_{11} = K_{22}$. We normalize the two and define a coupling constant discrepancy parameter A , $K_{11} + K_{12} = 1, A \equiv K_{11} - K_{12}$. It is convenient to shift the Sakaguchi-parameter β such that $\beta \rightarrow \pi/2 - \beta$ and define a second discrepancy parameter between the frequencies of the external




Figure 1: Bifurcations in the (A, β) space for fixed parameters $\omega_1 = 0.203, \omega_2 = 0.9, \Lambda = 0.5$, dotted lines mark unstable bifurcations. The subdiagram shows how the original Hopf cycle loses stability in subsequent period doublings in the $(A, L^2 - norm)$ space.

drive and the oscillators initial frequency $\omega_\sigma \equiv \Omega - \omega_{0\sigma}$. These constraints reduces 10 to

$$\begin{aligned}
 dr/dt &= \frac{1-r^2}{4} ((1-A) \sin(\phi_1 - \phi_2 + \beta) + (1+A)r \sin(\beta) + 2\Lambda \cos(\phi_2)) \\
 d\phi_1/dt &= \omega_1 - \frac{1-A}{2} r \cos(\phi_2 - \phi_1 + \beta) - \frac{1+A}{2} \cos(\beta) - \Lambda \sin(\phi_1) \\
 d\phi_2/dt &= \omega_2 - \frac{1+r^2}{4r} ((1-A) \sin(\phi_1 - \phi_2 + \beta) + (1+A)r \cos(\beta) + 2\Lambda \sin(\phi_2)).
 \end{aligned} \tag{11}$$


where $r = |r_2|$ in order to simplify the notation.

Solutions to this system exist in a torus-shaped subset of \mathbb{R}^3 , Chimera states are found in the interior of the torus, as solutions on the surface represent a completely synchronized system with $r = 1$. Fixed points of 11 can be found in the interior and are the basis for bifurcation diagrams as seen in Figure 1.

We use the software packages auto-07p [7] and xppaut [8] to continue bifurcations and investigate state space solutions. Figure 1 shows a complicated cut in the (A, β) parameter space which we will discuss.

Reading the diagram in a clockwise fashion a subcritical Hopf bifurcation originates from the Bogdanov-Takens (BT) point. When β is decreased the Hopf line eventually intersects the saddle-node bifurcation in a Generalized-Hopf (GH) point and becomes supercritical. The now supercritical Hopf line intersects the saddle-node a second time, this time tangentially in a Zero-Hopf (ZH) point. We also observe that this marks the beginning of a stable saddle-node bifurcation.

A fixed point attractor can be found in the parameter space between the stable saddle-node line



(a) Shilnikov repeller after homoclinic bifurcation. (b) Stable chaotic attractor after a period-doubling cascade.

Figure 2: Chaotic solutions.

and the supercritical Hopf line. This solution is a Chimera state, as the second community for this set of parameter values is in a stable desynchronized state. The degree of desynchronization, as measured by r , is however constant.

When the discrepancy A between the interior and exterior coupling constants is increased the fixed point attractor undergoes a Hopf bifurcation resulting in an oscillating Chimera state. A further increase leads to a loss of stability in a period-doubling (PD) bifurcation. For a region of $\beta \in [0.1, 0.14]$ we observe a cascade of period-doublings resulting in a stable chaotic attractor. The magnified diagram in Figure 1 shows this stability transition for increasing A and fixed $\beta = 0.13$. A Solution on the chaotic attractor which appears after the second period-doubling is shown in figure 2b.

We continue the original cycle that lost stability in the first period-doubling by increasing A further, and eventually observe a rapidly increasing period indicating the presence of a homoclinic orbit. We use auto-07p's Homcont module to continue a high period cycle towards the Bogdonov-Takens point. Unfortunately, Homcont was not able to continue all the way leading to a convergence failure (MX) when the homoclinic closes in on the subcritical Hopf line.

Homcont can be configured to detect test functions equivalent to definitions found in [11, p. 528]. The saddle point to the green homoclinic line is of the type saddle-focus, and for $\beta \gtrsim 0.15$ the eigenvalues λ_1, μ_1, μ_2 have values such that $\lambda_1 + \text{Re}(\mu_1) + \text{Re}(\mu_2) = 0$. To the left of this neutrally-divergent saddle-focus point, the homoclinic is a single orbit, and to the right, it is a chaotic repeller

of the Shilnikov type. See Figure 2a.

This concludes our numerical findings. We have shown how the Kuramoto model with communities and an external drive defined in 1 can be analyzed using Ott and Antonsens Ansatz. Using their method we derived the low-dimensional system, and analyzed it hoping to find Chimera states. We found that the model supports all the different types of Chimeras, including chaotic ones which came about from period-doubling cascade.

References

- [1] Daniel M. Abrams and Steven H. Strogatz. “Chimera States for Coupled Oscillators”. In: *Phys. Rev. Lett.* 93 (17 Oct. 2004), p. 174102. DOI: 10.1103/PhysRevLett.93.174102. URL: <https://link.aps.org/doi/10.1103/PhysRevLett.93.174102>.
- [2] K. Anesiadis and A. Provata. “Synchronization in Multiplex Leaky Integrate-and-Fire Networks With Nonlocal Interactions”. In: *Frontiers in Network Physiology* 2 (2022). ISSN: 2674-0109. DOI: 10.3389/fnetp.2022.910862. URL: <https://www.frontiersin.org/articles/10.3389/fnetp.2022.910862>.
- [3] Christian Bick, Mark J. Panaggio, and Erik A. Martens. “Chaos in Kuramoto oscillator networks”. In: *Chaos: An Interdisciplinary Journal of Nonlinear Science* 28.7 (July 2018). DOI: 10.1063/1.5041444. URL: <https://doi.org/10.1063%2F1.5041444>.
- [4] Michael Breakspear, Stewart Heitmann, and Andreas Daffertshofer. “Generative Models of Cortical Oscillations: Neurobiological Implications of the Kuramoto Model”. In: *Frontiers in Human Neuroscience* 4 (2010). ISSN: 1662-5161. DOI: 10.3389/fnhum.2010.00190. URL: <https://www.frontiersin.org/articles/10.3389/fnhum.2010.00190>.
- [5] Anna Büttner and Frank Hellmann. *Complex Couplings – A universal, adaptive and bilinear formulation of power grid dynamics*. 2023. arXiv: 2308.15285 [nlin.AO].
- [6] B. C. Daniels, S. T. M. Dissanayake, and B. R. Trees. “Synchronization of coupled rotators: Josephson junction ladders and the locally coupled Kuramoto model”. In: *Phys. Rev. E* 67 (2 Feb. 2003), p. 026216. DOI: 10.1103/PhysRevE.67.026216. URL: <https://link.aps.org/doi/10.1103/PhysRevE.67.026216>.
- [7] Eusebius J. Doedel et al. *AUTO-07P: Continuation and bifurcation software for ordinary differential equations*. Tech. rep. 2007.
- [8] Bard Ermentrout. *XPPAUT5.41*. Tech. rep. 2003.
- [9] Y. Kuramoto. “Chemical Oscillations, Waves and Turbulence”. In: *Springer* (1984).
- [10] Yoshiki Kuramoto and Dorjsuren Battogtokh. *Coexistence of Coherence and Incoherence in Nonlocally Coupled Phase Oscillators*. 2002. arXiv: cond-mat/0210694 [cond-mat.stat-mech].
- [11] Yuri A. Kuznetsov. *Elements of applied bifurcation theory*. 2nd ed. Applied Mathematical Sciences 112. Berlin: Springer, 1998. ISBN: 978-0387983820.
- [12] Géza Ódor and Jeffrey Kelling. “Critical synchronization dynamics of the Kuramoto model on connectome and small world graphs”. In: *Scientific Reports* 9 (Dec. 2019), p. 19621. DOI: 10.1038/s41598-019-54769-9.
- [13] Edward Ott and Thomas M Antonsen. “Low dimensional behavior of large systems of globally coupled oscillators”. In: *Chaos: An Interdisciplinary Journal of Nonlinear Science* 3 (2008).
- [14] Y. Sakaguchi. “Cooperative Phenomena in Coupled Oscillator Systems under External Fields”. In: *Prog. Theor. Phys* 79 (1988).
- [15] John Vandermeer et al. “New forms of structure in ecosystems revealed with the Kuramoto model”. In: *Royal Society Open Science* 8.3 (2021), p. 210122. DOI: 10.1098/rsos.210122. eprint: <https://royalsocietypublishing.org/doi/pdf/10.1098/rsos.210122>. URL: <https://royalsocietypublishing.org/doi/abs/10.1098/rsos.210122>.

- [16] Kurt Wiesenfeld, Pere Colet, and Steven H. Strogatz. “Synchronization Transitions in a Disordered Josephson Series Array”. In: *Phys. Rev. Lett.* 76 (3 Jan. 1996), pp. 404–407. DOI: 10.1103/PhysRevLett.76.404. URL: <https://link.aps.org/doi/10.1103/PhysRevLett.76.404>.