

An alternative evaluation of the leading-order hadronic contribution to the muon $g-2$ with MUonE

Fedor Ignatov^a, Riccardo Nunzio Pilato^a, Thomas Teubner^a, Graziano Venanzoni^{a,b}

^aUniversity of Liverpool, Foundation Building, Brownlow Hill, L69 3BX, Liverpool, United Kingdom

^bINFN Sezione di Pisa, Largo Bruno Pontecorvo 3, 56127, Pisa, Italy

Abstract

We propose an alternative method to extract the leading-order hadronic contribution to the muon $g-2$, a_μ^{HLO} , with the MUonE experiment. In contrast to the traditional method based on the integral of the hadronic contribution to the running of the effective fine-structure constant $\Delta\alpha_{\text{had}}$ in the space-like region, our approach relies on the computation of the derivatives of $\Delta\alpha_{\text{had}}(t)$ at zero squared momentum transfer t . We show that this approach allows to extract $\sim 99\%$ of the total value of a_μ^{HLO} from the MUonE data, while the remaining $\sim 1\%$ can be computed combining perturbative QCD and data on e^+e^- annihilation to hadrons. This leads to a competitive evaluation of a_μ^{HLO} which is robust against the parameterization used to model $\Delta\alpha_{\text{had}}(t)$ in the MUonE kinematic region, thanks to the analyticity properties of $\Delta\alpha_{\text{had}}(t)$, which can be expanded as a polynomial at $t \sim 0$.

Keywords: muon $g-2$, MUonE

1. Introduction

The muon anomalous magnetic moment, also known as the muon $g-2$, where g is the muon gyromagnetic ratio, exhibits a discrepancy between theory and experiment which persists for more than 20 years. It has received renewed interest, following the first measurement of the muon anomaly $a_\mu = (g-2)/2$ by the Muon $g-2$ Experiment at Fermilab [1], subsequently confirmed by the new result with a twofold improved precision [2]. The comparison with the Standard Model (SM) prediction a_μ^{SM} [3] is currently limited by the present tensions in the evaluation of the leading-order hadronic contribution to the muon anomaly, a_μ^{HLO} [4]. This term represents the main source of uncertainty of the theory prediction, due to the non-perturbative nature of QCD at low energy. A recent computation of a_μ^{HLO} based on lattice QCD, performed by the BMW Collaboration [5], indeed shows a 2.2σ tension with the one used in the SM evaluation of a_μ [3], which is instead based on a data-driven approach involving data for $e^+e^- \rightarrow$ hadrons cross sections. Moreover, a new experimental measurement of $e^+e^- \rightarrow \pi^+\pi^-$ channel from the CMD-3 experiment disagrees with the previous measurements [6].

New calculations from other lattice QCD groups and new results from e^+e^- colliders are expected to shed light on these tensions in the next few years [4]. Recently a new approach has been proposed to compute a_μ^{HLO} , based on the measurement of the hadronic contribution to the running of the electromagnetic coupling $\Delta\alpha_{\text{had}}$ in the space-like region [7]. The elastic scattering of high-energy muons on atomic electrons has been identified as an ideal process for this measurement and an experimental proposal, called MUonE, has been put forward at CERN to extract $\Delta\alpha_{\text{had}}$ from a precise measurement of the shape of the $\mu^+e^- \rightarrow \mu^+e^-$ elastic process [8]. The goal of MUonE is to determine a_μ^{HLO} with a $\sim 0.3\%$ statistical and a comparable systematic uncertainty, using the following integral [9]:

$$a_\mu^{\text{HLO}} = \frac{\alpha}{\pi} \int_0^1 dx (1-x) \Delta\alpha_{\text{had}}[t(x)], \quad t(x) = \frac{x^2 m_\mu^2}{x-1} < 0, \quad (1)$$

Email addresses: f.ignatov@liverpool.ac.uk (Fedor Ignatov), r.pilato@liverpool.ac.uk (Riccardo Nunzio Pilato), teubner@liverpool.ac.uk (Thomas Teubner), graziano.venanzoni@liverpool.ac.uk (Graziano Venanzoni)

where α is the fine structure constant, m_μ is the muon mass, and t is the space-like squared momentum transfer. The 160 GeV muon beam available at the M2 beamline at CERN allows to cover directly the momentum transfer range $-0.153 \text{ GeV}^2 < t < -0.001 \text{ GeV}^2$, which is equivalent to $0.258 < x < 0.936$. This corresponds to $\sim 86\%$ of the integral in Eq. 1, while the remaining fraction can be obtained by extrapolating $\Delta\alpha_{had}(t)$ outside the MUonE region by an appropriate analytical function or alternatively using lattice data. In the first case the space-like integral of Eq. 1 is sensitive to the behaviour of the parameterization chosen to model $\Delta\alpha_{had}(t)$ in the whole t -region, particularly in the asymptotic limit $t \rightarrow -\infty$, which could affect the extraction of a_μ^{HLO} ¹. In the following, we will discuss a different approach based on the evaluation of the derivatives of $\Delta\alpha_{had}(t)$ at zero momentum transfer. This leads to an evaluation of a_μ^{HLO} which is rather insensitive to the functional form adopted to describe the behaviour of $\Delta\alpha_{had}(t)$ and will provide an alternative and competitive way to determine a_μ^{HLO} with MUonE.

2. Description of the method

The new method is mainly based on [11, 12], where different quark flavours have been treated separately, and the dominant light-quark contributions to the hadronic vacuum polarization function $\Pi_{had}(s)$ have been computed either through a model-dependent approach [11], or using lattice QCD calculations [12]. The same strategy is not applicable to MUonE, since MUonE will provide an inclusive measurement of $\Pi_{had}(t)$ containing contributions from all the quark flavours. In the following, we summarize the original procedure and describe how it can be adapted to the MUonE case. We start from the well known dispersion relation [13–16]

$$a_\mu^{\text{HLO}} = \frac{\alpha^2}{3\pi^2} \int_{s_{\text{th}}}^{\infty} \frac{ds}{s} K(s) R(s), \quad (2)$$

where, due to contributions from the $\pi^0\gamma$ channel, the threshold s_{th} is usually identified as $m_{\pi^0}^2$, with m_{π^0} being the π^0 mass. The kernel function $K(s)$ is given by

$$K(s) = \int_0^1 dx \frac{x^2(1-x)}{x^2 + \frac{s}{m_\mu^2}(1-x)}, \quad (3)$$

and $R(s)$ is the ratio of the bare (i.e. excluding vacuum polarization effects) $e^+e^- \rightarrow$ hadrons annihilation cross section to the Born $e^+e^- \rightarrow \mu^+\mu^-$ pointlike one, $\sigma_{pt} = 4\pi\alpha^2/(3s)$. Taking advantage of the optical theorem, $R(s)$ can be related to the imaginary part of the hadronic vacuum polarization function $\Pi_{had}(s)$ as follows:

$$-\text{Im}\Pi_{had}(s) = \frac{\alpha}{3} R(s). \quad (4)$$

The dispersive integral in Eq. 2 can be split in two terms at a given value s_0 , above which $R(s)$ can be safely computed using perturbative QCD (pQCD). As originally proposed in [11]², it is convenient to approximate $K(s)$ by a meromorphic function $K_1(s)$ for $s \leq s_0$,

$$K_1(s) = c_0 s + \sum_{n=1}^3 \frac{c_n}{s^n}, \quad (5)$$

and the low energy part of the dispersive integral can be written as

$$-\frac{\alpha}{\pi} \int_{s_{\text{th}}}^{s_0} \frac{ds}{s} K(s) \frac{\text{Im}\Pi_{had}(s)}{\pi} = -\frac{\alpha}{\pi} \left[\int_{s_{\text{th}}}^{s_0} \frac{ds}{s} [K(s) - K_1(s)] \frac{\text{Im}\Pi_{had}(s)}{\pi} + \int_{s_{\text{th}}}^{s_0} \frac{ds}{s} K_1(s) \frac{\text{Im}\Pi_{had}(s)}{\pi} \right], \quad (6)$$

¹A convenient choice based on the analytic formula for the QED leading-order contribution to the running of α in the space-like region allows to compute a_μ^{HLO} at the required level of precision with negligible bias when time-like data are used as input [10] (see also Section 3).

²Different approaches to evaluate the dispersive integral by an approximate kernel function have been discussed in [17, 18].

where Eq. 4 was used to express $R(s)$ in terms of the hadronic vacuum polarization function. Cauchy's theorem can then be employed to handle the second term on the right-hand side [11, 12]:

$$\int_{s_{\text{th}}}^{s_0} \frac{ds}{s} K_1(s) \frac{\text{Im}\Pi_{had}(s)}{\pi} = \text{Res} \left[\Pi_{had}(s) \frac{K_1(s)}{s} \right]_{s=0} - \frac{1}{2\pi i} \oint_{|s|=s_0} \frac{ds}{s} K_1(s) \Pi_{had}(s) \Big|_{\text{pQCD}}. \quad (7)$$

Here, the contour integral around the circle of radius s_0 can be calculated using pQCD to evaluate the hadronic vacuum polarization function, whereas the residual can be written in terms of derivatives of $\Pi_{had}(s)$ at zero momentum transfer. Exploiting the functional form of the approximated kernel in Eq. 5,

$$\text{Res} \left[\Pi_{had}(s) \frac{K_1(s)}{s} \right]_{s=0} = \sum_{n=1}^3 \frac{c_n}{n!} \frac{d^{(n)}}{ds^n} \Pi_{had}(s) \Big|_{s=0} = \sum_{n=1}^3 \frac{c_n}{n!} \frac{d^{(n)}}{dt^n} \Delta\alpha_{had}(t) \Big|_{t=0}, \quad (8)$$

where the analyticity of Π_{had} and its derivatives at zero momentum transfer has been exploited to move the evaluation of the hadronic vacuum polarization from positive to negative momentum transfer. The relation $\Delta\alpha_{had}(t) = \text{Re}\Pi_{had}(t)$ has been used in the last step to link the hadronic vacuum polarization function to the hadronic contribution to the running of α .

The high energy tail of the dispersive integral can be treated in a similar way:

$$-\frac{\alpha}{\pi} \int_{s_0}^{\infty} \frac{ds}{s} K(s) \frac{\text{Im}\Pi_{had}(s)}{\pi} = -\frac{\alpha}{\pi} \left[\int_{s_0}^{\infty} \frac{ds}{s} [K(s) - \tilde{K}_1(s)] \frac{\text{Im}\Pi_{had}(s)}{\pi} + \int_{s_0}^{\infty} \frac{ds}{s} \tilde{K}_1(s) \frac{\text{Im}\Pi_{had}(s)}{\pi} \right], \quad (9)$$

where $\tilde{K}_1(s) = K_1(s) - c_0 s$. Following the same technique implemented for the low energy component, Cauchy's theorem can be applied to the second integral on the right hand side of Eq. 9, using the red closed path shown in Fig. 1. In this case, the integrand is free of poles and the contour integral over s with radius $|s| \rightarrow \infty$ is vanishing. This leads to

$$\int_{s_0}^{\infty} \frac{ds}{s} \tilde{K}_1(s) \frac{\text{Im}\Pi_{had}(s)}{\pi} = \frac{1}{2\pi i} \oint_{|s|=s_0} \frac{ds}{s} \tilde{K}_1(s) \Pi_{had}(s) \Big|_{\text{pQCD}}. \quad (10)$$




Figure 1: Blue (red): closed path in the complex s -plane used to calculate the contour integral in Eq. 7 (Eq. 10).

Rearranging Eqs. 6, 7, 9 and 10, a_μ^{HLO} can be calculated as the sum of four terms

$$a_\mu^{\text{HLO}} = a_\mu^{\text{HLO (I)}} + a_\mu^{\text{HLO (II)}} + a_\mu^{\text{HLO (III)}} + a_\mu^{\text{HLO (IV)}}, \quad (11)$$

where

$$a_{\mu}^{\text{HLO (I)}} = -\frac{\alpha}{\pi} \sum_{n=1}^3 \frac{c_n}{n!} \frac{d^{(n)}}{dt^n} \Delta\alpha_{had}(t) \Big|_{t=0}, \quad (12)$$

$$a_{\mu}^{\text{HLO (II)}} = \frac{\alpha}{\pi} \frac{1}{2\pi i} \oint_{|s|=s_0} \frac{ds}{s} c_0 s \Pi_{had}(s) \Big|_{\text{pQCD}}, \quad (13)$$

$$a_{\mu}^{\text{HLO (III)}} = \frac{\alpha^2}{3\pi^2} \int_{s_{\text{th}}}^{s_0} \frac{ds}{s} [K(s) - K_1(s)] R(s), \quad (14)$$

$$a_{\mu}^{\text{HLO (IV)}} = \frac{\alpha^2}{3\pi^2} \int_{s_0}^{\infty} \frac{ds}{s} [K(s) - \tilde{K}_1(s)] R(s). \quad (15)$$

$a_{\mu}^{\text{HLO (I)}}$ will be computed using MUonE data. This term represents $\sim 99\%$ of the total value of a_{μ}^{HLO} , as will be shown in the following. The other three terms contribute to the remaining $\sim 1\%$. $a_{\mu}^{\text{HLO (II)}}$ will be calculated via pQCD, $a_{\mu}^{\text{HLO (III)}}$ with e^+e^- data, while both e^+e^- data and pQCD will be used to compute $a_{\mu}^{\text{HLO (IV)}}$.

In the following sections, the calculation of the different contributions will be discussed in details. The robustness of the proposed method will be tested using three different values of s_0 : $(1.8 \text{ GeV})^2$, $(2.5 \text{ GeV})^2$ and $(12 \text{ GeV})^2$. Two different techniques will be used to determine the coefficients of $K_1(s)$. The first (called *Minimization 1*) consists in the least squares minimization of the difference of the approximated and the analytical kernel. The second (*Minimization 2*) is aimed at minimizing the contribution of e^+e^- data in the calculation of a_{μ}^{HLO} , and is carried out by fixing the coefficient c_3 to be 1/2 of its value obtained from *Minimization 1*, and then minimizing $\int_{s_{\text{th}}}^{s_0} \frac{ds}{s} |K(s) - K_1(s)| R(s)$ to find the other coefficients. Table 1 shows the coefficients for the two minimizations. Both methods provide a good approximation of $K(s)$ with a different sensitivity on the third derivative $\frac{d^3 \Delta\alpha_{had}(t)}{dt^3}$ at $t = 0$ (which will be shown to have the largest uncertainty, see Table 2).

Coefficients	Minimization 1			Minimization 2		
	$(1.8 \text{ GeV})^2$	$(2.5 \text{ GeV})^2$	$(12 \text{ GeV})^2$	$(1.8 \text{ GeV})^2$	$(2.5 \text{ GeV})^2$	$(12 \text{ GeV})^2$
$c_0 \cdot 10^5$	2.206	0.7326	0.002164	2.419	1.011	0.003743
$c_1 \cdot 10^3$	3.486	3.512	3.555	3.482	3.494	3.520
$c_2 \cdot 10^4$	-1.484	-1.554	-1.684	-1.402	-1.443	-1.564
$c_3 \cdot 10^6$	4.869	5.294	6.128	2.434	2.647	3.064

Table 1: Coefficients of the approximated kernel function $K_1(s)$ for the two minimizations and the three choices of s_0 .

3. Evaluation of $a_{\mu}^{\text{HLO (I)}}$

The first term $a_{\mu}^{\text{HLO (I)}}$ in Eq. 11 depends on the derivatives of $\Delta\alpha_{had}(t)$ at zero momentum transfer which can be obtained by fitting the MUonE data with a convenient functional form. Since space-like data in the MUonE range are not available, we use different data compilations of $\Delta\alpha_{had}(t)$ obtained from e^+e^- data in the time-like region using the dispersive integral:

$$\Delta\alpha_{had}(q^2) = -\frac{\alpha}{3\pi} q^2 \int_{s_{\text{th}}}^{\infty} ds \frac{R(s)}{s(s-q^2)} \quad (16)$$

where q^2 is a generic momentum transfer. The data compilations produced in [19], [20] and [21], indicated respectively as Dataset I, II and III will be used. Figure 2 represents $\Delta\alpha_{had}(q^2)$ computed from the three compilations both for positive and negative momentum transfer. The small deviations between the three compilations are due to different strategies adopted in combining data from various experiments. The influence of the hadronic resonances on the vacuum polarization effects is evident for positive momentum transfer, whereas $\Delta\alpha_{had}(q^2)$ is a smooth function for $q^2 < 0$.

To assess the sensitivity of our method on the parameterization adopted for $\Delta\alpha_{had}(t)$, we tested several possible parameterizations which are capable of reproducing the behaviour of $\Delta\alpha_{had}(t)$ in the MUonE kinematic range:




Figure 2: $\Delta\alpha_{had}$ as a function of $\sqrt{q^2}$ computed using Dataset I (blue) [19], II (red) [20] and III (green) [21]. The insertion shows the behaviour of $\Delta\alpha_{had}$ in the MUonE kinematic range.

1. "Lepton Like" (LL) parameterization. It is motivated by the one-loop QED calculation of vacuum polarisation induced by a lepton pair in the space-like region, and is also used in MUonE to calculate a_μ^{HLO} through the space-like integral in Eq. 1 [10]:

$$\Delta\alpha_{had}(t) = KM \left[-\frac{5}{9} - \frac{4M}{3t} + \left(\frac{4M^2}{3t^2} + \frac{M}{3t} - \frac{1}{6} \right) \frac{2}{\sqrt{1 - \frac{4M}{t}}} \log \left| \frac{1 - \sqrt{1 - \frac{4M}{t}}}{1 + \sqrt{1 - \frac{4M}{t}}} \right| \right], \quad (17)$$

where M is the squared lepton mass and $K = \alpha/(\pi M)$ in the leptonic case. On the contrary, these two parameters do not have a precise physics interpretation when modeling the hadronic running, since $\Delta\alpha_{had}(t)$ is not calculable in perturbation theory. In the limit of very small t , the LL function reduces to the following expansion:

$$\Delta\alpha_{had}(t) = -K \left(\frac{t}{15} + \frac{t^2}{140M} + \frac{t^3}{945M^2} + O[t^4] \right) \quad (18)$$

which corresponds to the dominant behaviour in the MUonE kinematic range. The first three derivatives at zero momentum transfer are:

$$D1 = \left. \frac{d\Delta\alpha_{had}(t)}{dt} \right|_{t=0} = -\frac{K}{15}; \quad D2 = \left. \frac{d^2\Delta\alpha_{had}(t)}{dt^2} \right|_{t=0} = -\frac{K}{70M}; \quad D3 = \left. \frac{d^3\Delta\alpha_{had}(t)}{dt^3} \right|_{t=0} = -\frac{2K}{315M^2}$$

2. A Padé approximant (Padé parameterization) with three free parameters $\{P_i\}_{i=1}^3$:

$$\Delta\alpha_{had}(t) = P_1 t \frac{1 + P_2 t}{1 + P_3 t} \quad (19)$$

The expansion in the limit of very small t gives the following expression:

$$\Delta\alpha_{had}(t) = P_1 t + P_1(P_2 - P_3)t^2 + P_1 P_3(-P_2 + P_3)t^3 + O[t^4], \quad (20)$$

with the first three derivatives at zero momentum transfer:

$$D1 = P_1; \quad D2 = 2P_1(P_2 - P_3); \quad D3 = -6P_1 P_3(P_2 - P_3)$$

3. A third order polynomial (Pol parameterization) with three free parameters $\{P_i\}_{i=1}^3$. Since $\Delta\alpha_{had}(t=0) = 0$, the P_0 term is fixed to be zero:

$$\Delta\alpha_{had}(t) = P_1 t + P_2 t^2 + P_3 t^3 \quad (21)$$

which gives the following derivatives:

$$D1 = P_1; \quad D2 = 2P_2; \quad D3 = 6P_3$$

4. "Reconstruction Approximants" (GdR) parameterization [22]. It is based on the analytic properties of the hadronic vacuum polarization function $\Pi_{had}(t)$ in QCD, which takes the form [22]:

$$\Delta\alpha_{had}(t) = \sum_{n=1}^N \mathcal{A}(n, L) \left(\frac{\sqrt{1 - \frac{t}{t_0}} - 1}{\sqrt{1 - \frac{t}{t_0}} + 1} \right)^n + \sum_{p=1}^{\lfloor \frac{L+1}{2} \rfloor} \mathcal{B}(2p-1) \text{Li}_{2p-1} \left(\frac{\sqrt{1 - \frac{t}{t_0}} - 1}{\sqrt{1 - \frac{t}{t_0}} + 1} \right) \quad (22)$$

where $\mathcal{A}(n, L)$ and $\mathcal{B}(2p-1)$ are free parameters which can be constrained by the theory [22], t_0 is an energy scale (set in this work to $4m_{\pi^\pm}^2$) and Li_{2p-1} are polylogs of $2p-1$ degree. In the following, we limit the expansion to the case with $L=1, N=3$:

$$\Delta\alpha_{had}(t) = A_1 \mathcal{S}_1 + A_2 \mathcal{S}_2 + A_3 \mathcal{S}_3 + B_1 \mathcal{L}_1 \quad (23)$$

where

$$\begin{aligned} \mathcal{S}_i &= \left(\frac{\sqrt{1 - \frac{t}{t_0}} - 1}{\sqrt{1 - \frac{t}{t_0}} + 1} \right)^i; & A_i &= \mathcal{A}(i, 1) \quad i = 1, 2, 3 \\ \mathcal{L}_1 &= \text{Li}_1 \left(\frac{\sqrt{1 - \frac{t}{t_0}} - 1}{\sqrt{1 - \frac{t}{t_0}} + 1} \right); & B_1 &= \mathcal{B}(1) \end{aligned}$$

In the limit of small momentum transfer, the GdR function can be approximated by:

$$\Delta\alpha_{had}(t) = -\frac{(A_1 + B_1)t}{4t_0} - \frac{(4A_1 - 2A_2 + 3B_1)t^2}{32t_0^2} - \frac{[3(5A_1 - 4A_2 + A_3) + 10B_1]t^3}{192t_0^3} + O[t^4]$$

The first derivatives at zero momentum transfer can be computed as:

$$D1 = -\frac{A_1 + B_1}{4t_0}; \quad D2 = \frac{-4A_1 + 2A_2 - 3B_1}{16t_0^2}; \quad D3 = \frac{-15A_1 + 12A_2 - 3A_3 - 10B_1}{32t_0^3}$$

Five different variants of this parameterization have been considered:

- GdR1: $\Delta\alpha_{had}(t) = A_1 \mathcal{S}_1 + B_1 \mathcal{L}_1$, where A_1 is a free parameter and B_1 is constrained to its 6 quarks asymptotic freedom value: $B_1 = 2 \frac{\alpha}{\pi} \frac{5}{3} = 0.00774273$ [22];
- GdR2: $\Delta\alpha_{had}(t) = A_1 \mathcal{S}_1 + B_1 \mathcal{L}_1$, where A_1 and B_1 are free parameters;
- GdR3: $\Delta\alpha_{had}(t) = A_1 \mathcal{S}_1 + A_2 \mathcal{S}_2 + B_1 \mathcal{L}_1$ where A_1, A_2 are free parameters and B_1 is constrained to its 6 quarks asymptotic freedom value;
- GdR4: $\Delta\alpha_{had}(t) = A_1 \mathcal{S}_1 + A_2 \mathcal{S}_2 + A_3 \mathcal{S}_3 + B_1 \mathcal{L}_1$ where A_1, A_2 are free parameters, A_3 is constrained to $A_3 = (2A_2 - A_1 - B/2)/3$ [22] and B_1 is constrained to its 6 quarks asymptotic freedom value;
- GdR5: $\Delta\alpha_{had}(t) = A_1 \mathcal{S}_1 + A_2 \mathcal{S}_2 + A_3 \mathcal{S}_3 + B_1 \mathcal{L}_1$ where A_1, A_2, A_3 are free parameters and B_1 is constrained to its 6 quarks asymptotic freedom value.

For each $\Delta\alpha_{had}(t)$ compilation, 10^4 pseudo-experiments were simulated in the MUonE momentum transfer region $-0.153 \text{ GeV}^2 < t < -0.001 \text{ GeV}^2$. The higher limit is needed to reproduce the geometric acceptance of MUonE, which will include all the elastic events where the electron angle is $< 32 \text{ mrad}$. Statistical fluctuations have been

added independently to each $\Delta\alpha_{had}(t)$ point according to the full integrated luminosity of MUonE, 15 fb^{-1} [8]. A χ^2 fit has been performed on each pseudo-experiment using all the parameterizations described above, and for each iteration both the values of a_μ^{HLO} computed according to Eq. 1 and $a_\mu^{\text{HLO (I)}}$ were calculated. In particular, $a_\mu^{\text{HLO (I)}}$ was calculated for the three choices of s_0 and for the two minimizations used to determine the approximated kernel $K_1(s)$. This allowed to test extensively the robustness of the proposed method.

As an example, Fig.3, Left, shows the fit results of few parameterizations of $\Delta\alpha_{had}(t)$ for a given pseudo-experiment generated with Dataset I. All the parameterizations describe well the MUonE simulated data. However, differences arise outside the MUonE experimental range, as shown in Fig.3, Right, which represents the integrand function in Eq.1. In particular, the Padé approximant and the third order polynomial fail to describe $\Delta\alpha_{had}(t)$ for $x \rightarrow 1$, which corresponds to $t \rightarrow -\infty$. As a consequence of this, values of a_μ^{HLO} computed from the integral in Eq. 1 using these two functions will not be in agreement with the expected (input) value. On the other hand, the results of the three derivatives are quite stable, as shown in Table 2, which compares the results of the three derivatives for all the different parameterizations used and the three $\Delta\alpha_{had}(t)$ compilations. These values must be compared with the ones obtained by the corresponding dispersive integrals obtained directly from the timelike data: $D1 = (-9.24 \pm 0.04) \cdot 10^{-3}$, $D2 = (-3.86 \pm 0.02) \cdot 10^{-2}$, $D3 = (-3.90 \pm 0.03) \cdot 10^{-1}$. Table 3 shows the values of $a_\mu^{\text{HLO (I)}}$ for the different parameterizations and different s_0 values for the two minimization using Eq.13 with Dataset I. As can be seen the results are quite stable independently by the parameterization used. This is a consequence of the analyticity of the hadronic vacuum polarization function $\Pi_{had}(t)$ which can be approximated by a polynomial at $t \sim 0$ (as can be obtained by a Taylor expansion of $\Delta\alpha_{had}(t)$ at $t \sim 0$ using Eq. 16).




Figure 3: Left: example of fit results for few parameterizations of $\Delta\alpha_{had}(t)$. The insertion shows the fit residuals for the LL best fit on the same pseudo-experiment. Right: integrand function of the MUonE integral in Eq. 1 computed using different different parameterizations fitted to the same pseudo-experiment. The insertion zooms on the kinematic range not accessible to MUonE. Note that the Padé approximant fails to follow the expected behaviour, while the third order polynomial diverges for $x \rightarrow 1$ ($t \rightarrow -\infty$).

4. Evaluation of $a_\mu^{\text{HLO (II, III, IV)}}$

The contour integral in $a_\mu^{\text{HLO (II)}}$ was calculated using the pQCD prediction of $\Pi_{had}(s)$. The QCD vector correlator Π_{QCD} , which is related to the vacuum polarization function via $\Pi_{had}(s) = -12/3\alpha\pi(\sum q_i^2)\Pi_{\text{QCD}}(s)$, is known up to five loops in the massless approximation, i.e. $O(\alpha_s^4)$ [23]. The mass terms up to $O(\alpha_s^2(m^2/q^2)^{30})$ are taken from [24], while the mass terms up to $O(\alpha_s^3(m^2/q^2))$ are taken from [25]. The strong coupling $\alpha_s(\mu^2)$ was determined from the PDG average for $\alpha_s(M_Z^2)$ using the CRunDec program [26]. The uncertainty of the contour integral was estimated including contributions from the uncertainty of the input parameter $\alpha_s(M_Z^2)$, the variation of the renormalisation scale μ^2 in the range from $s_0/2$ to $2s_0$ and the full value of the estimated duality violations [27, 28]. These three terms have been added in quadrature to estimate the total uncertainty on the contour integral. Note that for $s_0 = (1.8\text{ GeV})^2$, the error is dominated by the estimate of the duality violations, which amount to 1%, while the uncertainties from α_s and the scale variation are much smaller and of similar size. For $s_0 = (2.5\text{ GeV})^2$, duality violations are already suppressed, and they contribute at the level of 0.1%. They are negligible at $s_0 = (12\text{ GeV})^2$, where the error is dominated by the

Parameterization	Values	Dataset		
		I	II	III
LL	D1	-9.23 ± 0.05	-9.17 ± 0.05	-9.20 ± 0.05
	D2	-3.76 ± 0.22	-3.77 ± 0.22	-3.73 ± 0.22
	D3	-3.19 ± 0.35	-3.21 ± 0.35	-3.14 ± 0.35
Padé	D1	-9.24 ± 0.08	-9.18 ± 0.08	-9.20 ± 0.08
	D2	-3.81 ± 0.62	-3.81 ± 0.62	-3.75 ± 0.61
	D3	-2.14 ± 2.42	-2.21 ± 2.41	-2.03 ± 2.42
Pol	D1	-9.22 ± 0.07	-9.16 ± 0.07	-9.19 ± 0.07
	D2	-3.55 ± 0.45	-3.56 ± 0.45	-3.50 ± 0.44
	D3	-1.83 ± 0.89	-1.84 ± 0.88	-1.79 ± 0.88
GdR1	D1	-9.26 ± 0.03	-9.19 ± 0.03	-9.23 ± 0.03
	D2	-3.92 ± 0.04	-3.82 ± 0.04	-3.88 ± 0.04
	D3	-3.03 ± 0.10	-2.80 ± 0.10	-2.93 ± 0.10
GdR2	D1	-9.23 ± 0.06	-9.18 ± 0.06	-9.20 ± 0.06
	D2	-3.73 ± 0.31	-3.75 ± 0.31	-3.69 ± 0.31
	D3	-2.46 ± 0.94	-2.58 ± 0.94	-2.33 ± 0.95
GdR3	D1	-9.23 ± 0.06	-9.18 ± 0.06	-9.20 ± 0.06
	D2	-3.70 ± 0.36	-3.74 ± 0.36	-3.65 ± 0.36
	D3	-2.25 ± 1.30	-2.47 ± 1.31	-2.13 ± 1.29
GdR4	D1	-9.24 ± 0.06	-9.19 ± 0.06	-9.21 ± 0.06
	D2	-3.89 ± 0.31	-3.91 ± 0.32	-3.84 ± 0.32
	D3	-2.95 ± 1.11	-3.14 ± 1.11	-2.84 ± 1.11
GdR5	D1	-9.23 ± 0.09	-9.18 ± 0.09	-9.20 ± 0.09
	D2	-3.74 ± 1.23	-3.75 ± 1.23	-3.64 ± 1.24
	D3	-2.90 ± 3.23	-2.91 ± 3.37	-1.80 ± 4.52

Table 2: Values of the first three derivatives of $\Delta\alpha_{had}(t)$ at zero momentum transfer for all the parameterization considered in this study and the three different datasets. D1 is given in units of 10^{-3} , D2 in units of 10^{-2} , and D3 in units of 10^{-1} .

Minimization I		$a_\mu^{\text{HLO (I)}} (10^{-10})$						
s_0 values	LL	Padé	Pol	GdR1	GdR2	GdR3	GdR4	GdR5
$(1.8 \text{ GeV})^2$	688.7 ± 2.2	688.7 ± 2.9	688.9 ± 2.9	688.2 ± 2.2	688.0 ± 2.2	688.0 ± 2.2	687.0 ± 2.3	688.0 ± 2.6
$(2.5 \text{ GeV})^2$	691.7 ± 2.2	691.6 ± 3.0	691.8 ± 3.0	691.0 ± 2.2	690.8 ± 2.2	690.8 ± 2.2	689.8 ± 2.3	690.9 ± 2.9
$(12 \text{ GeV})^2$	696.3 ± 2.2	696.3 ± 3.0	696.3 ± 3.2	695.4 ± 2.2	695.3 ± 2.2	695.2 ± 2.2	694.1 ± 2.3	695.3 ± 3.7
Minimization II		$a_\mu^{\text{HLO (I)}} (10^{-10})$						
s_0 values	LL	Padé	Pol	GdR1	GdR2	GdR3	GdR4	GdR5
$(1.8 \text{ GeV})^2$	688.5 ± 2.2	688.1 ± 4.2	689.8 ± 3.3	688.3 ± 2.1	688.4 ± 2.1	688.6 ± 2.2	687.1 ± 2.1	688.4 ± 5.8
$(2.5 \text{ GeV})^2$	689.5 ± 2.2	689.1 ± 4.2	690.8 ± 3.3	689.3 ± 2.1	689.4 ± 2.1	689.6 ± 2.2	688.1 ± 2.1	689.4 ± 5.7
$(12 \text{ GeV})^2$	690.3 ± 2.1	689.9 ± 4.6	691.6 ± 3.6	689.8 ± 2.1	690.1 ± 2.2	690.2 ± 2.2	688.6 ± 2.1	690.0 ± 5.9

Table 3: Values of $a_\mu^{\text{HLO (I)}}$ for all the parameterizations considered in this study and the three choices of s_0 . Results using the two different minimization techniques for $K_1(s)$ are shown. Dataset I was used as input.

uncertainty of $\alpha_s(M_Z^2)$. To compute $a_\mu^{\text{HLO (III)}}$ and $a_\mu^{\text{HLO (IV)}}$, e^+e^- data from Dataset I³ were used for $R(s)$ up to $s = 10 \text{ GeV}$, while pQCD was used above that value. Table 4 shows the values of $a_\mu^{\text{HLO (II)}}$, $a_\mu^{\text{HLO (III)}}$ and $a_\mu^{\text{HLO (IV)}}$ for the three different s_0 values and the two minimization techniques for the approximated kernel.

³Similar values are obtained using Dataset II and III.

Minimization I			
s_0 values	$a_\mu^{\text{HLO (II)}} (10^{-10})$	$a_\mu^{\text{HLO (III)}} (10^{-10})$	$a_\mu^{\text{HLO (IV)}} (10^{-10})$
$(1.8 \text{ GeV})^2$	2.94 ± 0.04	0.43 ± 0.01	2.95 ± 0.05
$(2.5 \text{ GeV})^2$	1.84 ± 0.01	-0.34 ± 0.01	1.79 ± 0.02
$(12 \text{ GeV})^2$	0.208 ± 0.001	-1.695 ± 0.035	0.079 ± 0.001
Minimization II			
s_0 values	$a_\mu^{\text{HLO (II)}} (10^{-10})$	$a_\mu^{\text{HLO (III)}} (10^{-10})$	$a_\mu^{\text{HLO (IV)}} (10^{-10})$
$(1.8 \text{ GeV})^2$	3.23 ± 0.04	0.91 ± 0.02	3.00 ± 0.05
$(2.5 \text{ GeV})^2$	2.54 ± 0.01	1.52 ± 0.02	1.96 ± 0.02
$(12 \text{ GeV})^2$	0.360 ± 0.001	4.85 ± 0.05	0.096 ± 0.001

Table 4: Values of $a_\mu^{\text{HLO (II)}}$, $a_\mu^{\text{HLO (III)}}$, $a_\mu^{\text{HLO (IV)}}$ for the three choices of s_0 and the two different minimization techniques used to determine the approximated kernel $K_1(s)$. We use e^+e^- data from Dataset I.

5. Results

Table 5 shows the final results for a_μ^{HLO} obtained adding the four terms according to Eq. 11. They are compared to the values of a_μ^{HLO} obtained using the integral in Eq. 1. Using this method, the results obtained from the Padé and Polynomial parameterization are in strong disagreement with the reference input value. On the other hand, the same parameterizations can be safely adopted with our new method, since they allow to compute a_μ^{HLO} with no significant bias with respect to the reference value. For the sake of illustration, final results for $s_0 = (1.8 \text{ GeV})^2$, Dataset I and all the parameterizations are shown in Fig.4. Tables 6,7 and Figs. 5,6 show the same results for a_μ^{HLO} obtained with our methods when Datasets II and III are used.

6. Conclusions

We have shown a method to compute a_μ^{HLO} with MUonE data using a derivative approach based on Refs. [11, 12]. This method relies on the analyticity properties of the hadronic vacuum polarization function $\Pi_{had}(t)$ which allow to express $\Delta\alpha_{had}(t)$ as a polynomial by a Taylor expansion at $t \sim 0$. The results obtained for different parameterizations used to fit $\Delta\alpha_{had}(t)$ and different input datasets show that the method is competitive with the traditional one based on the integral of $\Delta\alpha_{had}(t)$ in the whole momentum transfer range. Moreover, the proposed method avoids the difficulties with the functional form of the parameterization used to extrapolate $\Delta\alpha_{had}(t)$ behaviour outside the MUonE range. We believe that by a convenient choice of the approximate kernel function this method can be applied to extract the NLO and NNLO hadronic vacuum polarization contributions to the muon $g-2$ in the space-like region [29].

Acknowledgements

We thank Stefano Laporta for useful suggestions and Massimo Passera for useful suggestions and contribution in a early phase of the project. This work was supported by the Leverhulme Trust, LIP-2021-01.

$a_\mu^{\text{HLO, INPUT}} = 695.1 (10^{-10})$								
$a_\mu^{\text{HLO (Der)}}$								
Minimization I	LL	Padé	Pol	GdR1	GdR2	GdR3	GdR4	GdR5
s_0 values								
$(1.8 \text{ GeV})^2$	695.0 ± 2.2	695.0 ± 2.9	695.2 ± 2.9	694.5 ± 2.2	694.3 ± 2.2	694.3 ± 2.2	693.3 ± 2.3	694.3 ± 2.6
$(2.5 \text{ GeV})^2$	695.0 ± 2.2	694.9 ± 3.0	695.1 ± 3.0	694.3 ± 2.2	694.1 ± 2.2	694.1 ± 2.2	693.1 ± 2.3	694.2 ± 2.9
$(12 \text{ GeV})^2$	694.9 ± 2.2	694.9 ± 3.0	694.9 ± 3.2	694.0 ± 2.2	693.9 ± 2.2	693.8 ± 2.2	692.7 ± 2.3	693.9 ± 3.7
Minimization II	LL	Padé	Pol	GdR1	GdR2	GdR3	GdR4	GdR5
s_0 values								
$(1.8 \text{ GeV})^2$	695.6 ± 2.2	695.2 ± 4.2	696.9 ± 3.3	695.4 ± 2.1	695.5 ± 2.1	695.7 ± 2.2	694.2 ± 2.1	695.5 ± 5.8
$(2.5 \text{ GeV})^2$	695.5 ± 2.2	695.1 ± 4.2	696.8 ± 3.3	695.3 ± 2.1	695.4 ± 2.1	695.6 ± 2.2	694.1 ± 2.1	695.4 ± 5.7
$(12 \text{ GeV})^2$	695.6 ± 2.1	695.2 ± 4.6	696.9 ± 3.6	695.1 ± 2.1	695.4 ± 2.2	695.5 ± 2.2	693.9 ± 2.1	695.3 ± 5.9
$a_\mu^{\text{HLO (Int)}}$	695.3 ± 2.1	702.6 ± 12.0	834.3 ± 63.2	696.6 ± 2.2	696.5 ± 2.2	696.3 ± 2.2	696.8 ± 2.2	696.5 ± 3.5

Table 5: Final results of a_μ^{HLO} calculated using the proposed method ($a_\mu^{\text{HLO (Der)}}$), for all the parameterizations considered in this study, with the three choices of s_0 and the two minimization techniques used to determine the approximated kernel $K_1(s)$. $a_\mu^{\text{HLO (Int)}}$ is the result of the integral in Eq.1, when a specific parameterization for $\Delta\alpha_{\text{had}}(t)$ is used as input. $a_\mu^{\text{HLO, INPUT}}$ is the reference value of a_μ^{HLO} obtained by integrating $\Delta\alpha_{\text{had}}(t)$ from Dataset I using Eq.1.




Figure 4: Values of a_μ^{HLO} for all the parameterizations considered in the text, obtained with different methods: MUonE integral from Eq. 1 (empty squares), our new method using Minimization 1 to get the approximated kernel (circles) our new method using Minimization 2 (diamond). Results for the Padé and Pol parameterizations computed using Eq. 1 are outside the plot range. The black dashed line represents the reference value obtained from Dataset I. Results are shown for $s_0 = (1.8 \text{ GeV})^2$.

$a_\mu^{\text{HLO, INPUT}} = 690.3 (10^{-10})$								
$a_\mu^{\text{HLO (Der)}}$								
Minimization I	LL	Padé	Pol	GdR1	GdR2	GdR3	GdR4	GdR5
s_0 values								
$(1.8 \text{ GeV})^2$	690.3 ± 2.2	690.2 ± 2.9	690.5 ± 2.9	689.7 ± 2.2	689.6 ± 2.2	689.6 ± 2.2	688.6 ± 2.3	689.6 ± 2.6
$(2.5 \text{ GeV})^2$	690.2 ± 2.2	690.1 ± 3.0	690.3 ± 3.0	689.5 ± 2.2	689.4 ± 2.2	689.4 ± 2.2	688.4 ± 2.3	689.4 ± 2.9
$(12 \text{ GeV})^2$	690.1 ± 2.2	690.0 ± 3.0	690.0 ± 3.2	689.1 ± 2.2	689.1 ± 2.2	689.1 ± 2.2	688.1 ± 2.3	689.1 ± 3.7
Minimization II	$a_\mu^{\text{HLO (Der)}}$							
s_0 values	LL	Padé	Pol	GdR1	GdR2	GdR3	GdR4	GdR5
$(1.8 \text{ GeV})^2$	690.8 ± 2.2	690.4 ± 4.1	692.1 ± 3.3	690.7 ± 2.1	690.7 ± 2.1	690.8 ± 2.2	689.4 ± 2.1	690.7 ± 5.9
$(2.5 \text{ GeV})^2$	690.8 ± 2.2	690.4 ± 4.2	692.0 ± 3.3	690.5 ± 2.1	690.6 ± 2.1	690.7 ± 2.2	689.3 ± 2.1	690.6 ± 5.7
$(12 \text{ GeV})^2$	690.8 ± 2.1	690.4 ± 4.6	692.1 ± 3.6	690.4 ± 2.1	690.5 ± 2.2	690.6 ± 2.2	689.1 ± 2.1	690.5 ± 5.9
$a_\mu^{\text{HLO (Int)}}$	690.5 ± 2.1	695.8 ± 8.8	724.2 ± 31.2	691.7 ± 2.2	691.7 ± 2.2	691.6 ± 2.2	692.0 ± 2.2	691.7 ± 3.5

Table 6: Final results of a_μ^{HLO} calculated using the proposed method ($a_\mu^{\text{HLO (Der)}}$), for all the parameterizations considered in this study, with the three choices of s_0 and the two minimization techniques used to determine the approximated kernel $K_1(s)$. $a_\mu^{\text{HLO (Int)}}$ is the result of the integral in Eq. 1, when a specific parameterization for $\Delta\alpha_{\text{had}}(t)$ is used as input. $a_\mu^{\text{HLO, INPUT}}$ is the reference value of a_μ^{HLO} obtained by integrating $\Delta\alpha_{\text{had}}(t)$ from Dataset II using Eq. 1.




Figure 5: Values of a_μ^{HLO} for all the parameterizations considered in the text, obtained with different methods: MUonE integral from Eq. 1 (empty squares), our new method using Minimization 1 to get the approximated kernel (circles) our new method using Minimization 2 (diamond). Results for the Padé and Pol parameterizations computed using Eq. 1 are outside the plot range. The black dashed line represents the reference value obtained from Dataset II. Results are shown for $s_0 = (1.8 \text{ GeV})^2$.

$a_\mu^{\text{HLO, INPUT}} = 693.0 (10^{-10})$								
$a_\mu^{\text{HLO (Der)}}$								
Minimization I	LL	Padé	Pol	GdR1	GdR2	GdR3	GdR4	GdR5
s_0 values								
$(1.8 \text{ GeV})^2$	693.0 ± 2.2	693.0 ± 2.9	693.2 ± 2.9	692.5 ± 2.2	692.2 ± 2.2	692.2 ± 2.2	691.2 ± 2.3	692.2 ± 2.6
$(2.5 \text{ GeV})^2$	692.9 ± 2.2	693.0 ± 3.0	693.1 ± 3.0	692.3 ± 2.2	692.1 ± 2.2	692.1 ± 2.2	691.0 ± 2.3	692.1 ± 2.9
$(12 \text{ GeV})^2$	692.9 ± 2.2	692.9 ± 3.0	692.9 ± 3.2	692.0 ± 2.2	691.8 ± 2.2	691.8 ± 2.2	690.7 ± 2.2	691.8 ± 3.7
Minimization II	LL	Padé	Pol	GdR1	GdR2	GdR3	GdR4	GdR5
s_0 values								
$(1.8 \text{ GeV})^2$	693.5 ± 2.2	693.3 ± 4.2	694.9 ± 3.3	693.4 ± 2.1	693.5 ± 2.1	693.7 ± 2.2	692.2 ± 2.1	693.7 ± 5.8
$(2.5 \text{ GeV})^2$	693.5 ± 2.2	693.3 ± 4.2	694.8 ± 3.3	693.3 ± 2.1	693.4 ± 2.1	693.6 ± 2.2	692.1 ± 2.1	693.6 ± 5.7
$(12 \text{ GeV})^2$	693.6 ± 2.1	693.4 ± 4.6	695.0 ± 3.6	693.1 ± 2.1	693.4 ± 2.1	693.6 ± 2.2	691.9 ± 2.1	693.7 ± 5.9
$a_\mu^{\text{HLO (Int)}}$	693.4 ± 2.1	698.6 ± 10.1	730.7 ± 32.9	694.5 ± 2.2	694.4 ± 2.2	694.3 ± 2.2	694.7 ± 2.2	694.3 ± 3.5

Table 7: Final results of a_μ^{HLO} calculated using the proposed method ($a_\mu^{\text{HLO (Der)}}$), for all the parameterizations considered in this study, with the three choices of s_0 and the two minimization techniques used to determine the approximated kernel $K_1(s)$. $a_\mu^{\text{HLO (Int)}}$ is the result of the integral in Eq. 1, when a specific parameterization for $\Delta\alpha_{\text{had}}(t)$ is used as input. $a_\mu^{\text{HLO, INPUT}}$ is the reference value of a_μ^{HLO} obtained by integrating $\Delta\alpha_{\text{had}}(t)$ from Dataset III using Eq. 1.




Figure 6: Values of a_μ^{HLO} for all the parameterizations considered in the text, obtained with different methods: MUonE integral from Eq. 1 (empty squares), our new method using Minimization 1 to get the approximated kernel (circles) our new method using Minimization 2 (diamond). Results for the Padé and Pol parameterizations computed using Eq. 1 are outside the plot range. The black dashed line represents the reference value obtained from Dataset III. Results are shown for $s_0 = (1.8 \text{ GeV})^2$.

References

- [1] B. Abi, et al., Measurement of the Positive Muon Anomalous Magnetic Moment to 0.46 ppm, Phys. Rev. Lett. 126 (14) (2021) 141801. [arXiv:2104.03281](#), [doi:10.1103/PhysRevLett.126.141801](#).
- [2] D. P. Aguillard, et al., Measurement of the Positive Muon Anomalous Magnetic Moment to 0.20 ppm (8 2023). [arXiv:2308.06230](#).
- [3] T. Aoyama, et al., The anomalous magnetic moment of the muon in the Standard Model, Phys. Rept. 887 (2020) 1–166. [arXiv:2006.04822](#), [doi:10.1016/j.physrep.2020.07.006](#).
- [4] G. Colangelo, et al., Prospects for precise predictions of a_μ in the Standard Model (3 2022). [arXiv:2203.15810](#).
- [5] S. Borsanyi, et al., Leading hadronic contribution to the muon magnetic moment from lattice QCD, Nature 593 (7857) (2021) 51–55. [arXiv:2002.12347](#), [doi:10.1038/s41586-021-03418-1](#).
- [6] F. V. Ignatov, et al., Measurement of the $e^+e^- \rightarrow \pi^+\pi^-$ cross section from threshold to 1.2 GeV with the CMD-3 detector (2 2023). [arXiv:2302.08834](#).
- [7] C. M. Carloni Calame, M. Passera, L. Trentadue, G. Venanzoni, A new approach to evaluate the leading hadronic corrections to the muon $g-2$, Phys. Lett. B 746 (2015) 325–329. [doi:10.1016/j.physletb.2015.05.020](#).
- [8] G. Abbiendi, et al., Measuring the leading hadronic contribution to the muon $g-2$ via μe scattering, Eur. Phys. J. C 77 (3) (2017) 139. [doi:10.1140/epjc/s10052-017-4633-z](#).
- [9] B. Lautrup, A. Peterman, E. de Rafael, Recent developments in the comparison between theory and experiments in quantum electrodynamics, Physics Reports 3 (4) (1972) 193–259. [doi:10.1016/0370-1573\(72\)90011-7](#).
- [10] G. Abbiendi, Status of the muone experiment, Physica Scripta 97 (5) (2022) 054007. [doi:10.1088/1402-4896/ac6297](#).
- [11] S. Bodenstein, C. A. Dominguez, K. Schilcher, Hadronic contribution to the muon $g-2$ factor: A theoretical determination, Phys. Rev. D 85 (2012) 014029. [doi:10.1103/PhysRevD.85.014029](#).
- [12] C. A. Dominguez, et al., Anomalous magnetic moment of the muon: A hybrid approach, Phys. Rev. D 96 (2017) 074016. [doi:10.1103/PhysRevD.96.074016](#).
- [13] M. Gourdin, E. De Rafael, Hadronic contributions to the muon g -factor, Nuclear Physics B 10 (4) (1969) 667–674. [doi:10.1016/0550-3213\(69\)90333-2](#).
- [14] L. Durand, Pionic contributions to the magnetic moment of the muon, Phys. Rev. 128 (1962) 441–448. [doi:10.1103/PhysRev.128.441](#).
- [15] L. Durand, Erratum: Pionic contributions to the magnetic moment of the muon, Phys. Rev. 129 (1963) 2835–2835. [doi:10.1103/PhysRev.129.2835.5](#).
- [16] J. Bell, E. de Rafael, Hadronic vacuum polarization and $g_\mu - 2$, Nuclear Physics B 11 (4) (1969) 611–620. [doi:10.1016/0550-3213\(69\)90250-8](#).
- [17] M. Davier, A. Hocker, New results on the hadronic contributions to $\alpha(M(Z)^{**2})$ and to $(g-2)(\mu)$, Phys. Lett. B 435 (1998) 427–440. [arXiv:hep-ph/9805470](#), [doi:10.1016/S0370-2693\(98\)00825-9](#).
- [18] S. Groote, J. G. Korner, K. Schilcher, N. F. Nasrallah, QCD sum rule determination of $\alpha(M(Z))$ with minimal data input, Phys. Lett. B 440 (1998) 375–385. [arXiv:hep-ph/9802374](#), [doi:10.1016/S0370-2693\(98\)01107-1](#).
- [19] F. Ignatov, Vacuum Polarization, <http://cmd.inp.nsk.su/~ignatov/vpl/> (2022).
- [20] F. Jegerlehner, Software packages alphaQED and pQCADAdler, <http://www-com.physik.hu-berlin.de/~fjeger/software.html> (2019).
- [21] A. Keshavarzi, D. Nomura, T. Teubner, $g-2$ of charged leptons, $\alpha(m_\tau^2)$, and the hyperfine splitting of muonium, Phys. Rev. D 101 (1 2020). [doi:10.1103/PhysRevD.101.014029](#).
- [22] D. Greynat, E. de Rafael, Hadronic vacuum polarization and the MUonE proposal, JHEP 2022 (5) (may 2022). [doi:10.1007/jhep05\(2022\)084](#).
- [23] P. A. Baikov, K. G. Chetyrkin, J. H. Kuhn, J. Rittinger, Vector Correlator in Massless QCD at Order $\mathcal{O}(\alpha_s^4)$ and the QED beta-function at Five Loop, JHEP 07 (2012) 017. [arXiv:1206.1284](#), [doi:10.1007/JHEP07\(2012\)017](#).
- [24] A. Maier, P. Marquard, Low- and High-Energy Expansion of Heavy-Quark Correlators at Next-To-Next-To-Leading Order, Nucl. Phys. B 859 (2012) 1–12. [arXiv:1110.5581](#), [doi:10.1016/j.nuclphysb.2012.01.021](#).
- [25] P. A. Baikov, K. G. Chetyrkin, J. H. Kuhn, $R(s)$ and hadronic tau-Decays in Order $\alpha_s^{*4}(s)$: Technical aspects, Nucl. Phys. B Proc. Suppl. 189 (2009) 49–53. [arXiv:0906.2987](#), [doi:10.1016/j.nuclphysbps.2009.03.010](#).
- [26] F. Herren, M. Steinhauser, Version 3 of RunDec and CRUnDec, Comput. Phys. Commun. 224 (2018) 333–345. [arXiv:1703.03751](#), [doi:10.1016/j.cpc.2017.11.014](#).
- [27] A. Pich, A. Rodríguez-Sánchez, Determination of the QCD coupling from ALEPH τ decay data, Phys. Rev. D 94 (3) (2016) 034027. [arXiv:1605.06830](#), [doi:10.1103/PhysRevD.94.034027](#).
- [28] D. Boito, M. Golterman, K. Maltman, S. Peris, M. V. Rodrigues, W. Schaaf, α_s from an improved τ vector isovector spectral function, EPJ Web Conf. 274 (2022) 03014. [arXiv:2210.13518](#), [doi:10.1051/epjconf/202227403014](#).
- [29] E. Balzani, S. Laporta, M. Passera, Hadronic vacuum polarization contributions to the muon $g-2$ in the space-like region, Physics Letters B 834 (2022) 137462. [doi:10.1016/j.physletb.2022.137462](#).