

Black hole charges in dark matter haloes and the seeds of cosmic magnetic fields

Nelson D. Padilla,^a Ignacio J. Araya^{b,c,d} and Federico Stasyszyn^{a,e}

^aInstituto de Astronomía Teórica y Experimental, IATE, CONICET-UNC, Laprida 854, X5000BGR, Córdoba, Argentina

^bInstituto de Ciencias Exactas y Naturales, Universidad Arturo Prat, Playa Brava 3256, 1111346, Iquique, Chile

^cFacultad de Ciencias, Universidad Arturo Prat, Avenida Arturo Prat Chacón 2120, 1110939, Iquique, Chile

^dSchool of Mathematics and Hamilton Mathematics Institute, Trinity College, Dublin 2, Ireland

^eObservatorio Astronómico de la Universidad Nacional de Córdoba, Laprida 854, X5000BGR, Córdoba, Argentina

E-mail: nelson.padilla@unc.edu.ar

Abstract. We present improved estimates of the electric charge that black holes could hold when these are embedded in the ionised plasma within dark matter haloes (or galaxies) in the Universe. We have implemented the spontaneous emission of charges of opposite sign to that of the black hole via athermal Hawking evaporation, including its dependence on black hole spin, and we have estimated the equilibrium charge that arises as this charge loss is balanced by the continuous accretion of charges from the surrounding plasma. The resulting charge can be several orders of magnitude lower than previously estimated upper limits, but it can surpass the pair production limit noted by Gibbons (1974) by a margin that increases with the amplitude of the black hole spin and the density of the plasma. We also implement a calculation for the net charge and magnetic moment of dark matter haloes when the dark matter is made of primordial black holes and also for astrophysical black holes that form part of the stellar halo of galaxies. We calculate the resulting magnetic fields of $z = 15$ haloes for these different cases, and show that both primordial black holes and stellar ones could provide the magnetic field that can seed the observed ones in present-day galaxies.

Contents

1	Introduction	1
1.1	Outline and conventions	3
2	Charge of black holes within dark matter haloes	3
2.1	Charge accretion and emission of non-maximally spinning black holes in an ionised plasma	4
2.2	Charging timescales of non-maximally spinning black holes embedded in a galaxy	8
2.3	Instantaneous Coulomb potential approximation	10
2.4	Black hole charges within DM haloes	10
2.5	Required charges to provide cosmic magnetic seeds	10
3	Net charge of dark matter haloes	12
3.1	Dark matter halo charge from maximally spinning primordial black holes	12
3.2	Net charges and magnetic moments in galaxies with non-maximally spinning black holes	14
4	Seeds for the cosmic magnetic fields	16
5	Discussion	18
6	Conclusions	20

1 Introduction

Our current picture of the Universe contains several unsolved mysteries, such as the existence and nature of the Dark Matter (DM) and the origin of magnetic fields.

The evidence for the existence of dark matter is quite varied but indirect, including early studies of the rotation curves of spiral galaxies [1, 2], and more recent results from the large-scale structure of the universe [3, 4] and cosmological parameter fits to the temperature fluctuation power spectrum of the cosmic microwave background [5, 6], among other probes.

Regarding the possible nature of dark matter, there are several candidates ranging from thermally produced fundamental particles that interact with other species mostly via gravity, as is the case of particles from extensions to the standard model of particle physics that produce the weakly interacting massive particles (WIMPs, [7–9]), to primordial black holes that formed very early in the Universe [10–12]. The latter are possible candidates as they form early enough, considering that dark matter should be present since very early times [5, 6, 13, 14].

The search for WIMPs in dark matter experiments has so far produced no direct detections [15–18], and the same is true of searches of effects due to black holes on different observables, which provide strong constraints on the fraction of dark matter that can be stored in black holes, which should be well below unity over most of the range of possible black hole masses, except in two windows around the asteroid mass and the solar mass [19–22].

In a recent paper, Araya et al. (2023, [23], A23 from this point on) showed that primordial black holes could hold a non-zero electric charge. Moreover, there are also particle

candidates that could be electrically charged such as milli-charged particles [24]. The charge associated to dark matter particles provides additional observables that could serve to prove in another, independent indirect way, the existence of dark matter. In A23, the total charge of black holes is in most cases much smaller than the extremal value for the Reissner-Nördstrom metric for charged black holes, meaning that the electromagnetic potential is always much smaller than that of gravity. This places charged black holes below the current limit for charges for dark matter particles [25]. The inferred black hole charge by A23 would depend on the black hole spin.

Even though there are multiple constraints on the abundance of primordial black holes as a function of their mass from observations, there are no constraints on the spin that these black holes could have. Given that the charge of black holes could depend on their spin, it is interesting to review the literature for the current limits to its possible values. Thorne (1974,[26]) investigated the possible spin-up of black holes as a result of accretion of material via a thin disc. They found that once the spin is high enough, radiation from the thin disc itself effectively removes angular momentum from the black hole limiting the maximum dimensionless spin parameter to $a_{max} = 0.998$. Several authors agree with this estimate using different methods, also providing further extraction of angular momentum via the production of a jet [27, 28]. However, there are numerical estimates that allow the spin to reach much higher values such as $a = 0.99997$ [29]. This was confirmed later by Benson & Babul [30], but they argue that an evolving spin will probably reach its maximum at the Thorne limit.

Observations of supermassive black holes do not seem to reach spins much higher than $a = 0.6$ [31], but semi-analytic simulations of the co-evolution of supermassive black holes and galaxies in a cosmological context, show that the spin could reach values $a \sim 1$ [32]. On a different mass scale, observations of high mass X-ray binaries point to an important increase in the spin of the black hole component [33]. These estimates of black hole spin are limited in precision which makes it difficult to estimate how close to the maximal one they actually are.

Opening the door for charged dark matter also brings along other possibilities, such as the production of magnetic fields. Magnetic fields are ubiquitous in astrophysics [34, 35], and are a necessary ingredient when trying to explain the structure, dynamics, and evolution of the Universe across all scales [36, 37]. They exist in large and small structures, impacting particle acceleration and the evolution of the primordial plasma [38]. Studying magnetic fields at high redshift remains challenging, but their influence on plasma physics may soon be observable. Primordial magnetic field astronomy is a highly intense and challenging field in Cosmology, offering insights into the Early Universe and its impact on CMB anisotropies [39, 40].

The origin, evolution, and amplification mechanisms of these fields are still unclear, leading to intriguing hypotheses [39, 41–43]. Generating mechanisms often rely on a "seed field", raising questions about their creation, amplification, and large-scale distribution. Various approaches have provided answers consistent with observations, following the principles of Magnetohydrodynamics. Magnetic seed fields may have originated from the Big Bang or the Early Universe, possibly through battery-like processes under specific conditions [36, 44, 45], during phase transitions [46–49], or spontaneously by turbulent motions of the plasma [50]. In particular, Araya et al. (2020, [51], A20 from this point on) show that primordial black holes with extended Press-Schechter like mass functions are in some cases able to provide the required seeds if black holes acquire a magnetic monopole charge as they form [52]. And in a companion letter, Padilla et al. (2023) we show that the required seeds can also be produced

by imbalances in the charged carriers of different sign at the epoch of galaxy formation.

In this paper we explore two main subjects. We first make an extension of the formalism of A23 to explore the late time charge in black holes embedded in a plasma, to make a more detailed mapping of the charges and their dependence on black hole parameters. A23 present two extreme cases of spin, namely maximally spinning ones with dimensionless spin parameter $a = 1$, which could preserve their initial charge, and black holes with zero spin. Here we explore the intermediate cases as there is ample evidence that black holes that accrete matter, even if their initial spin is null, can spin up to $a \sim 1$ [26]. We find that black holes with higher spin can hold larger electrical charges.

We then explore whether the charge in these black holes combined, in some cases, with the induced charge imbalance in the baryonic plasma in rotating proto-galaxies is able to produce the required seeds for the cosmic magnetic fields without resorting to magnetic monopole charges as in A20. We also look into the possibility that black holes of astrophysical origin can acquire electrical charge from the baryonic plasma of the first galaxies and can therefore provide the required seed field.

1.1 Outline and conventions

This paper is organised as follows. Section 2 expands the model for the charge accretion and emission by black holes in an ionised plasma within galaxies, it looks into the timescales involved in these two processes, presents an equilibrium charge, and compares it to the charge needed for black holes to explain the cosmic magnetic field seeds. Section 3 presents the final charge and magnetic moment of rotating dark matter haloes which host galaxies. Section 4 presents the resulting seed magnetic fields on scales of galaxies, at very high redshift, for black holes of primordial origin with different spins, and for black holes of astrophysical origin that populate the stellar haloes of galaxies. We present a discussion of our results and our assumptions in Section 5, and conclude in Section 6.

For our calculations we adopt a flat cosmology with $\Omega_{m,0} = 0.315$; $\Omega_{dm,0} = 0.264$; $\Omega_{r,0} = 9.237 \times 10^{-5}$ and $H_0 = 67 \frac{\text{km}}{\text{s}} \text{Mpc}^{-1}$, consistent with the latest measurements from the Planck Satellite [6]. When considering that a fraction of the dark matter is in primordial black holes, we assume that they all have equal mass, i.e. a monochromatic primordial black hole mass function. Throughout we assume that galaxies are hosted by dark matter haloes (sometimes referred to simply as haloes), and will refer to the total mass of galaxies including their dark matter haloes as the virial mass. We will frequently focus on Milky Way like galaxies since these are the typical galaxies of the Universe from the point of view of their stellar populations (e.g. [53]). We adopt the international system of units, and use the following notation for constants where e is the absolute value of the electron's charge, c stands for the speed of light, k_c is Coulomb's constant, G is the universal gravity constant, h is Planck's constant and \hbar its reduced version, m_p and m_e are the proton and electron masses, k_B is Boltzmann's constant and μ_0 is the magnetic permeability in vacuum.

2 Charge of black holes within dark matter haloes

We will consider that black holes can hold a charge that can come from two distinct origins; that it was acquired at the formation event of the black hole and that it lasted unperturbed to the present day, or that they acquired charge at late times. We will consider black holes of primordial and stellar origin indistinctly both of which can be present in galaxies, except that we will allow primordial ones to contain a non-zero initial electric charge and a dimensionless

spin parameter that can range from zero to maximal, whereas stellar ones will be assumed to hold no charge and to have no spin at their formation.

In the case of primordial black holes we will consider two different types. Ones that could form from overdensities left over from Inflation, that as they enter the horizon early during radiation domination they collapse and form a black hole that could hold a non-zero charge due to the unrelaxed spatial distribution of positive and negative charges at the horizon scale that is just coming into causal contact. In A23 this charge is referred to as the Poisson initial charge. For a Lagrangian volume, V_0 , that just enters the horizon the charge is,

$$Q_0 = \pm \sum_i |q_i| \sqrt{V_0 n_i}, \quad (2.1)$$

where the index i runs over all the particles present at temperature \bar{T}_0 , n_i is the number density of each particle,

$$V_0 = \frac{Mc^2}{\rho_{\text{uni}}(\bar{T}_0)}, \quad (2.2)$$

M if the mass of the formed black hole, and $\rho_{\text{uni}}(\bar{T}_0)$ is the average energy density of the Universe at \bar{T}_0 .

The second type of primordial black holes could be formed by the high energy collision of equal charge fundamental particles that could produce extremal Reissner-Nördstrom (RN) black holes of at least one Planck mass, with a charge that ensures extremality,

$$Q_{\text{max}} \sim \sqrt{\frac{G}{k_c}} m_{\text{Planck}}, \quad (2.3)$$

where m_{Planck} is the Planck mass. This charge could be achieved by the collision of two charged particles taking into account the running of the electric charge with the energy scale of the collision [54].

In A23 it is proposed that if the dimensionless spin parameter is maximal, $a = 1$ then the initial charge could be held within, or in close proximity of, the black hole such that for the distant observer the charge would be seen as remaining constant all the way to the present day.

For the case of non-maximal spin they show that the initial charge is lost by the epoch of recombination. However, they also show that later on when black holes find themselves within the ionised plasma of dark matter haloes/galaxies, they can acquire a charge since the late-time charge carriers with opposite charge sign have masses that differ by several orders of magnitude; i.e. electrons are more likely to be accreted. As a result primordial black holes with non-maximal spin should acquire a late time electric charge. Note that this particular case can also apply to black holes of stellar origin. The following subsection explores the charge that such non-maximally spinning black holes would be able to acquire as a function of black hole mass and spin.

2.1 Charge accretion and emission of non-maximally spinning black holes in an ionised plasma

In order to reach a reasonable estimate of the final black hole charge within a plasma in galaxies/haloes, A23 took into account processes of spontaneous charge production at the outer horizon of black holes, namely the Hawking radiation process [12] and the Schwinger mechanism [55, 56]. For the latter they did not include a detailed calculation of the emission

due to vacuum polarisation; instead they only checked whether the estimated charges were below the rapid discharge limit estimated by Gibbons (1975, [56]), which was satisfied by all their estimates. For the Hawking evaporation, A23 only made the calculation for black hole (BH) masses $M \lesssim 2 \times 10^{13} \text{kg}$ which corresponds to the BH mass that satisfies $k_B T_H \sim m_e c^2$, susceptible to charge emission by thermal Hawking radiation.

In this paper we improve the estimates of Hawking evaporation charge emission in higher mass black holes using Hawking radiation which, even when the Hawking temperature is much lower than the rest mass of electrons, is able to emit the latter due to the chemical potential produced by the black hole charge, as postulated by Lehman et al. [57], who term this emission as athermal. Furthermore, Lehmann et al. also estimate that the charge emitted via the Schwinger mechanism is of the same order of magnitude as the athermal Hawking emission, which allows us to use our Hawking athermal emission calculations as good measures of the total charge emission.

As in A23, we estimate the charge emission via Hawking radiation adopting the distribution function,

$$f(\epsilon(p), \mu) = \frac{1}{\exp\left(-\frac{\epsilon(p)-\mu}{k_B T_H}\right) + 1}, \quad (2.4)$$

where

$$T_H = \frac{\hbar}{2\pi G k_B} \frac{\sqrt{(1-a)M^2 - \frac{k_c}{G}Q^2}}{2M^2 + 2M\sqrt{(1-a)M^2 - \frac{k_c}{G}Q^2}}$$

is the Hawking temperature of the BH that takes into account the spin and charge of the black hole [58], $\epsilon(p) = \sqrt{p^2 c^2 + m_e^2 c^4}$ is the kinetic energy of a single particle with momentum p and mass m_e ¹. We introduce a modification in μ , the chemical potential that accounts for the Coulomb energy of the particle with electric charge $\pm e$ sourced by the BH charge Q , by allowing it to include the magnetic field due to the spin of the black hole,

$$\mu(\pm e) \simeq \frac{(1-a)k_c Q(\pm e)}{r_+}, \quad (2.5)$$

where we have made use of the simplification that the averaged amplitude of the dipolar magnetic field energy is simply that of the Coulomb one multiplied by the dimensionless spin parameter $a = cJ/(GM^2)$ (derived from [59]) valid when $a = 0$ or $a \sim 1$, with J the angular momentum of the black hole, and where r_+ is the outer horizon,

$$r_+ = \frac{r_s + \sqrt{r_s^2 - 4r_a^2 - 4r_Q^2}}{2}, \quad (2.6)$$

where $r_Q^2 = Q^2 G/(4\pi\epsilon_0 c^4)$ and $r_a = J/(Mc)$. This simple assumption allows us to explore the effect of black hole spin on athermal Hawking charge emission. Note though that even when the spin is very close to the maximal one, i.e. for very small $(1-a)$, the chemical potential is still non-zero and able to produce athermal discharge when $k_B T_H \ll m_e c^2$, although at a slower rate.

¹Note that the distribution is of the Fermi-Dirac type as indicated by the +1 in the denominator; the -1 in A23 is simply a typo present only in their manuscript and not in their calculations (Araya, private communication).

We follow A23, and use the charge density ρ of created electrons or positrons in the vicinity of the BH horizon to write the current $J = \rho v$ that can be used to obtain the rate of charged particle loss by the black hole,

$$\dot{Q}_H(\pm e) = \int \vec{J}(\pm e) \cdot d\vec{A} = A \int \frac{2d^3p}{(2\pi\hbar)^3} v(p) \cos(\theta(\vec{p}, \hat{n})) (\pm e) f(\epsilon(p), \mu(\pm e)). \quad (2.7)$$

Adding together the emission of electrons and positrons, regardless of the BH charge sign, the final discharge rate due to Hawking emission and its dependence on black hole mass, charge and spin, reads

$$-\dot{Q}_H(M, Q, a) = A \int_0^\infty \frac{dp p^2}{(2\pi)^2 \hbar^3} v(p) (-e) (f(\epsilon(p), \mu(-e)) - f(\epsilon(p), \mu(e))). \quad (2.8)$$

This approximation ignores several details such as grey body correction factors, but allows us to obtain order of magnitude estimates for the charge emission by massive black holes with non-zero spin. Note that the inclusion of spin will also introduce changes to the final average charge of low mass black holes with Hawking temperature below the electron mass.

After implementing Eq. 2.8 we find that if the black hole charge is larger than the Gibbons limit for pair production charge [56],

$$Q^{\text{pairs}} = \frac{2GMm_e}{k_c e}, \quad (2.9)$$

then the rate of discharge is positive. For $Q \leq Q^{\text{pairs}}$, the rate is null, i.e. a charge below Q^{pairs} is stable if no charges are accreted onto the black hole. In a companion letter, Padilla et al. (2023), we note that this charge corresponds to an excess of one charge every $\sim 10^{39}$ baryons, which corresponds to the maximum charge that would be able to be maintained by the gravitational pull of any massive object.

One can compare the rate of discharge to the rate of charge accretion from the plasma surrounding the black holes. A23 estimated the maximum charge that a black hole would acquire until its repulsive force would impede further charge accretion as,

$$Q_{\text{max}} = \frac{k_B T_{\text{vir}} \left(\frac{1}{r_+} - \frac{1}{r_+ + \ell_{\text{Debye}}} \right)^{-1} + GMm_e}{k_c e}. \quad (2.10)$$

where they consider that electrons can move much faster than protons, and freely within a Debye length of the plasma,

$$\ell_{\text{Debye}} = \left(\frac{k_B T_{\text{vir}}}{4\pi k_c n_e e^2} \right)^{1/2}, \quad (2.11)$$

where n_e is the average number density of electrons. We only take into account the number density of electrons since regardless of virial temperature ($T_{\text{vir}} \leq 10^{10}\text{K}$) they have a higher thermal velocity than protons.

The rate of charge emission when the charge is Q_{max} is effectively zero, but if this charge is larger than Q^{pairs} then the charge will drop due to Hawking athermal emission. This emission fueled drop will continue until the negative charge accretion equals the emission, or until the emission stops at Q^{pairs} .

We follow A23 and assume that the cross-section for neutral BH charge accretion is the geometrical one, which is adequate for black holes with no charge; here we add the modification

that if the black hole charge is already non-zero, instead of adopting the innermost stable circular orbit, we use the distance at which the Coulomb and gravitational forces cancel each other out,

$$r_E^i = \frac{-k_c Q(\pm e) + GMm_p}{3k_B T_{\text{vir}}}, \quad (2.12)$$

where i stands for protons or electrons (p or e). We limit this scale to two extreme values. First, for charges below the limit where particles are energetic enough to be able to cross the Coulomb barrier, i.e.

$$k_B T_{\text{vir}} > (GM_{\text{BH}}m_e + k_c Qe) \left(\frac{1}{r_{\text{isco}} + \ell_{\text{Debye}}} - \frac{1}{r_{\text{isco}}} \right), \quad (2.13)$$

we limit the accretion area to have a radius of the innermost stable circular orbit, r_{ISCO} since the resulting charges are low enough that electrons and protons are able to break the Coulomb barrier and reach the black hole from a ℓ_{Debye} distance away. On the upper side we limit the radius of the cross section to one Debye length away from r_{ISCO} , since beyond this separation the potential of the charged black hole is shielded by the plasma. We ignore the magnetic dipole force as it drops with an extra factor of $1/r$ compared to the Coulomb one. With this, the cross sections for protons and electrons read,

$$\sigma_{\text{BH}}^i(M) = \begin{cases} \pi(1.4r_+)^2 & 1.4r_+ > r_E^i \text{ or } |Q| < e \\ \pi(r_E^i)^2 & 1.4r_+ < r_E^i < 1.4r_+ + \ell_{\text{Debye}} \\ \pi(1.4r_+ + \ell_{\text{Debye}})^2 & r_E^i > 1.4r_+ + \ell_{\text{Debye}}. \end{cases} \quad (2.14)$$

Then, the rate of charge acquisition (or loss) by BHs of mass M reads [60],

$$\Gamma(M) = n_e (\sigma_{\text{BH}}^e(M)v_e(T_{\text{vir}}) - \sigma_{\text{BH}}^p(M)v_p(T_{\text{vir}})) + \dot{Q}_H(M, Q, a)/e, \quad (2.15)$$

where v_e and v_p are the electron and proton thermal velocity within the halo, and we have added the charge loss due to Hawking emission in the last term. Black hole stable charges that do not rise above $\sim Q^{\text{pairs}}$ are not affected by Hawking athermal discharge.

Notice that the cross-sections for accretion of electrons and protons can be quite different depending on the sign of the charge of the black hole. Even when electrons are thermally faster than protons, protons can have much larger cross sections for accretion, and in this case the initial negative charge acquired when the black hole was neutral, is rapidly lost, and possibly turns negative. However, electrons can be rapidly acquired again, and the charge can be lost. The case in which this does not happen is when the cross section radii are similar to $r_{\text{isco}} + \ell_{\text{Debye}}$. Depending on the plasma density, this condition is met for black holes with masses $M > 10^{22}\text{kg}$, and black holes with larger masses can in general hold a charge as long as they are embedded in a plasma (and do not encounter free charges).

We explore values of n_e that correspond to matter densities ranging from 1 to 10^6 times the virial one defined as 200 times the critical density, i.e., $n_e^{\text{vir}} = 200\Omega_b\rho_c/(1.4m_p)$ approximating the average ion nuclei weight to $1.4m_p$ taking into account the presence of Helium.

We integrate Equation 2.15 in time using time intervals that do not produce changes in the charge larger than 1%. We integrate up to a time such that the charge shows no further significant changes (if the rate is always of the same sign) or until an oscillatory behaviour is reached (when the rate alternates sign). We refer to this final charge as the equilibrium charge Q_{eq} . Note that higher baryon densities produce higher accretion rates and increase the resulting equilibrium value.

2.2 Charging timescales of non-maximally spinning black holes embedded in a galaxy

We calculate the $z = 0$ timescales of accretion and emission, Q/\dot{Q} , and show estimates as a function of black hole mass for different dimensionless spin parameters $a < 1$, and show the results in the top panel of Figure 1. As can be seen, a larger dimensionless spin parameter results in a slower Hawking discharge above solar masses (blue dashed). We specifically show the case with $a = 0.998$ corresponding to the Thorne limit [26] for illustrative purposes. The green solid lines show that the accretion times are lower than discharge ones for masses below $\sim 10^{30}kg$, which implies that such black holes acquire their Q_{\max} charge relatively fast and are able to keep it for long (modulo exchange of protons and electrons to maintain equilibrium), sometimes even for longer than the dynamical time of a typical halo (t_{vir} , shown as a horizontal dotted line); the charges of such black holes can be felt by a distant observer as soon as the black hole moves away from the region of the plasma where it acquired its charge. In some cases the black hole could take along a trailing group of protons that would effectively neutralise its charge (see Section 5).

We illustrate the black hole mass ranges that satisfy these conditions in the bottom panel of Fig. 1, which shows in grey and hatched regions the black hole masses that can hold a charge for longer than the dynamical time of the halo (negative and positive charges, respectively), i.e. that can exert a Coulomb force that a distant observer can detect with its full charge, as a function of plasma density. Above this mass black holes reach an equilibrium where the charging and Hawking discharge times are equal (to within the numerical accuracy of our calculations). In this case the black hole charge is stable, but charges constantly enter and exit the black hole at very high rates. This charge is probably not measurable with its full amplitude by distant observers as there is a trail of exchanged charges left behind as the black hole traverses the plasma. This trail should rapidly relax back with the plasma possibly leaving a thermal contribution to its temperature. However, even if their stable charge is larger, these black holes can in principle be detected as having $Q^{\text{pairs}}(M)$ as further discussed below. The figure shows the time required to obtain this latter charge as an orange line. As can be seen, this charge is acquired quite rapidly in comparison to the virial time.

The extent to which there are macroscopic, galaxy wide consequences of the charge of black holes can be tackled bearing in mind that the dynamics of a pressureless fluid composed of black holes and that of the plasma can be thought of as being independent. If a black hole retains its charge for a long timescale and it does not capture from the plasma a trailing group of particles with opposite charge sign that could neutralise the BH charge, its effect will be seen by a distant observer (say, at least hundreds of parsecs away, in a different component of the halo, or from outside the halo). On the other hand, if the charge is stable but a result of a continuous, almost instantaneous accretion and emission of charges (in terms of the dynamical time of the halo), then one possibility is that the net charge of the plasma and black holes together will be seen as zero throughout the halo/galaxy by an external observer. A second possibility is that, even though the net stable charge could be larger, since the initial stages of charging of a black hole up to $Q^{\text{pairs}}(M)$ were not subject to emission discharges, the distant observer will be able to detect $Q^{\text{pairs}}(M)$. The black hole charge in this case would have been acquired from the plasma quite rapidly at some early time, and moved away from the gas that provided the electrons, effectively imprinting a difference in the spatial distribution of charges in black holes and in the gas within a galaxy.

Notice that all black holes are in principle subject to these charging and emission discharge processes, including ones of astrophysical origin, which can make up to a significant




Figure 1. Top: $z = 0$ timescales of charge accretion and emission by black holes. The blue lines show charge emission due to Hawking radiation, for increasing values of the dimensionless spin parameter a as shown in the figure key (larger timescales for larger spin). Green lines show the recharge time for the equilibrium or stable charge. Notice that at high masses the green and blue lines match closely, indicating that the equilibrium charge has been reached; in these cases the charge is $Q > Q_{\text{pairs}}$. When the green line lies below the blue one, the charge is rapidly acquired and an equilibrium between electron and proton accretion is reached. The orange line shows the time required to acquire the limit of pair production discharge Q_{pairs} . The horizontal dotted line shows the typical virial time of a $10^{12} h^{-1} M_{\odot}$ dark matter halo, and the violet dotted line shows the free-fall timescale for the Bondi mass, for BH masses where mass accretion from a plasma is possible. The solid black line shows the light travel time to traverse the Schwarzschild radius of the black hole showing that up to $M \sim 10^{35} \text{kg}$ the instantaneous Coulomb potential approximation is valid. Bottom: ranges of black hole mass for Hawking stable (equal Hawking discharge and charge accretion times), negative stable charge (accretion of electrons is balanced by short bursts of accretion of protons, maintaining a stable charge), positive stable charge, and zero charge of $Q = e$, as a function of plasma density at $z = 0$.

percentage of the total stellar mass formed within galaxies and can also be considered to have

independent dynamics to that of the baryonic plasma (see for instance [61, 62]).

2.3 Instantaneous Coulomb potential approximation

This formalism assumes that the Coulomb potential produced by the black hole is instantaneously adjusted as individual charges are accreted or emitted by the black hole. We check that this assumption is valid by adding in the top panel of Fig. 1 the light travel time for the Schwarzschild radius (black). As can be seen up to masses well above the solar mass, the light travel times are below the charging and discharge timescales for most of the black hole masses we will explore here. This ensures that there is enough time for the potential to relax between individual accretion or emission events.

2.4 Black hole charges within DM haloes

The resulting $z = 0$ final charges for non-maximally spinning black holes are shown in the top panel of Figure 2 as solid, dashed and dot-dashed black and grey lines, for different dark matter halo masses M_{vir} , different baryon densities and different dimensionless spin parameters that span a range that brackets the Thorne limit. The halo masses, baryon densities and spins are presented in the figure key and in the caption. The black dotted diagonal line shows the A23 estimate for the maximum charge a non-maximally spinning BH could acquire from the halo plasma². The figure follows the style of A23 showing for reference the Schwinger charges for pair production and rapid discharge [56] in orange and light green dashed lines, the RN extremal charge (red), the electron charge, minimum mass for primordial black holes, evaporated mass by today and stellar mass remnants (dotted horizontal, and left, centre and right vertical lines, respectively). The black circle and cyan lines show the charge for maximally spinning black holes for the extremal RN and Poisson initial charge cases, respectively.

As can be seen, when the spin is non-maximal, as soon as the black hole mass is such that its charge rises a few orders of magnitude above the pair production charge (orange) the Hawking athermal emission kicks in and prevents the charge from increasing further for higher black hole masses, which makes the final equilibrium charge to depart from the A23 maximum charge. Black holes in halos/galaxies of lower mass acquire lower final charges (dashed black and grey lines), a difference that is only seen at lower black hole masses where the Hawking emission is thermal.

The bottom panel shows that when the plasma density is larger, the accretion increases and allows a larger black hole equilibrium charge. An increase in the baryon density of 6 orders of magnitude with respect to the average virial density, produces an increase in the charge of almost one order of magnitude (see also the dot-dashed lines in the upper panel).

2.5 Required charges to provide cosmic magnetic seeds

One can compare the black hole charges to the ones needed to produce a magnetic field at early times, $z \sim 15$, with the required amplitude to explain present-day galaxy-wide cosmic magnetic fields. The seeds need to reach roughly $B_{\text{seed}} \sim 10^{-30}G$ [63]. The astrophysical processes that take place afterwards are able to increase the magnetic field by more than 20 orders of magnitude, reaching the expected $B_{\text{fil}} \sim 10^{-9}G$ in cosmic filaments or $B_{\text{gal}} \sim 10^{-9}G$ in galactic halos [64].

²We have chosen not to consider the case when a fraction of black holes hold a charge $Q = e$ presented in A23 simply because it would not be able to produce a galaxy-wide magnetic field.




Figure 2. Top: black hole charge in units of e as a function of its mass at $z = 0$. The cyan line shows the initial Poisson charge for primordial black holes, the red line the maximum charge for a RN extremal black hole. The circle shows the RN extremal charge for Planck mass primordial black holes. The dashed orange line shows the charge for pair production by Schwinger and/or Hawking (athermal) radiation. The dashed light green line shows the rapid discharge limit of [56]. Dark green lines show the charge needed to produce a magnetic field seed of $10^{-30}G$ for different halo masses (decreasing towards the top). The horizontal dotted line shows $Q = e$, and the vertical dotted lines show the black hole mass formed at reheating (left) the mass of a primordial black hole that would evaporate by today (middle), and the approximate mass of black holes of stellar origin (right). The black solid and dashed lines show the black hole charge acquired from the ionised baryonic plasma (for haloes of virial mass $M_{\text{vir}} = 10^5$ and $10^{10}h^{-1}M_{\odot}$, respectively) for a dimensionless spin parameter of $a = 0$ and for the average virial overdensity. The grey lines show the same for $a = 0.999999$. Dot-dashed lines show the charge for plasma densities of 10^6 times the virial density. Bottom: Ratio of the black hole charge embedded in a plasma as a function of density in units of the virial one at $z = 0$ for $a = 0$ and 0.999999 (black and grey, respectively).

We make a simple calculation assuming that all of the dark matter is made of black holes

and that these are distributed in an NFW profile with dimensionless halo spin parameter $\lambda' = J_{\text{vir}}/(M_{\text{vir}}V_{\text{vir}})$, where J_{vir} and V_{vir} are the angular momentum and virial velocity of the halo, respectively [65], with which the amplitude of the magnetic field at the virial radius, as a function of the mass of the black holes that make up the halo, reads,

$$B(r_{\text{vir}}) = \frac{\mu_0}{4\pi} \frac{m(r_{\text{vir}})}{r_{\text{vir}}^3} = \frac{\mu_0}{4\pi} \frac{1}{2} \frac{Q}{M} \lambda' \sqrt{2} \frac{M_{\text{vir}} V_{\text{vir}}}{r_{\text{vir}}^2} \quad (2.16)$$

where μ_0 is the magnetic permeability of the vacuum, m is the magnetic moment, r_{vir} is the virial radius, and we have assumed $\log_{10} \lambda' = -1.5$ [65]. The required charge per mass can then be written as

$$Q/M = \frac{8\pi B_{\text{seed}} r_{\text{vir}}^2}{\sqrt{2} \mu_0 \lambda' M_{\text{vir}} V_{\text{vir}}}, \quad (2.17)$$

for black holes with equal sign charge (i.e. for comparison to halo acquired charges only). The dark green lines in the top panel of Fig. 2 show these charges for different dark matter halo masses ranging from 10^5 to $10^{14} h^{-1} M_{\odot}$ (top to bottom lines), which lie close and above Q_{pairs} and roughly among the charges that black holes can acquire within dark matter haloes (black and grey lines).

Notice though, that this simple calculation does not take into account whether black holes with these charges would be able to produce a galaxy/halo wide non-zero charge and magnetic fields, or take into account the population of haloes at redshifts as high as $z = 15$. We tackle these details next.

3 Net charge of dark matter haloes

We now explore galaxy wide consequences of black holes containing electric charge. We will treat separately the case of primordial black holes which are able to preserve the charge with which they are formed, that A23 propose as being the case for maximally rotating primordial black holes (see for instance [66] for a discussion of spin of primordial black holes) which could contain positive and negative charges of the amplitudes shown by the circle and cyan lines of the top panel of Fig. 2. In principle these black holes do not exchange charge with the ionised plasma, leaving the latter perfectly neutral in a global sense. On the other hand, non-maximally rotating black holes of primordial or astrophysical origin, could extract a negative charge from the plasma, leaving the latter positively charged, but still leaving each individual dark matter halo or galaxy perfectly neutral. We explore these possibilities in the following subsections.

3.1 Dark matter halo charge from maximally spinning primordial black holes

In this subsection we will assume that primordial black holes (PBH) of equal mass conform all of the dark matter, for simplicity. If primordial black holes form from the initial fluctuation field left over from Inflation (see for instance [14]), then they can be assumed to be randomly distributed at late times when dark matter halo formation starts. In this case, one immediate question is whether the overdense patch containing the black holes and baryons that will collapse to form a halo, is electrically neutral. This would indeed be the case if the charged black holes were able to conform a plasma.

In order to test this we make a simple calculation of the potential Gravitational and Coulomb energy exerted between PBHs of Planck mass holding the extremal RN charge,




Figure 3. Comparison between the available total (Coulomb and gravitational) and gravitational energy (black and red dashed lines, respectively) exerted by Planck mass black holes with maximal RN charge, that lie within overdense patches that will collapse to form dark matter haloes, and the energy equivalent of the expansion of the universe on the same scales (black solid), at a scale of the mean separation between black holes. Since Coulomb and gravity cannot overcome the expansion, it can be safely assumed that charged, maximally spinning black holes with positive and negative initial charges, cannot relax into a plasma and their charges will remain unrelaxed and essentially randomly distributed in space down to the present day.




Figure 4. Halo charge per mass in primordial or astrophysical black holes at $z = 0$, as a function of black hole mass. Red and cyan show the net halo charge due to Poisson noise in the number of positive and negative maximally spinning primordial black holes within the overdense patch that gave rise to the formation of the halo (solid and dashed correspond to different halo masses as shown in the key). Black and grey lines show the net charge in non-maximally spinning black holes alone, considering two different values for the dimensionless spin parameter. The orange line shows the halo charge in black holes considering the pair production black hole charge Q_{pairs} .

which are the ones that present the highest Coulomb to gravitational force ratios, and would therefore present the highest chances of forming a plasma. To do the calculation we take

into account the mean physical distance between these PBHs as a function of redshift, and compare the Coulomb and gravitational potentials at these separations with the required kinetic energy that the PBHs would need to acquire in order to offset the expansion of the Universe on the same scales. This is shown in Fig. 3 where regardless of the redshift, the Coulomb and gravitational potentials are at least a factor of a few below the needed levels to overcome expansion.

Armed with this result, we can postulate that regardless of origin and mass, primordial black holes will not be able to relax into a plasma configuration, and therefore dark matter haloes will be endowed with a net charge arising from the Poisson imbalance of primordial black holes with positive and negative charges, $Q(M)$, within the initial overdense patch from which the halo collapsed. We will refer to this halo charge as the halo initial Poisson charge,

$$Q_p^i(M_{\text{vir}}) \sim Q(M) \sqrt{M_{\text{vir}}/M}. \quad (3.1)$$

The resulting halo charges can be of both positive and negative sign, and are shown as red symbols and cyan lines in Fig. 4, for two different halo masses with different line types.

3.2 Net charges and magnetic moments in galaxies with non-maximally spinning black holes

In the case of non-maximally spinning black holes, their negative charge is acquired from the plasma, which makes the halo electrically neutral. Excluding any high energy processes that could expel baryon charges away from the halo, or any intrinsic sources of charge imbalance within galaxies [67, 68], the total charge in black holes within a halo equals that in the plasma or the baryons in general within that same halo with the opposite sign. In this case the total halo charge in black holes is simply the product of the number of black holes in the halo and of their charge, and is shown for $z = 0$ as black and grey lines in Fig. 4 for different halo masses and black hole dimensionless spin parameters, as shown in the key. As it can be seen, the charge is rather constant since when $Q > 0$ the scaling of the black hole charge is $\propto M$. The halo charge is higher for higher black hole spins at both the low and high black hole mass ends. The amplitude of the charge can be similar to that obtained with maximally spinning PBHs, depending on the halo mass and black hole spin. The orange dotted line shows the charge in black holes considering a charge of Q^{pairs} which, as postulated above, is the minimum charge that black holes with $M > 10^{22}kg$ (a value that depends on the plasma density) could be seen to hold from afar, even if their equilibrium charge was higher.

If black holes are extracting charge from the plasma, this could be at odds with the typically assumed global neutrality of the plasma in galaxies. Fig. 5 shows the ratio of charges in black holes taken from the plasma and the total number of protons in the plasma, as a function of black hole mass, for different black hole spins and halo masses (black and grey, solid and dotted lines as indicated in the key), and also for the case when black holes are seen from afar to hold only Q^{pairs} (orange). As can be seen, even in the most extreme cases this ratio is below 10^{-30} , which is a very low level of non-neutrality that does not preclude assuming neutrality for the plasma.

If the halo is neutral it is interesting to wonder if it would be possible for longevous charges in black holes (or simply Q^{pairs}), to produce a halo wide non-zero net magnetic field. Considering that black holes are collisionless, whereas the plasma is pressure supported, one can expect there to be a net magnetic field simply due to the different density profiles of collisionless and collisional fluids. Assuming a Navarro, Frenk and White [69] profile (NFW) with a concentration vs. halo mass and redshift from [70] for the case of primordial black




Figure 5. Ratio of charges stored in black holes and in the halo plasma, as a function of black hole mass, for different halo masses, black hole spins and for Q_{pairs} , as shown in the key.

holes, and a core profile for the baryons [61] assumed to form a singular isothermal sphere [71], we calculate the magnetic moment that each fluid would produce, as a function of distance from the halo center,

$$m(r) = \left| \frac{1}{2} \int_V \vec{r} \times (\rho_Q \vec{v}) dV \right|, \quad (3.2)$$

where we assume a specific angular momentum that is equal for ionised baryons and black holes, and simplify the calculation taking purely circular motions. Here we can ignore the actual value of the charges and assume that each fluid contains the same total charge but of opposite sign (i.e. magnetic moments pointing in exact opposite directions, with different amplitudes as a function of distance to the center of the halo/galaxy). The resulting magnetic moments show a very mild dependence with the distance to the galaxy center for the gas and dark matter. With the aim to find whether the total magnetic moment of a galaxy is nonzero, we show the resulting radial dependence of the modulus ratios of gas to black hole magnetic moments as the black solid and dotted lines in Fig. 6. We show this ratio for two redshifts for a final $z = 0$ halo mass similar to that of the Milky Way, and in both cases we find that the amplitude of the magnetic moment for each fluid is different, except slightly above the virial radius r_{vir} (equal amplitudes imply zero net magnetic field as the magnetic moments of both fluids are assumed to be directed in opposite directions).

This result allows us to simply report the magnetic moment of black holes with NFW profiles as an order of magnitude estimate of the net magnetic moment of the halo in the following Subsection. If needed, the reader can also obtain more accurate estimates as a function of distance to the halo/galaxy centre with the ratios shown in this figure.

We also explore the possibility that there is a net magnetic moment due to astrophysical black holes (SBH) in the stellar halo of a galaxy that acquire a charge via their interaction with the ionised plasma within the halo. In this case we adopt the measured logarithmic slope of the density profile of halo stars in the Galaxy from Slater et al. [72] of -3.5 . Again we assume that the amount of charge is the same but of opposite sign for astrophysical black holes and the plasma. Even though the spatial extent of the stellar halo is in principle more confined than that of the hot plasma or the dark matter, a large fraction of the stellar (and




Figure 6. Ratio of the magnetic moment produced by the baryon plasma and by the black holes, for a Milky Way sized halo at $z = 0$ and $z = 15$ (solid and dotted lines, respectively) that is overall electrically neutral, as a function of radius from the halo center. We show in black the case of primordial black holes that follow a Navarro, Frenk and White profile and, in grey, stellar black holes that follow a stellar halo profile. The plasma for the calculation of both ratios is assumed to follow a cored isothermal profile.

stellar products) content of the halo are thought to have formed *ex situ*, i.e. outside the galaxy, and to have been deposited in the halo as a result of mergers (see for instance, Pilepich & Madau [73]). Therefore, stellar halo black holes could have acquired a charge during their infall process as they came into contact with the ionised plasma of their new host galaxy. This justifies assuming the charge to have the full extent of the distribution of gas and stellar black holes. In this case the magnetic moment has a steeper drop ($\propto r^{-1.45}$) compared to NFW distributed BHs or the isothermal ionised gas of the galaxy. The ratio of magnetic moments between plasma and SBH shown as a grey line in Fig. 6 indicates that it is indeed possible for SBHs to produce a non-zero net magnetic moment within galaxies, except at roughly 1/10 of the virial radius.

4 Seeds for the cosmic magnetic fields

With the ability to obtain non-zero net magnetic moments for all black hole types considered here, we proceed to estimate the total magnetic fields that dark matter haloes/galaxy hosts could produce right after formation (we pick $z = 15$), when the gas has already thermalised and conforms a plasma from which non-maximally spinning black holes can acquire a charge. We also look at the possibility that dark matter haloes contain a net charge due to the Poisson-like distribution of maximally spinning primordial black holes. We assume that PBHs conform all of the dark matter whereas when considering SBH we assign them a fraction of 10^{-3} of the total mass of the halo. We adopt this fraction since we focus here on galaxies such as the Milky Way which are shown to hold the highest efficiency for star formation turning about 10 to 50% of their total baryonic matter into stars [74], a relation that should hold constant with redshift for the galaxy to be this efficient even today [75]. We further consider that stars with masses $> 20M_{\odot}$ end their lives leaving a black hole remnant which translates




Figure 7. Magnetic field amplitude in haloes as a function of black hole mass at $z = 15$, for Poisson halo charge from maximally rotating primordial black holes (cyan lines) and RN extremal PBHs (red symbols), and for haloes with non-maximally rotating primordial black holes that acquired charge from the halo plasma (black solid and black and grey dotted lines) for halo masses and black hole spins as indicated in the key. The orange line shows the case where the seed is provided by the spatially displaced Q_{pairs} charges by black holes. The violet symbol shows the seed that stellar black holes in the stellar halo could provide if their charge reached Q_{pairs} . The horizontal grey line shows a constant magnetic field amplitude at $10^{-30}G$, the required seed magnetic field to explain the observed present-day cosmic magnetic fields.

into $\sim 7\%$ of the stellar mass converted into the latter when adopting a Salpeter [76] initial mass function for stellar masses. These three factors together result in a fraction of mass in black holes in galaxies of $\sim 10^{-3}$ as adopted here, and we assume that this fraction holds since the moment the galaxy forms. This is conservative as the very first stars could have a top-heavy initial mass function producing even higher mass fractions of high mass stars and, consequently, black holes [77–79]. The resulting amplitude of the seed magnetic field is directly proportional to this fraction.

The approach we take here is to look at the seed magnetic field within haloes/galaxies, i.e. in peaks of the density field. This can then be used to infer the magnetic field seed on any scale following an approach similar to that presented in A20. Instead of looking at the field produced in peaks of the density field as we do here, A20 propose that the magnetic field is proportional to the density field of matter, which allows them to obtain the seeds for any particular scale. We leave the study of seed magnetic fields from charged BHs at different cosmological scales to a forthcoming paper and concentrate on galaxy-wide magnetic seeds here.

We will focus on (i) the magnetic fields produced by primordial black holes that make up all of the dark matter, distributed in an NFW profile with the characteristics laid out in Subsection 2.5, where Eq. 2.16 gives the resulting magnetic field dipole amplitude at the virial radius r_{vir} , and (ii) the ones produced by stellar black holes, for which we assume the density profile of a stellar halo, with logarithmic slope of -3.5 , and use the ratio between the magnetic moment of stellar black holes and that of the plasma to make an estimate of the magnetic field amplitude in this case, considering a fraction of the total mass of the galaxy in black holes of 10^{-3} .

The resulting seed magnetic field amplitudes in haloes/galaxies as a function of black hole mass can be seen in Figure 7, calculated at $z = 15$ and $r = r_{\text{vir}}$ for PBH and $r \sim 0.05r_{\text{vir}}$ for SBH. The red symbols and cyan lines show the magnetic field when the dark matter is fully made by maximally spinning primordial black holes (for black holes from high energy collisions and for collapse at horizon entry, respectively), for haloes of different mass shown with different line types, as indicated. We choose to show the results for haloes that bracket the possible progenitors of Milky Way-like galaxies, which from simulation results are expected to have a virial mass of $\sim 10^9 M_{\odot}$ at $z = 15$ [80]. Black and grey lines show the magnetic field amplitude for non-maximally spinning primordial black holes (that again are assumed to form all of the dark matter). These values should be compared to the horizontal grey line, at $B = 10^{-30} G$ which is the seed value reported for instance in Davis et al. (1999, [63], see also [81]), who also point out that an acceptable seed could be of lower amplitude. As can be seen, if maximally spinning PBHs were a large fraction of the dark matter, regardless of their mass they could provide reasonable seeds. Non-maximally spinning BHs could also provide the seeds for black hole masses $M > 10^{22} kg$ since these can hold a charge, specially in high plasma density in large haloes at early times, even if a distant observer could only detect a fraction of the black hole charge corresponding to Q^{pairs} (dotted orange line).

In the case of zero spin stellar black holes (shown by the open violet symbol, placed at $M = 10M_{\odot}$) the possible seed lies ~ 4 orders of magnitude below the Davis et al. seed value, and corresponds to a distance to the galaxy centre of $\sim 0.05r_{\text{vir}}$. Even though this seed value is rather low, Davis et al. point out that it is possible that the required seed can be lowered due to uncertainties in the e-folding timescale for the dynamo processes within galaxies. On the other, the magnetic moment for stellar halo black holes decreases with distance to the galaxy center roughly as $\propto r^{-1.45}$, i.e. slower than the drop of a dipolar field ($\propto r^{-3}$), which means that nearer the center of the galaxy the amplitude of the magnetic field from SBH can be a few orders of magnitude higher, and this inner magnetic field could be transported outwards via AGN radio lobes [82]. Furthermore, allowing for a fraction of black holes to be part, at some point in their evolution, of X-ray binaries would imply that their average spin would be higher [33], which would also increase their average charges and the magnetic field seed they would be able to produce. It would be quite interesting to be able to explain the seeds of cosmic magnetic fields with the acquired charge of relic black holes of stellar origin.

5 Discussion

In this section we explore in more detail some caveats that could affect our conclusions.

We start looking at the effect that accretion discs could have on the charge of the black holes, which we have ignored up to this point. We expect accretion discs to be able to form around black holes when the Bondi radius [83] $r_B = 2GM/\tilde{v}^2$ (where \tilde{v} is the sum in quadrature of the sound speed and the virial velocity) exceeds the Debye length of the plasma. Since the main conclusions here are centred on high redshift haloes, we compare these two quantities for $z = 15$ and $M_{\text{vir}} \sim 10^{10} M_{\odot}$. In this case, assuming a fully ionised plasma at the virial temperature, we find that black holes with $M > 10^{22} kg$ would be able to form accretion discs. If the radiation field within the halo, coming either from the accretion onto a budding supermassive black hole or from intense bursts of initial star formation, is able to increase the plasma temperature, this would increase the threshold mass for accretion discs. For a temperature 1000 times higher [84], the minimum mass becomes $M \sim 10^{27} kg$, which

is below the solar mass. Therefore, it is quite plausible that the charges of BHs below this threshold mass are free of the effect of an accretion disc.

We have also mentioned earlier that the charge of BHs could be neutralised by a trailing group of protons with equal charge. We remind the reader that we consider that BHs and the plasma are dynamically independent. Since BHs with $M > 10^{22} - 10^{27} \text{kg}$ and higher (depending on the plasma density and redshift) could form an accretion disc, they would in principle be able to take along charges of opposite sign after them simply because these would be confined within the Bondi mass bound to the BH. This could indeed neutralise the charge as seen by a distant observer. However, if charges were acquired faster than the binding time of the Bondi mass, then the gravitational field of the black hole would not have enough time to take along the compensating protons; they would be left behind trapped in the plasma. We take the free fall timescale for the Bondi mass for BHs in a plasma with ρ_{vir} , $t_B = \sqrt{r_B^3/(2GM)}$, as the binding timescale for the Bondi mass and show its dependence on black hole mass as the violet dotted line in Fig. 1 (the line starts at the minimum BH mass that can form an accretion disc for a surrounding plasma at T_{vir}). As it can be seen, Q_{pairs} can be acquired much faster for $M > 10^{28} \text{kg}$, suggesting that no compensating charges will be carried along by the BH for this charge. Longevous charges could be affected, as the Bondi timescale is shorter than the charging time for lower mass BHs.

One important possibility that could discharge black holes completely is that the accretion disc around stellar or primordial black holes were energetic enough to produce positron-electron pairs; the emitted positrons would be able to discharge the black hole. Laurent et al. [85] show that the emission in some cases is as high as $> 10^{35}$ positrons per second. However, Sarkar & Chattopadhyay [86] show that the positron emission is non-zero only when the Eddington normalised accretion rate is $\dot{M}/\dot{M}_{Edd} > 0.8$, where $\dot{M}_{Edd} = 4\pi GMm_p/(c\sigma_T)$, with σ_T the Thompson scattering cross-section.

We calculate the Bondi-Hoyle-Lyttleton [83, 87, 88] accretion rates,

$$\dot{M} = 4\pi r_B^2 \tilde{v} \rho = \frac{4\pi G^2 M^2 n \mu m_p}{\tilde{v}^3} \quad (5.1)$$

where μ is the mean molecular weight and n is the gas number density. We find that all black holes with $M < 10^{34} \text{kg}$ (i.e. $\sim 10^4 M_\odot$) in our calculations accrete well below 10 percent of the Eddington limit as long as the baryon density around black holes is $< 10^6 \Omega_b \rho_{vir}$ (the density of molecular clouds, for example), indicating that the charges and seed magnetic fields estimated in this paper should not be affected by positron emission in accretion discs, i.e., it is reasonable to assume that accretion discs do not affect the charges of BHs with $M \lesssim 10^4 M_\odot$. The exception would be stellar black holes in high mass X-ray binaries, but only while they accrete from their companion. Once this process stops, their accretion rates fall below the threshold for electron-positron spontaneous emission. On the other hand, supermassive black holes (even those orbiting within the galaxy, coming from merged satellites) would not be expected to hold a non-zero electrical charge.

Another point to discuss is the spatial scale of the magnetic field seeds inferred here. We have concentrated on virialised objects, i.e. on the magnetic fields within peaks in the density field. This can be used to obtain a power spectrum for the fluctuations of the magnetic field to compute the seeds on any scale, following for example A20. However, the estimates we presented here are useful for studies of the origin of the magnetic field within galaxies. Additionally, the fact that the seed fields are enclosed in virialised objects confirms the mea-

measurements by [89] and [90], allowing to reach the lower limit of 60% magnetic filling factor in cosmic voids.

We also point out that our calculation of magnetic field seeds assumes that the black holes hold charge throughout the spatial extent of a dark matter halo/galaxy. We have purposely ignored that the gas in a galaxy can be thought of as at least having two phases, one which is hot and ionised and a cold, neutral one. We have done so because we are assuming that it is enough for black holes to at least come into contact with the ionised phase at some point, and this is justified by concentrating on long-lived charges, on charges below the Gibbons limit, or on a fraction of the black hole charge corresponding to Q_{pairs} , given the fast rates of charge acquisition by black holes for all these cases. However, the effect of the BH charges on the frontier between neutral and ionised gas within galaxies needs to be modelled further. Our present calculations should be robust at least in regions lying within the ionised plasma, which can still imprint non-zero magnetic fields due to the different local steepness of the black hole and gas density profiles.

We end by noticing also that we have shown quite a broad range of halo masses in our estimates of magnetic field seeds and that the increase in magnetic field amplitude is smaller than that in halo mass. This indicates that our results are not too sensitive to the halo mass and, in turn, to the redshift where the seed is calculated. The magnetic field seed for a Milky Way-like progenitor would only grow by roughly three orders of magnitude from $z = 15$ to $z = 0$, whereas astrophysical processes can increase the amplitude of the magnetic field by tens of orders of magnitude [63]. This shows that charged black holes, even at the Gibbons limit, provide seeds of roughly the adequate amplitude independently of the exact values adopted for redshift and halo mass.

6 Conclusions

In this paper we extend the formalism presented in Araya et al. (2023, [23], A23) estimating the charge that non-maximally spinning black holes embedded in a baryonic plasma are able to attain. In particular we study the rates of Hawking thermal charge emission, and of charge accretion from the surrounding ionised plasma, which allows us to estimate an equilibrium charge and its dependence on the black hole spin for dimensionless spin parameters $a < 1$. We find that black holes either surrounded by high density plasma or with high spins, are indeed able to surpass the Gibbons 1975 [56] limit of discharge by pair particle production Q_{pairs} for black holes of $M > 10^{22}$ kg. This is mostly due to the limiting cross sections for charge carriers which on the lower limit have a radius of the black hole radius, and on the upper limit the Debye length; in a galactic plasma both are of the same order of magnitude for these black hole masses. We also find that lower mass black holes would not hold a charge for a significant fraction of the time.

We also note the distant observer would only be able to infer at most charges that are longevous or a charge of Q_{pairs} for a higher equilibrium charge, since this is the only charge that black holes would be able to effectively spatially displace from the baryon plasma. This formalism can be applied indistinctively to primordial and astrophysical black holes.

We also explore the net charges and magnetic moments of galaxies and the dark matter haloes that host them, considering that primordial black holes make up all of the dark matter. In this case, if the black holes are maximally spinning and retain the charge they formed with, or even if their spin is non-maximal and the charge is acquired from the plasma, then their charge combined with the typical angular momentum of galaxies and dark matter haloes is

enough to produce magnetic fields of $\sim 10^{-30}$ Gauss for black hole masses $M > 10^{26}$ kg (even for $M > 10^{22}$ kg for high density plasmas). Notice that the current mass windows where primordial black holes can make up all of the dark matter lie roughly at $M \sim 10^{10}$ and 10^{30} kg, i.e. asteroid and solar masses, respectively [20, 21], presenting primordial black holes of solar mass with any spin, and maximally spinning ones of asteroid mass, as possible candidates to produce the magnetic field seeds regardless of the existence of magnetic monopole charges within them (see also [51]).

Another interesting result is the following. Current numerical simulations of galaxy formation, as well as observational evidence, suggest that the stellar haloes of galaxies contain high fractions of ex-situ formed stars [72] and, consequently, black holes. Considering the leeway in the estimated magnetic field seeds needed to explain the observed ones today [63], we find that even astrophysical black holes in the stellar haloes of galaxies could provide the magnetic field seeds. To do so these black holes would indeed need to be accreted from merged galaxy satellites so that they would have been at some point embedded in the hot, ionised gas halo surrounding galaxies to acquire a charge; even if they are able to accrete just the Gibbons pair production charge Q_{pairs} . This charge is quite low, producing a Coulomb potential some ~ 20 orders of magnitude lower than the gravitational one regardless of black hole mass. This is probably the reason why it is commonly regarded that the black hole charge can be safely ignored. However, we have shown here that even such low level of stellar black hole charge is able to provide seeds of magnetic field close to the required values to explain the cosmic magnetic fields observed today.

Acknowledgments

We thank Juan Magaña, Joaquín Armijo, Marcelo Rubio, Anna Durrant and Patricio Colazo for helpful comments and discussions. NDP acknowledges support from a RAICES, a RAICES-Federal, and PICT-2021-I-A-00700 grants from the Ministerio de Ciencia, Tecnología e Innovación, Argentina. The work of IJA is funded by ANID FONDECYT grants No. 11230419 and 1231133, and by ANID Becas Chile grant No. 74220042. IJA also acknowledges funding by ANID, REC Convocatoria Nacional Subvención a Instalación en la Academia Convocatoria Año 2020, Folio PAI77200097. FAS thanks support by grants PIP 11220130100365CO, PICT-2016-4174, PICT-2021-GRF-00719 and Consolidar-2018-2020, from CONICET, FONCyT (Argentina) and SECyT-UNC. IJA is grateful to Andrei Parnachev and the School of Mathematics at Trinity College Dublin for their hospitality.

References

- [1] K. C. Freeman. On the Disks of Spiral and S0 Galaxies. *ApJ*, 160:811, June 1970.
- [2] V. C. Rubin, Jr. Ford, W. K., K. M. Strom, S. E. Strom, and W. Romanishin. Extended rotation curves of high-luminosity spiral galaxies. II. The anemic Sa galaxy NGC 4378. *ApJ*, 224:782–795, September 1978.
- [3] Shadab Alam, Metin Ata, Stephen Bailey, Florian Beutler, Dmitry Bizyaev, Jonathan A. Blazek, Adam S. Bolton, Joel R. Brownstein, Angela Burden, Chia-Hsun Chuang, Johan Comparat, Antonio J. Cuesta, Kyle S. Dawson, Daniel J. Eisenstein, Stephanie Escoffier, Héctor Gil-Marín, Jan Niklas Grieb, Nick Hand, Shirley Ho, Karen Kinemuchi, David Kirkby, Francisco Kitaura, Elena Malanushenko, Viktor Malanushenko, Claudia Maraston, Cameron K. McBride, Robert C. Nichol, Matthew D. Olmstead, Daniel Oravetz, Nikhil Padmanabhan, Nathalie Palanque-Delabrouille, Kaike Pan, Marcos Pellejero-Ibanez, Will J. Percival, Patrick

- Petitjean, Francisco Prada, Adrian M. Price-Whelan, Beth A. Reid, Sergio A. Rodríguez-Torres, Natalie A. Roe, Ashley J. Ross, Nicholas P. Ross, Graziano Rossi, Jose Alberto Rubiño-Martín, Shun Saito, Salvador Salazar-Albornoz, Lado Samushia, Ariel G. Sánchez, Siddharth Satpathy, David J. Schlegel, Donald P. Schneider, Claudia G. Scóccola, Hee-Jong Seo, Erin S. Sheldon, Audrey Simmons, Anže Slosar, Michael A. Strauss, Molly E. C. Swanson, Daniel Thomas, Jeremy L. Tinker, Rita Tojeiro, Mariana Vargas Magaña, Jose Alberto Vazquez, Licia Verde, David A. Wake, Yuting Wang, David H. Weinberg, Martin White, W. Michael Wood-Vasey, Christophe Yèche, Idit Zehavi, Zhongxu Zhai, and Gong-Bo Zhao. The clustering of galaxies in the completed SDSS-III Baryon Oscillation Spectroscopic Survey: cosmological analysis of the DR12 galaxy sample. *MNRAS*, 470(3):2617–2652, September 2017.
- [4] Agne Semenaite, Ariel G. Sánchez, Andrea Pezzotta, Jiamin Hou, Roman Scoccimarro, Alexander Eggemeier, Martin Crocce, Chia-Hsun Chuang, Alexander Smith, Cheng Zhao, Joel R. Brownstein, Graziano Rossi, and Donald P. Schneider. Cosmological implications of the full shape of anisotropic clustering measurements in BOSS and eBOSS. *MNRAS*, 512(4):5657–5670, June 2022.
- [5] Planck Collaboration. Planck 2015 results - xix. constraints on primordial magnetic fields. *A&A*, 594:A19, 2016.
- [6] N. Aghanim et al. Planck 2018 results. VI. Cosmological parameters. *Astron. Astrophys.*, 641:A6, 2020. [Erratum: *Astron. Astrophys.* 652, C4 (2021)].
- [7] Jonathan L. Feng. Dark Matter Candidates from Particle Physics and Methods of Detection. *ARAA*, 48:495–545, September 2010.
- [8] Giorgio Arcadi, Maíra Dutra, Pradipta Ghosh, Manfred Lindner, Yann Mambrini, Mathias Pierre, Stefano Profumo, and Farinaldo S. Queiroz. The waning of the WIMP? A review of models, searches, and constraints. *European Physical Journal C*, 78(3):203, March 2018.
- [9] Marc Schumann. Direct detection of WIMP dark matter: concepts and status. *Journal of Physics G Nuclear Physics*, 46(10):103003, October 2019.
- [10] Ya. B. Zel’dovich and I. D. Novikov. The Hypothesis of Cores Retarded during Expansion and the Hot Cosmological Model. *AZH*, 43:758, Jan 1966.
- [11] Stephen Hawking. Gravitationally collapsed objects of very low mass. *Mon. Not. Roy. Astron. Soc.*, 152:75, 1971.
- [12] S. W. Hawking. Particle Creation by Black Holes. *Commun. Math. Phys.*, 43:199–220, 1975. [Erratum: *Commun. Math. Phys.* 46, 206 (1976)].
- [13] A. Arbey and F. Mahmoudi. Dark matter and the early Universe: a review. *Prog. Part. Nucl. Phys.*, 119:103865, 2021.
- [14] Maxim Yu Khlopov. Primordial black holes. *Research in Astronomy and Astrophysics*, 10(6):495–528, May 2010.
- [15] R. Bernabei et al. First results from DAMA/LIBRA and the combined results with DAMA/NaI. *Eur. Phys. J. C*, 56:333–355, 2008.
- [16] Gregory D. Mack, John F. Beacom, and Gianfranco Bertone. Towards closing the window on strongly interacting dark matter: Far-reaching constraints from earth’s heat flow. *Phys. Rev. D*, 76:043523, Aug 2007.
- [17] Ivone F. M. Albuquerque and Carlos Pérez de los Heros. Closing the window on strongly interacting dark matter with icecube. *Phys. Rev. D*, 81:063510, Mar 2010.
- [18] Andrzej K. Drukier, Katherine Freese, and David N. Spergel. Detecting cold dark-matter candidates. *Phys. Rev. D*, 33:3495–3508, Jun 1986.

- [19] Bernard Carr, Kazunori Kohri, Yuuiti Sendouda, and Jun'ichi Yokoyama. Constraints on Primordial Black Holes. *arXiv e-prints*, page arXiv:2002.12778, February 2020.
- [20] Bernard Carr and Florian Kuhnel. Primordial Black Holes as Dark Matter: Recent Developments. *arXiv e-prints*, page arXiv:2006.02838, June 2020.
- [21] Joaquin Sureda, Juan Magana, Ignacio J. Araya, and Nelson D. Padilla. Press-Schechter primordial black hole mass functions and their observational constraints. *arXiv e-prints*, page arXiv:2008.09683, August 2020.
- [22] Nelson D Padilla, Juan Magaña, Joaquín Sureda, and Ignacio J Araya. Power spectrum of density fluctuations, halo abundances, and clustering with primordial black holes. *Monthly Notices of the Royal Astronomical Society*, 504(3):3139–3155, 04 2021.
- [23] I. J. Araya, N. D. Padilla, M. E. Rubio, J. Sureda, J. Magaña, and L. Osorio. Dark matter from primordial black holes would hold charge. *JCAP*, 2023(2):030, February 2023.
- [24] Julian B. Muñoz and Abraham Loeb. A small amount of mini-charged dark matter could cool the baryons in the early Universe. *Nature*, 557(7707):684–686, May 2018.
- [25] Andrea Caputo, Laura Sberna, Miguel Frías, Diego Blas, Paolo Pani, Lijing Shao, and Wenming Yan. Constraints on millicharged dark matter and axionlike particles from timing of radio waves. *PhRD*, 100(6):063515, September 2019.
- [26] Kip S. Thorne. Disk-Accretion onto a Black Hole. II. Evolution of the Hole. *ApJ*, 191:507–520, July 1974.
- [27] D. L. Meier. The Association of Jet Production with Geometrically Thick Accretion Flows and Black Hole Rotation. *ApJL*, 548(1):L9–L12, February 2001.
- [28] R. S. Nemmen, R. G. Bower, A. Babul, and T. Storchi-Bergmann. Models for jet power in elliptical galaxies: a case for rapidly spinning black holes. *MNRAS*, 377(4):1652–1662, June 2007.
- [29] Robert Popham and Charles F. Gammie. Advection-dominated Accretion Flows in the Kerr Metric. II. Steady State Global Solutions. *ApJ*, 504(1):419–430, September 1998.
- [30] Andrew J. Benson and Arif Babul. Maximum spin of black holes driving jets. *MNRAS*, 397(3):1302–1313, August 2009.
- [31] Alejo Martínez-Sansigre and Steve Rawlings. Observational constraints on the spin of the most massive black holes from radio observations. *MNRAS*, 414(3):1937–1964, July 2011.
- [32] Claudia Del P. Lagos, Nelson D. Padilla, and Sofía A. Cora. Black hole spin and radio loudness in a Λ cold dark matter universe. *MNRAS*, 395(2):625–636, May 2009.
- [33] Morgan MacLeod and Jonathan Grindlay. Observations of Stellar-Mass Black Holes in the Galaxy. *arXiv e-prints*, page arXiv:2304.09368, April 2023.
- [34] E. Battaner, E. Florido, and J. Jimenez-Vicente. Magnetic fields and large scale structure in a hot universe. I. General equations. *AAP*, 326:13–22, October 1997.
- [35] Kanhaiya L. Pandey, T. Roy Choudhury, Shiv K. Sethi, and Andrea Ferrara. Reionization constraints on primordial magnetic fields. *Monthly Notices of the Royal Astronomical Society*, 451(2):1692–1700, 06 2015.
- [36] Lawrence M. Widrow. Origin of galactic and extragalactic magnetic fields. *Rev. Mod. Phys.*, 74:775–823, Jul 2002.
- [37] A. M. Beck, M. Hanasz, H. Lesch, R. S. Remus, and F. A. Stasyszyn. On the magnetic fields in voids. *MNRAS*, 429:L60–L64, February 2013.
- [38] Philipp P. Kronberg. Intergalactic Magnetic Fields. *Physics Today*, 55(12):40, 2002.

- [39] John D. Barrow, Roy Maartens, and Christos G. Tsagas. Cosmology with inhomogeneous magnetic fields. *Physics Reports*, 449(6):131 – 171, 2007.
- [40] K. Subramanian. Primordial magnetic fields and cmb anisotropies. *Astronomische Nachrichten*, 327(5-6):403–409, 2006.
- [41] Alejandra Kandus, Kerstin E. Kunze, and Christos G. Tsagas. Primordial magnetogenesis. *Physics Reports*, 505(1):1 – 58, 2011.
- [42] Kandaswamy Subramanian. The origin, evolution and signatures of primordial magnetic fields. *Reports on Progress in Physics*, 79(7):076901, may 2016.
- [43] Dai G. Yamazaki, Toshitaka Kajino, Grant J. Mathews, and Kiyotomo Ichiki. The search for a primordial magnetic field. *Physics Reports*, 517(5):141 – 167, 2012. The search for a primordial magnetic field.
- [44] Dario Grasso and Hector R. Rubinstein. Magnetic fields in the early universe. *Physics Reports*, 348(3):163 – 266, 2001.
- [45] Arnon Dar and A. De Rújula. Magnetic field in galaxies, galaxy clusters, and intergalactic space. *Phys. Rev. D*, 72:123002, Dec 2005.
- [46] Jean M. Quashnock, Abraham Loeb, and David N. Spergel. Magnetic Field Generation during the Cosmological QCD Phase Transition. *ApJL*, 344:L49, September 1989.
- [47] Robi Banerjee and Karsten Jedamzik. Are cluster magnetic fields primordial? *Phys. Rev. Lett.*, 91:251301, Dec 2003.
- [48] Ruth Durrer. Is the mystery of cosmic magnetic fields solved? *Science*, 311(5762):787–788, 2006.
- [49] Shohei Saga, Kiyotomo Ichiki, Keitaro Takahashi, and Naoshi Sugiyama. Magnetic field spectrum at cosmological recombination revisited. *Phys. Rev. D*, 91:123510, Jun 2015.
- [50] Muni Zhou, Vladimir Zhdankin, Matthew W. Kunz, Nuno F. Loureiro, and Dmitri A. Uzdensky. Spontaneous magnetization of collisionless plasma. *Proceedings of the National Academy of Science*, 119(19):e2119831119, May 2022.
- [51] I. J. Araya, M. E. Rubio, M. San Martín, F. A. Stasyszyn, N. D. Padilla, J. Magaña, and J. Sureda. Magnetic field generation from PBH distributions. *MNRAS*, 503(3):4387–4399, May 2021.
- [52] Juan Maldacena. Comments on magnetic black holes. *Journal of High Energy Physics*, 2021(4):79, April 2021.
- [53] I. K. Baldry, S. P. Driver, J. Loveday, E. N. Taylor, L. S. Kelvin, J. Liske, P. Norberg, A. S. G. Robotham, S. Brough, A. M. Hopkins, S. P. Bamford, J. A. Peacock, J. Bland-Hawthorn, C. J. Conselice, S. M. Croom, D. H. Jones, H. R. Parkinson, C. C. Popescu, M. Prescott, R. G. Sharp, and R. J. Tuffs. Galaxy And Mass Assembly (GAMA): the galaxy stellar mass function at $z < 0.06$. *MNRAS*, 421(1):621–634, March 2012.
- [54] Michael Edward Peskin and Daniel V. Schroeder. *An Introduction to Quantum Field Theory*. Westview Press, 1995. Reading, USA: Addison-Wesley (1995) 842 p.
- [55] Julian Schwinger. On gauge invariance and vacuum polarization. *Phys. Rev.*, 82:664–679, Jun 1951.
- [56] G. W. Gibbons. Vacuum Polarization and the Spontaneous Loss of Charge by Black Holes. *Commun. Math. Phys.*, 44:245–264, 1975.
- [57] Benjamin V. Lehmann, Christian Johnson, Stefano Profumo, and Thomas Schwemberger. Direct detection of primordial black hole relics as dark matter. *JCAP*, 2019(10):046, October 2019.

- [58] O. Ruiz, U. Molina, and P. Vitoria. Thermodynamic analysis of Kerr-Newman black holes. In *Journal of Physics Conference Series*, volume 1219 of *Journal of Physics Conference Series*, page 012016, May 2019.
- [59] Brandon Carter. Global Structure of the Kerr Family of Gravitational Fields. *Physical Review*, 174(5):1559–1571, October 1968.
- [60] M.S. Longair. *Galaxy Formation*. Astronomy and Astrophysics Library. Springer Berlin Heidelberg, 1998.
- [61] Cedric G. Lacey, Carlton M. Baugh, Carlos S. Frenk, Andrew J. Benson, Richard G. Bower, Shaun Cole, Violeta Gonzalez-Perez, John C. Helly, Claudia D. P. Lagos, and Peter D. Mitchell. A unified multiwavelength model of galaxy formation. *MNRAS*, 462(4):3854–3911, November 2016.
- [62] Claudia Del P. Lagos, Nelson D. Padilla, Michael A. Strauss, Sofia A. Cora, and Lei Hao. Host galaxy-active galactic nucleus alignments in the Sloan Digital Sky Survey Data Release 7. *MNRAS*, 414(3):2148–2162, July 2011.
- [63] Anne-Christine Davis, Matthew Lilley, and Ola Törnkvist. Relaxing the bounds on primordial magnetic seed fields. *PhRD*, 60(2):021301, July 1999.
- [64] Franco Vazza, Nicola Locatelli, Kamlesh Rajpurohit, Serena Banfi, Paola Domínguez-Fernández, Denis Wittor, Matteo Angelinelli, Giannandrea Inchingolo, Marisa Brienza, Stefan Hackstein, Daniele Dallacasa, Claudio Gheller, Marcus Brüggem, Gianfranco Brunetti, Annalisa Bonafede, Stefano Ettori, Chiara Stuardi, Daniela Paoletti, and Fabio Finelli. Magnetogenesis and the Cosmic Web: A Joint Challenge for Radio Observations and Numerical Simulations. *Galaxies*, 9(4):109, November 2021.
- [65] Andrea V. Macciò, Aaron A. Dutton, Frank C. van den Bosch, Ben Moore, Doug Potter, and Joachim Stadel. Concentration, spin and shape of dark matter haloes: scatter and the dependence on mass and environment. *MNRAS*, 378(1):55–71, June 2007.
- [66] Mehrdad Mirbabayi, Andrei Gruzinov, and Jorge Noreña. Spin of primordial black holes. *Journal of Cosmology and Astroparticle Physics*, 2020(03):017–017, mar 2020.
- [67] C. Caprini and P. G. Ferreira. Constraints on the electrical charge asymmetry of the universe. *JCAP*, 2005(2):006, February 2005.
- [68] Cody Goolsby-Cole and Lorenzo Sorbo. On the electric charge of the observable Universe. *arXiv e-prints*, page arXiv:1511.07465, November 2015.
- [69] Julio F. Navarro, Carlos S. Frenk, and Simon D. M. White. A Universal Density Profile from Hierarchical Clustering. *ApJ*, 490(2):493–508, December 1997.
- [70] Aaron D. Ludlow, Sownak Bose, Raúl E. Angulo, Lan Wang, Wojciech A. Hellwing, Julio F. Navarro, Shaun Cole, and Carlos S. Frenk. The mass-concentration-redshift relation of cold and warm dark matter haloes. *MNRAS*, 460(2):1214–1232, August 2016.
- [71] Volker Springel, Simon D. M. White, Giuseppe Tormen, and Guinevere Kauffmann. Populating a cluster of galaxies - I. Results at $z=0$. *MNRAS*, 328(3):726–750, December 2001.
- [72] Colin T. Slater, David L. Nidever, Jeffrey A. Munn, Eric F. Bell, and Steven R. Majewski. The Stellar Density Profile of the Distant Galactic Halo. *ApJ*, 832(2):206, December 2016.
- [73] Annalisa Pillepich, Piero Madau, and Lucio Mayer. Building Late-type Spiral Galaxies by In-situ and Ex-situ Star Formation. *ApJ*, 799(2):184, February 2015.
- [74] Benjamin P. Moster, Thorsten Naab, and Simon D. M. White. Galactic star formation and accretion histories from matching galaxies to dark matter haloes. *MNRAS*, 428(4):3121–3138, February 2013.

- [75] Peter S. Behroozi, Risa H. Wechsler, and Charlie Conroy. On the Lack of Evolution in Galaxy Star Formation Efficiency. *ApJL*, 762(2):L31, January 2013.
- [76] Edwin E. Salpeter. The Luminosity Function and Stellar Evolution. *ApJ*, 121:161, January 1955.
- [77] D. Schaerer. On the properties of massive Population III stars and metal-free stellar populations. *AAP*, 382:28–42, January 2002.
- [78] D. Schaerer. The transition from Population III to normal galaxies: Ly α and He II emission and the ionising properties of high redshift starburst galaxies. *AAP*, 397:527–538, January 2003.
- [79] David Sobral, Jorryt Matthee, Behnam Darvish, Daniel Schaerer, Bahram Mobasher, Huub J. A. Röttgering, Sérgio Santos, and Shoubaneh Hemmati. Evidence for PopIII-like Stellar Populations in the Most Luminous Lyman- α Emitters at the Epoch of Reionization: Spectroscopic Confirmation. *ApJ*, 808(2):139, August 2015.
- [80] Onsi Fakhouri, Chung-Pei Ma, and Michael Boylan-Kolchin. The merger rates and mass assembly histories of dark matter haloes in the two Millennium simulations. *MNRAS*, 406(4):2267–2278, August 2010.
- [81] Amjad Ashoorioon and Robert B. Mann. Generation of cosmological seed magnetic fields from inflation with cutoff. *Phys. Rev. D*, 71:103509, May 2005.
- [82] P. P. Kronberg, Q. W. Dufton, H. Li, and S. A. Colgate. Magnetic Energy of the Intergalactic Medium from Galactic Black Holes. *ApJ*, 560(1):178–186, October 2001.
- [83] H. Bondi. On Spherically Symmetrical Accretion. *Monthly Notices of the Royal Astronomical Society*, 112(2):195–204, 04 1952.
- [84] Takumi Ohmura, Mami Machida, Kenji Nakamura, Yuki Kudoh, and Ryoji Matsumoto. Two-temperature magnetohydrodynamic simulations for sub-relativistic active galactic nucleus jets: dependence on the fraction of the electron heating. *MNRAS*, 493(4):5761–5772, April 2020.
- [85] Philippe Laurent and Lev Titarchuk. Electron-Positron Pair Creation Close to a Black Hole Horizon: Redshifted Annihilation Line in the Emergent X-Ray Spectra of a Black Hole. I. *ApJ*, 859(2):89, June 2018.
- [86] Shilpa Sarkar and Indranil Chattopadhyay. Pair production and annihilation in advective accretion disks around black holes. In *Journal of Physics Conference Series*, volume 1640 of *Journal of Physics Conference Series*, page 012022, October 2020.
- [87] H. Bondi and F. Hoyle. On the Mechanism of Accretion by Stars. *Monthly Notices of the Royal Astronomical Society*, 104(5):273–282, 10 1944.
- [88] F. Hoyle and R. A. Lyttleton. The effect of interstellar matter on climatic variation. *Mathematical Proceedings of the Cambridge Philosophical Society*, 35(3):405–415, 1939.
- [89] Andrii Neronov and Ievgen Vovk. Evidence for strong extragalactic magnetic fields from fermi observations of tev blazars. *Science*, 328(5974):73–75, 2010.
- [90] K. Dolag, M. Kachelriess, S. Ostapchenko, and R. Tomàs. Lower Limit on the Strength and Filling Factor of Extragalactic Magnetic Fields. *ApJL*, 727(1):L4, January 2011.