

Induced gravitational waves from flipped SU(5) superstring theory at nHz

Spyros Basilakos,^{1,2,3} Dimitri V. Nanopoulos,^{4,5,6} Theodoros Papanikolaou,^{7,1} Emmanuel N. Saridakis,^{1,8} and Charalampos Tzerefos^{1,9}

¹*National Observatory of Athens, Lofos Nymfon, 11852 Athens, Greece*

²*Academy of Athens, Research Center for Astronomy and Applied Mathematics, Soranou Efessiou 4, 11527, Athens, Greece*

³*School of Sciences, European University Cyprus, Diogenes Street, Engomi, 1516 Nicosia, Cyprus*

⁴*George P. and Cynthia W. Mitchell Institute for Fundamental Physics and Astronomy, Texas A&M University, College Station, Texas 77843, USA*

⁵*Astroparticle Physics Group, Houston Advanced Research Center (HARC), Mitchell Campus, Woodlands, Texas 77381, USA*

⁶*Academy of Athens, Division of Natural Sciences, Athens 10679, Greece*

⁷*Scuola Superiore Meridionale, Largo S. Marcellino 10, I-80138, Napoli, Italy*

⁸*CAS Key Laboratory for Researches in Galaxies and Cosmology, Department of Astronomy, University of Science and Technology of China, Hefei, Anhui 230026, P.R. China*

⁹*Department of Physics, National & Kapodistrian University of Athens, Zografou Campus GR 157 73, Athens, Greece*

The no-scale flipped SU(5) superstring framework constitutes a very promising paradigm for physics below the Planck scale providing us with a very rich cosmological phenomenology in accordance with observations. In particular, it can accommodate Starobinsky-like inflation, followed by a reheating phase, which is driven by a light "flaton" field, and during which the GUT phase transition occurs. In this Letter, we extract for the first time a gravitational-wave (GW) signal which naturally arises in the context of the flipped SU(5) cosmological phenomenology and is related to the existence of an early matter era (eMD) driven by the flaton field. Specifically, we study GWs non-linearly induced by inflationary perturbations and which are abundantly produced during a sudden transition from the flaton-driven eMD era to the late-time radiation-dominated era. Remarkably, we find a GW signal with a characteristic peak frequency $f_{\text{GW,peak}}$ depending only on the string slope α' and reading as $f_{\text{GW,peak}} \propto 10^{-9} \left(\frac{\alpha'}{\alpha'_*}\right)^4$ Hz, where α'_* is the fiducial string slope being related directly to the reduced Planck scale M_{Pl} as $\alpha'_* = 8/M_{\text{Pl}}^2$. Interestingly enough, $f_{\text{GW,peak}}$ lies within the nHz frequency range; hence rendering this primordial GW signal potentially detectable by SKA, NANOGrav and PTA probes at their very low frequency region of their detection bands.

Introduction – Superstring theory constitutes one of the most promising paradigms describing the underlying fundamental quantum "theory of everything" including gravity as well as the standard model (SM). If string theory describes nature unifying all fundamental interactions, it should be able to explain all physical phenomena at all scales. String phenomenology has made several steps towards this goal by string constructions of particle physics models and continue then with concrete models of cosmology [1–6]. Though, very little has been done in explicit constructions of both at the same time.

In this Letter, we focus on the flipped SU(5) superstring framework which can give us a simultaneous description of particle phenomenology and cosmology [7, 8]. In particular, it can naturally give rise to Starobinsky-like inflationary setups [9, 10], favored by CMB observations, and explain the neutrino mass generation through an inflaton-induced see-saw mechanism [11, 12], the baryon asymmetry in the Universe as well as the neutralino dark matter [13, 14].

Interestingly enough, during inflation, the vacuum remains in the unbroken GUT phase, and GUT symmetry breaking occurs later when a field with a flat direction (the flaton) acquires a vacuum expectation value. At the end, reheating is driven by the incoherent part of the flaton operating in the so-called strong

regime [12, 15, 16]. As the Universe cools down, the flaton becomes non-relativistic and eventually triggers an early matter-dominated era (eMD) [15–17].

During this eMD era [18, 19], the inflationary adiabatic perturbations can be significantly enhanced inducing at second order in cosmological perturbation theory an abundant production of gravitational waves [20–25]. These scalar-induced GWs (SIGWs), behaving as radiation, can contribute as well to the effective number of extra neutrino species ΔN_{eff} potentially relaxing in a natural way the Hubble tension [See [26, 27] for PBH induced GWs responsible for the alleviation of the Hubble tension].

It is these GWs that we study within this work. Remarkably, we find a peak GW frequency $f_{\text{GW,peak}}$ which depends only on the string slope α' and reading as $f_{\text{GW,peak}} \propto 10^{-9} \left(\frac{\alpha'}{\alpha'_*}\right)^4$ Hz, where α'_* is the fiducial string slope being related to the reduced Planck scale M_{Pl} as $\alpha'_* = 8/M_{\text{Pl}}^2$. Notably, $f_{\text{GW,peak}}$ lies within the nHz frequency range; hence rendering this primordial GW signal detectable by SKA, NANOGrav and PTA probes.

The flipped SU(5) superstring paradigm– The string-derived flipped SU(5) model, initially introduced in [1], constitutes a step towards a unified particle and cosmological phenomenology [7, 8] being built within the

framework of the free fermionic formulation (FFF) of four-dimensional (4d) heterotic superstrings [28–30]. The FFF has the advantage of calculability in perturbation theory around a vacuum where all moduli, except the dilaton, are fixed at the fermionic point by a set of gauge and discrete symmetries, such as flipped $SU(5) \times U(1)$ [31–33].

The choice of the sub-Planckian gauge group, i.e. flipped $SU(5) \times U(1)$, is not accidental. In weakly-coupled heterotic string compactifications the matter representations are limited in size, e.g. to $\bar{5}$ and $\bar{10}$ representations of $SU(5)$ [34]. This consideration motivates our choice for a GUT gauge group, namely the flipped $SU(5) \times U(1)$ [31–33] one, pretty uniquely as it can be broken down to the Standard Model (SM) $SU(3)_C \times SU(2)_L \times U(1)_Y$ group by a combination of $\bar{10}$ and 10 antisymmetric Higgs representations, whereas $SU(5)$ and larger GUT groups require adjoint or larger Higgs representations. The resemblance of the representation content of $SU(5) \times U(1)$ and that of the SM $SU(3)_C \times SU(2)_L \times U(1)_Y$ is rather striking. Interestingly enough, flipped $SU(5) \times U(1)$ offers a promising framework for supersymmetric grand unification that provides resolutions of several phenomenological issues in particle physics. For example, in addition to accommodating, naturally, light neutrino masses, it provides a minimal mechanism for splitting the masses of the triplet and doublet components of the fiveplets of Higgs fields and in such a way that avoids automatically the dimension-5 operator that leads to catastrophically fast proton decay, endemic in supersymmetric GUTs, like SUSY $SU(5)$ [33].

Recently, the particle physics and cosmology of the string derived flipped $SU(5) \times U(1)$ have been studied in great detail [7, 8]. Specifically, a set of vacuum expectation values (vevs) $\langle \phi_i \rangle$ are triggered by the breaking of an anomalous $U(1)_A$ gauge symmetry [35], $\langle \phi_i \rangle \sim \xi$ of roughly an order of magnitude below the string scale, which are induced in order to satisfy the F - and D -flatness superpotential conditions up to 6th order [8]. In particular, all extra color triplets become superheavy, guarantying observable proton decay stability while the Higgs doublet mass matrix has a massless pair eigenstate with realistic hierarchical Yukawa couplings to quarks and leptons [8]. Actually, more than thirty years ago, this string derived flipped $SU(5) \times U(1)$ model [1] predicted that the mass of the top quark would be around $\sim 170 - 180\text{GeV}$ (see fig.1 of [1]) as was observed at Fermilab in 1995. The stringy derived masses of quarks and leptons and their relations (at the string scale) are summarized below [8] e.g.

$$\begin{aligned} m_c &\sim \xi^2 m_t, \quad m_s \sim \xi^2 m_b, \\ m_u &\sim \xi^5 m_t, \quad m_d \sim \xi^3 m_b, \quad m_e \sim \xi^4 m_t \end{aligned} \quad (1)$$

with ξ dynamically calculated to be $\xi = M_s/2\pi$ with M_s the string scale $M_s \equiv 1/\sqrt{2a'}$, with a' the string slope,

and thus, in *string units*, $\xi \approx \mathcal{O}(1/10)$.

The flipped $SU(5)$ superstring cosmology– It has been shown [7] that the string derived no-scale flipped $SU(5) \times U(1)$ can accommodate also a successful cosmology, based on the no-scale supergravity [36–40] derived from string theory [7, 41–43] as well as an appropriate induced superpotential suppressed by five powers of the string scale [7]. It utilises two gauge singlet chiral superfields present in the low energy spectrum, namely the inflaton field y , identified as the superpartner of a state mixed with right-handed neutrinos, and the goldstino field z with a superpotential of the form $W_I = M_I z (y - \lambda y^2)$, where λ is a dimensionless $\mathcal{O}(1)$ parameter. $M_I \sim 10^{13}\text{GeV}$ is the mass scale of inflation, being generated at the 5th order by the breaking of an anomalous $U(1)_A$ gauge symmetry of the heterotic string chiral vacuum [7].¹ The resulting scalar potential leads to Starobinsky-type inflation with a scalar spectral index $n_s \simeq 0.965$ and a tensor-to-scalar ratio $r = 2.4 \times 10^{-3}$ [7] in excellent agreement with Planck [44].

Furthermore, it should be stressed that during inflation and subsequent reheating, the GUT symmetry $SU(5) \times U(1)$ is left unbroken [12, 15, 16, 45]. The $SU(5) \times U(1)$ symmetry is broken by a 10 and $\bar{10}$ Higgs fields, H and \bar{H} respectively, along the F - and D - flat directions $\langle \tilde{\nu}_H^c \rangle = \langle \tilde{\nu}_{\bar{H}}^c \rangle \neq 0$, where ν_H^c and $\nu_{\bar{H}}^c$ are the SM singlets in the flipped 10 and $\bar{10}$ representations respectively. These vevs, which can be naturally large thanks to the F - and D - flat directions, are induced by the soft supersymmetry breaking masses. The resulting symmetry breaking pattern is $SU(5) \times U(1) \rightarrow SU(3)_C \times SU(2)_L \times U(1)_Y$.² After H and \bar{H} develop vevs, a SM singlet, a linear combination of ν_H^c and $\nu_{\bar{H}}^c$, appears as a physical state, massless in the supersymmetric limit, due to the presence of the F - and D -flat direction in the potential, and has a mass of the order of the soft supersymmetric-breaking mass scale. We denote this combination by Φ and refer to it as the *flaton* [12, 15, 16]. At low energies the soft supersymmetric breaking scale, i.e. the flaton mass-squared, m_Φ^2 , is driven negative by renormalisation group equation effects (RGE) due to relevant Yukawa couplings [33]. This negative mass-squared term destabilizes the origin of the flat direction, and thus the flaton field develops a vev, breaking the $SU(5) \times U(1)$ GUT symmetry into the SM group.

¹ Note that $M_I \simeq C_6 a_s^5 M_s \sim 10^{13}\text{GeV}$ and $\lambda \simeq g_s \frac{M_I}{M_s} \sim \mathcal{O}(1)$, with C_6 the numerical value of the correlator associated to the $N = 6$ coupling and $a_s \equiv g_s/2\pi$, with g_s the string coupling constant.

² Notice that this symmetry breaking pattern is *unique*, contrary to the case of SUSY GUTs, which have degenerate vacua, e.g. $SU(5)$ may be broken into other gauge groups, as $SU(4) \times U(1)$ [46]. We also mention that flipped $SU(5) \times U(1)$ is free from any monopole problem, even if it is broken after inflation, since the $SU(5) \times U(1)$ group is not simple [32].

Notice that if the height of the potential barrier between the symmetric ($SU(5) \times U(1)$) GUT and the Higgs ($SU(3)_C \times SU(2)_L \times U(1)_Y$) phase of the theory is smaller than the thermal kinetic energy of Φ , the incoherent part of Φ will drive the transition [15, 17], destroying any coherent contribution and displacing Φ to the low-energy vacuum within a Hubble time [15, 17]. We call this scenario strong reheating [12, 15, 16], during which a substantial amount of entropy release of $\mathcal{O}(10^4)$ reduces the cosmological baryon asymmetry and the density of the neutralino dark matter so as to be compatible with Planck [44]. In particular, as the Universe cools down, the energy density of Φ will simply redshift as radiation, until the temperature of the Universe decreases at such level that the interactions keeping the flaton in thermal equilibrium cease to be efficient. At even later times, when $m_\Phi > T_\Phi$, the flaton becomes non-relativistic and eventually dominates the energy budget of the universe, leading to an early matter dominated era (eMD) [15, 16].

This transient eMD era lasts until the end of the flaton decay, which proceeds fast [15–17]. Consequently, the transition from the eMD era to the subsequent late radiation dominated era (IRD) is quite sudden. This means that the gravitational waves induced by inflationary adiabatic perturbations around the time of the flaton decay will become significantly enhanced due to the fast oscillations of the scalar metric perturbations well within the Hubble horizon [20]. It is the spectrum of these GWs that we are going to investigate in what follows.

Scalar induced gravitational waves– Having described above the flipped $SU(5)$ cosmological phenomenology, let us now review the basic formalism of the gravitational waves induced by second order gravitational interactions [47–50] [See [51] for a review] and which in our setup are abundantly produced due to the existence of an eMD era driven by the flaton field. We choose to work within the so-called Newtonian gauge ³, in which the perturbed metric is written as

$$ds^2 = a^2(\eta) \left\{ -(1 + 2\Psi)d\eta^2 + \left[(1 - 2\Psi)\delta_{ij} + \frac{h_{ij}}{2} \right] dx^i dx^j \right\}, \quad \Omega_{\text{GW}}(\eta_0, k) = \Omega_r^{(0)} \frac{g_{*\rho,*}}{g_{*\rho,0}} \left(\frac{g_{*S,0}}{g_{*S,*}} \right)^{4/3} \Omega_{\text{GW}}(\eta_*, k), \quad (7)$$

where Ψ is the first order Bardeen gravitational potential and h_{ij} denotes the second order tensor perturbation. Then, after a Fourier transformation of the tensor perturbation and minimisation of the tensor part of the cubic order gravitational action, one straightforwardly gets the equation of motion for $h_{\mathbf{k}}$ reading as

$$h_{\mathbf{k}}^{s,\prime\prime} + 2\mathcal{H}h_{\mathbf{k}}^{s,\prime} + k^2 h_{\mathbf{k}}^s = 4S_{\mathbf{k}}^s, \quad (3)$$

where \prime denotes differentiation with respect to the conformal time and $s = (+), (\times)$, stands for the two tensor

mode polarisation states in general relativity (GR), with \mathcal{H} is the conformal Hubble parameter, while the polarization tensors $e_{ij}^s(\mathbf{k})$ are the standard ones [56]. The source function $S_{\mathbf{k}}^s$ is given by ⁴

$$S_{\mathbf{k}}^s = \int \frac{d^3\mathbf{q}}{(2\pi)^{3/2}} e_{ij}^s(\mathbf{k}) q_i q_j \left[2\Psi_{\mathbf{q}} \Psi_{\mathbf{k}-\mathbf{q}} + \frac{4}{3(1+w_{\text{tot}})} (\mathcal{H}^{-1}\Psi'_{\mathbf{q}} + \Psi_{\mathbf{q}})(\mathcal{H}^{-1}\Psi'_{\mathbf{k}-\mathbf{q}} + \Psi_{\mathbf{k}-\mathbf{q}}) \right] \quad (4)$$

and as one can see is quadratically dependent on the first order scalar perturbation of the metric. After a lengthy but straightforward calculation, one obtains that the tensor power spectrum $\mathcal{P}_h^{(s)}(\eta, k)$ reads as [56, 59–61]

$$\mathcal{P}_h^{(s)}(\eta, k) = 4 \int_0^\infty dv \int_{|1-v|}^{1+v} du \left[\frac{4v^2 - (1+v^2 - u^2)^2}{4uv} \right]^2 \cdot I^2(u, v, x) \mathcal{P}_\Psi(kv) \mathcal{P}_\Psi(ku), \quad (5)$$

where the two auxiliary variables u and v are defined as $u \equiv |\mathbf{k} - \mathbf{q}|/k$ and $v \equiv q/k$, and the kernel function $I(u, v, x)$ is a complicated function containing information for the transition between the eMD era driven by the flaton field and the IRD era [20, 21, 25, 61]. Using Eq. (5), one can derive the GW spectral abundance, which is defined as follows as [62]

$$\Omega_{\text{GW}}(\eta, k) \equiv \frac{1}{\bar{\rho}_{\text{tot}}} \frac{d\rho_{\text{GW}}(\eta, k)}{d \ln k} = \frac{1}{24} \left(\frac{k}{\mathcal{H}(\eta)} \right)^2 \overline{\mathcal{P}_h^{(s)}(\eta, k)}, \quad (6)$$

where the bar stands for the oscillation average, i.e. the GW envelope. Finally, considering that the radiation energy density reads as $\rho_r = \frac{\pi^2}{30} g_{*\rho} T_r^4$ and that the temperature of the primordial plasma T_r scales as $T_r \propto g_{*S}^{-1/3} a^{-1}$, one finds that the GW spectral abundance at our present epoch reads as

$$\Omega_{\text{GW}}(\eta_0, k) = \Omega_r^{(0)} \frac{g_{*\rho,*}}{g_{*\rho,0}} \left(\frac{g_{*S,0}}{g_{*S,*}} \right)^{4/3} \Omega_{\text{GW}}(\eta_*, k), \quad (7)$$

where $g_{*\rho}$ and g_{*S} denote the energy and entropy relativistic degrees of freedom. Note that the reference conformal time η_* in the case of a sudden transition from the eMD to the IRD era should be of $\mathcal{O}(1)\eta_r$ [20].

The flipped $SU(5)$ induced gravitational wave signal– We are going to focus now on the induced gravitational waves generated within the flipped $SU(5)$ paradigm. Interestingly enough, these induced GWs are significantly enhanced in our case since during the sudden transition from the flaton-driven eMD era to the IRD era,

³ We work within the Newtonian gauge as it is standardly used in the literature related to SIGWs. The effect of the gauge dependence of the SIGWs is discussed in [52–55].

⁴ We mention here that in this work we neglect possible effects of non-Gaussianities [57] and one-loop corrections [58] to the induced GW background.

the time derivative of the Bardeen potential goes very quickly from $\Psi' = 0$ (since in a MD era $\Psi = \text{constant}$) to $\Psi' \neq 0$ [See [20, 51] for more details.]. This entails a resonantly enhanced production of GWs sourced mainly by the $\mathcal{H}^{-2}\Psi'^2$ term in Eq. (4). Furthermore, since the sub-horizon energy density perturbations during an MD era scale linearly with the scale factor, i.e. $\delta \propto a$, one should ensure working within the perturbative regime. Thus, we set a non-linear scale k_{NL} by requiring that $\delta_{k_{\text{NL}}}(\eta_r) = 1$. For inflationary power-low primordial curvature spectra, one finds that [20]

$$k \leq k_{\text{NL}} \simeq 470k_r, \quad (8)$$

where k_r is the comoving scale crossing the cosmological horizon at the onset of the IRD era. At the end, the relevant amplified GW signal is calculated as follows

$$\Omega_{\text{GW}}^{\text{res}}(\eta_0, k) = c_g \Omega_r^{(0)} \Omega_{\text{GW}}(\eta_{\text{IRD}}, k), \quad (9)$$

where $c_g = \frac{g_{**\rho,*}}{g_{**0}} \left(\frac{g_{**S,0}}{g_{**S,*}} \right)^{4/3} \sim 0.4$ and $\eta_{\text{IRD}} \sim O(\eta_r)$ stands for a time during the IRD era by which the curvature perturbations decouple from the tensor perturbations, thus one can assume freely propagating GWs while $\Omega_{\text{GW}}(\eta_{\text{IRD}}, k)$ is derived from Eq. (6) where $I(u, v, x) = I_{\text{IRD}}(u, v, x)$ [20].

Regarding the frequency of the GW signal, it can be computed as follows

$$\begin{aligned} f_{\text{GW}} &= \frac{k}{2\pi a_0} = \frac{k}{2\pi a_{\text{d}\Phi}} \frac{a_{\text{d}\Phi}}{a_{\text{eq}}} \frac{a_{\text{eq}}}{a_0} \\ &= \frac{k}{2\pi a_{\text{d}\Phi}} \left(\frac{\rho_{\text{eq}}}{\rho_{\text{d}\Phi}} \right)^{1/4} \left(\frac{\rho_0}{\rho_{\text{eq}}} \right)^{1/3}, \end{aligned} \quad (10)$$

where $\rho_{\text{eq}} = 1.096 \times 10^{-36} \text{GeV}^4$ and $\rho_0 = 3.6 \times 10^{-47} \text{GeV}^4$ are the background energy densities at the matter-radiation equality and today respectively. Lastly, $\rho_{\text{d}\Phi}$ and $a_{\text{d}\Phi}$ respectively stand for the energy density and the scale factor at the end of the eMD, namely at the time of the flaton decay.

In order to compute now $\rho_{\text{d}\Phi}$ and $a_{\text{d}\Phi}$, let us recap the basic physical quantities describing the dynamics of the flaton-driven eMD era. In particular, the Hubble parameter during the flaton-dominated era is given by [15]

$$H = \left(\frac{\rho_{\Phi}}{3M_P^2} \right)^{1/2} = \left(\frac{\zeta(3)m_{\Phi}T_{\Phi}^3}{3\pi^2 M_P^2} \right)^{1/2}. \quad (11)$$

With regard now to the decay rate Γ_{Φ} of the flaton, the latter was calculated in [17] via effective D -term diagrams, leading to

$$\Gamma_{\Phi} \simeq \frac{9\lambda_{1,2,3,7}^4}{2048\pi^5} \left(\frac{m_{\Phi} m_{F,\bar{f},\ell^c,\tilde{\phi}_a}^2}{M_{\text{GUT}}^2} \right) \quad (12)$$

where m_{Φ} is the mass of the flaton, M_{GUT} is the GUT energy scale and $m_{F,\bar{f},\ell^c,\tilde{\phi}_a}$ are the masses of the various

fields and $\lambda_{1,2,3,7}$ are appropriate Yukawa couplings of the order of $O(1/2)$ [See [15] for more details.]. Assuming to a very good approximation $m_{F,\bar{f},\ell^c,\tilde{\phi}_a} \simeq m_{\Phi}$ [15], one finds that the flaton decays approximately when $H \sim \Gamma_{\Phi}$, or equivalently when the flaton temperature is

$$T_{\text{d}\Phi} \simeq \frac{3\lambda_{1,2,3,7}^{8/3}}{128} \left(\frac{9}{2\zeta(3)\pi^8} \right)^{1/3} \left(\frac{m_{\Phi} m_{F,\bar{f},\ell^c,\tilde{\phi}_a}^4 M_P^2}{M_{\text{GUT}}^4} \right)^{1/3}. \quad (13)$$

Having reviewed above the dynamics of the flaton-dominated era, let us derive here the characteristic scales of the problem at hand. In particular, the mode k_{Φ} , which crosses the cosmological horizon at the onset of the flaton domination era at $T \sim m_{\Phi}$, will be written as $k_{\Phi} = a_{\Phi} H_{\Phi} = \left(\frac{\zeta(3)m_{\Phi}^4}{3\pi^2 M_P^2} \right)^{1/2}$, where we have normalised $a_{\Phi} = 1$ whereas the mode $k_{\text{d}\Phi}$ crossing the horizon at the time of the flaton decay when $H \sim \Gamma_{\Phi}$ will read as $k_{\text{d}\Phi} = a_{\text{d}\Phi} H_{\text{d}\Phi}$, where $H_{\text{d}\Phi} = \Gamma_{\Phi}$ and $a_{\text{d}\Phi} = (H_{\Phi}^2/H_{\text{d}\Phi}^2)^{1/3}$.

Finally, the peak frequency of our problem at hand, where one expects a resonantly amplified GW signal, is associated with the non-linear comoving cut-off scale, $k_{\text{NL}} = 470k_r$, where in our case $k_r = k_{\text{d}\Phi}$. At the end, plugging $k_{\text{d}\Phi}$ in Eq. (10), we obtain that the peak frequency $f_{\text{GW,peak}}$ reads as

$$\begin{aligned} f_{\text{GW,peak}} &= \frac{470k_r}{2\pi a_{\text{d}\Phi}} \left(\frac{\rho_{\text{eq}}}{\rho_r} \right)^{1/4} \left(\frac{\rho_0}{\rho_{\text{eq}}} \right)^{1/3} \\ &= 1.7 \times 10^{-9} \left(\frac{\lambda_{1,3,5,7}}{0.5} \right)^2 \left(\frac{m_{\Phi}}{10^4 \text{GeV}} \right)^{3/2} \left(\frac{10^{16} \text{GeV}}{M_{\text{GUT}}} \right). \end{aligned} \quad (14)$$

Remarkably, within the framework of flipped $\text{SU}(5)$ one can re-express the mass parameters of Eq. (14) in terms of solely the Planck Mass and ultimately directly in terms of only the string slope parameter α' . In particular, as it was shown in [15, 16], there is a relation between the flaton vev, $\langle \Phi \rangle$, and the flaton soft supersymmetry-breaking mass term m_{Φ}

$$|m_{\Phi}| \sim \frac{\langle \Phi \rangle^6}{M_{\text{Pl}}^5}. \quad (15)$$

Since the flaton vev is of order of the GUT scale $\langle \Phi \rangle \approx M_{\text{GUT}}$ [15], then, by substituting $M_{\text{GUT}} = 10^{16} \text{GeV}$ and $M_{\text{Pl}} = 4M_s$ where M_s is the heterotic string scale being defined as $M_s \equiv 1/\sqrt{2\alpha'}$ [7] one obtains that

$$f_{\text{GW,peak}} = 1.7 \times 10^{-9} \left(\frac{\lambda_{1,2,3,7}}{0.5} \right)^2 \left(\frac{\alpha'}{8/M_{\text{Pl}}^2} \right)^4 \text{Hz}. \quad (16)$$

We note here that in our analysis we ignored the GWs which are resonantly enhanced during the first reheating stage when the inflaton field oscillates at the bottom of its potential and during which the Universe effectively




FIG. 1. *The primordial GW background non-linearly induced by inflationary perturbations (solid green curve) arising from flipped SU(5) superstring theory for $m_{\Phi} = 10^4 \text{ GeV}$ and $M_{\text{GUT}} = 10^{16} \text{ GeV}$. On the top of our theoretical prediction for the induced stochastic GW background we show the 15-year NANOGrav GW data, as well as the sensitivities of SKA [63], LISA [64, 65], BBO [66] and ET [67] GW experiments. In the horizontal black dashed line, we show also the upper bound on $\Omega_{\text{GW}, \text{eff}} h^2 \leq 6.9 \times 10^{-6}$ coming from the upper bound constraint on ΔN_{eff} from CMB and BBN observations [68].*

behaves as pressureless matter [12, 15]. During this first eMD era before the flaton field dominates the Universe content, GWs can be abundantly produced at higher frequencies compared to the flaton-driven eMD era induced GWs. However, since after the end of the first reheating stage there follows a long second eMD era driven by the flaton, one expects that these GWs are sufficiently diluted since $\Omega_{\text{GW}} \propto \rho_{\text{GW}}/\rho_{\text{flaton}} \propto a^{-4}/a^{-3} \propto 1/a$. The relevant dilution factor of these primordial GWs generated during the first reheating stage will read as $\Delta \equiv \frac{\alpha_{\text{d}\Phi}}{\alpha_{\Phi}} \simeq 10^{10}$, for $m_{\Phi} = 10^4 \text{ GeV}$ and $M_{\text{GUT}} = 10^{16} \text{ GeV}$. Nevertheless, if one accounts for the non-linear growth of the perturbations entering the inflaton-driven eMD era, they may enhance the GW amplitude rendering it potentially detectable by the LISA, BBO and ET GW probes.

Conclusions – The flipped SU(5) superstring paradigm accommodates a very rich particle and cosmological phenomenology consistent with observations. In particular, it can naturally give rise to stable Starobinsky-type inflation preferred by Planck. What’s more, it can explain successfully the lepton and quark mass, while at the same time it can provide a realistic mechanism for the generation of the baryon asymmetry in the Universe and the production of neutralino dark matter. Among its most interesting aspects is its strong reheating scenario which proceeds via the inco-

herent component of the flaton field.

In this Letter, we focused on a particular phenomenology associated with the flaton field which naturally arises in the flipped SU(5) superstring paradigm. Specifically, as it was shown in [12, 15], one meets an eMD era driven by the flaton which transitions quite suddenly to the IRD era. This sudden transition leads to an abundant production of a GW signal non-linearly induced by inflationary curvature perturbations, constituting a pure observational GW signature of the flipped SU(5) superstring theory. Remarkably, we found that the peak frequency of this GW signal $f_{\text{GW, peak}}$ depends only on the string slope α' and reads as

$$f_{\text{GW, peak}} \propto 10^{-9} \left(\frac{\alpha'}{\alpha'_*} \right)^4 \text{ Hz}, \quad (17)$$

where α'_* is the fiducial string slope being related to the reduced Planck scale M_{Pl} as $\alpha'_* = 8/M_{\text{Pl}}^2$. Notably, as one may see from Eq. (17), the peak frequency of these GWs lies in the nHz frequency range and is very close to the NANOGrav/PTA data; hence rendering this primordial GW signal potentially detectable by SKA, NANOGrav and PTA probes at their very low frequency region of their detection bands.

If one now takes into account via GR numerical simulations the highly non-linear energy density perturbations present at smaller scales, i.e. at higher frequencies [69], the expected GW signal will be significantly enhanced at these scales compared to the one given here and displaced within the $10^{-9} - 10^{-8} \text{ nHz}$ frequency range, hence being potentially a good candidate to explain the NANOGrav/PTA GW signal [See here for recent works [70–86] on different cosmological models explaining the NANOGrav/PTA signal.]. Lastly, due to the linear growth of the sub-horizon energy density perturbations during the second reheating stage driven by the flaton, one expects the formation of flaton structures similar to the inflaton ones [87, 88] as well as primordial black holes (PBHs) [89, 90] with very interesting phenomenology. These aspects will be studied in future works.

Acknowledgements – The work of DVN was supported in part by the DOE grant DE-FG02-13ER42020 at Texas A&M University and in part by the Alexander S. Onassis Public Benefit Foundation. SB, ENS, TP and CT acknowledge the contribution of the LISA CosWG and the COST Actions CA18108 “Quantum Gravity Phenomenology in the multi-messenger approach” and CA21136 “Addressing observational tensions in cosmology with systematics and fundamental physics (CosmoVerse)”. TP and CT acknowledge as well financial support from the Foundation for Education and European Culture in Greece and A.G. Leventis Foundation respectively.

-
- [1] I. Antoniadis, J. R. Ellis, J. S. Hagelin, and D. V. Nanopoulos, *Phys. Lett. B* **231**, 65 (1989).
- [2] J. Rizos and K. Tamvakis, *Phys. Lett. B* **251**, 369 (1990).
- [3] I. Antoniadis, J. Rizos, and K. Tamvakis, *Phys. Lett. B* **278**, 257 (1992).
- [4] M. Cvetič, L. L. Everett, and J. Wang, *Phys. Rev. D* **59**, 107901 (1999), [arXiv:hep-ph/9808321](#).
- [5] G. Cleaver, M. Cvetič, J. R. Espinosa, L. L. Everett, and P. Langacker, *Phys. Rev. D* **57**, 2701 (1998), [arXiv:hep-ph/9705391](#).
- [6] M. Cicoli, J. P. Conlon, A. Maharana, S. Parameswaran, F. Quevedo, and I. Zavala, (2023), [arXiv:2303.04819 \[hep-th\]](#).
- [7] I. Antoniadis, D. V. Nanopoulos, and J. Rizos, *JCAP* **03**, 017 (2021), [arXiv:2011.09396 \[hep-th\]](#).
- [8] I. Antoniadis, D. V. Nanopoulos, and J. Rizos, *Eur. Phys. J. C* **82**, 377 (2022), [arXiv:2112.01211 \[hep-th\]](#).
- [9] J. Ellis, D. V. Nanopoulos, and K. A. Olive, *Phys. Rev. Lett.* **111**, 111301 (2013), [Erratum: *Phys.Rev.Lett.* 111, 129902 (2013)], [arXiv:1305.1247 \[hep-th\]](#).
- [10] J. Ellis, D. V. Nanopoulos, and K. A. Olive, *JCAP* **10**, 009 (2013), [arXiv:1307.3537 \[hep-th\]](#).
- [11] J. Ellis, D. V. Nanopoulos, and K. A. Olive, *Phys. Rev. D* **89**, 043502 (2014), [arXiv:1310.4770 \[hep-ph\]](#).
- [12] J. Ellis, M. A. G. Garcia, N. Nagata, D. V. Nanopoulos, and K. A. Olive, *JCAP* **07**, 006 (2017), [arXiv:1704.07331 \[hep-ph\]](#).
- [13] H. Goldberg, *Phys. Rev. Lett.* **50**, 1419 (1983), [Erratum: *Phys.Rev.Lett.* 103, 099905 (2009)].
- [14] J. R. Ellis, J. S. Hagelin, D. V. Nanopoulos, K. A. Olive, and M. Srednicki, *Nucl. Phys. B* **238**, 453 (1984).
- [15] J. Ellis, M. A. G. Garcia, N. Nagata, D. V. Nanopoulos, and K. A. Olive, *JCAP* **04**, 009 (2019), [arXiv:1812.08184 \[hep-ph\]](#).
- [16] J. Ellis, M. A. G. Garcia, N. Nagata, D. V. Nanopoulos, and K. A. Olive, *JCAP* **01**, 035 (2020), [arXiv:1910.11755 \[hep-ph\]](#).
- [17] B. Campbell, J. Ellis, J. Hagelin, D. Nanopoulos, and K. Olive, *Physics Letters B* **197**, 355 (1987).
- [18] A. G. Polnarev and M. Y. Khlopov, *Soviet Astronomy* **25**, 406 (1981).
- [19] A. G. Polnarev and I. M. Khlopov, *Astronomicheskii Zhurnal* **59**, 639 (1982).
- [20] K. Inomata, K. Kohri, T. Nakama, and T. Terada, *Phys. Rev. D* **100**, 043532 (2019), [arXiv:1904.12879 \[astro-ph.CO\]](#).
- [21] K. Inomata, K. Kohri, T. Nakama, and T. Terada, *JCAP* **10**, 071 (2019), [arXiv:1904.12878 \[astro-ph.CO\]](#).
- [22] K. Inomata, M. Kawasaki, K. Mukaida, T. Terada, and T. T. Yanagida, *Phys. Rev. D* **101**, 123533 (2020), [arXiv:2003.10455 \[astro-ph.CO\]](#).
- [23] T. Papanikolaou, V. Vennin, and D. Langlois, *JCAP* **03**, 053 (2021), [arXiv:2010.11573 \[astro-ph.CO\]](#).
- [24] G. Domènech, C. Lin, and M. Sasaki, *JCAP* **04**, 062 (2021), [arXiv:2012.08151 \[gr-qc\]](#).
- [25] T. Papanikolaou, *JCAP* **10**, 089 (2022), [arXiv:2207.11041 \[astro-ph.CO\]](#).
- [26] N. Bhaumik, A. Ghoshal, R. K. Jain, and M. Lewicki, *JHEP* **05**, 169 (2023), [arXiv:2212.00775 \[astro-ph.CO\]](#).
- [27] T. Papanikolaou, in *CORFU2022* (2023) [arXiv:2303.00600 \[astro-ph.CO\]](#).
- [28] I. Antoniadis, C. Bachas, and C. Kounnas, *Nuclear Physics B* **289**, 87 (1987).
- [29] I. Antoniadis and C. Bachas, *Nuclear Physics B* **298**, 586 (1988).
- [30] S. Kalara, J. L. Lopez, and D. V. Nanopoulos, *Nucl. Phys. B* **353**, 650 (1991).
- [31] S. Barr, *Physics Letters B* **112**, 219 (1982).
- [32] J.-P. Derendinger, J. E. Kim, and D. Nanopoulos, *Physics Letters B* **139**, 170 (1984).
- [33] I. Antoniadis, J. Ellis, J. Hagelin, and D. Nanopoulos, *Physics Letters B* **194**, 231 (1987).
- [34] J. R. Ellis, J. L. Lopez, and D. V. Nanopoulos, *Phys. Lett. B* **245**, 375 (1990).
- [35] M. Dine, N. Seiberg, and E. Witten, *Nucl. Phys. B* **289**, 589 (1987).
- [36] E. Cremmer, S. Ferrara, C. Kounnas, and D. Nanopoulos, *Physics Letters B* **133**, 61 (1983).
- [37] J. R. Ellis, A. B. Lahanas, D. V. Nanopoulos, and K. Tamvakis, *Phys. Lett. B* **134**, 429 (1984).
- [38] J. Ellis, C. Kounnas, and D. Nanopoulos, *Nuclear Physics B* **241**, 406 (1984).
- [39] J. Ellis, C. Kounnas, and D. Nanopoulos, *Nuclear Physics B* **247**, 373 (1984).
- [40] A. Lahanas and D. Nanopoulos, *Physics Reports* **145**, 1 (1987).
- [41] I. Antoniadis, J. Ellis, E. Floratos, D. Nanopoulos, and T. Tomaras, *Physics Letters B* **191**, 96 (1987).
- [42] S. Ferrara, L. Girardello, C. Kounnas, and M. Porrati, *Physics Letters B* **192**, 368 (1987).
- [43] S. Ferrara, L. Girardello, C. Kounnas, and M. Porrati, *Physics Letters B* **194**, 358 (1987).
- [44] N. Aghanim et al. (Planck), *Astron. Astrophys.* **641**, A6 (2020), [Erratum: *Astron.Astrophys.* 652, C4 (2021)], [arXiv:1807.06209 \[astro-ph.CO\]](#).
- [45] J. Ellis, M. A. Garcia, N. Nagata, D. V. Nanopoulos, and K. A. Olive, *Physics Letters B* **797**, 134864 (2019).
- [46] D. V. Nanopoulos and K. Tamvakis, *Phys. Lett. B* **110**, 449 (1982).
- [47] S. Matarrese, O. Pantano, and D. Saez, *Phys. Rev. D* **47**, 1311 (1993).
- [48] S. Matarrese, O. Pantano, and D. Saez, *Phys. Rev. Lett.* **72**, 320 (1994), [arXiv:astro-ph/9310036](#).
- [49] S. Matarrese, S. Mollerach, and M. Bruni, *Phys. Rev. D* **58**, 043504 (1998), [arXiv:astro-ph/9707278](#).
- [50] S. Mollerach, D. Harari, and S. Matarrese, *Phys. Rev. D* **69**, 063002 (2004), [arXiv:astro-ph/0310711](#).
- [51] G. Domènech, *Universe* **7**, 398 (2021), [arXiv:2109.01398 \[gr-qc\]](#).
- [52] J.-C. Hwang, D. Jeong, and H. Noh, *Astrophys. J.* **842**, 46 (2017), [arXiv:1704.03500 \[astro-ph.CO\]](#).
- [53] K. Tomikawa and T. Kobayashi, *Phys. Rev. D* **101**, 083529 (2020), [arXiv:1910.01880 \[gr-qc\]](#).
- [54] V. De Luca, G. Franciolini, A. Kehagias, and A. Riotto, *JCAP* **03**, 014 (2020), [arXiv:1911.09689 \[gr-qc\]](#).
- [55] K. Inomata and T. Terada, *Phys. Rev. D* **101**, 023523 (2020), [arXiv:1912.00785 \[gr-qc\]](#).
- [56] J. R. Espinosa, D. Racco, and A. Riotto, *JCAP* **1809**, 012 (2018), [arXiv:1804.07732 \[hep-ph\]](#).
- [57] R.-g. Cai, S. Pi, and M. Sasaki, *Phys. Rev. Lett.* **122**, 201101 (2019), [arXiv:1810.11000 \[astro-ph.CO\]](#).
- [58] C. Chen, A. Ota, H.-Y. Zhu, and Y. Zhu, *Phys. Rev. D* **107**, 083518 (2023), [arXiv:2210.17176 \[astro-ph.CO\]](#).
- [59] K. N. Ananda, C. Clarkson, and D. Wands, *Phys. Rev. D* **75**, 123518 (2007), [arXiv:gr-qc/0612013 \[gr-qc\]](#).

- [60] D. Baumann, P. J. Steinhardt, K. Takahashi, and K. Ichiki, *Phys. Rev.* **D76**, 084019 (2007), [arXiv:hep-th/0703290 \[hep-th\]](#).
- [61] K. Kohri and T. Terada, *Phys. Rev.* **D97**, 123532 (2018), [arXiv:1804.08577 \[gr-qc\]](#).
- [62] M. Maggiore, *Phys. Rept.* **331**, 283 (2000), [arXiv:gr-qc/9909001 \[gr-qc\]](#).
- [63] G. Janssen et al., *PoS AASKA14*, 037 (2015), [arXiv:1501.00127 \[astro-ph.IM\]](#).
- [64] P. Auclair et al. (LISA Cosmology Working Group), (2022), [arXiv:2204.05434 \[astro-ph.CO\]](#).
- [65] N. Karnesis et al., (2022), [arXiv:2209.04358 \[gr-qc\]](#).
- [66] G. M. Harry, P. Fritschel, D. A. Shaddock, W. Folkner, and E. S. Phinney, *Class. Quant. Grav.* **23**, 4887 (2006), [Erratum: *Class. Quant. Grav.* **23**, 7361 (2006)].
- [67] M. Maggiore et al., *JCAP* **03**, 050 (2020), [arXiv:1912.02622 \[astro-ph.CO\]](#).
- [68] T. L. Smith, E. Pierpaoli, and M. Kamionkowski, *Phys. Rev. Lett.* **97**, 021301 (2006), [arXiv:astro-ph/0603144](#).
- [69] K. Jedamzik, M. Lemoine, and J. Martin, *JCAP* **04**, 021 (2010), [arXiv:1002.3278 \[astro-ph.CO\]](#).
- [70] T. Bringmann, P. F. Depta, T. Konstandin, K. Schmidt-Hoberg, and C. Tassilo, (2023), [arXiv:2306.09411 \[astro-ph.CO\]](#).
- [71] K. Fujikura, S. Girmohanta, Y. Nakai, and M. Suzuki, (2023), [arXiv:2306.17086 \[hep-ph\]](#).
- [72] X. Niu and M. H. Rahat, (2023), [arXiv:2307.01192 \[hep-ph\]](#).
- [73] S. A. Hosseini Mansoori, F. Felegray, A. Talebian, and M. Sami, *JCAP* **08**, 067 (2023), [arXiv:2307.06757 \[astro-ph.CO\]](#).
- [74] S. Choudhury, (2023), [arXiv:2307.03249 \[astro-ph.CO\]](#).
- [75] Y.-M. Wu, Z.-C. Chen, and Q.-G. Huang, *Phys. Rev. D* **107**, 042003 (2023), [arXiv:2302.00229 \[gr-qc\]](#).
- [76] B. Das, N. Jaman, and M. Sami, (2023), [arXiv:2307.12913 \[gr-qc\]](#).
- [77] S. Basilakos, D. V. Nanopoulos, T. Papanikolaou, E. N. Saridakis, and C. Tzerefos, (2023), [arXiv:2307.08601 \[hep-th\]](#).
- [78] Z. Yi, Z.-Q. You, Y. Wu, Z.-C. Chen, and L. Liu, (2023), [arXiv:2308.14688 \[astro-ph.CO\]](#).
- [79] S. Datta, (2023), [arXiv:2309.14238 \[hep-ph\]](#).
- [80] S. Balaji, G. Domènech, and G. Franciolini, (2023), [arXiv:2307.08552 \[gr-qc\]](#).
- [81] Y. Gouttenoire, S. Trifinopoulos, G. Valogiannis, and M. Vanvlasselaer, (2023), [arXiv:2307.01457 \[astro-ph.CO\]](#).
- [82] N. Bhaumik, R. K. Jain, and M. Lewicki, (2023), [arXiv:2308.07912 \[astro-ph.CO\]](#).
- [83] G. Franciolini, D. Racco, and F. Rompineve, (2023), [arXiv:2306.17136 \[astro-ph.CO\]](#).
- [84] G. Franciolini, A. Iovino, Junior., V. Vaskonen, and H. Veermae, (2023), [arXiv:2306.17149 \[astro-ph.CO\]](#).
- [85] K. Inomata, M. Kawasaki, K. Mukaida, and T. T. Yanagida, (2023), [arXiv:2309.11398 \[astro-ph.CO\]](#).
- [86] K. Inomata, K. Kohri, and T. Terada, (2023), [arXiv:2306.17834 \[astro-ph.CO\]](#).
- [87] K. Jedamzik, M. Lemoine, and J. Martin, *JCAP* **1009**, 034 (2010), [arXiv:1002.3039 \[astro-ph.CO\]](#).
- [88] J. C. Hidalgo, L. E. Padilla, and G. German, *Phys. Rev. D* **107**, 063519 (2023), [arXiv:2208.09462 \[astro-ph.CO\]](#).
- [89] J. Martin, T. Papanikolaou, and V. Vennin, *JCAP* **01**, 024 (2020), [arXiv:1907.04236 \[astro-ph.CO\]](#).
- [90] J. Martin, T. Papanikolaou, L. Pinol, and V. Vennin, *JCAP* **05**, 003 (2020), [arXiv:2002.01820 \[astro-ph.CO\]](#).