# Unification of Gravity and Internal Interactions

Spyros Konitopoulos<sup>1</sup>, Danai Roumelioti<sup>2</sup>, George Zoupanos<sup>2,3,4,5</sup>

2023

<sup>1</sup>Institute of Nuclear and Particle Physics, NCSR Demokritos, Athens, Greece <sup>2</sup>Physics Department, National Technical University, Athens, Greece

<sup>3</sup>Theory Department, CERN

<sup>4</sup>Max-Planck Institut für Physik, München, Germany

5 Institut für Theoretische Physik der Universität Heidelberg, Germany

E-mails: [spykoni@gmail.com,](mailto:spykoni@gmail.com) [danai\\_roumelioti@mail.ntua.gr,](mailto:danai_roumelioti@mail.ntua.gr) [George.Zoupanos@cern.ch](mailto:George.Zoupanos@cern.ch)

#### Abstract

In the gauge theoretic approach of gravity, General Relativity is described by gauging the symmetry of the tangent manifold in four dimensions. Usually the dimension of the tangent space is considered to be equal to the dimension of the curved manifold. However, the tangent group of a manifold of dimension  $d$ is not necessarily  $SO_d$ . It has been suggested earlier that by gauging an enlarged symmetry of the tangent space in four dimensions one could unify gravity with internal interactions. Here we consider such a unified model by gauging the  $SO_{(1,17)}$  as the extended Lorentz group overcoming in this way some difficulties of the previous attempts of similar unification and eventually we obtain the  $SO_{10}$  GUT, supplemented by an  $SU_2 \times SU_2$  global symmetry.

## 1 Introduction

An ultimate aim of many theoretical physicists is the existence of a unification picture in which all known fundamental interactions are involved. A huge amount of serious research activity has been carried out, including works that elaborate the very interesting notion of extra dimensions. The earliest unification attempts of Kaluza and Klein [\[1,](#page-10-0) [2\]](#page-10-1) included gravity and electromagnetism, which were the established interactions at that time. The proposal was to reduce a pure gravity theory from five dimensions to four, which led to a  $U_1$ gauge theory, identified with electromagnetism, coupled to gravity. A revival of interest in the Kaluza-Klein scheme started after realizing [\[3](#page-10-2)[–5\]](#page-10-3) that non-abelian gauge groups appear naturally when one further extends the spacetime dimensions. With the assumption that the total spacetime manifold can be written as a direct product  $M_D = M_4 \times B$ , where B is a compact Riemannian space with a non-abelian isometry group S, dimensional reduction of the theory leads to gravity coupled to a Yang-Mills theory with a gauge group containing S and scalars in four dimensions. The main advantage of this picture is the geometrical unification of gravity with the other interactions and also the explanation of gauge symmetries. There exist serious problems though in the Kaluza-Klein framework, e.g. there is no classical ground state corresponding to the direct product structure of  $M_D$ . However, the most serious obstacle in obtaining a realistic model of the low-energy interactions seems to be that after adding fermions to the original action it is impossible to obtain chiral fermions in four dimensions [\[6\]](#page-10-4). Eventually, if one adds suitable matter fields to the original gravity action in particular Yang-Mills then most of the serious problems are resolved. Therefore one is led to introduce Yang-Mills fields in higher dimensions. In case the Yang-Mills are part of a Grand Unified Theory (GUT) together with a Dirac one [\[7](#page-10-5), [8\]](#page-10-6), the restriction to obtain chiral fermions in four dimensions is limited to the requirement that the total dimension of spacetime should be  $4k + 2$  (see e.g. [\[9](#page-10-7)]). During the last decades the Superstring theories (see e.g. [\[10](#page-10-8)[–12\]](#page-10-9)) dominated the research on extra dimensions consisting a solid framework. In particular the heterotic string theory [\[13](#page-10-10)] (defined in ten dimensions) was the most promising, since potentially it admits experimental compatibility, due to the fact that the Standard Model (SM) gauge group can be accommodated into those of GUTs that emerge after the dimensional reduction of the initial  $E_8 \times E_8$ . It is worth noting that even before the formulation of superstring theories, an alternative framework was developed that focused on the dimensional reduction of higher-dimensional gauge theories. This provided another venue for exploring the unification of fundamental interactions [\[9](#page-10-7), [14](#page-10-11)[–22](#page-11-0)]. The endeavor to unify fundamental interactions, which shared common objectives with the superstring theories, was first investigated by Forgacs-Manton (F-M) and Scherk-Schwartz (S-S). F-M explored the concept of Coset Space Dimensional Reduction (CSDR) [\[14\]](#page-10-11), which can lead naturally to chiral fermions while S-S focused on the group manifold reduction [\[18\]](#page-11-1), which does not admit chiral fermions. Recent attempts towards realistic models that can be confronted with experiment can be found in [\[15](#page-10-12), [23,](#page-11-2) [24\]](#page-11-3).

On the gravity side diffeomorphism-invariant gravity theory is obviously invariant with respect to transformations whose parameters are functions of spacetime, just as in the local gauge theories. Then, naturally, it has been long believed that general relativity (GR) can be formulated as a gauge theory [\[25](#page-11-4)[–27](#page-11-5)] with the spin connection as the corresponding gauge field which would enter in the action through the corresponding field strength. This idea was used heavily in supergravity (see e.g. [\[28](#page-11-6)]) while recently it was employed in non-commutative gravity too [\[29](#page-11-7)[–31](#page-11-8)]. Along the same lines rather recently was suggested a new idea for unification of all known interactions in four dimensions. Usually the dimension of the tangent space is taken to be equal to the dimension of the curved manifold. However, the tangent group of a manifold of dimension  $d$  is not necessarily  $SO_d$  [\[32\]](#page-11-9). It is possible to embed the coordinate tangent space in a higher-dimensional space, and therefore promote the gauge symmetry to a higher isometry group. In refs [\[33](#page-11-10)[–39\]](#page-11-11), the authors have considered higher-dimensional tangent spaces in 4−dimensional spacetime and managed in this way to achieve unification of internal interactions with gravity. The geometric unification of gravity and gauge internal interactions in refs [\[38,](#page-11-12) [39](#page-11-11)] is realized by writing the action of the full theory in terms only of the curvature invariants of the tangent group, which contain the Yang-Mills actions corresponding to the gauge groups describing in this way together the GR and the internal GUT in a unified manner. The best model found so far that unifies gravity and a chiral GUT is based on  $SO_{(1,13)}$  in a 14-dimensional tangent space leading to unification of gravity with  $SO_{10}$ . However as a drawback was considered the fact that fermions appear in double representations of the spinor 16 of  $SO_{10}$ , which only means that fermions appear in even families though [\[40](#page-11-13)]. Trying to resolve this

problem by imposing Majorana condition in addition to Weyl in ref [\[35](#page-11-14), [36](#page-11-15)] was proposed instead as a unifying group the  $SO_{(3,11)}$ , which leads to the unavoidable appearance of ghosts due to the more than one time-like coordinates of the Lorentz group. Here instead we propose as a unifying group the  $SO_{(1,17)}$ , in which one can impose both Weyl and Majorana conditions and the final group obtained in four dimensions is the ordinary  $SO_{10}$  GUT [\[8\]](#page-10-6), followed by a global  $SU_2 \times SU_2$  symmetry.

# 2 The  $SO_{(1,17)}$  as unifying group

## <span id="page-2-5"></span>2.1 Geometrical construction

Starting with  $SO_{(1,17)}$  as the initial gauge symmetry group, we wish to produce symmetry breakings that will lead to the product of two symmetries, one describing gravity as a gauge theory, and the other describing the internal interactions. These breakings can occur via a SSB mechanism, or by imposing constraints to the theory. In order for the presentation of the model to be self-contained and amplify the latter (let's call it "soldering mechanism") over the well-known SSB mechanism, here we lay out and follow the analysis of [\[38](#page-11-12), [39](#page-11-11)], implemented for a 18−dimensional extended tangent space. The breakings of the present model via the usual Higgs mechanism is also described in the subsection [2.3.](#page-7-0)

At every point of a curved 4−dimensional Lorenzian metric space, we erect an 18−dimensional extended tangent space, following a construction analogous to [\[38,](#page-11-12) [39\]](#page-11-11). The extended tangent space is spanned by the vectors  $\mathbf{v}_A$ , where  $A = 0, ..., 17$  in such a way that the coordinate tangent space, spanned by the coordinate vectors  $e_{\mu} \equiv \partial/\partial x^{\mu}$ ,  $\mu = 0, ..., 3$ , is fully embedded<sup>[1](#page-2-0)</sup>. Having chosen  $SO_{(1,17)}$  as the structure group of the extended tangent homomorphisms, the scalar product of the basis vectors  $\{v_A\}$  should be orthonormal with respect to the extended, 18−dim Minkowskian metric,  $\eta_{AB} = \text{diag}(-1, +1, ..., +1)$ ,

<span id="page-2-1"></span>
$$
\mathbf{v}_A \cdot \mathbf{v}_B = \eta_{AB}.\tag{1}
$$

It is clear that the orthogonality of the basis vectors  $\{v_A\}$  is preserved under the extended  $SO_{(1,17)}$  Lorentz transformations.

It will prove convenient to separate the tangent space spanned by the basis vectors  $\{v_A\}$  into two orthogonal subspaces. The first is identified with the coordinate tangent space and spanned by the coordinate basis vectors  ${e_u}$ , which are orthonormal with respect to the metric of the base manifold,

<span id="page-2-4"></span>
$$
\mathbf{e}_{\mu} \cdot \mathbf{e}_{\nu} = g_{\mu\nu} \left( x \right). \tag{2}
$$

The second, that will be called internal tangent space, will be the orthogonal complement to the first and spanned by the set of 14 basis vectors  $\{\mathbf{n}_i\}$ , where  $i = 4, ..., 17$ , which are orthonormal with respect to the Euclidean metric,

<span id="page-2-3"></span>
$$
\mathbf{n}_i \cdot \mathbf{n}_j = \delta_{ij}.\tag{3}
$$

The projections of the extended tangent space basis vectors,  $\{v_A\}$ , onto the embedded coordinate tangent space basis vectors  $\{e_{\mu}\}\text{, are performed via the soldering forms } e^{A}_{\mu}$ ,

$$
\mathbf{e}_{\mu} = \mathbf{v}_{A} e^{A}{}_{\mu}.\tag{4}
$$

Then, with the aid of [\(1\)](#page-2-1) one obtains

<span id="page-2-2"></span>
$$
e_{A\mu} = \mathbf{v}_A \cdot \mathbf{e}_\mu. \tag{5}
$$

Multiplying both sides of [\(5\)](#page-2-2) with  $g^{\mu\nu}$  yields,

$$
e_A^{\ \mu} = \mathbf{v}_A \cdot \mathbf{e}^\mu,\tag{6}
$$

<span id="page-2-0"></span><sup>&</sup>lt;sup>1</sup>This structure can be formulated more mathematically by the fibre bundle theory.

where the  $\mathbf{v}_A$  are now projected onto the co-tangent basis,  $\{e^{\mu}\}\equiv\{dx^{\mu}\}\.$  Naturally, the soldering forms for which both indices are coordinate, we have,  $e_{\nu}^{\ \mu} = \delta_{\nu}^{\ \mu}$ .

The projections of the extended tangent space basis vectors  $\{v_A\}$ , onto the basis vectors  $\{n_i\}$  are performed via  $n_{i}^{A}$ 

$$
\mathbf{n}_i = \mathbf{v}_A n^A{}_i,\tag{7}
$$

which with the aid of [\(3\)](#page-2-3) lead to

<span id="page-3-0"></span>
$$
n_{Ai} = \mathbf{v}_A \cdot \mathbf{n}_i. \tag{8}
$$

By multiplying both sides of [\(8\)](#page-3-0) with  $\delta^{ij}$  one obtains,

$$
n_A^i = \mathbf{v}_A \cdot \mathbf{n}^i. \tag{9}
$$

Hence, a vector  $\mathbf{v}_A$  can be decomposed into the basis vectors  $\mathbf{e}_{\mu}$  and  $\mathbf{n}_i$  as

<span id="page-3-3"></span>
$$
\mathbf{v}_A = e_A^{\ \mu} \mathbf{e}_{\mu} + n_A^{\ i} \mathbf{n}_i. \tag{10}
$$

Using [\(2\)](#page-2-4) and [\(3\)](#page-2-3) we obtain expressions for the base manifold metric exclusively in terms of the soldering forms  $e^{A}{}_{\mu}$ 

<span id="page-3-2"></span>
$$
g_{\mu\nu} = \eta_{AB} e^A{}_{\mu} e^B{}_{\nu} = e^A{}_{\mu} e_{A_{\nu}},\tag{11}
$$

and, respectively, for the Euclidean metric exclusively in terms of the forms  $n^A_{i}$ ,

$$
\delta_{ij} = \eta_{AB} n^A{}_i n^B{}_j = n^A{}_i n_{Aj}.\tag{12}
$$

As remarked in [\[38\]](#page-11-12), attention should be paid to the fact that  $e_A^{\mu}$  is not inverse to  $e^A_{\mu}$  when the dimensions of the tangent space and the base manifold don't match<sup>[2](#page-3-1)</sup>. In other words, although it is obvious from [\(11\)](#page-3-2) that

$$
e_A{}^{\mu}e^A{}_{\nu} = \delta^{\mu}_{\nu},\tag{13}
$$

when contracting with respect to the tangent indices, it is also clear from  $(1)$  and  $(10)$ , that

<span id="page-3-6"></span>
$$
e_A^{\ \mu} e^B_{\ \mu} = \delta^A_B - n_A^{\ j} n^B_{\ j},\tag{14}
$$

given the orthonormality relations

$$
n_j^A e_A^\mu = 0, \quad n_j^A n_A^i = \delta_j^i. \tag{15}
$$

Parallel transport is defined via the action of affine and spin connections, on the coordinate and extended tangent space basis vectors respectively,

<span id="page-3-4"></span>
$$
\nabla_{\nu} \mathbf{e}_{\mu} = \Gamma^{\lambda}{}_{\nu\mu} \mathbf{e}_{\lambda}, \quad \nabla_{\nu} \mathbf{v}_{A} = -\omega_{\nu A}{}^{B} \mathbf{v}_{B}, \tag{16}
$$

where  $\nabla_{\nu}$  is the covariant derivative along the direction of the tangent basis vector  $\mathbf{e}_{\nu}$ .

By defining the parallel transport of the coordinate basis vectors as above, a constraint has actually been imposed to the geometrical construction, as the most general form of it would be  $\nabla_{\nu} \mathbf{e}_{\mu} = \Gamma^{\lambda}{}_{\nu\mu} \mathbf{e}_{\lambda} + B^{i}{}_{\nu\mu} \mathbf{n}_{i}$ . The imposed constraint,  $B^i_{\ \nu\mu} = 0$ , causes the breaking

$$
SO_{(1,17)} \to SO_{(1,3)} \times SO_{14}.\tag{17}
$$

Having imposed this constraint, can be shown [\[39\]](#page-11-11) that the covariant derivative of the internal basis vectors is also an element of the internal subspace. Hence we have as well,

<span id="page-3-5"></span>
$$
\nabla_{\nu} \mathbf{n}_{i} = -A_{\nu i}^{j} \mathbf{n}_{j}.
$$
\n(18)

<span id="page-3-1"></span><sup>&</sup>lt;sup>2</sup>Recalling the definition of  $e^{A}{}_{\mu}$  as a projector of 18-dimensional vectors onto the 4-dimensional tangent space, it is clear that  $e_A^{\mu}$  cannot be considered as a reversed projector operator, since a lower dimensional vector cannot be projected onto a space of higher dimensionality. As already stated,  $e_A^{\mu}$  are projections of the vectors  $\mathbf{v$ 

The covariant derivative when acting on scalars, naturally coincides with the ordinary derivative<sup>[3](#page-4-0)</sup>. Therefore, the metricity condition,

<span id="page-4-2"></span>
$$
\nabla_{\nu}\eta_{AB}=0,\t\t(19)
$$

must hold. Following [\(16\)](#page-3-4), the action of the covariant derivative operator on a tangent vector  $V = V^{\mu} \mathbf{e}_{\mu}$  is,

$$
\nabla_{\nu} \left( V^{\mu} \mathbf{e}_{\mu} \right) = \left( \partial_{\nu} V^{\mu} \right) \mathbf{e}_{\mu} + V^{\mu} \nabla_{\nu} \mathbf{e}_{\mu} = \left( \partial_{\nu} V^{\mu} + V^{\lambda} \Gamma^{\mu}{}_{\nu \lambda} \right) \mathbf{e}_{\mu} = \left( \nabla_{\nu} V \right)^{\mu} \mathbf{e}_{\mu}.
$$
 (20)

It is convenient to use a different symbol to express the total action of the covariant derivative on a vector as an action upon its components<sup>[4](#page-4-1)</sup>. Guided from the above equation we define,

<span id="page-4-5"></span>
$$
D_{\nu}V^{\mu} \equiv (\nabla_{\nu}V)^{\mu} = \partial_{\nu}V^{\mu} + \Gamma^{\mu}{}_{\nu\lambda}V^{\lambda}.
$$
\n(21)

Similarly, for co-vectors  $W = W_{\mu} dx^{\mu}$ , we should have,

<span id="page-4-6"></span>
$$
D_{\nu}W_{\mu} = \partial_{\nu}W_{\mu} - W_{\lambda}\Gamma^{\lambda}{}_{\nu\mu}.
$$
\n(22)

Analogous equations hold for the action of the covariant derivative on the components of an extended tangent space spinor,  $\psi = \psi^A \mathbf{v}_A$ , now with the spin connection performing parallel transportation,

<span id="page-4-8"></span>
$$
D_{\nu}\psi^{A} = \partial_{\nu}\psi^{A} - \psi^{B}\omega_{\nu B}{}^{A},\tag{23}
$$

and for the the action of the covariant derivative on the components of coordinate space spinors,

<span id="page-4-9"></span>
$$
D_{\nu}X^{i} = \partial_{\nu}X^{i} - X^{j}A_{\nu j}^{i}.
$$
\n
$$
(24)
$$

Simultaneous validity of [\(1\)](#page-2-1) and [\(19\)](#page-4-2), implies that the spin connection is antisymmetric under the interchange of its extended tangent space indices,

<span id="page-4-7"></span>
$$
\omega_{\nu AB} = -\omega_{\nu BA}.\tag{25}
$$

Covariant differentiation of [\(5\)](#page-2-2) and taking into account [\(16\)](#page-3-4) implies,

<span id="page-4-3"></span>
$$
\partial_{\nu} e_{A\mu} = -\omega_{\nu A}{}^{B} e_{B\mu} + \Gamma^{\lambda}{}_{\mu\nu} e_{A\lambda}.
$$
\n(26)

In perfect analogy, covariant differentiation of [\(8\)](#page-3-0) yields the equation

<span id="page-4-10"></span>
$$
\partial_{\mu}n_{Ai} = -\omega_{\mu A}{}^{B}n_{Bi} - A_{\mu i}{}^{j}n_{Aj}.\tag{27}
$$

Equations [\(26\)](#page-4-3) are  $4 \times 18 \times 4 = 288$  in number and should be solved for the affine and spin connection components, in terms of the given soldering forms. In a torsion-free base manifold, the affine connection is symmetric over the interchange of its lower indices,

$$
\Gamma^{\lambda}{}_{\mu\nu} = \Gamma^{\lambda}{}_{\nu\mu}.\tag{28}
$$

This means that the number of independent affine connection components are  $4 \times 10 = 40$  and can be determined separately.

Let us first operate with the covariant derivative  $\nabla_{\lambda}$  on [\(11\)](#page-3-2),

<span id="page-4-4"></span>
$$
\nabla_{\lambda} g_{\mu\nu} = \partial_{\lambda} g_{\mu\nu} = \partial_{\lambda} \left( e^{A}{}_{\mu} e_{A\nu} \right) = \left( \partial_{\lambda} e^{A}{}_{\mu} \right) e_{A\nu} + e^{A}{}_{\mu} \partial_{\lambda} e_{A\nu}.
$$
\n(29)

Then, combining [\(26\)](#page-4-3) and [\(29\)](#page-4-4), we obtain

$$
\Gamma^{\rho}{}_{\mu\lambda}g_{\rho\nu} + \Gamma^{\rho}{}_{\nu\lambda}g_{\mu\rho} = \partial_{\lambda}g_{\mu\nu},\tag{30}
$$

<span id="page-4-0"></span><sup>3</sup> In the current analysis, tensor components are considered scalar functions. See also [\[41](#page-11-16)].

<span id="page-4-1"></span> $^{4}$ In general, the  $\nabla$  operator performs parallel transport in a passive way, acting on the basis vectors as denoted in [\(16\)](#page-3-4) and [\(18\)](#page-3-5), treating the tensor components as scalar functions. In contrast, the D operator performs parallel transport in an active way, acting as a covariant derivative upon the tensor components leaving the basis vectors intact, i.e. [\(21\)](#page-4-5), [\(22\)](#page-4-6).

which when inverted, gives the explicit expression for the torsion-free Christofell connection in terms of the metric,

$$
\Gamma^{\gamma}{}_{\mu\nu} = \frac{1}{2} g^{\lambda\rho} \left( g_{\mu\rho,\nu} + g_{\rho\nu,\mu} - g_{\mu\nu,\rho} \right). \tag{31}
$$

What we have actually done here was to employ 40 out of the 288 equations [\(26\)](#page-4-3) to fully determine  $\Gamma^{\gamma}{}_{\mu\nu}$ . Therefore, we are now left with 248 equations to determine the components of the spin connection which is antisymmetric over its last two indices [\(25\)](#page-4-7). Since the  $\nu$  index runs over the 4 base manifold dimensions, and the indices A, B over the 18 extended tangent space dimensions, we see that there are 612 components to be determined, while there are only 248 equations in our disposal. This implies that we are left with 612−248 = 364 undefined spin connection components, a number which matches the  $SO_{14}$  gauge fields (multiplied by 4).

The above results confirm that the initial group,  $SO_{(1,17)}$ , is defined to have an inner structure related to the geometry of the coordinate manifold. Its first 4 dimensions correspond to the tangent space of the manifold, while the rest 14 remain unmixed with them, showing that the initial gauge group has been reduced to the direct product  $SO_{(1,3)} \times SO_{14}$ . By the first group,  $SO_{(1,3)}$ , we are going to describe the spacetime geometry, while by  $SO_{14}$  the internal interactions.

The local Lorentz transformation law of a basis vector,  $\mathbf{v}_A$ , is

<span id="page-5-0"></span>
$$
\mathbf{v}_A \to \tilde{\mathbf{v}}_A = \Lambda_A{}^B \mathbf{v}_B,\tag{32}
$$

thus the soldering forms transform covariantly, as

$$
e_A^{\ \mu} \to \tilde{e}_A^{\ \mu} = \Lambda_A^{\ B} e_B^{\ \mu}.\tag{33}
$$

Using the local Lorentz transformation, [\(32\)](#page-5-0), and the parallel transportation rule, [\(16\)](#page-3-4), of the basis vectors, we can also show that

$$
\omega_{\nu A}{}^{B} \to \tilde{\omega}_{\nu A}{}^{B} = \left(\Lambda \omega_{\nu} \Lambda^{-1}\right)_{A}{}^{B} + \left(\Lambda \partial_{\nu} \Lambda^{-1}\right)_{A}{}^{B},\tag{34}
$$

which explicitly shows that the spin connection transforms under local Lorentz transformations as a Yang-Mills field, with  $SO_{(1,17)}$  in the role of the gauge group.

### 2.2 Constructing the total action

Taking into account the expressions [\(21\)](#page-4-5) and [\(22\)](#page-4-6), we can derive the explicit form for the affine curvature tensor, following the definition,

$$
[D_{\nu}, D_{\lambda}]V^{\mu} = V^{\rho}R^{\mu}_{\rho\nu\lambda}.
$$
\n(35)

Indeed, after some straightforward algebra we obtain,

$$
[D_{\nu}, D_{\lambda}]V^{\mu} = V^{\rho} \left( \partial_{\nu} \Gamma^{\mu}{}_{\rho\lambda} - \partial_{\lambda} \Gamma^{\mu}{}_{\rho\nu} + \Gamma^{\sigma}{}_{\rho\lambda} \Gamma^{\mu}{}_{\sigma\nu} - \Gamma^{\sigma}{}_{\rho\nu} \Gamma^{\mu}{}_{\sigma\lambda} \right), \tag{36}
$$

thus, it is clear that we can identify the affine curvature tensor,

$$
R^{\mu}{}_{\rho\nu\lambda} = \partial_{\nu}\Gamma^{\mu}{}_{\lambda\rho} - \partial_{\lambda}\Gamma^{\mu}{}_{\nu\rho} + \Gamma^{\mu}{}_{\nu\sigma}\Gamma^{\sigma}{}_{\lambda\rho} - \Gamma^{\mu}{}_{\lambda\sigma}\Gamma^{\sigma}{}_{\nu\rho}.
$$
 (37)

Similarly, following the definition for the spin curvature tensor,

$$
[D_{\mu}, D_{\nu}]\psi^{A} = \psi^{B} R_{\mu\nu}{}^{A}{}_{B}, \tag{38}
$$

and employing [\(23\)](#page-4-8) we obtain,

$$
R_{\mu\nu}^{\quad AB} = \partial_{\mu}\omega_{\nu}^{\quad AB} - \partial_{\nu}\omega_{\mu}^{\quad AB} + \omega_{\mu}^{\quad A}{}_{C}\omega_{\nu}^{\quad CB} - \omega_{\nu}^{\quad A}{}_{C}\omega_{\mu}^{\quad CB}.\tag{39}
$$

Finally, following the definition for the coordinate curvature tensor,

$$
[D_{\mu}, D_{\nu}]X^{i} = X^{j}F_{\mu\nu}{}_{j}^{i}, \qquad (40)
$$

and employing [\(24\)](#page-4-9) we get,

$$
F_{\mu\nu}^{i j} = \partial_{\mu} A_{\nu}^{i j} - \partial_{\nu} A_{\mu}^{i j} + A_{\mu}^{i}{}_{k} A_{\nu}^{k j} - A_{\nu}^{i}{}_{k} A_{\mu}^{k j}.
$$
\n(41)

Taking the partial derivative  $\partial_{\rho}$  of [\(26\)](#page-4-3) and substracting the same result but with the indices  $\nu$  and  $\rho$ interchanged, we end up to a relation between the spin and affine curvature tensors,

<span id="page-6-0"></span>
$$
R_{\mu\nu}{}^{AB}(\omega)e_{B\lambda} = e^A_{\ \rho}R_{\mu\nu}{}^{\rho}_{\lambda}(\Gamma). \tag{42}
$$

Following analogous procedure in [\(27\)](#page-4-10) we end up to a relation among the spin and coordinate curvature tensors,

<span id="page-6-1"></span>
$$
n_B{}^i R_{\mu\nu A}{}^B(\omega) = n_A{}^j F_{\mu\nu j}{}^i(A). \tag{43}
$$

Employing [\(14\)](#page-3-6), equation [\(42\)](#page-6-0) can be rewritten as an expansion on the basis vectors of the two subspaces of the extended tangent space,

$$
R_{\mu\nu}{}^{AB}(\omega) = R_{\mu\nu}{}^{AC}(\omega) n_C{}^i n^B{}_i + R^\rho{}_{\lambda\mu\nu}(\Gamma) e^A{}_{\rho} e^{B\lambda}.\tag{44}
$$

Substituting [\(43\)](#page-6-1) in the equation above, we obtain

$$
R_{\mu\nu}{}^{AB}(\omega) = n^A{}_i n^B{}_j F_{\mu\nu}{}^{ij}(A) + e^A{}_{\rho} e^{BA} R_{\mu\nu}{}^{\rho}{}_{\lambda}(\Gamma). \tag{45}
$$

We see that the spin curvature has been completely decomposed into the coordinate and affine curvatures. From this expression, all the invariants of the theory up to second order will be produced.

For the first order, the only contraction possible is

$$
R_{\mu\nu}^{\ A B}(\omega)e_A^{\ \mu}e_B^{\ \nu} = R(\Gamma),\tag{46}
$$

which produces the Ricci scalar of the theory. The second order invariants are

$$
R^2(\Gamma), R_{\mu\nu}(\Gamma)R^{\mu\nu}(\Gamma), R_{\mu\nu\lambda\delta}(\Gamma)R^{\mu\nu\lambda\delta}(\Gamma), \tag{47}
$$

which get produced by various combinations of soldering forms acting on the curvature 2-form. The kinetic terms are going to be produced by the contraction

$$
g^{\mu\lambda}g^{\nu\delta}R_{\mu\nu}{}^{AB}(\omega)R_{\lambda\delta AB}(\omega) = g^{\mu\lambda}g^{\nu\delta}\left(F_{\mu\nu}{}^{ij}(A)F_{\lambda\delta ij}(A)\right) + R_{\mu\nu\lambda\delta}(\Gamma)R^{\mu\nu\lambda\delta}(\Gamma). \tag{48}
$$

Now we have produced all the curvature invariants up to second order. In the general action, 512−dimensional Dirac spinor fields also have to be included,

$$
\int d^4x \sqrt{-g} \bar{\psi} i \Gamma^A e_A^{\ \mu} D_{\mu} \psi,\tag{49}
$$

where  $\Gamma_A$  matrices satisfy the Clifford algebra

$$
\left\{\Gamma^A, \Gamma^B\right\} = 2\eta^{AB},\tag{50}
$$

and

$$
D_{\mu} \equiv \partial_{\mu} + \frac{1}{4} \omega_{\mu}{}^{AB} S_{AB},\tag{51}
$$

where  $S_{AB} = \frac{1}{2} [\Gamma_A, \Gamma_B]$  are the generators of the  $SO_{(1,17)}$  algebra.

Hence, the expression of the general action of the theory is

$$
I_{SO_{(1,17)}} = \int d^4x \sqrt{-g} \left[ \frac{1}{16\pi G} R_{\mu\nu}{}^{AB}(\omega) e_A{}^{\mu} e_B{}^{\nu} + R_{\mu\nu}{}^{AB} R_{\lambda\delta}{}^{CD} \left( a e_A{}^{\mu} e_B{}^{\nu} e_C{}^{\lambda} e_D{}^{\delta} + b e_A{}^{\mu} e_C{}^{\nu} e_B{}^{\lambda} e_D{}^{\delta} + c e_C{}^{\mu} e_D{}^{\nu} e_A{}^{\lambda} e_B{}^{\delta} \right) \right]
$$
\n
$$
- \frac{1}{4} g^{\mu\lambda} g^{\nu\delta} R_{\mu\nu}{}^{AB}(\omega) R_{\lambda\delta AB}(\omega) + \bar{\psi} i \Gamma^A e_A{}^{\mu} D_{\mu} \psi \right] \Rightarrow \tag{52}
$$

$$
I_{SO_{(1,3)}\times SO_{14}} = \int d^4x \sqrt{-g} \left[ \frac{1}{16\pi G} R(\Gamma) + aR^2(\Gamma) - bR_{\mu\nu}(\Gamma)R^{\mu\nu}(\Gamma) + (c - \frac{1}{4})R_{\mu\nu\lambda\delta}(\Gamma)R^{\mu\nu\lambda\delta}(\Gamma) - \frac{1}{4}g^{\mu\lambda}g^{\nu\delta}F_{\mu\nu}{}^{ij}(A)F_{\lambda\delta ij}(A) + \phi_{SO_{(1,3)}}i\Gamma^{\mu}D_{\mu}\psi_{SO_{(1,3)}} + \bar{\psi}_{SO_{14}}i\Gamma^j e_j^{\mu}D_{\mu}\psi_{SO_{14}} \right],
$$
\n
$$
(53)
$$

where a, b, c are dimensionless constants, and the Weyl representation has been chosen for the Gamma matrices. The above action consists of  $SO_{(1,3)}$  and  $SO_{14}$  invariants, as expected. By setting  $a = \frac{b}{4} = c - \frac{1}{4}$ , the curvature terms form the integrand of the Gauss-Bonnet topological invariant, hence they do not contribute to the field equations, avoiding in this way the appearance of ghosts [\[42](#page-11-17)]. By the  $SO_{(1,3)}$  part we are able to retrieve Einstein's gravity as a gauge theory [\[25](#page-11-4)[–27](#page-11-5)], while by the  $SO_{14}$  we are going to describe internal interactions.

### <span id="page-7-0"></span>2.3 Breakings

According to the subsection [2.1,](#page-2-5) the original gauge symmetry,  $SO_{(1,17)}$ , of the theory, is being reduced to  $SO_{(1,3)} \times SO_{14}$  by employing the soldering mechanism presented above, i.e. the  $C_{SO_{(1,17)}}(SO_{(1,3)}) = SO_{14}$ remains as the gauge group that will describe the internal interactions. The same breaking can occur via Higgs mechanism, by introducing a scalar field in the 170 representation<sup>[5](#page-7-1)</sup> of  $SO_{18}$  [\[45\]](#page-11-18), and with the help of a Lagrange multiplier we can break the gauge symmetry non-linearly [\[31,](#page-11-8) [46](#page-11-19)],

$$
SO_{18} \to SU_2 \times SU_2 \times SO_{14}
$$
  
170 = (1,1,1) + (3,3,1) + (2,2,14) + (1,1,104). (54)

In order to break further the resulting  $SO_{14}$  gauge symmetry to a symmetry of a more familiar GUT, such as  $SO_{10}$ , we can employ a second Higgs mechanism by using the 104 representation<sup>[6](#page-7-2)</sup> of  $SO_{14}$  [\[47,](#page-11-20) [48\]](#page-11-21),

$$
SO_{14} \to SU_2 \times SU_2 \times SO_{10}
$$
  
104 = (1, 1, 1) + (3, 3, 1) + (2, 2, 10) + (1, 1, 54). (55)

Of the rest  $SU_2 \times SU_2 \times SU_2 \times SU_2$  symmetry that remains, one part  $SU_2 \times SU_2$  should be used for describing gauge gravity, while the other  $SU_2 \times SU_2$  should be broken. The irreducible spinor representation of  $SO_{18}$  is 256, which under  $SU_2 \times SU_2 \times SO_{14}$ , decomposes as

$$
256 = (2, 1, 64) + (1, 2, 64), \tag{56}
$$

while the irreducible spinor representation of  $SO_{14}$  is 64, that under  $SU_2 \times SU_2 \times SO_{10}$  decomposes as

$$
64 = (2, 1, 16) + (1, 2, \overline{16}). \tag{57}
$$

<span id="page-7-1"></span><sup>&</sup>lt;sup>5</sup>The representation 3060 can be also used for that purpose. The two breakings differ on the presence of a remaining unbroken parity symmetry in the case of 170, of which the  $SO_{10}$  analogous is discussed in detail in [\[43](#page-11-22), [44\]](#page-11-23).

<span id="page-7-2"></span><sup>&</sup>lt;sup>6</sup>Had we chosen the 3060 of  $SO_{18}$  for the previous breaking, the appropriate representation for the breaking of  $SO_{14}$  would be 1001 with analogous effects on the remaining symmetry.

By introducing further two scalars in the 256 representation of  $SO_{18}$ , when they take VEVs in their  $(\langle 2 \rangle, 1, 16)$ and  $(1,\langle 2\rangle, \overline{16})$  components of  $SO_{14}$  under the  $SU_2 \times SU_2 \times SO_{10}$  decomposition, the final unbroken gauge symmetry is  $SO_{10}$ . The final total symmetry that we are left with is

$$
[SU_2 \times SU_2]_{\text{Lorentz}} \times [SU_2 \times SU_2]_{\text{Global}} \times SO_{10 \text{Gauge group}}.\tag{58}
$$

# 3 Weyl and Majorana spinors

A Dirac spinor,  $\psi$ , has  $2^{D/2}$  independent components in D dimensions. The Weyl and Majorana constraints each divide the number of independent components by 2. The Weyl condition can be imposed only for D even, so a Weyl-Majorana spinor has  $2^{(D-4)/2}$  independent components (when D is even). Weyl-Majorana spinors can exist only for  $D = 4n + 2$ ; real Weyl-Majorana spinors can exist for  $D = 2$  modulo 8, and pseudoreal Weyl-Majorana spinors can exist for  $D = 6$  modulo 8.

The unitary representations of the Lorentz group  $SO_{(1,D-1)}$  are labeled by a continuous momentum vector k, and by a spin "projection", which in D dimensions is a representation of the compact subgroup  $SO_{(D-2)}$ . The Dirac, Weyl, Majorana, and Weyl-Majorana spinors carry indices that transform as finite-dimensional non-unitary spinor representations of  $SO_{(1,D-1)}$ .

It is well-known (see e.g. [\[49\]](#page-11-24) for a review, or [\[50\]](#page-11-25) for a more concise description) that the type of spinors one obtains for  $SO_{(p,q)}$  in the real case is governed by the signature  $(p-q)$  mod 8. Among even signatures, signature 0 gives a real representation, signature 4 a quaternionic representation, while signatures 2 and 6 give complex representations. In the case of  $SO_{(1,17)}$  the signature is zero, and the imposition of the Majorana condition on the spinors is permitted, in addition to Weyl.

Let us recall for completeness and fixing the notation, the otherwise well-known case of 4 dimensions. The  $SO_{(1,3)}$  spinors in the usual  $SU_2 \times SU_2$  basis transform as  $(2, 1)$  and  $(1, 2)$ , with representations labeled by their dimensionality. The 2–component Weyl spinors,  $\psi_L$  and  $\psi_R$ , transform as the irreducible spinors,

$$
\psi_L \sim (2, 1), \quad \psi_R \sim (1, 2), \tag{59}
$$

of  $SU_2 \times SU_2$  with "∼" meaning "transforms as". A Dirac spinor,  $\psi$ , can be made from the direct sum of  $\psi_L$ and  $\psi_R$ ,

$$
\psi \sim (2,1) \oplus (1,2). \tag{60}
$$

In 4-component notation the Weyl spinors in the Weyl basis are  $(\psi_L, 0)$  and  $(0, \psi_R)$ , and are eigenfunctions of  $\gamma^5$  with eigenvalues +1 and -1, respectively.

The usual Majorana condition for a Dirac spinor has the form,

<span id="page-8-0"></span>
$$
\psi = C\bar{\psi}^T,\tag{61}
$$

where  $C$  is the charge-conjugation matrix. In 4 dimensions  $C$  is off-diagonal in the Weyl basis, since it maps the components transforming as  $(2, 1)$  into  $(1, 2)$ . Therefore, if one tries to impose  $(61)$  on a Weyl spinor, there is no non-trivial solution and therefore Weyl-Majorana spinors do not exist in 4 dimensions.

For D even, it is always possible to define a Weyl basis where  $\Gamma^{D+1}$  (the product of all D  $\Gamma$  matrices) is diagonal, so

$$
\Gamma^{D+1}\psi_{\pm} = \pm \psi_{\pm}.\tag{62}
$$

We can express  $\Gamma^{D+1}$  in terms of the chirality operators in 4 and extra d dimensions,

$$
\Gamma^{D+1} = \gamma^5 \otimes \gamma^{d+1}.\tag{63}
$$

Therefore the eigenvalues of  $\gamma^5$  and  $\gamma^{d+1}$  are interrelated. However, clearly the choice of the eigenvalue of  $\Gamma^{D+1}$  does not impose the eigenvalues on the interrelated  $\gamma^5$  and  $\gamma^{d+1}$ .

Since  $\Gamma^{D+1}$  commutes with the Lorentz generators, then each of the  $\psi_+$  and  $\psi_-$  transforms as an irreducible spinor of  $SO_{(1,D-1)}$ . For D even, the  $SO_{(1,D-1)}$  always has two independent irreducible spinors; for  $D = 4n$ there are two self-conjugate spinors  $\sigma_D$  and  $\sigma_D'$ , while for  $D = 4n + 2$ ,  $\sigma_D$  is non-self-conjugate and  $\bar{\sigma}_D$  is the other spinor. By convention is selected  $\psi_+ \sim \sigma_D$  and  $\psi_- \sim \sigma_D'$  or  $\bar{\sigma}_D$ . Accordingly, Dirac spinors are defined as direct sum of Weyl spinors,

$$
\psi = \psi_+ \oplus \psi_- \sim \begin{cases} \sigma_D \oplus \sigma_D' & \text{for } D = 4n \\ \sigma_D \oplus \sigma_D' & \text{for } D = 4n + 2. \end{cases}
$$
(64)

When  $D$  is odd there are no Weyl spinors, as already mentioned.

The Majorana condition can be imposed in  $D = 2, 3, 4 + 8n$  dimensions and therefore the Majorana and Weyl conditions are compatible only in  $D = 4n + 2$  dimensions.

Let us limit ourselves here in the case that  $D = 4n + 2$ , while for the rest one can consult refs [\[9](#page-10-7), [16\]](#page-10-13). Starting with Weyl-Majorana spinors in  $D = 4n + 2$  dimensions, we are actually forcing a representation  $f_R$  of a gauge group defined in higher dimensions to be the charge conjugate of  $f_L$ , and we arrive in this way to a 4-dimensional theory with the fermions only in the  $f<sub>L</sub>$  representation of the gauge group.

In our case, we have for the Weyl spinor of  $SO_{(1,17)}$ :

$$
SO_{(1,17)} \rightarrow [SU_2 \times SU_2]_{\text{Lorentz}} \times SO_{14} \text{Gauge group}
$$

$$
\sigma_{18} = 256 = (2, 1; 64) + (1, 2; \bar{64}).\tag{65}
$$

Then, the Majorana condition maps the  $(2, 1; 64)$  into the  $(1, 2; 64)$ . Therefore in 4 dimensions, only the  $(2, 1; 64)$  remains from the spinor 256 of  $SO_{18}$ , after imposing the Majorana condition, i.e. we obtain  $SO_{14}$  with  $64L$ .

On the other hand, the spinor of  $SO_{14}$ ,  $64_L$ , has the following decomposition after the SSB of  $SU_2 \times SU_2$ , as described in [2.3](#page-7-0) under  $[SU_{2}\times SU_{2}]_{\rm Global}\times SO_{10}$ :

$$
SO_{14} \to [SU_2 \times SU_2]_{\text{Global}} \times SO_{10 \text{Gauge}} \tag{66}
$$

$$
64 = (2, 1; 16) + (1, 2; \overline{16}). \tag{67}
$$

Therefore, after imposing Weyl (by choosing  $\sigma_{18}$ ) and Majorana conditions in 4 dimensions, we obtain  $f_L = (2, 1, 16)_L$  and  $(1, 2, 16)_L = (1, 2, 16)_R = g_R$ . The  $f_L$ ,  $g_R$  are eigenfunctions of the  $\gamma^5$  matrix with eigenvalues +1 and −1 respectively, as already mentioned. Keeping only the +1 eigenvalue, i.e. imposing an additional discrete symmetry, we are left with  $(2, 1, 16)$ <sub>L</sub>.

## 4 Conclusions

In the present work we have constructed a realistic model based on the idea that unification of gravity and internal interactions in four dimensions can be achieved by gauging an enlarged tangent Lorentz group. The enlarged group used in our construction is originally the  $SO_{(1,17)}$ , which eventually, in the broken phase, leads to GR and the  $SO_{10}$  GUT accompanied by an  $SU_2 \times SU_2$  global symmetry. The latter leads to even number of families.

In the phenomenological analysis that will be presented in a future work, obviously we will include appropriate scalar fields that will (i) break the  $SO_{10}$  to the SM and (ii) make the fourth generation of fermions heavy in the minimal setting of fermions in the model.

The unifying group  $SO_{(1,17)}$  does not contain the whole Poincaré group, but rather the Lorentz rotations,  $SO_{(1,3)}$ . Therefore, a conflict with the Coleman-Mandula (C-M) theorem, which states that internal and spacetime symmetries cannot be mixed [\[51](#page-11-26)], is avoided. Recall that the C-M theorem has several hypotheses with the most relevant being that the theory is Poincaré invariant. It might be a challenge of a further study to start with a unifying group that includes translational symmetry and examine consistency with the C-M theorem.

Another point concerning the phenomenological analysis is that in the present scheme we will do a RG analysis, in which, in addition to considering the various spontaneous symmetry breakings of  $SO_{10}$ , the fact that the unification scale is the Planck scale should also be taken into account. Finally, we note that the use of the RG analysis is legitimate based on the theorem [\[52\]](#page-11-27) stating that if an effective 4−dimensional theory is renormalizable by power counting, then it is consistent to consider it as renormalizable a la Wilson [\[53](#page-11-28)[–55](#page-11-29)].

### Acknowledgments

It is a pleasure to thank Pantelis Manousselis for continuous discussions and suggestions on the content of the paper. We would like also to thank Ali Chamseddine, Slava Mukhanov and Roberto Percacci for several discussions on their work. Finally, we would like to thank Peter Forgacs, Dieter Lust and Roberto Percacci for reading the manuscript and for their comments. D.R. would like to thank NTUA for a fellowship for doctoral studies. D.R. and G.Z. have been supported by the Basic Research Programme, PEVE2020 of National Technical University of Athens, Greece. G.Z. would like to thank CERN-TH, MPP-Munich, ITP-Heidelberg and DFG Exzellenzcluster 2181:STRUCTURES of Heidelberg University for their hospitality and support.

## <span id="page-10-0"></span>References

- <span id="page-10-1"></span>[1] Th. Kaluza. Sitzungsber. Preuss. Akad. Wiss. Berlin (Math. Phys.) (1921) 966.
- <span id="page-10-2"></span>[2] O. Klein. Z. Phys. 37 (1926) 895.
- [3] R. Kerner. Ann. Inst. H. Poincare Phys. Theor. 9 (1968) 143.
- <span id="page-10-3"></span>[4] Y.M. Cho. J. Math. Phys. 16 (1975) 2029.
- <span id="page-10-4"></span>[5] Y.M. Cho and P.G.O. Freund. Phys. Rev. D 12 (1975) 1711.
- <span id="page-10-5"></span>[6] E. Witten. Conf. Proc. C 8306011 (1983) 227.
- <span id="page-10-6"></span>[7] H. Georgi and S.L. Glashow. Phys. Rev. Lett. 32 (1974) 438.
- <span id="page-10-7"></span>[8] H. Fritzsch and P. Minkowski. Ann. Phys. 93 (1975) 193.
- <span id="page-10-8"></span>[9] G. Chapline and R. Slansky. Nucl. Phys. B 209 (1982) 461.
- [10] M.B. Green et al. Superstring Theory. Vol. 1 & 2. Cambridge University Press, 1988.
- <span id="page-10-9"></span>[11] J. Polchinski. *String theory*. Vol. 1 & 2. Cambridge University Press, 1998.
- <span id="page-10-10"></span>[12] D. Lust and S. Theisen. *Lectures on String Theory*. Vol. 346. 1989.
- <span id="page-10-11"></span>[13] D.J. Gross, J.A. Harvey, E. Martinec, and R. Rohm. Nucl. Phys. B 256 (1985) 253.
- <span id="page-10-12"></span>[14] P. Forgåcs and N.S. Manton. Commun. Math. Phys. 72 (1980) 15.
- <span id="page-10-13"></span>[15] N. Irges and G. Zoupanos. Phys. Lett. B 698 (2011) 146.
- [16] D. Kapetanakis and G. Zoupanos. Phys. Rept. 219 (1992) 4.
- [17] Y.A. Kubyshin, I.P. Volobuev, J.M. Mourao, and G. Rudolph. Dimensional Reduction of Gauge Theories, Spontaneous Compactification and Model Building. Vol. 349. Springer, 1989.
- <span id="page-11-1"></span>[18] J. Scherk and J.H. Schwarz. Nucl. Phys. B 153 (1979) 61.
- [19] N.S. Manton. Nucl. Phys. B 193 (1981) 502.
- [20] D. Lust and G. Zoupanos. Phys. Lett. B 165 (1985) 309.
- <span id="page-11-0"></span>[21] P. Manousselis and G. Zoupanos. JHEP 2004.11 (2004) 025.
- <span id="page-11-2"></span>[22] A. Chatzistavrakidis and G. Zoupanos. JHEP 09 (2009) 077.
- <span id="page-11-3"></span>[23] G. Manolakos, G. Patellis, and G. Zoupanos. Phys. Lett. B 813 (2021).
- <span id="page-11-4"></span>[24] G. Patellis, W. Porod, and G. Zoupanos (2023). arXiv: [2307.10014 \[hep-ph\]](https://arxiv.org/abs/2307.10014).
- [25] R. Utiyama. Phys. Rev. 101 (1956) 1597.
- <span id="page-11-5"></span>[26] T.W.B. Kibble. J. Math. Phys. 2 (1961) 212.
- <span id="page-11-6"></span>[27] S.W. MacDowell and F. Mansouri. Phys. Rev. Lett. 38 (1977) 739.
- <span id="page-11-7"></span>[28] D.Z. Freedman and A. Van Proeyen. Supergravity. Cambridge University Press, 2012.
- [29] A. Chatzistavrakidis et al. Fortsch. Phys. 66 (2018) 1800047.
- <span id="page-11-8"></span>[30] G. Manolakos, P. Manousselis, and G. Zoupanos. JHEP 08 (2020) 001.
- <span id="page-11-9"></span>[31] G. Manolakos, P. Manousselis, and G. Zoupanos. Fortsch. Phys. 69 (2021) 2100085.
- <span id="page-11-10"></span>[32] S. Weinberg. Fifth Workshop on Grand Unification. 1984.
- [33] R. Percacci. Phys. Lett. B 144 (1984) 37.
- <span id="page-11-14"></span>[34] R. Percacci. Nucl. Phys. B 353 (1991) 271.
- <span id="page-11-15"></span>[35] F. Nesti and R. Percacci. J.Phys.A 41 (2008) 075405.
- [36] F. Nesti and R. Percacci. Phys. Rev. D 81 (2010) 025010.
- <span id="page-11-12"></span>[37] K. Krasnov and R. Percacci. Class. Quant. Grav. 35.14 (2018) 143001.
- <span id="page-11-11"></span>[38] A.H. Chamseddine and V. Mukhanov. *JHEP* 03 (2010) 033.
- <span id="page-11-13"></span>[39] A.H. Chamseddine and V. Mukhanov. *JHEP* 03 (2016) 020.
- [40] G. Manolakos, P. Manousselis, D. Roumelioti, S. Stefas, and G. Zoupanos. *Eur. Phys. J.ST* (2023). DOI: [https://doi.org/10.1140/epjs/s11734-023-00830-8](https://doi.org/https://doi.org/10.1140/epjs/s11734-023-00830-8). arXiv: [2305.11785](https://arxiv.org/abs/2305.11785).
- <span id="page-11-17"></span><span id="page-11-16"></span>[41] M. Nakahara. *Geometry, Topology and Physics*. CRC Press, 2003.
- <span id="page-11-22"></span>[42] L. Alvarez-Gaume, A. Kehagias, C. Kounnas, D. Lust, and A. Riotto. Fortsch. Phys. 64 (2016) 176.
- <span id="page-11-23"></span>[43] D. Chang, R.N. Mohapatra, and M.K. Parida. Phys. Rev. Lett. 52 (1984) 1072.
- <span id="page-11-18"></span>[44] D. Chang et al. *Phys. Rev. D* 31 (1985) 1718.
- <span id="page-11-19"></span>[45] R. Feger., T.W. Kephart, and R.J. Saskowski. Comput. Phys. Commun. 257 (2020) 107490.
- <span id="page-11-20"></span>[46] K.S. Stelle and P.C. West. *Phys. Rev. D* 21 (1980) 1466.
- <span id="page-11-21"></span>[47] R. Slansky. *Phys. Rept.* 79 (1981) 1.
- [48] W. McKay and J. Patera. Tables of Dimensions, Indices, and Branching Rules for Representations of Simple Lie Algebras. 1981.
- <span id="page-11-25"></span><span id="page-11-24"></span>[49] R. D'Auria, S. Ferrara, M.A. Lledó, and V.S. Varadarajan. J. Geom. Phys. 40 (2001) 101.
- <span id="page-11-26"></span>[50] J. Figueroa-O'Farrill. URL: <www.maths.ed.ac.uk/~jmf/Teaching/Lectures/Majorana.pdf>.
- <span id="page-11-27"></span>[51] S. Coleman and J. Mandula. Phys. Rev. 159 (1967) 1251.
- <span id="page-11-28"></span>[52] J. Polchinski. Nucl. Phys. B 231 (1984) 269.
- [53] K.G. Wilson. Phys. Rev. B 4 (1971) 3174.
- <span id="page-11-29"></span>[54] K.G. Wilson and John B. Kogut. Phys. Rept. 12 (1974) 75.
- [55] L.P. Kadanoff. Phys. Phys. Fiz. 2 (1966) 263.