Three-body decays of $\pi_2(1670)$ and $\rho_3(1690)$ via the Sill distribution

Shahriyar Jafarzade^{1,2} and Enrico Trotti¹

¹Institute of Physics, Jan Kochanowski University, ul. Universytecka 7, 25-406, Kielce, Poland,

² Institute of Radiation Problems, Ministry of Science and Education, B.Vahabzade St. 9, AZ1143, Baku, Azerbaijan

Abstract

We present the three-body decays for the resonances $\pi_2(1670)$ and $\rho_3(1690)$. We use an effective model based on the flavor symmetry and have assumed the Sill distribution for describing the lie-shape of intermediate unstable resonances. The vector-pseudoscalar meson decay channel is considered as an intermediate step to explain the decay of three-pseudoscalar meson channels.

1. INTRODUCTION

The light ground state mesons, such as $\pi_2(1670)$ and $\rho_3(1690)$ (with $J^{PC} = 2^{-+}$, 3^{--}), according to the Particle Data Group (PDG) [1], are well-established. Their total decay widths are $\Gamma_{\pi_2(1670)}^{\text{tot}} = 258^{+8}_{-9}$ MeV and $\Gamma_{\rho_3(1690)}^{\text{tot}} = 161 \pm 10$ MeV. Experimental data for the decay processes of these resonances was used in [2–5] to test the corresponding spontaneous and anomalous breaking features of QCD symmetries used in the extended Linear Sigma Model (eLSM) [6]. We list some of the vector-pseudoscalar meson decay data of these resonances in Tab. 1.

In this work, we present novel results for three-body decays of these resonances based on the effective model constructed in Refs. [2, 4]. To this end, we use the Sill distribution developed in Ref. [7] and successively used in the case of e.g. spin-2 mesons [8] and hybrid mesons in Ref. [9] (for other works that have applied the Sill distribution, see also [10–13]).

Decay process	Model $[2, 4]$	PDG [1]
$\pi_2(1670) \to ho(770) \pi$	80.6 ± 10.8	80.6 ± 10.8
$\pi_2(1670) \to \bar{K}^*(892) K + \mathbf{c.c}$	11.7 ± 1.6	10.9 ± 3.7
$\rho_3(1690) \to \omega(782) \pi$	35.8 ± 7.4	25.8 ± 9.8
$\rho_3(1690) \to \rho(770) \eta$	3.8 ± 0.8	_
$\rho_3(1690) \to \bar{K}^*(892) K + \mathbf{c.c.}$	3.4 ± 0.7	_

TABLE 1. Decay rates (MeV) of $\pi_2(1670)$ and $\rho_3(1690)$. (The first entry has been used to determine the model parameter.)

2. Effective Model

The following effective interaction term describes the decay of the pseudotensor mesons¹ $\left(P_2 = \left\{\pi_2(1670), K_2(1770), \eta_2(1870), \eta_2(1645)\right\}\right)$ into the vector $\left(V = \left\{\rho(770), K^*(892), \omega(782), \phi(1020)\right\}\right)$ and pseudoscalar $\left(P = \left\{\pi, K, \eta(547), \eta'(958)\right\}\right)$ mesons [2]:

$$\mathcal{L}_{p_2 v p} = g_{p_2 v p} \operatorname{tr} \left[P_2^{\mu \nu} \left[V_{\mu} , (\partial_{\nu} P) \right]_{-} \right] , \qquad (2.1)$$

which leads to the decay rate:

$$\Gamma_{P_2 \to V+P}(m_{p_2}, m_v, m_p) = \frac{g_{p_2vp}^2 \, |\vec{k}_{p_2,v,p}|^3}{120 \, \pi \, m_{p_2}^2} \left(5 + \frac{2 \, |\vec{k}_{p_2,v,p}|^2}{m_v^2} \right) \kappa_i \, \Theta(m_{p_2} - m_v - m_p) \,. \tag{2.2}$$

Above κ_i is a decay coefficient that takes into account flavor combinatoric and renormalization factors, extracted from the extended version of the lagrangian. The coupling constant is determined to be:

$$g_{p_2vp}^2 = (11.9 \pm 1.6).$$
 (2.3)

A similar Lagrangian describing the decay of spin-3 tensor mesons $\left(W_3 = \left\{\rho_3(1690), K_3^*(1780), \phi_3(1850), \omega_3(1670)\right\}\right)$ has the following form [4]:

$$\mathcal{L}_{w_3vp} = g_{w_3vp} \,\varepsilon^{\mu\nu\rho\sigma} \,\mathrm{tr} \Big[W_{3,\mu\alpha\beta} \left\{ (\partial_\nu V_\rho), \, (\partial^\alpha \partial^\beta \partial_\sigma P) \right\}_+ \Big] \,. \tag{2.4}$$

The tree-level decay rate is:

$$\Gamma_{W_3 \to V+P}(m_{w_3}, m_v, m_p) = g_{w_3 v p}^2 \frac{\left|\vec{k}_{w_3, v, p}\right|^7}{105} \kappa_i \Theta(m_{w_3} - m_v - m_p).$$
(2.5)

As a consequence of the parameter determination shown in Ref. [4], one has the following value for the coupling:

$$g_{w_3vp}^2 = (9.2 \pm 1.9) \cdot 10^{-16} \text{ MeV}^{-6}$$
. (2.6)

Some decay channels are listed in Tab.1 (see Ref. [4] for more details).

 $^{{}^{1}}P_{2}$ is a 3 × 3 matrix containing the nonet of listed states, see [2]. The same is true for the other quoted nonets.

3. Results for three-body decays

The PDG lists the following decay rates [1]:

$$\Gamma_{\pi_2(1670)\to\pi^{\pm}\pi^{+}\pi^{-}} = 137 \pm 11 \,\mathrm{MeV}\,, \qquad \Gamma_{\rho_3(1690)\to K\overline{K}\pi} = 6.1 \pm 2.0 \,\mathrm{MeV}\,. \tag{3.1}$$

In order to theoretically describe them, we introduce the "Sill" spectral function $d^{\text{Sill}}(x)$, which is chosen because of the following features: it is normalized even for broad states, it has a vanishing real part contribution to the loop of virtual particles, it takes into account the decay threshold, and is rather simple to use. For further details, see Ref. [7,13].

Considering the spectral function for $\rho(770)$, one has:

$$d_{\rho}^{\text{Sill}}(y) = \frac{2y}{\pi} \frac{\sqrt{y^2 - 4m_{\pi}^2} \,\tilde{\Gamma}_{\rho}}{(y^2 - m_{\rho}^2)^2 + (\sqrt{y^2 - 4m_{\pi}^2} \,\tilde{\Gamma}_{\rho})^2} \,\Theta(y - 2m_{\pi}) \,, \qquad \qquad \int_{2m_{\pi}}^{\infty} dy \, d_{\rho}^{\text{Sill}}(y) = 1 \,, \qquad (3.2)$$

where the definition of $\tilde{\Gamma}_{\rho}$ is

$$\tilde{\Gamma}_{\rho} \equiv \frac{\Gamma_{\rho \to 2\pi}^{\text{PDG}} m_{\rho}}{\sqrt{m_{\rho}^2 - 4m_{\pi}^2}} \quad , \tag{3.3}$$

with the PDG values $\Gamma_{\rho\to 2\pi}^{\text{PDG}} = 149.1 \text{ MeV}$ and $m_{\rho} = 775.11 \text{ MeV}$. Then, upon integrating over it, we obtain the following result for the three-body decay of π_2 :

$$\Gamma_{\pi_2 \to \rho \pi \to \pi \pi \pi} \simeq \int_{2m_{\pi}}^{\infty} dy \, d_{\rho}^{\text{Sill}}(y) \, \Gamma_{\pi_2 \to \rho \pi}(m_{\pi_2}, y, m_{\pi}) \simeq (73.9 \pm 9.9) \, \text{MeV} \,. \tag{3.4}$$

The Sill distribution for the vector $K^*(892)$ -meson reads:

$$d_{K^{\star}}^{\text{Sill}}(y) = \frac{2y}{\pi} \frac{\sqrt{y^2 - (m_K + m_\pi)^2} \,\tilde{\Gamma}_{K^{\star}}}{(y^2 - m_{K^{\star}}^2)^2 + (\sqrt{y^2 - (m_K + m_\pi)^2} \,\tilde{\Gamma}_{K^{\star}})^2} \,\Theta(y - m_K - m_\pi)\,,\tag{3.5}$$

where $\tilde{\Gamma}_{K^{\star}}$ is linked to the PDG values $\Gamma_{K^{\star} \to K\pi}^{\text{PDG}} = 51.4 \text{ MeV}$ and $m_{K^{\star}} = 890 \text{ MeV}$ according to:

$$\tilde{\Gamma}_{K^{\star}} \equiv \frac{\Gamma_{K^{\star} \to K\pi}^{\text{PDG}} m_{K^{\star}}}{\sqrt{m_{K^{\star}}^2 - (m_K - m_\pi)^2}} \quad . \tag{3.6}$$

The normalization

$$\int_{m_k+m_{\pi}}^{\infty} dy \, d_{K^{\star}}^{\text{Sill}}(y) = 1 \,, \tag{3.7}$$

holds also here.

In this case, the three-body decay of $\rho_3(1690)$ can be calculated as:

$$\Gamma_{\rho_3(1690)\to\bar{K}^*(892)\,K\to K\bar{K}\pi} \simeq \int_{m_k+m_\pi}^{\infty} dy \, d_{K^*}^{\text{Sill}}(y) \,\Gamma_{\rho_3\to\bar{K}^*\,K}(m_{\rho_3}, y, m_K) \simeq (3.43\pm0.70)\,\text{MeV} \,. \tag{3.8}$$

The decay channel $\Gamma_{\rho_3 \to \eta \pi \pi}$ has also been seen experimentally [14]; we estimate it as:

$$\Gamma_{\rho_3(1690)\to\rho(770)\eta\to\eta\pi\pi} \simeq \int_{2m_\pi}^\infty dy \, d_\rho^{\rm Sill}(y) \, \Gamma_{\rho_3\to\rho\eta}(m_{\rho_3}, y, m_\eta) \simeq (3.95\pm0.81) \, \text{MeV} \,. \tag{3.9}$$

By using the decay rates given in Tab.1, we present various three-body decay channels assuming the Sill distribution in Tab. 2.

Decay channel	Sill distribution	PDG
$\pi_2(1670) \to \rho(770)\pi \to \pi^{\pm}\pi^+\pi^-$	73.9 ± 9.9	137 ± 11
$\pi_2(1670) \to \overline{K}^*(892)\pi \to K\overline{K}\pi$	10.4 ± 1.4	
$ \rho_3(1690) \to \rho(770)\eta \to \eta\pi\pi $	3.88 ± 0.80	seen
$ \rho_3(1690) \to \overline{K}^*(892)\pi \to K\overline{K}\pi $	3.43 ± 0.70	6.1 ± 2.0

TABLE 2. Three body decay rates of $\pi_2(1670)$ and $\rho_3(1690)$ mesons (MeV).

4. Conclusion

In this paper, we used a hadronic model and the Sill distribution, introduced in the vector meson intermediate decays, to evaluate the three-body decays of $\pi_2(1670)$ and $\rho_3(1690)$. Results are compared to the ones listed in PDG: they agree within the error bars for $\rho_3(1690) \rightarrow K\overline{K}\pi$, while it is two times off the experimental result for $\pi_2(1670) \rightarrow 3\pi$. This implies the necessity of the consideration of multi-channel version of the Sill distribution [7] in the future.

Acknowledgement

We are thankful to Francesco Giacosa, Adrian Koenigstein, and Vanamali Shastry for useful discussions. We acknowledge financial support through the project "Development Accelerator of the Jan Kochanowski University of Kielce", co-financed by the European Union under the European Social Fund, with no. POWR.03.05.00-00-Z212 / 18. The work of Shahriyar Jafarzade is partially supported by the Polish National Science Centre (NCN) through the OPUS project 2019/33/B/ST2/00613.

References

- Particle Data Group Collaboration, R. L. Workman *et al.*, "Review of Particle Physics," *PTEP* 2022 (2022) 083C01.
- [2] A. Koenigstein and F. Giacosa, "Phenomenology of pseudotensor mesons and the pseudotensor glueball," *Eur. Phys. J. A* 52 no. 12, (2016) 356, arXiv:1608.08777 [hep-ph].
- [3] F. Giacosa, S. Jafarzade, and R. Pisarski, "Anomalous interactions for heterochiral mesons with $J^{PC} = 1^{+-}$ and 2^{-+} ," arXiv:2309.00086 [hep-ph].
- [4] S. Jafarzade, A. Koenigstein, and F. Giacosa, "Phenomenology of J^{PC} = 3⁻⁻ tensor mesons," *Phys. Rev. D* 103 no. 9, (2021) 096027, arXiv:2101.03195 [hep-ph].
- [5] V. Shastry, E. Trotti, and F. Giacosa, "Constraints imposed by the partial wave amplitudes on the decays of J=1, 2 mesons," *Phys. Rev. D* 105 no. 5, (2022) 054022, arXiv:2107.13501 [hep-ph].
- [6] D. Parganlija, P. Kovacs, G. Wolf, F. Giacosa, and D. H. Rischke, "Meson vacuum phenomenology in a three-flavor linear sigma model with (axial-)vector mesons," *Phys. Rev. D* 87 no. 1, (2013) 014011, arXiv:1208.0585 [hep-ph].
- [7] F. Giacosa, A. Okopińska, and V. Shastry, "A simple alternative to the relativistic Breit-Wigner distribution," *Eur. Phys. J. A* 57 no. 12, (2021) 336, arXiv:2106.03749 [hep-ph].
- [8] S. Jafarzade, A. Vereijken, M. Piotrowska, and F. Giacosa, "From well-known tensor mesons to yet unknown axial-tensor mesons," *Phys. Rev. D* 106 no. 3, (2022) 036008, arXiv:2203.16585 [hep-ph].

- [9] V. Shastry and F. Giacosa, "Radiative production and decays of the exotic η1'(1855) and its siblings," Nucl. Phys. A 1037 (2023) 122683, arXiv:2302.07687 [hep-ph].
- [10] Joint Physics Analysis Center Collaboration, D. Winney, A. Pilloni, V. Mathieu, A. N. Hiller Blin,
 M. Albaladejo, W. A. Smith, and A. Szczepaniak, "XYZ spectroscopy at electron-hadron facilities. II. Semi-inclusive processes with pion exchange," *Phys. Rev. D* 106 no. 9, (2022) 094009, arXiv:2209.05882 [hep-ph].
- [11] D. Yıldırım, "Spin partners of the B^(*) B^(*) resonances with a different approach than the Breit-Wigner parameterization," *Eur. Phys. J. A* 59 no. 7, (2023) 148, arXiv:2306.12403 [hep-ph].
- [12] ALICE Collaboration, S. Acharya *et al.*, "Accessing the strong interaction between Λ baryons and charged kaons with the femtoscopy technique at the LHC," *Phys. Lett. B* 845 (2023) 138145, arXiv:2305.19093 [nucl-ex].
- [13] F. Giacosa and V. Shastry, "Sill distribution: genesis and salient features," in 20th International Conference on Hadron Spectroscopy and Structure. 10, 2023. arXiv:2310.06346 [hep-ph].
- [14] S. Fukui et al., "Vector Resonances Around 1.6-GeV of the ηπ⁺π⁻ System in the π⁻p Charge Exchange Reaction at 8.95-GeV/c," Phys. Lett. B 202 (1988) 441–446.