Study of two-body doubly charmful baryonic B decays with SU(3) flavor symmetry

Yu-Kuo Hsiao^{1,*}

¹School of Physics and Information Engineering, Shanxi Normal University, Taiyuan 030031, China

(Dated: October 2, 2023)

Abstract

Within the framework of SU(3) flavor symmetry, we investigate two-body doubly charmful baryonic $B \to \mathbf{B}_c \bar{\mathbf{B}}'_c$ decays, where $\mathbf{B}_c \bar{\mathbf{B}}'_c$ represents the anti-triplet charmed dibaryon. We determine the $SU(3)_f$ amplitudes and calculate $\mathcal{B}(B^- \to \Xi_c^0 \bar{\Xi}_c^-) = (3.4^{+1.0}_{-0.9}) \times 10^{-5}$ and $\mathcal{B}(\bar{B}^0_s \to \Lambda_c^+ \bar{\Xi}_c^-) = (3.9^{+1.2}_{-1.0}) \times 10^{-5}$ induced by the single W-emission configuration. We find that the W-exchange amplitude, previously neglected in studies, needs to be taken into account. It can cause a destructive interfering effect with the W-emission amplitude, alleviating the significant discrepancy between the theoretical estimation and experimental data for $\mathcal{B}(\bar{B}^0 \to \Lambda_c^+ \bar{\Lambda}_c^-)$. To test other interfering decay channels, we calculate $\mathcal{B}(\bar{B}^0_s \to \Xi_c^{0(+)} \bar{\Xi}_c^{0(+)}) = (3.0^{+1.4}_{-1.1}) \times 10^{-4}$ and $\mathcal{B}(\bar{B}^0 \to$ $\Xi_c^0 \bar{\Xi}_c^0) = (1.5^{+0.7}_{-0.6}) \times 10^{-5}$. We estimate non-zero branching fractions for the pure W-exchange decay channels, specifically $\mathcal{B}(\bar{B}^0_s \to \Lambda_c^+ \bar{\Lambda}_c^-) = (8.1^{+1.7}_{-1.5}) \times 10^{-5}$ and $\mathcal{B}(\bar{B}^0 \to \Xi_c^+ \bar{\Xi}_c^-) = (3.0 \pm 0.6) \times 10^{-6}$. Additionally, we predict $\mathcal{B}(B_c^+ \to \Xi_c^+ \bar{\Xi}_c^0) = (2.8^{+0.9}_{-0.7}) \times 10^{-4}$ and $\mathcal{B}(B_c^+ \to \Lambda_c^+ \bar{\Xi}_c^0) = (1.6^{+0.5}_{-0.4}) \times 10^{-5}$, which are accessible to experimental facilities such as LHCb.

^{*} Email address: yukuohsiao@gmail.com

I. INTRODUCTION

The tree-level dominated two-body charmless baryonic B meson decays, $B \to \mathbf{B}\mathbf{B}'$, can proceed through the W-boson exchange (W_{ex}) , W-boson annihilation (W_{an}) , and W-boson emission (W_{em}) decay configurations. In analogy with leptonic B decay, where the W_{an} amplitude $\mathcal{M}_{\text{wan}}(B \to \ell \bar{\nu}) \propto m_{\ell} \bar{u}_{\ell} (1 + \gamma_5) v_{\bar{\nu}}$ involves a tiny lepton mass m_{ℓ} corresponding to helicity suppression [1, 2], $\mathcal{M}_{\text{wex}(\text{wan})}(B \to \mathbf{B}\mathbf{B}') \propto m_{-} \langle \mathbf{B}\mathbf{B}' | \bar{q}q' | 0 \rangle + m_{+} \langle \mathbf{B}\mathbf{B}' | \bar{q}\gamma_5 q' | 0 \rangle$ with $m_{\mp} = m_q \mp m_{q'}$ is considered to be more suppressed than $\mathcal{M}_{\text{wem}}(B \to \mathbf{B}\mathbf{B}')$ [3]. Hence, it raises the debate if one can really neglect the $W_{\text{ex}(\text{an})}$ contribution to the branching fractions [3–10].

In the study of singly charmful baryonic $B \to \mathbf{B}_c \bar{\mathbf{B}}'$ decays, the $W_{\text{ex(an)}}$ amplitude was also neglected [11, 12]. Nonetheless, it has been found that $\mathcal{M}_{\text{wex(wan)}}(B \to \mathbf{B}_c \bar{\mathbf{B}}') \propto$ $m_c \langle \mathbf{B}_c \bar{\mathbf{B}}' | \bar{c}(1+\gamma_5) q | 0 \rangle$ with $m_c \gg m_q$ can alleviate the helicity suppression [13]. This results in $\mathcal{B}_{\text{wex}}(\bar{B}_s^0 \to \Lambda_c^+ \bar{p})$ and $\mathcal{B}_{\text{wex}}(\bar{B}^0 \to \Xi_c^+ \bar{\Sigma}^-)$ being predicted to be of order 10⁻⁵, much more reachable than $\mathcal{B}(B \to \mathbf{B}\bar{\mathbf{B}}') \sim 10^{-8} - 10^{-7}$ for the test of a non-negligible $W_{\text{ex(an)}}$ contribution. However, until very recently, these observations have not been reported.

It is worth noting that two-body doubly charmful baryonic B decays, $B \to \mathbf{B}_c \mathbf{B}'_c$, have provided a possible experimental indication of a non-negligible contribution from the W_{ex} term. The measured branching fractions for $B \to \mathbf{B}_c \mathbf{\bar{B}}'_c$ are reported as follows:

$$\begin{aligned} \mathcal{B}(\bar{B}^{0} \to \Xi_{c}^{+} \bar{\Lambda}_{c}^{-}) &= (1.2 \pm 0.8) \times 10^{-3} [14], \\ \mathcal{B}(B^{-} \to \Xi_{c}^{0} \bar{\Lambda}_{c}^{-}) &= (9.5 \pm 2.3) \times 10^{-4} [14], \\ \mathcal{B}(\bar{B}^{0} \to \Lambda_{c}^{+} \bar{\Lambda}_{c}^{-}) &< 1.6 \times 10^{-5} [14, 15] \\ &= (2.2^{+2.2}_{-1.6} \pm 1.3) \times 10^{-5} [16], \\ \mathcal{B}(\bar{B}^{0}_{s} \to \Lambda_{c}^{+} \bar{\Lambda}_{c}^{-}) &< 9.9 \times 10^{-5} [14, 15]. \end{aligned}$$
(1)

Initially, it was considered that $B \to \mathbf{B}_c \mathbf{\bar{B}}'_c$ receives a single contribution from the $W_{\rm em}$ topology [16, 17]. In Eq. (1), $\mathcal{B}(\bar{B}^0 \to \Xi_c^+ \bar{\Lambda}_c^-) \simeq \mathcal{B}(\bar{B}^0 \to \Lambda_c^+ \bar{\Lambda}_c^-)$ seemly supporting this assumption. Nonetheless, it also leads to an estimation of $\mathcal{B}(\bar{B}^0 \to \Lambda_c^+ \bar{\Lambda}_c^-) \simeq$ $(V_{cd}/V_{cs})^2 (\tau_{\bar{B}^0}/\tau_{B^-}) \mathcal{B}(B^- \to \Xi_c^0 \bar{\Lambda}_c^-) = (4.7 \pm 1.1) \times 10^{-5}$ by utilizing the $\mathcal{B}(B^- \to \Xi_c^0 \bar{\Lambda}_c^-)$ value from Eq. (1). This clearly shows a significant deviation from the experimental upper limit of 1.6×10^{-5} by around 3 standard deviations. Therefore, it is reasonable to infer that the $W_{\rm ex}$ topology, overlooked in $\bar{B}^0 \to \Lambda_c^+ \bar{\Lambda}_c^-$, should be taken into account. It can




FIG. 1. Feynman diagrams of $B_q \to \mathbf{B}_c \mathbf{\bar{B}}'_c$ and $B_c^+ \to \mathbf{B}_c \mathbf{\bar{B}}'_c$ decays.

cause a destructive interfering effect, thus reducing the overestimated branching fraction. Additionally, the W_{ex} topology can induce a non-zero $\mathcal{B}(\bar{B}^0_s \to \Lambda_c^+ \bar{\Lambda}_c^-)$, warranting further examination.

For clarification, a careful study of $B \to \mathbf{B}_c \bar{\mathbf{B}}'_c$ is necessary. The SU(3) flavor symmetry can be a useful theoretical tool [3, 8, 18–31], allowing us to parameterize the amplitudes without involving the complexity of model calculations. Hence, we propose using the $SU(3)_f$ approach to specifically explore the W_{ex} and W_{em} contributions to $B \to \mathbf{B}_c \bar{\mathbf{B}}'_c$. The first observation of $B_c^+ \to J/\Psi p\bar{p}\pi^+$ by LHCb [32] indicates a potential test-bed for the baryonic phenomena in B_c^+ decays, such as the branching fraction [33–41], direct CP asymmetry [42, 43], triple product asymmetry [44], angular distribution [45], and exotic states [46–48] as studied in baryonic B decays. Therefore, we will estimate $\mathcal{B}(B_c^+ \to \mathbf{B}_c \bar{\mathbf{B}}'_c)$ to initiate a theoretical investigation.

II. FORMALISM

To study the two-body doubly charmful baryonic $B_{(c)} \to \mathbf{B}_c \bar{\mathbf{B}}'_c$ decays with B_c denoting $B_c^+(b\bar{c})$, the quark-level effective Hamiltonians for the $b \to c\bar{q}q'$ weak transitions are required, given by [49, 50]

$$\mathcal{H}_{eff}^{b \to c\bar{q}q'} = \frac{G_F}{\sqrt{2}} V_{cb} V_{qq'}^* \left[c_1(\bar{q}'q)(\bar{c}b) + c_2(\bar{q}'_\beta q_\alpha)(\bar{c}_\alpha b_\beta) \right], \tag{2}$$

where G_F is the Fermi constant, V_{cb} and $V_{qq'}$ with q = (u, c) and q' = (s, d) the Cabibbo-Kobayashi-Maskawa (CKM) matrix elements. In Eq. (2), we define $(\bar{q}_1q_2) = \bar{q}_1\gamma_{\mu}(1-\gamma_5)q_2$, and the subscripts (α, β) denote the color indices; moreover, $c_{1,2}$ are the scale (μ) -dependent Wilson coefficients with $\mu = m_b$ for the *b* decays. In the $SU(3)_f$ representation, $\mathcal{H}_{eff}^{b \to c\bar{c}q'}$ and $\mathcal{H}_{eff}^{b \to c\bar{u}q'}$ by omitting Lorentz structure are reduced as H^i and H^i_j , respectively, where *i* and *j* run from 1 to 3 to represent the flavor indices. Explicitly, the nonzero entries are given by [27]

$$H_1^2 = \lambda_{ud} , \ H_1^3 = \lambda_{us} , \ H^2 = \lambda_{cd} , \ H^3 = \lambda_{cs} ,$$
(3)

with $\lambda_{qq'} \equiv V_{cb}V_{qq'}^*$. Accordingly, we present the *B* meson and \mathbf{B}_c baryon in the $SU(3)_f$ forms:

$$B(B_{i}) = (B^{-}, \bar{B}^{0}, \bar{B}^{0}_{s}),$$

$$B_{c}(\mathbf{B}^{ij}_{c}) = \begin{pmatrix} 0 & \Lambda^{+}_{c} & \Xi^{+}_{c} \\ -\Lambda^{+}_{c} & 0 & \Xi^{0}_{c} \\ -\Xi^{+}_{c} & -\Xi^{0}_{c} & 0 \end{pmatrix},$$
(4)

whereas B_c is a singlet. By connecting the flavor indices of the initial state to those of the effective Hamiltonian and final states, the $SU(3)_f$ approach yields the amplitudes to be

$$\mathcal{M}(B \to \mathbf{B}_c \bar{\mathbf{B}}'_c) = e B_i H^i \mathbf{B}_{c \ jk} \bar{\mathbf{B}}'_c{}^{jk} + c' B_i H^j \mathbf{B}_{c \ jk} \bar{\mathbf{B}}'_c{}^{ik} ,$$

$$\mathcal{M}(B_c \to \mathbf{B}_c \bar{\mathbf{B}}'_c) = \bar{c}' H^i_j \mathbf{B}_{c \ ik} \bar{\mathbf{B}}'_c{}^{jk} , \qquad (5)$$

where the parameter e and $c'(\bar{c}')$ correspond to the W_{ex} and W_{em} configurations in Fig. 1a and Fig. 1b(c), respectively. For a later numerical analysis, we use the equation [14]:

$$\mathcal{B}(B_{(c)} \to \mathbf{B}_{c}\bar{\mathbf{B}}_{c}') = \frac{G_{F}^{2}|\vec{p}_{cm}|\tau_{B_{(c)}}}{16\pi m_{B_{(c)}}^{2}}|\mathcal{M}(B_{(c)} \to \mathbf{B}_{c}\bar{\mathbf{B}}_{c}')|^{2},$$
$$|\vec{p}_{cm}| = \frac{\sqrt{(m_{B_{(c)}}^{2} - M_{+}^{2})(m_{B_{(c)}}^{2} - M_{-}^{2})}}{2m_{B_{(c)}}},$$
(6)

to compute the branching fractions, where $M_{\pm} \equiv m_{\mathbf{B}_c} \pm m_{\mathbf{\bar{B}}'_c}$, \vec{p}_{cm} is the three-momentum of the \mathbf{B}_c baryon in the $B_{(c)}$ meson rest frame, and $\tau_{B_{(c)}}$ stands for the $B_{(c)}$ lifetime. The amplitude $\mathcal{M}(B_{(c)} \to \mathbf{B}_c \mathbf{\bar{B}}'_c)$ can be found in Table I.

III. NUMERICAL RESULTS

In the numerical analysis, the CKM matrix elements are adopted from PDG [14]:

$$(V_{cb}, V_{cs}, V_{ud}, V_{us}, V_{cd}) = (A\lambda^2, 1 - \lambda^2/2, 1 - \lambda^2/2, \lambda, -\lambda),$$
(7)

where A = 0.826 and $\lambda = 0.225$ in the Wolfenstein parameterization. In Eq. (5), the parameters e and c' are complex numbers, which we present as

$$|c'|, |e|e^{i\delta_e},\tag{8}$$

with δ_e a relative phase. By using the experimental data in Table II, we solve the parameters as

$$|c'| = (1.29 \pm 0.18) \text{ GeV}^3, \ |e| = (0.19 \pm 0.03) \text{ GeV}^3, \ \delta_e = 180^{\circ}.$$
 (9)

Moreover, we assume $\bar{c}' = c'$ due to the similarity of the Feynman diagrams in Figs. 1b and 1c. Subsequently, we calculate the branching fractions as provided in Table II using the determination in Eq. (9).

IV. DISCUSSION AND CONCLUSIONS

The $SU(3)_f$ approach enables us to explore all possible $B \to \mathbf{B}_c \mathbf{B}'_c$ decays, as summarized in Table I. Furthermore, it helps in deriving constraints on $SU(3)_f$ relations, facilitating the decomposition of amplitudes into e and c' terms. These terms parameterize the W_{ex} and W_{em} topologies depicted in Fig.1a and Fig.1b(c), respectively.

In $b \to c\bar{c}s$ induced decays, the $SU(3)_f$ symmetry unequivocally establishes that both $\bar{B}^0 \to \Xi_c^+ \bar{\Lambda}_c^-$ and $B^- \to \Xi_c^0 \bar{\Lambda}_c^-$ solely proceed through the $W_{\rm em}$ topology, supporting earlier considerations [16, 17]. The nearly identical branching fractions, as shown in Eq. (1), also provide consistent evidence. In our latest findings, we unveil additional insights. For

Decay modes	Amplitudes	Decay modes	Amplitudes
$\bar{B}^0 \to \Xi_c^+ \bar{\Lambda}_c^-$	$-\lambda_{cs}c'$	$\bar{B}^0\to \Xi^0_c \bar{\Xi}^0_c$	$-\lambda_{cd}(2e+c')$
$B^-\to \Xi^0_c \bar\Lambda^c$	$\lambda_{cs}c'$	$\bar{B}^0\to \Xi_c^+\bar{\Xi}_c^-$	$-\lambda_{cd}(2e)$
$\bar{B}^0_s\to \Xi^0_c \bar{\Xi}^0_c$	$-\lambda_{cs}(2e+c')$	$\bar{B}^0\to\Lambda_c^+\bar{\Lambda}_c^-$	$-\lambda_{cd}(2e+c')$
$\bar{B}^0_s\to \Xi_c^+\bar{\Xi}_c^-$	$-\lambda_{cs}(2e+c')$	$B^-\to \Xi_c^0 \bar \Xi_c^-$	$-\lambda_{cd}c'$
$\bar{B}^0_s\to\Lambda^+_c\bar{\Lambda}^c$	$-\lambda_{cs}(2e)$	$\bar{B}^0_s\to\Lambda^+_c\bar{\Xi}^c$	$-\lambda_{cd}c'$
$B_c^+ \to \Xi_c^+ \bar{\Xi}_c^0$	$-\lambda_{ud}\bar{c}'$	$B_c^+ \to \Lambda_c^+ \bar{\Xi}_c^0$	$\lambda_{us} \bar{c}'$

TABLE I. Amplitudes of $B_{(c)} \to \mathbf{B}_c \mathbf{\bar{B}}'_c$ with the $SU(3)_f$ parameters e and $c'(\vec{c}')$.

 $\bar{B}^0_s \to \Xi^{0(+)}_c \bar{\Xi}^{0(-)}_c$, the interference of the $W_{\rm ex}$ amplitude with the $W_{\rm em}$ amplitude adds a new contribution to the decay process. Furthermore, $\bar{B}^0_s \to \Lambda^+_c \bar{\Lambda}^-_c$ represents a pure $W_{\rm ex}$ decay, offering a clear and distinct case for experiments to clarify if the $W_{\rm ex}$ contribution can be neglected.

The $b \to c\bar{c}d$ induced decays with $|V_{cd}/V_{cs}| \simeq 0.05$ are more suppressed. Unlike $\mathcal{M}(\bar{B}^0_s \to \Xi^0_c \bar{\Xi}^0_c) = \mathcal{M}(\bar{B}^0_s \to \Xi^+_c \bar{\Xi}^-_c), \ \bar{B}^0 \to \Xi^+_c \bar{\Xi}^-_c$ as a pure W_{ex} decay does not share an isospin relation with $\bar{B}^0 \to \Xi^0_c \bar{\Xi}^0_c$, which is against our naive expectation. Instead, the equality relation arises from $\mathcal{M}(\bar{B}^0 \to \Lambda^+_c \bar{\Lambda}^-_c) = \mathcal{M}(\bar{B}^0 \to \Xi^0_c \bar{\Xi}^0_c)$. Moreover, an interesting triangle relation exists:

$$\mathcal{M}(\bar{B}^0 \to \Xi_c^+ \bar{\Xi}_c^-) + \mathcal{M}(B^- \to \Xi_c^0 \bar{\Xi}_c^-) = \mathcal{M}(\bar{B}^0 \to \Xi_c^0 \bar{\Xi}_c^0) \,. \tag{10}$$

If the W_{ex} contribution is negligible, the relation is simplified to $\mathcal{M}(B^- \to \Xi_c^0 \bar{\Xi}_c^-) \simeq \mathcal{M}(\bar{B}^0 \to \Xi_c^0 \bar{\Xi}_c^0)$, resulting in nearly equal branching fractions.

It can be challenging to compute the W_{ex} and W_{em} amplitudes. For example, the factorization approach derives the W_{ex} amplitude as $\mathcal{M}_{\text{wex}} \propto f_B q^{\mu} \langle \mathbf{B}_c \bar{\mathbf{B}}'_c | \bar{c} \gamma_{\mu} (1 - \gamma_5) c | 0 \rangle^{-1}$, where

decay channel	this work	experimental data
$10^4 \mathcal{B}(\bar{B}^0 \to \Xi_c^+ \bar{\Lambda}_c^-)$	$7.2^{+2.1}_{-1.9}$	12 ± 8 [14]
$10^4 \mathcal{B}(B^- \to \Xi_c^0 \bar{\Lambda}_c^-)$	$7.8^{+2.3}_{-2.0}$	9.5 ± 2.3 [14]
$10^4 \mathcal{B}(\bar{B}^0_s \to \Xi^0_c \bar{\Xi}^0_c)$	$3.0^{+1.4}_{-1.1}$	
$10^4 \mathcal{B}(\bar{B}^0_s \to \Xi_c^+ \bar{\Xi}_c^-)$	$3.0^{+1.4}_{-1.1}$	
$10^5 \mathcal{B}(\bar{B}^0_s \to \Lambda^+_c \bar{\Lambda}^c)$	$8.1^{+1.7}_{-1.5}$	$< 9.9 \ [14, \ 15]$
$10^4 \mathcal{B}(B_c^+ \to \Xi_c^+ \bar{\Xi}_c^0)$	$2.8^{+0.9}_{-0.7}$	
$10^5 \mathcal{B}(\bar{B}^0 \to \Xi_c^0 \bar{\Xi}_c^0)$	$1.5_{-0.6}^{+0.7}$	
$10^6 \mathcal{B}(\bar{B}^0 \to \Xi_c^+ \bar{\Xi}_c^-)$	3.0 ± 0.6	
$10^5 \mathcal{B}(\bar{B}^0 \to \Lambda_c^+ \bar{\Lambda}_c^-)$	$2.1^{+1.0}_{-0.8}$	$< 1.6 [14, 15] (2.2^{+2.6}_{-2.1} [16])$
$10^5 \mathcal{B}(B^- \to \Xi_c^0 \bar{\Xi}_c^-)$	$3.4^{+1.0}_{-0.9}$	
$10^5 \mathcal{B}(\bar{B}^0_s \to \Lambda^+_c \bar{\Xi}^c)$	$3.9^{+1.2}_{-1.0}$	
$10^5 \mathcal{B}(B_c^+ \to \Lambda_c^+ \bar{\Xi}_c^0)$	$1.6^{+0.5}_{-0.4}$	

TABLE II. Branching fractions of $B_{(c)} \to \mathbf{B}_c \mathbf{\overline{B}}'_c$ decays.

¹ Please consult the similar deviation for $\mathcal{M}(B \to \mathbf{B}_c \bar{\mathbf{B}})$ in Ref. [13].

 f_B is the B meson decay constant, q^{μ} the momentum transfer, and the matrix elements present the vacuum (0) to $\mathbf{B}_c \bar{\mathbf{B}}'_c$ production. As information on the $0 \to \mathbf{B}_c \bar{\mathbf{B}}'_c$ production is lacking, a model calculation is currently unavailable. For a calculation on \mathcal{M}_{wem} , one proposes a meson propagator to provide an additional quark pair, resulting in the branching fractions to be a few times 10^{-3} for $\bar{B}^0 \to \Xi_c^+ \bar{\Lambda}_c^-$ and $B^- \to \Xi_c^0 \bar{\Lambda}_c^-$ [17]. Additionally, a theoretical attempt incorporating final state interactions yields $\mathcal{B} \simeq \mathcal{O}(10^{-3})$ [51]. It appears that the above approaches may overestimate the W_{em} contribution.

Without invoking the model calculations, we determine e and c' with the experimental data based on the $SU(3)_f$ symmetry. Explicitly, we use the experimental results for $\mathcal{B}(\bar{B}^0 \to \Xi_c^+ \bar{\Lambda}_c^-)$ and $\mathcal{B}(B^- \to \Xi_c^0 \bar{\Lambda}_c^-)$ to fit |c'|. On the other hand, the experimental data have not been sufficient and accurate enough to simultaneously determine |e| and the relative phase δ_e , as indicted in Table II. For a practical determination, we fix $\delta_e = 180^\circ$ to cause a maximum destructive interference, where the fitted |c'| value has been used. As a consequence, the experimental upper bounds of $\mathcal{B}(\bar{B}^0_s \to \Lambda_c^+ \bar{\Lambda}_c^-)$ and $\mathcal{B}(\bar{B}^0 \to \Lambda_c^+ \bar{\Lambda}_c^-)$ can sandwich an allowed range for |e|, as given in Eq. (9).

As the numerical results, we obtain $\mathcal{B}(\bar{B}^0 \to \Xi_c^+ \bar{\Lambda}_c^-) = (7.2^{+2.1}_{-1.9}) \times 10^{-4}$ and $\mathcal{B}(B^- \to \Xi_c^0 \bar{\Lambda}_c^-) = (7.8^{+2.3}_{-2.0}) \times 10^{-4}$ utilizing $|c'| = (1.29 \pm 0.18)$ GeV³, in agreement with the experimental inputs. This demonstrates that c' can estimate the $W_{\rm em}$ contribution. Thus, we predict

$$\mathcal{B}(B^{-} \to \Xi_{c}^{0} \bar{\Xi}_{c}^{-}) = (3.4^{+1.0}_{-0.9}) \times 10^{-5} ,$$

$$\mathcal{B}(\bar{B}_{s}^{0} \to \Lambda_{c}^{+} \bar{\Xi}_{c}^{-}) = (3.9^{+1.2}_{-1.0}) \times 10^{-5} , \qquad (11)$$

which are also contributed by the single $W_{\rm em}$ amplitude, promising to be measured by experimental facilities such as LHCb.

The previous studies have assumed that $\bar{B}^0 \to \Lambda_c^+ \bar{\Lambda}_c^-$ receives the single $W_{\rm em}$ contribution [16, 17]. Consequently, the estimated branching fraction $\mathcal{B}(\bar{B}^0 \to \Lambda_c^+ \bar{\Lambda}_c^-) \simeq 5 \times 10^{-5}$ mentioned in the introduction significantly exceeds the experimental upper bound. In Table I, since $\mathcal{M}(\bar{B}^0 \to \Lambda_c^+ \bar{\Lambda}_c^-) = -\lambda_{cd}(2e + c')$ is found to include the $SU(3)_f$ parameter e, it suggests a non-negligible $W_{\rm ex}$ amplitude. By newly incorporating e, a destructive interference with c' could occur, effectively reducing the branching fraction. In fact, we estimate $|e| \simeq 0.2 \text{ GeV}^3$, and obtain $\mathcal{B}(\bar{B}^0 \to \Lambda_c^+ \bar{\Lambda}_c^-) = (2.1^{+1.0}_{-0.8}) \times 10^{-5}$, thus alleviating the discrepancy. To carefully test the W_{ex} contribution, we predict the branching fractions of the other interfering decay channels:

$$\mathcal{B}(\bar{B}^0_s \to \Xi^{0(+)}_c \bar{\Xi}^{0(+)}_c) = (3.0^{+1.4}_{-1.1}) \times 10^{-4} ,$$

$$\mathcal{B}(\bar{B}^0 \to \Xi^0_c \bar{\Xi}^0_c) = (1.5^{+0.7}_{-0.6}) \times 10^{-5} .$$
(12)

When |e| = 0, $\mathcal{B}(\bar{B}^0_s \to \Xi^{0(+)}_c \bar{\Xi}^{0(+)}_c)$ would be enhanced to $(6.3^{+1.9}_{-1.6}) \times 10^{-4}$; moreover, $\mathcal{B}(\bar{B}^0 \to \Xi^0_c \bar{\Xi}^0_c)$ would be enhanced to $(3.1^{+0.9}_{-0.8}) \times 10^{-5}$, making it close to $\mathcal{B}(B^- \to \Xi^0_c \bar{\Xi}^-_c)$, in accordance with the description for the triangle relation in Eq. (10). We also anticipate non-zero branching fractions of the pure $W_{\rm ex}$ decays, given by

$$\mathcal{B}(\bar{B}^0_s \to \Lambda^+_c \bar{\Lambda}^-_c) = (8.1^{+1.7}_{-1.5}) \times 10^{-5} ,$$

$$\mathcal{B}(\bar{B}^0 \to \Xi^+_c \bar{\Xi}^-_c) = (3.0 \pm 0.6) \times 10^{-6} , \qquad (13)$$

which serve to test the W-exchange mechanism in the $B \to \mathbf{B}_c \mathbf{B}'_c$ decays.

To initiate a theoretical investigation of baryonic B_c^+ decays, we derive the amplitudes for $B_c \to \mathbf{B}_c \mathbf{\bar{B}}'_c$ using the $SU(3)_f$ symmetry. This results in two possible decay channels: $B_c^+ \to \Xi_c^+ \overline{\Xi}_c^0$ and $B_c^+ \to \Lambda_c^+ \overline{\Xi}_c^0$, with \vec{c}' representing the sole contribution from the $W_{\rm em}$ term, as given in Table I. Since both $B_c \to \mathbf{B}_c \mathbf{\bar{B}}'_c$ and $B \to \mathbf{B}_c \mathbf{\bar{B}}'_c$ involve $c\bar{c}$ in the $\mathbf{B}_c \mathbf{\bar{B}}'_c$ formation, it is reasonable to assume that QCD effects cannot distinguish between the topology in Fig. 1b and the one in Fig. 1c in the hadronization process. Hence, we assume $\vec{c}' = c'$, and predict the following branching fractions:

$$\mathcal{B}(B_c^+ \to \Xi_c^+ \bar{\Xi}_c^0) = (2.8^{+0.9}_{-0.7}) \times 10^{-4} ,$$

$$\mathcal{B}(B_c^+ \to \Lambda_c^+ \bar{\Xi}_c^0) = (1.6^{+0.5}_{-0.4}) \times 10^{-5} , \qquad (14)$$

which can be probed by the LHCb experiment.

In summary, we have explored the two-body doubly charmful baryonic $B \to \mathbf{B}_c \mathbf{\bar{B}}'_c$ decays. Here, the W_{ex} and W_{em} amplitudes have been parametrized as e and c', respectively, using the $SU(3)_f$ approach. With the determination of the $SU(3)_f$ parameters, we have calculated the branching fractions $\mathcal{B}(B^- \to \Xi_c^0 \bar{\Xi}_c^-) = (3.4^{+1.0}_{-0.9}) \times 10^{-5}$ and $\mathcal{B}(\bar{B}^0_s \to \Lambda_c^+ \bar{\Xi}_c^-) = (3.9^{+1.2}_{-1.0}) \times 10^{-5}$. Considering that the single W_{em} contribution to $\bar{B}^0 \to \Lambda_c^+ \bar{\Lambda}_c^-$ has caused the branching fraction to significantly exceed the experimental upper bound, we have added the W_{ex} amplitude (or the $SU(3)_f$ parameter e), overlooked in previous studies, to account for a destructive interfering effect. Subsequently, we have alleviated the discrepancy. To further test the interfering decay channels, we have predicted $\mathcal{B}(\bar{B}^0_s \to \Xi^{0(+)}_c \bar{\Xi}^{0(+)}_c) = (3.0^{+1.4}_{-1.1}) \times 10^{-4}$ and $\mathcal{B}(\bar{B}^0 \to \Xi^0_c \bar{\Xi}^0_c) = (1.5^{+0.7}_{-0.6}) \times 10^{-5}$. For the pure W_{ex} decay channels, we have expected non-zero branching fractions, such as $\mathcal{B}(\bar{B}^0_s \to \Lambda^+_c \bar{\Lambda}^-_c) = (8.1^{+1.7}_{-1.5}) \times 10^{-5}$ and $\mathcal{B}(\bar{B}^0 \to \Xi^+_c \bar{\Xi}^-_c) = (3.0 \pm 0.6) \times 10^{-6}$, promising to be observed in near-future measurements. Additionally, we have predicted $\mathcal{B}(B^+_c \to \Xi^+_c \bar{\Xi}^0_c) =$ $(2.8^{+0.9}_{-0.7}) \times 10^{-4}$ and $\mathcal{B}(B^+_c \to \Lambda^+_c \bar{\Xi}^0_c) = (1.6^{+0.5}_{-0.4}) \times 10^{-5}$, which are accessible to the LHCb experiment.

ACKNOWLEDGMENTS

This work was supported by NSFC (Grants No. 11675030 and No. 12175128).

- [1] A. J. Bevan et al. [BaBar and Belle Collaborations], Eur. Phys. J. C 74, 3026 (2014).
- [2] W. S. Hou, M. Kohda, T. Modak and G. G. Wong, Phys. Lett. B 800, 135105 (2020).
- [3] C. K. Chua, Phys. Rev. D **106**, 036015 (2022).
- [4] V. L. Chernyak and I. R. Zhitnitsky, Nucl. Phys. B 345, 137 (1990).
- [5] P. Ball and H. G. Dosch, Z. Phys. C **51**, 445 (1991).
- [6] C. H. V. Chang and W. S. Hou, Eur. Phys. J. C 23, 691 (2002).
- [7] H. Y. Cheng and K. C. Yang, Phys. Rev. D 66, 014020 (2002).
- [8] C. K. Chua, Phys. Rev. D 89, 056003 (2014).
- [9] Y. K. Hsiao and C. Q. Geng, Phys. Rev. D **91**, 077501 (2015).
- [10] X. Huang, Y. K. Hsiao, J. Wang and L. Sun, Adv. High Energy Phys. 2022, 4343824 (2022).
- [11] X. G. He, T. Li, X. Q. Li and Y.M. Wang, Phys. Rev. D 75, 034011 (2007).
- [12] H. Y. Cheng and K. C. Yang, Phys. Rev. D 67, 034008 (2003).
- [13] Y. K. Hsiao, S. Y. Tsai, C. C. Lih and E. Rodrigues, JHEP 04, 035 (2020).
- [14] R. L. Workman et al. [Particle Data Group], PTEP 2022, 083C01 (2022).
- [15] R. Aaij *et al.* [LHCb], Phys. Rev. Lett. **112**, 202001 (2014).
- [16] Y. Uchida *et al.* [Belle], Phys. Rev. D 77, 051101 (2008).
- [17] H. Y. Cheng, C. K. Chua and Y. K. Hsiao, Phys. Rev. D 79, 114004 (2009).
- [18] M. J. Savage and M. B. Wise, Nucl. Phys. B 326, 15 (1989).

- [19] M. J. Savage and M. B. Wise, Phys. Rev. D 39, 3346 (1989); [erratum: Phys. Rev. D 40, 3127 (1989)].
- [20] M. J. Savage and R. P. Springer, Phys. Rev. D 42, 1527 (1990).
- [21] Y. Kohara, Phys. Rev. D 43, 2429 (1991).
- [22] X. G. He, Y. K. Hsiao, J. Q. Shi, Y. L. Wu and Y. F. Zhou, Phys. Rev. D 64, 034002 (2001).
- [23] H. K. Fu, X. G. He and Y. K. Hsiao, Phys. Rev. D 69, 074002 (2004).
- [24] Y. K. Hsiao, C. F. Chang and X. G. He, Phys. Rev. D 93, 114002 (2016).
- [25] H. J. Zhao, Y. L. Wang, Y. K. Hsiao and Y. Yu, JHEP 02, 165 (2020).
- [26] Y. K. Hsiao, Y. Yao and H. J. Zhao, Phys. Lett. B **792**, 35 (2019).
- [27] J. Pan, Y. K. Hsiao, J. Sun and X. G. He, Phys. Rev. D 102, 056005 (2020).
- [28] Y. K. Hsiao, Y. L. Wang and H. J. Zhao, JHEP 09, 035 (2022).
- [29] Z. P. Xing, X. G. He, F. Huang and C. Yang, Phys. Rev. D 108, 053004 (2023).
- [30] F. Huang, Z. P. Xing and X. G. He, JHEP 03, 143 (2022); [erratum: JHEP 09, 087 (2022)].
- [31] H. Zhong, F. Xu, Q. Wen and Y. Gu, JHEP **02**, 235 (2023).
- [32] R. Aaij *et al.* [LHCb], Phys. Rev. Lett. **113**, 152003 (2014).
- [33] W. S. Hou and A. Soni, Phys. Rev. Lett. 86, 4247 (2001).
- [34] C. K. Chua, W. S. Hou and S. Y. Tsai, Phys. Rev. D 65, 034003 (2002).
- [35] C. K. Chua, W. S. Hou and S. Y. Tsai, Phys. Rev. D 66, 054004 (2002).
- [36] C. K. Chua and W. S. Hou, Eur. Phys. J. C 29, 27 (2003).
- [37] C. Q. Geng and Y. K. Hsiao, Phys. Lett. B 619, 305 (2005).
- [38] C. Q. Geng, Y. K. Hsiao and J. N. Ng, Phys. Rev. D 75, 094013 (2007).
- [39] Y. K. Hsiao and C. Q. Geng, Phys. Lett. B 770, 348 (2017).
- [40] Y. K. Hsiao, Phys. Lett. B 845, 138158 (2023).
- [41] Y. K. Hsiao, Eur. Phys. J. C 83, 300 (2023).
- [42] C. Q. Geng, Y. K. Hsiao and J. N. Ng, Phys. Rev. Lett. 98, 011801 (2007).
- [43] Y. K. Hsiao, S. Y. Tsai and E. Rodrigues, Eur. Phys. J. C 80, 565 (2020).
- [44] C. Q. Geng and Y. K. Hsiao, Phys. Rev. D 72, 037901 (2005).
- [45] X. Huang, Y. K. Hsiao, J. Wang and L. Sun, Phys. Rev. D 105, 076016 (2022).
- [46] Y. K. Hsiao and C. Q. Geng, Phys. Lett. B 727, 168 (2013).
- [47] Y. K. Hsiao and C. Q. Geng, Phys. Lett. B **751**, 572 (2015).
- [48] Y. K. Hsiao and C. Q. Geng, Eur. Phys. J. C 75, 101 (2015).

- [49] G. Buchalla, A. J. Buras and M. E. Lautenbacher, Rev. Mod. Phys. 68, 1125 (1996).
- [50] A. J. Buras, hep-ph/9806471.
- [51] C. H. Chen, Phys. Lett. B 638, 214 (2006).