

# $B \rightarrow K^* M_X$ vs $B \rightarrow K M_X$ as a probe of a scalar-mediator dark matter scenario

Alexander Berezhnoy<sup>a</sup> and Dmitri Melikhov<sup>a,b,c</sup>

<sup>a</sup>*D. V. Skobeltsyn Institute of Nuclear Physics, M. V. Lomonosov Moscow State University, 119991, Moscow, Russia*

<sup>b</sup>*Joint Institute for Nuclear Research, 141980 Dubna, Russia*

<sup>c</sup>*Faculty of Physics, University of Vienna, Boltzmannngasse 5, A-1090 Vienna, Austria*

(Dated: November 7, 2023)

Recently, Belle II reported the observation of the decay  $B \rightarrow K M_X$ ,  $M_X$  the missing mass, with the branching ratio much exceeding  $\mathcal{B}(B \rightarrow K \nu \bar{\nu})$  which is the only Standard Model (SM) process contributing to this reaction. If confirmed, this might be an indication of new nonSM particles produced in this decay. One of the possible explanations of the observed effect could be light dark-matter (DM) particles produced via a scalar mediator field. We give simple arguments, that a combined analysis of the  $B \rightarrow K M_X$  and  $B \rightarrow K^* M_X$  reactions would be a clean probe of the scalar mediator scenario: (i) making use of an observed value  $\mathcal{B}(B \rightarrow K M_X) \simeq 5.4 \mathcal{B}(B \rightarrow K \nu \bar{\nu})_{\text{SM}}$  and (ii) assuming that the effect is due to the light dark matter coupling to the top quark via a scalar mediator field, one finds an upper limit  $\mathcal{B}(B \rightarrow K^* M_X) < 2.8 \mathcal{B}(B \rightarrow K^* \nu \bar{\nu})_{\text{SM}}$ . Within the discussed scenario, this upper limit does not depend on the mass of the scalar mediator nor on the specific details of the unobserved dark-matter particles in the final state.

1. *Introduction.* A recent Belle II observation [1] of  $B \rightarrow K M_X$  (for which also the notation  $B \rightarrow K \cancel{E}$ ,  $\cancel{E}$  the missing energy, is used) at the level much exceeding the Standard model (SM) prediction for  $B \rightarrow K \nu \bar{\nu}$ ,

$$\mathcal{B}(B^+ \rightarrow K^+ M_X) = (2.4 \pm 0.7) \times 10^{-5} \simeq (5.4 \pm 1.5) \mathcal{B}(B^+ \rightarrow K^+ \nu \bar{\nu})_{\text{SM}}, \quad (1)$$

opened a window for immediate discussions of possible new physics effects capable to explain this result (see e.g. recent publications [2–8]). One of the discussed options is the decay into Dark Matter (DM) particles [6] with multiple scenarios for the content of these particles and the possible mediators. The goal of this short communication is to remark that combining the present Belle II result for the  $B \rightarrow K M_X$  with the hypothesis of a DM origin of the enhancement of  $\Gamma(B \rightarrow K M_X)$  coupled to the SM particles via a scalar mediator field leads to interesting and rather robust constraints independent of further details of the DM model.

2. *The  $B \rightarrow (K, K^*) \bar{\chi} \chi$  decays via scalar mediator  $\phi$ .* As an example we consider a simple but rather general Lagrangian describing the interaction of DM particles  $\chi$  with the top quark via a scalar mediator field  $\phi$  [9, 10]:

$$\mathcal{L}_{\text{int}} = -\frac{y m_t}{v} \phi \bar{t} t - \kappa \phi \bar{\chi} \chi, \quad (2)$$

The corresponding effective Lagrangian describing the flavour-changing neutral current (FCNC)  $b \rightarrow s \phi$  vertex has the form [9, 10]

$$\mathcal{L}_{b \rightarrow s \phi} = g_{b \rightarrow s \phi} \phi \bar{s}_L b_R + \text{h.c.}, \quad g_{b \rightarrow s \phi} = \frac{y m_b}{v} \frac{3\sqrt{2} G_F m_q^2 V_{qs}^* V_{qb}}{16\pi^2} \quad (3)$$

This interaction leads to the following amplitudes of the  $B \rightarrow (K, K^*) \bar{\chi} \chi$  decay via the  $\phi$  mediator:

$$\begin{aligned} A(B(p) \rightarrow K(p-q) \bar{\chi}(k) \chi(q-k)) &= -i \langle K(p-q) \bar{\chi}(k) \chi(q-k) | L_{b \rightarrow s \phi} | B(p) \rangle \\ &= \langle \bar{\chi} \chi | \bar{\chi} \chi | 0 \rangle \kappa \frac{1}{M_\phi^2 - q^2} g_{b \rightarrow s \phi} \langle K(p-q) | \bar{s}_L b_R | B(p) \rangle \end{aligned} \quad (4)$$

$$\begin{aligned} A(B(p) \rightarrow K^*(p-q) \bar{\chi}(k) \chi(q-k)) &= -i \langle K^*(p-q) \bar{\chi}(k) \chi(q-k) | L_{b \rightarrow s \phi} | B(p) \rangle \\ &= \langle \bar{\chi} \chi | \bar{\chi} \chi | 0 \rangle \kappa \frac{1}{M_\phi^2 - q^2} g_{b \rightarrow s \phi} \langle K^*(p-q) | \bar{s}_L b_R | B(p) \rangle, \end{aligned} \quad (5)$$

where

$$\langle \bar{\chi} \chi | \bar{\chi} \chi | 0 \rangle \equiv \langle \bar{\chi}(k) \chi(q-k) | \bar{\chi}(0) \chi(0) | 0 \rangle, \quad (6)$$

and  $q$  is the momentum of the outgoing  $\bar{\chi} \chi$  pair of unobserved DM particles,  $M_X^2 \equiv q^2$  is the missing mass squared. Using these amplitudes, one can calculate  $d\Gamma(B \rightarrow (K, K^*) \bar{\chi} \chi) / dq^2$ , where the summation over polarizations of the DM particles  $\chi$  and the integration over their phase space are performed leading to

$$\sum_{\chi \text{ polar}} |\langle \bar{\chi}(k_1) \chi(k_2) | \bar{\chi}(0) \chi(0) | 0 \rangle|^2 \delta(m_\chi^2 - k_1^2) \delta(m_\chi^2 - k_2^2) \theta(k_{10}) \theta(k_{20}) \delta(q - k_1 - k_2) dk_1 dk_2 = \Pi_\chi(q^2). \quad (7)$$

We are interested in the ratio of the differential distributions in two reactions proceeding via the scalar mediator  $\phi$ :

$$R_{K^*/K}^{(\phi)}(q^2) = \frac{d\Gamma(B \rightarrow K^*\bar{\chi}\chi)/dq^2}{d\Gamma(B \rightarrow K\bar{\chi}\chi)/dq^2}. \quad (8)$$

For this ratio, the explicit form of  $\Pi_\chi(q^2)$  is of no importance: it is the same function in  $d\Gamma(B \rightarrow K\bar{\chi}\chi)/dq^2$  and  $d\Gamma(B \rightarrow K^*\bar{\chi}\chi)/dq^2$  and therefore drops out in the ratio. It is also obvious that the specific properties of the DM particles  $\chi$  are of no importance too: the final estimate holds for the DM particles independent of their spins. Moreover, the ratio is not sensitive to the properties of the scalar mediator  $\phi$  as its propagator also cancels in the ratio. Essential for the ratio (8) is merely *the operator structure of the vertex*,  $\phi\bar{s}_L b_R$ . Using the QCD equations of motion, it is straightforward to calculate the amplitudes

$$\begin{aligned} \langle K|\bar{s}_L b_R|B\rangle &= \frac{1}{2}\langle K|\bar{s}(1-\gamma_5)b|B\rangle = \frac{1}{2}\langle K|\bar{s}b|B\rangle = \frac{1}{2}\frac{M_B^2 - M_K^2}{m_b - m_s}f_0^{B\rightarrow K}(q^2), \\ \langle K^*|\bar{s}_L b_R|B\rangle &= \frac{1}{2}\langle K^*|\bar{s}(1-\gamma_5)b|B\rangle = -\frac{1}{2}\langle K^*|\bar{s}\gamma_5 b|B\rangle = -i(\epsilon q)\frac{M_{K^*}}{m_b + m_s}A_0^{B\rightarrow K^*}(q^2), \end{aligned} \quad (9)$$

where the dimensionless form factors  $f_0$  and  $A_0$  are well-known quantities parametrizing the  $\langle K|\bar{s}\gamma_\mu b|B\rangle$  and  $\langle K^*|\bar{s}\gamma_\mu\gamma_5 b|B\rangle$  amplitudes [11]. The decay rates of interest then take the form:

$$\frac{d\Gamma(B \rightarrow K\bar{\chi}\chi)}{dq^2} = \left| \frac{g_{b\rightarrow s\phi}\kappa}{M_\phi^2 - q^2} \right|^2 \frac{\lambda^{1/2}(M_B^2, M_K^2, q^2)}{16\pi M_B^3} |\langle K|\bar{s}_L b_R|B\rangle|^2, \quad (10)$$

$$\frac{d\Gamma(B \rightarrow K^*\bar{\chi}\chi)}{dq^2} = \left| \frac{g_{b\rightarrow s\phi}\kappa}{M_\phi^2 - q^2} \right|^2 \frac{\lambda^{1/2}(M_B^2, M_{K^*}^2, q^2)}{16\pi M_B^3} \sum_{K^*\text{polar}} |\langle K^*|\bar{s}_L b_R|B\rangle|^2. \quad (11)$$

Performing summation over the  $K^*$  polarizations, we find for the ratio (8):

$$R_{K^*/K}^{(\phi)}(q^2) = \frac{\lambda^{3/2}(M_B^2, M_{K^*}^2, q^2)}{\lambda^{1/2}(M_B^2, M_K^2, q^2)(M_B^2 - M_{K^*}^2)^2} \left[ \frac{m_b - m_s}{m_b + m_s} \right]^2 \left| \frac{A_0(q^2)}{f_0(q^2)} \right|^2. \quad (12)$$

*3. Numerical estimates.* For the form factors, we make use of the results from [12] and [13] in the form of convenient parametrizations of [14]:

$$f_0(q^2) = \frac{0.33}{1 - 0.7r_V + 0.27r_V^2}, \quad r_V = q^2/m_{B_s}^2, \quad (13)$$

$$A_0(q^2) = \frac{0.37}{(1 - 0.46r_P)(1 - r_P)}, \quad r_P = q^2/m_{B_s}^2. \quad (14)$$

We then come to the following numerical prediction for the ratio of the differential distributions in the reactions  $B \rightarrow K\phi \rightarrow K\chi\chi$  and  $B \rightarrow K^*\phi \rightarrow K^*\chi\chi$  shown in Fig. 1. Notice that using more recent parametrizations for the form factors, e.g., those from [15, 16], does not lead to any visible changes of the results in Fig. 1.

*4. Discussion.* As clear from the plot, one can obtain an upper limit:

$$\delta\Gamma(B \rightarrow K^*M_X)_\phi < 1.05 \delta\Gamma(B \rightarrow KM_X)_\phi. \quad (15)$$

This upper limit is obtained from Fig. 1 at  $M_X^2 = q^2 \rightarrow 0$ . The upper limit of Eq. (15) is a rather conservative estimate and in practice one may expect a larger suppression of  $\delta\Gamma(B \rightarrow K^*M_X)_\phi$  compared to  $\delta\Gamma(B \rightarrow KM_X)_\phi$ : The  $q^2$ -region providing the dominant contribution to the  $B \rightarrow (K, K^*)\bar{\chi}\chi$  cross sections is mainly determined by the resonant structure of the  $\phi$ -propagator and corresponds to nonzero  $q^2$  thus leading to a further suppression  $\delta\Gamma(B \rightarrow K^*M_X)_\phi < \delta\Gamma(B \rightarrow KM_X)_\phi$ .

Now, assuming that the observed enhancement of  $\Gamma(B \rightarrow K\nu\bar{\nu})_{\text{SM}}$  is due to unobserved DM particles interacting via a scalar mediator we can write

$$\Gamma(B \rightarrow KM_X) = \Gamma(B \rightarrow K\nu\bar{\nu})_{\text{SM}} + \delta\Gamma(B \rightarrow KM_X)_\phi, \quad (16)$$

$$\Gamma(B \rightarrow K^*M_X) = \Gamma(B \rightarrow K^*\nu\bar{\nu})_{\text{SM}} + \delta\Gamma(B \rightarrow K^*M_X)_\phi. \quad (17)$$




Fig. 1: The ratio  $R_{K^*/K}^{(\phi)} = d\Gamma(B \rightarrow K^* M_X) / d\Gamma(B \rightarrow K M_X)$  vs  $M_X$ ,  $M_X^2 = q^2$ ,  $q$  is the momentum of the outgoing  $\chi\bar{\chi}$  pair of DM particles interacting with the top quark via scalar mediator field. Phase-space restriction related to the finite masses of the DM  $\chi$  particles should be taken into account (i.e. not all values of  $M_X$  shown in the plot are accessible in the decay). We point out that all factors related to the  $\phi$ -propagator (including  $M_\phi$  and possible finite-width effects) and to the details of the  $\phi\chi\chi$  vertex cancel in the ratio.

For  $\mathcal{B}(B \rightarrow K^* \nu \bar{\nu}) / \mathcal{B}(B \rightarrow K \nu \bar{\nu})$  one obtains  $2.5 \pm 0.7$  making use of old estimates [19–21] or  $2.23 \pm 0.40$  using the recent analysis of [15]. Within the quoted uncertainties these two values are in excellent agreement with each other.

Collecting together the central values of the discussed constraints:

$$\Gamma(B \rightarrow K^* \nu \bar{\nu})_{\text{SM}} \simeq 2.5 \Gamma(B \rightarrow K \nu \bar{\nu})_{\text{SM}} \quad [\text{Theoretical estimates}] \quad (18)$$

$$\delta\Gamma(B \rightarrow K M_X)_\phi \simeq 4.4 \Gamma(B \rightarrow K \nu \bar{\nu})_{\text{SM}} \quad [\text{Belle measurement}] \quad (19)$$

$$\delta\Gamma(B \rightarrow K^* M_X)_\phi < 1.05 \delta\Gamma(B \rightarrow K M_X)_\phi \quad [\text{Our result here}], \quad (20)$$

we then obtain the following upper bound for the  $\Gamma(B \rightarrow K^* M_X)$  within the *scalar mediator scenario*:

$$\Gamma(B \rightarrow K^* M_X) < 2.8 \Gamma(B \rightarrow K^* \nu \bar{\nu})_{\text{SM}}. \quad (21)$$

Therefore, for the case of scalar mediator, an expected enhancement of the branching ratio  $\Gamma(B \rightarrow K^* M_X)$  compared to the SM value is approximately two times smaller than the enhancement for  $\Gamma(B \rightarrow K M_X)$ , Eq. (1). This constraint is well compatible with the present experimental limit [17]

$$\mathcal{B}(B \rightarrow K^* M_X) \leq 2.7 \times 10^{-5} \leq 2.7 \mathcal{B}(B \rightarrow K^* \nu \bar{\nu})_{\text{SM}}, \quad (22)$$

where at the second step we have used a recently updated estimate  $\mathcal{B}(B \rightarrow K^* \nu \bar{\nu})_{\text{SM}} = (1.03 \pm 0.15) \times 10^{-5}$  [15].

On the other hand, if the “mediator” is a spin-1 field, it is natural to expect a similar enhancement for both  $K$  and  $K^*$  mesons in the final state,  $\mathcal{B}(B \rightarrow K^* M_X) / \mathcal{B}(B \rightarrow K^* \nu \bar{\nu})_{\text{SM}} \simeq \mathcal{B}(B \rightarrow K M_X) / \mathcal{B}(B \rightarrow K \nu \bar{\nu})_{\text{SM}}$ .

We therefore conclude that the combined analysis of  $\mathcal{B}(B \rightarrow K^* M_X)$  and  $\mathcal{B}(B \rightarrow K M_X)$  may provide a useful probe of the scalar mediator scenario [e.g., may exclude this scenario if the obtained upper limit is violated]. Since the obtained constraints are not sensitive to the scalar mediator mass  $M_\phi$ , the latter should then be determined from other observables.

*Acknowledgments.* The research was carried out within the framework of the program “Particle Physics and Cosmology” of the National Center for Physics and Mathematics.

---

[1] A. Glazov (Belle II), *News from Belle II*, BELLE2-TALK-CONF-2023-123.

[2] P. Athron, R. Martinez, and C. Sierra, *B meson anomalies and large  $B^+ \rightarrow K^+ \nu \bar{\nu}$  in non-universal  $U(1)'$  models*, arXiv:2308.13426.

[3] R. Bause, H. Gisbert, and G. Hiller, *Implications of an enhanced  $B \rightarrow K \nu \bar{\nu}$  branching ratio*, arXiv:2309.00075.

[4] L. Allwicher, D. Becirevic, G. Piazza, S. Rosauero-Alcaraz, O. Sumensari, *Understanding the first measurement of  $Br(B \rightarrow K \nu \bar{\nu})$* , arXiv:2309.02246.

[5] T. Felkl, A. Giri, R. Mohanta, M. A. Schmidt, *When Energy Goes Missing: New Physics in  $b \rightarrow s \nu \bar{\nu}$  with Sterile Neutrinos*, arXiv:2309.02940.

[6] M. Abdughani and Y. Reyimuaji, *Constraining light dark matter and mediator with  $B^+ \rightarrow K^+ \nu \bar{\nu}$  data*, arXiv:2309.03706.

- [7] H. K. Dreiner, J. Y. Günther, and Z. S. Wang, *The Decay  $B \rightarrow K\nu\bar{\nu}$  at Belle II and a Massless Bino in R-parity-violating Supersymmetry*, arXiv:2309.03727.
- [8] X.-G. He, X.-D. Ma, and G. Valencia, *Revisiting models that enhance  $B \rightarrow K\nu\bar{\nu}$  in light of the new Belle II measurement*, arXiv:2309.12741.
- [9] B. Batell, M. Pospelov, and A. Ritz, *Multi-lepton Signatures of a Hidden Sector in Rare B Decays*, Phys. Rev. **D83**, 054005 (2011).
- [10] K. Schmidt-Hoberg, F. Staub, and M. W. Winkler, *Constraints on light mediators: confronting dark matter searches with B physics*, Phys. Lett. **B727**, 506 (2013).
- [11] M. Wirbel, B. Stech, and M. Bauer, *Exclusive Semileptonic Decays of Heavy Mesons*, Z. Phys. **C29**, 637 (1985).
- [12] P. Ball and R. Zwicky, *New results on  $B \rightarrow \pi, K, \eta$  decay formfactors from LCSRs*, Phys. Rev. **D71**, 014015 (2005).
- [13] P. Ball and R. Zwicky,  *$B_{d,s} \rightarrow \rho, \omega, K^*, \phi$  decay form-factors from LCSRs revisited*, Phys. Rev. **D71**, 014029 (2005).
- [14] D. Melikhov and B. Stech, *Weak form-factors for heavy meson decays: an update*, Phys. Rev. **D62**, 014006 (2000).
- [15] D. Becirevic, G. Piazza, and O. Sumensari, *Revisiting  $B \rightarrow K^{(*)}\nu\bar{\nu}$  decays in the Standard Model and beyond*, Eur. Phys. J. **C83**, 252 (2023).
- [16] W. G. Parrott, C. Bouchard, and C. T. H. Davies (HPQCD collaboration), *Standard Model predictions for  $B \rightarrow Kl^+l^-$ ,  $B \rightarrow Kl_1^-l_2^+$ , and  $B \rightarrow K\nu\bar{\nu}$  using form factors from  $N_f = 2 + 1 + 1$  lattice QCD*, Phys. Rev. **D107**, 014511 (2023); Phys. Rev. **D107**, 119903(E) (2023).
- [17] J. Grygier *et al.* [Belle], *Search for  $B \rightarrow h\nu\bar{\nu}$  decays with semileptonic tagging at Belle*, Phys. Rev. **D96**, 091101 (2017).
- [18] P. Colangelo, F. De Fazio, P. Santorelli, and E. Scrimieri, *Rare  $B \rightarrow (K, K^*)\bar{\nu}\nu$  decays at B-factories*, Phys. Lett. **B395**, 339 (1997).
- [19] D. Melikhov, N. Nikitin, and S. Simula, *Rare decays  $B \rightarrow (K, K^*)(l^+l^-, \nu\bar{\nu})$  in the quark model*, Phys. Lett. **B410**, 290 (1997).
- [20] D. Melikhov, N. Nikitin, and S. Simula, *Right-handed currents in rare exclusive  $B \rightarrow (K, K^*)\nu\bar{\nu}$  decays*, Phys. Lett. **B428**, 171 (1997).
- [21] D. Melikhov, N. Nikitin, and S. Simula, *Rare exclusive semileptonic  $b \rightarrow s$  transitions in the SM*, Phys. Rev. **D57**, 6814 (1998).