REFINED AND REFINED HARMONIC JACOBI–DAVIDSON METHODS FOR COMPUTING SEVERAL GSVD COMPONENTS OF A LARGE REGULAR MATRIX PAIR*

JINZHI HUANG[†] AND ZHONGXIAO JIA[‡]

Abstract. Three refined and refined harmonic extraction-based Jacobi–Davidson (JD) type methods are proposed, and their thick-restart algorithms with deflation and purgation are developed to compute several generalized singular value decomposition (GSVD) components of a large regular matrix pair. The new methods are called refined cross product-free (RCPF), refined cross productfree harmonic (RCPF-harmonic) and refined inverse-free harmonic (RIF-harmonic) JDGSVD algorithms, abbreviated as RCPF-JDGSVD, RCPF-HJDGSVD and RIF-HJDGSVD, respectively. The new JDGSVD methods are more efficient than the corresponding standard and harmonic extractionbased JDSVD methods proposed previously by the authors, and can overcome the erratic behavior and intrinsic possible non-convergence of the latter ones. Numerical experiments illustrate that RCPF-JDGSVD performs better for the computation of extreme GSVD components while RCPF-HJDGSVD and RIF-HJDGSVD suit better for that of interior GSVD components.

Key words. Generalized singular value decomposition, generalized singular value, generalized singular vector, standard extraction, harmonic extraction, refined extraction, refined harmonic extraction, Jacobi–Davidson type method

AMS subject classifications. 65F15, 15A18, 65F10

1. Introduction. The GSVD was initially established by Van Loan [37] and developed by Paige and Saunders [27], and it has soon become one of the most important matrix decompositions [4, 33, 34].

Let $A \in \mathbb{R}^{m \times n}$ and $B \in \mathbb{R}^{p \times n}$ with $m + p \ge n$. Suppose that $\mathcal{N}(A) \cap \mathcal{N}(B) = \{\mathbf{0}\}$, where $\mathcal{N}(\cdot)$ denotes the null space of a matrix. Then (A, B) is called a regular matrix pair. Write $q_1 = \dim(\mathcal{N}(A))$, $q_2 = \dim(\mathcal{N}(B))$ and $l_1 = \dim(\mathcal{N}(A^T))$, $l_2 = \dim(\mathcal{N}(B^T))$, respectively, where the superscript T denotes the transpose of a matrix and $\dim(\cdot)$ denotes the dimension of a subspace. Then the GSVD of (A, B) is as follows:

(1.1)
$$\begin{cases} A = U\Sigma_A X^{-1}, \\ B = V\Sigma_B X^{-1}, \end{cases} \text{ with } \begin{cases} \Sigma_A = \text{diag}\{C, \mathbf{0}_{l_1, q_1}, I_{q_2}\}, \\ \Sigma_B = \text{diag}\{S, I_{q_1}, \mathbf{0}_{l_2, q_2}\}, \end{cases}$$

where $U = [U_q, U_{l_1}, U_{q_2}]$ and $V = [V_q, V_{q_1}, V_{l_2}]$ are orthogonal, $X = [X_q, X_{q_1}, X_{q_2}]$ is nonsingular, and the diagonal $C = \text{diag}\{\alpha_1, \ldots, \alpha_q\}$ and $S = \text{diag}\{\beta_1, \ldots, \beta_q\}$ satisfy

$$0 < \alpha_i, \beta_i < 1$$
 and $\alpha_i^2 + \beta_i^2 = 1, \quad i = 1, ..., q$

with $q = n - q_1 - q_2$. Here the subscripts in the block submatrices of U, V, X are their column numbers, and I_i and $\mathbf{0}_{i,j}$ denote the *i*-by-*i* identity matrix and *i*-by-*j* zero matrix, respectively. The subscripts are dropped whenever their sizes are clear from the context. Let u_i, v_i and x_i be the *i*th columns of U_q, V_q and X_q , respectively,

^{*}The work of the first author was supported by the Youth Fund of the National Science Foundation of China (No. 12301485) and the Youth Program of the Natural Science Foundation of Jiangsu Province (No. BK20220482), and the work of the second author was supported by the National Science Foundation of China (No. 12171273).

[†]School of Mathematical Sciences, Soochow University, 215006 Suzhou, China (jzhuang21@suda.edu.cn)

[‡]Corresponding author. Department of Mathematical Sciences, Tsinghua University, 100084 Beijing, China (jiazx@tsinghua.edu.cn).

 $i = 1, \ldots, q$. Then the quintuples $(\alpha_i, \beta_i, u_i, v_i, x_i)$, $i = 1, \ldots, q$ are called *nontrivial* GSVD components of (A, B). In particular, the scalar pairs (α_i, β_i) or, equivalently, $\sigma_i = \frac{\alpha_i}{\beta_i}$ are called the nontrivial generalized singular values of (A, B), and u_i, v_i and x_i are the corresponding left and right generalized singular vectors, respectively.

From (1.1), for i = 1, 2, ..., q, the GSVD of (A, B) can be written in the form

(1.2)
$$\begin{cases} Ax_i = \alpha_i u_i, \\ Bx_i = \beta_i v_i, \\ \beta_i A^T u_i = \alpha_i B^T v_i \end{cases}$$

Denote by $(\alpha_i, \beta_i) = (0, 1)$ or $(\alpha_i, \beta_i) = (1, 0)$ a trivial zero or infinite generalized singular value. Then the above form still holds with u_i, v_i and x_i being the left and right generalized singular vectors corresponding to the zero or infinite generalized singular value. From (1.1), we have $X^T(A^TA + B^TB)X = I_n$. Therefore, X is $(A^TA + B^TB)$ -orthogonal, and its columns x_i 's are of $(A^TA + B^TB)$ -norm unit length. Naturally, we require that any approximation to x_i have the same length.

In this paper, we consider the following GSVD computational problem of a large and possibly sparse regular matrix pair (A, B).

Problem 1.1. For a given target $\tau > 0$, label all the nontrivial generalized singular values of (A, B) as

(1.3)
$$|\sigma_1 - \tau| \le \dots \le |\sigma_\ell - \tau| < |\sigma_{\ell+1} - \tau| \le \dots \le |\sigma_q - \tau|$$

We want to compute the GSVD components $(\alpha_i, \beta_i, u_i, v_i, x_i)$, $i = 1, \ldots, \ell$ associated with the ℓ generalized singular values σ_i , $i = 1, \ldots, \ell$ of (A, B) closest to τ .

If τ is inside the spectrum of the nontrivial generalized singular values of (A, B), then those $(\alpha_i, \beta_i, u_i, v_i, x_i)$'s are called interior GSVD components of (A, B); if τ is close to one of the ends of the nontrivial generalized singular spectrum, then they are called the extreme, i.e., largest or smallest, ones. In the sequel, we assume that the target τ is not equal to any generalized singular value of (A, B).

Hochstenbach [7] proposes a Jacobi–Davidson (JD) type GSVD method, called JDGSVD, to compute several extreme or interior GSVD components of (A, B) where B has full column rank. At the subspace expansion phase, an (m+n)-by-(m+n) linear system, called the correction equation, needs to be solved iteratively; for analysis and details on the accuracy requirement on the inner iterations of JD type methods for eigenvalue and SVD problems, see [9, 11, 12, 19, 20], where it is shown that it generally suffices to solve correction equations with low or modest accuracy; that is, the relative errors of approximate solutions lie in $[10^{-4}, 10^{-2}]$. More generally, the JDGSVD method formulates the GSVD of (A, B) either as the generalized eigendecomposition of the augmented matrix pair $\left(\begin{bmatrix} A^T & A \end{bmatrix}, \begin{bmatrix} I & B^T B \end{bmatrix}\right)$ for B of full column rank or that of $\left(\begin{bmatrix} B^T & B \end{bmatrix}, \begin{bmatrix} I & A^T A \end{bmatrix}\right)$ for A of full column rank, computes the corresponding generalized eigenpairs, and then reconstructs the desired approximate GSVD components from the relevant converged eigenpairs. For the second formulation, an (p+n)-by-(p+n)correction equation is solved iteratively at each subspace expansion step. However, as has been theoretically proven and numerically confirmed in [10, 12], a fairly ill conditioned B or A may make the corresponding JDGSVD method numerically backward unstable for the GSVD problem itself, even if the relative residuals of approximate eigenpairs of the underlying generalized eigenvalue problem are already at the level of machine precision.

Zwaan and Hochstenbach [41] present two GSVD methods, called the generalized Davidson (GDGSVD) and multidirectional (MDGSVD) methods, to compute several extreme GSVD components of (A, B). The right searching subspace for the GDGSVD method is spanned by the residuals of the generalized Davidson method [1, Sec. 11.2.4 and Sec. 11.3.6] applied to the eigenvalue problem of the cross-product matrix pair $(A^T A, B^T B)$; that for the MDGSVD method is first expanded by two dimensions with the vectors formed by premultiplying the best approximate right generalized singular vector with $A^T A$ and $B^T B$, and then truncated by one dimension so that an inferior search direction is discarded. The left searching subspaces for these two methods are formed by premultiplying the right one with A and B, respectively. These two methods make use of the standard extraction approach to compute the approximate GSVD components without explicitly forming $A^T A$, $B^T B$. Zwaan [40] utilizes the Kronecker canonical form of a matrix pair [34], and proves that the GSVD of (A, B)is equivalent to the generalized eigendecomposition of a matrix pair with much larger order 2m + p + n. Though the pair does not involve cross-products or any other matrix-matrix product, this formulation may be mainly of theoretical value since (i) the nontrivial generalized eigenvalues and eigenvectors of the larger structured matrix pair come in quadruples and are always complex, (ii) the conditioning of the structured generalized eigenvalue problem is unknown, and (iii) it is extremely hard to propose a numerically backward stable structure-preserving algorithm.

Adapted standard and harmonic Rayleigh–Ritz projections [4, 28, 30, 33], or called the standard and harmonic extraction approaches, for eigenvalue and generalized eigenvalue problems to GSVD problems, the authors have recently proposed the cross product-free (CPF) JDGSVD [12], the CPF-harmonic and inverse-free (IF) harmonic JDGSVD methods [11], written as CPF-JDGSVD, CPF-HJDGSVD and IF-HJDGSVD for short, to solve Problem 1.1. At the subspace expansion step, each of these methods requires an approximate solution of its own *n*-by-*n* correction equation, and uses it to expand the right searching subspace; the two left searching subspaces are formed by premultiplying the right one with A and B, respectively. The three methods are fundamentally different in the extraction phase. The CPF-JDGSVD method works on (A, B) directly, applies the standard extraction approach to the left and right searching subspaces, and computes the GSVD of a small projection matrix pair. It implicitly realizes the standard Rayleigh–Ritz projection of the generalized eigenvalue problem of $(A^T A, B^T B)$ onto the right searching subspace [12]. With B of full column rank, the CPF-HJDGSVD method implicitly realizes the harmonic extraction approach of the singular value decomposition (SVD) problem of AL^{-T} onto the right and one of the left searching subspaces, where $L^T \in \mathbb{R}^{n \times n}$ is the factor in the sparse Cholesky factorization $B^T B = L L^T$. At each extraction step, the method needs to solve the generalized eigenvalue problem of a small symmetric positive definite matrix pair. For a general and possibly rank deficient B, the IF-HJDGSVD method implicitly carries out the harmonic extraction of the generalized eigenvalue problem of $(A^T A, B^T B)$ onto the right searching subspace, and computes the generalized eigendecomposition of a small symmetric positive definite matrix pair; it is the inverses $(A^T A)^{-1}$ and $(B^T B)^{-1}$ -free, and works for a general matrix pair (A, B). For justifications and details, we refer the reader to [11]

Just like those standard Rayleigh–Ritz methods for the matrix eigenvalue problem and the SVD problem [2, 5, 7, 31, 35, 41], CPF-JDGSVD suits better for the computation of extreme generalized singular values of (A, B). However, we deduce from, e.g., [16, 21, 23], that the approximate generalized singular vectors obtained by it may converge erratically or even fail to converge even if the approximate generalized singular values converge. These phenomena have been numerically observed in [12].

The refined extraction approach or refined Rayleigh–Ritz projection was initially proposed by the second author in [13], and has been intensively studied and developed in, e.g., [6, 8, 14, 17, 20, 21, 22, 23, 24, 38, 39] for the eigenvalue and SVD problems. For the large matrix eigenvalue problem, it is systematically accounted for in the books [1, 33, 36]. As is shown, the refined extraction has better convergence, and fixes the erratic convergence behavior and possible non-convergence of standard and harmonic extractions [13, 14, 16, 17, 22, 23]; also see, e.g., [6, 8, 24, 38, 39]. Importantly, the basic convergence results in [16, 17, 23] adapted to CPF-JDGSVD, CPF-HJDGSVD and IF-HJDGSVD indicate that these three methods inherit those mentioned convergence deficiencies of the standard and harmonic extractions; that is, the three methods may work erratically and inefficiently.

In this paper, in order to fix the deficiency of CPF-JDGSVD, CPF-HJDGSVD and IF-HJDGSVD and to better solve Problem 1.1, we will propose three refined extraction-based JDGSVD methods. We first present a refined JDGSVD (RCPF-JDGSVD) method. It computes the approximate generalized singular values by the standard extraction-based JDGSVD method but nontrivially adapts the refined extraction of the generalized eigenvalue problem of (A^TA, B^TB) to the GSVD problem of (A, B), and seeks new approximate generalized singular vectors that are generally more and can be much more accurate than those obtained by the standard extraction.

Like the interior eigenvalue and SVD problems, for the computation of interior GSVD components, the standard extraction may produce spurious Ritz values and has difficulty to pick up good Ritz values, if any, correctly even if the searching subspaces are sufficiently good [20, 23, 33, 36], causing that the CPF-JDGSVD method may converge slowly or even fail to converge, as has been numerically confirmed in [11]. Whenever Ritz values are poor or, though good, they are selected incorrectly, a refined extraction-based method certainly delivers incorrect approximate GSVD components, which severely affects the correct expansion of underlying subspaces in JDGSVD type methods. As a result, the refined extraction-based method may perform poorly when computing interior GSVD components. Just as the harmonic extraction-based methods for the eigenvalue and SVD problems [33, 36] that suit better for computing interior eigenpairs and singular triplets [6, 8, 9, 15, 20, 22, 25, 26], our two harmonic extraction-based CPF-HJDGSVD and IF-HJDGSVD methods are more suitable for computing interior generalized singular values. Nevertheless, the harmonic Ritz vectors may have erratic convergence behavior and may even fail to converge because of spurious harmonic Ritz value(s) even if searching subspaces are sufficiently accurate [17, 20].

In order to overcome the aforementioned deficiency, for the computation of interior GSVD components, on the basis of CPF-HJDGSVD and IF-HJDGSVD, we propose refined harmonic JDGSVD type methods that retain the merits of the harmonic JDGSVD methods for computing generalized singular values but seek more accurate approximate generalized singular vectors by using the refined extraction in a proper way. The resulting methods are abbreviated as RCPF-HJDGSVD and RIF-HJDGSVD, respectively.

We first focus on the case $\ell = 1$, and propose basic refined and refined harmonic extraction-based JDGSVD methods for the GSVD problem of interest. Then combining the methods with appropriate restart, deflation and purgation, we develop thick-restart RCPF-JDGSVD, RCPF-HJDGSVD and RIF-HJDGSVD algorithms for Prob-

lem 1.1 with $\ell > 1$, with details on effective and efficient implementations described. We will numerically demonstrate that they have better convergence behavior and are considerably more efficient than the corresponding standard and harmonic extractionbased JDGSVD algorithms. We also illustrate that RCPF-JDGSVD performs better than RCPF-HJDGSVD and RIF-HJDGSVD for extreme GSVD components, but for interior GSVD problems the latter two ones are preferable and RIF-HJDGSVD has wider applicability than RCPF-HJDGSVD.

The rest of this paper is organized as follows. In Section 2, we review the CPF-JDGSVD method in [12] and the CPF-HJDGSVD and IF-HJDGSVD methods in [11]. In Section 3, we introduce a refined extraction approach for the GSVD computation; combining it with the standard, CPF-harmonic and IF-harmonic extractions, we propose the refined, refined CPF-harmonic and refined IF-harmonic extraction-based JDGSVD methods: RCPF-JDGSVD, RCPF-HJDGSVD, and RIF-HJDGSVD. In Section 4, we develop thick-restart schemes of these three JDGSVD algorithms with effective deflation and purgation for computing several GSVD components of (A, B). Numerical experiments are presented in Section 5 to illustrate the performance of the three refined JDGSVD algorithms and to make comparisons of them and the three standard and harmonic extraction-based JDGSVD algorithms. Finally, we conclude the paper in Section 6.

2. The standard and two harmonic extraction-based JDGSVD methods. We review CPF-JDGSVD, CPF-HJDGSVD and IF-HJDGSVD in [11, 12] for computing $(\alpha_*, \beta_*, u_*, v_*, x_*) := (\alpha_1, \beta_1, u_1, v_1, x_1)$. Section 2.1 is devoted to the construction and expansion of the searching subspaces, Section 2.2 reviews the standard extraction, Section 2.3 is on the CPF-harmonic extraction, and Section 2.4 describes the IF-harmonic extraction.

2.1. The construction and expansion of searching subspaces. Assume that a k-dimensional right searching subspace $\mathcal{X} \subset \mathbb{R}^n$ is available, from which an approximation to x_* is sought. Then we construct the two left searching subspaces

(2.1)
$$\mathcal{U} = A\mathcal{X} \quad \text{and} \quad \mathcal{V} = B\mathcal{X},$$

from which approximations to u_* and v_* are extracted, respectively. Theorem 2.1 of [12] shows that the distance between u_* and \mathcal{U} is as small as that between x_* and \mathcal{X} provided that α_* is not very small; analogously, the distance between v_* and \mathcal{V} is as small as that between x_* and \mathcal{X} if β_* is not very small. Therefore, for the GSVD components corresponding to not very large or small generalized singular values, the left searching subspaces \mathcal{U} and \mathcal{V} constructed by (2.1) are as good as \mathcal{X} .

Let $\widetilde{X} \in \mathbb{R}^{n \times k}$ be an orthonormal basis matrix of \mathcal{X} , and compute the thin QR factorizations

(2.2)
$$A\widetilde{X} = \widetilde{U}R_A$$
 and $B\widetilde{X} = \widetilde{V}R_B$

to obtain the orthonormal basis matrices $\widetilde{U} \in \mathbb{R}^{m \times k}$ and $\widetilde{V} \in \mathbb{R}^{p \times k}$ of \mathcal{U} and \mathcal{V} . With \mathcal{U}, \mathcal{V} and \mathcal{X} as well as their orthonormal bases available, we can use one of the following six extraction approaches to compute an approximation to the desired GSVD component $(\alpha_*, \beta_*, u_*, v_*, x_*)$: (i) the standard extraction, as done in the CPF-JDGSVD method [12] and will be reviewed in Section 2.2; (ii) the CPF-harmonic extraction, as exploited by the CPF-HJDGSVD method [11] and will be sketched in Section 2.3; (iii) the IF-harmonic extraction, as adopted in the IF-HJDGSVD method [11] and will be reviewed in Section 2.4; (iv) the refined CPF extraction; (v) the refined CPF-harmonic extraction; (vi) the refined IF-harmonic extraction.

In Section 3, we shall propose the three extraction approaches in (iv)–(vi). Together with their extensions and restart schemes for computing more than one GSVD components that will be presented in Section 4, we will set up our complete algorithms, which constitute our major contribution in this paper.

We temporarily denote by $(\tilde{\alpha}, \tilde{\beta}, \tilde{u}, \tilde{v}, \tilde{x})$ an approximation to the desired GSVD component $(\alpha_*, \beta_*, u_*, v_*, x_*)$ computed by any one of the six extraction approaches listed above, where the positive scalar pair $(\tilde{\alpha}, \tilde{\beta})$ is required to satisfy $\tilde{\alpha}^2 + \tilde{\beta}^2 = 1$, and the 2-norm unit length vectors $\tilde{u} \in \mathcal{U}$, $\tilde{v} \in \mathcal{V}$ and the $(A^T A + B^T B)$ -norm unit length $\tilde{x} \in \mathcal{X}$ are required to satisfy $A\tilde{x} = \tilde{\alpha}\tilde{u}$ and $B\tilde{x} = \tilde{\beta}\tilde{v}$. Therefore, for the GSVD problem of (A, B), in terms of (1.2), the GSVD residual of $(\tilde{\alpha}, \tilde{\beta}, \tilde{u}, \tilde{v}, \tilde{x})$ is

(2.3)
$$r = r(\tilde{\alpha}, \tilde{\beta}, \tilde{u}, \tilde{v}, \tilde{x}) = \tilde{\beta} A^T \tilde{u} - \tilde{\alpha} B^T \tilde{v}.$$

Clearly, $(\tilde{\alpha}, \tilde{\beta}, \tilde{u}, \tilde{v}, \tilde{x})$ is an exact GSVD component of (A, B) if and only if r = 0. Let tol > 0 be a user-prescribed stopping tolerance. If

(2.4)
$$||r|| \le (\hat{\beta} ||A||_1 + \tilde{\alpha} ||B||_1) \cdot tol,$$

we stop the iterations and accept $(\tilde{\alpha}, \tilde{\beta}, \tilde{u}, \tilde{v}, \tilde{x})$ as a converged approximation to the desired $(\alpha_*, \beta_*, u_*, v_*, x_*)$. Throughout the paper, we denote by $\|\cdot\|$ and $\|\cdot\|_1$ the 2- and 1-norms of a matrix or vector, respectively.

If $(\tilde{\alpha}, \tilde{\beta}, \tilde{u}, \tilde{v}, \tilde{x})$ does not yet converge, a JDGSVD type method first expands the right searching subspace \mathcal{X} , then updates the left searching subspaces \mathcal{U} and \mathcal{V} in the way (2.1). Specifically, notice that

(2.5)
$$\tilde{y} = (A^T A + B^T B)\tilde{x} = \tilde{\alpha}A^T \tilde{u} + \tilde{\beta}B^T \tilde{v}$$

satisfies $\tilde{y}^T \tilde{x} = 1$. Therefore, $I - \tilde{y}\tilde{x}^T$ and $I - \tilde{x}\tilde{y}^T$ are oblique projectors. We approximately solve the correction equation

(2.6)
$$(I - \tilde{y}\tilde{x}^T)(A^T A - \rho^2 B^T B)(I - \tilde{x}\tilde{y}^T)t = -r \quad \text{for} \quad t \perp \tilde{y}$$

using some Krylov subspace iterative method such as the MINRES method [29] with the fixed $\rho = \tau$ when the approximate GSVD component $(\tilde{\alpha}, \tilde{\beta}, \tilde{u}, \tilde{v}, \tilde{x})$ is not yet reasonably good, and then switch to solving problem (2.6) with the adaptively changing $\rho = \tilde{\theta} := \tilde{\alpha}/\tilde{\beta}$ if

(2.7)
$$\|r\| \le (\hat{\beta} \|A\|_1 + \tilde{\alpha} \|B\|_1) \cdot fixtol$$

for a user-prescribed tolerance fixed > 0, say, 10^{-4} . Criterion (2.7) means that $(\tilde{\alpha}, \tilde{\beta}, \tilde{u}, \tilde{v}, \tilde{x})$ is already a fairly good approximation to $(\alpha_*, \beta_*, u_*, v_*, x_*)$.

Iteratively solving the correction equations of form (2.6) in the JDGSVD type methods is called the inner iterations, and the extraction of approximate GSVD components with respect to \mathcal{U} , \mathcal{V} and \mathcal{X} is called the outer iterations. It has been shown in [12] that solving the correction equations with *low or modest* accuracy generally suffices to make the outer iterations of the resulting inexact JD type GSVD algorithms well mimic those of their exact counterparts where all the correction equations are solved accurately. Therefore, for the correction equations of form (2.6) in the JDGSVD methods proposed in [11, 12] and in the new JDGSVD methods to be proposed in this paper, we adopt the inner stopping criterion in [12], and stop the inner iterations when the inner relative residual norm $||r_{in}||$ of an approximate solution satisfies

$$||r_{in}|| \le \min\{2c\tilde{\varepsilon}, 0.01\},\$$

where $\tilde{\varepsilon} \in [10^{-4}, 10^{-3}]$ is a user-prescribed parameter and c is a constant depending on the value of ρ and all the approximate generalized singular values of (A, B) computed by the underlying JDGSVD method during the current outer iteration.

An approximate solution of (2.6), still denoted by t for brevity, is utilized to expand \mathcal{X} so as to obtain the new $\mathcal{X}_{new} = \operatorname{span}\{\widetilde{X}, t\}$, and the corresponding orthonormal basis matrix is updated by

- - -

(2.9)
$$\widetilde{X}_{\text{new}} = [\widetilde{X}, x_+] \quad \text{with} \quad x_+ = \frac{(I - \widetilde{X}\widetilde{X}^T)t}{\|(I - \widetilde{X}\widetilde{X}^T)t\|},$$

where x_+ is called an expansion vector. Making use of (2.1) and (2.2) gives rise to the expanded left searching subspaces

$$\mathcal{U}_{\text{new}} = A\mathcal{X}_{\text{new}} = \text{span}\{\widetilde{U}, Ax_+\} \text{ and } \mathcal{V}_{\text{new}} = B\mathcal{X}_{\text{new}} = \text{span}\{\widetilde{V}, Bx_+\}.$$

We obtain their orthonormal basis matrices $\widetilde{U}_{\rm new}$ and $\widetilde{V}_{\rm new}$ by updating the thin QR factorizations

(2.10)
$$A\widetilde{X}_{\text{new}} = \widetilde{U}_{\text{new}} \cdot R_{A,\text{new}} = [\widetilde{U}, \widetilde{u}_+] \cdot \begin{bmatrix} R_A & r_A \\ & \gamma_A \end{bmatrix},$$

(2.11)
$$B\widetilde{X}_{\text{new}} = \widetilde{V}_{\text{new}} \cdot R_{B,\text{new}} = [\widetilde{V}, \widetilde{v}_+] \cdot \begin{bmatrix} R_B & r_B \\ & \gamma_B \end{bmatrix}$$

where

(2.12)
$$r_A = \widetilde{U}^T A x_+, \qquad \gamma_A = \|A x_+ - \widetilde{U} r_A\|, \qquad \widetilde{u}_+ = \frac{A x_+ - \widetilde{U} r_A}{\gamma_A},$$

(2.13) $r_B = \widetilde{V}^T B x_+, \qquad \gamma_B = \|B x_+ - \widetilde{V} r_B\|, \qquad \widetilde{v}_+ = \frac{B x_+ - \widetilde{V} r_B}{\gamma_A}$

We then compute a new and hopefully better approximation $(\tilde{\alpha}, \tilde{\beta}, \tilde{u}, \tilde{v}, \tilde{x})$ with respect to \mathcal{X}_{new} and \mathcal{U}_{new} , \mathcal{V}_{new} , and repeat the above process until convergence occurs.

2.2. The standard extraction approach. Given k-dimensional right and left searching subspaces \mathcal{X} and \mathcal{U} , \mathcal{V} of form (2.1), the standard extraction approach finds nonnegative pairs $(\tilde{\alpha}, \tilde{\beta})$ with $\tilde{\alpha}^2 + \tilde{\beta}^2 = 1$, unit length $\tilde{u} \in \mathcal{U}$ and $\tilde{v} \in \mathcal{V}$, and $(A^T A + B^T B)$ -norm unit length $\tilde{x} \in \mathcal{X}$ satisfying the conditions

(2.14)
$$\begin{cases} A\tilde{x} = \tilde{\alpha}\tilde{u}, \\ B\tilde{x} = \tilde{\beta}\tilde{v}, \\ \tilde{\beta}A^T\tilde{u} - \tilde{\alpha}B^T\tilde{v} \perp \mathcal{X}. \end{cases}$$

Write $\tilde{\theta} = \tilde{\alpha} / \tilde{\beta}$. It is straightforward to justify that

$$(A^T A - \tilde{\theta}^2 B^T B) \tilde{x} \perp \mathcal{X},$$

which is exactly the standard Rayleigh–Ritz projection of the generalized eigenvalue problem of the matrix pair $(A^T A, B^T B)$ onto the subspace \mathcal{X} , and each of the k pairs $(\tilde{\theta}^2, \tilde{x})$ is a Ritz approximation. Therefore, we call $(\tilde{\alpha}, \tilde{\beta}, \tilde{u}, \tilde{v}, \tilde{x})$ a Ritz approximation to $(\alpha_*, \beta_*, u_*, v_*, x_*)$ with $(\tilde{\alpha}, \tilde{\beta})$ or $\tilde{\theta} = \frac{\tilde{\alpha}}{\tilde{\beta}}$ the Ritz value and \tilde{u}, \tilde{v} and \tilde{x} the left and right Ritz vectors of (A, B) with respect to the left and right subspaces, respectively.

It is known from (2.2) that the projection matrices $\tilde{U}^T A \tilde{X} = R_A$ and $\tilde{V}^T A \tilde{X} = R_B$. Write $\tilde{u} = \tilde{U}\tilde{e}, \ \tilde{v} = \tilde{V}\tilde{f}$ and $\tilde{x} = \tilde{X}\tilde{d}$. Then (2.14) reduces to

(2.15)
$$R_A \tilde{d} = \tilde{\alpha} \tilde{e}, \qquad R_B \tilde{d} = \tilde{\beta} \tilde{f}, \qquad \tilde{\beta} R_A^T \tilde{e} = \tilde{\alpha} R_B^T \tilde{f},$$

which is the vector form of GSVD of (R_A, R_B) . Therefore, in the extraction phase, the standard extraction-based CPF-JDGSVD method computes the GSVD of the k-by-k matrix pair (R_A, R_B) , picks up the GSVD component $(\tilde{\alpha}, \tilde{\beta}, \tilde{e}, \tilde{f}, \tilde{d})$ corresponding to the generalized singular value $\tilde{\theta} = \frac{\tilde{\alpha}}{\tilde{\beta}}$ closest to the target τ , and takes

$$(\tilde{\alpha}, \tilde{\beta}, \tilde{u}, \tilde{v}, \tilde{x}) = (\tilde{\alpha}, \tilde{\beta}, \widetilde{U}\tilde{e}, \widetilde{V}\tilde{f}, \widetilde{X}\tilde{d})$$

as an approximation to the desired GSVD component $(\alpha_*, \beta_*, u_*, v_*, x_*)$ of (A, B).

2.3. The CPF-harmonic extraction approach. For *B* of full column rank, let $B^T B = LL^T$ be the Cholesky factorization of $B^T B$. It is proven in [11] that (σ_*, u_*, z_*) with $z_* = \frac{1}{\beta_*} L^T x_*$ is a singular triplet of the matrix

Take the k-dimensional \mathcal{U} and $\mathcal{Z} = L^T \mathcal{X}$ as the left and right searching subspaces for the left and right singular vectors u_* and z_* of \check{A} , respectively. We note that $\widetilde{Z} = L^T X$ is a basis matrix of $\mathcal{Z} = L^T \mathcal{X}$. Then the CPF-harmonic extraction [11] finds positive scalars $\phi > 0$ and vectors $\check{u} \in \mathcal{U}$ and $\check{z} \in \mathcal{Z}$ such that

(2.17)
$$\begin{bmatrix} 0 & \check{A}^T \\ \check{A} & 0 \end{bmatrix} \begin{bmatrix} \check{z} \\ \check{u} \end{bmatrix} - \phi \begin{bmatrix} \check{z} \\ \check{u} \end{bmatrix} \perp \left(\begin{bmatrix} 0 & \check{A}^T \\ \check{A} & 0 \end{bmatrix} - \tau I \right) \cdot \mathcal{R} \left(\begin{bmatrix} \widetilde{Z} & \\ & \widetilde{U} \end{bmatrix} \right),$$

where $\mathcal{R}(\cdot)$ denotes the range space of a matrix. This is the harmonic extraction approach for the eigenvalue problem of the augmented matrix $\begin{bmatrix} \check{A}^{T} \end{bmatrix}$ with respect to the searching subspace $\mathcal{R}\left(\begin{bmatrix} \tilde{Z} \\ \tilde{U} \end{bmatrix}\right)$ and the given target $\tau > 0$; see [33, 36].

Write $\check{z} = \widetilde{Z}\check{d}$ and $\check{u} = \widetilde{U}\check{e}$. It is shown in [11] that (2.17) amounts to the following symmetric generalized eigenvalue problem:

$$(2.18) \begin{bmatrix} R_A^T R_A + \tau^2 R_B^T R_B & -2\tau R_A^T \\ -2\tau R_A & \widetilde{U}^T A (B^T B)^{-1} A^T \widetilde{U} + \tau^2 I \end{bmatrix} \begin{bmatrix} \check{d} \\ \check{e} \end{bmatrix} = (\phi - \tau) \begin{bmatrix} -\tau R_B^T R_B & R_A^T \\ R_A & -\tau I \end{bmatrix} \begin{bmatrix} \check{d} \\ \check{e} \end{bmatrix}.$$

Denote by H_c and G_c the $2k \times 2k$ symmetric matrices in the left and right hand sides of the above equation, respectively. Computationally, suppose that $(B^TB)^{-1} = (LL^T)^{-1}$ can be efficiently applied to obtain H_c . The CPF-harmonic extraction approach computes the generalized eigendecomposition of the symmetric positive definite matrix pair (G_c, H_c) , picks up the largest generalized eigenvalue ν in magnitude and the corresponding eigenvector $\begin{bmatrix} d\\ e \end{bmatrix}$, and takes

(2.19)
$$(\phi, \check{u}, \check{z}) = \left(\tau + \frac{1}{\mu}, \frac{\widetilde{U}\check{e}}{\|\check{e}\|}, \frac{\widetilde{Z}\check{d}}{\|\widetilde{Z}\check{d}\|}\right)$$

as an approximation to the singular triplet (σ_*, u_*, z_*) of A.

Note that the exact right generalized singular vector $x_* = \beta_* L^{-T} z_*$. Therefore, we take $L^{-T} \check{z} = L^{-T} \widetilde{Z} \check{d} = \widetilde{X} \check{d}$ as the approximation to x_* in direction. Concretely, we take the approximate right generalized singular vector \check{x} of (A, B) as

(2.20)
$$\check{x} = \frac{1}{\check{\delta}} \widetilde{X} \check{d} \quad \text{with} \quad \check{\delta} = \sqrt{\|\check{e}\|^2 + \|\check{f}\|^2},$$

where \check{e} is recomputed by $\check{e} = R_A \check{d}$ and $\check{f} = R_B \check{d}$. With such $\check{\delta}$, the approximate \check{x} is of $(A^T A + B^T B)$ -norm unit length [11]. We then take the new approximate generalized singular value and left generalized singular vectors as

(2.21)
$$\check{\alpha} = \frac{\|\check{e}\|}{\check{\delta}}, \quad \check{\beta} = \frac{\|\check{f}\|}{\check{\delta}} \quad \text{and} \quad \check{u} = \frac{U\check{e}}{\|\check{e}\|}, \quad \check{v} = \frac{V\check{f}}{\|\check{f}\|},$$

which are called the CPF-harmonic Ritz approximations and satisfy $A\check{x} = \check{\alpha}\check{u}$ and $B\check{x} = \check{\beta}\check{v}$ with $\check{\alpha}^2 + \check{\beta}^2 = ||\check{u}|| = ||\check{v}|| = 1$. Moreover, it is known from [11] that the new $\check{\theta} = \frac{\check{\alpha}}{\check{\beta}}$ is a better approximation to σ_* than ϕ in (2.19) in the sense that

(2.22)
$$\| (A^T A - \check{\theta}^2 B^T B) \check{x} \|_{(B^T B)^{-1}} \le \| (A^T A - \phi^2 B^T B) \check{x} \|_{(B^T B)^{-1}},$$

where $\|\cdot\|_M$ is the *M*-norm for a symmetric positive definite matrix *M*.

2.4. The IF-harmonic extraction approach. For a general and possibly rank deficient B, the CPF-harmonic extraction does not work. Alternatively, the IF-harmonic extraction [11] is for more general purpose. It finds approximate generalized singular values $\varphi > 0$ and approximate right generalized singular vectors $\hat{x} \in \mathcal{X}$ with $\|\hat{x}\|_{A^T A + B^T B} = 1$ such that

(2.23)
$$(A^T A - \varphi^2 B^T B) \hat{x} \perp (A^T A - \tau^2 B^T B) \mathcal{X}.$$

This is precisely the harmonic Rayleigh–Ritz projection on the generalized eigenvalue problem of $(A^T A, B^T B)$ with respect to the subspace \mathcal{X} and the given target τ^2 .

Write $\hat{x} = \frac{1}{\hat{\delta}} \tilde{X} \hat{d}$ with $\|\hat{d}\| = 1$ and $\hat{\delta}$ a normalizing parameter to be determined. Then requirement (2.23) is equivalent to

(2.24)
$$\widetilde{X}^T (A^T A - \tau^2 B^T B)^2 \widetilde{X} \hat{d} = (\varphi^2 - \tau^2) \widetilde{X}^T (A^T A - \tau^2 B^T B) B^T B \widetilde{X} \hat{d}.$$

Denote by H_{τ} and G_{τ} the k-by-k matrices in the left and right hand sides of the above equation, respectively. Then $(\varphi^2 - \tau^2)$ is a generalized eigenvalue of the matrix pair (H_{τ}, G_{τ}) with \hat{d} the corresponding unit length generalized eigenvector. The IF-harmonic extraction approach computes the generalized eigendecomposition of (G_{τ}, H_{τ}) , and picks up the generalized eigenpair (ν, \hat{d}) corresponding to $\varphi = \sqrt{\tau^2 + \frac{1}{\nu}}$ closest to τ among all the eigenpairs of (G_{τ}, H_{τ}) . Then φ is the IF-harmonic Ritz value that approximates the desired σ_* and

(2.25)
$$\hat{x} = \frac{1}{\hat{\delta}} \widetilde{X} \hat{d} \quad \text{with} \quad \hat{\delta} = \sqrt{\|\hat{e}\|^2 + \|\hat{f}\|^2}$$

is the corresponding right IF-harmonic Ritz vector that approximates the desired x_* , where $\hat{e} = R_A \hat{d}$ and $\hat{f} = R_B \hat{d}$. Such $\hat{\delta}$ guarantees that \hat{x} is of $(A^T A + B^T B)$ -norm

unit length. The corresponding IF-harmonic Ritz values and left IF-harmonic Ritz vectors are computed by

(2.26)
$$\hat{\alpha} = \frac{\|\hat{e}\|}{\hat{\delta}}, \quad \hat{\beta} = \frac{\|\hat{f}\|}{\hat{\delta}} \quad \text{and} \quad \hat{u} = \frac{\widetilde{U}\hat{e}}{\|\hat{e}\|}, \quad \hat{v} = \frac{\widetilde{V}\hat{f}}{\|\hat{f}\|}$$

It is known from [11] that the IF-harmonic Ritz approximation $(\hat{\alpha}, \hat{\beta}, \hat{u}, \hat{v}, \hat{x})$ satisfies $A\hat{x} = \hat{\alpha}\hat{u}$ and $B\hat{x} = \hat{\beta}\hat{v}$ with $\hat{\alpha}^2 + \hat{\beta}^2 = \|\hat{u}\| = \|\hat{v}\| = 1$. The IF-harmonic approximate generalized singular value $\hat{\theta} = \frac{\hat{\alpha}}{\hat{\beta}}$ is better than the above φ in the sense of (2.22) where $\phi, \check{\theta}$ and \check{x} are replaced by $\varphi, \hat{\theta}$ and \hat{x} , respectively.

3. The refined JDGSVD type methods. For an approximation to the desired GSVD component $(\alpha_*, \beta_*, u_*, v_*, x_*)$ obtained by the standard or harmonic extraction approaches described in Section 2, in this section we will unify the notation and denote the approximation by (α, β, u, v, x) with $\theta = \frac{\alpha}{\beta}$. We now propose a refined extraction approach: Find a unit length vector $x_r \in \mathcal{X}$ satisfying the optimal requirement

(3.1)
$$\| (A^T A - \theta^2 B^T B) x_r \| = \min_{w \in \mathcal{X}, \|w\| = 1} \| (A^T A - \theta^2 B^T B) w \|,$$

rescale x_r to \bar{x} with $\|\bar{x}\|_{A^TA+B^TB} = 1$, and use \bar{x} to approximate x_* . We call \bar{x} a refined approximate right generalized singular vector, or simply the refined or refined harmonic right Ritz vector, of (A, B) over the subspace \mathcal{X} if θ is a Ritz value or harmonic Ritz value obtained by the standard or harmonic extraction approaches.

Jia [16] has proven that if (θ^2, \bar{x}) is not an exact eigenpair of $(A^T A, B^T B)$ then $||(A^T A - \theta^2 B^T B)\bar{x}|| < ||(A^T A - \theta^2 B^T B)x||$ strictly; moreover, if there is another standard or harmonic Ritz value close to θ , then

$$\|(A^T A - \theta^2 B^T B)\bar{x}\| \ll \|(A^T A - \theta^2 B^T B)x\|$$

generally holds, meaning that \bar{x} may be much more accurate than x as an approximation to x_* . Most importantly, the fundamental convergence results in [17, 23] indicate that there is a Ritz value $\theta \to \sigma_*$ unconditionally and the convergence of the refined Ritz or harmonic Ritz vector $\bar{x} \to x_*$ is guaranteed too once the distance between x_* and \mathcal{X} tends to zero, while the Ritz or (CPF or IF-)harmonic Ritz vector x may converge erratically and even may fail to converge even if \mathcal{X} contains sufficiently accurate approximations to x_* .

We now consider the accurate and efficient computation of \bar{x} . For an arbitrary unit length vector $w \in \mathcal{X}$, write $w = \tilde{X}d$ with ||d|| = 1. Then $x_r = \tilde{X}\bar{d}$ with $||\bar{d}|| = 1$, and the minimization problem (3.1) is equivalent to

(3.2)
$$\| (A^T A - \theta^2 B^T B) \widetilde{X} \overline{d} \| = \min_{d \in \mathbb{R}^k, \|d\| = 1} \| (A^T A - \theta^2 B^T B) \widetilde{X} d \|.$$

Therefore, \overline{d} is the right singular vector of $G_{\theta} = (A^T A - \theta^2 B^T B) \widetilde{X}$ corresponding to its smallest singular value, and it is also the eigenvector of the cross-product matrix

(3.3)
$$H_{\theta} = G_{\theta}^T G_{\theta} = \widetilde{X}^T (A^T A - \theta^2 B^T B)^2 \widetilde{X}$$

associated with its smallest eigenvalue. Jia [18] has proposed and developed a variant of the cross product-based QR algorithm for the accurate SVD computation of a general matrix. In our specific context, we only need to use the standard QR algorithm to obtain \overline{d} . Remarkably, Jia [18] has shown that the cross product-based QR algorithm is much more efficient than the standard Golub–Kahan and Chan SVD algorithms [4, 33] applied to G_{θ} for $n \gg k$.

More precisely, in finite precision arithmetic, Jia [18] has proven that, provided that the smallest singular value of G_{θ} is well separated from its second smallest one then the computed \bar{d} 's by the cross product-based QR algorithm and the Golub– Kahan or Chan SVD algorithm essentially have the same accuracy; when the computed smallest singular value of G_{θ} is taken as the square root of the Rayleigh quotient $x^T H_{\theta} x$ that is calculated by the formula $(G_{\theta} \bar{d})^T (G_{\theta} \bar{d})$ with \bar{d} the computed eigenvector of H_{θ} , it has the same accuracy as the smallest singular value computed by the standard Golub–Kahan or Chan SVD algorithm applied to G_{θ} ; see [18] for a detailed analysis and comparison. We should particularly remind that in our context the smallest singular value of G_{θ} tends to zero as $\theta \to \sigma_*$ but the second smallest one of G_{θ} is typically not small and thus is well separated from the smallest one.

In view of the above, rather than computing the SVD of G_{θ} at expensive cost, we compute the eigendecomposition of H_{θ} cheaply, and pick up the desired eigenvector \bar{d} . We remark that, as the subspaces are expanded, we can efficiently form H_{θ} by

(3.4)
$$H_{\theta} = H_A + \theta^4 H_B - \theta^2 (H_{A,B}^T + H_{A,B}),$$

where the intermediate matrices

(3.5)
$$H_A = \widetilde{X}^T (A^T A)^2 \widetilde{X}, \qquad H_B = \widetilde{X}^T (B^T B)^2 \widetilde{X}, \qquad H_{A,B} = \widetilde{X}^T A^T A B^T B \widetilde{X}$$

can be efficiently updated at each step and they are also used to efficiently form the projected matrices $G_{\tau} = H_{A,B} - \tau^2 H_B$ and $H_{\tau} = H_A + \tau^4 H_B - \tau^2 (H_{A,B}^T + H_{A,B})$ involved in the IF-harmonic extraction approach; see (2.24). Also, it is important to notice that when ℓ GSVD components are required, which will be considered in the next section, we can efficiently form all the cross-product matrices H_{θ} in (3.4) for different θ 's.

By definition, we need to rescale x_r in (3.1) to obtain the refined or refined harmonic Ritz vector \bar{x} with $\|\bar{x}\|_{A^TA+B^TB} = 1$. Write

$$\bar{x} = \frac{1}{\bar{\delta}} \widetilde{X} \bar{d},$$

where $\bar{\delta}$ is a normalizing factor to be determined. Following the same derivations as in Section 2, we have

$$\bar{\delta} = \sqrt{\|\bar{e}\|^2 + \|\bar{f}\|}$$
 with $\bar{e} = R_A \bar{d}$ and $\bar{f} = R_B \bar{d}$,

where R_A and R_B are defined in (2.2). Analogously to those done in the CPF- and IF-harmonic extraction approaches, we compute the refined or refined harmonic Ritz values and the two refined or refined harmonic left Ritz vectors by

(3.6)
$$\bar{\alpha} = \frac{\|\bar{e}\|}{\bar{\delta}}, \quad \bar{\beta} = \frac{\|\bar{f}\|}{\bar{\delta}} \quad \text{and} \quad \bar{u} = \frac{\widetilde{U}\bar{e}}{\|\bar{e}\|}, \quad \bar{v} = \frac{\widetilde{V}\bar{f}}{\|\bar{f}\|}.$$

It is easily verified that

$$A\bar{x} = \bar{\alpha}\bar{u}$$
 and $B\bar{x} = \beta\bar{v}$

with $\bar{\alpha}^2 + \bar{\beta}^2 = \|\bar{u}\| = \|\bar{v}\| = 1$. The quintuple $(\bar{\alpha}, \bar{\beta}, \bar{u}, \bar{v}, \bar{x})$ is called a refined or refined CPF-harmonic or refined IF-harmonic approximation to the desired GSVD component $(\alpha_*, \beta_*, u_*, v_*, x_*)$ of (A, B), depending on by which extraction approach θ in (3.1) is computed: the standard extraction, the CPF-harmonic or IF-harmonic extraction. Particularly, $(\bar{\alpha}, \bar{\beta})$ or $\bar{\theta} = \frac{\bar{\alpha}}{\bar{\beta}}$ is called the refined, refined CPF-harmonic, or refined IF-harmonic Ritz value and, correspondingly, \bar{u}, \bar{v} and \bar{x} are called the refined, refined CPF-harmonic or IF-harmonic left and right Ritz vectors of (A, B).

Combining the refined, refined CPF-harmonic and refined IF-harmonic extraction approaches with the subspace expansion approach described in Section 2.1, we have now proposed the refined CPF, refined CPF-harmonic and refined IF-harmonic JDGSVD methods, i.e., RCPF-JDGSVD, RCPF-HJDGSVD and RIF-HJDGSVD, respectively. For each of these methods, the corresponding residual and stopping criterion are the same as (2.3) and (2.4).

4. Thick-restart refined JDGSVD algorithms with deflation and purgation. In this section, by introducing appropriate deflation and purgation techniques, we develop practical thick-restart refined JDGSVD algorithms for solving Problem 1.1.

4.1. Thick-restart. As the dimension k of searching subspaces increases, the storage requirements and computational costs of the previous basic JDGSVD type algorithms become unaffordable. When k reaches the maximum number k_{max} allowed but the algorithms do not yet converge, it is necessary to restart them. To this end, we adopt the thick-restart technique, which was first advocated in [32] for the eigenvalue problem and has been nontrivially extended to the SVD and GSVD problems in [9, 11, 12, 32, 38, 39]. Specifically, the thick-restart takes certain k_{\min} -dimensional subspaces of the current left and right searching subspaces as the initial left and right ones for the next cycle, so that they retain as much information as possible on the desired and $k_{\min} - 1$ nearby GSVD components of (A, B). We then expand them step by step in the way described in Section 2, compute new approximate GSVD components with respect to the expanded subspaces at each step, and check the convergence. If converged, we stop; otherwise, we repeat the same process until the dimension of expanded subspaces reaches k_{\max} .

We will present an efficient and stable computational procedure for the thickrestart. For each of RCPF-JDGSVD, RCPF-HJDGSVD and RIF-HJDGSVD, when $k = k_{\text{max}}$ but the method does not yet converge, we compute k_{min} approximate right generalized singular vectors $\bar{x}_i = \tilde{X}\bar{d}_i$, $i = 1, \ldots, k_{\text{min}}$ associated with the k_{min} approximate singular values closest to τ , where \bar{x}_1 is selected as the approximation to the desired x_* . Then the thick-restart takes the new initial right searching subspace $\mathcal{X}_{\text{new}} = \text{span}\{\bar{x}_1, \ldots, \bar{x}_{\min}\}.$

Denote $D_1 = [\bar{d}_1, \ldots, \bar{d}_{k_{\min}}]$. Next we show how to efficiently obtain the k_{\min} -dimensional new right and left subspaces \mathcal{X}_{new} , \mathcal{U}_{new} and \mathcal{V}_{new} as well as their orthonormal bases, whose computational details were not described in [11, 12].

Compute the thin QR factorization $D_1 = Q_d R_d$ using $\mathcal{O}(k_{\max} k_{\min}^2)$ flops. Then

(4.1)
$$\widetilde{X}_{\text{new}} = \widetilde{X}Q_d$$

is the orthonormal basis matrix of \mathcal{X}_{new} , whose computation costs $2nk_{\max}k_{\min}$ flops. As for the new $\mathcal{U}_{\text{new}} = A\mathcal{X}_{\text{new}}$ and $\mathcal{V}_{\text{new}} = B\mathcal{X}_{\text{new}}$, we compute the thin QR factorizations of the small sized matrices

(4.2)
$$R_A Q_d = Q_e R_{A,\text{new}}$$
 and $R_B Q_d = Q_f R_{B,\text{new}}$

using $\mathcal{O}(k_{\max}^2 k_{\min})$ flops, where R_A , R_B are defined as in (2.2). Exploiting (2.2) and (4.1)–(4.2), we obtain the thin QR factorizations

$$A\widetilde{X}_{\text{new}} = A\widetilde{X}Q_d = \widetilde{U}R_AQ_d = (\widetilde{U}Q_e)R_{A,\text{new}},$$

$$B\widetilde{X}_{\text{new}} = B\widetilde{X}Q_d = \widetilde{V}R_BQ_d = (\widetilde{V}Q_f)R_{B,\text{new}},$$

showing that the columns of

$$\widetilde{U}_{\text{new}} = \widetilde{U}Q_e$$
 and $\widetilde{V}_{\text{new}} = \widetilde{V}Q_f$

form orthonormal bases of \mathcal{U}_{new} and \mathcal{V}_{new} , respectively, whose computation costs $2(m+p)k_{\max}k_{\min}$ flops. Together with the computation of $\widetilde{X}_{\text{new}}$, the above whole process approximately costs $2(m+n+p)k_{\max}k_{\min}$ flops since $m, p, n \gg k_{\max} > k_{\min}$.

For the intermediate matrices in (3.5) used to form H_{θ} defined by (3.4), as has been done in the IF-harmonic JDGSVD algorithm, we efficiently update them by

(4.3) $H_{A,\text{new}} = Q_d^T H_A Q_d, \qquad H_{B,\text{new}} = Q_d^T H_B Q_d, \qquad H_{A,B,\text{new}} = Q_d^T H_{A,B} Q_d$

at cost of $\mathcal{O}(k_{\max}^2 k_{\min})$ flops.

Summarizing the above, the construction cost of orthonormal bases of the k_{\min} dimensional \mathcal{U}_{new} , \mathcal{V}_{new} , \mathcal{X}_{new} and the matrices in (4.3) is approximately $2(m + p + n)k_{\max}k_{\min}$ flops. Typically, in computation, one takes $k_{\max} = 20 \sim 30$ and $k_{\min} = 3 \sim 5$; see [11, 12]. Therefore, forming the restarting initial left and right searching subspaces in the thick-restart is very cheap.

4.2. Deflation and purgation. We can adapt the effective and efficient deflation techniques proposed in [11, 12] to the thick-restart RCPF-JDGSVD, RCPF-HJDGSVD and RIF-HJDGSVD. We will elaborate some key elements of subspaces during deflation that were not given adequate and very clear arguments in [11, 12], which turn out to play a crucial role in both mathematics and effective and efficient implementations of JDGSVD type algorithms for computing more than one GSVD components.

Suppose that one of the RCPF-JDGSVD, RCPF-HJDGSVD and RIF-HJDGSVD algorithms has computed $j(< \ell)$ converged approximations $(\alpha_{i,c}, \beta_{i,c}, u_{i,c}, v_{i,c}, x_{i,c})$ to $(\alpha_i, \beta_i, u_i, v_i, x_i)$, whose residual norms satisfy

$$(4.4) \quad \|r_i\| = \|\beta_{i,c}A^T u_{i,c} - \alpha_{i,c}B^T v_{i,c}\| \le (\beta_{i,c}\|A\|_1 + \alpha_{i,c}\|B\|_1) \cdot tol, \quad i = 1, \dots, j.$$

Denote

(4.5)

$$U_{c} = [u_{1,c}, \dots, u_{j,c}], \qquad C_{c} = \text{diag}\{\alpha_{1,c}, \dots, \alpha_{j,c}\}, \\ V_{c} = [v_{1,c}, \dots, v_{j,c}], \qquad S_{c} = \text{diag}\{\beta_{1,c}, \dots, \beta_{j,c}\}, \\ Y_{c} = [x_{1,c}, \dots, x_{j,c}], \qquad Y_{c} = (A^{T}A + B^{T}B)X_{c}.$$

Then

$$AX_{c} = U_{c}C_{c}, \quad BX_{c} = V_{c}S_{c}, \quad C_{c}^{2} + S_{c}^{2} = I_{j}, \quad Y_{c} = A^{T}U_{c}C_{c} + B^{T}V_{c}S_{c},$$

and the F-norm of the residual matrix satisfies

$$||R_c||_F = ||A^T U_c S_c - B^T V_c C_c||_F \le \sqrt{j(||A||_1^2 + ||B||_1^2)} \cdot tol$$

Suppose that the current X_c and Y_c are bi-orthogonal, i.e., $Y_c^T X_c = I_j$. We point out that this bi-orthogonality is fulfilled in our six JDGSVD type algorithms. It is known from Proposition 4.1 of [12] that $(\alpha_i, \beta_i, u_i, v_i, x_i)$, $i = j + 1, \ldots, q$ are the exact GSVD components of the deflated matrix pair

$$(4.6) \qquad (A(I - X_c Y_c^T), B(I - X_c Y_c^T))$$

restricted to the range space of the oblique projector $(I - X_c Y_c^T)$ if tol = 0 in (4.4). Therefore, we can apply any one of RCPF-JDGSVD, RCPF-HJDGSVD and RIF-HJDGSVD to the deflated matrix pair in (4.6) to compute the next desired GSVD component $(\alpha_*, \beta_*, u_*, v_*, x_*) := (\alpha_{j+1}, \beta_{j+1}, u_{j+1}, v_{j+1}, x_{j+1})$ of (A, B).

Remarkably, when the converged $(\alpha_{j,c}, \beta_{j,c}, u_{j,c}, v_{j,c}, x_{j,c})$ has been found, the current subspaces usually contain reasonably rich information on u_*, v_*, x_* . To make full use of such available information, we present an effective and efficient purgation technique in the thick-restart RCPF-JDGSVD, RCPF-HJDGSVD and RIF-HJDGSVD algorithms. Instead of constructing initial searching subspaces from scratch, we purge the newly converged

$$x_{i,c} := x = \widetilde{X}d$$

from the current \mathcal{X} , and take the reduced subspace, denoted by \mathcal{X}_{new} , as an initial right searching subspace when extracting an approximation to $(\alpha_*, \beta_*, u_*, v_*, x_*)$.

Concretely, denote

$$\widetilde{X}_{\text{new}} = \widetilde{X}Q_D$$

with some orthonormal matrix $Q_D \in \mathbb{R}^{k \times (k-1)}$ to be determined. We require that $\widetilde{X}_{\text{new}}$ be orthogonal to $Y_{c,\text{new}} = [Y_c, y]$, where $y = (A^T A + B^T B)x$. Suppose that the current \widetilde{X} is orthogonal to Y_c . Then we only need to make $\widetilde{X}_{\text{new}}$ orthogonal to y. By (2.2), this amounts to

$$\widetilde{X}_{\text{new}}^T y = Q_D^T \widetilde{X}^T (A^T A + B^T B) \widetilde{X} d = Q_D^T (R_A^T R_A + R_B^T R_B) d = \mathbf{0}.$$

Therefore, the columns of Q_D form an orthonormal basis of the orthogonal complement of $\mathcal{R}(d')$ with respect to \mathbb{R}^k , where $d' = (R_A^T R_A + R_B^T R_B)d$. In computation, we compute the full QR factorization of the k-by-1 matrix d' using approximately $4k^2$ flops [4, p.249-250], take the second to last columns of its Q-factor to form Q_D , and obtain the desired \mathcal{X}_{new} . Then following the analogous process to those in Section 4.1, by replacing Q_d with Q_D , we can carry out this purgation strategy to obtain the reduced \mathcal{U}_{new} and \mathcal{V}_{new} with little extra cost. We then extract an approximation to $(\alpha_*, \beta_*, u_*, v_*, x_*)$ with respect to them, and expand the searching subspaces in the thick-restart way described previously.

We now unify to denote by (α, β, u, v, x) the approximate GSVD component obtained by one of RCPF-JDGSVD, RCPF-HJDGSVD and RIF-HJDGSVD with respect to the left and right searching subspaces \mathcal{U}, \mathcal{V} and \mathcal{X} . If the current \mathcal{X} is orthogonal to $\mathcal{R}(Y_c)$, then

$$\mathcal{U} = (A - X_c Y_c^T) \mathcal{X} = A \mathcal{X} \text{ and } \mathcal{V} = (B - X_c Y_c^T) \mathcal{X} = B \mathcal{X}.$$

Therefore, for such an \mathcal{X} , in the JDGSVD type algorithms of [11, 12] and this paper, computationally, we never work on the matrix pair in (4.6) explicitly, instead we

always work on (A, B) directly. The gains are twofold: the resulting JDGSVD type algorithms are more efficient at each step; in finite precision arithmetic, they enable us to compute the approximations more accurately. The first point is straightforward. But the second point is subtle and quite complicated, and its arguments and details are out of the scope of this paper.

Recall that the reduced \mathcal{X}_{new} can be made orthogonal to $\mathcal{R}(Y_c)$ and is an instance of the current subspace \mathcal{X} when computing $(\alpha_*, \beta_*, u_*, v_*, x_*)$. We present the following important result.

Theorem 4.1. Suppose that the current right subspace \mathcal{X} is orthogonal to $\mathcal{R}(Y_c)$. Then the expanded \mathcal{X} 's are also orthogonal to $\mathcal{R}(Y_c)$ at subsequent expansion steps.

Proof. We only need to prove the assertion for one expansion step. For the current \mathcal{X} , if (α, β, u, v, x) does not converge yet, then in the expansion phase, the original correction equation (2.6) for $\ell > 1$ becomes (cf. [11, 12])

(4.7)
$$(I - Y_p X_p^T) (A^T A - \rho^2 B^T B) (I - X_p Y_p^T) t = -(I - Y_c X_c^T) r$$
 for $t \perp Y_p$,

where r is the residual of (α, β, u, v, x) defined by (2.3), $\rho = \tau$ or $\theta = \frac{\alpha}{\beta}$ (see (2.6) and the paragraph that follows), and $X_p = [X_c, x]$, $Y_p = [Y_c, y]$ with X_c and Y_c defined in (4.5) and y defined by (2.5). It follows from $x \in \mathcal{X} \perp \mathcal{R}(Y_c)$ that X_p and Y_p are bi-orthogonal: $Y_p^T X_p = I_{j+1}$, meaning that $I - Y_p X_p^T$ and $I - X_p Y_p^T$ are oblique projectors. With an approximate solution t of (4.7) found, we orthonormalize it against the orthonormal basis matrix \widetilde{X} of \mathcal{X} to obtain the expansion vector x_+ and update the basis matrices \widetilde{X} , \widetilde{U} and \widetilde{V} by (2.9)–(2.13). By the way that \widetilde{X} is augmented, the resulting expanded \mathcal{X} is automatically orthogonal to $\mathcal{R}(Y_c)$ at the expansion step.

The assertion in this theorem is crucial and ensures that we always work on (A, B) rather than the explicitly deflated matrix pair in (4.6) when using each of the six JDGSVD type algorithms to computing more than one GSVD components.

Once (α, β, u, v, x) has converged, we add it to the previous converged partial GSVD $(C_c, S_c, U_c, V_c, X_c)$ of (A, B), and set j := j + 1. The RCPF-JDGSVD, RCPF-HJDGSVD and RIF-HJDGSVD algorithms proceed in this way until all the ℓ desired GSVD components of (A, B) are found, and the computed $X_c = [x_{1,c}, \ldots, x_{\ell,c}]$ and $Y_c = [y_{1,c}, \ldots, y_{\ell,c}]$ satisfy $Y_c^T X_c = I_\ell$.

4.3. Thick restart refined and refined harmonic JDGSVD algorithms. Algorithm 1 summarizes the thick-restart RCPF-JDGSVD, RCPF-HJDGSVD and RIF-HJDGSVD algorithms with deflation and purgation for computing a partial GSVD (C_c, S_c, U_c, V_c, X_c) of (A, B) with the ℓ generalized singular values closest to a given target τ . Each of these three algorithms demands the modules to perform matrix-vector multiplications with A, B and A^T, B^T , the target $\tau > 0$, the number ℓ of desired GSVD components, a unit-length vector x_0 for the initial right searching subspace, and the stopping tolerance tol for the outer iterations. RCPF-HJDGSVD needs a device to act $(B^TB)^{-1}$ so as to form and update the projection matrix H_c in (2.18). Other parameters include the minimum and maximum dimensions k_{\min} and k_{\max} of searching subspaces, the switching tolerance fixtol > 0 for the correction equations (2.6) and (4.7) with the fixed shift $\rho = \tau$ to their counterparts with the adaptively changing shift $\rho = \theta$, and the stopping tolerance $\tilde{\varepsilon} > 0$ in (2.8) for the inner iterations. By default, we set them as 3, 30, 10^{-4} and 10^{-4} , respectively. Algorithm 1 The thick-restart RCPF-JDGSVD, RCPF-HJDGSVD, RIF-HJDGSVD algorithms with deflation and purgation for the target τ .

- 1: **Initialization:** Set k = 1, $k_c = 0$, and initialize $C_c = [], S_c = [], U_c = [], V_c = [], V_c = [], X_c = [], Y_c = []$. Set $\tilde{U} = [], \tilde{V} = [], \tilde{X} = []$, and choose a unit-length starting vector $x_+ = x_0$.
- 2: while $k \ge 0$ do
- 3: Set $\widetilde{X} = [\widetilde{X}, x_+]$, and update the QR factorizations $A\widetilde{X} = \widetilde{U}R_A, B\widetilde{X} = \widetilde{V}R_B$.
- 4: **RCPF-JDGSVD:** Compute the GSVD of (R_A, R_B) , and pick up the generalized singular value θ closest to the target τ . **RCPF-HJDGSVD:** Form H_c and G_c defined in (2.18). Compute the general-

ized eigendecomposition of (G_c, H_c) , pick up the eigenvector d corresponding to the largest eigenvalue μ in magnitude, and compute the approximate $\theta = \frac{\|R_A d\|}{\|R_B d\|}$. **RIF-HJDGSVD:** Form G_{τ} and H_{τ} defined in (2.24). Compute the generalized eigendecomposition of (G_{τ}, H_{τ}) , pick up the eigenvector d associated with the eigenvalue ν such that $\sqrt{\tau^2 + \frac{1}{\nu}}$ is closest to τ , and compute $\theta = \frac{\|R_A d\|}{\|R_B d\|}$.

- 5: Form H_{θ} defined by (3.4), compute the eigendecomposition of H_{θ} , and pick up the eigenvector \bar{d} corresponding to the smallest eigenvalue.
- 6: Form $\bar{e} = R_A \bar{d}$, $\bar{f} = \hat{R}_B \bar{d}$, and calculate $\bar{\delta} = \sqrt{\|\bar{e}\|^2 + \|\bar{f}\|^2}$. Compute the approximate GSVD component $(\bar{\alpha}, \bar{\beta}, \bar{u}, \bar{v}, \bar{x}) = \left(\frac{\|\bar{e}\|}{\bar{\delta}}, \frac{\|\bar{f}\|}{\bar{\delta}}, \frac{1}{\bar{\delta}}\tilde{X}\bar{d}, \frac{\tilde{U}\bar{e}}{\|\bar{e}\|}, \frac{\tilde{V}\bar{f}}{\|\bar{f}\|}\right)$, the vector $\bar{y} = \bar{\alpha}A^T\bar{u} + \bar{\beta}B^T\bar{v}$, and the residual $r = \bar{\beta}A^T\bar{u} - \bar{\alpha}B^T\bar{v}$.
- 7: **if** $||r|| \le (\bar{\beta} ||A||_1 + \bar{\alpha} ||B||_1) \cdot tol$ **then**
- 8: Update the matrices $C_c = \text{diag}\{C_c, \bar{\alpha}\}, S_c = \text{diag}\{S_c, \bar{\beta}\}, U_c = [U_c, \bar{u}], V_c = [V_c, \bar{v}], X_c = [X_c, \bar{x}], Y_c = [Y_c, \bar{y}], \text{ and set } k_c = k_c + 1.$
- 9: **if** $k_c = \ell$ **then** return $(C_c, S_c, U_c, V_c, X_c)$ and stop. **fi**
- 10: Purge $\bar{x}, \bar{u}, \bar{v}$ from $\mathcal{X}, \mathcal{U}, \mathcal{V}$, update the orthonormal basis matrices $\widetilde{X}, \widetilde{U}, \widetilde{V}$ and projection matrices R_A, R_B . Set k = k - 1, and go to step 4.
- 11: end if
- 12: Form $X_p = [X_c, \bar{x}]$ and $Y_p = [Y_c, \bar{y}]$. Iteratively solve the correction equation

$$(I - Y_p X_p^T)(A^T A - \rho^2 B^T B)(I - X_p Y_p^T)t = -(I - Y_c X_c^T)r \quad \text{for} \quad t \perp Y_p$$

with $\rho = \tau$ or $\rho = \bar{\alpha}/\bar{\beta}$ until the relative residual norm fulfills (2.8).

13: if $k = k_{\text{max}}$ then perform the thick-restart, and set $k = k_{\text{min}}$. fi

The authors in [11] have made a detailed analysis on the cost of the thick-restart CPF-HJGSVD and IF-HJDGSVD algorithms, which straightforwardly applies to Algorithm 1. The conclusion is that, for $k_{\max} \ll n$, the cost of each algorithm is dominated by the matrix-vector multiplications with A, B and A^T, B^T , assuming that the MINRES method is used to solve all the correction equations. We remind that, for the thick-restart CPF-HJGSVD and RCPF-HJGSVD, this conclusion requires that each linear system with the coefficient matrix $B^T B$ can be solved efficiently at cost of $\mathcal{O}(n)$ flops.

5. Numerical experiments. We illustrate the performance of thick-restart RCPF-JDGSVD, RCPF-HJDGSVD and RIF-HJDGSVD algorithms on several prob-

^{14:} Orthonormalize t against X to obtain the expansion vector x_+ , and set k = k+1. 15: end while

lems, and compare them with the standard and harmonic JDGSVD algorithms [11, 12]: CPF-JDGSVD, CPF-HJDGSVD and IF-HJDGSVD, showing the considerable advantage of the refined and refined harmonic algorithms over their respective unrefined counterparts. All the numerical experiments were implemented on a 13th Gen Intel (R) Core (TM) i9-13900KF CPU 3.00 GHz with 64 GB RAM using the MAT-LAB R2022b with the machine precision $\epsilon_{\rm mach} = 2.22 \times 10^{-16}$ under the Miscrosoft Windows 11 Pro 64-bit system.

 $\begin{array}{c} {\rm TABLE \ 1} \\ {\rm Test \ matrix \ pairs \ with \ their \ basic \ properties.} \end{array}$

| A | В | m | p | n | nnz | $\kappa\bigl(\bigl[\begin{smallmatrix} A\\B \end{smallmatrix}\bigr]\bigr)$ | $\sigma_{ m max}$ | $\sigma_{ m min}$ |
|----------------------------|---|-------|-------|-------|---------|--|-------------------|-------------------|
| $dano3mip^T$ | T | 15851 | 3202 | 3202 | 91237 | 1.81e + 3 | 1.28e + 3 | 1.00e-16 |
| $\mathrm{plddb}^{\bar{T}}$ | T | 5049 | 3069 | 3069 | 20044 | 91.9 | 61.2 | 3.65e-3 |
| barth | T | 6691 | 6691 | 6691 | 46510 | 5.56 | 3.16 | 7.61e-19 |
| $large^{T}$ | T | 8617 | 4282 | 4282 | 33479 | 3.53e + 3 | 2.42e + 3 | 2.25e-3 |
| ns3Da | T | 20414 | 20414 | 20414 | 1740839 | 5.01 | 4.02e-1 | 1.86e-4 |
| e40r0100 | T | 17281 | 17281 | 17281 | 605403 | 11.2 | 9.56 | 2.62e-8 |
| rat^T | T | 9408 | 3136 | 3136 | 278314 | 3.06 | 1.42 | 2.85e-1 |
| Kemelmacher | T | 28452 | 9693 | 9693 | 129952 | 68.0 | 2.35e+2 | 2.02e-3 |
| lpi_gosh^T | T | 13455 | 3792 | 3792 | 111327 | $3.09e{+}2$ | 1.60e+2 | 3.31e-16 |
| rdist2 | T | 3198 | 3198 | 3198 | 66426 | 1.17e + 3 | 4.50e+2 | 2.64e-5 |
| $deter7^T$ | T | 18153 | 6375 | 6375 | 56254 | 6.89 | 7.97 | 6.97e-3 |
| shyy41 | T | 4720 | 4720 | 4720 | 34200 | 1.89e + 2 | 1.85e+2 | 8.26e-21 |
| nemeth01 | D | 9506 | 9505 | 9506 | 744064 | 32.0 | 7.61e+4 | 2.06e-1 |
| raefsky1 | D | 3242 | 3241 | 3242 | 299891 | 6.81 | 8.13e + 2 | 2.18e-4 |
| $r05^T$ | D | 9690 | 5189 | 5190 | 114523 | 62.4 | $1.19e{+4}$ | 2.91e-1 |
| $p010^T$ | D | 19090 | 10089 | 10090 | 138178 | 60.2 | $1.95e{+}4$ | 2.92e-1 |
| $scagr7-2b^T$ | D | 13847 | 9742 | 9742 | 55369 | 2.60e + 2 | 8.93e + 3 | 9.20e-3 |
| cavity16 | D | 4562 | 4561 | 4562 | 147009 | $2.19e{+}2$ | 1.32e + 3 | 9.85e-7 |
| utm5940 | D | 5940 | 5939 | 5940 | 95720 | 52.5 | $8.91e{+}2$ | 4.10e-9 |

In Table 1 we list all the test problems and some of their basic properties, where for each matrix pair (A, B), A or A^T (so that $m \ge n$) is a sparse matrix from the University of Florida Sparse Matrix Collection [3] and B is either

| | 3 | 1 | | | | | F 4 | | | - | |
|-----|---|----|-------|---|----|-----|------------|----|---|----|--|
| | 1 | ۰. | ۰. | | | | 1 | -1 | | | |
| T = | 1 | · | · | | or | D = | | · | · | | |
| | | · | · · . | 1 | | | | | 1 | -1 | |
| | L | | 1 | 3 | | | L | | - | | |

with p = n or n - 1, the latter of which is the scaled discrete approximation of the first order derivative operator in dimension one, nnz is the total number of nonzero elements in A and B, $\kappa(\begin{bmatrix} A \\ B \end{bmatrix})$ is the condition number of $\begin{bmatrix} A \\ B \end{bmatrix}$, and σ_{\max} and σ_{\min} are the largest and smallest nontrivial generalized singular values of (A, B) computed, for experimental purpose, by the MATLAB built-in function gsvd. Note that for the matrix pairs (A, B) with B = T, all generalized singular values are nontrivial ones while for those with B = D, there exists at least one infinite generalized singular value. We take B = T in Experiments 5.1–5.5 and B = D in the remaining experiments.

For all the six algorithms, we take the outer stopping tolerance $tol = 10^{-8}$, the initial vector $x_0 = \text{mod}(1:n,4)/||\text{mod}(1:n,4)||$ with mod the MATLAB built-in

function, and all the other parameters by default as described in Section 4.3. For the correction equations (2.6) and (4.7), we take the *n*-dimensional zero vectors as initial guesses, and use the MATLAB built-in function minres to solve them until it converges with the prescribed tolerance $\tilde{\varepsilon} = 10^{-4}$ in (2.8) or maximum *n* inner iteration steps have been consumed. We stop each algorithm and output the computed partial GSVD of (A, B) if all the ℓ desired GSVD components of (A, B) have been found or the total *n* correction equations have been solved.

In all the tables, we abbreviate the thick-restart CPF-JDGSVD, CPF-HJDGSVD, IF-HJDGSVD and RCPF-JDGSVD, RCPF-HJDGSVD, RIF-HJDGSVD algorithms as CPF, CPFH, IFH, and RCPF, RCPFH, RIFH, respectively. We denote by $I_{\rm out}$ and $I_{\rm in}$ the total numbers of outer and inner iterations used, and by $T_{\rm cpu}$ the total CPU time in second counted by the MATLAB built-in commands tic and toc.

Experiment 5.1. We compute one GSVD component of $(A, B) = (\text{dano3mip}^T, T)$ with the target $\tau = 3.75e + 2$. The desired σ_* is one of the largest generalized singular values of (A, B) and is clustered with its nearby ones.



FIG. 1. Computing one GSVD component of $(A, B) = (\text{dano3mip}^T, T)$ with $\tau = 3.75e + 2$.

For the matrix pairs in this and the next four experiments, the matrices $B^T B$'s are symmetric positive definite and well conditioned, whose Cholesky factorizations can be efficiently computed using $\mathcal{O}(n)$ flops, meaning that matrix-vector multiplications with its inversion $(B^T B)^{-1}$ can be applied at cost of $\mathcal{O}(n)$ flops for each linear system. For this reason, during the subspace expansion phase of the CPF-HJDGSVD and RCPF-HJDGSVD algorithms, we exploit the MATLAB built-in command \setminus to implement matrix-vector multiplications with $(B^T B)^{-1}$ and to update the intermediate matrix H_c in (2.18). Moreover, for experimental purpose, in this and the next experiments we solve all the correction equations (2.6) and (4.7) accurately by using the LU factorizations of $(A^T A - \rho^2 B^T B)$'s in order to exhibit how RCPF-JDGSVD, RCPF-HJDGSVD and RIF-HJDGSVD truly behave.

Figure 1 depicts the outer convergence curves of CPF-JDGSVD and RCPF-JDGSVD for computing the desired GSVD component of (A, B). Clearly, CPF-JDGSVD converges slowly and irregularly and even suffers from two sharp oscillations at outer iterations 22 and 40, respectively, and it uses 42 outer iterations to converge. In contrast, RCPF-JDGSVD exhibits far smoother outer convergence behavior, and uses only 16 outer iterations to attain the convergence, thereby saving 62% outer iterations. This demonstrates the considerable superiority of RCPF-JDGSVD to CPF-JDGSVD when computing an extreme GSVD component of (A, B).

Experiment 5.2. We compute one GSVD component of $(A, B) = (\text{plddb}^T, T)$ with the interior generalized singular value σ_* closest to the target $\tau = 10$, which is clustered

with its neighbors.



FIG. 2. Computing one GSVD component of $(A, B) = (\text{plddb}^T, T)$ with $\tau = 10$.

Figure 2 draws the outer convergence curves of the two harmonic extractionbased algorithms CPF-HJDGSVD and IF-HJDGSVD as well as the two refined harmonic extraction-based algorithms RCPF-HJDGSVD and RIF-HJDGSVD. We can see that RCPF-HJDGSVD and RIF-HJDGSVD converge more quickly and much more smoothly than CPF-HJDGSVD and IF-HJDGSVD, and the former two algorithms use 4 and 7 fewer outer iterations than the latter two ones to reach the convergence, respectively. We also observe that RCPF-HJDGSVD and RIF-HJDGSVD perform equally well for this problem.

Experiment 5.3. We compute one, five and ten largest GSVD components of (A, B) = (barth, T) with the target $\tau = 20$. The desired generalized singular values are clustered one another.

| l | Algorithm | $I_{\rm out}$ | $I_{\rm in}$ | $T_{\rm cpu}$ | Algorithm | $I_{\rm out}$ | $I_{\rm in}$ | $T_{\rm cpu}$ |
|----|----------------|---------------|--------------|---------------|-----------|---------------|--------------|---------------|
| | CPF | 36 | 655 | 0.18 | RCPF | 33 | 673 | 0.18 |
| 1 | CPFH | 36 | 653 | 0.17 | RCPFH | 33 | 673 | 0.19 |
| | \mathbf{IFH} | 35 | 697 | 0.18 | RIFH | 33 | 673 | 0.18 |
| | CPF | 86 | 2515 | 0.56 | RCPF | 67 | 2543 | 0.55 |
| 5 | CPFH | 86 | 2560 | 0.58 | RCPFH | 67 | 2543 | 0.59 |
| | \mathbf{IFH} | 81 | 2609 | 0.56 | RIFH | 61 | 2466 | 0.51 |
| | CPF | 166 | 5575 | 1.25 | RCPF | 91 | 4834 | 0.94 |
| 10 | CPFH | 173 | 5634 | 1.30 | RCPFH | 91 | 4834 | 0.97 |
| | IFH | 155 | 5603 | 1.25 | RIFH | 109 | 5269 | 1.09 |

TABLE 2 Computing the ℓ GSVD components of (A, B) = (barth, T) with $\tau = 20$.

Table 2 reports the results obtained. As we can see, for $\ell = 1$, the refined and refined harmonic extraction-based JDGSVD algorithms use slightly fewer outer iterations and comparable inner iterations and CPU time to achieve the convergence, compared with the standard and harmonic extraction-based JDGSVD algorithms. However, for $\ell = 5$, RCPF-JDGSVD, RCPF-HJDGSVD and RIF-HJDGSVD are considerably faster and use nearly 20 fewer outer iterations than CPF-JDGSVD, CPF-HJDGSVD and IF-HJDGSVD, respectively, which save 22.1% ~ 24.5% of the outer iterations, though the six JDGSVD algorithms use almost the same inner iter-

ations and CPU time to converge. For $\ell = 10$, RIF-HJDGSVD uses 29% fewer outer iterations, 6% fewer inner iterations and 13% less CPU time than IF-HJDGSVD; RCPF-JDGSVD and RCFP-HJDGSVD save more than 45% of the outer iterations, 13% of the inner iterations, and 24% of the CPU time, respectively, relative to CPF-JDGSVD and CPF-HJDGSVD. Of the three refined and refined harmonic algorithms, RCPF-JDGSVD and RCPF-HJDGSVD are better than RIF-HJDGSVD due to their faster outer convergence, and RCPF-JDGSVD is the most efficient due to its least CPU time.

Clearly, for the sake of faster outer convergence and higher overall efficiency, RCPF-JDGSVD is the most recommended algorithm for this problem. This should be expected as the generalized singular values of interest are extreme ones, for which the standard extraction suits better than the harmonic extraction. As a consequence, RCPF-JDGSVD performs better than RCPF-HJDGSVD and RIF-HJDGSVD, though its advantage is not so obvious relative to RCPF-HJDGSVD.

Experiment 5.4. We compute one, five and ten interior GSVD components of the matrix pair $(A, B) = (\text{large}^T, T)$ with the generalized singular values closest to $\tau = 14$ highly clustered.



FIG. 3. Computing one GSVD component of $(A, B) = (\text{large}^T, T)$ with $\tau = 14$.

| TABLE 3 | |
|---|--|
| Computing the ℓ GSVD components of $(A, B) = (\text{large}^T, T)$ with $\tau = 14$. | |
| | |

| l | Algorithm | $I_{\rm out}$ | I_{in} | $T_{\rm cpu}$ | Algorithm | $I_{\rm out}$ | I_{in} | $T_{\rm cpu}$ |
|----|----------------------|---------------|-------------------|---------------|-----------|---------------|-------------------|---------------|
| | CPF | 13 | 3266 | 0.23 | RCPF | 33 | 6068 | 0.44 |
| 1 | CPFH | 16 | 2677 | 0.18 | RCPFH | 12 | 3109 | 0.22 |
| | IFH | 12 | 3544 | 0.24 | RIFH | 11 | 3332 | 0.22 |
| | CPF | 65 | 42115 | 2.84 | RCPF | 720 | 635121 | 43.5 |
| 5 | CPFH | 43 | 13255 | 0.93 | RCPFH | 25 | 16434 | 1.11 |
| | IFH | 29 | 15602 | 1.06 | RIFH | 27 | 23767 | 1.61 |
| | CPF | 4291 | 4084862 | 358 | RCPF | 771 | 720147 | 50.7 |
| 10 | CPFH | 61 | 29032 | 2.27 | RCPFH | 39 | 33313 | 2.48 |
| | IFH | 43 | 30135 | 2.23 | RIFH | 36 | 30570 | 2.26 |

Figure 3 depicts the outer convergence curves of the six JDGSVD algorithms for computing one desired GSVD component of (A, B), and Table 3 displays all the results obtained.

For $\ell = 1$, all the six algorithms succeed to compute the desired GSVD component. As is seen from Figure 3 and Table 3, RCPF-JDGSVD uses much more outer iterations and thus much more inner iterations and CPU time to converge than CPF-JDGSVD because of the outer convergence stagnation of the former. Notice that this is an interior GSVD problem. The stagnation implies that the expanded subspaces have little improvements at those stagnation steps, causing that the accuracy of selected approximate generalized singular values and right generalized singular vectors remain almost the same. This is because RCPF-JDGSVD selects approximate generalized singular values incorrectly or there is no good Ritz value at all at those steps, which is an intrinsic deficiency of the standard extraction for interior eigenvalue, SVD and GSVD problems, as we have pointed out in the introduction: If this selection is done incorrectly or there are spurious values at some step, the searching subspaces in the refined extraction are expanded wrongly at that step and provide little information on the desired generalized singular vectors. In contrast, RCPF-HJDGSVD and RIF-HJDGSVD outperform CPF-HJDGSVD and IF-HJDGSVD respectively, due to the smoother and faster outer convergence and/or better overall efficiency. This is because the harmonic extraction approach suits better for interior GSVD problems and can pick up the approximate generalized singular values correctly.

For $\ell = 5$, it seems from Table 3 that RCPF-JDGSVD uses significantly more outer and inner iterations and much more CPU time than CPF-JDGSVD. As a matter of fact, both CPF-JDGSVD and RCPF-JDGSVD are unreliable and fail to deliver all the desired GSVD components in this case: They find only the first three desired GSVD components but recompute each of the first two twice. RCPF-HJDGSVD uses nearly half of the outer iterations though a little bit more inner iterations and CPU time relative to CPF-HJDGSVD. RIF-HJDGSVD uses slightly fewer and considerably more inner iterations and CPU time than IF-HJDGSVD. Obviously, regarding outer convergence, RCPF-HJDGSVD and RIF-HJDGSVD are the best.

For $\ell = 10$, only CPF-HJDGSVD, IF-HJDGSVD and RCPF-HJDGSVD, RIF-HJDGSVD succeed to compute all the desired GSVD components of (A, B), while CPF-JDGSVD computes the first three desired ones three times in the beginning and fails to compute the fourth one when *n* correction equations are solved, and RCPF-JDGSVD only computes the first six desired GSVD components and recomputes the first three ones twice or three times. As a consequence, CPF-JDGSVD and RCPF-JDGSVD fail to solve the concerning GSVD problem correctly and are thus unreliable for this interior GSVD problem. On the contrary, RCPF-HJDGSVD and RIF-HJDGSVD outperform CPF-HJDGSVD and IF-HJDGSVD as they use considerably fewer outer iterations and comparable inner iterations; these four algorithms outperform CPF-JDGSVD and RCPF-JDGSVD significantly.

Clearly, due to the smoother and faster outer convergence and the reliability, RCPF-HJDGSVD and RIF-HJDGSVD are the most suitable choices for this interior GSVD problem.

Experiment 5.5. We compute ten GSVD components of the matrix pairs (A, B) = (ns3Da, T), (e40r0100, T), (Kemelmacher, T), (rat^T, T) , $(lpi_{-}gosh^T, T)$, (rdist2, T), (deter7^T, T) and (shyy41, T) with the targets $\tau = 2$, 11, 0.2, 0.15, 14.3, 19, 4.5 and 0.8, respectively. The desired generalized singular values of (ns3Da, T), (e40r0100, T) and (Kemelmacher, T), (rat^T, T) are extreme, i.e., the largest or smallest, and are clustered, and those of the remaining four matrix pairs are both interior and clustered.

Table 4 reports all the results obtained. For the computation of the largest GSVD components of (ns3Da, T) and (e40r0100, T), we observe from Table 4 that the

| A | Algorithm | $I_{\rm out}$ | $I_{ m in}$ | $T_{\rm cpu}$ | Algorithm | $I_{\rm out}$ | $I_{ m in}$ | $T_{\rm cpu}$ |
|------------------------|----------------------|---------------|-------------|---------------|-----------|---------------|-------------|---------------|
| | CPF | 126 | 3823 | 6.81 | RCPF | 89 | 3458 | 6.03 |
| ns3Da | CPFH | 119 | 3858 | 6.87 | RCPFH | 90 | 3471 | 6.17 |
| | IFH | 113 | 3776 | 6.85 | RIFH | 100 | 3646 | 6.49 |
| | CPF | 141 | 9507 | 5.20 | RCPF | 58 | 7656 | 3.86 |
| e40r0100 | CPFH | 91 | 9094 | 4.74 | RCPFH | 58 | 7549 | 3.87 |
| | IFH | 84 | 9020 | 4.74 | RIFH | 58 | 7615 | 3.83 |
| | CPF | 45 | 48004 | 9.82 | RCPF | 34 | 45001 | 9.09 |
| Kemelm | CPFH | 39 | 48669 | 10.1 | RCPFH | 37 | 48271 | 10.0 |
| | IFH | 35 | 45056 | 9.25 | RIFH | 36 | 46594 | 9.60 |
| | CPF | 172 | 2364 | 0.70 | RCPF | 81 | 2289 | 0.54 |
| rat^T | CPFH | 94 | 2191 | 0.54 | RCPFH | 80 | 2232 | 0.56 |
| | IFH | 105 | 2118 | 0.54 | RIFH | 81 | 2289 | 0.54 |
| | CPF | 52 | 6042 | 0.81 | RCPF | 43 | 6255 | 0.77 |
| lpi_gosh^T | CPFH | 44 | 5661 | 0.71 | RCPFH | 37 | 5203 | 0.63 |
| | IFH | 39 | 5405 | 0.67 | RIFH | 36 | 5023 | 0.60 |
| | CPF | 68 | 5397 | 0.48 | RCPF | 46 | 4476 | 0.39 |
| rdist2 | CPFH | 51 | 4485 | 0.40 | RCPFH | 44 | 4348 | 0.40 |
| | IFH | 54 | 5238 | 0.45 | RIFH | 47 | 4734 | 0.41 |
| | CPF | 87 | 7208 | 1.12 | RCPF | 68 | 6820 | 1.04 |
| $deter7^T$ | CPFH | 78 | 6876 | 1.07 | RCPFH | 55 | 6716 | 0.97 |
| | IFH | 69 | 7430 | 1.11 | RIFH | 57 | 7418 | 1.06 |
| | CPF | 4727 | 7644960 | 639 | RCPF | 4724 | 3555937 | 263 |
| shyy41 | CPFH | 4727 | 4493658 | $1.13e{+}3$ | RCPFH | 66 | 81061 | 6.35 |
| | IFH | 4728 | 17581871 | 1.49e+3 | RIFH | 63 | 79204 | 6.29 |

 TABLE 4

 Results on test matrix pairs in Example 5.5, where Kemelm is the abbreviation of Kemelmacher.

refined and refined harmonic extraction-based RCPF-JDGSVD, RCPF-HJDGSVD, RIF-HJDGSVD algorithms outperform the standard and harmonic extraction-based CPF-JDGSVD, CPF-HJDGSVD, IF-HJDGSVD algorithms as they use moderately to substantially fewer outer iterations, slightly to moderately fewer inner iterations and less CPU time. For (ns3Da, T), RCPF-JDGSVD and RCPF-HJDGSVD are very comparable, and they both are superior to RIF-HJDGSVD due to the faster outer convergence and higher overall efficiency. For (e40r0100, T), all the three refined and refined harmonic JDGSVD algorithms are suitable.

For the computation of the smallest GSVD components, we see from Table 4 that for (Kemelmacher, T), RCPF-JDGSVD and RCPF-HJDGSVD use moderately or slightly fewer outer and inner iterations and less CPU time than CPF-JDGSVD and CPF-HJDGSVD, respectively, while RIF-HJDGSVD and IF-HJDGSVD work equally well. Obviously, except for CPF-JDGSVD, all the other five algorithms are appropriate for the concerning GSVD problem of this matrix pair. Nonetheless, as far as both the outer convergence and overall efficiency are concerned, RCPF-JDGSVD is the most recommended. For (rat^T, T) , the refined and refined harmonic extraction-based JDGSVD algorithms outperform the standard and harmonic extraction-based

JDGSVD algorithms substantially, as outer iterations indicate. Regarding the overall efficiency, RCPF-JDGSVD outmatches CPF-JDGSVD considerably as it uses slightly fewer inner iterations and much less CPU time. On the other hand, RCPF-HJDGSVD and RIF-HJDGSVD are competitive with CPF-HJDGSVD and IF-HJDGSVD in terms of inner iterations and CPU time. For the sake of better convergence behavior, RCPF-JDGSVD, RCPF-HJDGSVD and RIF-HJDGSVD are all proper choices for this problem.

For the computation of interior GSVD components of $(\text{lpi}_{\text{gosh}}^T, T)$, (rdist2, T)and $(\text{deter7}^T, T)$, as far as the outer convergence is concerned, RCPF-JDGSVD, RCPF-HJDGSVD and RIF-HJDGSVD surpass CPF-JDGSVD, CPF-HJDGSVD and IF-HJDGSVD, respectively. Regarding the overall efficiency, the three refined and refined harmonic JDGSVD algorithms are superior to the standard and harmonic JDGSVD algorithms as they use less CPU time and/or fewer inner iterations.

For (shyy41, T), since the desired generalized singular values are truly interior and clustered, as we have elaborated in the introduction, the standard, harmonic and even refined extraction approach-based JDGSVD algorithms may face severe difficulties, which is confirmed by the results in Table 4. In fact, we observe very erratic convergence behavior of CPF-JDGSVD, CPF-HJDGSVD, IF-HJDGSVD and RCPF-JDGSVD; when n correction equations are solved, they fail to compute all the desired GSVD components and only output seven, seven, eight and four converged ones, respectively. On the contrary, RCPF-HJDGSVD and RIF-HJDGSVD are very successful to compute all the desired GSVD components quickly. Undoubtedly, RCPF-HJDGSVD and RIF-HJDGSVD are the only two proper choices for this difficult problem; they are competitive in terms of both outer convergence and overall efficiency.

Experiment 5.6. We compute one GSVD component of (A, B) = (nemeth01, D) corresponding to the interior generalized singular value closest to the target $\tau = 6.5$.



FIG. 4. Computing one GSVD component of (A, B) = (nemeth01, D) with $\tau = 6.5$.

For the matrix pairs in this and the next experiments, the matrices B's are rank deficient, so that CPF-HJDGSVD and RCPF-HJDGSVD are not applicable. We compute the desired GSVD components of (A, B) using CPF-JDGSVD, IF-HJDGSVD and RCPF-JDGSVD, RIF-HJDGSVD. Specifically, for the problem in this experiment, for the illustration of true outer convergence of these four algorithms, we use the LU factorizations of the matrices $(A^T A - \rho^2 B^T B)$'s to solve all the correction equations involved accurately, and draw their outer convergence curves in Figure 4.

As is seen from Figure 4, in contrast to CPF-JDGSVD and IF-HJDGSVD whose convergence is delayed because of the frequent oscillations and stagnation, respectively, RCPF-JDGSVD and RIF-HJDGSVD converge faster and much more smoothly, and use 23 and 6 fewer outer iterations to achieve the convergence, respectively. Moreover, since the standard extraction approach may pick up approximate generalized singular values incorrectly for interior GSVD problems, RCPF-JDGSVD may suffer from oscillations, as shown by Figure 4 at iterations 4–9 and 19–21. These phenomena confirm the intrinsic shortcoming of RCPF-JDGSVD for computing truly interior GSVD components. As a consequence, although RCPF-JDGSVD and RIF-HJDGSVD use the same number of outer iterations to converge and outperform CPF-JDGSVD and IF-HJDGSVD, we recommend RIF-HJDGSVD for this problem.

Experiment 5.7. We compute ten GSVD components of (A, B) = (raefsky1, D), $(\text{r05}^T, D)$, $(\text{p010}^T, D)$ and $(\text{scagr7-2b}^T, D)$, (cavity16, D), (utm5940, D) with the targets $\tau = 57, 61, 80$ and 32.5, 20.8, 58, respectively. All the desired generalized singular values are interior and clustered ones. This implies that all the correction equations may be hard to solve by the MINRES method.

| Α | Algorithm | $I_{\rm out}$ | $I_{ m in}$ | $T_{\rm cpu}$ | Algorithm | $I_{\rm out}$ | I_{in} | $T_{\rm cpu}$ |
|------------------------------|---|--|----------------------------|--------------------|--------------|--|---|---|
| raefsky1 | $\begin{array}{c} \mathrm{CPF} \\ \mathrm{IFH} \end{array}$ | $\frac{52}{37}$ | $91560 \\ 67609$ | $16.0 \\ 13.5$ | RCPF RIFH | $\begin{array}{c} 42\\ 35 \end{array}$ | $76689 \\ 66005$ | $\begin{array}{c} 13.4\\ 11.9 \end{array}$ |
| $r05^{T}$ | CPF IFH | 100 91 | $60409 \\ 64795$ | $8.71 \\ 19.2$ | RCPF RIFH | 99 69 | $\begin{array}{c} 65947 \\ 53142 \end{array}$ | $9.76 \\ 7.73$ |
| $p010^T$ | $\begin{array}{c} \mathrm{CPF} \\ \mathrm{IFH} \end{array}$ | $\begin{array}{c} 104 \\ 87 \end{array}$ | $104797 \\ 132654$ | $28.8 \\ 55.2$ | RCPF RIFH | $\begin{array}{c} 107 \\ 63 \end{array}$ | $\frac{113825}{81573}$ | $31.4 \\ 22.4$ |
| $\operatorname{scagr7-2b}^T$ | CPF IFH | $\begin{array}{c} 185 \\ 65 \end{array}$ | $\frac{130938}{141444}$ | $22.1 \\ 23.3$ | RCPF RIFH | $\begin{array}{c} 294\\ 46 \end{array}$ | $232935 \\ 112561$ | $\begin{array}{c} 39.0 \\ 18.4 \end{array}$ |
| cavity16 | $\begin{array}{c} \mathrm{CPF} \\ \mathrm{IFH} \end{array}$ | 122 101 | $95455 \\ 85234$ | $13.5 \\ 23.7$ | RCPF RIFH | $4565 \\ 68$ | $2107949 \\ 63650$ | $\begin{array}{c} 292 \\ 8.95 \end{array}$ |
| utm5940 | CPF IFH | $\begin{array}{c} 4255\\ 1951 \end{array}$ | $\frac{14983907}{8178349}$ | 2.13e+3 2.36e+3 | RCPF RIFH | $\begin{array}{c} 5949\\ 87\end{array}$ | $10827153 \\ 266135$ | 1.55e+3 35.8 |

TABLE 5Results on test matrix pairs in Example 5.7.

Table 5 lists the results. For (raefsky1, D), we see that RCPF-JDGSVD and RIF-HJDGSVD respectively outperform CPF-JDGSVD and IF-HJDGSVD by using significantly or slightly fewer outer and inner iterations and fairly less CPU time.

For $(r05^T, D)$ and $(p010^T, D)$, RIF-HJDGSVD uses substaintially fewer outer and inner iterations and considerably less CPU time than CPF-JDGSVD and RCPF-JDGSVD, which themselves behave similarly and outperform IF-HJDGSVD significantly with much less CPU time. Obviously, RIF-HJDGSVD is the best choice for these two problems in terms of both outer convergence and overall efficiency.

For (scagr7-2b^T, D), CPF-JDGSVD and RCPF-JDGSVD consume lots of outer iterations to converge. Actually, RCPF-JDGSVD fails to solve the problem, and it repeatedly computes the first two desired GSVD components. In contrast, IF-HJDGSVD and RIF-HJDGSVD use substantially fewer outer iterations to compute all the desired GSVD components. Between them, RIF-HJDGSVD wins IF-HJDGSVD with obviously fewer outer and inner iterations and less CPU time. Clearly, for this problem, RIF-HJDGSVD is the best choice.

For (cavity16, D) and (utm5940, D), the desired generalized singular values are

truly interior and clustered, causing that the standard, harmonic and even refined extraction approaches may not perform well, as is confirmed and shown in Table 5. We observe sharp oscillations in the outer convergence curves of CPF-JDGSVD, IF-HJDGSVD and RCPF-JDGSVD, of which the last one even fails to compute all the desired GSVD components when we use up the solutions of n correction equations. On the other hand, RIF-HJDGSVD converges smoothly and uses marvelously fewer outer and inner iterations and extremely less CPU time than the other three algorithms. Clearly, RIF-HJDGSVD is the most suitable choice for these two problems.

Summarizing all the numerical experiments, we can draw two conclusions: (i) For extreme GSVD components, refined and refined harmonic RCPF-JDGSVD, RCPF-HJDGSVD and RIF-HJDGSVD algorithms generally perform better than the standard and harmonic CPF-JDGSVD, CPF-HJDGSVD and IF-HJDGSVD algorithms, respectively; RCPF-JDGSVD is the best. (ii) For interior GSVD components, RCPF-HJDGSVD and RIF-HJDGSVD outmatch CPF-JDGSVD, CPF-HJDGSVD and IF-HJDGSVD, RCPF-JDGSVD with smoother and faster outer convergence and higher overall efficiency; if *B* has full column rank, then both RCPF-HJDGSVD and RIF-HJDGSVD are suitable choices; otherwise RIF-HJDGSVD is the most recommended algorithm due to its wider applicability.

6. Conclusions. The reliable and efficient computation of a partial GSVD of a large regular matrix pair (A, B) is vital in extensive applications, and has attracted much attention in recent years. In this paper, we have proposed three refined and refined harmonic extraction-based JDGSVD methods: RCPF-JDGSVD, and RCPF-HJDGSVD, RIF-HJDGSVD, and have developed practical thick-restart algorithms with deflation and purgation that can compute several GSVD components. They fix erratic convergence behavior and intrinsic possible non-convergence of the standard and harmonic JDGSVD methods: CPF-JDGSVD, CPF-HJDGSVD and IF-HJDGSVD proposed and developed in [11, 12]. The three new JDGSVD algorithms suit better for the computation of extreme and interior GSVD components of large regular matrix pairs, respectively. Numerical experiments have demonstrated that RCPF-HJDGSVD and RIF-HJDGSVD are generally the best choices for computing interior GSVD components, while RCPF-JDGSVD suits best for the extreme GSVD components. These confirm our elaborations in the introduction and the necessity and superiority of refined and refined harmonic extraction-based JDGSVD methods.

There remain some important issues that should be given special considerations. When the desired generalized singular values are truly interior and clustered, the correction equations (2.6) and (4.7) involved in JDGSVD algorithms are highly indefinite and ill conditioned, causing that the MINRES method may be very costly even if the relative residual of an approximate solution is only required to be fairly small. Unfortunately, we have observed that commonly used incomplete LU preconditioners generally work poorly and have no acceleration effect. Therefore, the efficient solutions of correction equations constitute the bottleneck of all JDGSVD algorithms. How to propose and develop specific preconditioners for the correction equations is extremely important and definitely deserves enough attention. This constitutes our future work.

Funding The first author was supported by the Youth Fund of the National Science Foundation of China (No. 12301485) and the Youth Program of the Natural Science Foundation of Jiangsu Province (No. BK20220482), and the second author was supported by the National Science Foundation of China (No.12171273).

Data Availability Enquires about data availability should be directed to the authors.

Declarations. The two authors declare that they have no financial interests, and the two authors read and approved the final manuscript.

REFERENCES

- Z. BAI, J. DEMMEL, J. DONGARRA, A. RUHE, AND H. A. VAN DER VORST, Templates for the Solution of Algebraic Eigenvalue Problems: A Practical Guide, SIAM, Philadelphia, PA, 2000.
- [2] J. K. CULLUM AND R. A. WILLOUGHBY, Lanczos Algorithms for Large Symmetric Eigenvalue Computations: Vol. I: Theory, SIAM, Philadelphia, PA, 2002.
- [3] T. A. DAVIS AND Y. HU, The University of Florida sparse matrix collection, ACM Trans. Math. Software, 38 (2011), pp. 1-25. Data available online at http://www.cise.ufl.edu/ research/sparse/matrices/.
- [4] G. H. GOLUB AND C. F. VAN LOAN, Matrix Computations, 4th Ed., The John Hopkins University Press, Baltimore, 2012.
- [5] M. E. HOCHSTENBACH, A Jacobi-Davidson type SVD method, SIAM J. Sci. Comput., 23 (2001), pp. 606–628.
- [6] M. E. HOCHSTENBACH, Harmonic and refined extraction methods for the singular value problem, with applications in least squares problems, BIT, 44 (2004), pp. 721–754.
- M. E. HOCHSTENBACH, A Jacobi-Davidson type method for the generalized singular value problem, Linear Algebra Appl., 431 (2009), pp. 471–487.
- [8] M. E. HOCHSTENBACH AND G. L. SLEIJPEN, Harmonic and refined Rayleigh-Ritz for the polynomial eigenvalue problem, Numer. Linear Algebra Appl., 15 (2008), pp. 35–54.
- [9] J. HUANG AND Z. JIA, On inner iterations of Jacobi-Davidson type methods for large SVD computations, SIAM J. Sci. Comput., 41 (2019), pp. A1574-A1603.
- [10] J. HUANG AND Z. JIA, On choices of formulations of computing the generalized singular value decomposition of a matrix pair, Numer. Algorithms, 87 (2021), pp. 689–718.
- [11] J. HUANG AND Z. JIA, Two harmonic Jacobi-Davidson methods for computing a partial generalized singular value decomposition of a large matrix pair, J. Sci. Comput., 93 (2022). article no.41.
- [12] J. HUANG AND Z. JIA, A cross-product free Jacobi-Davidson type method for computing a partial generalized singular value decomposition of a large matrix pair, J. Sci. Comput., 94 (2023). article no.3.
- [13] Z. JIA, Refined iterative algorithms based on Arnoldi's process for large unsymmetric eigenproblems, Linear Algebra Appl., 259 (1997), pp. 1–23.
- [14] Z. JIA, Polynomial characterizations of the approximate eigenvectors by the refined Arnoldi method and an implicitly restarted refined Arnoldi algorithm, Linear Algebra Appl., 287 (1999), pp. 191–214.
- [15] Z. JIA, The refined harmonic Arnoldi method and an implicitly restarted refined algorithm for computing interior eigenpairs of large matrices, Appl. Numer. Math., 42 (2002), pp. 489– 512.
- [16] Z. JIA, Some theoretical comparisons of refined Ritz vectors and Ritz vectors, Sci. China Ser. A, 47 (2004), pp. 222–233.
- [17] Z. JIA, The convergence of harmonic Ritz values, harmonic Ritz vectors and refined harmonic Ritz vectors, Math. Comput., 74 (2005), pp. 1441–1456.
- [18] Z. JIA, Using cross-product matrices to compute the SVD, Numer. Algorithms, 42 (2006), pp. 31–61.
- [19] Z. JIA AND C. LI, Inner iterations in the shift-invert residual Arnoldi method and the Jacobi-Davidson method, Sci. China Math., 57 (2014), pp. 1733–1752.
- [20] Z. JIA AND C. LI, Harmonic and refined harmonic shift-invert residual Arnoldi and Jacobi-Davidson methods for interior eigenvalue problems, J. Comput. Appl. Math., 282 (2015), pp. 83–97.
- [21] Z. JIA AND D. NIU, An implicitly restarted refined bidiagonalization Lanczos method for computing a partial singular value decomposition, SIAM J. Matrix Anal. Appl., 25 (2003), pp. 246–265.
- [22] Z. JIA AND D. NIU, A refined harmonic Lanczos bidiagonalization method and an implicitly restarted algorithm for computing the smallest singular triplets of large matrices, SIAM J. Sci. Comput., 32 (2010), pp. 714–744.

26

- [23] Z. JIA AND G. W. STEWART, An analysis of the Rayleigh-Ritz method for approximating eigenspaces, Math. Comput., 70 (2001), pp. 737-747.
- [24] E. KOKIOPOULOU, C. BEKAS, AND E. GALLOPOULOS, Computing smallest singular triplets with implicitly restarted Lanczos bidiagonalization, Appl. Numer. Math., 49 (2004), pp. 39–61.
- [25] R. B. MORGAN AND M. ZENG, Harmonic projection methods for large non-symmetric eigenvalue problems, Numer. Linear Algebra Appl., 5 (1998), pp. 33–55.
- [26] R. B. MORGAN AND M. ZENG, A harmonic restarted Arnoldi algorithm for calculating eigenvalues and determining multiplicity, Linear Algebra Appl., 415 (2006), pp. 96–113.
- [27] C. C. PAIGE AND M. A. SAUNDERS, Towards a generalized singular value decomposition, SIAM J. Numer. Anal., 18 (1981), pp. 398–405.
- [28] B. N. PARLETT, The Symmetric Eigenvalue Problem, SIAM, Philadelphia, PA, 1998.
- [29] Y. SAAD, Iterative Methods for Sparse Linear Systems, 2nd Ed., SIAM, Philadelphia, PA, 2003.
- [30] Y. SAAD, Numerical Methods for Large Eigenvalue Problems, SIAM, Philadelphia, PA, 2011.
- [31] H. D. SIMON AND H. ZHA, Low-rank matrix approximation using the Lanczos bidiagonalization process with applications, SIAM J. Sci. Comput., 21 (2000), pp. 2257–2274.
- [32] A. STATHOPOULOS, Y. SAAD, AND K. WU, Dynamic thick restarting of the Davidson, and the implicitly restarted Arnoldi methods, SIAM J. Sci. Comput., 19 (1998), pp. 227–245.
- [33] G. W. STEWART, Matrix Algorithms II: Eigensystems, SIAM, Philadelphia, PA, 2001.
- [34] G. W. STEWART AND J.-G. SUN, Matrix Perturbation Theory, Acadmic Press, Inc., Boston, 1990.
- [35] M. STOLL, A Krylov-Schur approach to the truncated SVD, Linear Algebra Appl., 436 (2012), pp. 2795–2806.
- [36] H. VAN DER VORST, Computational Methods for Large Eigenvalue Problems, Elsvier, Holland, 2002.
- [37] C. F. VAN LOAN, Generalizing the singular value decomposition, SIAM J. Numer. Anal., 13 (1976), pp. 76–83.
- [38] L. WU, E. ROMERO, AND A. STATHOPOULOS, PRIMME_SVDS: A high-performance preconditioned SVD solver for accurate large-scale computations, SIAM J. Sci. Comput., 39 (2017), pp. S248–S271.
- [39] L. WU AND A. STATHOPOULOS, A preconditioned hybrid SVD method for accurately computing singular triplets of large matrices, SIAM J. Sci. Comput., 37 (2015), pp. S365–S388.
- [40] I. N. ZWAAN, Cross product-free matrix pencils for computing generalized singular values, (2019). arXiv:1912.08518 [math.NA].
- [41] I. N. ZWAAN AND M. E. HOCHSTENBACH, Generalized Davidson and multidirectional-type methods for the generalized singular value decomposition, (2017). arXiv:1705.06120 [math.NA].