

# Regularity criteria of 3D generalized magneto-micropolar fluid system in terms of the pressure

Jae-Myoung Kim<sup>1</sup>

<sup>1</sup>Department of Mathematics Education, Andong National University,

Andong 36729, Republic of Korea,

Email: jmkim02@anu.ac.kr

## Abstract

This work focuses on regularity criteria of 3D generalized magneto-micropolar fluid system in terms of the pressure in Lorentz spaces inspired by the recent works in [10] and [15].

*Mathematics Subject Classification(2000):* 35Q35, 35B65, 76D05

*Key words:* Regularity criteria; Weak solution, generalized magneto-micropolar fluid system

## 1 Introduction

This paper is concerned about the regularity conditions of the weak solutions to generalized magneto-micropolar fluid equations in  $\mathbb{R}^3$ , which are described by

$$\begin{cases} \partial_t u + (u \cdot \nabla) u + \nabla \pi = -(\mu + \chi) \Lambda^{2\alpha} u + \chi \nabla \times w + (b \cdot \nabla) b, \\ \partial_t w + (u \cdot \nabla) w = -\kappa \Lambda^{2\gamma} w + \eta \nabla (\nabla \cdot w) + \chi \nabla \times u - 2\chi w, \\ \partial_t b + (u \cdot \nabla) b = -\nu \Lambda^{2\beta} b + (b \cdot \nabla) u, \\ \nabla \cdot u(\cdot, t) = \nabla \cdot b(\cdot, t) = 0, \end{cases} \quad (1.1)$$

where  $u = u(x, t)$ ,  $w = w(x, t)$ ,  $b = b(x, t)$  and  $\pi := \pi(x, t) = \mathcal{P} + \frac{|b|^2}{2}$  denote the fluid velocity, the micro-rotation velocity (angular velocity of the rotation of the fluid particles), the magnetic and total pressure fields respectively. The notation  $\Lambda := (-\Delta)^{1/2}$  stands for positive constants. The positive constant  $\kappa$  in (1.1) correspond to the angular

viscosity,  $\nu$  is the inverse of the magnetic Reynolds number and  $\chi$  is the micro-rotational viscosity. We consider the initial value problem of (1.1), which requires initial conditions

$$u(x, 0) = u_0(x), \quad w(x, 0) = w_0(x) \quad \text{and} \quad b(x, 0) = b_0(x), \quad x \in \mathbb{R}^3, \quad (1.2)$$

and we also assume that  $\operatorname{div} u_0 = 0 = \operatorname{div} b_0$ .

The authors in [4, Theorem 2.2] construct the existence of Leray-Hopf solution on the whole space  $\mathbb{R}^3 \times (0, T)$  to the generalized Navier-Stokes equation. It is worth pointing that  $\dot{H}^{\frac{5-4\alpha}{2}}$  is a critical space, that is,  $\dot{H}^{\frac{5-4\alpha}{2}}$  norm is scaling invariant is also a solution, where  $u_\lambda(x, t) = \lambda^{2\alpha-1} u(\lambda x, \lambda^{2\alpha} t)$  with any  $\lambda > 0$ . By Sobolev embedding theorem, it is checked that  $\dot{H}^{\frac{5-3\alpha}{2}}(\mathbb{R}^3) \hookrightarrow L^{\frac{6}{3-2\alpha}}(\mathbb{R}^3)$ . Then

$$\|u\|_{L^4(0, T; L^{\frac{6}{3-2\alpha}})} \leq C \|u\|_{L^\infty(0, T; \dot{H}^{\frac{5-4\alpha}{2}})}^2 \|u\|_{L^2(0, T; \dot{H}^{\frac{5-2\alpha}{2}})}^2, \quad \frac{2\alpha}{4} + \frac{3}{\frac{6}{3\alpha-2}} = 2\alpha - 1.$$

For the regularity issues to weak solutions to the generalized Navier-Stokes equation, we refer to [3], [8], [9].

Recently, Deng and Shang [6] obtained global-in-time existence and uniqueness of smooth solutions to the problem (1.1)–(1.2) if  $\alpha \geq \frac{1}{2} + \frac{n}{4}$ ,  $\alpha + \gamma \geq \max(2, \frac{n}{2})$ , and  $\alpha + \beta \geq 1 + \frac{n}{2}$ . On the other hands, Fan and Zhong [10] established local-in-time existence and uniqueness of smooth solutions to the problem (1.1)–(1.2) for  $\alpha + \gamma > 1$  and furthermore they gave some regularity criteria via the gradient of velocity in a meaningful and appropriate spaces.

For the regularity criteria in Lorentz space, Li and Niu [15] proved that a weak solution  $(u, w, b)$  for the standard 3D MHD equations become regular under the scaling invariant conditions for the total pressure, in particualr, so called Serrin's conditions,  $\pi \in L^{q, \infty}(0, T; L^{p, \infty}(\mathbb{R}^3))$  with  $3/p + 2/q \leq 2$  and  $p > \frac{3}{2}$  (compare to [1], [21], [22], [23], [2], [19], [20] for Navier–Stokes equations). For  $p, q \in [1, \infty]$ , we define

$$\|f\|_{L^{p,q}(\Omega)} = \begin{cases} \left( p \int_0^\infty \alpha^q |\{x \in \Omega : |f(x)| > \alpha\}|^{\frac{q}{p}} \frac{d\alpha}{\alpha} \right)^{\frac{1}{q}}, & q < \infty, \\ \sup_{\alpha > 0} \alpha |\{x \in \Omega : |f(x)| > \alpha\}|^{\frac{1}{p}}, & q = \infty. \end{cases}$$

(see e.g. [17], [11], [14]) Motivated by the recent works in [10] and [15], the purpose of this note is to establish regularity criteria of 3D generalized magneto-micropolar fluid system (1.1) in terms of the pressure in Lorentz space. In this paper, we assume that  $\gamma = \nu = \chi = 1$  and  $\alpha = \beta$ .

Our results are stated as follows.

**Theorem 1.1.** Let  $0 < T < \infty$  and  $u_0, b_0, w_0 \in H^m(\mathbb{R}^3)$  with  $m > \frac{5}{2}$  and  $1 \leq \alpha, \gamma \leq \frac{5}{4}$ . There exists a sufficient constant  $\epsilon > 0$  such that if  $\pi$  or  $\nabla\pi$  satisfy

(A)  $\pi \in L^{q,\infty}(0, T; L^{p,\infty}(\mathbb{R}_+^3))$  and

$$\|\pi\|_{L^{p,\infty}(0,T;L^{q,\infty}(\mathbb{R}^3))} \leq \varepsilon, \text{ with } \frac{3}{p} + \frac{2\alpha}{q} = 2(2\alpha - 1), \frac{3}{2(\alpha - 1)} < q < \infty;$$

(B)  $\nabla\pi \in L^{\frac{2r\alpha}{2r\alpha-3}}(0, T; L^{r,\infty}(\mathbb{R}^3))$  with  $\frac{3}{2\alpha} < r \leq \infty$ ,

then the weak  $(u, b)$  is regular on  $(0, T]$ .

**Theorem 1.2.** Let  $\alpha = \beta = \gamma = 1$ . Assume  $(u_0, b_0) \in L_\sigma^2(\mathbb{R}^3) \cap L_\sigma^4(\mathbb{R}^3)$  and  $w_0 \in L^2(\mathbb{R}^3) \cap L^4(\mathbb{R}^3)$ . Let the triple  $(u, b, w)$  be a weak solution to system (1.1) on some time interval  $[0, T)$  with  $0 < T < \infty$ . There exists a sufficient constant  $\epsilon > 0$  such that if  $\nabla\mathcal{P}$  and  $b$  satisfy  $\nabla\mathcal{P} \in L^{\frac{2r}{3r-3},\infty}(0, T; L^{r,\infty}(\mathbb{R}^3))$  with

$$\|\nabla\mathcal{P}\|_{L^{\frac{2r}{3r-3},\infty}(0,T;L^{r,\infty})} \leq \epsilon, \quad \frac{3}{2} < r \leq \infty,$$

and

$$\|b\|_{L^{\frac{2a_1}{a_1-3},\infty}(0,T;L^{r,\infty})} \leq \infty, \quad 3 < a_1 \leq \infty,$$

then the weak solutions  $(u, b, w)$  is regular on  $(0, T]$ .

## 2 Proof of Theorem 1.1

To control the fractional diffusion term, we recall the following result (see e.g. [5]).

**Lemma 2.1.** With  $0 < \alpha < 2$ ,  $v, \Lambda^\alpha v \in L^p(\mathbb{R}^3)$  with  $p = 2k$ ,  $k \in \mathbb{N}$ , we obtain

$$\int |v|^{p-2} v \Lambda^\alpha v \, dx \geq \frac{1}{p} \int |\Lambda^{\frac{\alpha}{2}} v^{\frac{p}{2}}|^2 \, dx.$$

Also, we recall the following nonlinear Gronwall-type inequality established in [18] (see also [2] and [16]).

**Lemma 2.2.** Let  $T > 0$  and  $\varphi \in L_{loc}([0, T))$  be non-negative function. Assume further that

$$\varphi(t) \leq C_0 + C_1 \int_0^t \mu(s) \varphi(s) \, ds + \kappa \int_0^t \lambda(s)^{1-\epsilon} \varphi(s)^{1+A(\epsilon)} \, ds, \quad \forall 0 < \epsilon < \epsilon_0.$$

Where  $\kappa, \epsilon_0 > 0$  are constants,  $\mu \in L^1(0, T)$  and  $A(\epsilon) > 0$  satisfies  $\lim_{\epsilon \rightarrow 0} \frac{A(\epsilon)}{\epsilon} = c_0 > 0$ . Then  $\varphi$  is bounded on  $[0, T]$  if  $\|\lambda\|_{L^{1,\infty}(0,T)} < c_0^{-1} \kappa^{-1}$ .

In order to derive the regularity criteria of weak solutions to the system (1.1), we introduce the definition of weak solution. Let us denote

$$z^+ = u + b, \quad z^- = u - b. \quad (2.1)$$

Then system (1.1) can be reformulated as

$$\left\{ \begin{array}{l} \partial_t z^+ - \Lambda^{2\alpha} z^+ + (z^- \cdot \nabla) z^+ - \chi \nabla \times w + \nabla \pi = 0, \\ \partial_t z^- - \Lambda^{2\alpha} z^- + (z^+ \cdot \nabla) z^- - \chi \nabla \times w + \nabla \pi = 0, \\ \partial_t w - \kappa \Lambda^{2\gamma} w + (\frac{z^+ + z^-}{2}) \cdot \nabla w - \eta \nabla \operatorname{div} w + w - \nabla \times (\frac{z^+ + z^-}{2}) = 0, \\ \nabla \cdot z^+ = \nabla \cdot z^- = 0, \\ z^+(x, 0) = z_0^+(x), \quad z^-(x, 0) = z_0^-(x), \end{array} \right. \quad (2.2)$$

It is easy to show the following global  $L^2$ -bound,

$$\|(u, b, w)((\tau))\|_{L^2}^2 + \int_0^t (\|\Lambda^\alpha u(\tau)\|_{L^2}^2 + \|\Lambda^\gamma w(\tau)\|_{L^2}^2 + \|\Lambda^\alpha b(\tau)\|_{L^2}^2) d\tau \leq C.$$

**Proof:** Multiplying the first and the second equations of (2.2) by  $|z^+|^2 z^+$  and  $|z^-|^2 z^-$ , respectively, integrating by parts and summing up, we have

$$\begin{aligned} & \frac{1}{4} \frac{d}{dt} \| (z^+, z^-) \|_{L^4}^4 + \|\Lambda^\alpha (|z^+|^2, |z^-|^2) \|_{L^2}^2 \\ &= - \underbrace{\int_{\mathbb{R}^3} \nabla \pi \cdot (z^+ |z^+|^2 + z^- |z^-|^2) dx}_{\mathcal{J}_1} + \underbrace{\int_{\mathbb{R}^3} (|z^+|^2 z^+ + |z^-|^2 z^-) \cdot (\nabla \times w) dx}_{\mathcal{J}_2}. \end{aligned} \quad (2.3)$$

Taking the operator  $\operatorname{div}$ , to the first equation of (4.1), and using the facts  $\operatorname{div}(\nabla \times w) = 0$ , we see

$$-\Delta \pi = \operatorname{div} \operatorname{div}(z^+ \otimes z^-),$$

and thus,

$$\begin{aligned} \|\pi\|_{L^{q,2}} &\leq C \|z^+ \otimes z^-\|_{L^{q,2}} \leq C \|z^+\|_{L^{2q,4}} \|z^-\|_{L^{2q,4}} = C \||z^+|^2\|_{L^{q,2}}^{1/2} \||z^-|^2\|_{L^{q,2}}^{1/2} \\ &\leq C (\||z^+|^2\|_{L^{q,2}} + \||z^-|^2\|_{L^{q,2}}). \end{aligned}$$

Using integration by parts, Hölder's inequality in Lorentz space, Young's inequalities and Sobolev embedding for the fractional power, we note that for  $p > 1$ ,  $q > 2$  and  $\frac{5}{2} > \alpha > 0$  with

$$\frac{1}{q} + \frac{1}{2p} + \frac{5-2\alpha}{6} = 1, \quad (2.4)$$

$$\begin{aligned}
\int_{\mathbb{R}^3} \nabla \pi \cdot |z^+|^2 z^+ dx &= \int_{\mathbb{R}^3} \nabla(z^+ |z^+|^2) \pi dx \leq C \int_{\mathbb{R}^3} |z^+| |\nabla|z^+|^2| |\pi| dx \\
&\leq C \|z^+\|_{L^{2q,4}} \|\nabla|z^+|^2\|_{L^{\frac{6}{5-2\alpha},2}} \|\pi\|_{L^{2p,\infty}}^{1/2} \|\pi\|_{L^{2q,4}}^{1/2}, \\
&= C \||z^+|^2\|_{L^{q,2}}^{1/2} \|\nabla|z^+|^2\|_{L^{\frac{6}{5-2\alpha},2}} \|\pi\|_{L^{p,\infty}}^{1/2} \|\pi\|_{L^{q,2}}^{1/2} \\
&\leq C \||z^+|^2\|_{L^{q,2}}^{1/2} \|\nabla|z^+|^2\|_{L^{\frac{6}{5-2\alpha},2}} \|\pi\|_{L^{p,\infty}}^{1/2} \left( \||z^+|^2\|_{L^{q,2}}^{1/2} + \||z^-|^2\|_{L^{q,2}}^{1/2} \right) \\
&\leq C \||z^+|^2\|_{L^{q,2}} \|\nabla|z^+|^2\|_{L^{\frac{6}{5-2\alpha},2}} \|\pi\|_{L^{p,\infty}}^{1/2} \||z^-|^2\|_{L^{q,2}}^{1/2} \\
&\quad + C \||z^+|^2\|_{L^{q,2}}^{1/2} \|\nabla|z^+|^2\|_{L^{\frac{6}{5-2\alpha},2}} \|\pi\|_{L^{p,\infty}}^{1/2} \||z^-|^2\|_{L^{q,2}}^{1/2} \\
&\leq C \|\pi\|_{L^{p,\infty}}^{1/2} \left( \||z^+|^2\|_{L^{q,2}} + \||z^-|^2\|_{L^{q,2}} \right) \|\nabla|z^+|^2\|_{L^{\frac{6}{5-2\alpha},2}}.
\end{aligned}$$

In the same way,  $\int_{\mathbb{R}^3} \nabla \pi \cdot z^- |z^-|^2 dx$  can be bounded by

$$C \|\pi\|_{L^{p,\infty}}^{1/2} \left( \||z^+|^2\|_{L^{q,2}} + \||z^-|^2\|_{L^{q,2}} \right) \|\nabla|z^-|^2\|_{L^{\frac{6}{5-2\alpha},2}}.$$

And thus, we get

$$\begin{aligned}
\mathcal{J}_1 &\leq C \|\pi\|_{L^{p,\infty}}^{1/2} \left( \||z^+|^2\|_{L^{q,2}} + \||z^-|^2\|_{L^{q,2}} \right) \left( \|\nabla|z^+|^2\|_{L^{\frac{6}{5-2\alpha},2}} + \|\nabla|z^-|^2\|_{L^{\frac{6}{5-2\alpha},2}} \right) \\
&\leq C \|\pi\|_{L^{p,\infty}}^{1/2} \left( \||z^+|^2\|_{L^2}^{1-\left(\frac{3}{2\alpha}+\frac{3}{q\alpha}\right)} + \||z^-|^2\|_{L^2}^{1-\left(\frac{3}{2\alpha}+\frac{3}{q\alpha}\right)} \right) \\
&\quad \times \left( \|\nabla|z^+|^2\|_{L^{\frac{6}{5-2\alpha},2}}^{1+\left(\frac{3}{2\alpha}+\frac{3}{q\alpha}\right)} + \|\nabla|z^-|^2\|_{L^{\frac{6}{5-2\alpha},2}}^{1+\left(\frac{3}{2\alpha}+\frac{3}{q\alpha}\right)} \right) \\
&\leq C \|\pi\|_{L^{p,\infty}}^{\frac{2\alpha p}{4\alpha p-2p-3}} \left( \||z^+|^2\|_{L^2}^2 + \||z^-|^2\|_{L^2}^2 \right) + \frac{1}{16} \left( \|\Lambda^\alpha|z^+|^2\|_{L^2}^2 + \|\Lambda^\alpha|z^-|^2\|_{L^2}^2 \right).
\end{aligned}$$

For  $\mathcal{J}_2$ , integrating by parts, we note that

$$\begin{aligned}
\int_{\mathbb{R}^3} |z^+|^2 z^+ \cdot (\nabla \times w) dx &\leq \|w\|_{L^4} \|\nabla|z^+|^2\|_{L^2} \|z^+\|_{L^4} \leq \|w\|_{L^4}^2 \|z^+\|_{L^4}^2 + \frac{1}{16} \|\nabla|z^+|^2\|_{L^2}^2 \\
&\leq \|w\|_{L^4}^2 \|z^+\|_{L^4}^2 + \frac{1}{16} \|z^+\|_{L^2}^{2\theta} \|\Lambda^\alpha|z^+|^2\|_{L^2}^{2(1-\theta)}, \quad \theta = \frac{\alpha-1}{\alpha} \\
&\leq C(\|w\|_{L^4}^4 + \|z^+\|_{L^4}^4) + \frac{C}{16} \|z^+\|_{L^2}^2 + \frac{1}{16} \|\Lambda^\alpha|z^+|^2\|_{L^2}^2 \\
&\leq C(\|w\|_{L^4}^4 + \|z^+\|_{L^4}^4) + \frac{1}{16} \|\Lambda^\alpha|z^+|^2\|_{L^2}^2.
\end{aligned}$$

And thus,  $\mathcal{J}_2$  is bounded by

$$\mathcal{J}_2 \leq C(\|w\|_{L^4}^4 + \|z^+\|_{L^4}^4 + \|z^-\|_{L^4}^4) + \frac{1}{16} \left( \|\Lambda^\alpha|z^+|^2\|_{L^2}^2 + \|\Lambda^\alpha|z^-|^2\|_{L^2}^2 \right).$$

To get  $L^4$ -estimate for  $w$ , as before, multiplying the third equation of (2.2) by  $|w|^2 w$ , integrating by parts and summing up, we have

$$\begin{aligned} & \frac{1}{4} \frac{d}{dt} \|w\|_{L^4}^4 + \|\Lambda^\gamma |w|^2\|_{L^2}^2 + 2\chi \|w\|_{L^4}^4 + \kappa \|w \operatorname{div} w\|_{L^2}^2 \\ &= \underbrace{\frac{\chi}{2} \int_{\mathbb{R}^3} |w|^2 w \cdot (\nabla \times (z^+ + z^-)) dx}_{\mathcal{J}_3} - \underbrace{\int_{\mathbb{R}^3} \operatorname{div} w (w \cdot \nabla |w|^2) dx}_{\mathcal{J}_4}. \end{aligned} \quad (2.5)$$

As same manner as  $\mathcal{J}_2$ ,  $\mathcal{J}_3$  is bounded by

$$\mathcal{J}_3 \leq C(\|w\|_{L^4}^4 + \|z^+\|_{L^4}^4 + \|z^-\|_{L^4}^4) + \frac{1}{16} \|\Lambda^\gamma |w|^2\|_{L^2}^2, \quad \gamma \geq 1.$$

In a similar way,  $\mathcal{J}_4$  is also bounded by

$$\begin{aligned} \mathcal{J}_4 &\leq \kappa \|w \operatorname{div} w\|_{L^2} \|\nabla |w|^2\|_{L^2} \leq C \|\nabla |w|^2\|_{L^2}^2 + \frac{\kappa}{16} \|w \operatorname{div} w\|_{L^2}^2 \\ &\leq C \|w\|_{L^4}^4 + \frac{1}{16} \|\Lambda^\gamma |w|^2\|_{L^2}^2 + \frac{\kappa}{16} \|w \operatorname{div} w\|_{L^2}^2, \quad \gamma \geq 1. \end{aligned}$$

Let  $Y(t) := \|(z^+, z^-, w)\|_{L^4(\mathbb{R}^3)}^4$  and thus (2.8) becomes

$$\frac{d}{dt} Y(t) \lesssim \|\pi\|_{L^{q,\infty}(\mathbb{R}^3)}^q Y(t), \quad q = \frac{2\alpha p}{4\alpha p - 2p - 3}. \quad (2.6)$$

Now, we use an argument similar to the one used in the work of Bosia et al. [2]. For  $\epsilon > 0$ , Choose  $q_\epsilon = q - \epsilon(q + \frac{\alpha}{2\alpha-1} - \frac{3c_0}{4(2\alpha-1)})$  and  $r_\epsilon := \frac{q-\epsilon(q+\frac{\alpha}{2\alpha-1}-\frac{3c_0}{4(2\alpha-1)})}{\frac{2}{3}(q(2\alpha-1)-\alpha)(1-\epsilon)+\frac{c_0\epsilon}{2}}$  with

$$\begin{cases} \frac{3}{p_\kappa} + \frac{2\alpha}{q_\kappa} = 2(2\alpha-1), \\ \frac{q_\kappa}{p_\kappa} = \frac{q(1-\kappa)}{p} + \frac{c_0\kappa}{2}. \end{cases}$$

Due to the above relation, we get

$$\begin{aligned} \|\pi\|_{L^{p_\kappa,\infty}(\mathbb{R}^3)}^{q_\kappa} &\lesssim \|\pi\|_{L^{p,\infty}(\mathbb{R}^3)}^{q(1-\kappa)} \|\pi\|_{L^2,\infty}^{4\kappa} \lesssim \|\pi\|_{L^{p,\infty}(\mathbb{R}^3)}^{q(1-\kappa)} \|\pi\|_{L^2(\mathbb{R}^3)}^{4\kappa} \\ &\lesssim \|\pi\|_{L^{p,\infty}(\mathbb{R}^3)}^{q(1-\kappa)} \left( \|u\|_{L^2(\mathbb{R}^3)}^{4\kappa} + \|b\|_{L^2(\mathbb{R}^3)}^{4\kappa} \right). \end{aligned} \quad (2.7)$$

Since the pair  $(p_\kappa, q_\kappa)$  also meets  $3/p_\kappa + 2\alpha/q_\kappa = 2(2\alpha-1)$ . Using the estimate (2.7), (2.6) becomes

$$\frac{d}{dt} Y(t) \lesssim \|\pi\|_{L^{q_\kappa,\infty}(\mathbb{R}^3)}^{p_\kappa} \|u\|_{L^2(\mathbb{R}^3)}^2 \leq C \|\pi\|_{L^{q,\infty}(\mathbb{R}^3)}^{p(1-\kappa)} Y(t)^{1+2\kappa}.$$

And then integrating with respect to time from 0 to  $t$  with  $0 \leq t < T$ ,

$$Y(t) \leq CY(0) + C \int_0^t \|\pi\|_{L^{q,\infty}(\mathbb{R}^3)}^{p(1-\kappa)} Y(t)^{1+2\kappa} ds,$$

or equivalently,

$$\begin{aligned} \|w^+(t)\|_{L^4(\mathbb{R}^3)}^4 + \|w^-(t)\|_{L^4(\mathbb{R}^3)}^4 &\leq C\|w_0^+\|_{L^4(\mathbb{R}^3)}^4 + \|w_0^-\|_{L^4(\mathbb{R}^3)}^4 \\ &+ C \int_0^t \|\pi\|_{L^{q,\infty}(\mathbb{R}^3)}^{p(1-\kappa)} \|w^+\|_{L^4(\mathbb{R}^3)}^4 + \|w^-\|_{L^4(\mathbb{R}^3)}^{4(1+2\kappa)} ds. \end{aligned}$$

Due to Lemma 2.2, we are now able to complete the proof of Theorem 1.1 under the assumption (A) in Theorem 1.1.

*Part (B): Indeed, a proof is almost same to that in the argument in [7] or [13], however, for the reader's convenience, a sketch of the proof will be given. Multiplying both side of (2.2)<sub>1</sub> by  $z^+|z^+|^{3r-4}$  and then integrating them over  $\mathbb{R}^3$  we conclude that*

$$\begin{aligned} &\frac{1}{3r-2} \frac{d}{dt} \int_{\mathbb{R}^3} |z^+|^{3r-2} dx + \frac{4(3r-4)}{(3r-2)^2} \int_{\mathbb{R}^3} |\Lambda^\alpha z^+|^{\frac{3r-2}{2}}|^2 dx \\ &\lesssim \underbrace{\int_{\mathbb{R}^3} \nabla \pi \cdot |z^+|^{3r-4} z^+ dx}_{\mathcal{J}_5} + \underbrace{\frac{1}{2} \int_{\mathbb{R}^3} |z^+|^{3r-4} z^+ \cdot (\nabla \times w) dx}_{\mathcal{J}_6} \end{aligned} \quad (2.8)$$

By the integration by parts and Hölder inequality,  $\mathcal{J}_5$  is also written by

$$\begin{aligned} \mathcal{J}_1 &\leq (3r-4) \int_{\mathbb{R}^3} |\pi| |\nabla|z^+||z^+|^{3r-4} dx \\ &\leq \frac{2(3r-4)}{(3r-2)} \left( \int_{\mathbb{R}^3} |\pi|^2 |z^+|^{3r-4} dx \right)^{\frac{1}{2}} \left( \int_{\mathbb{R}^3} |\nabla|z^+|^{\frac{3r-2}{2}} dx \right)^{\frac{1}{2}}. \end{aligned} \quad (2.9)$$

Note that  $0 \leq I \leq a$  and  $0 \leq I \leq b$ , then  $I \leq \sqrt{ab}$ . Combining  $\mathcal{J}_5$  in (2.8) and (2.9), we get

$$\begin{aligned} \mathcal{J}_5 &\lesssim \left( \int_{\mathbb{R}^3} |\nabla \pi| |z^+|^{3r-3} dx \right)^{1/2} \left( \int_{\mathbb{R}^3} |\pi|^2 |z^+|^{3r-4} dx \right)^{\frac{1}{4}} \left( \int_{\mathbb{R}^3} |\nabla|z^+|^{\frac{3r-2}{2}}| dx \right)^{\frac{1}{4}} \\ &\leq C \left( \int_{\mathbb{R}^3} \left( |\nabla \pi| (|z^+|^2 + |z^-|^2) dx \right)^{(3r-3)/2} \right)^{2/3} \left( \int_{\mathbb{R}^3} \left( |\pi| (|z^+|^2 + |z^-|^2) dx \right)^{(3r-4)/2} \right)^{1/3} \\ &\quad + \frac{3r-4}{(3r-2)^2} \left( \int_{\mathbb{R}^3} |\nabla|z^+|^{\frac{3r-2}{2}}|^2 dx \right). \end{aligned}$$

Due to

$$\int_{\mathbb{R}^3} |\pi|^2 (|z^+|^2 + |z^-|^2)^{\frac{3r-4}{2}} dx \lesssim \|\pi\|_{L^{\frac{3r}{2}, 6r-6}}^2 \| |z^+|^2 + |z^-|^2 \|_{L^{\frac{3r}{2}, \frac{3r-3}{2}}}^{\frac{3r-4}{2}}$$

$$\begin{aligned}
&\lesssim \||z^+|^2 + |z^-|^2\|_{L^{\frac{3r}{2}, 6r-6}}^2 \||z^+|^2 + |z^-|^2\|_{L^{\frac{3r}{2}, \frac{3r-3}{2}}}^{\frac{3r-4}{2}} \\
&\lesssim \||z^+|^2 + |z^-|^2\|_{L^{\frac{3r}{2}, \frac{3r-3}{2}}}^2 \||z^+|^2 + |z^-|^2\|_{L^{\frac{3r}{2}, \frac{3r-3}{2}}}^{\frac{3r-4}{2}} = \||z^+|^2 + |z^-|^2\|_{L^{\frac{3r}{2}, \frac{3r-3}{2}}}^{\frac{3r}{2}},
\end{aligned}$$

and

$$\int_{\mathbb{R}^3} |\nabla \pi| \left( |z^+|^2 + |z^-|^2 \right)^{\frac{3r-3}{2}} dx \lesssim \|\nabla \pi\|_{L^{r,\infty}} \||z^+|^2 + |z^-|^2\|_{L^{\frac{3r}{2}, \frac{3r-3}{2}}}^{\frac{3r-3}{2}},$$

$\mathcal{J}_5$  is estimated by

$$\mathcal{J}_5 \leq C \|\nabla \pi\|_{L^{r,\infty}}^{\frac{2}{3}} \||z^+|^2 + |z^-|^2\|_{L^{\frac{3r}{2}, \frac{3r-2}{2}}}^{\frac{3r-2}{2}} + \frac{3r-4}{(3r-2)^2} \left( \int_{\mathbb{R}^3} |\nabla|z^+|^{\frac{3r-2}{2}}|^2 dx \right). \quad (2.10)$$

Next, for  $\mathcal{J}_6$ , using the Hölder and Young inequalities, we have

$$\begin{aligned}
\mathcal{J}_6 &\lesssim \|w\|_{L^{3r-2}} \||z^+|^{\frac{3r-4}{2}}\|_{L^{\frac{2(3r-2)}{3r-4}}} \|\nabla|z^+|^{\frac{3r-2}{2}}\|_{L^2} \\
&\lesssim \|w\|_{L^{3r-2}}^2 \||z^+|^{\frac{3r-4}{2}}\|_{L^{\frac{2(3r-2)}{3r-4}}}^2 + \|\nabla|z^+|^{\frac{3r-2}{2}}\|_{L^2}^2 \lesssim (\|w\|_{L^{3r-2}}^{3r-2} + \||z^+|\|_{L^{3r-2}}^{3r-2}) + \|\nabla|z^+|^{\frac{3r-2}{2}}\|_{L^2}^2.
\end{aligned} \quad (2.11)$$

And then, considering the estimates (2.10) and (2.11), (2.8) reduces

$$\begin{aligned}
&\frac{d}{dt} \int_{\mathbb{R}^3} |z^+|^{3r-2} dx + \int_{\mathbb{R}^3} |\Lambda^\alpha|z^+|^{\frac{3r-2}{2}}|^2 dx \\
&\lesssim \|\nabla \pi\|_{L^{r,\infty}}^{\frac{2}{3}} \||z^+|^2 + |z^-|^2\|_{L^{\frac{3r}{2}, \frac{3r-2}{2}}}^{\frac{3r-2}{2}} + C(\|w\|_{L^{3r-2}}^{3r-2} + \||z^+|\|_{L^{3r-2}}^{3r-2}) + \|\nabla|z^+|^{\frac{3r-2}{2}}\|_{L^2}^2 \\
&\leq C \|\nabla \pi\|_{L^{r,\infty}}^{\frac{2}{3}} \||z^+|^2 + |z^-|^2\|_{L^{\frac{3r}{2}, \frac{3r-2}{2}}}^{\frac{3r-2}{2}} + C(\|w\|_{L^{3r-2}}^{3r-2} + \||z^+|\|_{L^{3r-2}}^{3r-2}) + \frac{1}{256} \|\Lambda^\alpha|z^+|^{\frac{3r-2}{2}}\|_{L^2}^2.
\end{aligned}$$

where we use the estimate

$$\|\nabla|z^+|^{\frac{3r-2}{2}}\|_{L^2}^2 \leq C \|z^+\|_{L^{3r-2}}^{3r-2} \|\Lambda^\alpha|z^+|^{\frac{3r-2}{2}}\|_{L^2}^{2(1-\theta)} \leq \||z^+|^{\frac{3r-2}{2}}\|_{L^2}^{2\theta} + \frac{1}{256} \|\Lambda^\alpha|z^+|^{\frac{3r-2}{2}}\|_{L^2}^2.$$

In a similar fashion, if you do it for the equation (2.2)<sub>2</sub>, we have

$$\begin{aligned}
&\frac{d}{dt} \int_{\mathbb{R}^3} |z^-|^{3r-2} dx + \int_{\mathbb{R}^3} |\Lambda^\alpha|z^-|^{\frac{3r-2}{2}}|^2 dx \\
&\leq C \|\nabla \pi\|_{L^{r,\infty}}^{\frac{2}{3}} \||z^+|^2 + |z^-|^2\|_{L^{\frac{3r}{2}, \frac{3r-2}{2}}}^{\frac{3r-2}{2}} + C(\|w\|_{L^{3r-2}}^{3r-2} + \||z^-|\|_{L^{3r-2}}^{3r-2}) + \frac{1}{256} \|\Lambda^\alpha|z^-|^{\frac{3r-2}{2}}\|_{L^2}^2.
\end{aligned} \quad (2.13)$$

After summing up (2.12) and (2.13), using Sobolev embedding and Young's inequality,

we obtain

$$\begin{aligned}
& \frac{d}{dt} \int_{\mathbb{R}^3} (|z^+|^{3r-2} + |z^-|^{3r-2}) dx + \int_{\mathbb{R}^3} (|\Lambda^\alpha z^+|^{\frac{3r-2}{2}}|^2 + |\Lambda^\alpha z^-|^{\frac{3r-2}{2}}|^2) dx \\
& \lesssim \|\nabla \pi\|_{L^{r,\infty}}^{\frac{2}{3}} \left( \|z^+\|_{L^{\frac{6r}{3r-2},1}}^{\frac{3r-2}{2}} + \|z^-\|_{L^{\frac{6r}{3r-2},1}}^{\frac{3r-2}{2}} \right) \\
& \quad + C(\|w\|_{L^{3r-2}}^{3r-2} + \|z^+\|_{L^{3r-2}}^{3r-2} + \|z^-\|_{L^{3r-2}}^{3r-2}) \\
& \lesssim \|\nabla \pi\|_{L^{r,\infty}}^{\frac{2}{3}} \left( \|z^+\|_{L^2}^{\frac{3r-2}{2}} \|\Lambda^\alpha z^+|^{\frac{3r-2}{2}}_{L^2}^{\frac{2}{r\alpha}} + \|z^-\|_{L^2}^{\frac{3r-2}{2}} \|\Lambda^\alpha z^-|^{\frac{3r-2}{2}}_{L^2}^{\frac{2}{r\alpha}} \right) \\
& \quad + C(\|w\|_{L^{3r-2}}^{3r-2} + \|z^+\|_{L^{3r-2}}^{3r-2} + \|z^-\|_{L^{3r-2}}^{3r-2}) \\
& \lesssim \|\nabla \pi\|_{L^{r,\infty}}^{\frac{2r\alpha}{3r\alpha-3}} \left( \|z^+\|_{L^2}^{\frac{3r-2}{2}} + \|z^-\|_{L^2}^{\frac{3r-2}{2}} \right) + \frac{1}{8} \int_{\mathbb{R}_+^3} (|\Lambda^\alpha z^+|^{\frac{3r-2}{2}}|^2 + |\Lambda^\alpha z^-|^{\frac{3r-2}{2}}|^2) dx \\
& \quad + C(\|w\|_{L^{3r-2}}^{3r-2} + \|z^+\|_{L^{3r-2}}^{3r-2} + \|z^-\|_{L^{3r-2}}^{3r-2}) \\
& \lesssim \|\nabla \pi\|_{L^{r,\infty}}^{\frac{2r\alpha}{3r\alpha-3}} \left( \|z^+\|_{L^{3r-2}(\mathbb{R}^3)}^{3r-2} + \|z^-\|_{L^{3r-2}}^{3r-2} \right) + \frac{1}{8} \int_{\mathbb{R}^3} (|\Lambda^\alpha z^+|^{\frac{3r-2}{2}}|^2 + |\Lambda^\alpha z^-|^{\frac{3r-2}{2}}|^2) dx \\
& \quad + C(\|w\|_{L^{3r-2}}^{3r-2} + \|z^+\|_{L^{3r-2}}^{3r-2} + \|z^-\|_{L^{3r-2}}^{3r-2}). \tag{2.14}
\end{aligned}$$

To control  $L^{3r-2}$ -estimate for  $w$ , multiplying both side of (2.2)<sub>3</sub> by  $w|w|^{3r-4}$ , we have

$$\begin{aligned}
& \frac{1}{3r-2} \frac{d}{dt} \int_{\mathbb{R}^3} |w|^{3r-2} dx + \frac{4(3r-4)}{(3r-2)^2} \int_{\mathbb{R}^3} |\Lambda^\gamma w|^{\frac{3r-2}{2}}|^2 dx \\
& \quad + \int_{\mathbb{R}^3} |w|^{3r-2} dx + \int_{\mathbb{R}^3} |w|^{3r-4} |\operatorname{div} w|^2 dx \\
& = \underbrace{\frac{\chi}{2} \int_{\mathbb{R}^3} |w|^{3r-4} w \cdot (\nabla \times (z^+ + z^-)) dx}_{\mathcal{J}_7} + \underbrace{\int_{\mathbb{R}^3} \operatorname{div} w \cdot w \operatorname{div} (|w|^{3r-4}) dx}_{\mathcal{J}_8}. \tag{2.15}
\end{aligned}$$

As same manner as  $\mathcal{J}_2$ , the term  $\mathcal{J}_7$  is bounded by

$$\mathcal{J}_7 \lesssim (\|w\|_{L^{3r-2}}^{3r-2} + \|z^+\|_{L^{3r-2}}^{3r-2} + \|z^-\|_{L^{3r-2}}^{3r-2}) + \frac{1}{256} \|\Lambda^\gamma w|^{\frac{3r-2}{2}}\|_{L^2}^2. \tag{2.16}$$

For  $\mathcal{J}_8$ , we get

$$\begin{aligned}
\mathcal{J}_8 & \leq C \|\nabla w|^{\frac{3r-2}{2}}\|_{L^2}^2 + \frac{1}{256} \int_{\mathbb{R}^3} |w|^{3r-4} |\operatorname{div} w|^2 dx \\
& \leq C \|w\|_{L^{3r-2}}^{3r-2} + \frac{1}{256} \left( \|\Lambda^\gamma w|^{\frac{3r-2}{2}}\|_{L^2}^2 + \|w|^{\frac{3r-4}{2}} |\operatorname{div} w|\|_{L^2}^2 \right).
\end{aligned} \tag{2.17}$$

Summing up all estimates (2.14)–(2.17), we have

$$\begin{aligned}
& \frac{d}{dt} (\|z^+\|_{L^{3r-2}(\mathbb{R}^3)}^{3r-2} + \|z^-\|_{L^{3r-2}(\mathbb{R}^3)}^{3r-2} + \|w\|_{L^{3r-2}(\mathbb{R}^3)}^{3r-2}) \\
& \lesssim \|\nabla \pi\|_{L^{r,\infty}}^{\frac{2r\alpha}{3r\alpha-3}} \left( \|z^+\|_{L^{3r-2}(\mathbb{R}^3)}^{3r-2} + \|z^-\|_{L^{3r-2}}^{3r-2} \right) + (\|w\|_{L^{3r-2}}^{3r-2} + \|z^+\|_{L^{3r-2}}^{3r-2} + \|z^-\|_{L^{3r-2}}^{3r-2}).
\end{aligned}$$

Let  $\mathcal{Y}(t) := \|z^+\|_{L^{3r-2}(\mathbb{R}^3)}^{3r-2} + \|z^-\|_{L^{3r-2}(\mathbb{R}^3)}^{3r-2} + \|w\|_{L^{3r-2}(\mathbb{R}^3)}^{3r-2}$  and then (2) becomes

$$\mathcal{Y}(t) \leq C \|\nabla \pi\|_{L^{r,\infty}(\mathbb{R}^3)}^{\frac{2r\alpha}{3r\alpha-3}} \mathcal{Y}(t) + \mathcal{Y}(t).$$

As the previous way, it allow us to finish the proof of Theorem 1.1.  $\square$

### 3 Proof of Theorem 1.2

For this, according to the argument in [22] or [12], we can establish a Serrin's type regularity criterion on the gradient of pressure function  $\pi$ . Indeed, from (2.8), we know

$$\begin{aligned} & \frac{1}{4} \frac{d}{dt} \|(u, b, w)\|_{L^4}^4 + \|\nabla(|u|^2, |b|^2, |w|^2)\|_{L^2}^2 \\ & + \|u\|\|\nabla u\|_{L^2}^2 + \|b\|\|\nabla b\|_{L^2}^2 + \|w\|\|\nabla w\|_{L^2}^2 + 2\chi\|w\|_{L^4}^4 + \kappa\|w\|\operatorname{div} w\|_{L^2}^2 \\ & \lesssim \underbrace{\int_{\mathbb{R}^3} |\nabla \pi| |u|^3 |\nabla|u|^2| dx}_{\mathcal{J}_1} + \underbrace{\int_{\mathbb{R}^3} (b \cdot \nabla) b \cdot |u|^2 u dx}_{\mathcal{J}_2} \\ & + \underbrace{\frac{1}{2} \int_{\mathbb{R}^3} \nabla(|b|^2) \cdot |u|^2 u dx}_{\mathcal{J}_3} + \underbrace{\int_{\mathbb{R}^3} (b \cdot \nabla) u \cdot |b|^2 b dx dt}_{\mathcal{J}_4} + \underbrace{\frac{\chi}{2} \int_{\mathbb{R}^3} |w|^2 w \cdot (\nabla \times u) dx}_{\mathcal{J}_5} \\ & + \underbrace{\frac{\chi}{2} \int_{\mathbb{R}^3} |w|^2 w \cdot (\nabla \times u) dx}_{\mathcal{J}_6} - \underbrace{\int_{\mathbb{R}^3} \operatorname{div} w (w \cdot \nabla|w|^2) dx}_{\mathcal{J}_7} \end{aligned} \quad (3.1)$$

For the result,  $\mathcal{J}_1$  only has been changed as follows: for  $\gamma > 1$

$$\begin{aligned} & \int_{\mathbb{R}^3} \nabla \pi \cdot |u|^2 u dx \leq \|\nabla \pi\|_{L^{4,4}}^{1/2} \|\nabla \pi\|_{L^{2\gamma,\infty}}^{1/2} \|u^3\|_{L^{\frac{4\gamma}{3\gamma-2},4}} \\ & = \|\nabla \pi\|_{L^{2,2}}^{1/2} \|\nabla \pi\|_{L^{\gamma,\infty}}^{1/2} \|u\|_{L^{\frac{12\gamma}{3\gamma-2},4}}^3 \leq \frac{1}{4} \|\nabla \pi\|_{L^2}^2 + C \|\nabla \pi\|_{L^{\gamma,\infty}}^{3/2} \|u\|_{L^{\frac{12\gamma}{3\gamma-2},4}}^4 \\ & \leq \frac{1}{4} \|\nabla \pi\|_{L^2}^2 + C \|\nabla \pi\|_{L^{\gamma,\infty}}^{\frac{2}{3}} \|u\|_{L^{\frac{6\gamma}{3\gamma-2},2}}^2 \\ & \leq \frac{1}{4} \|\nabla \pi\|_{L^2}^2 + C \|\nabla \pi\|_{L^{\gamma,\infty}}^{\frac{2}{3}} \|u\|_{L^{2,2}}^{2(1-\frac{1}{\gamma})} \|\nabla|u|^2\|_{L^{2,2}}^{\frac{2}{\gamma}} \\ & \leq \frac{1}{16} \|\nabla \pi\|_{L^2}^2 + \frac{1}{8} \|\nabla|u|^2\|_{L^2}^2 + C \|\nabla \pi\|_{L^{\gamma,\infty}}^{\frac{2\gamma}{3(\gamma-1)}} \|u\|_{L^4}^4, \end{aligned}$$

and thus

$$\mathcal{J}_1 \leq \frac{1}{4} \|\nabla \pi\|_{L^2}^2 + \frac{1}{16} \|\nabla|u|^2\|_{L^2}^2 + C \|\nabla \pi\|_{L^{\gamma,\infty}}^{\frac{2\gamma}{3(\gamma-1)}} \|u\|_{L^4}^4$$

Using the following estimate,

$$\|\nabla v\|_{L^2}^2 \lesssim \|(u \cdot \nabla)u + b \cdot \nabla)b\|_{L^2}^2 \lesssim \|u\|\|\nabla u\|_{L^2}^2 + \|b\|\|\nabla b\|_{L^2}^2.$$

we get

$$\mathcal{J}_1 \leq C\|\nabla\pi\|_{L^{\gamma,\infty}}^{\frac{2\gamma}{3(\gamma-1)}}\|u\|_{L^4}^4 + \frac{1}{8}(\|u\|\|\nabla u\|_{L^2}^2 + \|b\|\|\nabla b\|_{L^2}^2).$$

Using the integration by parts,  $\mathcal{J}_2$ ,  $\mathcal{J}_3$  and  $\mathcal{J}_4$  is bounded by

$$\begin{aligned} \int_0^T \int_{\mathbb{R}^3} |u||b|^2(\nabla|u|^2 + \nabla|b|^2)dxdt &\leq C(\|(|u|^2 + |b|^2)b\|_{L^2}^2 + \frac{1}{16}(\|\nabla|u|^2\|_{L^2}^2 + \|\nabla|b|^2\|_{L^2}^2) \\ &\leq C\|b\|_{L^{a_1}}^{\frac{2a_1}{a_1-3}}(\|u\|^2\|_{L^2}^2 + \|b\|^2\|_{L^2}^2) + \frac{1}{16}(\|\nabla|u|^2\|_{L^2}^2 + \|\nabla|b|^2\|_{L^2}^2) \end{aligned}$$

where we use the following inequality:

$$\|b\|_{L^{a_1}}^2\|u\|^2\|_{L^{\frac{2a_1}{a_1-2}}}^2 \lesssim \|b\|_{L^{a_1}}^2\|u\|^2\|_{L^2}^{2(1-\frac{3}{a_1})}\|\nabla|u|^2\|_{L^2}^{\frac{6}{a_1}} \leq C\|b\|_{L^{a_1}}^{\frac{2a_1}{a_1-3}}\|u\|^2\|_{L^2}^2 + \frac{1}{16}\|\nabla|u|^2\|_{L^2}^2$$

In a similar way, for  $\mathcal{J}_5$  and  $\mathcal{J}_6$ , it shows

$$|J_5| \leq C \int_{\mathbb{R}^3} (|u|^4 + |w|^4) dx + \frac{1}{16} \int_{\mathbb{R}^3} \|w\|\|\nabla w\|^2 dx,$$

and

$$|J_6| \leq C \int_{\mathbb{R}^3} |\operatorname{div} w|^2 dx + \frac{1}{16} \int_{\mathbb{R}^3} |\nabla|w|^2|^2 dx.$$

Plugging this into (3.1), we get

$$\begin{aligned} \frac{d}{dt} \|(u, b, w)\|_{L^4}^4 + \|\nabla(|u|^2, |b|^2, |w|^2)\|_{L^2}^2 + \|u\|\|\nabla u\|_{L^2}^2 + \|b\|\|\nabla b\|_{L^2}^2 + \|w\|\|\nabla w\|_{L^2}^2 \\ \lesssim \|\nabla\pi\|_{L^{q,\infty}(\mathbb{R}^3)}^p \|(u, b, w)\|_{L^4}^4 + \|b\|_{L^{a_1}}^{\frac{2a_1}{a_1-3}}\|(u, b, w)\|_{L^4}^4 \\ \lesssim \|\nabla\pi\|_{L^{q_\kappa,\infty}(\mathbb{R}^3)}^{p_\kappa} \|(u, b, w)\|_{L^4}^4 + \|b\|_{L^{a_1}}^{\frac{2a_1}{a_1-3}}\|(u, b, w)\|_{L^4}^4 \\ \lesssim \|\nabla\pi\|_{L^{q,\infty}(\mathbb{R}^3)}^{p(1-\kappa)} \|\nabla\Pi\|_{L^2(\mathbb{R}^3)}^{c_1\kappa} \|(u, b, w)\|_{L^4}^4 + \|b\|_{L^{a_1}}^{\frac{2a_1}{a_1-3}}\|(u, b, w)\|_{L^4}^4 + \|b\|_{L^{a_1}}^{\frac{2a_1}{a_1-3}}\|(u, b, w)\|_{L^4}^4 \\ \leq C\|\nabla\pi\|_{L^{q,\infty}(\mathbb{R}^3)}^{p(1-\kappa)} \left\| |u|\|\nabla u\| + |b|\|\nabla b\| \right\|_{L^2(\mathbb{R}^3)}^{c_1\kappa} \|(u, b, w)\|_{L^4}^4 \\ \leq C\|\nabla\pi\|_{L^{q,\infty}(\mathbb{R}^3)}^{\frac{2p(1-\kappa)}{2-c_1\kappa}} \|u\|_{L^4(\mathbb{R}^3)}^{\frac{8}{2-c_1\kappa}} + \|b\|_{L^{a_1}}^{\frac{2a_1}{a_1-3}}\|(u, b, w)\|_{L^4}^4 + \frac{1}{8} \left( \|u\|\|\nabla u\|_{L^2(\mathbb{R}^3)}^2 + \|b\|\|\nabla b\|_{L^2(\mathbb{R}^3)}^2 \right) \\ \leq C\|\nabla\pi\|_{L^{q,\infty}(\mathbb{R}^3)}^{p(1-\delta)} \|(u, b, w)\|_{L^4(\mathbb{R}^3)}^{4(1+2\delta)} + \|b\|_{L^{a_1}}^{\frac{2a_1}{a_1-3}}\|(u, b, w)\|_{L^4}^4 \end{aligned}$$

Notice that  $2/p_\kappa + 3/q_\kappa = 3$ . Choosing  $\delta = \frac{(2-c_1)\kappa}{2-c_1\kappa}$ ,  $c_1 = \frac{4}{3}$ , it finally follows

$$\frac{d}{dt} \|(u, b, w)\|_{L^4}^4 \lesssim \|\nabla\pi\|_{L^{q,\infty}(\mathbb{R}^3)}^{p(1-\delta)} \|(u, b, w)\|_{L^4(\mathbb{R}^3)}^{4(1+2\delta)}.$$

As the previous way, it allow us to finish the proof of Theorem 1.2.

## References

- [1] L.C. Berselli and G.P. Galdi, Regularity criteria involving the pressure for the weak solutions of the Navier-Stokes equations, Proc. Amer. Math. Soc., 130 (2002) 3585–3595.
- [2] S. Bosia, V. Pata and J. C. Robinson, A weak- $L^p$  Prodi-Serrin type regularity criterion for the Navier-Stokes equations, J. Math. Fluid Mech., 16 (2014), 721–725.
- [3] D. Chae, On the regularity conditions for the Navier–Stokes and related equations, Rev. Mat. Iberoam. 23 (2007), 371–384.
- [4] M. Colombo, S. Haffter, Global regularity for the hyperdissipative Navier-Stokes equation below the critical order, ArXiv e-prints, November 2019.
- [5] A. Córdoba, D. Córdoba, A maximum principle applied to quasi-geostrophic equations, Commun. Math. Phys. 249 (2004), 511–528.
- [6] L. Deng, H. Shang, Global well-posedness for n-dimensional magneto-micropolar equations with hyperdissipation, Appl. Math. Lett. 111 (2021) 106610.
- [7] H. Duan, On regularity criteria in terms of pressure for the 3D viscous MHD equations. Appl. Anal. 91 (2012) 947–952.
- [8] J. Fan, T. Ozawa, On the regularity criteria for the generalized Navier–Stokes equations and Lagrangian averaged Euler equations, Differ. Integral Equ., 21 (2008), 443–457.
- [9] J. Fan, Y. Fukumoto, Y. Zhou, Logarithmically improved regularity criteria for the generalized Navier-Stokes and related equations. Kinet. Relat. Models 6 (2013), 545–556.
- [10] J. Fan, X. Zhong, Regularity criteria for 3D generalized incompressible magneto-micropolar fluid equations. Appl. Math. Lett. 127 (2022), Paper No. 107840, 5 pp.
- [11] L. Grafakos, Classical Fourier analysis. 2nd Edition, Springer, 2008
- [12] X. Ji, Y. Wang, W. Wei, New regularity criteria based on pressure or gradient of velocity in Lorentz spaces for the 3D Navier-Stokes equations. J. Math. Fluid Mech. 22 (2020), Art. 13, 8 pp.

- [13] J.-M. Kim. On regularity criteria via pressure for the 3D MHD equations in a half space. *Adv. Math. Phys.* 2022, Art. ID 6954802, 7 pp.
- [14] J. Malý, Advanced theory of differentiation–Lorentz spaces, March 2003  
<http://www.karlin.mff.cuni.cz/~maly/lorentz.pdf>.
- [15] Z. Li, Zhouyu, P. Niu, New regularity criteria for the 3D magneto-micropolar fluid equations in Lorentz spaces. *Math. Methods Appl. Sci.* 44 (2021), no. 7, 6056–6066.
- [16] M. Loayza, M. A. Rojas-Medar. A weak- $L^p$  Prodi-Serrin type regularity criterion for the micropolar fluid equations. *J. Math. Phys.* 57 (2016) 021512, 6 pp.
- [17] R. O’Neil, Convolution operators and  $L^{p,q}$  spaces. *Duke Math J.*, 30 (1963), 129–142.
- [18] B. Pineau, X. Yu, On Prodi-Serrin type conditions for the 3D Navier-Stokes, *Nonlinear Anal.* 190 (2020), 111612, 15 pp.
- [19] T. Suzuki, Regularity criteria of weak solutions in terms of the pressure in Lorentz spaces to the Navier-Stokes equations. *J. Math. Fluid Mech.*, 14 (2012), 653–660.
- [20] T. Suzuki, A remark on the regularity of weak solutions to the Navier-Stokes equations in terms of the pressure in Lorentz spaces. *Nonlinear Analysis: Theory, Methods & Applications*, 75 (2012 ), 3849–3853.
- [21] Y. Zhou, On regularity criteria in terms of pressure for the Navier-Stokes equations in  $\mathbb{R}^3$ , *Proc. Amer. Math. Soc.*, 134 (2005) 149–156.
- [22] Y. Zhou, Regularity criteria for the 3D MHD equations in terms of the pressure. *Internat. J. Non-Linear Mech.* 41 (2006), 1174-1180.
- [23] Y. Zhou, Regularity criteria for the generalized viscous MHD equations. *Ann. Inst. H. Poincare Anal. Non Lineaire* 24 (2007), 491–505.