

# Uncertainty relations from state polynomial optimization

Moisés Bermejo Morán<sup>1</sup> and Felix Huber<sup>1,2</sup>

<sup>1</sup>*Faculty of Physics, Astronomy and Applied Computer Science,  
Institute of Theoretical Physics, Jagiellonian University, 30-348 Kraków, Poland\**

<sup>2</sup>*Bordeaux Computer Science Laboratory (LaBRI), University of Bordeaux,  
351 cours de la Liberation, 33405 Talence, France*

(Dated: August 7, 2024)

Uncertainty relations are a fundamental feature of quantum mechanics. How can these relations be found systematically? Here we develop a semidefinite programming hierarchy for additive uncertainty relations in the variances of non-commuting observables. Our hierarchy is built on the state polynomial optimization framework, also known as scalar extension. The hierarchy is complete, in the sense that it converges to tight uncertainty relations. We improve upon upper bounds for all 1292 additive uncertainty relations on up to nine operators for which a tight bound is not known. The bounds are dimension-free and depend entirely on the algebraic relations among the operators. The techniques apply to a range of scenarios, including Pauli, Heisenberg-Weyl, and fermionic operators, and generalize to higher order moments and multiplicative uncertainty relations.

## INTRODUCTION

Quantum mechanical particles exhibit a fundamental uncertainty relation between conjugate observables. Similar to probability distributions that are related by a Fourier transform, conjugate observables do not allow for simultaneous measurements with vanishing variances. This feature both enables as well as hinders certain applications of quantum technologies [1–6]. While commuting measurements can be jointly diagonalized, anti-commuting observables are incompatible: This is taken advantage of in quantum error correction, where errors that anti-commute with some stabilizer elements are detected by the change they induce in the respective error syndromes [5]. Also the security of quantum key distribution relies crucially on the fact that a basis change leads to an unavoidable uncertainty in measurement outcomes [6]. In particular, additive uncertainty relations find applications in spectroscopy and atomic clocks [7], and quantum metrology [8]. Considerable effort has been made to derive state-independent bounds [9, 10].

To introduce our setting, consider a two-dimensional spin system with  $\sigma_x, \sigma_y, \sigma_z$  the Pauli matrices and denote  $\langle \sigma_i \rangle_\rho := \text{tr}(\rho \sigma_i)$  the expectation value of  $\sigma_i$  on state  $\rho$ . The set of a qubit density matrices is described by the Bloch ball, satisfying

$$\langle \sigma_x \rangle_\rho^2 + \langle \sigma_y \rangle_\rho^2 + \langle \sigma_z \rangle_\rho^2 \leq 1. \quad (1)$$

A straightforward consequence of this constraint is an additive uncertainty relation of the form

$$\Delta^2 \sigma_x + \Delta^2 \sigma_y + \Delta^2 \sigma_z \geq 2, \quad (2)$$

where  $\Delta^2 \sigma_i := \langle \sigma_i^2 \rangle_\rho - \langle \sigma_i \rangle_\rho^2$  and we used that  $\sigma_i^2 = 1$ . More generally, for a set of anti-commuting operators the relation  $\sum_{i=1}^n \langle \sigma_i \rangle_\rho^2 \leq 1$  holds. This follows from the decomposition  $1 - \sum_{i=1}^n \langle \sigma_i \rangle_\rho^2 = \langle (1 - \sum_{i=1}^n \sigma_i \langle \sigma_i \rangle_\rho)^2 \rangle \geq 0$  as sum of squares. A natural question is then what happens

when some of the variables commute, while some others anti-commute.

To answer this question we investigate the following quantity: Given a family of hermitian and unitary operators  $\{A_i\}_{i=1}^n$  satisfying commutation relations of the form  $A_i A_j = \pm A_j A_i$ , define the quantity

$$\beta = \sup_{\rho} \sum_{i=1}^n \langle A_i \rangle_{\rho}^2. \quad (3)$$

Here the maximization is over all states  $\rho$  and bounded operators  $A_i$  on Hilbert spaces that support  $\rho$  and the commutation relations between the  $A_i$ . Bounds on  $\beta$  have several applications in quantum information, including the characterization of entanglement [11], nonlocality [12], measurement compatibility [13], and to estimate ground state energies [14]. Alternatively, Eq. (3) can in the spirit of Eq. (2) tight additive uncertainty relation in the variances  $\Delta^2 A_i := \langle A_i^2 \rangle_{\rho} - \langle A_i \rangle_{\rho}^2$ ,

$$\sum_{i=1}^n \Delta^2 A_i \geq n - \beta. \quad (4)$$

In a recent work [11], Gois et al. provided the upper bound  $\beta \leq \vartheta(G)$ . Here  $\vartheta(G)$  is the Lovász theta number of the observables anti-commutativity graph. This graph encodes the observables' relations, with two vertices joined by an edge, if the corresponding observables anti-commute; i.e.  $i \sim j$  if  $A_i A_j = -A_j A_i$  and  $i \not\sim j$  else. The related bound by Hastings and O'Donnell [15],  $\langle \sum_{i=1}^n a_i A_i \rangle_{\rho}^2 \leq \vartheta(G)$  for all  $\|a\| \leq 1$ , has been discovered in the context of optimizing fermionic Hamiltonians. In turn,  $\beta$  is lower bounded by the independence number  $\alpha$ , that is the maximal number of disconnected vertices of  $G$ . The appearance of the Lovász theta number in this context is intriguing, as it has already been linked to nonlocality, contextuality, and quantum zero-error communication [16–19].

	# 2	# 3	# 4	# 5
$\vartheta_1$	3.2361	2.2361	2.2361	2.2361
$\vartheta_2$	3.0000	2.0000	2.0000	2.0000
$\alpha$	3	2	2	2

TABLE I. **Tight uncertainty relations.** The commutation relations of operators are encoded in a graph: two vertices are connected if the corresponding operators anti-commute and disconnected otherwise. Our hierarchy [Eq. (15)] upper bounds  $\beta = \sup_{\varrho} \sum_{i=1}^n \langle A_i \rangle_{\varrho}^2$  as  $\beta \leq \dots \leq \vartheta_2 \leq \vartheta_1$ , from which one obtains the additive uncertainty relation  $\sum_{i=1}^n \Delta^2 A_i \geq n - \beta$ . The independence number  $\alpha$  is the size of the largest set of disconnected vertices and lower bounds  $\beta$ , while the Lovász number  $\vartheta$  [Eq. (6)] provides an upper bound and coincides with the first level of our hierarchy. For the graphs shown above, the second level gives a tight uncertainty relation as  $\alpha = \beta = \vartheta_2$ , whereas  $\vartheta_1$  does not. The labeling is identical to that of Table IV.

The aim of this work is to provide a semidefinite programming hierarchy that upper bounds Eq. (3) and improves upon the Lovász bound. The first level of our hierarchy coincides with the Lovász theta number, and thus with the result of Gois et al. Furthermore, if at any level of the hierarchy the upper bound coincides with  $\alpha$  or the rank loop condition is met, then the corresponding uncertainty relation in Eq. (4) is tight. Numerical tests show that already the second level of our hierarchy improve upon the previous best upper bounds for all 1292 non isomorphic graphs with up to nine vertices for which  $\beta$  is unknown [20]. In particular, Table IV shows that the second and third levels often close the gap left by the Lovász theta number to a tight bound. We also introduce a second semidefinite programming hierarchy [Eq. (19)] that is complete and thus converges to  $\beta$ . With this we partially answer an open question by Xu et al. on how to obtain efficient upper bounds on  $\beta$  [14].

The central tool we use is state polynomial optimization by Klep et al. [21], also known as scalar extension [22, 23], and which can be thought of as a variant of the non-commutative optimization framework by Navascués, Pironio, and Acín [24] [25]. This makes the resulting bounds both independent of the local dimension and applicable to a wide range of situations. Different normalization conditions (e.g. projectors, unitaries, unipotents), commutation relations, and additive as well as multiplicative uncertainty relations in higher order moments can be treated. For example, Table II shows values of  $\beta$  for a set of Heisenberg-Weyl operators that satisfy commutation relations of the form  $A_i A_j = \zeta_{ij} A_j A_i$  with  $\zeta_{ij}$  a root of unity.

In this Letter, we will restrict the exposition to unitary

operators and quadratic expressions as in Eq. (3). In what follows, upper case letters will refer to observables as in  $\langle A \rangle_{\varrho} := \text{tr}(\varrho A)$ , while the lower case refers to letters  $a$  and words  $w$  in the state polynomial framework, whose evaluation on a state is referred to as  $\langle a \rangle$  and  $\langle w \rangle$ .

## LOVÁSZ BOUND

We introduce the upper bound on  $\beta$  in Eq. (3) given by the Lovász theta number [11]. Let  $\{A_1, \dots, A_n\}$  be a collection of hermitian unitary operators that pairwise either commute or anti-commute, for example, a collection of  $n$ -qubit Pauli operators. We encode the commutation relations  $A_i A_j = \zeta_{ij} A_j A_i$  with  $\zeta_{ij}$  in a graph  $G$  with adjacency matrix  $\Gamma_{ij} = (1 - \zeta_{ij})/2$ , where  $\zeta_{ij} \in \{+1, -1\}$ . In this way, every vertex represents an operator, with two vertices connected whenever the corresponding operators anti-commute. The Lovász bound then constrains  $\beta$  as [11, 15]

$$\beta \leq \vartheta(G). \quad (5)$$

Here  $\vartheta(G)$  is the Lovász theta number of the graph  $G$ .

A for us convenient definition of  $\vartheta(G)$  is through the optimal value of the following semidefinite program [26],

$$\begin{aligned} \vartheta(G) = \max_M \quad & \sum_{i=1}^n M_{ii} \\ \text{s.t.} \quad & M_{ii} = a_i, \\ & M_{ij} = 0 \quad \text{if } \zeta_{ij} = -1, \\ & \Delta = \begin{pmatrix} 1 & a^T \\ a & M \end{pmatrix} \geq 0. \end{aligned} \quad (6)$$

To see how Eq. (6) gives an upper bound on Eq. (3), that is  $\beta \leq \vartheta(G)$ , consider the following relaxation: For any  $\varrho$  construct a moment matrix  $\Gamma$  indexed by the set  $\{\mathbb{1}, \langle A_1^\dagger \rangle_{\varrho} A_1, \dots, \langle A_n^\dagger \rangle_{\varrho} A_n\}$  where  $\langle A_i \rangle_{\varrho} = \text{tr}(\varrho A_i)$  and  $A_0 = \mathbb{1}$ ,

$$\Gamma_{ij} = \text{tr}(\varrho \langle A_i \rangle_{\varrho} \langle A_j^\dagger \rangle_{\varrho} A_i^\dagger A_j). \quad (7)$$

Such matrix  $\Gamma$  has the form

$$\begin{matrix} & \mathbb{1} & \langle A_1^\dagger \rangle_{\varrho} A_1 & \dots & \langle A_n^\dagger \rangle_{\varrho} A_n \\ \mathbb{1} & \begin{bmatrix} 1 & \Gamma_{01} & \dots & \Gamma_{0n} \\ & \Gamma_{11} & \dots & \Gamma_{1n} \\ & & \ddots & \vdots \\ & & & \Gamma_{nn} \end{bmatrix} & & \end{matrix}. \quad (8)$$

In particular,  $\Gamma_{0i} = \Gamma_{ii} = \langle A_i \rangle_{\varrho}^2$  since  $A_i^\dagger A_i = \mathbb{1}$ , and  $\Gamma$  is positive semidefinite by construction. Also, we can set  $\Gamma_{ij} = 0$  when  $\zeta_{ij} = -1$ , because if  $\Gamma$  is feasible, so is  $(\Gamma + \Gamma^T)/2$  with the objective value unchanged. Note that

these last three properties coincide with the constraints imposed on  $\Delta$  in Eq. (6). However, a moment matrix  $\Gamma$  must additionally arise from some quantum state, and generally the set of matrices  $\Delta$  can be larger than the set of matrices  $\Gamma$  arising from states. Consequently, the optimum of Eq. (6) upper bounds  $\beta$  and the Lovász bound holds,  $\beta \leq \vartheta(G)$ .

While this approach already gives tight upper bounds in several settings, the pentagon graph in Table II with  $\beta = 2 < \sqrt{5} = \vartheta(G)$  shows that this Lovász bound is not tight in general. It turns out that a slight modification of the program in Eq. (6) can significantly strengthen the bounds. The key idea is to consider larger moment matrices, so to incorporate additional commutativity constraints between products of operators.

### A LOVÁSZ-TYPE HIERARCHY

To obtain stronger bounds, consider the set of moment matrices indexed by  $\{\langle A_i^\dagger \rangle_\rho \langle A_j^\dagger \rangle_\rho A_i A_j\}_{0 \leq i < j \leq n}$  [27],

$$\Gamma_{ij,kl} = \text{tr} \left( \rho \langle A_i \rangle_\rho \langle A_j \rangle_\rho \langle A_k^\dagger \rangle_\rho \langle A_l^\dagger \rangle_\rho (A_i A_j)^\dagger A_k A_l \right). \quad (9)$$

These have the form

$$\begin{array}{c} \mathbb{1} \\ \vdots \\ \langle A_i \rangle_\rho \langle A_j \rangle_\rho (A_i A_j)^\dagger \\ \vdots \end{array} \begin{bmatrix} \mathbb{1} & \cdots & \langle A_k^\dagger \rangle_\rho \langle A_l^\dagger \rangle_\rho A_k A_l & \cdots \\ 1 & \cdots & \Gamma_{00,kl} & \cdots \\ \ddots & \ddots & \vdots & \ddots \\ \langle A_i \rangle_\rho \langle A_j \rangle_\rho (A_i A_j)^\dagger & & \Gamma_{ij,kl} & \cdots \\ \vdots & & & \ddots \end{bmatrix}, \quad (10)$$

The submatrix  $(\Gamma_{i0,j0})_{0 \leq i < j \leq n}$  coincides with Eq. (8), while the complete matrix satisfies additional second-order commutation relations,

$$\Gamma_{ij,kl} = \zeta_{ji} \Gamma_{\underline{j},\underline{i},kl} = \zeta_{kl} \Gamma_{ij,\underline{l},\underline{k}} = \zeta_{ik} \Gamma_{\underline{k},\underline{j},il}. \quad (11)$$

Again,  $\Gamma$  is positive semidefinite.

Thus in analogy to Eq. (6), we formulate the relaxation

$$\begin{aligned} \vartheta_2(G) &= \max_M \sum_{i=1}^n M_{i0,i0} & (12) \\ \text{s.t.} \quad & M_{ij,kl} = \zeta_{ji} M_{j,i,kl}, \\ & M_{ij,kl} = \zeta_{kl} M_{i,j,lk}, \\ & M_{ij,kl} = \zeta_{ik} M_{k,j,il}, \\ & M_{ii,kl} = M_{0i,kl}, \\ & M \geq 0. \end{aligned}$$

As in the previous section, we can restrict to real matrices: if  $M$  is feasible, then so is  $(M + M^\dagger)/2$  with unchanged objective value. Thus we can set  $M_{ij,kl} = 0$  if  $(A_i A_j)^\dagger A_k A_l = -(A_k A_l)^\dagger A_i A_j$ . Later on, we cover the

general case with non-hermitian operators and complex phases [Eq. (19)], which requires complex matrices.

In analogy to the argument made for the Lovász bound, any  $\Gamma$  matrix arising from a state also satisfies the constraints imposed on the matrix  $M$  in Eq. (12). Thus the set of valid moment matrices  $\Gamma$  is contained in the set of feasible matrices  $M$ . Consequently,  $\beta \leq \vartheta_2(G)$ . Additionally, as the submatrix  $(M_{i0,j0})_{i,j=0}^n$  coincides with  $\Delta$  of Eq. (6), we have the strengthening

$$\beta \leq \vartheta_2(G) \leq \vartheta(G). \quad (13)$$

For the pentagon graph in Table II, this enlarged program already yields a tight bound on  $\beta = 2 = \vartheta_2(G)$ . For the remaining graphs with five vertices the Lovász number equals the independence number, and thus  $\vartheta_2(G)$  is tight for all graphs with up to five vertices.

The argument above easily generalizes to larger indexing sequences that incorporate commutation relations between products of up to  $k$  operators. Denote  $\vec{i} = (i_1, \dots, i_k)$ , a moment matrix  $\Gamma$  indexed by  $\{\langle A_{i_1}^\dagger \rangle_\rho \dots \langle A_{i_k}^\dagger \rangle_\rho A_{i_1} \dots A_{i_k}\}_{i_1, \dots, i_k=0}^n$  where all non-zero indices are pairwise distinct and ordered increasingly. This has entries

$$\begin{aligned} \Gamma_{\vec{i},\vec{j}} &= \langle A_{i_1} \rangle_\rho \dots \langle A_{i_k} \rangle_\rho \langle A_{j_1}^\dagger \rangle_\rho \dots \langle A_{j_k}^\dagger \rangle_\rho \\ &\quad \times (\langle A_{i_1} \dots A_{i_k} \rangle^\dagger A_{j_1} \dots A_{j_k})_\rho \end{aligned} \quad (14)$$

satisfies the semidefinite program

$$\begin{aligned} \vartheta_k(G) &= \max_M \sum_{i=1}^n M_{0\dots 0i,0\dots 0i} & (15) \\ \text{s.t.} \quad & M_{\pi(a,b)(\vec{i},\vec{j})} = \xi_{ab} M_{\vec{i},\vec{j}}, \\ & M_{i\dots i_k,\vec{j}} = M_{0i\dots i_k,\vec{j}}, \\ & M \geq 0. \end{aligned}$$

Here,  $\pi(a, b)$  is the transposition exchanging the  $a$ 'th and  $b$ 'th elements in the joint sequence  $(i_k, \dots, i_1, j_1, \dots, j_k)$  (note the reversed indexing in the index  $i$  arising from the dagger in the moment matrix) and

$$\xi_{ab} = \prod_{a < c \leq b} \zeta_{ac} \zeta_{cb} \quad (16)$$

is the factor that appears from the exchange.

By construction, Eq. (15) defines a hierarchy of upper bounds  $\vartheta_k(G)$

$$\beta(G) \leq \dots \leq \vartheta_k(G) \leq \dots \leq \vartheta_1(G) = \vartheta(G). \quad (17)$$

We show in the next section that the relaxations  $\vartheta_k$  in Eq. (15), in particular  $\vartheta$  in Eq. (6), can be enlarged to a complete hierarchy of semidefinite programs converging to the optimal value of Eq. (3).

## A COMPLETE HIERARCHY

We now present a complete hierarchy that converges to  $\beta$ . Let  $\zeta$  be a matrix encoding the commutation relations between a collection of operators  $\{A_1, \dots, A_n\}$ . The upper bound  $\beta$  in Eq. (3) can be formulated as the following optimization problem over expectations  $\langle \cdot \rangle_\rho$ ,

$$\begin{aligned} \sup_{\rho} \quad & \sum_{i=1}^n |\langle A_i \rangle_\rho|^2 \\ \text{s.t.} \quad & A_i^\dagger A_i = A_i A_i^\dagger = 1, \\ & A_i A_j = \zeta_{ij} A_j A_i, \\ & \langle \mathbf{1} \rangle_\rho = 1. \end{aligned} \quad (18)$$

We now show how to tackle this problem with a variant of non-commutative polynomial optimization [24], allowing for non-linear expressions in the expectations, known as state polynomial optimization [21] or scalar extension [28].

To solve this type of optimization problem, consider words (or non-commutative monomials)  $w = a_{j_1} \cdots a_{j_p}$  built from letters  $\{a_i\}_{i=1}^n$  and expectation value of words denoted by  $\langle w \rangle = \langle a_{j_1} \cdots a_{j_p} \rangle$ . The most general monomials in words and expectations are then of the form  $w = w_0 \langle w_1 \rangle \cdots \langle w_m \rangle$ . Such formal state monomials can be added and multiplied to form state polynomials. The involution of a word is defined by  $(a_1 \dots a_n)^* = a_n^* \cdots a_1^*$ , which plays the role of the adjoint of an operator in the formal setting. In particular,  $a_i^* = a_i$  when the operators are required to be hermitian. Also, expectations behave as scalars, so that  $v \langle w \rangle = \langle w \rangle v$  for all  $v, w$ . Finally,  $\langle v \langle w \rangle \rangle = \langle \langle v \rangle \langle w \rangle \rangle = \langle v \rangle \langle w \rangle$  and  $(v \langle w \rangle)^* = v^* \langle w \rangle^* = \langle w \rangle^* v^*$ , while  $e$  is the identity (or empty) word satisfying  $we = w = ew$  for all words  $w$ .

For our problem, one additionally imposes the constraints  $a_i a_j = \zeta_{ij} a_j a_i$  arising from the commutation relations  $A_i A_j = \zeta_{ij} A_j A_i$ . Likewise, the constraint  $A_i^\dagger A_i = A_i A_i^\dagger = \mathbf{1}$  gives rise to  $a_i^* a_i = a_i a_i^* = e$ . For Pauli matrices, this reduces all monomials to square-free monomials.

Now consider a moment matrix  $M_\ell$ , indexed by all state monomials of degree at most  $\ell$ , whose entries are  $M_\ell(v, w) = \langle v^* w \rangle$ . Then, the optimal solution of Eq. (18) is approximated from above with the following hierarchy of semidefinite programs with  $\ell \in \mathbb{N}$  [21, Lemma 6.5],

$$\begin{aligned} \nu_\ell(\zeta) = \max \quad & \sum_{i=1}^n |\langle a_i \rangle|^2 \\ \text{s.t.} \quad & \langle \text{id} \rangle = 1, \\ & M_\ell(v, w) = M_\ell(x, y) \quad \text{when} \quad \langle v^* w \rangle = \langle x^* y \rangle, \\ & M_\ell \geq 0, \end{aligned} \quad (19)$$

where one imposes on the entries  $M_\ell$  all relations arising from  $a_i a_j = \zeta_{ij} a_j a_i$  and  $a^* a = a a^* = e$ . By increasing

the degree  $\ell$ , these relaxations converge to the optimal solution of Eq. (3) [21, Theorem 5.5 and Proposition 6.7],

$$\lim_{\ell \rightarrow \infty} \nu_\ell(\zeta) = \beta. \quad (20)$$

For this result to hold, it is necessary that the optimization is over a bounded set of operators. The technical condition is that the set is Archimedean, that is, there exists a constant  $C > 0$ , such that  $\sum_{i=1}^n a_i a_i^* \leq C$ . In our problem this is satisfied with  $C = n$ . Additionally, when the rank loop condition is met, that is  $\text{rank}(M_\ell) = \text{rank}(M_{\ell+1})$  (this is called a flat extension), then the optimum has been reached [21, Proposition 6.10].

It can now be seen that  $\vartheta_k(G)$  of Eq. (15) arises from the state polynomial optimization framework [Eq. (19)] where one considers only state monomials of the form

$$\{\langle a_{i_1}^* \rangle \cdots \langle a_{i_k}^* \rangle a_{i_1} \cdots a_{i_k}\}_{i_1, \dots, i_k=0}, \quad (21)$$

with all non-zero indices pairwise distinct.

## NON-HERMITIAN OPERATORS

Note that both the complete and relaxed hierarchies can easily deal with operators that neither commute nor anti-commute, but for which there is a complex phase  $\zeta_{ij} \in \mathbb{C}$ . Then for every letter  $a$  we define two new symbols to denote its real  $\Re w$  and imaginary  $\Im w$  part. Additionally, for each word  $w$  impose the constraints  $\Re w = \Re w^*$  and  $\Im w = -\Im w^*$ . The positive semidefinite constraint on the complex moment matrix  $M_\ell = \Re M_\ell + i \Im M_\ell \geq 0$  is equivalent to

$$\begin{pmatrix} \Re M_\ell & -\Im M_\ell \\ \Im M_\ell & \Re M_\ell \end{pmatrix} \geq 0. \quad (22)$$

Table II shows the first two levels of the hierarchy of Lovász relaxations for the Heisenberg-Weyl operators computed through the complex to real mapping of Eq. (22). These operators are not hermitian in dimensions  $d$  greater than two, and we use weighted graphs to specify their commutation relations where  $\zeta_{ij}$  is a  $d$ 'th root of unity. Note that the cost of solving the problem in Eq. (19) does not increase with  $d$  but only with the number of operators and the level of the hierarchy  $k$ .

## CUTTING PLANES

The Lovász bound in Eq. (6) can be strengthened with additional constraints, for instance with so-called odd-hole inequalities [26]. Let  $G$  be the anti-commutation graph of some hermitian operators, and let  $H$  be a subset of vertices of  $G$  inducing a cycle (i.e. a hole). Then

$$\sum_{i \in H} \langle A_i \rangle_\rho^2 \leq \left\lfloor \frac{|H|}{2} \right\rfloor. \quad (23)$$




	$d = 2$	$d = 3$	$d = 5$	
	$\vartheta_1$	1.0000	1.0000	1.4906
	$\vartheta_2$	1.0000	1.0000	1.4575
	lb	1.0000	1.0000	1.4500
	$\vartheta_1$	2.2361	2.0000	2.3058
	$\vartheta_2$	2.0000	2.0000	2.2001
	lb	2.0000	1.7770	1.2690
	$\vartheta_1$	1.0000	2.0000	2.0000
	$\vartheta_2$	1.0000	2.0000	2.0000
	lb	1.0000	2.0000	1.9940

TABLE II. **Upper bounds  $\vartheta_k$  for higher dimensional spin operators**, obtained with an adaptation of Eq. (15) for commutation relations given by  $d$ 'th roots of unity,  $A_i A_j = e^{2\pi i/d} A_j A_i$  when  $i \sim j$ . This requires complex moment matrices that can be evaluated through Eq. (22). Lower bounds (lb) are obtained from sampling Haar random states and evaluating them on Heisenberg-Weyl operators [Eq. (36)]. Displacement operators [Eq. (40)] give the same bounds.


		
$\vartheta_1$	2.2361	2.2356
$\vartheta_2$	2.0000	2.0000
$\alpha$	2	

TABLE III. **Cutting planes in graph #3**. Left:  $\vartheta_k$  of Eq. (15). Right:  $\vartheta_k$  strengthened with two intertwined 5-holes inequalities [Eq. (23)].

This constraint acts as a half-plane in the semidefinite program, and can strengthen the Lovász bound for odd  $|H|$ . For graphs with up to seven vertices with non-overlapping holes, these additional constraints are enough to tighten the bound for  $\beta$ . However, these are not enough when the holes are intertwined as in Table III (#3 in Table IV). More generally, one can impose constraints that arise from higher levels in the hierarchy [Eq. (19)] on selected subgraphs.

## CONCLUSIONS

We presented in Eq. (19) a complete semidefinite programming hierarchy converging to tight uncertainty relations. The reduced formulation of this hierarchy in Eq. (15) can be seen as a natural generalization of the Lovász bound [11]. Interestingly, the second level of this reduced hierarchy already provides tight bounds on

$\beta$  for all graphs with no more than six vertices. This answers an open question by Xu et al. on how to efficiently bound  $\beta$  [14]. Additionally, our hierarchy applies to a wide range of scenarios, including the  $n$ -qubit Pauli group, generalised Pauli operators for higher-dimensional systems, Clifford algebras, and fermionic operators. It also applies to higher order moments and multiplicative uncertainty relations. Several questions remain:

- Can a similar approach be derived for entropic uncertainty relations [29–34]?
- Can uncertainty relations for Gell-Mann matrices be found that make use of the structure constants of  $SU(3)$ , in analogy to the Lovász bound?
- Is the hierarchy of relaxations  $\vartheta_k(G)$  in Eq. (15) complete, in the sense that it converges to  $\beta$ ?
- Can sparsity [35] or symmetry [36–40] reductions boost efficiency of the SDP?
- Which hyperplane constraints improve the Lovász bound most without incurring a too high computational overhead [26]? How do these constraints differ from the classical case?
- The weight enumerators describing quantum error correcting codes can be expressed as sums over  $|\langle A_i \rangle|^2$ . Can also here a Lovász bound be established to upper bound the size of a quantum code?
- Can structure theorems for quasi-Clifford algebras or quantum tori [41, 42] help to derive algebraic bounds? Are there any significant changes with respect to the choice of the matrix basis [43], for example when considering nice error bases [44]?

*Acknowledgements.*— MBM and FH were supported by the Foundation for Polish Science through TEAM-NET (POIR.04.04.00-00-17C1/18-00). We thank Carlos de Gois, Otfried Gühne, Kiara Hansenne, Igor Klep, Victor Magron, Stefano Pironio, Alex Pozas, René Schwonnek, Jurij Volčič, Zhen-Peng Xu, and Karol Życzkowski for helpful comments and fruitful discussions. We thank the organizers of the Quantum Corona 2022 workshop for hospitality and the friendly atmosphere.

## Further applications

Already for hermitian operators, Eq. (3) finds applications in the characterization of entanglement and non-locality. For example, Eq. (1) gives the entanglement witness [11]

$$W = \mathbf{1} \otimes \mathbf{1} - \sigma_x \otimes \sigma_x - \sigma_y \otimes \sigma_y + \sigma_z \otimes \sigma_z. \quad (24)$$

It holds that  $\langle W \rangle_\rho \geq 0$  for all separable states  $\rho$ , whereas a joint eigenstate of  $\sigma_x \otimes \sigma_x$ ,  $\sigma_y \otimes \sigma_y$ , and  $\sigma_z \otimes \sigma_z$  achieves a value  $\langle W \rangle_\rho = -2$ .

Similar constraints involving Pauli strings bound the quantum value of Bell inequalities [45] and Bell monogamy [12]. For instance, let  $\mathcal{L}$  be a Bell inequality with two measurements with two outcomes between



two parties. Normalizing the value over classical correlations as  $\mathcal{L} \leq 1$ , the optimal quantum value is obtained by the value in Eq. (3) for the following Pauli strings

$$\mathcal{L}^2 \leq \langle \sigma_x \otimes \sigma_x \rangle^2 + \langle \sigma_x \otimes \sigma_y \rangle^2 + \langle \sigma_y \otimes \sigma_x \rangle^2 + \langle \sigma_y \otimes \sigma_y \rangle^2 \leq 2. \quad (25)$$

This gives the Tsirelson's bound for the CHSH inequality. More generally, upper bounds on Eq. (3) for Pauli strings yield upper bounds for the quantum value of Bell inequalities and Bell monogamy involving only dichotomic measurements [12, 45]. It is an interesting question whether the bound in Eq. (25) extends to measurements with more than two outcomes.

As noticed in Ref. [14], Eq. (3) also bounds the ground state energy of the Hamiltonian  $H = \sum_i^n A_i$ , since  $\langle H \rangle^2 \leq n \sum_i^n \langle A_i \rangle^2$ . These bounds are determined by the algebraic relations between the terms and thus are independent of the physical dimension.

### The Lovász theta number

A graph  $G = (V, E)$  consists of a set  $V$  of *vertices* and a set  $E \subset V \times V$  of *edges* between them. Two vertices  $u, v \in V$  are connected ( $u \sim v$ ) when  $(u, v) \in E$ , and disconnected or *independent* otherwise. The *complement* of  $G$  is  $\bar{G} = (V, \bar{E})$ , where  $(u, v) \in \bar{E}$  if and only if  $(u, v) \notin E$ .

The *independence number*  $\alpha(G)$  is the maximum number of pairwise independent vertices of  $G$ . The *chromatic number*  $\chi(G)$  is the minimum number of different colors needed to assign to the vertices of  $G$ , such that no connected vertices share the same color. These quantities, also called *graph invariants*, encode key combinatorial properties of  $G$  and are NP-complete [46].

However, some graph invariants capture properties of the graph that are easier to compute. Of particular interest is the *Lovász number*  $\vartheta(G)$ , which is sandwiched by the two quantities described above [47],

$$\alpha(G) \leq \vartheta(G) \leq \chi(\bar{G}). \quad (26)$$

It can be computed in polynomial time since it can be defined through a semidefinite program [48],

$$\begin{aligned} \vartheta(G) = \max_M \quad & \sum_{i,j=1}^n M_{ij} \\ \text{s.t.} \quad & \text{tr}(M) = 1 \\ & M_{ij} = 0 \quad \text{if } i \sim j \\ & M \geq 0. \end{aligned} \quad (27)$$

Equivalent definitions for  $\vartheta(G)$  are [26],

$$\begin{aligned} \vartheta(G) = \max_M \quad & \sum_{i=1}^n M_{ii} \\ \text{s.t.} \quad & M_{ii} = a_i \\ & M_{ij} = 0 \quad \text{if } i \sim j \\ & \Delta = \begin{pmatrix} 1 & a^T \\ a & M \end{pmatrix} \geq 0, \end{aligned} \quad (28)$$

and [47]

$$\begin{aligned} \vartheta(G) = \max_M \quad & \lambda_{\max}(M) \\ \text{s.t.} \quad & M_{ii} = 1 \\ & M_{ij} = 0 \quad \text{if } i \sim j \\ & M \geq 0, \end{aligned} \quad (29)$$

where  $\lambda_{\max}(M)$  denotes the maximum eigenvalue of  $M$ .

Given a collection of hermitian unitary operators  $A_1, \dots, A_n$  that mutually either commute or anti-commute, define the *anti-commutativity graph*  $G$  on vertices  $\{1, \dots, n\}$  where  $i \sim j$  if  $A_i A_j = -A_j A_i$ . The quantity

$$\beta = \sup_{\rho} \sum_{i=1}^n \langle A_i \rangle_{\rho}^2. \quad (30)$$

is then a graph invariant that is defined solely through the commutation relations encoded in  $G$ . Commuting operators can be jointly diagonalized, thus it is clear that  $\alpha(G) \leq \beta(G)$ : a joint eigenstate simultaneously achieves the value  $\langle A_i \rangle_{\rho}^2 = 1$  for each term in an independent set. On the other hand, Ref. [11] showed that  $\beta(G) \leq \vartheta(G)$  by using the definition in Eq. (29).

As an example, take the following five two-qubit Pauli observables

$$\sigma_x \otimes \sigma_x, \sigma_x \otimes \sigma_y, \text{id} \otimes \sigma_x, \sigma_y \otimes \sigma_z, \sigma_y \otimes \sigma_x. \quad (31)$$

Their anti-commutativity graph is the pentagon  $G = C_5$  shown in Table IV. The second level of the hierarchy improves on the Lovász bound, closing the gap with the independence number and thus giving the tight bound

$$2 = \alpha(C_5) = \beta(C_5) = \vartheta_2(C_5) < \vartheta_1(C_5) = \sqrt{5}. \quad (32)$$

### Comparison with previous work

Upper bounds in Ref. [14] are obtained fixing a representation of the operators  $(A_1, \dots, A_n)$  as Pauli strings in dimension  $2^N$ , through the separability problem

$$\begin{aligned} \max_{\sigma} \quad & \sum_{i=1}^n \langle A_i \otimes A_i \rangle_{\sigma} \\ \text{s.t.} \quad & \sigma \in \text{SEP}. \end{aligned} \quad (33)$$

Here the maximization runs over separable states on  $\mathbb{C}^{2^{\otimes N}} \otimes \mathbb{C}^{2^{\otimes N}}$ . This problem can be outer-approximated with semidefinite programs based on symmetric extension. Level  $k \in \mathbb{N}$  of this hierarchy corresponds to

$$\begin{aligned} \max_{\sigma} \quad & \sum_{i=1}^n \langle A_i \otimes A_i \rangle_{\sigma} \\ \text{s.t.} \quad & \sigma \in \text{Sym}_k. \end{aligned} \quad (34)$$

Here, the maximization runs over all states  $\sigma = \sigma_{AB_1 \dots B_k}$  on  $(\mathbb{C}^2)^{\otimes N} \otimes (\mathbb{C}^2)^{\otimes (N(1+k))}$  that are invariant under exchange of the subsystems  $B_1, \dots, B_k$ , with positive partial transpose across all bipartitions. The dimension of this hierarchy grows as  $2^{N(2+k)}$ , which renders the relaxations intractable quickly.

For instance, take the graph  $G = \bar{C}_7$ , the complement of the cycle with seven vertices. These operators can be represented with Pauli strings of  $N = 3$  qubits. The first extension has size  $2^9 = 512$ , which is already quite large for a desktop computer to solve.

Our approach based on state polynomial optimization [Eq. (18)] does not fix a representation, and is thus dimension-independent. The size of the relaxations is given by the size of the indexing sequence for the moment matrix in Eq. (19), which allows a finer control and flexibility in the size. In particular, our hierarchy  $\vartheta_k(G)$  in Eq. (15) requires for a graph  $G$  with  $n$  vertices to solve SDPs of size  $1+n$  for  $\vartheta_1(G)$  [Eq. (6)], size  $1+n(n+1)/2$  for  $\vartheta_2(G)$  [Eq. (12)], and size  $2^n + 1$  for  $\vartheta_n(G)$ . For a graph  $G$  with seven vertices such as  $\bar{C}_7$ ,  $\vartheta_7(G)$  has size  $2^7 = 128$ , which can easily be solved on a desktop computer.

In the approach by Ref. [14] the size of the resulting SDPs depends on whether the commutation relations can be represented on a small number of qubits, while it only depends on the number of observables for our hierarchies. However, with the symmetric extension approach one in principle is able to give approximation guarantees through the quantum de Finetti theorem [49].

### Small graphs

Table IV shows all 43 non-isomorphic graphs with up to seven vertices for which  $\beta < \vartheta_1$ . Strengthening  $\vartheta_1$  with odd-hole inequalities as in Eq. (23), we obtain tight bounds on  $\beta$  for 18 graphs by matching the upper bounds with the independence number. For all but 10 of these graphs it holds that  $\beta = \vartheta_2$ , and  $\beta = \vartheta_3$  holds for all but 7 graphs.

Our hierarchy also improves on the best known upper bounds for all 36 non-isomorphic graphs with eight vertices and all 1256 non-isomorphic graphs with nine vertices for which the value of  $\beta$  is unknown [20]. Already our second level [Eq. (12)] gives tight bounds

$\beta(G) = \vartheta_2(G)$  for four of those graphs, up to numerical precision.

### Non-hermitian operators

**Heisenberg-Weyl basis.** The Heisenberg-Weyl operators can be seen as a generalization of the Pauli matrices. They play a central role in the description of  $d$ -dimensional quantum spin systems [43] and in the construction of non-binary quantum codes [50]. On the space  $\mathbb{C}^d$  define the operators  $X$  and  $Z$  through [51]

$$X|j\rangle = |j+1\rangle, \quad Z|j\rangle = \omega^j|j\rangle, \quad (35)$$

where  $\omega = \exp(2\pi i/d)$  is the principal root of unity and the addition is taken modulo  $d$ . These operators are related by the quantum Fourier transform in dimension  $d$ . The collection of operators

$$\sigma(k, l)|i\rangle = X^k Z^l |i\rangle = \omega^{il} |i+k\rangle, \quad (36)$$

forms the Heisenberg-Weyl basis for the space of complex  $d \times d$  matrices. These satisfy the commutativity relations

$$\sigma(k, l)\sigma(m, n) = \omega^{lm-kn}\sigma(m, n)\sigma(k, l). \quad (37)$$

Their adjoints are given by

$$\sigma(k, l)^\dagger = \omega^{kl}\sigma(d-k, d-l), \quad (38)$$

and they form an orthogonal basis

$$\text{tr}[\sigma(k, l)^\dagger \sigma(m, n)] = \delta_{km} \delta_{ln} d. \quad (39)$$

**Displacement basis.** Introducing a phase [51],

$$D(k, l) = \omega^{kl/2} \sigma(k, l), \quad (40)$$

the adjoint operators are given by

$$D(k, l)^\dagger = (-1)^{k+l+d} D(d-k, d-l). \quad (41)$$

These are called displacement operators, and the algebra they generate is isomorphic to the one generated by Heisenberg-Weyl operators. Thus, the bounds for the non-hermitian version of Eq. (18) coincide for both choices of generators (c.f. Table II).

---

\* moises.moran@uj.edu.pl

- [1] C. A. Fuchs and A. Peres, Phys. Rev. A **53**, 2038 (1996).
- [2] O. Gühne, Phys. Rev. Lett. **92**, 117903 (2004).
- [3] E. G. Cavalcanti and M. D. Reid, J. Mod. Opt. **54**, 2373 (2007).
- [4] J. M. Renes and J.-C. Boileau, Phys. Rev. Lett. **103**, 020402 (2009).












































								
$\vartheta_1$	2.2361	3.2361	2.2361	2.2361	2.2361	4.2361	3.2361	3.2361
$\vartheta_2$	2.0000	3.0000	2.0000	2.0000	2.0000	4.0000	3.0000	3.0000
$\alpha$	2	3	2	2	2	4	3	3
								
$\vartheta_1$	3.2361	3.2361	3.2361	3.2361	3.2361	3.2361	3.2361	3.2361
$\vartheta_2$	3.0000	3.0000	3.0000	3.0000	3.0000	3.0000	3.0000	3.0000
$\alpha$	3	3	3	3	3	3	3	3
								
$\vartheta_1$	3.2361	3.1966	3.0642	3.3177	3.2361	3.1966	3.2361	3.1966
$\vartheta_2$	3.0000	3.0000	3.0000	3.0000	3.0000	3.0000	3.0000	3.0000
$\alpha$	3	3	3	3	3	3	3	3
								
$\vartheta_1$	2.2361	2.2361	2.2361	2.2361	2.2361	2.2361	2.2361	2.2361
$\vartheta_2$	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000	2.0000
$\alpha$	2	2	2	2	2	2	2	2
								
$\vartheta_1$	2.2361	2.2361	2.2361	2.2361	2.2361	2.2361	2.2361	2.2361
$\vartheta_2$	2.0363	2.0056	2.0085	2.0033	2.0249	2.0392	2.0121	2.0024
$\vartheta_3$	2.0067	2.0004	2.0017	2.0000	2.0047	2.0052	2.0006	2.0000
$\vartheta_7$	2.0013	2.0000	2.0003	2.0000	2.0014	2.0011	2.0002	2.0000
$\alpha$	2	2	2	2	2	2	2	2
								
$\vartheta_1$	2.2361	2.2361	2.1099					
$\vartheta_2$	2.0910	2.0000	2.0950					
$\vartheta_3$	2.0076	2.0000	2.0938					
$\vartheta_7$	2.0024	2.0000	2.0938					
$\alpha$	2	2	2					

TABLE IV. **Hierarchy of upper bounds** for all non-isomorphic graphs up to seven vertices for which  $\beta < \vartheta_1$ . Our hierarchy  $\vartheta_k$  in Eq. (15) improves on the bounds given by the Lovász number  $\vartheta_1$  in Eq. (6) and closes the gap for all but 7 graphs: 33, 35, 37, 38, 39, 41 and 43.



- [5] D. Gottesman, in *Quantum information science and its contributions to mathematics, Proceedings of Symposia in Applied Mathematics*, Vol. 68 (2010) pp. 13–58.
- [6] C. H. Bennett and G. Brassard, *Theor. Comput. Sci.* **560**, 7 (2014).
- [7] A. S. Sørensen and K. Mølmer, *Phys. Rev. Lett.* **86**, 4431 (2001).
- [8] G. Tóth and F. Fröwis, *Phys. Rev. Res.* **4**, 013075 (2022).
- [9] L. Dammeier, R. Schwonnek, and R. F. Werner, *New J. Phys.* **17**, 093046 (2015).
- [10] K. Szymański and K. Życzkowski, *J. Phys. A Math. Theor.* **53**, 015302 (2019).
- [11] C. de Gois, K. Hansenne, and O. Gühne, *Phys. Rev. A* **107**, 062211 (2023).
- [12] P. Kurzyński, T. Paterek, R. Ramanathan, W. Laskowski, and D. Kaszlikowski, *Phys. Rev. Lett.* **106**, 180402 (2011).
- [13] F. Loulidi and I. Nechita, *PRX Quantum* **3**, 040325 (2022).
- [14] Z.-P. Xu, R. Schwonnek, and A. Winter, “Bounding the joint numerical range of Pauli strings by graph parameters,” (2023), arXiv:2308.00753.
- [15] M. B. Hastings and R. O’Donnell, in *Proceedings of the 54th Annual ACM SIGACT Symposium on Theory of Computing* (2022) pp. 776–789.
- [16] A. Cabello, S. Severini, and A. Winter, *Phys. Rev. Lett.* **112**, 040401 (2014).
- [17] A. Acín, T. Fritz, A. Leverrier, and A. B. Sainz, *Commun. Math. Phys.* **334**, 533 (2015).
- [18] A. Acín, R. Duan, D. E. Roberson, A. B. Sainz, and A. Winter, *Discrete Appl. Math.* **216**, 489 (2017).
- [19] R. Duan, S. Severini, and A. Winter, *IEEE Trans. Inf. Theory* **59**, 1164 (2013).
- [20] Private communication with the authors of Ref. [14].
- [21] I. Klep, V. Magron, J. Volčič, and J. Wang, “State polynomials: positivity, optimization and nonlinear Bell inequalities,” (2023), arXiv:2301.12513.
- [22] A. Pozas-Kerstjens, R. Rabelo, L. Rudnicki, R. Chaves, D. Cavalcanti, M. Navascués, and A. Acín, *Phys. Rev. Lett.* **123**, 140503 (2019).
- [23] A. Pozas-Kerstjens, *Quantum information outside quantum information*, Ph.D. thesis, Universitat Politècnica de Catalunya (2019).
- [24] S. Pironio, M. Navascués, and A. Acín, *SIAM J. Optim.* **20**, 2157 (2010).
- [25] While the scalar extension hierarchy was formulated earlier, Ref. [21] shows also completeness of the hierarchy. We here use the term state polynomial optimization, as it highlights the connections to polynomial optimization best. In any case, the two terms can be used interchangeably.
- [26] L. Galli and A. N. Letchford, *Discret. Optim.* **25**, 159 (2017).
- [27] Interestingly, we observe a similar numerical behaviour with the sequence  $\{\langle (A_i A_j)^\dagger \rangle_e A_i A_j\}_{0 \leq i < j \leq n}$ .
- [28] A. Tavakoli, A. Pozas-Kerstjens, M.-X. Luo, and M.-O. Renou, *Rep. Prog. Phys.* **85**, 056001 (2022).
- [29] H. Maassen and J. B. M. Uffink, *Phys. Rev. Lett.* **60**, 1103 (1988).
- [30] S. Wehner and A. Winter, *J. Math. Phys.* **49**, 062105 (2008).
- [31] Z. Puchała, L. Rudnicki, and K. Życzkowski, *J. Phys. A Math. Theor.* **46**, 272002 (2013).
- [32] K. Korzekwa, M. Lostaglio, D. Jennings, and T. Rudolph, *Phys. Rev. A* **89**, 042122 (2014).
- [33] Y.-Y. Zhao, G.-Y. Xiang, X.-M. Hu, B.-H. Liu, C.-F. Li, G.-C. Guo, R. Schwonnek, and R. Wolf, *Phys. Rev. Lett.* **122**, 220401 (2019).
- [34] A. F. Rotundo and R. Schwonnek, “An entropic uncertainty principle for mixed states,” (2023), arXiv:2303.11382.
- [35] J. Wang and V. Magron, *Comput. Optim. Appl.* **80**, 483 (2021).
- [36] M. Ioannou and D. Rosset, “Noncommutative polynomial optimization under symmetry,” (2021), arXiv:2112.10803.
- [37] C. Bachoc, D. C. Gijswijt, A. Schrijver, and F. Vallentin, “Invariant semidefinite programs,” in *Handbook on semidefinite, conic and polynomial optimization* (Springer, 2012) pp. 219–269.
- [38] F. Permenter and P. A. Parrilo, *Math. Program.* **181**, 51 (2020).
- [39] D. Brosch and E. de Klerk, *Optim. Methods Softw.* **37**, 2001 (2022).
- [40] L. T. Ligthart and D. Gross, *J. Math. Phys.* **64**, 072201 (2023).
- [41] H. Gastineau-Hills, *J. Aust. Math. Soc.* **32**, 1 (1982).
- [42] A. N. Panov, *Mathematical Notes* **69**, 537 (2001).
- [43] A. Vourdas, *Rep. Prog. Phys.* **67**, 267 (2004).
- [44] A. Klappenecker and M. Rötteler, *IEEE Trans. Inf. Theory* **48**, 2392 (2002).
- [45] M. Żukowski and Č. Brukner, *Phys. Rev. Lett.* **88**, 210401 (2002).
- [46] R. M. Karp, “Reducibility among combinatorial problems,” in *Complexity of Computer Computations: Proceedings of a symposium on the Complexity of Computer Computations, held March 20–22, 1972, at the IBM Thomas J. Watson Research Center, Yorktown Heights, New York, and sponsored by the Office of Naval Research, Mathematics Program, IBM World Trade Corporation, and the IBM Research Mathematical Sciences Department* (Springer US, Boston, MA, 1972) pp. 85–103.
- [47] D. E. Knuth, “The sandwich theorem,” (1993), arXiv:math/9312214.
- [48] L. Lovász, *IEEE Trans. Inf. Theory* **25**, 1 (1979).
- [49] M. Christandl, R. König, G. Mitchison, and R. Renner, *Commun. Math. Phys.* **273**, 473 (2007).
- [50] A. Ketkar, A. Klappenecker, S. Kumar, and P. K. Sarvepalli, *IEEE Trans. Inf. Theory* **52**, 4892?4914 (2006).
- [51] A. J. Scott, *Phys. Rev. A* **69**, 052330 (2004).