Defining Millisecond Pulsars

Priyam Halder,¹ Satyaki Goswami,² Protyusha Halder,³ Uday Ghosh,⁴ and Sushan Konar⁵

¹SASTRA University, Thanjavur 613401, India
²IIT Madras, Chennai 600036, India
³Lovely Professional University, Jalandhar 144411, India
⁴ISI-North-East Centre, Tezpur 784501, India
⁵NCRA-TIFR, SP Pune University Campus, Pune 411007, India

ABSTRACT

Millisecond pulsars (MSP) are an important subclass of rotation powered pulsars (RPP), traditionally defined as those with $P_s < 20 - 30$ ms and $B_s \leq 10^{10}$ G. We re-examine this definition by applying Gaussian mixture model (GMM) analysis to identify distinct clusters within the RPP population and find that the MSPs appear to be better demarcated by the condition $\mathbf{P}_s \leq 16$ ms.

Approximately 4000 neutron stars have been observed since their serendipitous discovery more than fifty years ago (Hewish et al. 1968). Despite widely divergent observational characteristics and their consequent grouping in a somewhat large number of observational classes, it is possible to classify these neutron stars into three main types depending on their mode of energy generation. Accordingly, they can either be **a**) rotation powered pulsars (RPP), or **b**) accretion powered pulsars or **c**) internal energy powered objects (Konar 2013; Konar et al. 2016; Konar 2017).

However, most of the observed neutron stars belong to the RPP class and are powered by the loss of rotational energy due to magnetic braking. The basic observed quantities of a pulsar are its spin-period (P_s) and the period derivative (\dot{P}) , and the most important derived quantity is the dipolar component of the surface magnetic field given by (Manchester & Taylor 1977) -

$$B_{\rm s} \simeq 3.2 \times 10^{19} \left(P_{\rm s}/s \right)^{\frac{1}{2}} \left(\dot{P}/ss^{-1} \right)^{\frac{1}{2}} \,\,{\rm G}\,.$$
 (1)

An important sub-class of RPPs is the rapidly spinning *Millisecond Pulsars* (MSP). Starting with the discovery of B1937+21 ($P_s = 1.558$ ms) by Backer et al. (1982), initially there were only a handful of pulsars with $P_s < 10$ ms which were called MSPs (Bhattacharya & van den Heuvel 1991). As the number of ultra-fast pulsars increased, MSPs came to be known as those with $P_s < 20 - 30$ ms (Cordes & Chernoff 1997; Lorimer 2008), a group that currently accounts for $\sim 15\%$ of the Galactic RPPs. These MSPs, recycled pulsars with both ultra-short spin-periods and weak magnetic fields $(B_s \leq 10^{10} \text{ G})$, are understood to have different evolutionary histories. The two characteristic properties, namely P_s and B_s , get inter-related through the process of MSP formation either by *recycling* of ordinary radio pulsars or by *accretion-induced collapse* of white dwarfs, during long-lived accretion phases of the low-mass or the intermediate mass X-ray binaries (LMXB, IMXB) (Radhakrishnan & Srinivasan 1982; Bhattacharva & van den Heuvel 1991: Tauris et al. 2000).

The direct support for MSP formation in X-ray binaries came from the discovery of coherent millisecond Xray pulsations in SAX J1808.4 (Chakrabarty & Morgan 1998; Wijnands & van der Klis 1998) and subsequently in many similar objects (Wijnands et al. 2005; Watts 2012; Patruno & Watts 2021). On the other hand, the Galactic centre gamma-ray excess appears to have an easy explanation if the MSPs there have formed due to accretion-induced-collapse (Gautam et al. 2022).

In order to understand these different formation scenarios and their influences on the characteristics properties of neutron stars, it is important to study the population of MSPs, requiring a well-defined criterion for identifi-

Corresponding author: Sushan Konar sushan.konar@gmail.com



Figure 1. Top : 1D GMM decomposition of the P_s data. Bottom : 2D GMM decomposition of the combined P_s, B_s data. The ellipses denote the 2σ and 3σ contours of the cluster components. The pink and the green lines show the P_s maxima of the corresponding contours of the MSP cluster. [Data - ATNF Pulsar catalog]

cation of MSPs. So far the classification of MSPs has depended upon the fact that the P_s data showed a clear gap around $P_s \sim 20 - 30$ ms in the beginning. With a much larger RPP population, this demarcation is not so obvious any more.

But defining MSPs with accuracy is not easy and a rigorous definition must involve both P_s and \dot{P}_s (or, equivalently $P_s \& B_s$) since these parameters get intertwined during MSP formation. An attempt to do so was made by Story et al. (2007), assuming the Galactic disc MSPs to form through direct recycling in LMXBs. Clearly, this has limited applicability. For example, in Globular Clusters, the MSPs form through complex routes of binary evolution, resulting in somewhat different distribution of P_s and B_s compared to those in the disc (Konar 2010; Konar & Chahal 2019).

In this work, we attempt to arrive at a semi-rigorous definition of MSPs, using both the P_s and B_s data. An earlier GMM analysis showed that the P_s data is best fit by three components - a) the MSPs, b) the pulsars recycled in high-mass X-ray binaries and c) the ordinary radio pulsars (Konar 2017), though this work did not attempt to quantitatively demarcate individual Gaussian components. In the current work - a) we redo the 1D GMM as the current data has much larger size and find a qualitatively similar result (Fig.1, top); b) - we perform a 2D GMM analysis on the combined P_s , B_s set. The 2D analysis brings out 6 independent Gaussian clusters as shown in Fig.1 (bottom), similar to the results obtained by a recent work on pulsar classification (Reddy et al. 2022).

Ideally, a 2D analysis should be more reliable as it includes both the characteristic parameters. Unfortunately, the 2D case suffers from a serious drawback due to the unavailability of B_s values for a large number of RPPs. Currently, ATNF (Manchester et al. 2005) has P_s values for 3295 objects, but B_s values for only 2514 of them. This is why, it is essential to combine the results from both 1D and 2D GMM analysis to arrive at a 'good' definition of MSPs.

The P_s values obtained from 1D GMM at 95% (2 σ) and 99.7% (3 σ) population cut-off are 10 ms and 16 ms respectively; whereas those obtained from 2D GMM are 8 ms (2 σ) and 12 ms (3 σ). Clearly, the 2D GMM limits are well within those obtained from 1D GMM. On the other hand, 95% cutoff for the middle component of 1D GMM is 27 ms which allows for a robust upper cutoff for MSPs at 16 ms. Therefore, even keeping in mind that GMM analysis is an unsupervised machine learning algorithm which does not allow for very high accuracy, it would perhaps not be too unreasonable to define MSPs as the *subclass of rotation powered pulsars with* $\mathbf{P}_s \lesssim \mathbf{16} \ ms.$

Excellent hospitality at MANUU, Hyderabad during ADAP'23, organised by Priya & Najam Hasan, allowed us to finally settle this question.

REFERENCES

- Backer, D. C., Kulkarni, S. R., Heiles, C., Davis, M. M., & Goss, W. M. 1982, Nature, 300, 615
- Bhattacharya, D., & van den Heuvel, E. P. J. 1991, Phys. Rep., 203, 1
- Chakrabarty, D., & Morgan, E. H. 1998, Nature, 394, 346
- Cordes, J. M., & Chernoff, D. F. 1997, ApJ, 482, 971
- Gautam, A., Crocker, R. M., Ferrario, L., et al. 2022, Nature Astronomy, 6, 703
- Hewish, A., Bell, S. J., Pilkington, J. D. H., Scott, P. F., & Collins, R. A. 1968, Nature, 217, 709
- Konar, S. 2010, MNRAS, 409, 259
- —. 2013, in Astronomical Society of India Conference Series, ed. S. Das, A. Nandi, & I. Chattopadhyay, Vol. 8, 89–95
- 2017, Journal of Astrophysics and Astronomy, 38, 47
- Konar, S., Bagchi, M., Bandyopadhyay, D., et al. 2016, Journal of Astrophysics and Astronomy, 37, 36
- Konar, S., & Chahal, M. 2019, Research Notes of the American Astronomical Society, 3, 88
- Lorimer, D. R. 2008, Living Reviews in Relativity, 11, 8

- Manchester, R. N., & Taylor, J. H. 1977, Pulsars (San Francisco: W. H. Freeman)
- Manchester, R. N., Hobbs, G. B., Teoh, A., & Hobbs, M. 2005, AJ, 129, 1993
- Patruno, A., & Watts, A. L. 2021, in Astrophysics and Space Science Library, ed. T. M. Belloni, M. Méndez, & C. Zhang, Vol. 461, 143–208
- Radhakrishnan, V., & Srinivasan, G. 1982, Current Science, 51, 1096
- Reddy, C., Tej, T., & Desai, S. 2022, New A, 91, 101673
- Story, S. A., Gonthier, P. L., & Harding, A. K. 2007, ApJ, 671, 713
- Tauris, T. M., van den Heuvel, E. P. J., & Savonije, G. J. 2000, ApJ, 530, L93
- Watts, A. L. 2012, ARA&A, 50, 609
- Wijnands, R., & van der Klis, M. 1998, Nature, 394, 344
- Wijnands, R., Homan, J., Heinke, C. O., Miller, J. M., & Lewin, W. H. G. 2005, ApJ, 619, 492