

Revealing characteristics of dark GRB 150309A: dust extinguished or high- z ?

A. J. Castro-Tirado^{1,2}, Rahul Gupta^{3,4}, S. B. Pandey³, A. Nicuesa Guelbenzu⁵, S. Eikenberry⁶, K. Ackley⁶, A. Gerarts⁷, A. F. Valeev^{8,9}, S. Jeong^{1,10}, I. H. Park¹⁰, S. R. Oates¹¹, B.-B. Zhang^{12,13}, R. Sánchez-Ramírez^{1,14}, A. Martín-Carrillo¹⁵, J. C. Tello¹, M. Jelínek¹⁶, Y.-D. Hu^{1,17}, R. Cunniffe¹, V. V. Sokolov⁸, S. Guziy^{1,18,19}, P. Ferrero¹, M. D. Caballero-García¹, A. K. Ror³, A. Aryan^{3,4}, M. A. Castro Tirado¹, E. Fernández-García¹, M. Gritsevich^{20,21}, I. Olivares¹, I. Pérez-García¹, J. M. Castro Cerón²², and J. Cepa²³

(Affiliations can be found after the references)

Received 2023; accepted

ABSTRACT

Context. Dark Gamma-Ray Bursts (GRBs) constitute a significant fraction of the GRB population. In this paper, we present the multiwavelength analysis (both prompt emission and afterglow) of an intense (3.98×10^{-5} erg cm⁻² using *Fermi*-Gamma-Ray Burst Monitor) two-episodic GRB 150309A observed early on to ~ 114 days post-burst. Despite the strong gamma-ray emission, no optical afterglow was detected for this burst. However, we discovered near-infrared (NIR) afterglow (K_S -band), ~ 5.2 hours post burst, with the CIRCE instrument mounted at the 10.4m Gran Telescopio Canarias (hereafter GTC).

Aims. We aim to examine characteristics of GRB 150309A as a dark burst and to constrain other properties using multiwavelength observations.

Methods. We used *Fermi* observations of GRB 150309A to understand the prompt emission mechanisms and jet composition. We performed the early optical observations using the BOOTES robotic telescope and late-time afterglow observations using the GTC. A potential faint host galaxy is also detected at optical wavelength using the GTC. We modelled the potential host galaxy of GRB 150309A in order to explore the environment of the burst.

Results. The time-resolved spectral analysis of *Fermi* data indicates a hybrid jet composition consisting of a matter-dominated fireball and magnetic-dominated Poynting flux. GTC observations of the afterglow revealed that the counterpart of GRB 150309A was very red, with $H-K_S > 2.1$ mag (95 % confidence). The red counterpart was not discovered in any bluer filters of *Swift* UVOT/BOOTES, indicative of high redshift origin. This possibility was discarded based on multiple arguments, such as spectral analysis of X-ray afterglow constrain $z < 4.15$ and a moderate redshift value obtained using spectral energy distribution (SED) modelling of the potential galaxy. The broadband (X-ray to NIR bands) afterglow SED implies a very dusty host galaxy with deeply embedded GRB (suggesting $A_V \gtrsim 35$ mag).

Conclusions. The environment of GRB 150309A demands a high extinction towards the line of sight, demanding dust obscuration is the most probable origin of optical darkness and the very red afterglow of GRB 150309A. This result makes GRB 150309A the highest extinguished GRB known to date.

Key words. Gamma-ray burst: general – Gamma-ray burst: individual: GRB 150309A– techniques: photometric–dust, extinction

arXiv:2310.08503v1 [astro-ph.HE] 12 Oct 2023

1. Introduction

Gamma-ray bursts (GRBs) are the most powerful and sudden explosions of gamma-ray radiation, occurring at cosmological distances (Piran 2004; Kumar & Zhang 2015). GRBs can generally be categorised into two families: long-duration (lasting more than two seconds) and short-duration (lasting less than two seconds) burst (Kouveliotou et al. 1993). Long-duration GRBs are believed to originate from the collapse of massive stars (collapsar model). The core collapse of a massive star leads to the formation of a black hole or a neutron star and the succeeding release of a powerful and ultra-relativistic GRB jet (MacFadyen & Woosley 1999; Woosley & Bloom 2006; Heger et al. 2003; Obergaulinger & Aloy 2020). Additionally, the observational evidence of their progenitors comes from late-time multi-band observations of their afterglows and host galaxies. Some nearby long GRBs have been observed with the photometric and spectroscopic evidence of supernova signatures (Stanek et al. 2003; Cano et al. 2017), further supporting the collapsar model and a dense environment. Long-duration GRBs are generally found in star-forming regions of their host galaxies (Bloom et al. 1998). On the other hand, Short-duration GRBs are believed to originate from the merger of compact objects and are generally found in older and less active host galaxies (Perna & Belczynski 2002; Belczynski et al. 2002; Abbott et al. 2017; Goldstein et al. 2017). However, in recent times, some of the hybrid events (long-duration GRB from merger and short-duration GRB from collapsar origin) were observed, which further challenges our classical understanding (Della Valle et al. 2006; Ahumada et al. 2021; Zhang et al. 2021; Troja et al. 2022; Rastinejad et al. 2022; Yang et al. 2022; Rossi et al. 2022).

There is a subclass of GRBs afterglows, known as ‘dark bursts’ for which no optical counterpart is detected in the face of deep follow-up (fainter than ~ 21 - 22 mag) in < 24 hours after the trigger, the number of similar sources has been increasing in the present era of GRB research (Jakobsson et al. 2004; Greiner et al. 2011). These days, dark bursts appear to make up a significant part of the overall GRB population. From 1997 to 2021, about 66 % (1526/2311) of well-localised bursts are observed with an X-ray afterglow; however, only 38 % (874/2311) bursts are observed with an optical afterglow¹. In some cases, optical afterglows could not be detected due to a lack of early observations (Pandey et al. 2003). But in other cases, deep and early observations reveal that true dark GRBs indeed exist and have a typical fraction of around 25-40 % of all the bursts (Perley et al. 2016). In the case of GRB 150309A, an X-ray counterpart of the burst was discovered by *Swift* X-ray telescope (XRT; Burrows et al. 2005) but no optical afterglow was discovered in spite of deep search using the Burst Observer and Optical Transient Exploring System (BOOTES) and Gran Telescopio Canarias (GTC) telescopes. However, observations in the near-infrared (NIR) bands using the Canarias InfraRed Camera Experiment (CIRCE) instrument mounted on the GTC telescope reveal the detection of a very red afterglow. These observations suggest that GRB 150309A belongs to the ‘dark bursts’ subclass.

Several methods exist in the literature to classify these dark GRBs. At the early stage of their studies, GRBs without an optical afterglow were classified as dark GRBs (Groot et al. 1998). In later studies (De Pasquale et al. 2003; Jakobsson et al. 2004), the analogy between the X-ray and optical/near-IR afterglow properties were used for their classification. De Pasquale et al. (2003) used the flux ratio of optical and X-ray afterglows, and Jakobsson et al. (2004) used the spectral index ($\beta_{\text{OX}} < 0.5$) obtained

using optical to X-ray spectral energy distribution (SED) to classify these bursts. van der Horst et al. (2009) suggested a more enlightened method and considered dark GRBs by β_{OX} (joint X-ray and optical spectral index) $< \beta_{\text{X}}$ (X-ray spectral index) -0.5 in the framework of the fireball model.

There may be many potential causes for the optical darkness of GRB afterglows. 1) Dark GRB afterglows could be intrinsically faint (Fynbo et al. 2001), for example, if the relativistic fireball is decelerated in a low-density surrounding medium (Sari et al. 1998). 2) The high redshift origin (Taylor et al. 1998) of GRB afterglows, in such case the Ly- α forest will influence the optical wavelength (Lamb & Reichart 2000). The cause of optical darkness due to high redshift origin is low and expected for ~ 10 - 20 % of dark bursts (Perley et al. 2016). 3) Obscuration scenario: This could be because of i) a high column density of gas or ii) dust in their host galaxies at larger distances so that the optical afterglow is very reddened (Fynbo et al. 2001). The last scenario is particularly expected to be the most favoured cause for the optical darkness of GRB afterglows, as most of the host galaxies of dark bursts are detected at the optical wavelength.

The outline of the present work is as follows: In § 2, we present the multiwavelength observations and data reduction of GRB 150309A. In § 3, we have given the results of this work. Finally, a discussion and a brief summary and conclusion of the work are given in § 4 and § 5, respectively. The errors given in the paper are mentioned at 1σ level unless stated otherwise.

2. Observations and Data Reduction

2.1. High energy gamma-ray observations

On 9th March 2015 at 22:59:50.66 UT (hereafter T_0), Gamma-Ray Burst Monitor (GBM, Meegan et al. 2009) instrument of *Fermi* Gamma-ray space mission triggered a bright GRB 150309A with a T_{90} duration of about 52 s (Roberts & Stanbro 2015). The GBM light curve consists of two distinct emission episodes, the main episode followed by a soft, weak emission episode with a temporal gap of ~ 200 s in between (see Fig. 2). After ~ 200 s (at 23:03:06 UT, $T_{\text{sw},0}$), the Burst Alert Telescope (BAT; Barthelmy et al. 2005) instrument of *Swift* mission also triggered GRB 150309A during the second weak emission episode (Cummings et al. 2015). The mask-weighted light curve obtained using BAT observations also shows that GRB 150309A consists of two separate episodes with a T_{90} duration of 242 ± 6 s (15-350 keV). The prompt emission of GRB 150309A also was independently discovered by *Konus-Wind* mission (Golenetskii et al. 2015).

We retrieved photon events received by three sodium iodide (NaI) and one bismuth germanium oxide (BGO) detector (having the largest number of photon counts) of the GBM instrument onboard *Fermi*. These NaI and BGO detectors have the smallest boresight angle with respect to the GRB location. We carried out the temporal and spectral analysis of high energy γ -ray data as described in Caballero-García et al. (2023).

2.2. X-ray and Ultra-Violet observations

The XRT instrument of *Swift* mission started follow-up observations of the BAT localisation region at 23:05:18.2 UT (131.5 s after the BAT trigger) to search for X-ray afterglow. An uncatalogued counterpart (X-ray) candidate was discovered at RA, DEC = 18h 28m 24.81, +86d 25' 43.6" (J2000) within the *Swift*

¹ <https://www.mpe.mpg.de/~jcg/grbgen.html>

BAT error circle with a 90% uncertainty radius of 1.4 arcsec² in the initial window timing (WT) mode exposure (Cummings et al. 2015). The position of this fading afterglow was observed up to $\sim 4 \times 10^5$ s post-BAT detection. This work utilised X-ray afterglow data products, including both light curve and spectrum, obtained from the *Swift* online repository³ hosted and managed by the University of Leicester (Evans et al. 2007, 2009). We conducted the analysis of the X-ray afterglow spectra acquired from the *Swift* XRT using the X-Ray Spectral Fitting Package (XSPEC) package (Arnaud 1996). The XRT spectra were analysed within the energy range of 0.3–10 keV. For the XRT spectral analysis, we used an absorbed power-law model to explain the spectral properties of the X-ray afterglow of GRB 150309A. We considered both the photoelectric absorption from the Milky Way galaxy (phabs model in XSPEC) and the host galaxy of the GRB (zphabs model in XSPEC) along with the power-law model for the afterglow. The absorption due to our Galaxy was set as a fixed spectral parameter with a hydrogen column density of $N_{\text{HGal}} = 9.05 \times 10^{20} \text{ cm}^{-2}$ (Willingale et al. 2013). However, the intrinsic absorption ($N_{\text{H}z}$) at the redshift value equal to two (the mean redshift z value for a typical long-duration GRB) was permitted to vary freely. In addition, the photon index value of the power-law component was left to vary. We have used C-Stat statistics for the spectral fitting of XRT data.

The Ultra-Violet and Optical Telescope (UVOT; Roming et al. 2005) onboard *Swift* began observing the XRT localisation region 140 s after the BAT trigger to search for UV/Optical afterglow. However, no credible Ultra-Violet (UV) or optical counterpart candidate was discovered within the *Swift* XRT error circle (Cummings et al. 2015; Oates & Cummings 2015). We extracted the UV afterglow data obtained using *Swift* mission following the method described in Gupta et al. (2021).

2.3. Optical and near-IR afterglow observations

For GRB 150309A, soon after the detection of X-ray counterpart by *Swift* XRT, many ground-based optical observatories (including BOOTES robotic telescope) started follow-up observations to search for the optical counterpart of GRB 150309A, although no optical afterglow (from early to late phases) candidate consistent with the BAT position was detected. Our later optical non-detections are consistent with the optical limits given by (Rumyantsev et al. 2015) and (Mazaeva et al. 2015).

We triggered the Target of opportunity (ToO) follow-up observations in the near-IR (JHK) starting 5.0 hr after the prompt discovery with the GTC equipped with the CIRCE instrument at the Spanish Observatory of La Palma. We detected a potential NIR counterpart candidate using GTC, although we could not find any afterglow emission at optical wavelengths. Additional near-IR observations were made at GTC on July 3 2015 (i.e. 114 days post-burst). Table 1 displays the optical and NIR observing log of GRB 150309A; the magnitude values listed in the table are given in the AB system. The K_s -filter image of the GRB 150309A field taken with the GTC (+CIRCE) in 2015 is shown in Fig. 1.

To determine the magnitudes from the optical and NIR frames, we utilised the DAOPHOT routine under Image Reduction and Analysis Facility (IRAF)⁴.




Fig. 1. The field of GRB 150309A. Top panel: The K_s -band discovery image (in green circle) of the afterglow of GRB 150309A taken at the GTC (+CIRCE) in 2015. The near-IR afterglow is close to the centre of the image. Bottom panel: The late-time g -filter image of the field observed using the GTC in July 2016. The potential host galaxy is close to the centre of the image (within the XRT error circle of radius 1.4 arcsec, shown with a blue circle). North is shown in the upward direction, and East is shown towards the left direction.

2.4. Search for potential Host Galaxy of GRB 150309A

We performed deep optical imaging in *griz* filters using the GTC telescope (see Table 1). The deeper late-time optical photometric observations were gathered on July 19, August 24, August 25 (2015), July 07, July 30 (2016), and March 11 (2021). We carried out image pre-processing, such as dark subtraction and flat-fielding, using IRAF routines. Further, we performed the photometry of cleaned GTC images using standard IRAF software. To calibrate the instrumental magnitude, we completed the field calibration using the Sloan Digital Sky Survey (SDSS) DR12 catalogue Alam et al. (2015). At the position RA= 18:28:24.67, and DEC= +86:25:44.16 (J2000), we detected a faint galaxy with the following magnitude values $g = 25.56 \pm 0.27$, $r = 25.26 \pm 0.27$, $i = 24.80 \pm 0.22$, and $z \geq 24.4$. The position of the candidate host galaxy is consistent with the XRT position (as shown in Fig. 1). We also performed spectroscopy observations (covering the 3700 – 7750 Å wave range) with an exposure time of 3×1200 s on July 07 2016, using GTC. We used standard procedures for analysing the Optical System for Imaging and low-Intermediate-Resolution Integrated Spectroscopy (OSIRIS) spectra. Table 1 shows the optical and NIR photometry of the potential host. The reported magnitude values are not corrected for Galactic reddening. The calculated Galactic reddening value is $E(B-V) = 0.1206$ (Schlafly & Finkbeiner 2011). The potential host galaxy of GRB 150309A is shown in Fig. 1. A detailed method of host galaxy data analysis of GTC data is described in Gupta et al. (2021).

ties for Research in Astronomy, Inc., under cooperative agreement with the US National Science Foundation. NSF.

² https://www.swift.ac.uk/xrt_positions/00634200/

³ <https://www.swift.ac.uk/>

⁴ IRAF is distributed by the NOAO, National Optical Astronomy Observatories, which are operated by USRA, the Association of Universi-

Table 1. The log of optical and near-infrared images obtained on the GRB 150309A field (top panel for the afterglow and bottom panel for the potential host galaxy). The magnitude values in different filters are uncorrected for the reddening due to our galaxy. The non-detection upper limits values are listed at 3σ . The magnitudes listed are given in the AB system. * marker denotes the magnitudes in the Vega system.

Date (UT, mid)	T-T ₀ (s)	Telescope/ Instrument	Filter/ Grism	Exposure Time (s)	Magnitude/ Upper limit
09 Mar 2015, 23:05:53	362	0.6m BOOTES-2/TELMA	--	60	≥18.0
09 Mar 2015, 23:05:53	362	0.3m BOOTES-1	--	60	≥16.5
09 Mar 2015, 23:11:08	677	0.6m BOOTES-2/TELMA	--	120	≥20.2
10 Mar 2015, 00:03:21	3810	0.6m BOOTES-2/TELMA	<i>i</i>	5400	≥21.0
10 Mar 2015, 01:22:34	8563	0.6m BOOTES-2/TELMA	<i>z</i>	3600	≥19.2
10 Mar 2015, 03:21:26	15695	0.6m BOOTES-2/TELMA	<i>Y</i>	9300	≥18.0
10 Mar 2015, 04:15	18909	GTC (CIRCE)	<i>K_S</i>	300	19.28 ± 0.11*
10 Mar 2015, 04:25	19509	GTC (CIRCE)	<i>H</i>	300	≥ 21.4*
10 Mar 2015, 04:35	20109	GTC (CIRCE)	<i>J</i>	300	≥21.3*
10 Mar 2015, 05:09	22149	GTC (CIRCE)	<i>K_S</i>	300	19.50 ± 0.20*
03 Jul 2015, 00:30	115.0625 (days)	GTC (CIRCE)	<i>K_S</i>	1800	≥21.5*
03 Jul 2015, 00:40	115.0696 (days)	GTC (CIRCE)	<i>H</i>	1800	≥22.0*
Date (UT, mid)	T-T ₀ (days)	Telescope/ Instrument	Filter/ Grism	Exposure Time (s)	Magnitude/ Upper limit
19 Jul 2015, 01:31:50	131.1054	GTC (OSIRIS)	<i>r</i>	1800 (15x120 s)	25.26 ± 0.27
24 Aug 2015, 00:34:18	167.0654	GTC (OSIRIS)	<i>i</i>	2160 (24x90 s)	24.89 ± 0.16
25 Aug 2015, 23:33:37	169.0233	GTC (OSIRIS)	<i>i</i>	2160 (24x90 s)	25.09 ± 0.39
07 Jul 2016, 01:32:06	485.1058	GTC (OSIRIS)	R1000B	3600 (3x1200 s)	–
30 Jul 2016, 00:26:21	508.0600	GTC (OSIRIS)	<i>g</i>	1200 (8x150 s)	25.56 ± 0.27
30 Jul 2016, 01:20:48	508.0979	GTC (OSIRIS)	<i>i</i>	900 (10x90 s)	24.80 ± 0.22
11 Mar 2021, 05:54:10	2193.2879	GTC (OSIRIS)	<i>z</i>	1890 (42x45 s)	≥24.4

3. Results

3.1. Prompt characterise and classification

In the following subsections, we present the results of the prompt analysis of GRB 150309A using high-energy observations.

3.1.1. Light curve and spectral analysis:

The energy-resolved γ -ray light curve for GRB 150309A, along with the evolution of hardness ratio (HR), is presented in Fig. 2. The light curves show two episodic emissions with a temporal gap in between. The first main episode is followed by a soft, weak episode with a very large (~ 200 s) temporal gap in between. Generally, two-episodic GRBs' prompt emission light curves consist of three different sub-classes. They follow a soft, weak emission episode preceding the main episode, the main emission component followed by a soft, weak emission component, or both the emission episodes have comparable intensity. The second sub-class (the main emission component followed by a soft, weak emission component) is very rare Lan et al. (2018) and GRB 150309A belongs to this rare class. The evolution of HR shows that for GRB 150309A, the first episode is harder than the subsequent episode, which is also seen from the very low signal for the second episode in the BGO 0.3-1 MeV light curve.

We performed the time-integrated and time-resolved spectral analysis using data from GBM detectors with the strongest signal: NaI 0, NaI 1, and NaI 2 (energy range 8-900 keV). We also selected the BGO detector (energy range 250-40000 keV) closest to the GRB direction (BGO 0). We generated the custom response matrices and spectra files using the publicly

available `gtburst`⁵ software. We have used the Multi-Mission Maximum Likelihood framework (Vianello et al. 2015, 3ML⁶) software. A detailed spectral analysis method is described in Caballero-García et al. (2023). The time-integrated spectra of the first episode ($T_0-1.35$ to $T_0 + 60.00$ s) and second episode ($T_0 + 215.72$ to $T_0 + 272.95$ s) are best fitted by the Band function with following model parameters: E_p (peak energy) = $169.53^{+5.08}_{-5.11}$, $33.07^{+5.30}_{-5.47}$ keV, α_{pt} (low energy spectral index) = $-0.60^{+0.03}_{-0.03}$, $-0.59^{+0.38}_{-0.38}$ and β_{pt} (high energy spectral index) = $-2.86^{+0.20}_{-0.20}$, $-2.66^{+0.29}_{-0.31}$, respectively. The calculated values of E_p also support the harder nature of the first episode.

Time-resolved spectroscopy of prompt emission of GRBs is a propitious method to explore the emission mechanisms (thermal or non-thermal) and to probe the correlations between spectral parameters of GRBs, which are yet an open question of GRB prompt emission physics. We divided the total emission duration of the first and second episodes of GRB 150309A based on the Bayesian Block method, selected the energy channel from 8 - 900 keV in the sodium iodide detector having a maximum count rate (NaI 1) as the Bayesian Block algorithm is believed to be the most acceptable method to recognise the intrinsic intensity variation in the prompt emission light curve of GRBs (Scargle et al. 2013; Burgess 2014). We further consider only those Bayesian bins with statistical significance (S) ≥ 30 to ensure an excellent signal-to-noise ratio for the spectral analysis. These selection criteria provided the nine spectra for the first episodes; however, no bins for the second episode fulfilled our binning criteria. We used two models (Band, and Cutoff-power law) for time-resolved analysis. We notice that most of the time-resolved spectra (5/9)

⁵ <https://fermi.gsfc.nasa.gov/ssc/data/analysis/scitools/gtburst.html>

⁶ <https://three.ml.readthedocs.io/en/latest/>




Fig. 2. Energy-resolved *Fermi* GBM and *Swift* BAT gamma-ray light curves of GRB 150309A: The multi-channel background-subtracted prompt emission light curve (top five panels) consists of two emission episodes with a significant temporal gap. The first and second episodes are shown with solid black and blue vertical lines. The peak times during both episodes are shown using black and blue vertical dashed-dotted lines, respectively. The insets in panels four and five show the zoomed version of the second episode based on GBM and BAT observations. The last panel shows the evolution of the hardness ratio during both episodes. The green horizontal solid line indicates the unit value of the hardness ratio.

for the first episode are best fitted with the Band function, and the remaining spectrum for the first episode is described with Cutoff-power law model. The best-fit model parameters and associated errors are given in Table A. The evolution of spectral parameters along with the prompt emission light curve is shown in Fig. 3. As it can be seen from Fig. 3 that E_p is changing throughout the emission episodes, resulting in spectral evolution. The E_p evolution seems to follow an intensity-tracking trend for the first episode. On the other hand, the evolution of α_{pt} has a decreasing pattern with time. This signifies that the spectrum changes from harder to softer (a hard to soft trend), and the spectrum has a different origin than the synchrotron at the beginning (it exceeds the line of death for synchrotron slow cooling), but later synchrotron emission may dominate.

We look for the correlations between spectral parameters obtained from time-resolved spectroscopy as they play an essential role in finding the emission mechanism of the prompt emission of GRBs. We look for the correlation between time-resolved E_p -flux, α_{pt} -flux, and E_p - α_{pt} spectral parameters obtained using the Band function based on GBM observations for each temporal bin of the first episode of GRB 150309A. We found a strong correlation between the E_p and the flux in 8 keV to 30 MeV energy channels with a Pearson coefficient (r) and p -value of 0.94 and 1.9×10^{-4} . We found no correlation between α of Band function and flux. Furthermore, we investigated the correlation between E_p and α_{pt} . In this case, also, we did not find any correlation. Therefore, the first episodes of GRB 150309A have the characteristics of an ‘intensity-tracking’ GRB (the peak




Fig. 3. Evolution of spectral parameters for GRB 150309A. (Top) The peak energy evolves with time and follows an intensity-tracking trend during the first episode. (Middle) The evolution of low energy spectral index α_{pt} using *Fermi* GBM data. α_{pt} seems to follow a hard to a soft trend during the first episode. The two horizontal lines are lines of death for synchrotron fast cooling ($\alpha_{pt} = -3/2$, dotted dashed line) and synchrotron slow cooling ($\alpha_{pt} = -2/3$, solid red line). (Bottom) The evolution high energy spectral index β_{pt} using *Fermi* GBM data.

energy follows the intensity-tracking pattern) similar to many multi-episodic GRBs (Li et al. 2020).

3.1.2. Classification of GRB 150309A:

We calculated the prompt properties of GRB 150309A (see Table 2) and compared them with a large population of GRBs to know the intrinsic class of GRB 150309A. We mainly use four classifiers: spectral hardness-duration distribution, minimum variability timescale-duration distribution, spectral lag-luminosity correlation, and Amati correlation.

To find the time-integrated HR of individual episodes, we divided the counts in soft (10 - 50 keV) and hard (50-300 keV) energy ranges using the three most brightest sodium iodide detectors of GRB 150309A. We also calculated the T_{90} duration of both the episodes and compared it with other two-episodic *Fermi* GRBs (with known redshift) studied by Chand et al. (2020). The spectral hardness-duration distribution diagram (see Fig. 4 (top left)) shows that both the episodes of GRB 150309A have properties of long GRBs. The probabilities of a burst classified as a long burst are also shown in the background (obtained from Goldstein et al. 2017).

The minimum variability time scales (t_{mvs}) for long GRBs are typically longer than short GRBs. We calculated t_{mvs} for GRB 150309A utilising the Bayesian block method on the GBM light curve in 8-900 keV. A more detailed method to determine t_{mvs} is described in Vianello et al. (2018). The minimum variability timescale-duration distribution diagram for GRB 150309A (red square) along with other GRBs sample Golkhou et al. (2015) is shown in Fig. 4 (top right). The diagram shows that GRB 150309A belongs to long GRBs. The probabil-

ities of a burst classified as a long burst in the sample studied by Golkhou et al. (2015) are also shown.

Further, we calculated the spectral lag for GRB 150309A following the method discussed in Caballero-García et al. (2023). We calculated the lag using prompt emission light curves ($T_0 - 1.35$ to $T_0 + 60.00$ s) in two energy ranges (15–25 keV and 50–100 keV). We obtained a negative lag for GRB 150309A (possibly due to the superposition effect). The observed negative lag indicates that the low-energy photons were observed before the high-energy photons. We also calculated the isotropic peak luminosity of GRB 150309A and plotted the spectral lag-luminosity correlation diagram (Fig. 4 (bottom left)) for GRB 150309A along with other samples studied by Ukwatta et al. (2010). We found that GRB 150309A does not follow the spectral lag-luminosity anti-correlation.

According to the ‘Amati correlation’ (Amati 2006), the time-integrated peak energies corrected to the rest frame ($E_{p,i}$) are correlated with the prompt emission isotropic equivalent γ -ray energy ($E_{\gamma,iso}$). This correlation is also studied and found valid for an episode-wise sample of GRBs Basak & Rao (2013); Chand et al. (2020). Amati correlation is useful for classifying the GRBs. We studied the correlation for each episode of GRB 150309A. To calculate $E_{\gamma,iso}$ for both episodes, we calculated the time-integrated total energy fluence in $1-10^4$ keV energy channels in source rest frames. We have shown the episode-wise GRB 150309A (red and blue squares) in the Amati correlation plane along with the other data points for long GRBs and short bursts studied in Minaev & Pozanenko (2020) and Chand et al. (2020) (two-episodic GRBs). We noticed that both the episodes of GRB 150309A are consistent with the Amati correlation of long bursts (see bottom right panel of Fig. 4).

Table 2. Prompt emission properties of individual episodes of GRB 150309A. The episode-wise peak fluxes are measured in the 1–10,000 keV energy range in the source frame. t_{mvtS} , HR, E_p , peak flux (F_p), $E_{\gamma,iso}$, and $L_{\gamma,iso}$ denote the minimum variability time scales, hardness ratio, peak energy, peak flux, isotropic equivalent γ -ray energy and isotropic equivalent γ -ray luminosity, respectively.

Prompt Emission Properties	Episode 1	Episode 2
T_{90} (s)	37.43	11.27
t_{mvtS} (s)	~ 0.7	
HR	1.14	0.20
E_p (keV)	$169.53^{+5.08}_{-5.11}$	$33.07^{+5.30}_{-5.47}$
F_p (10^{-7} erg cm^2 s^{-1})	7.11	0.27
$E_{\gamma,iso}$ (erg)	43.09×10^{52}	1.68×10^{52}
$L_{\gamma,iso}$ (erg s^{-1})	8.72×10^{52}	-
Redshift z	2.0	
Spectral lag (s)	$-0.56^{+0.49}_{-0.50}$	-

3.2. Afterglow Characteristics

In the following subsections, we present the results of the afterglow analysis of GRB 150309A using X-ray to NIR band observations.

3.2.1. The nature of X-ray afterglow of GRB 150309A

The X-ray afterglow observations using *Swift*/XRT shows a very steep decline $\alpha_{x1} = -2.13^{+0.09}_{-0.09}$ from $T_{sw,0} + 123$ s to $T_{sw,0} + 397$

s followed by a shallower decline rate of $-1.05^{+0.03}_{-0.03}$ after $T_{sw,0} + 397$ s (see Fig. A1 of the appendix). Further, we carried out *Swift* XRT spectral analysis for the temporal bins selected before and after the temporal break following (Gupta et al. 2021). We calculated the spectral parameters for all the Photon Counting (PC) mode observations (interval after the temporal break) as the photon index value of $\Gamma_{XRT} = 1.78^{+0.10}_{-0.10}$, and the intrinsic hydrogen density column $N_{H,z} = 2.43^{+0.62}_{-0.57} \times 10^{22}$ cm^{-2} , considering $z = 2$. Our intrinsic hydrogen density column measurement shows clear evidence of excess absorption over Galactic hydrogen column density. For the interval before the break, we have frozen the $N_{H,z}$ from the PC mode data and calculated $\Gamma_{XRT} = 1.90^{+0.05}_{-0.05}$. Further, we also checked the evolution of photon indices from the *Swift* burst analyser page⁷ and noted that photon indices do not change significantly during the entire emission phase. The unavailability of obvious spectral evolution between the data before and after the break plus the large change in the temporal index rules out the possibility of relating the temporal break at $T_{sw,0} + 397 \pm 48.60$ s to the crossing of the cooling break frequency (ν_c) across the X-ray wavelength.

The early steep decay emission observed before the break time (< 400 s) could be interpreted as the tail (high-latitude emission) of the prompt emission. In this case, if the emission is purely non-thermal and consists of a single power-law, the temporal index can be expressed as $\alpha = 2 + \beta$ (Kumar & Panaitescu 2000). For the present burst, β_1 (X-ray spectral index during step decay phase) = 0.90 ± 0.05 , which results in an expected temporal index of ~ 2.90 , inconsistent with the observed decay (2.13 ± 0.09). Although, α - β relation during the early steep decay phase can be explained considering that the peak energy (E_p) and flux during this phase rely on the viewing angle Rossi et al. (2011). This multi-peaked behaviour observed during the prompt emission can also cause the temporal indices inconsistent with the simple correlation between α - β during the high-latitude emission (Genet & Granot 2009).

At later times (> 400 s), the observed temporal and spectral indices ($\alpha_2 = 1.05 \pm 0.03$, $\beta_2 = 0.78 \pm 0.10$) are dominated by non-thermal emission because of the interaction of ejecta with the ambient medium (see section 3.3). To investigate the nature of the circumburst medium (ISM or wind), we apply the closure relations between the temporal and spectral index during the shallow decay phase.

1.- Case when $\nu_X < \nu_c$:

$$\begin{aligned} - \alpha_{ISM} &= 3\beta_2/2 = 1.17 \pm 0.15 \\ - \alpha_{wind} &= (3\beta_2 + 1)/2 = 1.67 \pm 0.15 \end{aligned}$$

2.- Case when $\nu_X > \nu_c$ (both mediums are indistinguishable):

$$- \alpha = (3\beta_2 - 1)/2 = 0.67 \pm 0.15$$

The case for an ISM environment with $\nu_X < \nu_c$ seems to be the best fit for our data. All these relations assume an electron spectral index, $p > 2$. The electron spectral index obtained through the closure relations is $p = 2.56 \pm 0.20$. Based on closure relations, all the cases with the wind ($\rho \propto k^{-2}$) environment seem to be completely ruled out. Further, using the calculated values of temporal ($\alpha_2 = 1.05 \pm 0.03$) and spectral ($\beta_2 = 0.78 \pm 0.10$) indices during $\nu_X < \nu_c$ spectral regime and equation 1 of Martin-Carrillo et al. (2014), we computed the density profile ($k = 0$ for ISM and $k = 2$ for wind) of the environment of GRB 150309A. We obtained $k \sim -1.3$, suggesting an intermediate density between ISM and wind ambient medium.

⁷ https://www.swift.ac.uk/burst_analyser/00634200/




Fig. 4. Classification of GRB 150309A: The conventional scheme of GRB classification based on the spectral hardness-duration distribution (top left), minimum variability timescale-duration distribution (top right), spectral lag-luminosity correlation (bottom left), and episode-wise Amati correlation (bottom right). The dashed lines show the boundary between long and short families of GRBs. The high-energy properties of GRB 150309A (displayed with red/blue square symbol) are common among long soft bursts.

Additionally, we computed the value of jet kinetic energy ($E_{K,iso}$) for an ISM environment with $v_X < v_c$ spectral regime utilising the equation 11 of Li et al. (2018). We assumed typical micro-physical parameters (jet is due to Wiebel shocks and $\epsilon_e \approx \sqrt{\epsilon_B}$) to those found in the case of other well-studied GRBs, such as ϵ_B (energy fraction in the magnetic field) = 0.01, ϵ_e (energy fraction in electrons) = 0.1, and number density $n_0 = 1$ (Panaitescu & Kumar 2002). We obtained $E_{K,iso} = 5.36 \times 10^{53}$ erg for GRB 150309A.

3.2.2. A potential near-IR afterglow of GRB 150309A and its confirmation

The fact that the XRT discovered the X-ray counterpart of GRB 150309A (Cummings et al. 2015) enabled us to quickly identify a potential near-IR afterglow within the XRT X-ray error box, at coordinates RA (J2000) = 18 28 25.00, Dec(J2000) = +86 25 45.10 ($\pm 0''.4$). The source remained stable (within errors) in brightness during the CIRCE observation window, with $K_S = 19.28 \pm 0.11$ (Vega) and $K_S = 19.50 \pm 0.20$ (Vega) at the beginning and end of the observation. The object was very red (Fig. 1), and only upper limits were derived at the bluest NIR bands: $J \geq 21.3$ (Vega) and $H \geq 21.4$ (Vega). We calibrated the NIR observations utilising the Two Micron All Sky Survey (2MASS) catalogue.

In order to confirm whether the highly reddened object ($H - K_S \geq 2$) is the afterglow or an extremely reddened host galaxy, a

second epoch observation at GTC (+CIRCE) was conducted on July 3. No source was detected with the following upper limits: $H \geq 22.0$ (Vega) and $K_S \geq 21.5$ (Vega), implying that the afterglow faded by more than 2 mag in the K_S -band after 114 days. This fading behaviour confirms that the highly reddened object is the NIR afterglow of GRB 150309A.

3.3. Spectral Energy Distribution of GRB 150309A

SED analysis using multiwavelength data is very useful in constraining the location of synchrotron break frequencies and host extinction. We created the SED at $T_0 + 5.2$ hours at the time of the GTC (+CIRCE) observations (to maximise the near-simultaneous broadband coverage). We have shown the SED for GRB 150309A in Fig. 5. The closure relations for the X-ray afterglow support that the cooling break frequency is beyond the X-ray spectral coverage (> 10 keV) for an ISM-like ($\rho = \text{constant}$) ambient medium and slow-cooling regime at the epoch of the SED, supporting that no break in SED between X-ray and optical/NIR energies. Therefore, we extrapolated the unabsorbed spectral index of X-ray afterglow towards NIR wavelengths to calculate the extinction in the host galaxy. We noted that K_S band Galactic extinction corrected detection and upper limits in J and H filters using the GTC telescope are situated much below the extrapolated X-ray spectral slope (see Fig. 5). Our SED analysis suggests that the host galaxy of the burst is highly extinguished, and GRB 150309A is a dark burst. We de-




Fig. 5. The SED of GRB 150309A afterglow. Top panel: The SED at $T_0 + 5.2$ hours using the simultaneous X-ray data as well as the near-IR upper limits (and the K-band detection) reported in this paper. The solid red line indicates the X-ray spectral slope obtained using the best fit time-averaged PC mode spectrum, and the pink shaded region within the red dashed lines indicates the associated uncertainty with the X-ray spectral slope. The solid black line indicates the joint NIR-X-ray spectral index (β_{K-X}). The left and right sides of the Y-axis, representing the flux density and magnitude values, are in the AB system. Bottom panel: Ratio of X-ray observed data and best-fit model. The horizontal cyan line indicates the ratio equal to one.

terminated the K_S band host extinction ($A_{K_S} = 3.60^{+0.80}_{-0.76}$ mag) utilizing the X-ray to NIR SED. The calculated host extinction in the K_S filter corresponds to $A_V = 34.67^{+7.70}_{-7.32}$ mag, suggesting that GRB 150309A is one of the most extinguished bursts to date. We constrain the host galaxy extinction considering the Milky Way extinction law. We noted that such a high value of optical extinction ($A_V = 34.67^{+7.70}_{-7.32}$ mag) had been previously observed in the case of a few dark bursts, for example, GRB 051022 (Castro-Tirado et al. 2007), GRB 070306 (Jaunsen et al. 2008), GRB 080325 (Hashimoto et al. 2010), GRB 100614A, GRB 100615A (D’Elia & Stratta 2011).

4. Discussion

4.1. GRB 150309A: A dark burst

There are many possible explanations proposed for GRBs to be dark. In the case of GRB 150309A, the optical non-detections up to deeper limits indicate dark behaviour following the definition proposed at the early stages of their discovery (Groot et al. 1998). In addition, we calculated the joint NIR-X-ray spectral index (β_{K-X}) for GRB 150309A (the closure relation support for no spectral break between NIR to X-ray and cooling frequency lies beyond the X-ray energies) using the spectral energy distribution at $T_{sw,0} + 5.2$ hour and noted $\beta_{K-X} = 0.42 \pm 0.02$ (see Fig. 3.3). However, we could not estimate the optical-to-X-ray spectral index (β_{OX}) at 11 hours post-trigger due to the unavailability of the optical observations near this epoch for GRB 150309A. The calculated value of β_{K-X} also supports that GRB 150309A is a dark burst such as GRB 051022 (Castro-Tirado et al. 2007) and follows the definition described in Jakobsson et al. (2004). Further, we plotted the evolution of NIR-X-ray spectral index vs X-ray spectral index for GRB 150309A (see Fig. 6). We have

also shown the β_{OX} - β_X data points for other well-studied samples of *Swift* GRBs Melandri et al. (2012); Littlejohns et al. (2015) for the comparison. Furthermore, van der Horst et al. (2009) suggested the possible range of joint optical-X-ray spectral slope in different possible regimes: $\beta_X - 0.5 \leq \beta_{OX} \leq \beta_X$ following the external forward shock synchrotron model. We found GRB 150309A well satisfy the definition of darkness suggested by van der Horst et al. (2009) in $\nu < \nu_c$ spectral regime.




Fig. 6. The evolution of NIR-X-ray spectral index as a function of X-ray spectral index for GRB 150309A (presented with a red square). We have also shown the β_{OX} - β_X data points for other well-studied samples of *Swift* GRBs Melandri et al. (2012); Littlejohns et al. (2015) for the comparison. The horizontal black dashed line shows the $\beta_{OX} = 0.5$.

4.2. Origin of the optical darkness of GRB 150309A

We detect a source at the edge of the *Swift* XRT error circle (1.4 arcsec) with $K_S = 19.28 \pm 0.11$ mag (Vega) in the first set of K_S -band observations. We do not detect the afterglow either in the H or the J band data. We detected it again at $K_S = 19.50 \pm 0.20$ mag (Vega) in the second set of K_S data (the magnitude difference is ≤ 1 -sigma, so no apparent fading on this timescale of 20 minutes). The position offset of the K_S -band afterglow is 1.1 -arcsec \pm 1.5 -arcsec (including 1.4 -arcsec error circle from *Swift*/XRT, and about 0.25 -arcsec from CIRCE/2MASS astrometry), so this is definitely in the positional uncertainty region. The clear evidence of the fading nature of the NIR source and its position consistent with the X-rays afterglow suggest that it is certainly the counterpart of the burst. Perhaps most interesting is the non-detection in H (bracketed in time by detection in the K_S -band). The 2-sigma upper limit in H band observations is $H \geq 21.4$ mag (Vega). This means that the counterpart is very red, with $H-K \geq 2.1$ mag (95 % confidence). Only two scenarios are possible: i) either a very dusty host galaxy with deeply embedded GRB ($A_V = 34.67^{+7.70}_{-7.32}$ mag) or a very high-redshift Ly-alpha dropout ($z \geq 10$).

4.2.1. Dust extinguished scenario

The observed characteristic of GRB 150309A indicates that it is positioned undoubtedly in the dark burst region of the $\beta_{OX} - \beta_X$ diagram (see Fig. 6). It is clear that the optical afterglow of GRB 150309A could not be detected because of the obstruction

of sight. We carried out SED analysis (at $T_{\text{sw},0} + 5.2$ hours) to determine the host extinction at NIR wavelengths. Our SED analysis suggests K_S band brightness equal to ~ 17.49 mag (the intrinsic afterglow brightness without any extinction obtained from the extrapolation of the unabsorbed spectral index of X-ray afterglow towards K_S band), from which we found a large host extinction at NIR wavelengths, $A_{K_S} = 3.60^{+0.80}_{-0.76}$ mag (see section 3.3).

4.2.2. A high redshift scenario

From our deep afterglow/host observations using GTC, no optical counterpart or confirmed associated host galaxy (see section 4.3) is detected. Therefore, we could not calculate the exact redshift of the burst. To determine the redshift and to examine the high redshift possibility for optical darkness, we utilised prompt emission Amati correlation (the correlation between rest frame E_p and isotropic gamma-ray energy). We have used time-averaged spectral parameters and bolometric fluence in the source frame for different values of redshift ranging from $z=0.1$ to $z=10$. The Amati correlation for GRB 150309A at different redshift values is shown in Fig. 7. For the comparison, we have also shown the other long bursts studied by Nava et al. (2012). This analysis indicates that the isotropic gamma-ray energy of GRB 150309A is such that it would not be excessive even for $z \geq 10$. The very faint/non-detection of the afterglow in NIR/UV/optical filters also supports a high- z origin. In addition, the prompt emission duration of GRB 150309A would be ~ 5 s in the source frame at $z \geq 10$, consistent with a long GRB.




Fig. 7. Amati correlation for GRB 150309A. As there is no redshift measurement is available, we have to vary the redshift from 0.1 to 10. For the comparison, we have also shown the other long bursts studied by Nava et al. (2012). The solid red line shows the best-fit line, and the pink shaded band shows the associated $2\text{-}\sigma$ uncertainty using the study of a sample of long bursts by Nava et al. (2012).

However, the high redshift origin of GRB 150309A is challenged based on the following argument: considering a high redshift origin of GRB 150309A, the soft X-rays photons in the source frame will be shifted out of the XRT energy coverage (0.3-10 keV). Therefore, even a large column density in the source will result in little attenuation. Furthermore, we utilised the X-ray afterglow spectrum to constrain the redshift of GRB 150309A. We fitted the X-ray PC mode spectral data and

derived (observer frame) considering $z = 0$. The measured column density is higher than that of Galactic column density, and this excess column density is useful to estimate the limit on z . We have used the following equation to constrain the limit on z Grupe et al. (2007):

$$\log(1+z) < 1.3 - 0.5 \log_{10}(1 + \Delta N_H), \quad (1)$$

where ΔN_H represents the difference between column density measured considering $z = 0$ ($N_H \sim 23.06 \times 10^{20} \text{cm}^{-2}$) and Galactic ($N_H \sim 9.05 \times 10^{20} \text{cm}^{-2}$) taken from Willingale et al. (2013). The above equation indicates that GRB 150309A has a redshift value $z < 4.15$.

4.3. Nature of the potential host galaxy of GRB 150309A

Deeper observations searching the host of GRB 150309A found a potential host with an angular separation of $\sim 1.1''$ from the NIR afterglow position (see section 2.4). The potential host is a faint ($r \sim 25.26 \pm 0.27$ mag) one with typical $g-i$ colour measured for galaxies at moderate redshift values detected in the optical bands (see Table 1). The observed spectrum of the potential host galaxy taken using GTC was very noisy, and there is no emission-like feature up to $z=1.08$, discarding the low-redshift possibility (the non-detection of UV emission also discards the low redshift possibility). So, we executed the SED modelling of the potential galaxy of GRB 150309A utilising photometric data and Prospector software. Prospector is a python-based stellar population code developed for the host galaxy SED modelling using both photometric and spectroscopic observations (Leja et al. 2017; Johnson et al. 2021). In order to model the observed data, Prospector applies a library of FSPS (Flexible Stellar Population Synthesis) models (Conroy et al. 2009). We used the dynamic nested sampling fitting routine dynesty on the observed photometry of the potential host galaxy and calculated the posterior distributions of host galaxy parameters. To determine the stellar population properties of the potential host galaxy, we used the `parametric_sfh` model. This model enabled us to calculate several key host properties, including the stellar mass formed (M_\star) in solar mass units, host galaxy age (t), stellar metallicity ($\log Z/Z_\odot$), dust attenuation (A_V), and the star formation timescale (τ). In our SED analysis, we used these host galaxy model parameters as free variables. A detailed method of host galaxy SED modelling is described in Gupta et al. (2022). Due to the limited number of data points, we also included the limiting mag values of z , H , and K_S filters for the SED analysis. We assumed redshift as a free parameter to constrain the photometric redshift of the potential host galaxy. Fig. 8 and Fig. A2 (in the appendix) show the SED fitting and corresponding corner plot of the potential host galaxy of GRB 150309A, respectively. We determined the following parameters using SED modelling: stellar mass formed (M_\star) = $10.66^{+0.44}_{-0.54}$, stellar metallicity ($\log Z/Z_\odot$) = $-2.27^{+1.12}_{-0.83}$, age of the galaxy (t) = $5.80^{+4.77}_{-3.66}$, dust attenuation in rest-frame (A_V) = $0.97^{+0.34}_{-0.37}$ mag, and star formation timescale (τ) = $1.68^{+0.82}_{-0.83}$. Considering the redshift 1.88 obtained from SED modelling, we calculated the physical offset ~ 9.5 kpc between the centre of the potential host galaxy and the NIR afterglow position. The measured physical distance is very large compared to those typically observed for long GRBs, suggesting that the galaxy might not be related with GRB 150309A. To further explore the nature of the potential galaxy, we calculated the chance coincidence probability of the candidate galaxy following the method described in Bloom et al. (2002). Using the

measured brightness of the galaxy in the r-band and the observed offset values, we derived a chance alignment (P_{cc}) of about 5%. The derived P_c value is small but still significantly high, indicating the candidate galaxy to be the host. In literature, authors have used a diverse range of P_{cc} values, that is, ~ 1 -10%, to establish the association of faint galaxies with such transients. For example, Bloom et al. (2002); Berger (2010); O'Connor et al. (2022), use $P_{cc} < 10\%$, Fong et al. (2013) use $P_{cc} < 5\%$, and Tunnicliffe et al. (2014) use 1-2%, to identify if a galaxy is related to afterglow or not. In light of the above analysis, it is hard to decipher whether the observed candidate galaxy is the host of GRB 150309A.

5. Conclusions

We investigated detailed prompt emission properties of GRB 150309A detected by *Swift* and *Fermi*. The high energy light curve of GRB 150309A consists of a rarely observed two-episodic emission phase intersected by a quiescent gap, the first brighter main episode prior to the second fainter and softer episode. The episode-wise time-averaged as well as time-resolved spectral analysis of high energy observations of GRB 150309A exhibit intensity-tracking peak energy E_p evolution. The evolution of the low-energy index (α_{pt}) of Band function overshoot the synchrotron limits in early temporal bins of the first episode posing challenges for the thin shell synchrotron origin and demands for photospheric contribution; however, more complex synchrotron emission models (such as synchrotron emission in decaying magnetic field or time-dependent cooling of electrons) may produce harder α_{pt} . Our detailed time-resolved spectral analysis of high energy data helps us to understand the radiation physics and jet composition of GRB 150309A. The evolution of α_{pt} indicates a hybrid jet composition consisting of a matter-dominated fireball and magnetic-dominated Poynting flux. Considering the value of $E_{\gamma,iso}$ for an assumed value of $z=2$, and derived value of $E_{K,iso}$, we constrained the radiative efficiency ($\eta = E_{\gamma,iso}/(E_{\gamma,iso} + E_{K,iso})$) < 0.4 , typically similar to those found in case of other long GRBs.

Detection of the very red afterglow is crucial to explore the early Universe as they are probes to look for very high redshifts or test beds studying dusty environments surrounding GRBs. In this work, we apprise the discovery of a very red counterpart (K_S -band) of GRB 150309A ~ 5.2 hours post burst with the CIRCE instrument mounted at the GTC, but it was not detected in any bluer filters of *Swift* UVOT/BOOTES, suggesting for a very high redshift origin. However, our present analysis discarded this possibility based on a few arguments, including spectral analysis of X-ray afterglow, constrain $z < 4.15$; SED modelling of the potential nearby galaxy constraining moderate redshift values and offset analysis between the centroid of the potential host to the location of the afterglow etc. Furthermore, we also performed the SED modelling of the potential host galaxy. Our results demand a relatively lower value of extinction and redshift along with typical physical parameters with a rather large physical offset from the galaxy's centre. The considerable difference between A_V values obtained from the host galaxy SED modelling and the one estimated from afterglow SED analysis of GRB 150309A is an indicative of local dust surrounding the progenitor or anomalous extinction within the host galaxy as described in Perley et al. (2009, 2013). Based on our analysis of the potential host, it is hard to decipher whether the observed candidate galaxy is the host of GRB 150309A.

Our analysis of the afterglow SED shows that significant reddening is required to describe the observed K_S band (NIR) afterglow of GRB 150309A, assuming the cooling frequency of external synchrotron model is beyond the optical/NIR and X-rays frequencies. Following this method, we calculated the K_S band host reddening (A_{K_S}) = $3.60^{+0.80}_{-0.76}$ mag, equivalent to $A_V = 34.67^{+7.70}_{-7.32}$ mag. Our analysis also indicates that GRB 150309A is one of the most intense dark GRBs detected so far. Our results suggest that the environment of GRB 150309A demands a high extinction towards the line of sight. Hence, dust obscuration is the most probable reason for the optical darkness of GRB 150309A.

The synergy between state-of-the-art NIR cameras such as CIRCE and the largest diameter optical telescope available so far (the GTC), along with the recently launched James Webb Space Telescope (JWST), makes an ideal combination for studying the population of dark GRBs to a great extent and determining if extinction in the host galaxy or very high redshift is the reason for a significant fraction of the afterglows being beyond the reach of optical telescopes in about 20% of events (Greiner et al. 2011; Tanvir et al. 2008; D'Elia & Stratta 2011). Furthermore, the community is developing several larger optical telescopes, for example, the Extremely Large Telescope (ELT) and Thirty Meter Telescope (TMT). Our study provides insights for future observations of similar fainter afterglows of dark GRBs using upcoming larger telescopes (Pandey 2013).

Acknowledgements. We thank the anonymous referee for providing constructive and positive comments. This work is based partly on observations carried out with the 10.4m Gran Telescopio Canarias (GTC) at the Spanish island of La Palma. This work has partially made use of data products from the Two Micron All Sky Survey (2MASS), which is a joint project of the Univ. of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation. This work has made use of data obtained from the High Energy Astrophysics Science Archive Research Center (HEASARC) and the Leicester Database and Archive Service (LEDAS), provided by NASA's Goddard Space Flight Center and the Department of Physics and Astronomy, Leicester University, UK, respectively. RG and SBP acknowledge the financial support of ISRO under AstroSat archival Data utilisation program (DS_2B-13013(2)/1/2021-Sec.2). AJCT acknowledges support from the Spanish Ministry project PID2020-118491GB-I00 and Junta de Andalucía grant P20_010168 and from the Severo Ochoa grant CEX2021-001131-S funded by MCIN/AEI/ 10.13039/501100011033. MCG acknowledges support from the Ramón y Cajal Fellowship RYC2019-026465-I (funded by the MCIN/AEI/10.13039/501100011033 and the European Social Funding). YDH acknowledges support under the additional funding from the RYC2019-026465-I. RS-R acknowledges support under the CSIC-MURALES project with reference 20215AT009. MG acknowledges the Academy of Finland project No. 325806. Research in UrFU is supported by the Program of Development within the Priority-2030.

References

- Abbott, B. P., Abbott, R., Abbott, T. D., et al. 2017, *The Astrophysical Journal*, 848, L13
- Ahumada, T., Singer, L. P., Anand, S., et al. 2021, *Nature Astronomy*, 5, 917
- Alam, S., Albareti, F. D., Allende Prieto, C., et al. 2015, *ApJS*, 219, 12
- Amati, L. 2006, *MNRAS*, 372, 233
- Arnaud, K. A. 1996, in *Astronomical Society of the Pacific Conference Series*, Vol. 101, *Astronomical Data Analysis Software and Systems V*, ed. G. H. Jacoby & J. Barnes, 17
- Barthelmy, S. D., Barbier, L. M., Cummings, J. R., et al. 2005, *Space Science Reviews*, 120, 143
- Basak, R. & Rao, A. R. 2013, *MNRAS*, 436, 3082
- Belczynski, K., Bulik, T., & Rudak, B. 2002, *ApJ*, 571, 394
- Berger, E. 2010, *ApJ*, 722, 1946
- Bloom, J. S., Djorgovski, S. G., Kulkarni, S. R., & Frail, D. A. 1998, *ApJ*, 507, L25
- Bloom, J. S., Kulkarni, S. R., & Djorgovski, S. G. 2002, *AJ*, 123, 1111
- Burgess, J. M. 2014, *MNRAS*, 445, 2589




Fig. 8. Spectral energy distribution modelling of the potential host galaxy (from the g -band to the K_S band) of GRB 150309A using Prospector software. The SED fitting shows no evidence of internal reddening. The bottom plot shows the transmission curves of the corresponding filters.

- Burrows, D. N., Hill, J. E., Nousek, J. A., et al. 2005, *Space Sci. Rev.*, 120, 165
- Caballero-García, M. D., Gupta, R., Pandey, S. B., et al. 2023, *MNRAS*, 519, 3201
- Cano, Z., Wang, S.-Q., Dai, Z.-G., & Wu, X.-F. 2017, *Advances in Astronomy*, 2017, 8929054
- Castro-Tirado, A. J., Bremer, M., McBreen, S., et al. 2007, *A&A*, 475, 101
- Chand, V., Banerjee, A., Gupta, R., et al. 2020, *ApJ*, 898, 42
- Conroy, C., Gunn, J. E., & White, M. 2009, *ApJ*, 699, 486
- Cummings, J. R., Barthelmy, S. D., Page, K. L., Palmer, D. M., & Ukwatta, T. N. 2015, *GRB Coordinates Network*, 17553, 1
- De Pasquale, M., Piro, L., Perna, R., et al. 2003, *ApJ*, 592, 1018
- D’Elia, V. & Stratta, G. 2011, *A&A*, 532, A48
- Della Valle, M., Chincarini, G., Panagia, N., et al. 2006, *Nature*, 444, 1050
- Evans, P. A., Beardmore, A. P., Page, K. L., et al. 2009, *MNRAS*, 397, 1177
- Evans, P. A., Beardmore, A. P., Page, K. L., et al. 2007, *A&A*, 469, 379
- Fong, W., Berger, E., Chornock, R., et al. 2013, *ApJ*, 769, 56
- Fynbo, J. U., Gorosabel, J., Dall, T. H., et al. 2001, *A&A*, 373, 796
- Genet, F. & Granot, J. 2009, *MNRAS*, 399, 1328
- Goldstein, A., Veres, P., Burns, E., et al. 2017, *ApJ*, 848, L14
- Golenetskii, S., Aptekar, R., Frederiks, D., et al. 2015, *GRB Coordinates Network*, 17566, 1
- Golkhou, V. Z., Butler, N. R., & Littlejohns, O. M. 2015, *ApJ*, 811, 93
- Greiner, J., Krühler, T., Kloke, S., et al. 2011, *A&A*, 526, A30
- Groot, P. J., Galama, T. J., van Paradijs, J., et al. 1998, *ApJ*, 493, L27
- Grupe, D., Nousek, J. A., vanden Berk, D. E., et al. 2007, *AJ*, 133, 2216
- Gupta, R., Oates, S. R., Pandey, S. B., et al. 2021, *MNRAS*, 505, 4086
- Gupta, R., Pandey, S. B., Kumar, A., et al. 2022, *Journal of Astrophysics and Astronomy*, 43, 82
- Hashimoto, T., Ohta, K., Aoki, K., et al. 2010, *ApJ*, 719, 378
- Heger, A., Fryer, C. L., Woosley, S. E., Langer, N., & Hartmann, D. H. 2003, *ApJ*, 591, 288
- Jakobsson, P., Hjorth, J., Fynbo, J. P. U., et al. 2004, *ApJ*, 617, L21
- Jaunsen, A. O., Rol, E., Watson, D. J., et al. 2008, *ApJ*, 681, 453
- Johnson, B. D., Leja, J., Conroy, C., & Speagle, J. S. 2021, *ApJS*, 254, 22
- Kouveliotou, C., Meegan, C. A., Fishman, G. J., et al. 1993, *ApJ*, 413, L101
- Kumar, P. & Panaitescu, A. 2000, *ApJ*, 541, L51
- Kumar, P. & Zhang, B. 2015, *Phys. Rep.*, 561, 1
- Lamb, D. Q. & Reichart, D. E. 2000, *ApJ*, 536, 1
- Lan, L., Lü, H.-J., Zhong, S.-Q., et al. 2018, *ApJ*, 862, 155
- Leja, J., Johnson, B. D., Conroy, C., van Dokkum, P. G., & Byler, N. 2017, *ApJ*, 837, 170
- Li, L., Ryde, F., Pe’er, A., Yu, H.-F., & Acuner, Z. 2020, *arXiv e-prints*, arXiv:2012.03038
- Li, L., Wu, X.-F., Lei, W.-H., et al. 2018, *ApJS*, 236, 26
- Littlejohns, O. M., Butler, N. R., Cucchiara, A., et al. 2015, *MNRAS*, 449, 2919
- MacFadyen, A. I. & Woosley, S. E. 1999, *ApJ*, 524, 262
- Martin-Carrillo, A., Hanlon, L., Topinka, M., et al. 2014, *A&A*, 567, A84
- Mazaeva, E., Inasaridze, R., Makandarashvili, S., et al. 2015, *GRB Coordinates Network*, 17570, 1
- Meegan, C., Lichti, G., Bhat, P. N., et al. 2009, *ApJ*, 702, 791
- Melandri, A., Sbarufatti, B., D’Avanzo, P., et al. 2012, *MNRAS*, 421, 1265
- Minaev, P. Y. & Pozanenko, A. S. 2020, *MNRAS*, 492, 1919
- Nava, L., Salvaterra, R., Ghirlanda, G., et al. 2012, *MNRAS*, 421, 1256
- Oates, S. R. & Cummings, J. R. 2015, *GRB Coordinates Network*, 17559, 1
- Obergaulinger, M. & Aloy, M. A. 2020, *MNRAS*, 492, 4613
- O’Connor, B., Troja, E., Dichiara, S., et al. 2022, *MNRAS*, 515, 4890
- Panaitecu, A. & Kumar, P. 2002, *ApJ*, 571, 779
- Pandey, S. B. 2013, *Journal of Astrophysics and Astronomy*, 34, 157
- Pandey, S. B., Anupama, G. C., Sagar, R., et al. 2003, *A&A*, 408, L21
- Perley, D. A., Cenko, S. B., Bloom, J. S., et al. 2009, *AJ*, 138, 1690
- Perley, D. A., Krühler, T., Schulze, S., et al. 2016, *ApJ*, 817, 7
- Perley, D. A., Levan, A. J., Tanvir, N. R., et al. 2013, *ApJ*, 778, 128
- Perna, R. & Belczynski, K. 2002, *ApJ*, 570, 252
- Piran, T. 2004, *Reviews of Modern Physics*, 76, 1143
- Rastinejad, J. C., Gompertz, B. P., Levan, A. J., et al. 2022, *Nature*, 612, 223
- Roberts, O. J. & Stanbro, M. 2015, *GRB Coordinates Network*, 17561, 1
- Roming, P. W. A., Kennedy, T. E., Mason, K. O., et al. 2005, *Space Sci. Rev.*, 120, 95
- Rossi, A., Rothberg, B., Palazzi, E., et al. 2022, *ApJ*, 932, 1
- Rossi, A., Schulze, S., Kloke, S., et al. 2011, *A&A*, 529, A142
- Rumyantsev, V., Antoniuik, K., Mazaeva, E., & Pozanenko, A. 2015, *GRB Coordinates Network*, 17565, 1
- Sari, R., Piran, T., & Narayan, R. 1998, *ApJ*, 497, L17
- Scargle, J. D., Norris, J. P., Jackson, B., & Chiang, J. 2013, *arXiv e-prints*, arXiv:1304.2818
- Schlafly, E. F. & Finkbeiner, D. P. 2011, *ApJ*, 737, 103
- Stanek, K. Z., Matheson, T., Garnavich, P. M., et al. 2003, *ApJ*, 591, L17
- Tanvir, N. R., Levan, A. J., Rol, E., et al. 2008, *MNRAS*, 388, 1743
- Taylor, G. B., Frail, D. A., Kulkarni, S. R., et al. 1998, *ApJ*, 502, L115
- Troja, E., Fryer, C. L., O’Connor, B., et al. 2022, *Nature*, 612, 228
- Tunnicliffe, R. L., Levan, A. J., Tanvir, N. R., et al. 2014, *MNRAS*, 437, 1495
- Ukwatta, T. N., Stamatikos, M., Dhuga, K. S., et al. 2010, *The Astrophysical Journal*, 711, 1073–1086
- van der Horst, A. J., Kouveliotou, C., Gehrels, N., et al. 2009, *ApJ*, 699, 1087
- Vianello, G., Gill, R., Granot, J., et al. 2018, *ApJ*, 864, 163
- Vianello, G., Lauer, R. J., Younk, P., et al. 2015, *arXiv e-prints*, arXiv:1507.08343
- Willingale, R., Starling, R. L. C., Beardmore, A. P., Tanvir, N. R., & O’Brien, P. T. 2013, *MNRAS*, 431, 394
- Woosley, S. E. & Bloom, J. S. 2006, *ARA&A*, 44, 507
- Yang, J., Ai, S., Zhang, B.-B., et al. 2022, *Nature*, 612, 232
- Zhang, B. B., Liu, Z. K., Peng, Z. K., et al. 2021, *Nature Astronomy*, 5, 911

-
- ¹ Instituto de Astrofísica de Andalucía (IAA-CSIC), Glorieta de la Astronomía s/n, E-18008, Granada, Spain
e-mail: ajct@iaa.es
 - ² Departamento de Ingeniería de Sistemas y Automática, Escuela de Ingenierías, Universidad de Málaga, C. Dr. Ortiz Ramos sn, E-29071, Málaga, Spain
 - ³ Aryabhata Research Institute of Observational Sciences (ARIES), Manora Peak, Nainital-263002, India
e-mail: rahulbhu.c157@gmail.com
 - ⁴ Department of Physics, Deen Dayal Upadhyaya Gorakhpur University, Gorakhpur-273009, India
 - ⁵ Thüringer Landessternwarte Tautenburg, Sternwarte 5, 07778 Tautenburg, Germany
 - ⁶ Department of Astronomy, University of Florida, Bryant Space Science Center, Gainesville, FL 32611, USA
 - ⁷ Grantecan, Santa Cruz de la Palma, Tenerife, Spain.
 - ⁸ Special Astrophysical Observatory of Russian Academy of Sciences, Nizhniy Arkhyz 369167, Russia
 - ⁹ Crimean Astrophysical Observatory, Russian Academy of Sciences, Nauchnyi 298409, Russia
 - ¹⁰ Institute for Science and Technology in Space, SungKyunKwan University, Suwon 16419, Korea
 - ¹¹ School of Physics and Astronomy, University of Birmingham, Birmingham B15 2TT, UK
 - ¹² School of Astronomy and Space Science, Nanjing University, Nanjing 210093, China
 - ¹³ Key Laboratory of Modern Astronomy and Astrophysics (Nanjing University), Ministry of Education, Nanjing 210093, China
 - ¹⁴ INAF, Istituto di Astrofisica e Planetologia Spaziali, via Fosso del Cavaliere 100, I-00133 Rome, Italy
 - ¹⁵ School of Physics, O'Brien Centre for Science North, University College Dublin, Belfield, Dublin 4, Ireland
 - ¹⁶ Astronomical Institute of the Czech Academy of Sciences (ASU-CAS), Fričova 298, 251 65 Ondřejov, CZ
 - ¹⁷ Universidad de Granada, Facultad de Ciencias Campus Fuentenueva s/n, E-18071 Granada, Spain
 - ¹⁸ Nikolaev National University, Nikolska 24, Nikolaev 54030, Ukraine
 - ¹⁹ Nikolaev Astronomical Observatory, Nikolaev 54030, Ukraine
 - ²⁰ Department of Physics, University of Helsinki, Gustav Hällströmin katu 2, FI-00014 Helsinki, Finland
 - ²¹ Institute of Physics and Technology, Ural Federal University, Mira street 19, 620002 Ekaterinburg, Russia
 - ²² European Space Astronomy Centre (ESA-ESAC), Camino bajo del Castillo, s/n, Villafranca del Castillo, E-28692 Villanueva de la Cañada (Madrid), Spain
 - ²³ Instituto de Astrofísica de Canarias, Via Láctea s/n, La Laguna (Tenerife), Spain.




Fig. A.1. X-ray flux light curve of GRB 150309A. Top panel shows the XRT flux light curves at 2 keV energy range. The light curve has been fitted with power-law and broken power-law (best fit) models. Bottom panel: Evolution of hardness ratio in the XRT energy channel. The vertical red dashed line indicates the mean value of hardness ratio for GRB 150309A.

Appendix A: Additional material

Table A.1. Time-resolved spectral fitting of the first episode of GRB 150309A using Band and CPL models. The listed flux values are estimated in 8 keV–40 MeV.

T_{start} (s)	T_{stop} (s)	Significance	α_{pt}	β_{pt}	E_{p} (keV)	(Flux $\times 10^{-06}$)	DIC _{Band}	Γ_{CPL}	E_{c} (keV)	(Flux $\times 10^{-06}$)	DIC _{CPL}	Δ DIC
2.119	9.975	40.59	$-0.15^{+0.08}_{-0.08}$	$-3.07^{+0.22}_{-0.23}$	$114.09^{+4.34}_{-4.54}$	0.58	4572.74	$-0.23^{+0.07}_{-0.07}$	$68.02^{+4.52}_{-4.67}$	0.49	4570.91	1.83
9.975	11.875	33.09	$-0.36^{+0.10}_{-0.10}$	$-2.63^{+0.24}_{-0.25}$	$149.37^{+11.85}_{-11.59}$	1.37	2593.98	$-0.48^{+0.08}_{-0.08}$	$112.10^{+11.79}_{-12.12}$	1.02	2593.37	0.61
11.875	16.349	74.58	$-0.38^{+0.05}_{-0.05}$	$-2.44^{+0.12}_{-0.11}$	$172.78^{+7.96}_{-8.02}$	2.8	3898.49	$-0.50^{+0.03}_{-0.03}$	$134.78^{+6.75}_{-6.69}$	1.87	3922.39	-23.9
16.349	20.547	82.42	$-0.54^{+0.04}_{-0.04}$	$-2.63^{+0.17}_{-0.17}$	$215.39^{+8.90}_{-8.98}$	3.12	3731.73	$-0.60^{+0.03}_{-0.03}$	$168.88^{+8.03}_{-8.17}$	2.37	3741.33	-9.6
20.547	21.861	33.15	$-0.51^{+0.08}_{-0.09}$	$-2.63^{+0.25}_{-0.25}$	$176.76^{+14.22}_{-14.07}$	1.95	2035.34	$-0.60^{+0.07}_{-0.07}$	$142.79^{+16.30}_{-16.29}$	1.47	2036.18	-0.84
21.861	23.799	48.52	$-0.61^{+0.06}_{-0.06}$	$-2.31^{+0.16}_{-0.16}$	$198.38^{+16.64}_{-16.56}$	3.15	2656.74	$-0.71^{+0.05}_{-0.05}$	$185.58^{+17.54}_{-17.42}$	1.93	2671.61	-14.87
23.799	25.994	34.11	$-0.77^{+0.08}_{-0.08}$	$-2.75^{+0.34}_{-0.33}$	$187.26^{+18.14}_{-18.09}$	1.3	2701.82	$-0.81^{+0.06}_{-0.06}$	$169.40^{+21.18}_{-21.11}$	1.03	2700.68	1.14
25.994	29.582	29.29	$-0.75^{+0.09}_{-0.09}$	$-2.50^{+0.38}_{-0.38}$	$161.17^{+18.08}_{-17.65}$	0.87	3385.32	$-0.82^{+0.08}_{-0.08}$	$157.01^{+23.98}_{-24.00}$	0.6	3387.92	-2.6
31.088	60.001	18.84	$-0.95^{+0.15}_{-0.15}$	$-2.83^{+0.33}_{-0.33}$	$87.04^{+11.25}_{-11.39}$	0.12	6215.44	$-1.03^{+0.12}_{-0.12}$	$100.06^{+21.49}_{-21.18}$	0.09	6212.99	2.45




Fig. A.2. The posterior distributions for the SED model parameters of the potential host galaxy of GRB 150309A obtained using nested sampling via dynesty using Prospector software.