

Predictions for Electromagnetic Counterparts to Neutron Star Mergers Discovered during LIGO-Virgo-KAGRA Observing Runs 4 and 5

Ved G. Shah^{1,2} ,  Gautham Narayan^{1,4,5} , Haille M. L. Perkins^{1,4,5} , Ryan J. Foley⁶ ,
Deep Chatterjee⁷ , Bryce Cousins^{3,5} , Phillip Macias⁶ 

¹ Department of Astronomy, University of Illinois at Urbana-Champaign, 1002 West Green Street, Urbana, IL 61801, USA

² Department of Computer Science, University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA

³ Department of Physics, University of Illinois Urbana-Champaign, Urbana, IL 61801, USA

⁴ Center for AstroPhysical Surveys, National Center for Supercomputing Applications, Urbana, IL, 61801, USA

⁵ Illinois Center for Advanced Studies of the Universe, University of Illinois Urbana-Champaign, Urbana, IL 61801, USA

⁶ Department of Astronomy and Astrophysics, University of California Santa Cruz, Santa Cruz, CA 95064, USA

⁷ LIGO Laboratory and Kavli Institute for Astrophysics and Space Research, Massachusetts Institute of Technology, 185 Albany Street, Cambridge, Massachusetts 02139, USA

Accepted XXX. Received YYY; in original form ZZZ

ABSTRACT

We present a comprehensive, configurable open-source software framework for estimating the rate of electromagnetic detection of kilonovae (KNe) associated with gravitational wave detections of binary neutron star (BNS) mergers. We simulate the current LIGO-Virgo-KAGRA (LVK) observing run (O4) using current sensitivity and uptime values as well as using predicted sensitivities for the next observing run (O5). We find the number of discoverable kilonovae during LVK O4 to be 1_{-1}^{+4} or 2_{-2}^{+3} , (at 90% confidence) depending on the distribution of NS masses in coalescing binaries, with the number increasing by an order of magnitude during O5 to 19_{-11}^{+24} . Regardless of mass model, we predict at most five detectable KNe (at 95-per cent confidence) in O4. We also produce optical and near-infrared light curves that correspond to the physical properties of each merging system. We have collated important information for allocating observing resources for search and follow-up observations, including distributions of peak magnitudes in several broad bands and timescales for which specific facilities can detect each KN. The framework is easily adaptable, and new simulations can quickly be produced in response to updated information such as refined merger rates and NS mass distributions. Finally, we compare our suite of simulations to the thus-far completed portion of O4 (as of October 14, 2023), finding a median number of discoverable KNe of 0 and a 95-percentile upper limit of 2, consistent with no detections so far in O4.

Key words: gravitational waves; stars: neutron; methods: statistical;

1 INTRODUCTION

Currently, observable gravitational waves are primarily produced by the coalescence of binary compact objects (Abbott et al. 2016; Abbott et al. 2019b, 2021b; The LIGO Scientific Collaboration et al. 2021). Specifically, binary neutron star (BNS) mergers, like GW170817 (Abbott et al. 2017a), are of interest as these events can yield a post-merger, electromagnetic counterpart known as a kilonova (Abbott et al. 2017c). These transient events are fueled by the radioactive decay of heavy nuclei which are synthesized through r-process nucleosynthesis reactions possible given the neutron rich environment. Under certain conditions, black hole-neutron star mergers can produce kilonovae as well; however, it is much less likely (Fragione 2021), so we focus on BNS mergers here.

As the two neutron stars inspiral, they become tidally disrupted, causing neutron-rich material to be ejected from the system. The amount of material ejected depends, among other factors, on the

equation of state (EOS) (Sekiguchi et al. 2015; Lattimer & Prakash 2016) being "stiff" or "soft" (Lattimer & Prakash 2016; Shibata 2016). A neutron star with a stiff EOS exhibits greater pressure, for a given density, and has larger radius causing it to experience greater tidal forces from its companion. In this work we use the SFHo EOS (Steiner et al. 2013a) used in Setzer et al. (2023) for modeling the kilonova population. Several spectral-energy density (SED) models exist that are parameterized by, for example, the mass and velocity of the ejecta, electron fraction, or opacity (e.g., Barnes & Kasen 2013; Kasen et al. 2015; Metzger 2017; Radice et al. 2018b). For this work we use the *bns_m3_3comp* model grid developed in Bulla (2019); Dietrich et al. (2020), henceforth referred to as the Bulla (2019) grid since it has consideration for observing constraints like viewing angles in its parameter space.

SSS2017a or AT2017gfo is the first optically confirmed kilonova from a binary neutron star merger (Coulter et al. 2017; Lipunov et al. 2017; Soares-Santos et al. 2017; Valenti et al. 2017), which was detected in conjunction with the gravitational-wave event GW170817 (Abbott et al. 2017b) and the gamma ray

* E-mail: vedgs2@illinois.edu




Figure 1. Schematic of the pipeline used to generate synthetic observables for BNS mergers and determine which mergers will produce detectable gravitational waves and electromagnetic counterparts.

burst GRB170717A (Abbott et al. 2017d). This was a landmark discovery for the field of multi-messenger astronomy (MMA) since it was the first detection of a cosmic event via gravitational-waves, a kilonova, and gamma rays.

However, GW170817 remains the only such KN discovery to-date. This is in part due to the current limitations in GW event localization, the coordination required to perform proper follow-up, and the expected rarity of such events. Nonetheless, these events promise many scientific opportunities, such as studying the neutron stars and their EOS (Kilpatrick et al. 2017; Margalit & Metzger 2017a; Siebert et al. 2017; Radice et al. 2018a; Coughlin et al. 2019; Dietrich et al. 2020), understanding r-process nucleosynthesis (Chornock et al. 2017; Cowperthwaite et al. 2017a; Drout et al. 2017; Murguía-Berthier et al. 2017; Shappee et al. 2017), and measuring the expansion of the Universe (Abbott et al. 2018, 2019a; Coughlin et al. 2020; Dietrich et al. 2020). But, to capitalize on these scientific promises, observers must be prepared to discover and follow-up future BNS events. Understanding the number of observable kilonovae expected during gravitational-wave observing runs would provide critical input for the follow-up efforts within the MMA community.

To address this need, we present here a new methodology to quantify the rates of observable kilonovae during the LVK’s ongoing and future observing runs, complimenting similar analysis done recently (Colombo et al. 2022; Frostig et al. 2022; Weizmann Kiendrebeogo et al. 2023). We base our calculation on a number of factors to obtain realistic estimates of these rates (summarized in Figure 1). First, we sample from the appropriate distributions of BNS masses, astronomical extinction, merger rates, and distances adopted from the literature. We use these sampled parameters, either directly or as inputs to compute flux parameters, to perform interpolation on radiative

transfer SED models that we then use to determine the likelihood of electromagnetic counterpart detection. We implement Monte Carlo trials to sample from the distributions in our parameter space to get the distributions of discovery and peak magnitudes, the distances of detected events, and the number of counterpart detections expected in the LVK O4 and O5 observing runs. The framework is also expandable and can support new parameter models, telescopes, and PSDs from future observing runs can be added as they become available.

The paper is outlined as follows. In Section 2, we detail our usage of existing SED models to build synthetic photometry. In Section 3, we describe the BNS parameter distributions that we use in our analysis. In Section 4, we explain our process of sampling from these distributions while accounting for instrumental downtime and other observational constraints. We present the resulting kilonova detection rates in Section 5.

2 SED APPROXIMATION

Running comprehensive, independent simulations (Kasen et al. 2017; Bulla 2019) to produce SEDs for each merger over all trials is computationally unfeasible. Thus, we use interpolation methods over existing SED grids to approximate the EM radiation. Bulla (2019) produced a model (PSSIS) for a grid of kilonovae SEDs simulated using three-dimensional Monte Carlo radiative transfer. These models are parameterised by two different components of the ejecta matter, m_{ej}^{total} : the lanthanide-rich dynamical component, m_{ej}^{dyn} which is released during the merger and the typically larger, lanthanide-free wind component, m_{ej}^{wind} released after the merger as a result of unbinding disk matter. Another parameter is the half-opening angle of

Parameter	List of grid values
Φ	15, 30, 45, 60, 75
$\cos \Theta$	0, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1
$m_{\text{ej}}^{\text{wind}}$	0.01, 0.03, 0.05, 0.07, 0.09, 0.11, 0.13 M_{\odot}
$m_{\text{ej}}^{\text{dyn}}$	0.001, 0.005, 0.01, 0.02 M_{\odot}

Table 1. List of Bulla SED grid values for each of the 4 parameters.

the lanthanide-rich component of the dynamical ejecta, Φ , and the model has a dependence on the cosine of the observing angle, $\cos \Theta$.

Dietrich et al. (2020) further improved the model to account for thermal efficiencies and time dependence for the temperature. We choose to use this model in our work. Thus, our SED model is parameterised by $\{m_{\text{ej}}^{\text{dyn}}, m_{\text{ej}}^{\text{wind}}, \Phi, \cos \Theta\}$. It is important to note that our SED model does not have spin parameter. Although high spin values will have an effect on the resultant kilonova (Raaijmakers et al. 2021), the vast majority of Milky Way neutron stars have very low spin (Zhu et al. 2018), suggesting that high-spin systems are uncommon.

2.1 Interpolation method

All SEDs from the above mentioned model, except those with $\Phi = 0$ and $\Phi = 90$, were used to create a 4 dimensional grid. These two Φ values were excluded as they either lack SEDs for different observing angles or are not available for all permutations of the $m_{\text{ej}}^{\text{wind}}$ and $m_{\text{ej}}^{\text{dyn}}$.

Table 1 describes the discrete points at which SEDs are computed using radiative transfer. While computing the SED for parameters not on the model grid, linear interpolation was used via a regular grid interpolator. The motivation here was that since the flux at every wavelength is known at several finely spaced points in our parameter space through robust simulations, it is reasonable to interpolate between two known points. We use linear interpolation since it is very fast to generate new SEDs on the fly which eliminates the need for pre-computing them, however other interpolation methods with different speed trade-offs also exist within packages like NMMA (Pang et al. 2022).

Distributions of the ejecta masses computed during our trials (Figure 2) indicate that a non-negligible fraction of binary neutron star mergers will produce $m_{\text{ej}}^{\text{dyn}}$ that is greater than the maximum value on the grid (0.02 M_{\odot}) or $m_{\text{ej}}^{\text{wind}}$ that is less than the minimum value on the grid (0.01 M_{\odot}), when sampling component BNS masses from realistic distributions. This necessitates some method for estimating SEDs when the m_{ej} parameters fall outside the grid range.

Given the linear relationship between energy radiated and ejecta mass (Section 3.1 Barnes (2020); Equation 4 Li & Paczyński (1998)), we have computed scaling laws for the total energy radiated for each $\cos \Theta$ and Φ pair. If the $m_{\text{ej}}^{\text{total}}$ from our BNS merger exceeds the grid limit, we use these linear laws to scale the closest grid SED.

If our $m_{\text{ej}}^{\text{total}}$ is lower than the minimum $m_{\text{ej}}^{\text{total}}$ value on the grid, we scale down the closest grid SED using a power law fit since it has the additional benefit of predicting zero flux when the $m_{\text{ej}}^{\text{total}} = 0$, according to:

$$\text{SED} = \alpha \text{SED}_{\text{nn}} \quad (1)$$

where α is the scaling factor and SED_{nn} is the nearest neighboring




Figure 2. Kernel density estimate contours (corresponding to 20%, 50%, and 80% of the probability mass) for dynamical and wind ejecta mass from LVK O4 Monte Carlo trials. The dotted lines show the range of grid values for the two components of the ejecta (Bulla 2019). Data points beyond the grid limits demonstrate the need for an extrapolation method. The SSS17a fit parameters ($\log_{10}(m_{\text{ej}}^{\text{wind}}/M_{\odot}) = -1.28^{+0.42}_{-0.35}$, $\log_{10}(m_{\text{ej}}^{\text{dyn}}/M_{\odot}) = -2.27^{+1.01}_{-0.54}$) were first computed by Dietrich et al. (2020). Both the Farrow et al. (2019) and Galaudage et al. (2021) mass models were used for this analysis.

SED:

$$\alpha = \begin{cases} \left(\frac{m_{\text{ej}}^{\text{total}}}{m_{\text{ej-nn}}^{\text{total}}} \right)^n & \text{if } m_{\text{ej}}^{\text{total}} < \text{lowest grid } m_{\text{ej}}^{\text{total}} \\ \left(\frac{m \cdot m_{\text{ej}}^{\text{total}} + c}{m \cdot m_{\text{ej-nn}}^{\text{total}} + c} \right) & \text{otherwise} \end{cases} \quad (2)$$

where m and c denote the slope and intercept for the linear fit respectively and n denotes the exponent for the power law fit. Note that all the best fit scaling parameters (i.e. m , c , and n) were pre-computed for every pair of $(\Phi, \cos \Theta)$. Figure D1 shows the best fits for the linear and power scaling laws along with the relative errors for some pairs of $(\Phi, \cos \Theta)$. Table D2 and table D3 document all the parameters for linear and power law scaling respectively.

We use this $m_{\text{ej}}^{\text{total}}$ -dependent interpolation scheme instead of linear extrapolation beyond the regular grid range since a small negative slope over a large extrapolated grid range eventually result in negative fluxes at many wavelengths. These extrapolation artifacts result in non-physical SEDs.

Since we want to sample the $(\Phi, \cos \Theta)$ parameters from a continuous range rather than the discrete points computed above to avoid quantization, we fit a spline function to our data for all three scaling parameters (namely m , c , and n) using the smooth bivariate spline. Figure 3 shows the spline surfaces fit to the discrete points. We compute our scaling parameters from this surface for all our Monte Carlo trials. The sum of residuals from the surfaces are provided in Table D1.

Using this piecewise extrapolation method ensures that we can always get SEDs that have reliable total flux since we are scaling the SEDs based on the ejecta mass. However, this method fails to take into account any changes in color as a function of the $m_{\text{ej}}^{\text{total}}$ since there was no obvious statistical trend for how the spectrum shifted. Doing this correctly will require updating the original radiative transfer simulations for a larger range of m_{ej} values which is outside the scope of this paper.




Figure 3. **Left:** Spline surfaces for the slopes (m) of the linear scaling laws for KN SEDs. **Middle:** Spline surfaces for the intercept (c) of the linear scaling laws. **Right:** Spline surfaces for the exponents (n) of the power scaling laws. These laws are used to scale SEDs in cases where the ejecta masses exceed the grid limits of the SED model.

2.2 Redshift and Extinction

Since most binary neutron star mergers are expected to be of extragalactic origin, we treat our SEDs for both host and Milky Way extinction. For the host galaxy, considered to be at rest, we use the CCM89Dust effect based on work from [Cardelli et al. \(1989\)](#). The E_{B-V}^{host} is computed for each SED using the A_V sampled from the distribution described in Equation 10 and $R_V = 3.1$ using the following equation

$$E_{B-V}^{\text{host}} = \frac{A_V}{R_V} \quad (3)$$

For the Milky Way galaxy, considered to be the observing frame, we use the F99Dust effect to redden the SED based on work from [Fitzpatrick \(1999\)](#). Here, we use the SFD dust map [Schlegel et al. \(1998\)](#) to find the E_{B-V}^{MW} based on the RA and Dec of each event.

Finally, we redshift the SED based on the luminosity distance of our kilonova. Details about how this distance is sampled are provided in the parameter distribution section (Section 3). All effects are applied to the SED within SNCOSMO ([Barbary et al. 2016](#)).

3 PARAMETER DISTRIBUTION

Our Monte Carlo simulations use the aforementioned pipeline to estimate SEDs and produce the associated light curves. We sample the component neutron star masses in order to determine if the merger can be detected via gravitational waves. The component masses are also used to compute the $m_{\text{ej}}^{\text{wind}}$ and $m_{\text{ej}}^{\text{dyn}}$ which, in conjunction with the sampled Φ and $\cos \Theta$, are used to estimate the SEDs. Finally we sample A_V and the event coordinates to treat the SEDs and generate the synthetic observables. The distributions for all of these inputs are described below. Table 3 summarizes these distributions.

3.1 BNS mass distribution and computing ejecta mass

For our BNS pairs, we consider the standard formation scenario where binaries consist of a first-born recycled neutron star spun up from accretion (with mass M_{recycled}) and a second-born slow neutron star (with mass M_{slow}).

Analyzing the the mass distributions of these two distinct NS populations has been the subject of numerous studies. [Farrow et al. \(2019\)](#) used a two-peak Gaussian for the recycled NS (Table 2) and

a flat distribution with the range $[1.16, 1.42] M_{\odot}$ for the slow NS, with further analysis done by [Golomb & Talbot \(2022\)](#). [Galauage et al. \(2021\)](#) used a two-peak Gaussian to describe both the slow and recycled NS mass distributions (Table 2). The motivation behind the two peak Gaussian model for slow neutron stars is to reconcile the disagreement between the empirical galactic data on BNS pairs from radio sources ([Farrow et al. 2019](#)) and the distribution that would be required to explain gravitational wave events like GW190425 ([Abbott et al. 2020](#)). It is worth noting that both results use the same two peak Gaussian to explain the recycled NS distribution.

Our simulations support three different BNS mass models: the [Farrow et al. \(2019\)](#) and [Galauage et al. \(2021\)](#) models discussed above (Figure 4) and a flat distribution with some astrophysical priors from the KILOROP package ([Setzer et al. 2023](#)). While presenting the results, we only use the [Farrow et al. \(2019\)](#) and [Galauage et al. \(2021\)](#) models since the uniform distribution is not empirically motivated. Ultimately, we find that while the choice of mass distribution changes the distribution of ejecta properties, it makes little difference in the number of discoverable KNe estimated by our simulations (Table 5).

3.1.1 Discussion of EOS

In addition to the components masses, the equation of state employed will also have an impact on the both the M_{TOV} ([Oppenheimer & Volkoff 1939](#)) and the ejected matter. Consequently, the post merger remnant and the resulting kilonova also depend on the assumed EOS. A neutron star with a stiff EOS exhibits greater pressure, for a given density, and has larger radius causing it to experience greater tidal forces from its companion. This results in greater dynamical ejecta. The post-merger remnant influences the opacity of the ejecta ([Kasen et al. 2015](#); [Radice et al. 2018b](#)). A long-lived neutron star remnant will emit neutrinos, causing the electron fraction to increase and thus reduce the production of lanthanides. In the case of GW170817, [Margalit & Metzger \(2017b\)](#) were able to constrain the nature of the merger remnant to be a short-lived hyper-massive neutron star through the gravitational wave and electromagnetic signals.

While there is still no clear EOS that is most favorable, significant work concerned with fitting functions ([Setzer et al. 2023](#)) to ejecta properties is done using the SFHo EOS ([Steiner et al. 2013b](#)). Since we make use of these fitting functions in our work, we opt to use the same EOS. It is important to note that the equation of state used in

Parameter	Recycled	Slow
ζ	0.68	0.5
μ_1	$1.34M_\odot$	$1.29M_\odot$
σ_1	$0.02M_\odot$	$0.09M_\odot$
μ_2	$1.47M_\odot$	$1.8M_\odot$
σ_2	$0.15M_\odot$	$0.15M_\odot$
M_{low}	—	$1.16M_\odot$
M_{high}	—	$1.42M_\odot$

Table 2. BNS population parameters. ζ defines the fraction of binaries in the low mass peak.

this work places constraints on both minimum and maximum masses for neutron stars. For this reason, we cut off the tails of our mass distributions at $1M_\odot$ and $2.05M_\odot$ respectively.

There is some evidence that the SFHo EOS results in low M_{TOV} compared to empirical data (Foley et al. 2020) and thus may not capture the population diversity. However, recomputing the ejecta fits for different EOS models is outside the scope of this paper.

3.1.2 Computing ejecta masses

While the two NS masses are not themselves parameters for the chosen SED model, we use the masses to compute the ejecta parameters. Although we are not introducing any new ejecta fits in this work, we use this section to recapitulate the methodology for computing these ejecta parameters.

Setzer et al. (2023) used the fitting function introduced by Coughlin et al. (2019) and data from 259 Numerical Relativity (NR) simulations (Radice et al. 2018c) to come up with the final fitting function for the dynamical ejecta mass which depends on the masses ($M_{1,2}$) and compactness ($C_{1,2}$) of the merging neutron stars and is given by

$$\log_{10}(m_{\text{ej}}^{\text{dyn fit}}) = \left[a \frac{(1 - 2C_1)M_1}{C_1} + bM_2 \left(\frac{M_1}{M_2} \right)^n + \frac{d}{2} \right] + [1 \leftrightarrow 2], \quad (4)$$

where $a = -0.0719$, $b = 0.2116$, $d = -2.42$, and $n = -2.905$, and $[1 \leftrightarrow 2]$ refers to repetition of the preceding fit with the indices interchanged. The compactness of the neutron star is given by

$$C = \frac{GM}{c^2 R}. \quad (5)$$

Due to the logarithmic nature of our fit, two heavy neutron stars ($\sim 2M_\odot$ each) result in dynamical ejecta mass that can exceed $1M_\odot$. Metzger (2019) (Section 3.1.1) and work referenced within find that the total dynamical ejecta from BNS mergers lie in the range $10^{-4}M_\odot - 10^{-2}M_\odot$. Thus, we limit the maximum dynamical ejecta from our mergers to $0.09M_\odot$ which means our final $m_{\text{ej}}^{\text{dyn}}$ is

$$m_{\text{ej}}^{\text{dyn}} = \min(0.09, m_{\text{ej}}^{\text{dyn fit}}). \quad (6)$$

The other ejecta parameter, $m_{\text{ej}}^{\text{wind}}$, is some fraction of the disk mass, m_{disk} , where the two are related by

$$m_{\text{ej}}^{\text{wind}} = \zeta m_{\text{disk}}, \quad (7)$$

and ζ is the unbinding efficiency which is sampled uniformly from the range 10–40% and m_{disk} is computed as follows

$$\log_{10}(m_{\text{disk}}) = \max\left(-3, a \left(1 + b \tanh\left[\frac{c - (M_1 + M_2)/M_{\text{thr}}}{d} \right] \right)\right),$$




Figure 4. Distribution of sampled masses of BNS populations for LVK O4 and O5 simulation. The uniform distribution for the slow NS (blue) is from Farrow et al. (2019). The two peak Gaussian for the slow NS (orange) is from Galaudage et al. (2021). The distribution for the recycled NS (green) is shared by both models.

(8)

where $a = -31.335$, $b = -0.9760$, $c = 1.0474$, and $d = 0.05957$.

Finally, M_{thr} , the mass threshold for prompt black hole collapse, is computed using the following (Bauswein et al. 2013),

$$M_{\text{thr}} = \left(2.38 - 3.606 \frac{M_{\text{TOV}}}{R_{1.6M_\odot}} \right) M_{\text{TOV}}. \quad (9)$$

If $m_1 + m_2 \geq M_{\text{thr}}$, then we set both dynamical and wind ejecta to zero, ensuring there is no luminous remnant.

3.2 A_V distribution

Modeling the extinction of kilonovae host galaxies based on empirical data is unlikely to yield accurate results given the single data point - host NGC 4993 for GW170817 (Pan et al. 2017). For this reason, we sample from an extinction distribution for galaxies known to host supernovae.

Kessler et al. (2009) (Equation 18) inferred the mean reddening parameter for host galaxies using a SDSS-II sample of Type 1a supernovae and found that their extinction can be well-explained by an exponential function of the form

$$P(A_V) = \exp\left(\frac{-A_V}{\tau_V}\right), \quad (10)$$

where $\tau_V = 0.334 \pm 0.088$. We sample from this distribution with a fixed $\tau_V = 0.334$.

3.3 Spatial and distance distributions

For each set of trials, we first define a cube of length l within which we simulate the events. We sample x , y , and z coordinates from a uniform, random distribution in the range of $[-\frac{l}{2}, +\frac{l}{2}]$ where l is specific to the simulation and detailed in Section 4. The Cartesian coordinates are converted to spherical coordinates to compute the event RA and Dec values. The euclidean distance is computed using the following:

$$D = 0.05 \text{ Mpc} + \sqrt{x^2 + y^2 + z^2} \quad (11)$$




Figure 5. Light curves for GW170817 constructed using interpolated spectra with fit parameters computed by Dietrich et al. (2020), overplotted with real photometry. Appendix B provides a complete list of the sources for all the photometry used in this plot.

The additional $0.05 Mpc$ term is added to ensure a minimum distance for events. We use these distances to compute the redshifts, assuming a flat Lambda-CDM cosmology¹. Since we are only simulating mergers $\leq 500 Mpc$, sampling distances is roughly equivalent to sampling redshifts.

3.4 Φ distribution

Φ describes the half-opening angle at which the lanthanide-rich component of the dynamical ejecta is distributed close to the equatorial plane during the merger giving rise to the red kilonova. The remainder of the ejecta is expelled further from the equator resulting in the blue kilonova. The lack of comprehensive data on kilonovae means that we do not have reliable distributions for Φ . For this reason, we choose to sample from a uniform continuous distribution of values for Φ in the range $[15, 75]$, the entire range for values for which complete simulations exist (See Section 2.1).

3.5 $\cos \Theta$ distribution

Given the random distribution of BNS mergers in space, it is safe to assume that the cosine of the observing angle will be uniformly-distributed. Specifically, we sample from a continuous distribution of $\cos \Theta$ values in the range $[0, 1]$.

We also use the observing angle to compute the inclination, defined as the angle between the line of sight and the total angular momentum, (Chen et al. 2019) (Ω) which is used as a parameter in the GW waveform generation (Section 4.2).

$$\Omega = \min(\Theta, 180 - \Theta) \quad (12)$$

Parameter	Distribution
A_V	$\exp\left(\frac{-A_V}{\tau_V}\right)$
Φ	$U(15, 75)$
$\cos \Theta$	$U(0, 1)$

Table 3. Table of input parameters distributions used for LVK O4.

4 MONTE CARLO TRIALS

For each trial, we sample the BNS component masses, distances, coordinates, A_V , Φ , and $\cos \Theta$ from the distributions mentioned above. The criteria used for determining gravitational wave and electromagnetic detections are specified below. We found that the expected numbers of discoverable kilonovae begin to converge after a few hundred iterations of our simulations for both the O4 and O5 observing runs. For this reason, we run all our simulations for 1000 iterations since we do not expect the results to change significantly with further increase in the number of trials.

4.1 Finding the number of events

For each trial, we first need to find the number of events, henceforth called n_{events} where:

$$n_{\text{events}} = \text{rate} \cdot \text{volume} \cdot \text{time} \quad (13)$$

The time is computed from the overlap duration of our chosen optical survey and the LVK observing run. Based on Table C2, we know that the maximum distances for BNS GW detections is $\sim 253 Mpc$ and $\sim 449 Mpc$ for LVK observing runs O4 and O5, respectively. Thus, the lengths of our event cube for simulations, l , are set to 510 Mpc and 910 Mpc respectively. This ensures that the limiting factor for kilonova discovery is always either the sensitivity of the LVK detectors or the limiting magnitude of our survey.

The merger rate model is another configurable option in our simulations. Considerable work has been done to understand the frequency of BNS mergers (Nitz et al. 2021; Abbott et al. 2023). For this work, we set the BNS merger rate to $210_{-120}^{+240} \text{Gpc}^{-3} \text{yr}^{-1}$ with log-normal uncertainties, in accordance with the LVK user guide²(Figure 6) (Abbott et al. 2023).

4.2 Detecting gravitational waves from mergers

The maximum distance values at which a BNS merger is detectable is a function of the component masses of our binary system and its inclination (Chen et al. 2021).

Once we sample our mass distributions to find the component masses, we compute its gravitational waveform which acts as our signal. This waveform is parameterized by the masses of our coalescing binary and the inclination angle, Ω . Since we are dealing with binary neutron stars, we use the TaylorF2 waveform (Buonanno et al. 2009; Messina et al. 2019), which assumes neutron stars are non-spinning point masses and has been used for BNS merger rate modeling before (Nitz et al. 2021). Given that we only use the waveform to determine the maximum distance at which a detector would be able to discover a merger, the TaylorF2 waveform is sufficiently

¹ <https://docs.astropy.org/en/stable/api/astropy.cosmology.FlatLambdaCDM.html> ² <https://emfollow.docs.ligo.org/userguide/capabilities.html>




Figure 6. Distribution of BNS merger rates approximates the log normal distribution mentioned in the LVK user guide.

accurate (compared to a more comprehensive model like IMRPhe-nomPv2_NRTidal) and much faster to compute, a key advantage for the speed of our Monte Carlo trials.

For each instrument, we use the PSD² which describes the noise at a given frequency. We then use a signal to noise ratio threshold of 8 (to remain consistent with LVK²) (Petrov et al. 2022) to determine the maximum distance at which such a merger would produce a detection. Since we already know the luminosity distances for each of our mergers, we can determine if the event would produce a detection for each of the instruments.

Another aspect to take into account is the correlation in uptimes between the LVK detectors and their respective duty cycles. We used data on the correlation between different detectors from the LIGO O3a run³ to create a correlation matrix where the rows and columns are ordered by LIGO Hanford, LIGO Livingston, Virgo, and KAGRA respectively. Since we do not have duty cycle correlation data for KAGRA during O3, we assume the same $\sim 56 - 58\%$ correlation as Virgo. These values are consistent with the current $\sim 58\%$ duty cycle correlation between LIGO - Livingston and LIGO - Hanford reported for LVK O4 during the September 21, 2023 LVEM call⁴.

$$\text{COR} = \begin{bmatrix} 1.0 & 0.56 & 0.56 & 0.56 \\ 0.56 & 1.0 & 0.58 & 0.58 \\ 0.56 & 0.58 & 1.0 & 0.56 \\ 0.56 & 0.58 & 0.56 & 1.0 \end{bmatrix} \quad (14)$$

We use this to create a detector uptime correlation matrix of dimension 4×4 . For each trial we also create a matrix of random numbers in the range $[0,1]$ of dimension $n_{\text{events}} \times 4$. We multiply the random numbers with the correlation matrix and scale all values to the range $[0,1]$, using a min-max scaler, resulting in a matrix of dimensions $n_{\text{events}} \times 4$. Each column of this matrix is a series that represents the probability the detector is active when each merger in the trial is taking place. Since we already know the duty cycles for each detector (Table 4), we can set the detector to observe during an event if this probability during the event is less than the value of the duty cycle for that instrument. We can repeat this for all four




Figure 7. Predictions for BNS mergers detected by LVK during O4 using either the Farrow et al. (2019) (blue curves) or Galaudage et al. (2021) (orange curves) mass models. **Top:** Distribution of the number of GW-detected BNS mergers for our Monte-Carlo trials. The average number of mergers is represented by a vertical dashed line. **Middle:** Distribution of chirp masses for all BNS mergers with GW detections. **Bottom:** Kernel density estimate contours (corresponding to 20%, 50%, and 80% of the probability mass) for the luminosity distance as a function of chirp mass for all BNS mergers with GW detections. GW170817 and GW190425 are represented by red and black points (dashed vertical lines), respectively, in the bottom (middle) panel.

³ https://gwosc.org/detector_status/O3a/

⁴ <https://wiki.gw-astronomy.org/OpenLVEM/Telecon20230921>

Detector	Operating Months	O4 Duty cycle
LIGO - Livingston	18	0.7
LIGO - Hanford	18	0.7
Virgo	12	0.47
KAGRA	7	0.27

Table 4. Duty cycles for detectors used for the LVK O4 Monte Carlo simulations

detectors and determine which instruments would be on during each of our mergers in a given trial.

If a detector is on and observing during a merger and the merger is within the detection range for the component masses at their inclination, then we consider the event to be detected since we are not modeling any coincidental terrestrial noise, antenna patterns on the detector, or accounting for software failures. We then determine how many instruments will detect the merger by checking each detector’s status and its detection capability.

It is important to note that non-detections, in the context of GW events, can encode vital information that can help improve sky map localization. For instance, the non-detection of GW170817 by Virgo, despite observing during the event, helped narrow down the localization (Abbott et al. 2017a). However, since we are not generating sky maps for this work, we choose to ignore antenna patterns while considering detections.

A duty cycle of 70% was used based on the observing capabilities as reported in the LVK Userguide². However, both Virgo and KAGRA will not be operating for the entirety of the 18 month period, so we encode this information into their duty cycles.

$$\text{duty cycle} = 0.7 \frac{\text{operating months}}{18} \quad (15)$$

Based on the latest observing plan⁵ available at this time (dated October 14, 2023), we assume that Virgo will operating at optimum sensitivity for 12 months and KAGRA for 7 months. Table 4 details the duty cycles for all 4 detectors in the LVK network used in our simulations.

At this stage of the pipeline, we already have data on the BNS mergers that were detected in our simulation. Fig 7 shows the properties of these mergers. We find that the median number of GW detections for merging neutron stars over LVK O4 is $\sim 3 - 4$, depending on the mass model used.

4.3 Detecting EM counterparts from mergers

As mentioned in the Section 3, we can find the $m_{\text{ej}}^{\text{wind}}$ and $m_{\text{ej}}^{\text{dyn}}$ for each merger. If the $m_{\text{ej}}^{\text{total}} > 0$, then we conclude that the merger has left behind a kilonova. Thus, we use the SED approximation method described in Section 2 to produce synthetic light curves for all mergers that have non-zero $m_{\text{ej}}^{\text{total}}$. We can use the synthetic photometry and the survey’s detection thresholds to find the discovery magnitude and peak magnitude of each detectable kilonova.

Next, we need to account for the fraction of events that would be lost to light from the sun, called $F_{\text{sun loss}}$. This value changes for different surveys and is a configurable option. We uniformly sample a number between 0 and 1 for each of the n_{events} ; if this number is greater than $F_{\text{sun loss}}$, then the event is not lost to the sun.

⁵ <https://observing.docs.ligo.org/plan/>




Figure 8. Distribution of apparent peak magnitude for all kilonovae simulated, including those without an EM or GW detection (up to 40 magnitudes) for LVK O4.

Finally, we label an event as n_{good} if it was detected by n GW detector(s), has non-zero ejecta, has a peak magnitude that can be detected by the survey, and is not lost to the sun since these filters provide all the necessary, but not sufficient, conditions for the discovery of the kilonova. Based on our simulations, typically $\sim 1 - 3\%$ of the BNS merger have a prompt collapse to black holes, resulting in zero ejecta.

Since there are 4 detectors in the LVK network we will be doing our analysis for $n = 1, 2, 3, \text{ and } 4$. It is worth noting that a single instrument detection will typically yield very poorly localized sky maps (on the order of ten thousand sq degrees) and lower network SNR which makes targeted search for kilonovae difficult.

This concludes the entire methodology we use in order to estimate the rate of discoverable BNS kilonovae.

5 RESULTS

In this section we will discuss the results of our Monte Carlo trials. It is important to note that these results indicate the best case scenario for kilonova detection since they do not account for observing inefficiencies (like weather, tiling etc.) or poor gravitational wave skymap localizations.

5.1 LVK O4 observing run

For this simulation we use both the Galaudage et al. (2021) and Farrow et al. (2019) mass distribution, a detection threshold of 23 magnitude, the DECam r passband for detections, the LVK User guide rates for BNS mergers, and a sun loss fraction of 0.5. In reality, the sun loss fraction is dependent on the specific follow-up survey we consider, the site(s) for ground-based facilities, and the location of the KN on the sky (and in particular, for space-based facilities the overlap between the GW localization and the allowed viewing area). Determining if an event will be lost to light from the sun must be done on an event by event basis, taking into account the survey strategy. Additionally, given that the search for real KNe will be done by a network of both public and private telescopes, which may elect to not share information about a counterpart for several hours

Run	Survey	BNS mass model	GW only	1 GW + EM	2 GW + EM	3 GW + EM	4 GW + EM	All GW + EM	\mathbb{E} (KN)
O4	DECAM	Galaudage et al. (2021)	4^{+7}_{-3}	0^{+2}_{-0}	1^{+3}_{-1}	0^{+1}_{-0}	—	2^{+3}_{-2}	2.2
		Farrow et al. (2019)	3^{+6}_{-3}	0^{+2}_{-0}	1^{+2}_{-1}	0^{+1}_{-0}	—	1^{+4}_{-1}	1.8
O5	LSST	Galaudage et al. (2021)	42^{+47}_{-25}	5^{+7}_{-4}	13^{+16}_{-8}	0^{+2}_{-0}	1^{+2}_{-1}	19^{+24}_{-11}	21.6
		Farrow et al. (2019)	42^{+54}_{-24}	5^{+8}_{-4}	13^{+16}_{-9}	0^{+2}_{-0}	1^{+2}_{-1}	19^{+24}_{-11}	21.7

Table 5. Summary of kilonova discovery estimates over LVK O4 and O5 from this work. Last column presents the expected number for KNe. All other columns in this table represent the middle 90% credible intervals with the median, 5^{th} , and 95^{th} percentile numbers being reported.

Mean luminosity distances (in Mpc)

Run	BNS mass model	1 GW	2 GW	3 GW	4 GW
O4	Galaudage et al. (2021)	124	130	83	—
	Farrow et al. (2019)	118	122	76	—
O5	Galaudage et al. (2021)	228	231	110	89
	Farrow et al. (2019)	228	231	108	90

Table 6. Summary of mean luminosity distances (in Mpc) of discoverable KNe for LVK O4 and O5 based on simulations with 1, 2, 3, or 4 coincidental GW detection(s) and an EM detection. The missing statistics for the 4-detector events during LVK O4 is due to a negligible number of mergers having coincidental detections on 4 instruments.

Mean peak magnitudes (AB)

Run	BNS mass model	1 GW	2 GW	3 GW	4 GW
O4	Galaudage et al. (2021)	20.5	20.6	19.5	—
	Farrow et al. (2019)	20.5	20.6	19.5	—
O5	Galaudage et al. (2021)	21.9	21.9	20.4	20.8
	Farrow et al. (2019)	21.9	21.9	20.3	20.8

Table 7. Summary of mean apparent AB magnitudes of discoverable KNe for LVK O4 and O5 based on simulations with 1, 2, 3, or 4 coincidental GW detection(s) and an EM detection. The missing statistics for the 4-detector events during LVK O4 is due to a negligible number of mergers having coincidental detections on 4 instruments.

after discovery, makes modeling this effect infeasible without several assumptions. For the sake of simplicity, we choose to encode this information using a constant 0.5 fraction.

As evident from Table 5 and Figure 9, the median number of mergers with detectable electromagnetic counterpart over all of the LVK O4 is $\sim 1 - 2$, depending on the mass model used. The figure also shows the expected values and distributions for event distances and magnitudes while Table 5 breaks down the events by the number coincidental GW detections.

Additionally, we found the distribution for the discovery window, defined as the time for which the kilonova is brighter than the limiting magnitude of the survey, for 1, 2, and 3-detector events (Table 8). Even though the times shown are shorter than the 10+ days of observations obtained for GW170817 (Figure 5), we should still, on average, have several days to get detections of the EM counterpart.

Discovery windows (days)

BNS mass model	N	90% credible	Mean
Galaudage et al. (2021)	1	$3.4^{+3.4}_{-2.0}$	3.60
	2	$3.2^{+3.4}_{-2.0}$	3.45
Farrow et al. (2019)	3	$4.4^{+3.8}_{-2.2}$	4.82
	1	$4.0^{+3.0}_{-2.0}$	4.29
	2	$4.0^{+3.0}_{-2.2}$	4.13
	3	$5.8^{+3.2}_{-3.4}$	5.82

Table 8. Discovery windows (in days) for KNe during LVK O4 for N = 1, 2, and 3 detector events in DECAM r-band. The missing statistics for the 4-detector events is due to a negligible number of mergers having coincidental detections on 4 instruments. All these discovery windows are significantly shorter than the 10+ days for which SSS17a was discoverable.

5.2 Looking ahead - LVK O5

LVK O5 presents the next opportunity for finding kilonovae, post-GW trigger. With the observing run slated to begin at the end of 2026 with a proposed end in the middle of 2029, we expect the Vera Rubin Observatory to be operational for the entirety of O5. This section aims to paint a picture of what the next ~ 5 years of kilonova discovery could look like. Once again, we use the most updated PSDs for the O5 run from LVK which, notably, represent the high end targets of BNS ranges for LIGO and KAGRA and the low end for Virgo.

The predictions presented in this section must be assessed with the added context that the sensitivities used are the targeted, optimistic values and real PSDs during O5 might not achieve these goals. For this reason, another analysis for LVK O5 will likely be required once the PSDs and observing plans are defined more concretely. Regardless, we present tentative numbers here since they will be useful for planning and forecasting.

With this caveat, we predict the median number of mergers with detectable electromagnetic counterpart over all of the LVK O5 to be ~ 19 . As evident from Figure 10 and Table 5, an updated LVK network during O5 with significantly improved sensitivities may present the first opportunity to discover a small sample of kilonovae which would enable exciting new population studies furthering both transient astronomy and cosmology.

5.3 Comparison with current LVK O4 results

To test the consistency of our pipeline with empirical observations, we simulate a partial LVK O4 run to compare with the actual ongoing




Figure 9. LVK O4 Top pane shows results from using the [Farrow et al. \(2019\)](#) mass model while the bottom pane shows results from using the [Galaudage et al. \(2021\)](#) mass model. **Left:** Distribution of the number of EM detectable events for 1, 2, and 3 GW detector events. Solid black line shows the distribution of the total number of discoverable kilonovae. **Center:** Distribution of distances of EM detectable events for 1, 2, and 3 GW detector events. **Right:** Distribution of peak and discovery magnitudes of EM detectable events for 1, 2, and 3 GW detector events. Solid lines represent distributions of peak magnitude, while dotted lines represent distributions of discovery magnitude. Table 5, Table 6, and Table 7 summarizes the results.

LVK O4 run. We adjust the simulation parameters to match the current O4 run by setting:

- the duration to be ~ 4.67 months, reflecting the current O4 period of May 24, 2023 to October 14, 2023;
- the duty cycles for Virgo and KAGRA to zero, since they are not observing during the current period;
- the duty cycles of the two LIGO detectors to 70% ².

All other configurable parameters mirror the full O4 simulation discussed in section 5.1.

This procedure allows us to assess the validity of our model, given that we have not detected any BNS mergers during the first ~ 4.67 months of LVK O4 (as of October 14, 2023). Using the [Galaudage et al. \(2021\)](#) mass model, we found that the median number of discoverable kilonovae in the first ~ 4.67 months of LVK O4, to be 0^{+2}_{-0} with 1^{+3}_{-1} BNS merger detections.

5.4 Comparison with complimentary work

Complimentary work has been done in the past to understand the detection rates of KN for surveys like the Zwicky Transient Facility (ZTF), the Wide-Field Infrared Transient Explorer (WINTER), and the Vera Rubin Observatory ([Colombo et al. 2022](#); [Frostig et al. 2022](#); [Weizmann Kiendrebeogo et al. 2023](#)). Table 9 summarizes these results while Table 5 reports the results from this work. Our independent analysis with the ZTF - r band, a limiting magnitude

Work	Band	KN Detections
Frostig et al. (2022)	J	1^{+2}_{-1} (over LVK O4)
Weizmann Kiendrebeogo et al. (2023)	r	$0.43^{+0.58}_{-0.26}$ (per year)
Colombo et al. (2022)	J	$2.4^{+3.6}_{-1.8}$ (per year)
Colombo et al. (2022)	r	$5.1^{+7.8}_{-3.8}$ (per year)

Table 9. Comparison of results from analysis for kilonova detection rates from complimentary work for LVK O4.

of 21.4, and the [Galaudage et al. \(2021\)](#) mass model predicts the number of discoverable kilonovae to be 1^{+3}_{-1} over the 18 month LVK O4.

5.5 Retrospective analysis for LVC O2 and O3

Since our flexible framework can easily adopt different PSDs over arbitrary observing durations to simulate discovery rates, we have performed our analysis for the LVC O2 and O3 observing campaigns. We note that this since the BNS merger rates used in this work are themselves inferred from the results of the O2 and O3 runs, these simulations are somewhat self fulfilling. However, to the extent that this model accurately represent the true BNS merger rate, we can




Figure 10. LVK O5 Top pane shows results from using the Farrow et al. (2019) mass model while the bottom pane shows results from using the Galaudage et al. (2021) mass model. **Left:** Distribution of the number of EM detectable events for 1, 2, 3, and 4 GW detector events. Solid black line shows the distribution of the total number of discoverable kilonovae. **Center:** Distribution of distances of EM detectable events for 1, 2, 3, and 4 GW detector events. **Right:** Distribution of peak and discovery magnitudes of EM detectable events for 1, 2, 3, and 4 GW detector events. Solid lines represent distributions of peak magnitude, while dotted lines represent distributions of discovery magnitude. Table 5, Table 6, and Table 7 summarizes the results.




Figure 11. Comparison of our results to different KNe detection rate analysis done in the past for LVK O4. MM1 and MM2 correspond to the Galaudage et al. (2021) and Farrow et al. (2019) BNS mass models respectively

still gain valuable insights into the predictive capabilities of our framework.

For this analysis, we use the PSDs from the GWTC-1 (Abbott

et al. 2019b)⁶ and GWTC-2 catalogs (Abbott et al. 2021a)⁷. The O2 campaign ran from November 30, 2016 to August 25, 2017 while the O3 campaign ran from April 1, 2019 to March 27, 2020. With the exception of the campaign duration, PSDs, and the absence of KAGRA, all other simulation configurations were identical to the ones used for LVK O4. Additionally, we exclusively used the Galaudage et al. (2021) BNS mass model for this exercise.

We predict the number of detectable BNS mergers over the O2 run to be 1_{-1}^{+2} with 0_{-0}^{+2} discoverable KNe (90% credible). Only $\sim 32\%$ of our trials had ≥ 1 discoverable KNe during this period, consistent with the single discovery of GW170817 (Abbott et al. 2017a) and SSS17a (Coulter et al. 2017) during O2.

For the O3 run, we predict the number of detectable BNS mergers to be 1_{-1}^{+3} with 0_{-0}^{+2} discoverable KNe (90% credible). Despite the increased sensitivity from O2, only $\sim 49\%$ of our trials had ≥ 1 discoverable KNe during this period. Nevertheless, these results are also consistent with the single discovery of GW190425 (Abbott et al. 2020) and no KN discovery during O3.

⁶ <https://dcc.ligo.org/LIGO-P1800374/public>

⁷ <https://dcc.ligo.org/LIGO-P2000251/public>

6 CONCLUSION

These results paint kilonovae like SSS2017a with confirmed GW detections, gamma ray bursts, counterpart photometry, and spectroscopy as incredibly rare. The low intrinsic rate of detectable kilonovae is compounded by difficulties in locating the counterparts in poorly localized skymaps that, at this time, routinely have 90% confidence intervals that span on the order of 10^3 sq deg. Such large localization are simply impractical to probe efficiently without wide field of view surveys (Weizmann Kiendrebeogo et al. 2023). These inefficiencies are difficult to model accurately and thus the numbers presented here are upper limits.

Moreover, Table 8 illustrates how our window of opportunity for finding future kilonovae will likely be significantly shorter than GW170817’s counterpart. These factors demonstrate the need for improved tooling, infrastructure, and search strategies for future KN discovery, such as TEGLOP⁸ and the systems described in Almualla et al. (2021), Bom et al. (2023) and Chatterjee et al. (2022).

Finally, a prompt chirp mass estimate from LVK, even if provided only to a tenth of a solar mass or with a small random offset applied, would allow forecasting of the electromagnetic signal. This, in turn, would enable observers to prioritize and coordinate follow-up resources more effectively, improving the yield of counterpart discoveries. Our synthetic photometry pipeline can be integrated into alerts systems, like the one described in Section A, to inform discovery and follow up strategies.

7 DATA AVAILABILITY

Software: This work makes use of NUMPY (Harris et al. 2020), ASTROPY (Astropy Collaboration et al. 2013, 2018, 2022), SNCOSMO (Barbary et al. 2016), KILOPOP (Setzer et al. 2023), SCIPY (Virtanen et al. 2020), LIGO EM BRIGHT⁹, INSPIRAL RANGE¹⁰, POSSIS¹¹, MATPLOTLIB (Hunter 2007), and PANDAS (pandas development team 2020; Wes McKinney 2010).

Data availability: All the code used in this work is publicly available at <https://github.com/uiucsn/KNmodel>.

8 ACKNOWLEDGMENTS

VS acknowledges the support of the LSST Corporation’s 2021 Enabling Science award for undergraduates for making this work possible, as well as travel support to present this research at the LSST Project and Community Workshop in 2023. GN gratefully acknowledges NSF support from AST-2206195, and a CAREER grant, supported in-part by funding from Charles Simonyi. HP is supported by an Illinois Center for AstroPhysical Surveys Graduate Student Fellowship. DC would like to acknowledge support from the NSF grants OAC-2117997 and PHY-1764464. BC is supported by the NSF Graduate Research Fellowship Program under Grant No. DGE 21-46756.

The UCSC team is supported in part by NASA grant NNG17PX03C, NSF grants AST-1815935 and AST-2307710, the

Gordon & Betty Moore Foundation, and by a fellowship from the David and Lucile Packard Foundation to R.J.F.

The SLACK application developed by this work is hosted by the Scalable Cyberinfrastructure for Multi-Messenger Astrophysics group (SCIMMA, <https://scimma.org>, PI: Narayan), which is supported by the National Science Foundation through the Office of Advanced Cyberinfrastructure awards OAC-1841625, OAC-1934752, and OAC-2311355. It is also deployed on the ANTARES broker system (<https://antares.noirlab.edu>) hosted by NSF’s National Optical and Infrared Research Laboratory (NOIRLab). This work made use of the Illinois Campus Cluster, a computing resource that is operated by the Illinois Campus Cluster Program (ICCP) in conjunction with the National Center for Supercomputing Applications (NCSA) and which is supported by funds from the University of Illinois at Urbana-Champaign. This work was partially supported by the Center for AstroPhysical Surveys (CAPS) at the National Center for Supercomputing Applications (NCSA), University of Illinois Urbana-Champaign. This work makes extensive use of software and public data products produced by the LIGO Scientific Collaboration. The work of the LSC is supported by NSF’s LIGO Laboratory which is a major facility fully funded by the National Science Foundation.

REFERENCES

- Abbott B. P., et al., 2016, *Phys. Rev. Lett.*, 116, 061102
 Abbott B. P., et al., 2017a, *Phys. Rev. Lett.*, 119, 161101
 Abbott B. P., et al., 2017b, *Phys. Rev. Lett.*, 119, 161101
 Abbott B. P., et al., 2017c, *Astrophys. J. Lett.*, 848, L12
 Abbott B. P., et al., 2017d, *Astrophys. J. Lett.*, 848, L13
 Abbott B. P., et al., 2018, *Phys. Rev. Lett.*, 121, 161101
 Abbott B. P., et al., 2019a, *Phys. Rev. X*, 9, 011001
 Abbott B. P., et al., 2019b, *Physical Review X*, 9, 031040
 Abbott B. P., et al., 2020, *ApJ*, 892, L3
 Abbott R., et al., 2021a, *Physical Review X*, 11, 021053
 Abbott R., et al., 2021b, *ApJ*, 913, L7
 Abbott R., et al., 2023, *Physical Review X*, 13, 011048
 Almualla M., et al., 2021, *MNRAS*, 504, 2822
 Andreoni I., et al., 2017, *Publ. Astron. Soc. Australia*, 34, e069
 Arcavi I., et al., 2017, *Nature*, 551, 64
 Astropy Collaboration et al., 2013, *A&A*, 558, A33
 Astropy Collaboration et al., 2018, *AJ*, 156, 123
 Astropy Collaboration et al., 2022, *apj*, 935, 167
 Barbary K., et al., 2016, SNCosmo: Python library for supernova cosmology, Astrophysics Source Code Library, record ascl:1611.017 (ascl:1611.017)
 Barnes J., 2020, *Frontiers in Physics*, 8, 355
 Barnes J., Kasen D., 2013, *ApJ*, 775, 18
 Bauswein A., Baumgarte T. W., Janka H. T., 2013, *Phys. Rev. Lett.*, 111, 131101
 Bom C. R., et al., 2023, *arXiv e-prints*, p. arXiv:2302.04878
 Bulla M., 2019, *MNRAS*, 489, 5037
 Buonanno A., Iyer B. R., Ochsner E., Pan Y., Sathyaprakash B. S., 2009, *Phys. Rev. D*, 80, 084043
 Cardelli J. A., Clayton G. C., Mathis J. S., 1989, *ApJ*, 345, 245
 Chatterjee D., Narayan G., Aleo P. D., Malanchev K., Muthukrishna D., 2022, *MNRAS*, 509, 914
 Chen H.-Y., Vitale S., Narayan R., 2019, *Physical Review X*, 9, 031028
 Chen H.-Y., Holz D. E., Miller J., Evans M., Vitale S., Creighton J., 2021, *Classical and Quantum Gravity*, 38, 055010
 Chornock R., et al., 2017, *ApJ*, 848, L19
 Colombo A., Salafia O. S., Gabrielli F., Ghirlanda G., Giacomazzo B., Perego A., Colpi M., 2022, *ApJ*, 937, 79
 Coughlin M. W., Dietrich T., Margalit B., Metzger B. D., 2019, *MNRAS*, 489, L91
 Coughlin M. W., et al., 2020, *Phys. Rev. Res.*, 2, 022006
 Coulter D. A., et al., 2017, *Science*, 358, 1556

⁸ <https://github.com/davecoulter/teglon>

⁹ <https://git.ligo.org/emfollow/em-properties/em-bright>

¹⁰ <https://git.ligo.org/gwinc/inspiral-range>

¹¹ https://github.com/mbulla/kilonova_models

- Cowperthwaite P. S., et al., 2017a, *The Astrophysical Journal. Letters (Online)*, 848
- Cowperthwaite P. S., et al., 2017b, *ApJ*, 848, L17
- Díaz M. C., et al., 2017, *ApJ*, 848, L29
- Dietrich T., Coughlin M. W., Pang P. T. H., Bulla M., Heinzl J., Issa L., Tews I., Antier S., 2020, *Science*, 370, 1450
- Drout M. R., et al., 2017, *Science*, 358, 1570
- Evans P. A., et al., 2017, *Science*, 358, 1565
- Farrow N., Zhu X.-J., Thrane E., 2019, *ApJ*, 876, 18
- Fitzpatrick E. L., 1999, *PASP*, 111, 63
- Foley R. J., Coulter D. A., Kilpatrick C. D., Piro A. L., Ramirez-Ruiz E., Schwab J., 2020, *MNRAS*, 494, 190
- Fragione G., 2021, *The Astrophysical Journal Letters*, 923, L2
- Frostig D., et al., 2022, *ApJ*, 926, 152
- Galaudage S., Adamcewicz C., Zhu X.-J., Stevenson S., Thrane E., 2021, *ApJ*, 909, L19
- Golomb J., Talbot C., 2022, *ApJ*, 926, 79
- Harris C. R., et al., 2020, *Nature*, 585, 357
- Hu L., et al., 2017, *Science Bulletin*, 62, 1433
- Hunter J. D., 2007, *Computing In Science & Engineering*, 9, 90
- Kasen D., Fernández R., Metzger B. D., 2015, *Monthly Notices of the Royal Astronomical Society*, 450, 1777
- Kasen D., Metzger B., Barnes J., Quataert E., Ramirez-Ruiz E., 2017, *Nature*, 551, 80
- Kasliwal M. M., et al., 2017, *Science*, 358, 1559
- Kessler R., et al., 2009, *ApJS*, 185, 32
- Kilpatrick C. D., et al., 2017, *Science*, 358, 1583
- Kilpatrick C. D., et al., 2021, *ApJ*, 923, 258
- Lattimer J. M., Prakash M., 2016, *Phys. Rep.*, 621, 127
- Li L.-X., Paczyński B., 1998, *ApJ*, 507, L59
- Lipunov V. M., et al., 2017, *ApJ*, 850, L1
- Margalit B., Metzger B. D., 2017a, *ApJ*, 850, L19
- Margalit B., Metzger B. D., 2017b, *ApJ*, 850, L19
- Matheson T., et al., 2021, *AJ*, 161, 107
- Messina F., Dudi R., Nagar A., Bernuzzi S., 2019, *Phys. Rev. D*, 99, 124051
- Metzger B. D., 2017, *Living Reviews in Relativity*, 20, 3
- Metzger B. D., 2019, *Living Reviews in Relativity*, 23, 1
- Murguia-Berthier A., et al., 2017, *ApJ*, 848, L34
- Nitz A. H., Kumar S., Wang Y.-F., Kastha S., Wu S., Schäfer M., Dhurkunde R., Capano C. D., 2021, *arXiv e-prints*, p. arXiv:2112.06878
- Oppenheimer J. R., Volkoff G. M., 1939, *Phys. Rev.*, 55, 374
- Pan Y. C., et al., 2017, *ApJ*, 848, L30
- Pang P. T. H., et al., 2022, NMMA: A nuclear-physics and multi-messenger astrophysics framework to analyze binary neutron star mergers ([arXiv:2205.08513](https://arxiv.org/abs/2205.08513))
- Petrov P., et al., 2022, *ApJ*, 924, 54
- Pian E., et al., 2017, *Nature*, 551, 67
- Pozanenko A. S., et al., 2018, *ApJ*, 852, L30
- Raaijmakers G., et al., 2021, *ApJ*, 922, 269
- Radice D., Perego A., Zappa F., Bernuzzi S., 2018a, *ApJ*, 852, L29
- Radice D., Perego A., Hotokezaka K., Fromm S. A., Bernuzzi S., Roberts L. F., 2018b, *The Astrophysical Journal*, 869, 130
- Radice D., Perego A., Hotokezaka K., Fromm S. A., Bernuzzi S., Roberts L. F., 2018c, *ApJ*, 869, 130
- Schlegel D. J., Finkbeiner D. P., Davis M., 1998, *ApJ*, 500, 525
- Sekiguchi Y., Kiuchi K., Kyutoku K., Shibata M., 2015, *Phys. Rev. D*, 91, 064059
- Setzer C. N., Peiris H. V., Korobkin O., Rosswog S., 2023, *MNRAS*, 520, 2829
- Shappee B. J., et al., 2017, *Science*, 358, 1574
- Shibata M., 2016, *Numerical Relativity*, doi:10.1142/9692.
- Siebert M. R., et al., 2017, *ApJ*, 848, L26
- Smartt S. J., et al., 2017, *Nature*, 551, 75
- Soares-Santos M., et al., 2017, *ApJ*, 848, L16
- Steiner A. W., Hempel M., Fischer T., 2013a, *The Astrophysical Journal*, 774, 17
- Steiner A. W., Hempel M., Fischer T., 2013b, *ApJ*, 774, 17
- Tanvir N. R., et al., 2017, *ApJ*, 848, L27
- The LIGO Scientific Collaboration et al., 2021, *arXiv e-prints*, p. arXiv:2111.03606
- Troja E., et al., 2017, *Nature*, 551, 71
- Utsumi Y., et al., 2017, *PASJ*, 69, 101
- Valenti S., et al., 2017, *ApJ*, 848, L24
- Villar V. A., et al., 2017, *The Astrophysical Journal*, 851, L21
- Virtanen P., et al., 2020, *Nature Methods*, 17, 261
- Weizmann Kiendrebeogo R., et al., 2023, *arXiv e-prints*, p. arXiv:2306.09234
- Wes McKinney 2010, in Stéfan van der Walt Jarrod Millman eds, *Proceedings of the 9th Python in Science Conference*. pp 56 – 61, doi:10.25080/Majora-92bf1922-00a
- Zhu X., Thrane E., Osłowski S., Levin Y., Lasky P. D., 2018, *Phys. Rev. D*, 98, 043002
- pandas development team T., 2020, pandas-dev/pandas: Pandas, doi:10.5281/zenodo.3509134, <https://doi.org/10.5281/zenodo.3509134>

bot-alerts-bbh ▾

**Scimma-Alert-Bot** APP 6:14 PM

Alert Type: PRELIMINARY

Superevent ID: S230814ah

Group: CBC

Event Time: 2023-08-14T23:09:01.810Z

Alert Time: 2023-08-14T23:14:08Z

FAR [1/yr]: 5.827976020973662e-14

Detectors: ['L1']

Terrestrial : 0.000

BNS: 0.000

NSBH: 0.000

BBH: 1.000

Has NS: 0.000

Has Remnant: 0.000

Has Mass Gap: 0.000


Distance (Mean): 405.201 +/- 123.899 Mpc

Distance modulus: 38.038

Join related channel: #s230814ah

[Skymap Link](#) | [Grace DB](#)

(163 kB) ▾

**Figure A1.** Screen capture of LVK alert streamed to SLACK workspace**APPENDIX A: SLACK BOT**

In order to facilitate future kilonovae discovery, we created a bot that streams LVK compact binary coalescence (CBC) and burst alerts to SLACK workspaces (Figure A1) using Scimma’s Hopskotch¹². This bot can be configured to filter alerts by the false alarm rate, likelihood of being a BNS or NSBH merger, having a luminous remnant, distance etc. It can also create different channels for events to facilitate event specific discussion. The bot is open source and publicly available at <https://github.com/scimma/slackbot> and is already operational on The Gravity Collective (Kilpatrick et al. 2021) and ANTARES (Matheson et al. 2021)¹³ workspaces.

¹² <https://scimma.org/hopskotch.html>¹³ <https://antares.noirlab.edu/loci>

Instrument	O4 PSD File	O5 PSD File
LIGO - Livingston	aligo_O4high.txt	AplusDesign.txt
LIGO - Hanford	aligo_O4high.txt	AplusDesign.txt
Virgo	avirgo_O4high_NEW.txt	avirgo_O5low_NEW.txt
KAGRA	KAGRA_10Mpc.txt	kagra_128Mpc.txt

Table C1. PSDs used for LVK O4 and O5 simulation work.

Observing Run	Instrument	Min HD (Mpc)	Max HD (Mpc)
O4	LIGO	140.68	252.23
	Virgo	90.13	162.41
	KAGRA	7.78	14.13
O5	LIGO	253.39	449.47
	Virgo	113.00	203.16
	KAGRA	100.12	180.22

Table C2. Minimum and maximum horizon distances for LVK O4 and O5 observing runs.

Parameter	Name	Sum of residuals
c	Intercept	0.0043
m	Slope	9.5348
n	Exponent	$8.2920e - 05$

Table D1. Sum of residuals for the spline surfaces created for all three scaling parameters**APPENDIX B: PHOTOMETRY CREDIT**

Figure 5 uses photometry that was originally collected by Andreoni et al. (2017), Arcavi et al. (2017), Coulter et al. (2017), Cowperthwaite et al. (2017b), Díaz et al. (2017), Drout et al. (2017), Evans et al. (2017), Hu et al. (2017), Kasliwal et al. (2017), Lipunov et al. (2017), Pian et al. (2017), Pozanenko et al. (2018), Shappee et al. (2017), Smartt et al. (2017), Tanvir et al. (2017), Troja et al. (2017), Utsumi et al. (2017), and Valenti et al. (2017). As requested in the paper with the combined data, please ensure that any use of this photometry includes appropriate citation to the original papers, in addition to the paper that compiled all the data (Villar et al. 2017).

APPENDIX C: BNS HORIZON DISTANCES

In order to compute the BNS horizon distances (HD) we use the instrument PSD files mentioned in Table C1. Table C2 show the minimum ($m_1 = 1M_\odot, m_2 = 1M_\odot$) and maximum ($m_1 = 2.05M_\odot, m_2 = 2.05M_\odot$) BNS horizon distances for the LVK O4 and O5 observing runs. These were also used to determine the dimensions of the box in which the mergers would take place.

APPENDIX D: SED SCALING PARAMETERS

This section documents the scaling parameters used for the SED extrapolation process described in Section 2.




Figure D1. Scaling laws for all Φ and $\cos \Theta = 0$ or 0.1 pairs along with the $\Delta F/F$ errors. All parameters for the linear scaling laws and the power scaling laws are provided in table D2 and D3 respectively.

cos_theta	phi	slope	intercept
0.0	15	31.676920	0.711811
0.0	30	25.882260	0.574318
0.0	45	27.711210	0.600171
0.0	60	31.520312	0.668417
0.0	75	34.086420	0.714488
0.1	15	33.154611	0.732685
0.1	30	26.505415	0.588698
0.1	45	27.910461	0.606310
0.1	60	31.571041	0.668550
0.1	75	34.043173	0.715819
0.2	15	36.546367	0.800088
0.2	30	28.526260	0.634880
0.2	45	28.587370	0.626385
0.2	60	31.732794	0.674077
0.2	75	34.009856	0.715993
0.3	15	39.916188	0.868458
0.3	30	32.279970	0.715132
0.3	45	30.015271	0.663718
0.3	60	32.140097	0.685910
0.3	75	34.059853	0.716032
0.4	15	43.143247	0.931342
0.4	30	37.554645	0.822008
0.4	45	32.495046	0.719034
0.4	60	32.918693	0.703164
0.4	75	34.180384	0.717938
0.5	15	46.681507	0.999113
0.5	30	43.192080	0.933819
0.5	45	36.254005	0.798361
0.5	60	34.184250	0.731180
0.5	75	34.385484	0.721941
0.6	15	50.388877	1.070547
0.6	30	48.115833	1.026022
0.6	45	41.308726	0.905888
0.6	60	36.168610	0.775022
0.6	75	34.729490	0.728454
0.7	15	54.182222	1.144727
0.7	30	53.329365	1.121405
0.7	45	47.011114	1.029205
0.7	60	39.336503	0.837144
0.7	75	35.277649	0.738860
0.8	15	58.138435	1.215019
0.8	30	58.757791	1.219488
0.8	45	52.297437	1.127138
0.8	60	43.892658	0.936113
0.8	75	36.259997	0.755234
0.9	15	62.179061	1.283004
0.9	30	64.414780	1.315016
0.9	45	58.136458	1.228854
0.9	60	48.871449	1.035869
0.9	75	38.187313	0.793950
1.0	15	67.452758	1.331387
1.0	30	71.137258	1.398322
1.0	45	65.139861	1.323359
1.0	60	54.699193	1.110350
1.0	75	41.802789	0.863606

Table D2. Parameters for linear scaling

cos_theta	phi	coefficient	exponent
0.0	15	21.147741	0.730248
0.0	30	17.341877	0.736150
0.0	45	18.692025	0.740490
0.0	60	21.357779	0.742585
0.0	75	23.180504	0.744509
0.1	15	22.212722	0.733037
0.1	30	17.774688	0.735778
0.1	45	18.815112	0.739729
0.1	60	21.402487	0.742854
0.1	75	23.162582	0.744420
0.2	15	24.495838	0.734150
0.2	30	19.144498	0.734297
0.2	45	19.232806	0.737476
0.2	60	21.488825	0.742021
0.2	75	23.121571	0.743968
0.3	15	26.746196	0.735150
0.3	30	21.720584	0.734577
0.3	45	20.162467	0.735113
0.3	60	21.747593	0.741105
0.3	75	23.139055	0.743795
0.4	15	28.823373	0.735425
0.4	30	25.218208	0.735069
0.4	45	21.794859	0.733628
0.4	60	22.239784	0.740152
0.4	75	23.207371	0.743607
0.5	15	31.089173	0.735924
0.5	30	28.803779	0.734589
0.5	45	24.244122	0.732731
0.5	60	23.046518	0.738911
0.5	75	23.334746	0.743398
0.6	15	33.400762	0.735875
0.6	30	31.937504	0.735144
0.6	45	27.451223	0.731421
0.6	60	24.333332	0.737719
0.6	75	23.555430	0.743193
0.7	15	35.777313	0.736108
0.7	30	35.236191	0.735871
0.7	45	31.016546	0.730061
0.7	60	26.383905	0.737269
0.7	75	23.906473	0.742878
0.8	15	38.294384	0.737310
0.8	30	38.685709	0.736969
0.8	45	34.413624	0.731455
0.8	60	29.287982	0.735747
0.8	75	24.538580	0.742689
0.9	15	40.957331	0.739484
0.9	30	42.340656	0.739015
0.9	45	38.211447	0.733714
0.9	60	32.539923	0.736270
0.9	75	25.795598	0.741992
1.0	15	44.993768	0.748327
1.0	30	47.201948	0.746468
1.0	45	43.143204	0.740656
1.0	60	36.690299	0.743248
1.0	75	28.286827	0.743213

Table D3. Parameters for power scaling