

Gauge equivalence and transdimensional perturbation: point vortices

George W. Patrick
 Department of Mathematics and Statistics
 University of Saskatchewan Saskatoon, SK S7N 5E6
 November 2, 2023

Abstract

There is an explicit resolution of the Poisson reduction of four planar point vortices, in the case that three of the vortex strengths are equal and the total vorticity is zero. The resolution, a Hamiltonian system on a unified symplectic phase space with a symmetry breaking parameter, is obtained by appending redundant states. Though single point vortices do not have the attribute of mass, there are circular assemblages with the collective dynamics of free massive particles, demonstrating a finite dimensional dynamics where mass emerges from a gauge symmetry breaking. The internal vibration of these assemblages is coupled to their collective motion and has the same functional form as the de Broglie wavelength.

1 Introduction

As is very well known, equilibria of Hamiltonian systems occur at critical points of the energy function, and their stability may be established by utilizing energy as a Lyapunov function. The stability is not asymptotic, but rather sufficiently close initial conditions yield arbitrarily near solutions. At an equilibrium the Hamiltonian vector field is zero and so admits a linearization. Energy stability implies spectral stability and KAM theory may be applicable in the converse.




Figure 1: Collective motion of O_{α_e} .

In the case of a continuous symmetry, reductions involve both quotient and momentum, with purpose to arrive at a generic Hamiltonian system, and again there may be equilibria, (relative equilibria of the original system where the solution is by a one parameter subgroup of the symmetry group). Here, though, the symplectic leaves may vary and energy stability does not directly account for nearby solutions if momentum is altered [9]. For example, with the compact symmetry group $SO(3)$, the generic reduced spaces have dimension 4 less, but 6 less at zero momentum, and the orientation acquires a slow dynamics which can be modeled as a Lagrangian system [10, 11]. If the symmetry is noncompact then solutions near a singular leaf might escape in nearby leaves (Example 5 of [15]), and in the physically relevant model of the Kirchhoff approximation for the dynamics of submerged rigid objects [3, 4], dynamic stability is established instead by KAM methods after a *resolution* of the small symplectic leaves [14, 16]. The planar 4-vortex system [12, 13], has relative equilibria (here designated O_{α_e}) where three vortices of strength $-\Gamma/3$ in the shape of an equilateral triangle rotate in a circle radius α_e around a central vortex of strength Γ . The symmetry is the (noncompact) planar Euclidean group $SE(2)$, and the translational momentum p at O_{α_e} is zero. At small $p \neq 0$, the shape of O_{α_e} becomes vibrational and is observed in simulation (Fig. 1) to crawl at velocity p/m_e , where $m_e = 8\pi\alpha_e^2/3$. The stability of this motion is an open problem.

The meaning of *resolution* may be illustrated by $SO(2) = \{A \in \mathbb{C} \mid |A| = 1\}$ acting by multiplication on




Figure 2: Resolution of the standard $SO(2)$ action on \mathbb{C} .

$\mathbb{C} = \{z\}$ (Fig. 2). The *singularity* at $z = 0$ has isotropy $SO(2)$, and the complement $z \neq 0$ (*regular sector*) has trivial isotropies. The resolution consists of the manifold with boundary $\mathbb{R}_{\geq 0} \times S^1 = \{(r, u) \mid r \geq 0, |u| = 1\}$ together with the maps $r = |z|, u = z/|z|$ and $z = ru$. The first map is a diffeomorphism to the interior $r > 0$, while the second is a surjection restricting to the inverse of the first. A space of physical states $\mathbb{C} = \{z\}$ is replaced by one on $\mathbb{R}_{\geq 0} \times S^1$ via covariance followed by smooth extension from $r > 0$ to all of $r \geq 0$, and $z = 0$ becomes multiply represented by the circle of states on the boundary $r = 0$. Physical conclusions associated to any two boundary states must agree (*gauge equivalence*), a redundancy which can be represented by the obvious action of $SO(2)$ on the boundary (*gauge symmetry*).

Realizations of Poisson manifolds occur as far back as Lie (see [19]), and in [2, Def. 1.9.1] they are surjective Poisson submersions $(M, \omega) \rightarrow (P, \{\})$ from a symplectic manifold. Here, in the 4-vortex problem, and denoting the (open dense) regular part of P as P° and the complementary singular part as P^\dagger , a resolution is a manifold with boundary A , a symplectic manifold (M, ω) , and a surjective Poisson map $\varphi: A \times M \rightarrow P$, such that $\varphi(\partial A \times M) \subseteq P^\dagger$ and $\varphi: (\text{int } A) \times M \rightarrow P^\circ$ is a Poisson isomorphism. A dynamics generated on P by a Hamiltonian H is then covered by the dynamics on $A \times M$ generated by $H \circ \varphi$, and if φ is not injective on $(\text{int } A) \times M$ then states in P^\dagger are multiply represented. If P^\dagger is an embedded submanifold of P then it is naturally a Poisson manifold by function extension and $\varphi|_{\partial A \times M}$ is a Poisson submersion ie it is the reduction of $\partial A \times M$ by the level-set equivalence relation of φ in P^\dagger .

2 Explicit resolution

The planar Euclidean group $SE(2) := \{(A, a) \mid A \in \mathbb{C}, |A| = 1, a \in \mathbb{C}\}$ with

$$\text{identity} = (1, 0), \quad (A, a)(B, b) := (AB, a + Ab), \quad (A, a)^{-1} = (A^{-1}, -A^{-1}a),$$

acts on \mathbb{C} (*standard action*) by $(A, a)z = Az + a$ (the $SE(2)$ multiplication is determined by the requirement that this is a group action). The manifold structure of $SE(2)$ is as a submanifold of $\mathbb{R}^4 \cong \mathbb{C}^2$. The Lie algebra and corresponding identifications [6] are

$$\begin{aligned} se(2) &= \{(u, v) \mid u \in \mathbb{R}, v \in \mathbb{R}^2\}, & (u, v) &\equiv \left. \frac{d}{dt} \right|_{t=0} (1 + iut, tv) \equiv (iu, v), \\ se(2)^* &= \{(\mu, \nu) \mid \mu \in \mathbb{R}, \nu \in \mathbb{C}\}, & \langle (\mu, \nu), (u, v) \rangle &:= \mu u + \nu \cdot v, \end{aligned}$$

the infinitesimal generator and exponential map are

$$\begin{aligned} \left. \frac{d}{dt} \right|_{t=0} (1 + iutz + tv) &= iuz + v, & (1) \\ \exp(u, v) &= (A, a), \quad A = e^{iu}, \quad a = \begin{cases} \frac{(e^{iu} - 1)v}{iu}, & u \neq 0, \\ v, & u = 0, \end{cases} \end{aligned}$$

and the adjoint and coadjoint actions are

$$\begin{aligned} \text{Ad}_{(A,a)}(u', v') &= (u', Av' - iu'a), & \text{ad}_{(u,v)}(u', v') &= (0, i(uv' - u'v)), \\ \text{CoAd}_{(A,a)}(\mu, \nu) &= (\mu + a \wedge Av, A\nu), & \text{coad}_{(u,v)}(\mu, \nu) &= (v \wedge \nu, iu\nu), \end{aligned}$$

where $z \wedge w := -\text{Im}(z\bar{w})$ (and also there is the notation $z, w \in \mathbb{C} \ z \cdot w := \text{Re}(z\bar{w})$). The coadjoint isotropy groups are

$$SE(2)_{(\mu,\nu)} = \begin{cases} \{A = 1, a = t\nu \mid t \in \mathbb{R}\}, & \nu \neq 0, \\ SE(2), & \nu = 0, \end{cases}$$

and these are two $SE(2)$ conjugacy classes, so there are two isotypic subsets of momenta $se(2)^{*o} = \{\mu, \nu \mid \nu \neq 0\}$ (*generic*, or *regular*) and $se(2)^{*†} = \{\mu, \nu \mid \nu = 0\}$ (*rotational* or *singular*).




Figure 3: Standard and adjoint actions.

The flow lines of the exponential of $SE(2)$ through the standard action (Fig. 3 left) are rotations of z , radius $r = |z - iv/u|$, with center of rotation at iv/u , and angular frequency u , or if $u = 0$ then translations at velocity v . For constant v and $u \rightarrow 0$, the circle centers limit to infinity in the direction iv , the velocity at z to v , and the motion to straight-line with velocity v , thus smoothly interpolating plane rotation and translation. An element $(u, v) \in se(2)$ may be thought of as a mixture of rotation and translation, with $u = 0$ corresponding to pure translation. The adjoint action is a similar picture in each constant u' plane, except in $u' = 0$ (pure translation) there are only rotations about the origin (pure translational velocity has simple rotational covariance). The flow lines of the coadjoint action (Fig. 3 right) are intersections of planes with the constant $|\nu|$ cylinders and $\nu = 0$ is an axis of fixed points. If $SE(2)$ is brought to act on a physical state corresponding to one parameter subgroup of $SE(2)$, then the generator changes similarly to the standard action (above left) while the momentum changes in an entirely different manner eg translation can unboundedly increase moment arms. This does not occur for compact symmetry groups because their Lie algebras have an invariant metric and the two actions are isomorphic (and the adjoint and coadjoint orbits are compact). As is well known, in the case of noncompact symmetry, generator and momentum covariance are distinct. Given an index set \mathcal{N} , and nonzero Γ_n , $n \in \mathcal{N}$, the *plane vortex system* [8] is a Hamiltonian system with phase space

$$\begin{aligned} P &:= \{z \in \mathbb{C}^{\mathcal{N}} \mid z_n \neq z_m \ \forall m, n\}, \\ z_n &= x_n + iy_n, \\ \omega &:= \sum_n \Gamma_n dx_n \wedge dy_n = \frac{i}{2} \sum_n \Gamma_n dz_n \wedge d\bar{z}_n, \\ \{f, g\} &= \sum_n \frac{1}{\Gamma_n} \left(\frac{\partial f}{\partial x_n} \frac{\partial g}{\partial y_n} - \frac{\partial f}{\partial y_n} \frac{\partial g}{\partial x_n} \right) = \sum_n \frac{1}{\Gamma_n} \{f, g\}_{z_n}, \\ H &:= -\frac{1}{8\pi} \sum_{m \neq n} \Gamma_m \Gamma_n \ln |z_m - z_n|^2, \\ \frac{dz_n}{dt} &= -\frac{2i}{\Gamma_n} \frac{\partial H}{\partial \bar{z}_n} = \frac{i}{2\pi} \sum_{n \neq m} \frac{\Gamma_m (z_n - z_m)}{|z_n - z_m|^2}, \end{aligned} \tag{2}$$

(see Appendix A for a summary of Wirtinger derivatives as they are used here). The system is symmetric under the diagonal action of $SE(2)$, and a momentum mapping is

$$J: P \rightarrow se(2)^*, \quad \mu = -\frac{1}{2} \sum_n \Gamma_n |z_n|^2, \quad \nu = -i \sum_n \Gamma_n z_n. \quad (3)$$

J is equivariant with respect to the action $(\mu, \nu) \mapsto \text{CoAd}_{(A,a)}(\mu, \nu) + \sigma(A, a)$, where the *cocycle* $\sigma: SE(2) \rightarrow se(2)^*$ is [1, Def. 4.2.4]

$$\sigma(A, a) = \left(J((Az_n + a)_n) - \text{CoAd}_{(A,a)} J(z) \right) \Big|_{z=0} = - \left(\sum_n \Gamma_n \right) \left(\frac{1}{2} |a|^2, ia \right).$$

Assume n from $0, \dots, N$, $\Gamma \neq 0$, $\Gamma_0 = \Gamma$, and $\Gamma_n = -\Gamma/N$, $n \geq 1$, so that $\sum_n \Gamma_n = 0$ and the momentum is CoAd -equivariant. The symmetric group S_n acts symplectically by permutation of z_1, \dots, z_n . Define the $SE(2)$ and flow invariant submanifolds $P^\dagger := \{z \in P \mid \nu = 0\}$ (*rotational sector*) and $P^\circ := P \setminus P^\dagger$ (*regular sector*), note that

$$\nu = -i\Gamma z_0 + \sum_{n=1} \frac{i\Gamma}{N} z_n = \frac{i\Gamma}{N} \sum_{n=1} (z_n - z_0) = 0 \quad \Leftrightarrow \quad \sum_{n=1} (z_n - z_0) = 0,$$

and define

$$\begin{aligned} P_1 &:= \{u = (u_0, u_1, \dots, u_N) \mid u_0 \geq 0 \text{ and } \sum_{n=1} u_n = 0, u_m \neq u_n \forall m, n \geq 1\}, \\ P_1^\dagger &= \{u \mid u_0 = 0\}, \\ P_1^\circ &:= P_1 \setminus P_1^\dagger. \end{aligned}$$

P_1 is a manifold with boundary P_1^\dagger and interior P_1° . Define the $SE(2)$ invariant map $\pi_{P^\circ}: P^\circ \rightarrow P_1^\circ$ by

$$\pi_{P^\circ}: \quad u_0 = \left| \sum_{m=1} (z_m - z_0) \right|, \quad u_n = u_0 (z_n - z_0) \left(\sum_{m=1} (z_m - z_0) \right)^{-1} - \frac{u_0}{N}. \quad (4)$$

The map π_{P° translates z_0 to the origin, the division removes rotation, and P° and P^\dagger are mapped into P_1° and P_1^\dagger respectively.

To find a right inverse of π_{P° , assume $u \in P_1$, choose $z_0 = 0$, and obtain from the second of (4) the eigenvalue problem

$$\left(u_n + \frac{u_0}{N} \right) \sum_{m=1} z_m = u_0 z_n.$$

The matrix corresponding to the left has the $N - 1$ dimensional nullspace $z_1 + \dots + z_N = 0$, and there is one u_0 eigenvector, namely $z_n = u_n + u_0/N$ because

$$\left(u_n + \frac{u_0}{N} \right) \sum_{m=1} \left(u_m + \frac{u_0}{N} \right) = \left(u_n + \frac{u_0}{N} \right) u_0,$$

so $z_n = a(u_n + u_0/N)$ where a is an undetermined complex number. Substituting into the first of (4) gives

$$u_0 = \left| \sum_{n=1} a \left(u_n + \frac{u_0}{N} \right) \right| = u_0 |a|,$$

so $|a| = 1$, and choosing $a = 1$ obtains

$$\iota_{P_1}: P_1 \rightarrow P, \quad z_0 = 0, \quad z_n = u_n + \frac{u_0}{N}. \quad (5)$$

Although π_{P° is not defined on the rotational sector P^\dagger , the map ι_{P_1} is smooth on all of P_1 . The restriction $\iota_{P_1}: P_1^\circ \rightarrow P^\circ$ is a right inverse for π_{P° and a global section to the $SE(2)$ action on P^0 ie for $u_0 > 0$ the image of the right inverse is an embedded submanifold meeting every regular sector $SE(2)$ orbit exactly once and

$\pi_{P^\circ} : P^\circ \rightarrow P_1^\circ$ is a trivial left principle bundle. The map ι_{P_1} is not unique: if $g(u)$ is any smooth $SE(2)$ -valued function then $u \mapsto g(u) \iota_{P_1}(u)$ is another. Following general practice, such a map is a *gauge*, (5) will be called the *standard gauge*, a $g(u)$ is a *local gauge transformation*, and a *gauge transformation* is the special case of where $g(u)$ is constant. Evolutions $c(t)$ on P_1° may be *reconstructed* [1, §4.3] to evolutions on P° as $g(t) \iota_{P_1}(c(t))$ where $g(t)$ is a curve in $SE(2)$, and reconstruction does not depend on gauge (the dynamics defined by (2) does not).

A dynamics on all of P_1 will be established by imposing that $\pi_{P^\circ} : P^\circ \rightarrow P_1^\circ$ is Poisson and, descending the Hamiltonian (2), and then extending the bracket and Hamiltonian to all of P_1 by continuity. The Poisson bracket of

$$f(u_0, \dots, u_N, \bar{u}_0, \dots, \bar{u}_N), \quad g(u_0, \dots, u_N, \bar{u}_0, \dots, \bar{u}_N),$$

is obtained by composition

$$\begin{aligned} \{f, g\} &= -2i \sum_{p=0} \frac{1}{\Gamma_p} \{f, g\}_{z_p} \\ &= -2i \sum_{p=0} \sum_{m,n} \frac{1}{\Gamma_p} \left(\{u_m, u_n\}_{z_p} \frac{\partial f}{\partial u_m} \frac{\partial g}{\partial u_n} + \{u_m, \bar{u}_n\}_{z_p} \frac{\partial f}{\partial u_m} \frac{\partial g}{\partial \bar{u}_n} \right. \\ &\quad \left. + \{\bar{u}_m, u_n\}_{z_p} \frac{\partial f}{\partial \bar{u}_m} \frac{\partial g}{\partial u_n} + \{\bar{u}_m, \bar{u}_n\}_{z_p} \frac{\partial f}{\partial \bar{u}_m} \frac{\partial g}{\partial \bar{u}_n} \right) \\ &= \sum_{m,n} \{u_m, u_n\} \frac{\partial f}{\partial u_m} \frac{\partial g}{\partial u_n} + \{u_m, \bar{u}_n\} \frac{\partial f}{\partial u_m} \frac{\partial g}{\partial \bar{u}_n} + \{\bar{u}_m, u_n\} \frac{\partial f}{\partial \bar{u}_m} \frac{\partial g}{\partial u_n} + \{\bar{u}_m, \bar{u}_n\} \frac{\partial f}{\partial \bar{u}_m} \frac{\partial g}{\partial \bar{u}_n}. \end{aligned}$$

Since

$$\sum_{m=1} (z_m - z_0) = -Nz_0 + \sum_{m=1} z_m = -\frac{N}{\Gamma} \left(\Gamma z_0 - \sum_{m=1} \frac{\Gamma}{N} z_m \right) = \frac{-iN\nu}{\Gamma},$$

(4) may be written

$$u_0 = \frac{N|\nu|}{|\Gamma|}, \quad u_n = \frac{u_0}{N\nu} (i\Gamma(z_n - z_0) - \nu). \quad (6)$$

The computation is simplified by the expressions (6) because the Poisson bracket is zero for any function obtained by composition with (3) (ie a function $se(2)^* = \{(\mu, \nu)\}$) and any $SE(2)$ invariant function. In particular, $\{u_m, u_0\} = 0$, $m = 0, \dots, N$, and for $m, n \neq 0$

$$\{u_m, u_n\} = -2i \sum_{p=0} \frac{1}{\Gamma_p} \frac{-u_0^2 \Gamma^2}{N^2 \nu^2} \{z_m - z_0, z_n - z_0\}_{z_p} = 0,$$

and similarly $\{\bar{u}_m, \bar{u}_n\} = 0$, while

$$\{u_m, \bar{u}_n\} = -2i \sum_{p=0} \frac{1}{\Gamma_p} \frac{-u_0^2 \Gamma^2}{N^2 |\nu|^2} \{z_m - z_0, \bar{z}_n - \bar{z}_0\}_{z_p} = 2i \left(\frac{1}{\Gamma} + \frac{1}{\Gamma_n} \delta_{nm} \right)$$

where δ_{nm} is the Kronecker delta, and also similarly $\{\bar{u}_m, u_n\} = -\{u_m, \bar{u}_n\}$, giving the reduced Poisson bracket

$$\{f, g\} = -\sum_{p=1} \frac{N}{\Gamma} \{f, g\}_{u_p} + \frac{1}{\Gamma} \sum_{n,m} \left(\frac{\partial f}{\partial u_n} \frac{\partial g}{\partial \bar{u}_m} - \frac{\partial f}{\partial \bar{u}_n} \frac{\partial g}{\partial u_m} \right). \quad (7)$$

This must be regarded as a bracket on functions on the reduced space $u_1 + \dots + u_N = 0$, which are in correspondence with functions which satisfy

$$f(u_1 + a, \dots, u_N + a) = f(u_1, \dots, u_N) \quad (8)$$

ie are invariant under the diagonal addition action of \mathbb{C} on \mathbb{C}^N . But for such invariant functions the second term in (7) is zero because

$$\begin{aligned} \sum_n \frac{\partial f}{\partial u_n} &= \frac{d}{da} \Big|_{a=0} f(u_1 + a, \dots, u_N + a) = \frac{d}{da} \Big|_{a=0} f(u_1, \dots, u_N) = 0, \\ \sum_{n,m} \left(\frac{\partial f}{\partial u_n} \frac{\partial g}{\partial \bar{u}_m} - \frac{\partial f}{\partial \bar{u}_n} \frac{\partial g}{\partial u_m} \right) &= \sum_n \frac{\partial f}{\partial u_n} \times \sum_m \frac{\partial f}{\partial \bar{u}_m} - \sum_n \frac{\partial f}{\partial \bar{u}_n} \times \sum_m \frac{\partial f}{\partial u_m} = 0, \end{aligned}$$

so, under the identification, (7) is the same bracket as the system of $N - 1$ vortices with strength $\Gamma_p = -\Gamma/N$. Use of the simpler first part of (7) (as is done below) is valid for the bracket of functions on the reduced Poisson space $u_1 + \dots + u_N = 0$ that are extended so as to satisfy (8).

Since u_0 does not evolve it can be regarded as a parameter in a 6 dimensional symplectic space, with canonical symplectic form if $\Gamma > 0$ and the negative of that if $\Gamma < 0$. Even though the quotient map is undefined on the rotational sector, the symplectic form and Hamiltonian have well defined continuous limits at $u_0 = 0$.

The Hamiltonian may be pulled back to the reduced space by composition with (5), and the sum splits over $m = 0$ (in which $z_m = 0$) and $m \geq 1$, the multiplicative constants within the logarithm may be discarded, and rescaling $u := u_0/N$, with result

$$\begin{aligned} H &= -\frac{1}{4\pi} \sum_{1 \leq m < n} \Gamma_m \Gamma_n \ln |u_m - u_n|^2 - \frac{1}{4\pi} \sum_{m=1} \Gamma \Gamma_n \ln |u_m + u|^2 \\ &= -\frac{1}{4\pi} \sum_{1 \leq m < n} \left(\frac{\Gamma}{N} \right)^2 \ln |u_m - u_n|^2 + \frac{\Gamma^2}{4\pi N} \sum_{m=1} \ln |u_m + u|^2 \end{aligned} \quad (9)$$

ie the energy of the N vortex system (but again the phase space is constrained) plus an $SO(2)$ symmetry breaking perturbation.

By way of summary, the above may be regarded as a form of Poisson reduction of the point vortex system, where the regular part of the Poisson manifold is realized, while the singular part is the boundary of that and retains a symmetry. To compare with existing well-developed Poisson reduction theory, see [2, 5, 20].

Theorem 2.1. *The phase space P is the disjoint union of the $SE(2)$ invariant Poisson submanifolds P° and P^\dagger , and the phase space P_1 is the disjoint union of P_1° and P_1^\dagger . The map $\pi_{P^\circ} : P^\circ \rightarrow P_1^\circ$ defined by (4) is a Poisson quotient for the action of $SE(2)$ on P° ie π_{P° is a Poisson submersion such that $\pi_{P^\circ}(z) = \pi_{P^\circ}(z')$ if and only if there is a $(A, a) \in SE(2)$ such that $(A, a)z = z'$. The map $\iota_P : P_1 \rightarrow P$ is an injective immersion which respects the decompositions $P = P^\dagger \cup P^\circ$ and $P_1 = P_1^\dagger \cup P_1^\circ$, the $SE(2)$ stabilizer of P^\dagger is the natural copy of $SO(2)$ in $SE(2)$, ι_P is $SO(2)$ intertwining on P_1° , and induces a diffeomorphism $P_1^\dagger/SO(2) \simeq P^\dagger/SE(2)$. Moreover,*

1. *for all integral curves $c(t) \in P_1^\dagger$, $t \in (a, b)$, there is a curve $g(t) \in SE(2)$ such that $g(t) \iota_{P_1}(c(t))$ is an integral curve in P^\dagger ; and*
2. *for all integral curves $d(t) \in P^\dagger$, $t \in (a, b)$, there is an integral curve $c(t) \in P_1^\dagger$ and a curve $g(t) \in SE(2)$ such that $d(t) = g(t) \iota_{P_1} \circ c(t)$.*

For (1), let $t_0 \in (a, b)$, $p = c(t_0)$, $\hat{c}(t) := \iota_{P_1} \circ c$, and $\hat{p} := \hat{c}(t_0)$. Choose a sequence $\langle p_i \in P_1^\circ \rangle$ converging to p and let $\hat{p}_i := \iota_{P_1}(p_i)$. Since the flow on P_1 has open domain and is continuous, for large enough i there are integral curves $c_i : (a, b) \rightarrow P_1^\circ$ such that $c_i(t_0) = p_i$, and $c_i \rightarrow c$ pointwise. Since π_{P° is Poisson, there are integral curves $\hat{d}_i : (a, b) \rightarrow P$ such that $\hat{d}_i(t_0) = \hat{p}_i$ and $\pi_{P^\circ} \circ \hat{d}_i = c_i$, and then there is an integral curve $d : (a, b) \rightarrow P^\dagger$ such that $d(t_0) = \hat{p}$. $\pi_{P^\circ} \circ \hat{d}_i = \pi_{P^\circ} \circ \hat{c}_i$, so for any particular t there is a sequence $\langle g_i \rangle$ such that $\hat{d}_i(t) = g_i \hat{c}_i(t)$. $SE(2)$ acts freely and properly, so some subsequence of g_i converges, hence $d(t)$ and $\hat{c}(t)$ are in the same $SE(2)$ orbit, and there is a unique curve $g(t)$ such that $g(t) \hat{c}(t) = d(t)$. The converse (2) is similar. ■

For $N = 3$ the reduced phase space may be canonicalized using

$$u_1 = v_1 + v_2, \quad u_2 = \theta v_1 + \theta^2 v_2, \quad u_3 = \theta^2 v_1 + \theta v_2, \quad \theta := e^{2\pi i/3}, \quad (10)$$

which is from $\mathbb{C}^2 = \{(v_1, v_2)\}$ into $u_1 + u_2 + u_3 = 0$ because the third primitive root of unity satisfies

$$1 + \theta + \theta^2 = \frac{1 - \theta^3}{1 - \theta} = 0.$$

Altering (10) to the equations

$$u_1 = v_0 + v_1 + v_2, \quad u_2 = v_0 + \theta v_1 + \theta^2 v_2, \quad u_3 = v_0 + \theta^2 v_1 + \theta v_2,$$

creates an invertible linear operator on \mathbb{C}^3 because v_0 moves orthogonally off the reduced space along its normal $(1, 1, 1)$. The inverse is therefore a diffeomorphism from $u_1 + u_2 + u_3 = 0$ to $v_0 = 0$ and it corresponds to the lowerhand 2×3 matrix of

$$\begin{bmatrix} 1 & 1 & 1 \\ 1 & \theta & \theta^2 \\ 1 & \theta^2 & \theta \end{bmatrix}^{-1} = \frac{1}{3} \begin{bmatrix} 1 & 1 & 1 \\ 1 & \theta^2 & \theta \\ 1 & \theta & \theta^2 \end{bmatrix},$$

and so

$$v_1 = \frac{1}{3}(u_1 + \theta^2 u_2 + \theta u_3), \quad v_2 = \frac{1}{3}(u_1 + \theta u_2 + \theta^2 u_3),$$

together with (10) identifies the reduced space as $\mathbb{C}^2 = \{v_1, v_2\}$. The Poisson bracket of $f(v_1, v_2)$ and $g(v_1, v_2)$ (v_1 and v_2 satisfy (8))

$$\begin{aligned} \{v_1, \bar{v}_1\} &= \frac{1}{9} \{u_1 + \theta^2 u_2 + \theta u_3, \bar{u}_1 + \bar{\theta}^2 u_2 + \bar{\theta} u_3\} \\ &= \frac{1}{9} (\{u_1, \bar{u}_1\} + \{u_2, \bar{u}_2\} + \{u_3, \bar{u}_3\}) \\ &= \frac{1}{9} \times \frac{-3}{\Gamma} \times 3 \\ &= -\frac{1}{\Gamma} \\ \{v_1, \bar{v}_2\} &= \frac{1}{9} \{u_1 + \theta^2 u_2 + \theta u_3, \bar{u}_1 + \bar{\theta} u_2 + \bar{\theta}^2 u_3\} \\ &= \frac{1}{9} (\{u_1, \bar{u}_1\} + \theta \{u_2, \bar{u}_2\} + \theta^2 \{u_3, \bar{u}_3\}) \\ &= \frac{1}{9} \times \frac{-3}{\Gamma} \times 0 \\ &= 0, \end{aligned}$$

and similarly $\{v_2, \bar{v}_2\} = -1/\Gamma$ and $\{v_2, \bar{v}_1\} = 0$, so

$$\{f, g\} = -\frac{1}{\Gamma} \{f, g\}_{v_1} - \frac{1}{\Gamma} \{f, g\}_{v_2}.$$

The Hamiltonian is by (9) and (10) (up to a constant)

$$(u_1 - u_2)(u_1 - u_3)(u_2 - u_3) = 3\sqrt{3}(v_1^3 - v_2^3) i,$$

$$(u_1 + u)(u_2 + u)(u_3 + u) = v_1^3 + v_2^3 - 3uv_1v_2 + u^3,$$

$$\begin{aligned} H &= -\frac{1}{4\pi} \sum_{1 \leq m < n} \left(\frac{\Gamma}{N} \right)^2 \ln |u_m - u_n|^2 + \frac{\Gamma^2}{4\pi N} \sum_{m=1} \ln |u_m + u|^2 \\ &= -\frac{\Gamma^2}{36\pi} \ln (|(u_1 - u_2)(u_1 - u_3)(u_2 - u_3)|^2) + \frac{\Gamma^2}{12\pi} \ln (|(u_1 + u)(u_2 + u)(u_3 + u)|^2) \\ &= -\frac{\Gamma^2}{36\pi} \left(\ln (|v_1^3 - v_2^3|^2) - 3 \ln (|v_1^3 + v_2^3 - 3uv_1v_2 + u^3|^2) \right) \\ &= -\frac{\Gamma^2}{18\pi} \left(\ln \left| \frac{v_1^3 - v_2^3}{(v_1^3 + v_2^3)^3} \right| - 3 \ln \left| 1 - \frac{(3v_1v_2 - u^2)u}{v_1^3 + v_2^3} \right| \right). \end{aligned} \tag{11}$$

The permutation group S_3 acts on the space P^1 by fixing u_0 and permuting (u_1, u_2, u_3) , and both π^0 and i^0 intertwine with the the action of S^3 on P^0 . From (10) the action of S_3 on (v_1, v_2) , determined by the generators $(1, 2, 3)$ and $(2, 3)$ is

$$(1, 2, 3)(v_1, v_2) = (\theta v_1, \theta^{-1} v_2), \quad (2, 3)(v_1, v_2) = (v_2, v_1). \tag{12}$$

If $u = 0$ then (11) is

$$H = -\frac{\Gamma^2}{18\pi} \ln \left| \frac{v_1^3 - v_2^3}{(v_1^3 + v_2^3)^3} \right| \quad (13)$$

and this is invariant under the diagonal action of $SO(2)$ ie $e^{i\theta}(v_1, v_2) = (e^{i\theta}v_1, e^{i\theta}v_2)$ because

$$\left| \frac{(e^{i\theta}v_1)^3 - (e^{i\theta}v_2)^3}{((e^{i\theta}v_1)^3 + (e^{i\theta}v_2)^3)^3} \right| = \left| e^{6i\theta} \frac{v_1^3 - v_2^3}{(v_1^3 + v_2^3)^3} \right| = \left| \frac{v_1^3 - v_2^3}{(v_1^3 + v_2^3)^3} \right|.$$

The Hamiltonian vector field of (13) is

$$\begin{aligned} \frac{dv_1}{dt} &= \frac{-1}{\Gamma} \times -2i \frac{\partial H}{\partial \bar{v}_1} = \frac{2i}{\Gamma} \frac{\partial H}{\partial \bar{v}_1} \\ H &= -\frac{\Gamma^2}{36\pi} \ln \left| \frac{v_1^3 - v_2^3}{(v_1^3 + v_2^3)^3} \right|^2, \\ \frac{\partial H}{\partial \bar{v}_1} &= -\frac{\Gamma^2}{36\pi} \left| \frac{(v_1^3 + v_2^3)^3}{v_1^3 - v_2^3} \right|^2 \frac{v_1^3 - v_2^3}{(v_1^3 + v_2^3)^3} \frac{\partial}{\partial \bar{v}_1} \left(\frac{\bar{v}_1^3 - \bar{v}_2^3}{(\bar{v}_1^3 + \bar{v}_2^3)^3} \right) \\ &= -\frac{\Gamma^2}{36\pi} \left| \frac{(v_1^3 + v_2^3)^3}{v_1^3 - v_2^3} \right|^2 \frac{v_1^3 - v_2^3}{(v_1^3 + v_2^3)^3} \left(\frac{6\bar{v}_1^2(2\bar{v}_2^3 - \bar{v}_1^3)}{(\bar{v}_1^3 + \bar{v}_2^3)^4} \right) \\ &= -\frac{\Gamma^2}{6\pi} \left| \frac{(v_1^3 + v_2^3)^3}{v_1^3 - v_2^3} \right|^2 \frac{\bar{v}_1^2(v_1^3 - v_2^3)(v_1^3 + v_2^3)(2\bar{v}_2^3 - \bar{v}_1^3)}{|v_1^3 + v_2^3|^8} \\ &= -\frac{\Gamma^2}{6\pi} \frac{v_1^6 - v_2^6}{|v_1^6 - v_2^6|^2} \bar{v}_1^2(2\bar{v}_2^3 - \bar{v}_1^3) \\ \frac{dv_1}{dt} &= -\frac{i\Gamma}{3\pi} \frac{v_1^6 - v_2^6}{|v_1^6 - v_2^6|^2} \bar{v}_1^2(2\bar{v}_2^3 - \bar{v}_1^3) \end{aligned}$$

and dv_2/dt is the same after exchange of v_1 and v_2 .

Using (5) and (10) to pull back the $SE(2)$ momentum μ (see (3))

$$\begin{aligned} \mu &= -\frac{\Gamma}{6}(3|z_0|^2 - |z_1|^2 - |z_2|^2 - |z_3|^2), \\ z_0 &= 0, \quad z_n = a \left(u_n + \frac{u_0}{N} \right) = a(u_n + u), \quad |a| = 1, \\ u &= 0, \quad u_1 = v_1 + v_2, \quad u_2 = \theta v_1 + \theta^2 v_2, \quad u_3 = \theta^2 v_1 + \theta v_2, \quad \theta = e^{2\pi i/3}, \\ \mu &= \frac{\Gamma}{6}(|u_1|^2 + |u_2|^2 + |u_3|^2) \\ &= \frac{\Gamma}{6}(|v_1 + v_2|^2 + |v_1 + \theta v_2|^2 + |\theta v_1 + v_2|^2) \\ &= \frac{\Gamma}{6}(3|v_1|^2 + 3|v_2|^2 + 2\operatorname{Re}((1 + \theta + \bar{\theta})v_1\bar{v}_2)) \\ &= \frac{\Gamma}{2}(|v_1|^2 + |v_2|^2), \end{aligned} \quad (14)$$

and the corresponding vector field is

$$\frac{dv_n}{dt} = \frac{-1}{\Gamma} \times -2i \frac{\partial \mu}{\partial \bar{v}_n} = \frac{2i}{\Gamma} \times \frac{\Gamma}{2} = iv_n$$

ie μ is a momentum map for the diagonal action of $SO(2)$.

3 Relative equilibria

Generally, for a dynamical system with phase space $\{p\}$ and symmetry $\{g\}$, an evolution $c(t)$ of the form $g(t)p_e$, $g(0) = \mathbf{1}$, is by a one-parameter subgroup: for all $s, t \mapsto c(t+s)$ is the evolution starting at $c(s) = g(s)p_e$ at time

$t = 0$, so by equivariance $c(t + s) = g(s)(g(t)p_e)$, while $c(t + s) = g(t + s)p_e$, hence $g(t + s) = g(t)g(s)$. Thus by Theorem 2.1, the relative equilibria in P^1 may be found from the $SO(2)$ relative equilibria in P_1^1 .

Theorem 3.1. *The rotational relative equilibria of the Hamiltonian system $P^0 = \{z\}$ are rotations, translations, and (z_1, z_2, z_3) permutations of (with corresponding equilibria on $\partial P^1 = \{v\}$ the $SO(2)$ orbits of)*

$$O_{\alpha_e} : z = \alpha_e(0, 1, e^{2\pi i/3}, e^{4\pi i/3}), \quad v = (\alpha_e, 0),$$

$$Y_{\alpha_e} : z = \frac{\alpha_e(1 + r_1)}{2\sqrt{1 + r_1^2}} \left((-1, 1, -2, -2) + \frac{\sqrt{3}i}{2} \frac{1 - r_1}{1 + r_1} (0, 0, 1, -1) \right), \quad v = \frac{\alpha_e}{\sqrt{1 + r_1^2}} (1, r_1),$$

$$r_1 = (1 + \sqrt{3} - \sqrt[4]{12})/2,$$

and $u = \Gamma/3\pi\alpha_e^2$, $\mu = \Gamma\alpha_e^2/2$, $v = 0$, and $\nu = 0$, for both O_{α_e} and Y_{α_e} .

The equations for the $SO(2)$ relative equilibria on P_1^1 are

$$\frac{dv_1}{dt} = -\frac{i\Gamma}{3\pi} \frac{v_1^6 - v_2^6}{|v_1^6 - v_2^6|^2} \bar{v}_1^2 (2\bar{v}_2^3 - \bar{v}_1^3) = iu_e v_1, \quad (15)$$

$$\frac{dv_2}{dt} = -\frac{i\Gamma}{3\pi} \frac{v_2^6 - v_1^6}{|v_2^6 - v_1^6|^2} \bar{v}_2^2 (2\bar{v}_1^3 - \bar{v}_2^3) = iu_e v_2. \quad (16)$$

where $u_e > 0$.

Substituting $v_2 = 0$ into (15)

$$\frac{i\Gamma}{3\pi v_1} = iu_e v_1 \quad \Leftrightarrow \quad u_e = \frac{\Gamma}{3\pi v_1^2}.$$

and (16) becomes $0 = 0$. Without loss of generality $v_1 = \alpha_e$, $\alpha_e > 0$ (since $v_2 = 0$ and $SO(2)$ acts diagonally on (v_1, v_2)), so

$$u_e = \frac{\Gamma}{3\pi\alpha_e^2}, \quad \mu_e = \frac{\Gamma\alpha_e^2}{2},$$

and substituting into (10) and using the standard gauge (5) obtains O_{α_e}

$$z_0 = 0, \quad z_1 = \alpha_e, \quad z_2 = \alpha_e e^{2\pi i/3}, \quad z_3 = \alpha_e e^{4\pi i/3}.$$

From (1)

$$v_e = \frac{dz_0}{dt} = -\frac{i\Gamma}{6\pi} \left(-\frac{z_1}{|z_1|^2} - \frac{z_2}{|z_2|^2} - \frac{z_3}{|z_3|^2} \right) = 0$$

so the center of rotation is the origin.

Without loss of generality assume $v_2 \neq 0$ and $v_1 > 0$. (15) and (16) imply $v_2 dv_1/dt - v_1 dv_2/dt = 0$, and removing nonzero factors and conjugating is

$$\bar{v}_2 v_1^2 (2v_2^3 - v_1^3) = \bar{v}_1 v_2^2 (2v_1^3 - v_2^3),$$

which after substituting $v = v_1/v_2$ is

$$v^5 - 2\bar{v}v^3 - 2v^2 + \bar{v} = 0,$$

and then substituting $v = re^{i\theta}$, $r > 0$, and separating into real and imaginary parts

$$r^4 \cos 6\theta - 2r(r^2 + 1) \cos 3\theta + 1 = 0, \quad (17)$$

$$r^3 \sin 6\theta - 2(r^2 + 1) \sin 3\theta = 0. \quad (18)$$

If $\cos 3\theta \neq 0$ and $\sin 3\theta \neq 0$ then the powers of r may be iteratively eliminated (eg form $\sin 6\theta$ times the first minus $r \cos 6\theta$ times the second) with result

$$r^3 - \cos 3\theta + r = 0, \quad r^3 \cos 3\theta - r^2 - 1 = 0,$$

$$r^2 + r \cos 3\theta - \cos^2 3\theta + 1 = 0, \quad r^2 - r \cos 3\theta + 1 = 0,$$

$$2r - \cos 3\theta = 0, \quad 3r \cos^2 3\theta - 2 \cos^2 3\theta + 2 = 0,$$

so $\cos 3\theta = 0$ or $\sin 3\theta = 0$ because the last pair obtains the contradiction $\cos^2 3\theta = 4$. Substituting $\cos 3\theta = 0$ into (18) obtains the contradiction $2(r^2 + 1) = 0$, so $\sin 3\theta = 0$, $\cos 3\theta = \pm 1$, and (17) becomes

$$r^4 - 2r(r^2 + 1)\cos 3\theta + 1 = 0.$$

This is palindromic and the substitution $r + 1/r = R$ obtains a quadratic equation

$$R^2 - 2R\cos 3\theta - 2 = 0, \quad R = \cos 3\theta \pm \sqrt{3}.$$

Now $r + 1/r = R$ is the quadratic $r^2 - Rr + 1 = 0$, which can have positive real roots only for $R > 0$ (so $\cos 3\theta = 1$) and nonnegative discriminant $8(1 \pm \sqrt{3}\cos 3\theta)$ (so $R = 1 + \sqrt{3}$), after which the solutions are among the following six:

$$\theta = 0, 2\pi, 4\pi; \quad r_1 := \frac{1}{2} + \frac{\sqrt{3}}{2} - \frac{\sqrt{12}}{2}, \quad r_2 := \frac{1}{2} + \frac{\sqrt{3}}{2} + \frac{\sqrt{12}}{2},$$

Without loss of generality $\theta = 0$, $v_1 > 0$, and $v_2 = r_1 v_1$, because $r_1 r_2 = 1$ and by (12) the permutation group S_3 acts on v as

$$(1, 2, 3)v = e^{2\pi i/3}v, \quad (2, 3)v = 1/v,$$

Using the standard gauge, the computations are

$$\begin{aligned} z_0 &= 0, \\ z_1 &= v_1 + v_2 = (r_1 + 1)v_1, \\ z_2 &= \theta v_1 + \theta^2 v_2 = \theta v_1 + \bar{\theta} v_2, \\ \operatorname{Re} z_2 &= -\frac{1}{2}v_1 - \frac{1}{2}v_2 = -\frac{1}{2}(r_1 + 1)v_1, \\ \operatorname{Im} z_2 &= \frac{\sqrt{3}}{2}v_1 - \frac{\sqrt{3}}{2}v_2 = \frac{\sqrt{3}(1 - r_1)v_1}{2}, \\ z_2 &= \left(-\frac{1}{2}(r_1 + 1) + i\frac{\sqrt{3}(1 - r_1)}{2} \right) v_1, \\ z_3 &= \theta^2 v_1 + \theta v_2 = \bar{\theta} v_1 + \theta v_2 = \bar{z}_2, \\ v_e &= \frac{dz_0}{dt} = -\frac{i\Gamma}{6\pi} \left(-\frac{z_1}{|z_1|^2} - \frac{z_2}{|z_2|^2} - \frac{z_3}{|z_3|^2} \right) = \frac{i\Gamma}{6\pi} \left(\frac{1}{z_1} + \frac{2\operatorname{Re} z_2}{|z_2|^2} \right), \\ |z_2|^2 &= (\theta v_1 + \bar{\theta} v_2)(\bar{\theta} v_1 + \theta v_2) = v_1^2 - v_1 v_2 + v_2^2 = (1 - r_1 + r_1^2)v_1^2, \\ v_e &= \frac{i\Gamma}{6\pi} \left(\frac{1}{(r_1 + 1)v_1} - \frac{r_1 + 1}{(r_1^2 - r_1 + 1)v_1} \right) = \frac{-i\Gamma r_1}{2\pi(r_1^3 + 1)v_1}, \\ \frac{2r_1^3 - 1}{r_1^6 - 1} &= \frac{1}{r_1^2 + 1}, \\ u_e &= \frac{1}{iv_1} \frac{dv_1}{dt} = \frac{(2r_1^3 - 1)\Gamma}{3\pi(r_1^6 - 1)v_1^2} = \frac{\Gamma}{3\pi(1 + r_1^2)v_1^2}, \\ \frac{iv_e}{u_e} &= \frac{\Gamma r_1}{2\pi(r_1^3 + 1)v_1} \times \frac{\pi(r_1 + 1)(r_1^3 + 1)v_1^2}{\Gamma r_1} = \frac{1}{2}(r_1 + 1)v_1. \end{aligned}$$

After the scaling $v_1 = \alpha_e / \sqrt{1 + r_1^2}$, the $SO(2)$ momentum becomes

$$\mu = \frac{\Gamma}{2}(|v_1|^2 + |v_2|^2) = \frac{\Gamma}{2(1 + r_1)^2}(\alpha_e^2 + r_1^2 \alpha_e^2) = \frac{\Gamma \alpha_e^2}{2},$$

with advantage that same α_e correspond same $SO(2)$ symplectic reduced space as O_{α_e} , and then translating the center of rotation to the origin obtains Y_{α_e} . \blacksquare

The characteristic polynomial of the linearization of O_{α_e} is $\lambda^2(\lambda^2 + u_e^2)$ ie spectrally stable. The center of rotation of Y_{α_e} is at

$$\frac{iv_e}{u_e} = \frac{3\pi\alpha_e^2 i}{\Gamma} \times \frac{-3^{1/4}\Gamma i}{6\pi\alpha} = \frac{\alpha\sqrt[4]{3}}{2}$$




Figure 4: The relative equilibria O_{α_e} and Y_{α_e} as in the phase space $P = \{z\}$. Both assemblages rotated rigidly as shown.

ie half way between z_0 and z_1 , and so z_0 and z_1 rotate diametrically opposite on an inner circle while z_2 and z_4 rotate in the same circle on opposite sides of the line between the first two. The ratio of the radius of the two circles is $\sqrt{1 + 2\sqrt{3}} = 2.113$, and the angle between the two outer vortices to the center of rotation is 37.62 degrees, and the characteristic polynomial is $\lambda^2(\lambda^2 - (2\sqrt{3} - \sqrt{3}u)^2)$ ie spectrally unstable. The heteroclinic orbits attached to Y_{α_e} are exchanges of the outer vortices.

4 Transdimensional perturbation




Figure 5: Energy levels of the $SO(2)$ reduction of the Hamiltonian system (13).

The 4-vortex $SE(2)$ symplectic reduced systems at zero translation momentum have dimension 2 (the phase space has dimension $2 \times 4 = 8$, while the momentum levels corresponding to the zero dimensional coadjoint orbits in $se(2)^{* \dagger}$ and the $SE(2)$ quotient subtract 3 each). By Thm. 2.1, these (completely integrable) reduced systems are the $SO(2)$ reductions of $\mathbb{C}^2 = \{(v_1, v_2)\}$, with Hamiltonian (13) and momentum (14). Perturbative analyses to nearby 4 dimensional symplectic spaces are through (11) with small $u > 0$; the dimension jump from 2 to 4 is spanned by the gauge group $SO(2)$ (and its momentum). The reduced spaces may be realized by the substitution $v = v_2/v_1$ into (13) and then elimination of v_1 using (14)

$$H = -\frac{\Gamma^2}{18\pi} \ln \left| \left(\frac{\Gamma}{2\mu} \right)^3 \frac{(1 + |v|^2)^3 (1 - v^3)}{(1 + v^3)^3} \right|,$$

so the energy level sets are those of the function of v inside the logarithm (Fig. 5). O_{α_e} is the equilibrium in the

center and that is surrounded by three heteroclinic orbits between the three Y_{α_e} (the other O_{α_e} is at infinity), and there are six collision states. The periodic orbits near O_{α_e} may be regarded as its internal dynamics.

Perturbation of O_{α_e} can be accomplished by slice coordinates ie coordinates near $v_1 = \alpha_e, v_2 = 0$, which split into a part within momentum level sets and transverse to the $SO(2)$ orbit (and so coordinatizing the $SO(2)$ reduced space), and another part for the momentum and the group directions (and so coordinatizing $T^*SO(2)$) [10,11,17]. Here, these are explicit, as follows: Seeking an submanifold transverse to the orbit and within the momentum level set, is is natural to posit a graph of the form

$$(q, p) \mapsto (v_1(q, p), q - ip),$$

where $v_1(q, p)$ is real (the negative on p is to remove and unwanted negative on the coordinate symplectic form). Substituting this into the momentum and equating to the O_{α_e} momentum

$$\frac{\Gamma}{2}v_1^2 + q^2 + p^2 = \frac{\Gamma}{2}\alpha_e^2, \quad v_1 = \sqrt{\alpha_e^2 - q^2 - p^2},$$

where the positive root is so that the graph passes through the O_{α_e} at $q = 0, p = 0$. To extend to other momenta, replace α_e with $2j + \alpha_e^2$

$$v_1 = \sqrt{\alpha_e^2 + 2j - q^2 - p^2}$$

so that

$$\frac{1}{2}(|v_1|^2 + |v_2|^2) = \frac{1}{2}(\alpha_e^2 + 2j - q^2 - p^2) + \frac{1}{2}(q^2 + p^2) = j + \frac{1}{2}\alpha_e^2,$$

hence j is the momentum relative the O_{α_e} momentum. Using the group action, the slice becomes

$$v_1 = \sqrt{\alpha_e^2 + 2j - q^2 - p^2} e^{i\theta}, \quad v_2 = (q - ip) e^{i\theta},$$

and an explicit local inverse at O_{α_e} is

$$q + ip = \frac{v_1 \bar{v}_2}{|v_1|}, \quad j = \frac{1}{2}(|v_1|^2 + |v_2|^2 - \alpha_e^2), \quad e^{i\theta} = \frac{v_1}{|v_1|}, \quad |v_1 - \alpha_e| < \alpha_e, \quad -\pi < \theta < \pi.$$

The symplectic form in the slice coordinates becomes $\Gamma(d\theta \wedge dj + dq \wedge dp)$, because

$$\begin{aligned} f &:= \sqrt{\alpha_e^2 + 2j - q^2 - p^2}, \\ f df &= f \times \frac{1}{2f} \times (2dj - 2q dq - 2p dp) = dj - q dq - p dp, \\ v_1 &= f \cos \theta + if \sin \theta, \\ v_2 &= (q \cos \theta + p \sin \theta) + i(q \sin \theta - p \cos \theta), \\ d(f \cos \theta) \wedge d(f \sin \theta) &= (df \cos \theta - f \sin \theta d\theta) \wedge (df \sin \theta + f \sin \theta d\theta) = f df \wedge d\theta, \\ d(q \cos \theta + p \sin \theta) \wedge d(q \sin \theta - p \cos \theta) &= -q d\theta \wedge dq - p d\theta \wedge dp - dq \wedge dp, \\ -\Gamma d(f \cos \theta) \wedge d(f \sin \theta) - \Gamma d(q \cos \theta + p \sin \theta) \wedge d(q \sin \theta - p \cos \theta) \\ &= -\Gamma(dj - q dq - p dp) \wedge d\theta - \Gamma(-q d\theta \wedge dq - p d\theta \wedge dp - dq \wedge dp) \\ &= \Gamma(d\theta \wedge dj + dq \wedge dp). \end{aligned}$$

In slice coordinates O_{α_e} corresponds to $q = p = j = 0$ and the Hamiltonian can be expanded near there using

the substitution $v_1 = fe^{i\theta}$, $v_2 = \bar{z}e^{i\theta}$, $z = q + ip$, and the rescaling ϵq , ϵp , and $\epsilon^2 j$, with result

$$\begin{aligned}
H = & \frac{\Gamma^2 \ln \alpha_e}{3\pi} \\
& - \frac{\Gamma^2 \operatorname{Re}(e^{i\theta} z)}{2\pi\alpha_e^2} u \\
& + \frac{\Gamma^2(2j - |z|^2)}{6\pi\alpha_e^2} - \frac{3\Gamma^2 \operatorname{Re}(z^2 e^{2i\theta})}{4\pi\alpha_e^4} u^2 \\
& + \frac{2\Gamma^2 \operatorname{Re}(z^3)}{9\pi\alpha_e^3} + \frac{\Gamma^2(2j - |z|^2) \operatorname{Re}(e^{i\theta} z)}{2\pi\alpha_e^4} u \\
& - \frac{\Gamma^2(2j - |z|^2)^2}{12\pi\alpha_e^4} + \frac{\Gamma^2 \operatorname{Re}(z^4 e^{i\theta})}{2\pi\alpha_e^5} u + \frac{3\Gamma^2(2j - |z|^2) \operatorname{Re}(z^2 e^{2i\theta})}{2\pi\alpha_e^6} u^2 \\
& + O(z^5, jz^3, j^2 z, u^3),
\end{aligned} \tag{19}$$

as has been obtained by Taylor expansion to order ϵ^4 followed by expansion to u^2 . In the remainder term, z^r denotes the set of homogeneous degree r polynomials in $\operatorname{Re} z$ and $\operatorname{Im} z$, and $O(S)$ denotes the ideal generated by S in the ring of smooth functions. The $SO(2)$ symmetry breaking at $u = 0$ corresponds to an absence of θ in the expansion at $u = 0$. Terms involving powers of $e^{i\theta} z$ have an unexpected $SO(2)$ *low order symmetry* of addition to θ and simultaneous reverse rotation of z (this is not the $SO(2)$ symmetry at $u = 0$), arising because $v_1 v_2 / (v_1^3 + v_2^3)$ has degree -1 and at low order is multiplied by z^1 . The expansion has been carried to ϵ^4 so as demonstrate low order symmetry breaking by the term $z^4 e^{2i\theta}$.




Figure 6: Poincaré map near O_{α_e} corresponding to $\alpha_e = 2$ and $u = .075$.

At $u = 0$, and truncating (19) at quadratic z , the corresponding differential equations are

$$\frac{d\theta}{dt} = \frac{\Gamma}{3\pi\alpha_e^2} = \xi_e, \quad \frac{dj}{dt} = 0, \quad \frac{dz}{dt} = -2i \frac{\partial H}{\partial \bar{z}} = \frac{i\Gamma}{3\pi\alpha_e^2} z = i\xi_e z.$$

Viewing O_{α_e} as an orbit, a Poincare section is $j = 0$, $\theta = 0$, and the Poincare return map is the identity since the periods of θ and z are the same. Perturbation of such a degenerate object is unlikely to yield useful information eg stability using the KAM invariant curve theorem would require a twist map [7, 18]. Numerical computations of the Poincare map do indicate stability but the usual invariant curves appear to be replaced with concentric overlapping zones (Fig. 6).

The low order symmetry can be exploited to obtain a nondegenerate estimate of the Poincare map. Removing constant terms, dividing the symplectic form and Hamiltonian by Γ , scaling time as $t' := (\Gamma/3\pi\alpha_e^2)t$, and setting $\epsilon = 1$, and truncating, obtains the $SO(2)$ -symmetric canonical Hamiltonian system

$$H = j - \frac{1}{2}|z|^2 - \frac{3u \operatorname{Re}(e^{i\theta}z)}{2} + \frac{2 \operatorname{Re}(z^3)}{3\alpha_e} + O(u^2z^2, uz^3, z^4, juz, jz^2, u^3, j^2). \quad (20)$$

Change to coordinates to (θ, k, Q, P) by

$$k = j + \frac{1}{2}|z|^2, \quad w = Q + iP = e^{i\theta}z + \frac{3u}{4}.$$

The function k is the low order momentum, and, like j , scales as ϵ^2 , and these are canonical coordinates because

$$\begin{aligned} d\theta \wedge dk &= d\theta \wedge dj + q d\theta \wedge dq + p d\theta \wedge dp, \\ dQ \wedge dP &= d(q \cos \theta - p \sin \theta + 3u/4) \wedge (q \sin \theta + p \cos \theta) = dq \wedge dp + q dq \wedge d\theta + p dp \wedge d\theta, \\ d\theta \wedge dk + dQ \wedge dP &= d\theta \wedge dj + dq \wedge dp. \end{aligned}$$

The Hamiltonian (20) becomes (the remainder is abbreviated in the first line and a constant is deleted)

$$\begin{aligned} H &= k - \left| w - \frac{3u}{4} \right|^2 - \frac{3u}{2} \operatorname{Re} \left(w - \frac{3u}{4} \right) + \frac{2}{3\alpha_e} \operatorname{Re} \left(e^{-3i\theta} \left(w - \frac{3u}{4} \right)^3 \right) + \text{h.o.t.} \\ &= k - |w|^2 - \frac{3u \operatorname{Re}(e^{-3i\theta}w^2)}{2\alpha_e} + O(ku^2, kuw, kw^2, u^3, u^2w, w^3, k^2), \end{aligned} \quad (21)$$

and the Hamiltonian vector field of (21) (the remainder is suppressed) is

$$\begin{aligned} H &= k - w\bar{w} - \frac{3u(e^{-3i\theta}w^2 + e^{3i\theta}\bar{w}^2)}{4\alpha_e}, \\ \frac{d\theta}{dt} = \frac{\partial H}{\partial k} &= 1, \quad \frac{dk}{dt} = -\frac{\partial H}{\partial \theta} = -\frac{9iu \operatorname{Re}(e^{-3i\theta}w^2)}{2\alpha_e}, \quad \frac{dw}{dt} = 2iw + \frac{3iue^{3i\theta}\bar{w}}{\alpha_e}. \end{aligned} \quad (22)$$

Use the initial condition $\theta(0) = 0$, so that $\theta = t$, and solving the last equation of (22) (a routine Fourier transform exercise), obtains

$$w(t) = (A\sqrt{\omega^-} e^{-i\omega^+t} - \bar{A}\sqrt{\omega^+} e^{-i\omega^-t})e^{2it}, \quad \omega^\pm = \frac{1 \pm \sqrt{1 - 36u^2/\alpha_e^2}}{2}, \quad (23)$$

where A is a complex constant of integration, and after substituting $t = 2\pi$, (23) implies a linear map estimate with matrix conjugate to an $O(u^2)$ rotation ie a matrix PAP^{-1} where

$$P := \begin{bmatrix} \sqrt{1 - 6u/\alpha_e} & 0 \\ 0 & \sqrt{1 + 6u/\alpha_e} \end{bmatrix}, \quad A := \begin{bmatrix} \cos 2\pi\omega^- & -\sin 2\pi\omega^- \\ \sin 2\pi\omega^- & \cos 2\pi\omega^- \end{bmatrix}.$$

5 Conclusions

That O_{α_e} can exhibit a collective motion of a massive point particle, both on the plane and the sphere, and that a Lagrangian model near the $SE(2)$ group orbit may be used to describe that, is observed in [12], and the symplectic reduced space at zero translational momentum is computed in [13]. The main contribution here is, by resolution of the Poisson singularity, that the 2-dimensional Hamiltonian dynamics at the singularity can be glued smoothly as a boundary of the regular sector of 4-dimensional ones, at the cost of an $SO(2)$ symmetry which describes a redundancy of states. Since perturbation to nonzero translational momentum is necessarily from the

singular sector to the regular sector, the emergent mass is inextricably linked to an $SO(2)$ symmetry breaking. Paradoxically, the resolved O_{α_e} becomes a relative equilibrium (so a motion) of *redundant states* ie in that view the mass emerges from a motion that is *not really there*.

Once the dimensions have been equalized, the questions become ones of ordinary Hamiltonian perturbation theory and slice coordinates at O_{α_e} and the expansion (19). However, the usual stability by confinement of KAM invariant curves requires robustly incommensurate frequencies and that fails in the extreme because of a 1-1 resonance in the associated Poincare return map. Thus the stability problem has been illuminated but remains open. Notably, the expansion (19) cannot continue convergently for small O_{α_e} radius because of the divisions by α_e . By (23), the resonance gives rise to an emergent frequency ω of second order in the translational momentum. The wavelength of this as O_{α_e} translates is

$$\lambda = \text{period} \times \text{speed} = \frac{\text{momentum}}{\text{frequency} \times \text{mass}} = \frac{2\pi p}{m\omega}.$$

As ω is proportional to the square of the momentum perturbation ie $\omega = kp^2$, the wavelength is

$$\lambda = \frac{2\pi p}{m kp^2} = \frac{2\pi/mk}{p},$$

which is the same form as the de Broglie wavelength $\lambda = 2\pi\hbar/p$.

It should be mentioned that the identification of mass by the division (as observed in simulation) of translational momentum by velocity is spurious because it involves divisions of quantities obtained from the Lie algebra $se(2)$ and its dual $se(2)^*$. For a compact group this has more substance since there is an invariant metric and the two spaces are naturally identified. For the noncompact group $SE(2)$ the identification of the dual is an arbitrary theoretical input and only mass ratios between different O_{α_e} are valid predictions.

Finally, from a purely theoretical perspective, the regular sector of the resolution of the N -vortex system is fully Poisson reduced, leaving a symplectic boundary with an addition symmetry. The resulting *Poisson manifold with boundary* may be regarded as a partial reduction. In principle, the process may be applied to the boundary and iterated until no symmetry remains, and thus viewed as a replacement for reduction itself. At the beginning of modern Poisson geometry, from Alan Weinstein [19],

The aim... is to develop the theory of Poisson manifolds with an eye toward these applications and also a new application — the study of singular limits of hamiltonian systems.

While the N -vortex system is a coarse approximation of the full hydrodynamics, the resolution of the N -vortex system derived here does seem to be consistent with this vision.

Appendix A Wirtinger derivatives

A complex valued function of two real variables $f(x, y)$ may be exchanged with a complex valued function of two complex variables $f(z, \bar{z})$ by the substitutions

$$z = x + iy, \quad \bar{z} = x - iy; \quad x = \frac{1}{2}(z + \bar{z}), \quad y = \frac{1}{2i}(z - \bar{z}),$$

and the *Wirtinger derivatives* are by definition

$$\frac{\partial f}{\partial z} := \frac{1}{2} \left(\frac{\partial f}{\partial x} - i \frac{\partial f}{\partial y} \right), \quad \frac{\partial f}{\partial \bar{z}} := \frac{1}{2} \left(\frac{\partial f}{\partial x} + i \frac{\partial f}{\partial y} \right), \quad (24)$$

regarded as expressions in z and \bar{z} eg if $f(x, y) := xy$ then $f(z, \bar{z}) = (z^2 - \bar{z}^2)/4i$ and

$$\frac{\partial f}{\partial z} = \frac{1}{2}(y - ix) = \frac{1}{2i}z, \quad \frac{\partial f}{\partial \bar{z}} = \frac{1}{2}(y + ix) = -\frac{1}{2i}\bar{z}.$$

If $f(x, y) = u(x, y) + iv(x, y)$ then

$$\begin{aligned} \frac{\partial f}{\partial z} &= \frac{1}{2} \left(\frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right) (u + iv) = \frac{1}{2} \left(\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} \right) + \frac{i}{2} \left(\frac{\partial v}{\partial x} - \frac{\partial u}{\partial y} \right), \\ \frac{\partial f}{\partial \bar{z}} &= \frac{1}{2} \left(\frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right) (u + iv) = \frac{1}{2} \left(\frac{\partial u}{\partial x} - \frac{\partial v}{\partial y} \right) + \frac{i}{2} \left(\frac{\partial v}{\partial x} + \frac{\partial u}{\partial y} \right), \end{aligned}$$

so, if $f(x, y)$ is differentiable, then \bar{z} or z is missing after conversion to $f(z, \bar{z})$ if and only if $f(z) := f(\operatorname{Re} z, \operatorname{Im} z)$ is holomorphic or antiholomorphic respectively, and then the Wirtinger derivatives are same as the usual complex analysis derivatives df/dz and $df/d\bar{z}$. Wirtinger derivatives extend analogously to any number of variables, and they satisfy the usual calculus rules, and also from (24)

$$\frac{\partial f}{\partial x} = \frac{\partial f}{\partial z} + \frac{\partial f}{\partial \bar{z}}, \quad \frac{\partial f}{\partial y} = i \left(\frac{\partial f}{\partial z} - \frac{\partial f}{\partial \bar{z}} \right).$$

This can be efficient in an application that lends itself to complex arithmetic eg

$$f(x, y) := \frac{\bar{z}}{z}, \quad \frac{\partial f}{\partial x} = \frac{\partial f}{\partial z} + \frac{\partial f}{\partial \bar{z}} = -\frac{\bar{z}}{z^2} + \frac{1}{z} = \frac{2i \operatorname{Im} z}{z^2}$$

as compared to computing $\partial f/\partial x$ from

$$f(x, y) = \frac{x^2 - y^2}{x^2 + y^2} - \frac{2ixy}{x^2 + y^2}.$$

A formula may once be converted to refer to derivatives wrt z and \bar{z} and then a given function of z and \bar{z} may not need to be converted to x and y eg the Poisson bracket of $f(z)$ and $g(z)$ is

$$\{f, g\} = \frac{\partial f}{\partial x} \frac{\partial g}{\partial y} - \frac{\partial f}{\partial y} \frac{\partial g}{\partial x} = i \left(\frac{\partial f}{\partial z} + \frac{\partial f}{\partial \bar{z}} \right) \left(\frac{\partial g}{\partial z} - \frac{\partial g}{\partial \bar{z}} \right) - f \leftrightarrow g = -2i \left(\frac{\partial f}{\partial z} \frac{\partial g}{\partial \bar{z}} - \frac{\partial f}{\partial \bar{z}} \frac{\partial g}{\partial z} \right).$$

For example, if $f = z\bar{z} = x^2 + y^2$, $g = (z^2 + \bar{z}^2)/2 = x^2 - y^2$ then

$$\begin{aligned} \{f, g\} &= -2i \left(\frac{\partial f}{\partial z} \frac{\partial g}{\partial \bar{z}} - \frac{\partial g}{\partial z} \frac{\partial f}{\partial \bar{z}} \right) \\ &= -2i((\bar{z})(\bar{z}) - (z)(z)) \\ &= -2i((x - iy)(x - iy) - (x + iy)(x + iy)) \\ &= -2i(-4ixy) \\ &= -8xy, \end{aligned}$$

$$\{f, g\} = \frac{\partial f}{\partial x} \frac{\partial g}{\partial y} - \frac{\partial f}{\partial y} \frac{\partial g}{\partial x} = (2x)(-2y) - (2y)(2x) = -8xy.$$

If $f = u + iv$ is complex valued and H is real then

$$\frac{df}{dt} = \frac{du}{dt} + i \frac{dv}{dt} = \{u, H\} + i\{v, H\} = \{u + iv, H\} = \{f, H\}, \quad \frac{dz}{dt} = \{z, H\} = -2i \frac{\partial H}{\partial \bar{z}},$$

and also

$$\{f, g\}^- = 2i \left(\frac{\partial f}{\partial z} \frac{\partial g}{\partial \bar{z}} - \frac{\partial g}{\partial z} \frac{\partial f}{\partial \bar{z}} \right)^- = 2i \left(\frac{\partial \bar{f}}{\partial \bar{z}} \frac{\partial \bar{g}}{\partial z} - \frac{\partial \bar{g}}{\partial \bar{z}} \frac{\partial \bar{f}}{\partial z} \right) = \{\bar{f}, \bar{g}\}.$$

Given

$$f(w_1, \dots, w_n, \bar{w}_1, \dots, \bar{w}_N), \quad g(w_1, \dots, w_n, \bar{w}_1, \dots, \bar{w}_N), \quad w_1 = w_1(z, \bar{z}), \dots, w_n = w_n(z, \bar{z}),$$

$$\begin{aligned} \{f, g\} &= -2i \left(\frac{\partial f}{\partial z} \frac{\partial g}{\partial \bar{z}} - \frac{\partial f}{\partial \bar{z}} \frac{\partial g}{\partial z} \right) \\ &= -2i \sum_{i,j} \left(\frac{\partial f}{\partial w_i} \frac{\partial w_i}{\partial z} + \frac{\partial f}{\partial \bar{w}_i} \frac{\partial \bar{w}_i}{\partial z} \right) \left(\frac{\partial g}{\partial w_j} \frac{\partial w_j}{\partial \bar{z}} + \frac{\partial g}{\partial \bar{w}_j} \frac{\partial \bar{w}_j}{\partial \bar{z}} \right) - (f \leftrightarrow g) \\ &= -2i \sum_{i,j} \left(\frac{\partial f}{\partial w_i} \frac{\partial g}{\partial w_j} \frac{\partial w_i}{\partial z} \frac{\partial w_j}{\partial \bar{z}} + \frac{\partial f}{\partial w_i} \frac{\partial g}{\partial \bar{w}_j} \frac{\partial w_i}{\partial z} \frac{\partial \bar{w}_j}{\partial \bar{z}} \right. \\ &\quad \left. + \frac{\partial f}{\partial \bar{w}_i} \frac{\partial g}{\partial w_j} \frac{\partial \bar{w}_i}{\partial z} \frac{\partial w_j}{\partial \bar{z}} + \frac{\partial f}{\partial \bar{w}_i} \frac{\partial g}{\partial \bar{w}_j} \frac{\partial \bar{w}_i}{\partial z} \frac{\partial \bar{w}_j}{\partial \bar{z}} \right) - (f \leftrightarrow g, i \leftrightarrow j) \\ &= \sum_{i,j} \left(\{w_i, w_j\} \frac{\partial f}{\partial w_i} \frac{\partial g}{\partial w_j} + \{w_i, \bar{w}_j\} \frac{\partial f}{\partial w_i} \frac{\partial g}{\partial \bar{w}_j} + \{\bar{w}_i, w_j\} \frac{\partial f}{\partial \bar{w}_i} \frac{\partial g}{\partial w_j} + \{\bar{w}_i, \bar{w}_j\} \frac{\partial f}{\partial \bar{w}_i} \frac{\partial g}{\partial \bar{w}_j} \right), \end{aligned}$$

If the w_i are holomorphic then $\{w_i, w_j\} = \{\bar{w}_i, \bar{w}_j\} = 0$ and

$$\begin{aligned} \{f, g\} &= \sum_{i,j} \{w_i, \bar{w}_j\} \frac{\partial f}{\partial w_i} \frac{\partial g}{\partial \bar{w}_j} + \sum_{i,j} \{\bar{w}_i, w_j\} \frac{\partial f}{\partial \bar{w}_i} \frac{\partial g}{\partial w_j} \\ &= \sum_{i,j} \{w_i, \bar{w}_j\} \frac{\partial f}{\partial w_i} \frac{\partial g}{\partial \bar{w}_j} + \sum_{i,j} \{\bar{w}_j, w_i\} \frac{\partial f}{\partial \bar{w}_j} \frac{\partial g}{\partial w_i} \\ &= \sum_{i,j} \{w_i, \bar{w}_j\} \left(\frac{\partial f}{\partial w_i} \frac{\partial g}{\partial \bar{w}_j} - \frac{\partial f}{\partial \bar{w}_j} \frac{\partial g}{\partial w_i} \right). \end{aligned}$$

References

- [1] R. Abraham and J. E. Marsden. *Foundations of mechanics*. Benjamin/Cummings Publishing Co., Inc., Advanced Book Program, Reading, Mass., second edition, 1978.
- [2] J.-P. Dufour and N. T. Zung. *Poisson structures and their normal forms*, volume 242 of *Progress in Mathematics*. Birkhäuser Verlag, Basel, 2005.
- [3] N. E. Leonard. Stability of a bottom-heavy underwater vehicle. *Automatica*, 33:331–346, 1997.
- [4] N. E. Leonard and J. E. Marsden. Stability and drift of underwater vehicle dynamics: mechanical systems with rigid motion symmetry. *Physica D*, 105:130–162, 1997.
- [5] J. E. Marsden and T. S. Ratiu. Reduction of Poisson manifolds. *Lett. Math. Phys.*, 11(2), 1986.
- [6] J. E. Marsden and T. S. Ratiu. *Introduction to mechanics and symmetry*, volume 17 of *Texts in Applied Mathematics*. Springer-Verlag, New York, second edition, 1999.
- [7] K. R. Meyer, G. R. Hall, and D. Offin. *Introduction to Hamiltonian dynamical systems and the N-body problem*, volume 90 of *Applied Mathematical Sciences*. Springer, New York, second edition, 2009.
- [8] P. K. Newton. *The N-vortex problem*, volume 145 of *Applied Mathematical Sciences*. Springer-Verlag, New York, 2001. Analytical techniques.
- [9] G. W. Patrick. Relative equilibria in Hamiltonian systems: The dynamic interpretation of nonlinear stability on the reduced phase space. *J. Geom. Phys.*, 9:111–119, 1992.
- [10] G. W. Patrick. Relative equilibria of Hamiltonian systems with symmetry: linearization, smoothness, and drift. *J. Nonlin. Sci.*, 5:373–418, 1995.
- [11] G. W. Patrick. Dynamics near relative equilibria: Nongeneric momenta at a 1:1 group–reduced resonance. *Math. Z.*, 232:747–788, 1999.
- [12] G. W. Patrick. Dynamics of perturbed relative equilibria of point vortices on the sphere or plane. *J. Nonlin. Sci.*, 10:401–415, 2000.
- [13] G. W. Patrick. Reduction of the planar 4-vortex system at zero momentum. In B. Fiedler, K. Groger, and J. Sprekels, editors, *International conference on differential equations*, volume 2, pages 1000–1008. World Scientific, 2000.
- [14] G. W. Patrick. Stability by KAM confinement of certain wild, nongeneric relative equilibria of underwater vehicles with coincident centers of mass and bouyancy. *SIAM J. Appl. Dyn. Syst.*, 2:36–52, 2003.
- [15] G. W. Patrick, R. M. Roberts, and C. Wulff. Stability of Poisson equilibria and Hamiltonian relative equilibria by energy methods. *Arch. Rational Mech. Anal.*, 174:301–344, 2004.
- [16] G. W. Patrick, R. M. Roberts, and C. Wulff. Stability transitions for axisymmetric relative equilibria of Euclidean symmetric Hamiltonian systems. *Nonlinearity*, 21:325–352, 2008.

- [17] M. Roberts, C. Wulff, and J. S. W. Lamb. Hamiltonian systems near relative equilibria. *J. Differential Equations*, 179(2):562–604, 2002.
- [18] C. L. Siegel and J. K. Moser. *Lectures on celestial mechanics*. Classics in Mathematics. Springer-Verlag, Berlin, 1995. Translated from the German by C. I. Kalme, reprint of the 1971 translation.
- [19] A. Weinstein. The local structure of Poisson manifolds. *J. Differential Geom.*, 18(3):523–557, 1983.
- [20] A. Weinstein. Poisson geometry of the principal series and nonlinearizable structures. *Journal of Differential Geometry*, 25(1):55–73, 1987.