

THE FREE BOUNDARY FOR A SEMILINEAR NON-HOMOGENEOUS BERNOULLI PROBLEM

LILI DU AND CHUNLEI YANG

ABSTRACT. In the classical homogeneous one-phase Bernoulli-type problem, the free boundary consists of a "regular" part and a "singular" part, as Alt and Caffarelli have shown in their pioneer work (*J. Reine Angew. Math.*, 325, 105-144, 1981) that regular points are $C^{1,\gamma}$ in two-dimensions. Later, Weiss (*J. Geom. Anal.*, 9, 317-326, 1999) first realized that in higher dimensions a critical dimension d^* exists so that the singularities of the free boundary can only occur when $d \geq d^*$.

In this paper, we consider a non-homogeneous semilinear one-phase Bernoulli-type problem, and we show that the free boundary is a disjoint union of a regular and a singular set. Moreover, the regular set is locally the graph of a $C^{1,\gamma}$ function for some $\gamma \in (0, 1)$. In addition, there exists a critical dimension d^* so that the singular set is empty if $d < d^*$, discrete if $d = d^*$ and of locally finite \mathcal{H}^{d-d^*} Hausdorff measure if $d > d^*$. As a byproduct, we relate the existence of viscosity solutions of a non-homogeneous problem to the Weiss-boundary adjusted energy, which provides an alternative proof to existence of viscosity solutions for non-homogeneous problems.

1. INTRODUCTION AND MAIN RESULTS

Free boundary problems are a particular type of boundary value problem where the domain is a part of the solution that is up to be determined. Typically, the classical *one-phase Bernoulli-type problem* deals with the minimization problem

$$\min \left\{ \int_{\Omega} |\nabla u|^2 + Q^2(x) \chi_{\{u>0\}} dx : u \in H^1(\Omega), u = u^0 \text{ on } S \right\}. \quad (1.1)$$

Here $\Omega \subset \mathbb{R}^d$ ($d \geq 2$) is a bounded open set and locally $\partial\Omega$ is a Lipschitz graph. $S \subset \partial\Omega$ is measurable and $\mathcal{H}^{d-1}(S) > 0$. The Dirichlet data on S is given by $u^0 \in H^1(\Omega)$, $u^0 \geq 0$. The given force function $Q(x) : \Omega \rightarrow \mathbb{R}$ is assumed to be non-negative and measurable. $\chi_{\{u>0\}}$ denotes the indicator function of the set $\{u > 0\}$. Minimizers of problem (1.1) are solutions to the following problem:

$$\Delta u = 0 \quad \text{in} \quad \Omega^+(u), \quad u = 0, \quad |\nabla u| = Q(x) \quad \text{on} \quad \partial\Omega^+(u), \quad (1.2)$$

where $\Omega^+(u) := \Omega \cap \{u > 0\}$ and $\partial\Omega^+(u) := \Omega \cap \partial\{u > 0\}$.

In this paper, we investigate a natural generalization of the above one-phase Bernoulli-type problem in the presence of a right-hand side which depends on the solution itself. Let \mathbb{R}^d , Ω , $\partial\Omega$, S , u^0 and $Q(x)$ be the same mathematical objects as in (1.1) and (1.2). Let $F(t) \in C^{2,\beta}(\mathbb{R})$ be a

The first author is supported by National Nature Science Foundation of China Grant 11971331, 12125102, and Sichuan Youth Science and Technology Foundation 2021JDTD0024.

function for some $\beta \in (0, 1)$, and let us define

$$f(t) := -\frac{1}{2}F'(t). \quad (1.3)$$

Consider the functional

$$J(u) = \int_{\Omega} |\nabla u|^2 + F(u) + Q^2(x)\chi_{\{u>0\}} dx \quad (1.4)$$

over the class of admissible functions

$$\mathcal{K} := \{u \in H^1(\Omega) : u \leq \Psi \text{ a.e. in } \Omega, \quad u = u^0 \text{ on } S\}. \quad (1.5)$$

Here $\Psi \in C^0(\bar{\Omega}) \cap C^{2,\alpha}(\Omega)$ is a given function, satisfying

$$\Delta \Psi + f(\Psi) \leq 0 \quad \text{in } \Omega, \quad \Psi > 0 \quad \text{in } \Omega, \quad 0 \leq u^0 \leq \Psi \quad \text{on } S. \quad (1.6)$$

Observe that when the free boundary $\partial\Omega^+(u)$ is smooth, the Euler-Lagrange equations of (1.4) are given by

$$-\Delta u = f(u) \quad \text{in } \Omega^+(u), \quad u = 0, \quad |\nabla u| = Q(x) \quad \text{on } \partial\Omega^+(u). \quad (1.7)$$

Our main purposes of this article is to investigate the impact of non-zero right-hand side $f(u)$ on the structure and the regularity of the free boundary. This is because that the system (1.7) has numerous applications in physics and the right-hand side $f(u)$ has many physical explanations. For instance when $d = 2$, the unknown variable u can be regarded as the scalar stream function and the function $f(u)$ stands for vorticity strength and the system (1.7) is used for modeling the incompressible inviscid jet and cavity flow with general vorticity ([11, 12]), and for the periodic water waves with non-zero vorticity ([10, 32, 33, 34]).

Let us shortly summarize the mathematical results relevant in the context of homogeneous one-phase problem (1.2). In their seminal paper [1], Alt and Caffarelli derived the smoothness of the free boundary for weak solutions. After that, Caffarelli introduced the notion of viscosity solutions to the problem (1.2) in a series of artworks [3, 4, 5], and proved the smoothness of the free boundary for a larger class of solutions. These approaches appears to be universal and were used to study quasilinear (even degenerate) Bernoulli-type problems [2, 14]. This paper aims to extend the main theorem in the paper by Alt and Caffarelli to more general semilinear situations as well as the higher-dimensional case. However, our paper replace the core of that paper, which relies on a Weiss-type monotonicity formula and a viscosity approach developed by De Silva, which promises to be applicable to one-phase problems governed by more complex operators [17].

It should be also worth noting that the appearance of singularities and the structure of the free boundary in higher dimensions ($d \geq 3$) play a significant role in the study of minimal surfaces. The first far-reaching observation from this perspective was dedicated to Weiss [35], who revealed that for minimizers of the problem (1.1), there is a critical dimension $d^* \geq 3$ such that the singular set of the free boundary is empty if $d < d^*$, and is a discrete set of isolated points if $d = d^*$, and is a closed set of Hausdorff dimension $d - d^*$ if $d > d^*$. To the best of our knowledge, $d^* \in \{5, 6, 7\}$ and we refer readers to [9, 15, 22] for detail. In this article, we also study the properties of the singular

set. We will prove a similar result as Weiss in [35] for our semilinear case. The main obstacle here is that the blow-up limit of our problem appears not to minimize the original functional so that we can no longer follow directly the classical approach of Weiss. As a final twist, we add, is the precise value of d^* , which is still an open problem, even to the homogeneous problem (1.2).

1.1. Assumptions, notations and definitions. Let $F(t) \in C^{2,\beta}(\mathbb{R})$ be a convex function for some $\beta \in (0, 1)$, satisfying $F''(t) \leq F_0$ for all $t \in \mathbb{R}$, and

$$F(0) = 0, \quad -F_0 \leq F'(t) \leq 0 \quad \text{for } t \leq 0, \quad (1.8)$$

where $F_0 \geq 0$ is a given constant. Assume that $Q(x) \in C^{0,\beta}(\bar{\Omega})$ is a function which satisfies

$$0 < Q_{\min} \leq Q(x) \leq Q_{\max} < +\infty \quad \text{for each } x \in \bar{\Omega}. \quad (1.9)$$

Here $Q_{\min} < Q_{\max}$ are two given positive constants. Note that in view of the assumptions we imposed on F , it is obvious that $f(t) \in C^{1,\beta}(\mathbb{R})$ defined in (1.3) is function which satisfies

$$0 \leq f(t) \leq \frac{F_0}{2} \quad \text{for } t \leq 0, \quad -\frac{F_0}{2} \leq f'(t) \leq 0 \quad \text{for } t \in \mathbb{R}. \quad (1.10)$$

It should be worth noting that if $x^0 \in \partial\Omega^+(u)$ is a free boundary point, then by (1.10), $f(u(x^0)) = f(0) \geq 0$ on $\partial\Omega^+(u)$. This implies that the non-linear force term is non-negative on the free boundary. This is the main difference between the one considered in [23], whose right-hand side is vanishing on the free boundary.

We further make some comments to our monotone assumptions (1.8) on $F(t)$. Firstly, $F(0) = 0$ is quite natural since $F(t)$ is actually the primitive of $f(t)$ and $F(t) := -2 \int_0^t f(s) ds$. $F'(t) \leq 0$ for $t \leq 0$ was applied to prove that minimizers are non-negative in Ω . The convexity of $F''(t) \geq 0$ is needed for applying the comparison principle. These assumptions are imposed because that all the results demonstrated in this paper are *not the conditional regularity* of the free boundary and there are no apriori assumptions on the regularity of the solutions and topological properties of the free boundary. This is the main difference between our work and other non-homogeneous problems [16].

Throughout this paper \mathbb{R}^d will be the usual Euclidean space of dimension d . We denote by c_d or C_d the constants depending only on the dimensions. For any $t \in \mathbb{R}$, we denote the *positive* and *negative* parts of t by $t^\pm := \max\{\pm t, 0\}$, so that $t = t^+ - t^-$ and $|t| = t^+ + t^-$. The open d -dimensional ball with its center at $x^0 = (x_1^0, \dots, x_d^0) \in \mathbb{R}^d$ of radius $r > 0$ and volume $\omega_d r^d$ will be denoted by $B_r(x^0) := \{x \in \mathbb{R}^d : |x - x^0| < r\}$. If in particular $x^0 = 0$, we denote by $B_r := B_r(0)$ for simplicity. The unit sphere of d -dimensional will be denoted by \mathbb{S}^{d-1} . If $u: B_R \rightarrow \mathbb{R}$ is a function expressed in polar coordinates as $u = u(\rho, \theta)$, then

$$|\nabla u|^2 = (\partial_\rho u)^2 + \frac{1}{\rho^2} |\nabla_\theta u|^2, \quad \Delta u = \partial_{\rho\rho} u + \frac{d-1}{\rho} \partial_\rho u + \frac{1}{\rho^2} \Delta_{\mathbb{S}} u, \quad (1.11)$$

where $\nabla_\theta u$ is the sphere gradient and $\Delta_{\mathbb{S}} u$ is the spherical Laplacian.

Given a set $E \subset \mathbb{R}^d$, we denote by χ_E the characteristic function of the set E . We denote by $D \subset\subset E$ the compact subsets D of E . In the text we use the d -dimensional *Lebesgue-measure* $|\cdot|$ and the m -dimensional *Hausdorff measure* \mathcal{H}^m . Sometimes, $|\cdot|$ will also represent the absolute value, depending on the context. The *Lebesgue density* for E is defined to be the limit

$$\vartheta := \lim_{r \rightarrow 0} \frac{|E \cap B_r(x^0)|}{|B_r(x^0)|}, \quad \text{where } x^0 = (x_1^0, \dots, x_d^0) \in \mathbb{R}^d.$$

It is easy to see that $\vartheta \in [0, 1]$. For any $E \subset \mathbb{R}^d$ and $\vartheta \in [0, 1]$, the set $E^{(\vartheta)}$ will be the set of points at which E has Lebesgue density equal to ϑ , namely,

$$E^{(\vartheta)} = \left\{ x^0 = (x_1^0, \dots, x_d^0) \in \mathbb{R}^d : \lim_{r \rightarrow 0^+} \frac{|E \cap B_r(x^0)|}{|B_r(x^0)|} = \vartheta \right\}. \quad (1.12)$$

Let $\partial_{\text{red}}(E) := \{x \in \mathbb{R}^d : \text{there exists } \nu \in \partial B_1 \text{ such that } r^{-d} \int_{B_r(x)} |\chi_E - \chi_{\{y: (y-x) \cdot \nu(x) < 0\}}| \rightarrow 0 \text{ as } r \rightarrow 0\}$ be the *reduced boundary* of E .

We will use the notation $J(u, D)$ for any $D \subset \Omega$ to highlight the domain of the functional which was restricted to. Sometimes we use the notation $B_1^+(u) := B_1 \cap \{u > 0\}$ and $\partial B_1^+(u) := B_1 \cap \partial\{u > 0\}$. For each $r > 0$ and $x^0 \in \mathbb{R}^d$, we denote by $u_{x^0, r}(x)$ and $u_r(x)$ the functions $u_{x^0, r}(x) := \frac{1}{r}u(x^0 + rx)$ and $u_r := \frac{1}{r}u(rx)$, respectively. Let $\{r_n\}_{n=1}^\infty$ be a vanishing sequence of positive numbers, and we denote by $u_n(x) := u_{x^0, r_n}(x)$ the *blow-up sequence* for every $x \in B_R(x^0)$ and $R > 0$. We say that a function $u_0: \mathbb{R}^d \rightarrow \mathbb{R}$ is a *blow-up limit* of u at $x^0 \in \Omega$ if for every $r_n \rightarrow 0$ as $n \rightarrow \infty$ and for every $R > 0$, u_n converges uniformly to u_0 in B_R , that is, $\lim_{n \rightarrow \infty} \|u_n - u_0\|_{L^\infty(B_R)} = 0$. We say that $x^0 \in \partial\Omega^+(u)$ is a *regular point* if there exists a blow-up limit u_0 of u at x^0 which has the form

$$u_0(x) = Q(x^0)(x \cdot \nu)^+, \quad \text{for every } x \in \mathbb{R}^d, \quad \text{and some } \nu \in \mathbb{R}^d.$$

The set of regular points will be denoted by $\partial_{\text{reg}}\Omega^+(u)$ and we define the *singular part* of the free boundary as the complementary of $\partial_{\text{reg}}\Omega^+(u)$

$$\partial_{\text{sing}}\Omega^+(u) := \partial\Omega^+(u) \setminus \partial_{\text{reg}}\Omega^+(u).$$

In the majority of this article, we work with the so-called *minimizers*.

Definition 1.1 (Minimizers). *We say that $u \in \mathcal{K}$ is a minimizer of J in Ω , provided that*

$$J(u) \leq J(v) \quad \text{for all } v \in \mathcal{K}. \quad (1.13)$$

In some contexts, u is called an absolute minimizer of J in Ω if (1.13) is satisfied.

Remark 1.1. *Most of our results in this paper can be generalized also for local minimizers, for which (1.13) is satisfied for $v \in u + H_0^1(D)$ and for any $D \subset\subset \Omega$.*

The notion of viscosity solutions was first systematically introduced by Lions and his cooperators in [8], who established an integrated theory on the existence and uniqueness theory for fully-nonlinear partial differential equations. The viscosity approach to the associated free boundary

problem was later developed by Caffarelli. We now define the notion of viscosity solutions to our problem and we first introduce the concept of "touch".

Definition 1.2 (Touch). *Given $u, \phi \in C^0(\Omega)$, we say that ϕ touches u from below (above) at $x^0 \in \Omega$ if $u(x^0) = \phi(x^0)$ and $u(x) \geq \phi(x)$ ($u(x) \leq \phi(x)$) in a small neighborhood of x^0 .*

With the aid of "touch", we formally define viscosity solutions of problem (1.7).

Definition 1.3 (Viscosity solutions). *Let $u \in C^0(\Omega)$ be a non-negative function in Ω . Then u is a solution to the problem (1.7) in the sense of viscosity, provided that for every $x^0 \in \overline{\Omega^+(u)}$, the following does hold.*

- (1) If $x^0 \in \Omega^+(u)$ and
 - if $\phi \in C^2(\Omega^+(u))$ touches u from below at x^0 , then $\Delta\phi(x^0) + f(\phi(x^0)) \leq 0$,
 - if $\phi \in C^2(\Omega^+(u))$ touches u from above at x^0 , then $\Delta\phi(x^0) + f(\phi(x^0)) \geq 0$.
- (2) If $x^0 \in \partial\Omega^+(u)$ and
 - if $\phi \in C^2(\Omega)$ touches u from below at x^0 , then $|\nabla\phi(x^0)| \leq Q(x^0)$,
 - if $\phi \in C^2(\Omega)$ touches u from above at x^0 , then $|\nabla\phi(x^0)| \geq Q(x^0)$.

1.2. Main results and plan of the paper. The main results in this paper state that the free boundary of the problem (1.7) can be decomposed into a disjoint union of a regular set and a singular set (presented in Theorem 1.2), the regular set of the free boundary is the graph of a $C^{1,\gamma}$ function locally, while for the singular set, there exists a critical dimension $d^* \in \{5, 6, 7\}$ so that the singularities can only occur when $d > d^*$ (presented in Theorem 1.3).

In proving these two results we need an array of technique tools including a Weiss-type monotonicity formula (presented in Proposition 5.3), optimal linear growth of solutions (presented in Proposition 2.5) and an optimal linear non-degeneracy property (presented in Proposition 3.2). In addition, due to the Lipschitz continuity of minimizers (presented in Proposition 3.1), a blow-up analysis (presented in Proposition 4.5) was derived for every free boundary point.

The monotonicity formula for a homogeneous one-phase Bernoulli-type problem (1.10) was first obtained in [35]. There special cases were considered for $f(t) \equiv 0$. In this paper, we generalize it to more general settings. It should be noted that in Proposition 5.9 we establish a monotonicity formula relating the rescaling $u_{x^0,r}$ with its one-homogeneous extension $z_{x^0,r}$ (See (5.81) for the precise definition).

Theorem 1.2 (Structure of the free boundary). *Let J and \mathcal{K} be given as in (1.4) and (1.5), respectively. Then there is a solution to the minimization problem $\min_{v \in \mathcal{K}} J(v)$. Every solution u is a Lipschitz viscosity solution to the problem (1.7) in the sense of Definition 1.3.*

The set $\Omega^+(u)$ has locally finite perimeter in Ω and the free boundary $\partial\Omega^+(u)$ can be decomposed into a disjoint union of a regular set $\partial_{\text{reg}}\Omega^+(u)$ and a singular set $\partial_{\text{sing}}\Omega^+(u)$. Namely,

$$\partial\Omega^+(u) = \bigcup_{\frac{1}{2} \leq \vartheta < 1} (\Omega^+(u))^{(\vartheta)},$$

where $(\Omega^+(u))^{(\vartheta)}$ is defined in the sense of (1.12). The regular set consists of points in which the Lebesgue density of the set $\Omega^+(u)$ is precisely $\frac{1}{2}$, and the singular set consists of points in which the Lebesgue density of $\Omega^+(u)$ is strictly between $\frac{1}{2}$ and 1.

$$\partial_{\text{reg}}\Omega^+(u) = (\Omega^+(u))^{(1/2)} \quad \text{and} \quad \partial_{\text{sing}}\Omega^+(u) = \bigcup_{\frac{1}{2} < \vartheta < 1} (\Omega^+(u))^{(\vartheta)}.$$

Moreover, for every $\vartheta \in [\frac{1}{2}, 1)$ and every blow-up limit u_0 , we have

$$(\Omega^+(u))^{(\vartheta)} = \left\{ x \in \partial\Omega^+(u) : \vartheta = \frac{|B_1 \cap \{u_0 > 0\}|}{|B_1|} \right\}.$$

Theorem 1.3 (Regularity of the regular set and the dimension for the singular set). *Let $u \in \mathcal{K}$ be a minimizer of J in Ω , then the regular part of the free boundary $\partial_{\text{reg}}\Omega^+(u)$ is an open subset of $\partial\Omega^+(u)$ and is locally a graph of a $C^{1,\gamma}$ function for $\gamma \in (0, \beta)$. The singular part of the free boundary $\partial_{\text{sing}}\Omega^+(u)$ is a closed subset of $\partial\Omega^+(u)$ and*

$$\mathcal{H}^{d-1}(\partial_{\text{sing}}\Omega^+(u)) = 0.$$

Moreover, there exists $d^* \in \{5, 6, 7\}$ such that

- If $d < d^*$, then $\partial_{\text{sing}}\Omega^+(u) = \emptyset$ and therefore all free boundary points are regular and $\partial\Omega^+(u) = \partial_{\text{reg}}\Omega^+(u)$.
- If $d = d^*$, then the singular set $\partial_{\text{sing}}\Omega^+(u)$ contains at most a finite number of isolated points,
- If $d > d^*$, then the Hausdorff dimension of $\partial_{\text{sing}}\Omega^+(u)$ is less than $d - d^*$.

2. PRELIMINARIES

In this section, we establish some fundamental facts about minimizers of J in \mathcal{K} , including the non-negativity and the governing equation (weak sense) for u in Ω . Note that since $u^0 \in H^1(\Omega)$, it is easy to check that $u^0 \in \mathcal{K}$ and thus $\mathcal{K} \neq \emptyset$. Therefore, the existence of minimizers can be obtained by a compactness argument as in [1, Lemma 3.1].

Proposition 2.1 (Hölder regularity). *Let $u \in \mathcal{K}$ be a minimizer of J in $B_r(x^0) \subset \Omega$, then there exists a constant C , so that*

$$|u(x) - u(y)| \leq C|x - y| \log \left(\frac{1}{|x - y|} \right) \quad \text{in } B_{r/2}(x^0).$$

In particular, $u \in C^{0,\alpha}(B_{r/2}(x^0))$ for some $\alpha \in (0, 1)$.

Proof. Assume without loss of generality that $x^0 = 0$. Let $v \in B_r$ be a function given by $\Delta v + f(v) = 0$ in B_r and $v = u$ on ∂B_r . The maximum principle gives that $v \leq \Psi$ in Ω and thus $v \in \mathcal{K}$. By the minimality of u in B_r , one has $J(u; B_r) \leq J(v; B_r)$ and this implies

$$\begin{aligned} 0 &\leq \int_{B_r} -|\nabla(v-u)|^2 + 2\nabla(v-u) \cdot \nabla v + (F(v) - F(u)) dx + \int_{B_r} Q^2(x)(\chi_{\{v>0\}} - \chi_{\{u>0\}}) dx \\ &\leq \int_{B_r} -|\nabla(v-u)|^2 dx + Q_{\max}^2 \int_{B_r} \chi_{\{v>0\}} dx, \end{aligned} \tag{2.14}$$

where we have used the assumption (1.9) in the second inequality. Therefore,

$$\int_{B_r} |\nabla(u-v)|^2 dx \leq Q_{\max}^2 |B_r \cap \{v > 0\}| \leq Q_{\max}^2 |B_r| \leq Cr^d,$$

where C depends only on d and Q_{\max} . Proceeding as in [6, Theorem 2.1] we complete the proof of the proposition. \square

Proposition 2.2 (Properties of minimizers). *Let $u \in \mathcal{K}$ be a minimizer of J in Ω , then $u \geq 0$ in Ω and u satisfies the equation*

$$\Delta u + f(u) \geq 0 \quad \text{in } \Omega, \tag{2.15}$$

in the weak sense. Moreover, u satisfies the equation

$$\Delta u + f(u) = 0 \quad \text{in } \Omega^+(u), \tag{2.16}$$

in the classical sense. That is, $u \in C^{2,\alpha}(D)$ satisfies (2.16) for any $D \subset\subset \Omega^+(u)$.

Proof. Choose $u_\varepsilon := u - \varepsilon \min\{u, 0\}$ for any $\varepsilon \in (0, 1)$. Then $u_\varepsilon \in \mathcal{K}$ is admissible. By the minimality of u , we have

$$\int_{\Omega} |\nabla \min\{u, 0\}|^2 dx \leq - \int_{\Omega} \frac{F'((1-\varepsilon)\min\{u, 0\})}{2} \min\{u, 0\} dx. \tag{2.17}$$

Since $F'(t) \leq 0$ for $t \leq 0$, we have from (2.17) that $\int_{\Omega} |\nabla \min\{u, 0\}|^2 dx \leq 0$ and thus $u \geq 0$ in Ω . Consider now the first variation $\lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} (J(u - \varepsilon \xi) - J(u)) \geq 0$ for any $\xi \geq 0$, $\xi \in C_0^\infty(\Omega)$, and we have that

$$0 \leq - \int_{\Omega} \nabla u \cdot \nabla \xi + \frac{F'(u)}{2} \xi dx,$$

which gives (2.15). By Proposition 2.2, u is a continuous function in Ω and this implies that $\Omega^+(u)$ is an open subset of Ω . For any $\xi \in C_0^\infty(\Omega^+(u))$, it is easy to check that the function $u - \varepsilon \xi \in \mathcal{K}$ and $\Omega^+(u - \varepsilon \xi) \subset \Omega^+(u)$ for $|\varepsilon| > 0$ small. Thus, by the minimality of u we have

$$0 = \int_{\Omega} \nabla u \cdot \nabla \xi + \frac{F'(u - \varepsilon \xi)}{2} \xi dx \quad \text{for any } \varepsilon. \tag{2.18}$$

Passing to the limit as $\varepsilon \rightarrow 0$ in (2.18) gives (2.16). It now follows from the Schauder interior estimate in [21] that $u \in C^{2,\alpha}(D)$ for any compact subset D of $\Omega^+(u)$. \square

We now introduce the following Lemma, which helps us to establish the optimal linear growth of minimizers up to the free boundary.

Lemma 2.3. *Let Ψ be given in (1.6), then for every non-negative function $u \in H^1(B_r(x^0))$ where $B_r(x^0) \subset \Omega$ the following estimates does hold:*

$$|B_r(x^0) \cap \{u = 0\}| \left(\frac{1}{r} \int_{\partial B_r(x^0)} u \, d\mathcal{H}^{d-1} - r \frac{F_0 M_0}{2} \right)^2 \leq C_d \int_{B_r(x^0)} |\nabla(u - h)|^2. \quad (2.19)$$

Here h is a function defined by $\Delta h + f(h) = 0$ in $B_r(x^0)$ with boundary value $h = u$ on $\partial B_r(x^0)$ and $M_0 := \sup_{\bar{\Omega}} \Psi$.

Remark 2.4. *An estimate of the type (2.19) was first brought by Alt and Caffarelli [Lemma 3.2, [1]], and we generalize the result of Lemma 3.2 in [1] to our semilinear case.*

Proof. Assume without loss of generality $x^0 = 0$. Let $v \in H^1(B_r)$ be the solution to the problem

$$\min \left\{ \int_{B_r} |\nabla v|^2 + F(v) \, dx : v - u \in H_0^1(B_r), \quad v \geq u \right\}.$$

Then v satisfies $\Delta v + f(v) \leq 0$ in B_r and $\Delta v + f(v) = 0$ in $B_r \cap \{v > u\}$. Now for each $|z| \leq \frac{1}{2}$ we consider a transformation $T: B_r \rightarrow B_r$ given by $T_z(x) := (r - |x|)z + x$ for each $x \in B_r$. Define

$$u_z(x) := u(T_z(x)) \quad \text{and} \quad v_z(x) := v(T_z(x)).$$

Let

$$S_z := \left\{ \xi \in \partial B_1 : u_z(\rho\xi) = 0 \quad \text{for some} \quad \rho \in \left[\frac{r}{8}, r \right] \right\}.$$

For $\xi \in S_z$, we define

$$r_\xi := \inf \left\{ \rho \in \left[\frac{r}{8}, r \right] : u_z(\rho\xi) = 0, \quad \xi \in S_z \right\}.$$

By a slicing argument for \mathcal{H}^1 a.e. $\xi \in S_z$ the function $\rho \mapsto \nabla u_z(\rho\xi)$ and $\rho \mapsto \nabla v_z(\rho\xi)$ are square integrable. Define a function $g(\rho) := u_z(\rho\xi) - v_z(\rho\xi)$ for $0 < \rho \leq r$, then $g(\rho)$ is absolutely continuous in $(0, r)$, and so, using also the fact that $g(r) = 0$ (since $u = v$ on ∂B_r) and $u_z(r_\xi\xi) = 0$, we have $v_z(r_\xi\xi) = g(r) - g(r_\xi) = \int_{r_\xi}^r \nabla(v_z(\rho\xi) - u_z(\rho\xi)) \cdot \xi \, d\rho$. Using Hölder inequality we have

$$v_z(r_\xi\xi) \leq \sqrt{r - r_\xi} \left(\int_{r_\xi}^r |\nabla(v_z - u_z)(\rho\xi)|^2 \, d\rho \right)^{1/2}. \quad (2.20)$$

Since $\Delta h = -f(h) \leq -f(0) + \frac{F_0 M_0}{2} \leq \frac{F_0 M_0}{2}$ for any $x \in B_r$, where we have used the fact $f(0) \geq 0$ in the last inequality. Since $h \geq 0$ on ∂B_r satisfying $\Delta h \leq M$ in B_r for $M := \frac{F_0 M_0}{2} \geq 0$, it follows from Lemma 2.4 (a) in [20] that

$$h(x) \geq r^d \frac{r - |x|}{(r + |x|)^{d-1}} \left(\frac{1}{r^2} \int_{\partial B_r} u \, d\mathcal{H}^{d-1} - \frac{2^{d-1} M}{d} \right) \geq C_d \frac{r - |x|}{r} \left(\int_{\partial B_r} u \, d\mathcal{H}^{d-1} - r^2 M \right).$$

Since v is a super-solution in B_r with the same boundary value of h , we have $v(x) \geq h(x)$ in B_r . Taking $x = (r - r_\xi)z + r_\xi\xi$ yields

$$v_z(r_\xi\xi) = v((r - r_\xi)z + r_\xi\xi) \geq C_d \frac{r - r_\xi}{r} \left(\int_{\partial B_r} u d\mathcal{H}^{d-1} - r^2 M \right). \quad (2.21)$$

Combined (2.21) with (2.20) together gives

$$(r - r_\xi) \left(\frac{1}{r} \int_{\partial B_r} u d\mathcal{H}^{d-1} - rM \right)^2 \leq C_d \int_{r_\xi}^r |\nabla(v_z - u_z)(\rho\xi)|^2 d\rho.$$

Integrating over $\xi \in S_z \subset \mathbb{S}^{d-1}$, we obtain the inequality

$$\left(\int_{S_z} (r - r_\xi) d\xi \right) \left(\frac{1}{r} \int_{\partial B_r} u d\mathcal{H}^{d-1} - rM \right)^2 \leq C_d \int_{\partial B_1} \int_{r_\xi}^r |\nabla(v_z - u_z)(\rho\xi)|^2 d\rho d\xi.$$

By the uniform estimate that $r/8 \leq r_\xi \leq r$, we have

$$\begin{aligned} |B_r \setminus B_{r/4}(rz) \cap \{u = 0\}| \left(\frac{1}{r} \int_{\partial B_r} u d\mathcal{H}^{d-1} - rM \right)^2 &\leq C_d \int_{B_r} |\nabla(v_z - u_z)|^2 dx \\ &\leq C_d \int_{B_r} |\nabla(v - u)|^2 dx, \end{aligned}$$

where we have used the fact that the Jacobian of T is bounded by a dimensional constant which is independent of r . The desired result (2.19) follows by integrating over z . \square

With Lemma 2.3 at hand, we immediately have

Proposition 2.5 (Optimal linear growth of minimizers). *Let $u \in \mathcal{K}$ be a minimizer of J in Ω , and let $x^0 \in \Omega$ be such that $|B_r(x^0) \cap \{u = 0\}| \neq 0$ for $B_r(x^0) \subset \Omega$, then*

$$\frac{1}{r} \int_{\partial B_r(x^0)} u d\mathcal{H}^{d-1} \leq \sqrt{C_d} Q_{\max} + r \frac{F_0 M_0}{2}, \quad (2.22)$$

where F_0 is given in (1.8) and $M_0 = \sup_{\bar{\Omega}} \Psi$.

Proof. Assume $x^0 = 0$ and let h be a function defined by $\Delta h + f(h) = 0$ in B_r and $h = u$ on ∂B_r . The maximum principle gives that $h \leq \Psi$ in B_r and thus $h \in \mathcal{K}$ is admissible. By the optimality $J(u) \leq J(h)$ and a similar calculation as in (2.14) gives

$$\int_{B_r} |\nabla(u - h)|^2 \leq Q_{\max}^2 |B_r \cap \{u = 0\}|. \quad (2.23)$$

Applying Lemma 2.3 and it follows from (2.19) and (2.23) that

$$|B_r \cap \{u = 0\}| \left(\frac{1}{r} \int_{\partial B_r} u d\mathcal{H}^{d-1} - r \frac{F_0 M_0}{2} \right)^2 \leq C_d Q_{\max}^2 |B_r \cap \{u = 0\}|.$$

By our assumption $|B_r \cap \{u = 0\}| \neq 0$ and we obtain (2.22). \square

Remark 2.6. *In view of Proposition 2.5 and if in particular $x^0 \in \partial\Omega^+(u)$ is a free boundary point, then choose $B_r(x^0)$ be a small ball with $r \leq \min\{\frac{2Q_{\max}}{F_0M_0}, 1\}$, we have*

$$\frac{1}{r} \int_{\partial B_r(x^0)} u \, d\mathcal{H}^{d-1} \leq C_d^* Q_{\max},$$

where $C_d^* := \sqrt{C_d} + 1$.

3. OPTIMAL REGULARITY AND NON-DEGENERACY

The first goal of this section is to prove that minimizers are locally Lipschitz functions in Ω . Precisely, we found that if a minimizer grows like its distance to the free boundary, then it is Lipschitz continuous near the free boundary. We summarize it as follows:

Proposition 3.1 (Lipschitz regularity of minimizers). *Let $u \in \mathcal{K}$ be a minimizer of J in Ω , then $u \in C^{0,1}(\Omega)$. Moreover, for every $D \subset\subset \Omega$ containing a free boundary point, one has*

$$\sup_D |\nabla u(x)| \leq C(d, F_0, D, \Omega)(Q_{\max} + F_0), \quad (3.24)$$

where F_0 is given in (1.8).

Proof. Suppose that $x^0 \in \Omega$ with $\rho(x^0) < \frac{1}{2} \min\{\text{dist}(x^0, \partial\Omega), 1\}$ where $\rho(x^0) := \text{dist}(x^0, \{u = 0\})$. Then we claim that

$$\frac{u(x^0)}{\rho(x^0)} \leq C(d, F_0)(Q_{\max} + F_0), \quad (3.25)$$

where F_0 is given in (1.8). Assume $\frac{u(x^0)}{\rho(x^0)} > M > 0$ for some $M > 0$, and we derive an upper bound for M . Consider the function $u_\rho(x) := \frac{u(x^0 + \rho(x - x^0))}{\rho}$ for $\rho := \rho(x^0)$ and $x \in B_1(x^0)$. Then u_ρ satisfies $\Delta u_\rho + \rho f(\rho u_\rho) = 0$ in $B_1(x^0)$. A direct calculation gives that

$$|\rho f(\rho u_\rho)| \leq \rho |f(0)| + \rho^2 \max_{t \geq 0} |f'(t)| u_\rho \leq \frac{\rho}{2} F_0 + \frac{\rho^2}{2} F_0 u_\rho. \quad (3.26)$$

Thanks to the Harnack inequality [30, Corollary 1.1], we have

$$\inf_{B_{3/4}(x^0)} u_\rho \geq cM - CF_0, \quad (3.27)$$

where c and C are constants depending on d and F_0 . Let $y \in \partial B_1(x^0) \cap \{u_\rho = 0\}$ be a point and v be a function defined by $\Delta v + \rho f(\rho v) = 0$ in $B_1(y)$ and $v = u_\rho$ on $\partial B_1(y)$. Since u_ρ is a minimizer, we have from $J(u_\rho; B_1(y)) \leq J(v; B_1(y))$, the convexity of F , and a similar calculation as in (2.14) that

$$\int_{B_1(y)} |\nabla(u_\rho - v)|^2 \, dx \leq Q_{\max}^2 \int_{B_1(y)} \chi_{\{u_\rho=0\}} \, dx. \quad (3.28)$$

It then follows from the maximum principle that $v \geq u_\rho$ in $B_1(y)$. This combined with (3.27) yields $v \geq u_\rho \geq cM - CF_0 > 0$ in $B_{3/4}(x^0) \cap B_1(y)$. Using Harnack inequality [30, Corollary 1.1] for the

function v once again gives

$$v(x) \geq cM - CF_0 \quad \text{in} \quad B_{1/2}(y). \quad (3.29)$$

Take $y = 0$ for simplicity and define

$$\varphi(x) := C_0(e^{-\beta|x|^2} - e^{-\beta}) \quad C_0 := cM - CF_0.$$

After a direct computation, $\Delta\varphi + \rho f(\rho\varphi) > 0$ for $\frac{1}{2} < |x| < 1$, provided that $\beta > 0$ is sufficiently large. This implies that $\Delta(\varphi - v) + \rho f(\rho\varphi) - \rho f(\rho v) > 0$ in $B_1 \setminus B_{1/2}$. It follows from the maximum principle that $v(x) \geq \varphi(x) = C_0(e^{-\beta|x|^2} - e^{-\beta}) \geq cC_0(1 - |x|)$ in $B_1 \setminus B_{1/2}$, this together with (3.29) gives

$$v(x) \geq (cM - CF_0)(1 - |x|) \quad \text{in} \quad B_1. \quad (3.30)$$

With the aid of (3.28) and (3.30), it follows from a similar argument as in [1, Lemma 3.2] and [2, Lemma 2.2] that $(cM - CF_0)^2 \leq CQ_{\max}^2$. This implies $M \leq C(Q_{\max} + F_0)$ and proves (3.25).

We now prove that $u \in C^{0,1}(\Omega)$. Let $\rho(x^0) < \frac{1}{3} \min\{\text{dist}(x^0, \partial\Omega), 1\}$ where $\rho(x^0) = \text{dist}(x^0, \{u = 0\})$, and let us define $\tilde{u}(\tilde{x}) := \frac{u(x^0 + \rho(x^0)\tilde{x})}{\rho(x^0)}$. Then it follows from (3.25) that $\tilde{u}(\tilde{x}) \leq C(Q_{\max} + F_0)$ in B_1 for C depends on d and F_0 . Since $\Delta\tilde{u} + \rho(x^0)f(\rho(x^0)\tilde{u})$, we have from (3.25), (3.26) and elliptic estimates in [21] that

$$|\nabla\tilde{u}(0)| \leq C(\|u\|_{C^0(B_1)} + \rho(x^0)\|f(\rho(x^0)u)\|_{L^\infty(B_1)}) \leq C(Q_{\max} + F_0),$$

where C depends only on d and F_0 . Therefore

$$|\nabla u(x^0)| = |\nabla\tilde{u}(0)| \leq C(Q_{\max} + F_0).$$

Thus, for any $D \subset\subset \Omega$, $|\nabla u(x)|$ is bounded in $D \cap \Omega^+(u) \cap N$, where N is a small neighborhood of the free boundary. Since $u \in C^{2,\alpha}(\Omega^+(u))$ and $\nabla u = 0$ a.e. in $\{u = 0\} \cap \Omega$, we have $u \in C^{0,1}(\Omega)$. The rest of the proof is standard and can be obtained by proceeding as in [Page 11, Theorem 2.3, [2]]. \square

The second goal of this section is to establish a non-degeneracy property for minimizers, which gives us a uniform L^∞ bound of minimizers up to the free boundary $\partial\Omega^+(u)$.

Proposition 3.2 (Non-degeneracy of minimizers). *For any $\kappa \in (0, 1)$, there exists a positive constant $c_d^* = c_d^*(\kappa)$, so that for any $B_r(x^0) \subset \Omega$ with $r < \frac{1}{F_0} \max\{c_d^*Q_{\min}, 1\}$, the following does hold:*

$$\frac{1}{r} \int_{\partial B_r(x^0)} u \, d\mathcal{H}^{d-1} \leq c_d^* Q_{\min} \quad (3.31)$$

implies that

$$u(x) \equiv 0 \quad \text{in} \quad B_{\kappa r}(x^0).$$

Proof. Assume $x^0 = 0$ and consider the function u_r for some $r > 0$ so that $\{x \in \Omega: rx\} \subset \Omega$. Then obviously, $\Delta u_r + rf(ru_r) = 0$ in $\Omega^+(u_r)$. Define

$$\ell_u := \sup_{B_{\sqrt{\kappa}}} u_r \quad \text{for some } \kappa \in (0, 1). \quad (3.32)$$

Since $\Delta u_r \geq -rf(ru_r) \geq -rf(0) \geq -\frac{rF_0}{2}$ in Ω , it follows from the maximum principle that

$$u_r(x) \leq \frac{rF_0}{2} \frac{1 - |x|^2}{2d} + \frac{1 - |x|^2}{d\omega_d} \int_{\partial B_1} \frac{u_r}{|x - y|^d} d\mathcal{H}^{d-1} \quad \text{for every } x \in B_{\sqrt{\kappa}}. \quad (3.33)$$

Taking supremum on the both sides of (3.33) and using (3.32), we have

$$\ell_u \leq \frac{rF_0}{2} + C_d(\kappa) \int_{\partial B_1} u_r d\mathcal{H}^{d-1}. \quad (3.34)$$

Let v be a function defined by $\Delta v + rf(rv) = 0$ in $B_{\sqrt{\kappa}} \setminus B_\kappa$, $v = 0$ in B_κ and $v = \ell_u$ outside $B_{\sqrt{\kappa}}$. Then the standard elliptic estimates yields

$$\sup_{\partial B_\kappa} |\nabla v| \leq C(\sup_{B_{\sqrt{\kappa}}} v + r\|f(rv)\|_{L^\infty(B_{\sqrt{\kappa}})}) \leq C(\ell_u + rF_0), \quad (3.35)$$

where C is a constant depending on κ and F_0 . Note that the function $w := \min\{u_r, v\}$ is an admissible function and so $J(u_r; B_{\sqrt{\kappa}}) \leq J(w; B_{\sqrt{\kappa}})$. On the other hand, $\int_{B_\kappa} |\nabla w|^2 + F(rw) + Q^2(rx)\chi_{\{w>0\}} dx = 0$, due to the fact that $v = 0$ in B_κ . Thus

$$\begin{aligned} J(u_r; B_\kappa) &= \int_{B_\kappa} |\nabla u_r|^2 + F(ru_r) + Q^2(rx)\chi_{\{u_r>0\}} dx \\ &\leq \int_{B_{\sqrt{\kappa}} \setminus B_\kappa} |\nabla w|^2 - |\nabla u_r|^2 + F(rw) - F(ru_r) + Q^2(rx) (\chi_{\{w>0\}} - \chi_{\{u_r>0\}}) dx \\ &\leq \int_{B_{\sqrt{\kappa}} \setminus B_\kappa} -|\nabla(w - u_r)|^2 + 2\nabla w \cdot \nabla(w - u_r) + rF'(rw)(w - u_r) dx \\ &\leq \int_{B_{\sqrt{\kappa}} \setminus B_\kappa} -|\nabla \min\{v - u_r, 0\}|^2 + 2\nabla v \cdot \nabla \min\{v - u_r, 0\} + rF'(rv) \min\{v - u_r, 0\} dx \\ &\leq 2 \int_{\partial B_\kappa} \min\{v - u_r, 0\} \frac{\partial v}{\partial \nu} d\mathcal{H}^{d-1}, \end{aligned} \quad (3.36)$$

where ν is the unit normal vector. Here we have used the fact that $\int_{B_{\sqrt{\kappa}} \setminus B_\kappa} \chi_{\{w>0\}} - \chi_{\{u_r>0\}} dx \leq 0$ in the third inequality since $\{w > 0\} \subset \{u_r > 0\}$. By the convexity of F , we have $F(ru_r) \geq F'(0)ru_r \geq -rF_0u_r$ and this gives

$$\int_{B_\kappa} |\nabla u_r|^2 + Q^2(rx)\chi_{\{u_r>0\}} dx \leq J(u_r; B_\kappa) + \int_{B_\kappa} rF_0u_r dx. \quad (3.37)$$

It follows from (3.34), (3.35), (3.36), (3.37) and the trace theorem that

$$\begin{aligned}
& \int_{B_\kappa} |\nabla u_r|^2 + Q^2(rx)\chi_{\{u_r>0\}} dx \\
& \leq 2 \int_{\partial B_\kappa} \min\{v - u_r, 0\} \frac{\partial v}{\partial \nu} d\mathcal{H}^{d-1} + \int_{B_\kappa} rF_0 u_r dx \\
& \leq C(\ell_u + rF_0) \left(\int_{\partial B_\kappa} u_r d\mathcal{H}^{d-1} + \int_{B_\kappa} u_r dx \right) \\
& \leq C(\ell_u + rF_0) \left(\int_{B_\kappa} u_r \chi_{\{u_r>0\}} dx + C \int_{B_\kappa} |\nabla u_r| \chi_{\{u_r>0\}} dx \right) \\
& \leq C(\ell_u + rF_0) \left(\frac{\ell_u + rF_0}{Q_{\min}^2} \int_{B_\kappa} \chi_{\{u_r>0\}} dx + \frac{C}{Q_{\min}} \int_{B_\kappa} |\nabla u_r|^2 + Q^2(rx)\chi_{\{u_r>0\}} dx \right) \\
& \leq \frac{C(\ell_u + rF_0)}{Q_{\min}} \left[\frac{(\ell_u + rF_0)}{Q_{\min}} + 1 \right] \int_{B_\kappa} |\nabla u_r|^2 + Q^2(rx)\chi_{\{u_r>0\}} dx,
\end{aligned} \tag{3.38}$$

where C is a constant depending on d , F_0 and κ . Thanks to (3.31) and (3.34), we have

$$\frac{\ell_u + rF_0}{Q_{\min}} \leq C \left(\frac{1}{Q_{\min}} \int_{\partial B_1} u_r d\mathcal{H}^{d-1} + \frac{rF_0}{Q_{\min}} \right) \leq Cc_d^*.$$

This implies that $\frac{\ell_u + rF_0}{Q_{\min}}$ is sufficiently small, provided that c_d^* is sufficiently small. This together with the last inequality in (3.38) implies that

$$\int_{B_\kappa} |\nabla u_r|^2 + Q^2(rx)\chi_{\{u_r>0\}} dx = 0,$$

for c_d^* small enough. Thus $u_r = 0$ in B_κ , provided that c_d^* is small enough. This concludes the proof. \square

Remark 3.3. *Proposition 3.2 remains true if $B_r(x^0)$ is not contained in Ω , provided that $u = 0$ in $B_r(x^0) \cap \partial\Omega$.*

The following corollaries are direct applications of Proposition 3.2 and Remark 2.6.

Corollary 3.4. *Let $u \in \mathcal{K}$ be a minimizer of J in Ω , then for any $D \subset\subset \Omega$, there are constants $0 \leq c_d^* \leq C_d^*$ so that for every $B_r(x^0) \subset D$ with $x^0 \in \partial\Omega^+(u)$*

$$c_d^* Q_{\min} \leq \frac{1}{r} \int_{\partial B_r(x^0)} u d\mathcal{H}^{d-1} \leq C_d^* Q_{\max}.$$

Corollary 3.5 (optimal linear nondegeneracy). *Let $u \in \mathcal{K}$ be a local minimizer of J in $B_r(x^0) \subset \Omega$ with $x^0 \in \partial\Omega^+(u)$. Then*

$$\sup_{B_r(x^0)} |u| \geq rc_d^* Q_{\min}.$$

In the end of this section, we prove a density estimate that will be used in our forthcoming study.

Proposition 3.6 (Density estimates). *Let $D \subset\subset \Omega$, then there exists a positive constant $c \in (0, 1)$, such that for any $B_r(x^0) \subset D$, centered at $x^0 \in \partial\Omega^+(u)$ with $r > 0$ small,*

$$c < \frac{|B_r(x^0) \cap \{u > 0\}|}{|B_r(x^0)|} < 1 - c. \quad (3.39)$$

Proof. Without loss of generality, we assume that $x^0 = 0$, and we first prove the estimate by below. It follows from Corollary 3.5 that there exists a point $y \in \partial B_r$ such that $u(y) > \eta r$ for some constant $\eta > 0$. Define a function $v(x) = u(x) + \frac{F_0|x|^2}{4d}$. A direct calculation gives that $\Delta v = \Delta u + \frac{F_0}{2} \geq f(0) - f(u) \geq 0$ in B_r . This implies that for $\kappa > 0$ small,

$$\frac{F_0 r}{2d\kappa} + \frac{1}{\kappa r} \int_{\partial B_{\kappa r}(y)} u \, d\mathcal{H}^{d-1} \geq \frac{1}{\kappa r} \int_{\partial B_{\kappa r}(y)} v \, d\mathcal{H}^{d-1} \geq \frac{v(y)}{\kappa r} \geq \frac{\eta}{\kappa},$$

for sufficiently small $\kappa > 0$. It follows that

$$\frac{1}{\kappa r} \int_{\partial B_{\kappa r}(y)} u \, d\mathcal{H}^{d-1} \geq \frac{\eta}{\kappa} - \frac{F_0 r}{2d\kappa} > \frac{\eta}{4\kappa},$$

provided that $r > 0$ is sufficiently small. By Proposition 3.2, we have that $u > 0$ in $B_{\kappa r}(y)$, which gives the lower bound.

Consider a function v defined in \bar{B}_r by $\Delta v + f(v) = 0$ in B_r and $v = u$ on ∂B_r . The maximum principle implies that $u \leq v \leq \Psi$ in B_r where Ψ is defined in (1.6). Therefore $v \in \mathcal{X}$ is admissible. The minimality of u and a similar calculation as in (2.14) gives $\int_{B_r} |\nabla(u-v)|^2 dx \leq Q_{\max}^2 |B_r \cap \{u = 0\}|$. Thanks to Poincaré and Hölder inequality, one has

$$Q_{\max}^2 |B_r \cap \{u = 0\}| \geq \int_{B_r} |\nabla(v-u)|^2 dx \geq \frac{c}{r^2} \int_{B_r} |v-u|^2 dx. \quad (3.40)$$

For any $y \in B_{\kappa r}$, since $\Delta v \leq \frac{M_0 F_0}{2} - f(0)$ with $M_0 = \sup_{\bar{\Omega}} \Psi$, a direct calculation gives

$$\begin{aligned} v(y) &\geq \frac{r^2 - |y|^2}{d\omega_d r} \int_{\partial B_r} \frac{u}{|x-y|^d} \, d\mathcal{H}^{d-1} + (M_0 F_0 - f(0)) \frac{|y|^2 - r^2}{2d} \\ &\geq (1 - C\kappa) \int_{\partial B_r} u \, d\mathcal{H}^{d-1} - Cr^2. \end{aligned} \quad (3.41)$$

It follows from (3.41), Proposition 3.1 Proposition 3.2 that

$$\begin{aligned} v(y) - u(y) &\geq v(u) - C\kappa r \\ &\geq (1 - C\kappa) \int_{\partial B_r} u \, d\mathcal{H}^{d-1} - C\kappa r - Cr^2 \\ &\geq cQ_{\min} r, \end{aligned} \quad (3.42)$$

provided that κ and r are sufficiently small.

Combining with (3.40) and (3.42), one has

$$Q_{\max}^2 |B_r \cap \{u = 0\}| \geq \frac{c}{r^2} \int_{B_r} |v-u|^2 dx \geq \frac{c}{r^2} \int_{B_{\kappa r}} |v-u|^2 dx \geq c(\kappa) Q_{\min}^2 r^d, \quad (3.43)$$

for $r > 0$ sufficiently small. Thus, in view of (3.43), we have

$$\frac{|B_r \cap \{u = 0\}|}{|B_r|} \geq c \frac{Q_{\min}}{Q_{\max}},$$

where c is independent of r . This gives the upper bound in (3.39) since $\frac{Q_{\min}}{Q_{\max}} > 0$. \square

4. LOCALLY FINITE PERIMETER OF THE POSITIVITY SET AND BLOW-UP LIMITS

In this section we prove that the set $\Omega^+(u)$ has (locally) finite perimeter in Ω . The first step is a measure estimate of the free boundary. To begin with, let us define

$$\lambda := \Delta u + f(u), \quad (4.44)$$

and

$$\lambda_0 := \Delta u + f(u)\chi_{\{u > 0\}}. \quad (4.45)$$

Then we immediately have

Lemma 4.1. *Let λ and λ_0 be defined as in (4.44) and (4.45), respectively, then λ is a Radon measure supported on $\Omega \cap \{u = 0\}$ and its singular point is contained in the free boundary. λ_0 is a positive Radon measure supported in $\partial\Omega^+(u)$.*

Proof. For any $\xi \in C_0^\infty(\Omega)$ with $\xi \geq 0$, define $\eta(t) := \max\{\min\{2 - t, 1\}, 0\}$ for any $t > 0$. Then

$$\begin{aligned} \int_{\Omega} \nabla u \cdot \nabla(\xi\eta(tu)) \, dx &= \int_{\Omega} \nabla u \cdot \nabla \xi \, dx + \int_{\Omega \cap \{u \geq \frac{1}{t}\}} \nabla u \cdot \nabla(\xi\eta(tu) - \xi) \, dx \\ &= \int_{\Omega} \nabla u \cdot \nabla \xi \, dx - \int_{\Omega^+(u)} f(u)\xi \, dx \\ &\quad + \int_{\Omega \cap \{\frac{1}{t} \leq u \leq \frac{2}{t}\}} (2 - ut)f(u)\xi \, dx + \int_{\Omega \cap \{u \leq \frac{2}{t}\}} f(u)\xi \, dx \\ &\geq \int_{\Omega} \nabla u \cdot \nabla \xi \, dx - \int_{\Omega^+(u)} f(u)\xi \, dx, \end{aligned} \quad (4.46)$$

for $t > 0$ large enough, where we have used the fact $f(u) \geq \frac{f(0)}{2} \geq 0$ in $\Omega \cap \{u \leq \frac{2}{t}\}$ for $t > 0$ large enough. On the other hand, one has

$$\begin{aligned} \int_{\Omega} \nabla u \cdot \nabla(\xi\eta(tu)) \, dx &= \int_{\Omega \cap \{\frac{1}{t} \leq u \leq \frac{2}{t}\}} (2 - ut)\nabla u \cdot \nabla \xi - \xi t |\nabla u|^2 \, dx + \int_{\Omega \cap \{0 < u \leq \frac{1}{t}\}} \nabla u \cdot \nabla \xi \, dx \\ &\leq \int_{\Omega \cap \{0 < u \leq \frac{2}{t}\}} |\nabla u| |\nabla \xi| \, dx. \end{aligned} \quad (4.47)$$

It follows from (4.46) and (4.47) that

$$\int_{\Omega} \nabla u \cdot \nabla \xi \, dx - \int_{\Omega^+(u)} f(u)\xi \, dx \leq \int_{\Omega \cap \{0 < u \leq \frac{2}{t}\}} |\nabla u| |\nabla \xi| \, dx. \quad (4.48)$$

Passing to the limit as $n \rightarrow \infty$ in (4.48), we conclude that $\Delta u + f(u)\chi_{\{u>0\}} \geq 0$ in the sense of distributions. Consequently, there exists a positive Radon measure λ_0 supported on $\partial\Omega^+(u)$ so that $\lambda_0 = \Delta u + f(u)\chi_{\{u>0\}}$. Recalling (1.10) that $f(0) \geq 0$ there exists a positive Radon measure λ supported on $\Omega \cap \{u = 0\}$ so that $\lambda = \lambda_0 + f(0)\chi_{\{u=0\}} = \Delta u + f(u)$. \square

Compare to [1, Theorem 4.3], we have the following result adjusted to our semilinear settings.

Proposition 4.2. *Let λ and λ_0 be defined as in (4.44) and (4.45), respectively, and let $D \subset\subset \Omega$. Then there exist positive constants c and C , depending on d , F_0 and the Lipschitz coefficient of u in $B_r(x^0)$, so that for any $B_r(x^0) \subset G$ with $x^0 \in \partial\Omega^+(u)$ and $r < r_0$,*

$$cr^{d-1} \leq \int_{B_r(x^0)} d\lambda \leq Cr^{d-1}. \quad (4.49)$$

Moreover,

$$cr^{d-1} \leq \int_{B_r(x^0) \cap \partial\{u>0\}} d\lambda = \int_{B_r(x^0)} d\lambda_0 \leq Cr^{d-1}. \quad (4.50)$$

Proof. For any $\varepsilon > 0$, introduce the function $d_{\varepsilon, B_r(x^0)} = \min\{\frac{\text{dist}(x, \mathbb{R}^d \setminus B_r(x^0))}{\varepsilon}, 1\}$ as a test function for $B_r(x^0) \subset D$. Then

$$\int_{\Omega} d_{\varepsilon, B_r(x^0)} d\lambda = \int_{\Omega} \nabla u \cdot \nabla d_{\varepsilon, B_r(x^0)} + f(u)d_{\varepsilon, B_r(x^0)} dx. \quad (4.51)$$

Due to the fact that $d_{\varepsilon, B_r(x^0)}$ converges to $\chi_{B_r(x^0)}$ as $\varepsilon \rightarrow 0$, we take the limit $\varepsilon \rightarrow 0$ in (4.51) and obtain

$$\int_{B_r(x^0)} d\lambda = \int_{\partial B_r(x^0)} \nabla u \cdot \nu d\mathcal{H}^{d-1} + \int_{B_r(x^0)} f(u) dx \leq Cr^{d-1} + CF_0r^d \leq Cr^{d-1}, \quad (4.52)$$

where the constant C depends on d , F_0 and the Lipschitz constant of u in $B_r(x^0)$. This gives the right hand side of (4.49).

Let $y \in B_r(x^0)$ and $G_y(x)$ be the Green function for Laplacian in $B_r(x^0)$ with pole y . If $u(y) > 0$, then y is outside the support of the Radon measure λ , and this implies

$$\int_{B_r(x^0)} G_y(x) d\lambda = -u(y) - \int_{B_r(x^0)} f(u)G_y(x) dx + \int_{\partial B_r(x^0)} u\partial_{-\nu}G_y(x) d\mathcal{H}^{d-1}. \quad (4.53)$$

With the help of Corollary 3.5 there exists a point $y \in \partial B_{\kappa r}$ so that $u(y) \geq c_d^*(\kappa)Q_{\min}r > 0$ for $\kappa > 0$ small. On the other hand, since u is Lipschitz continuous, we have $u(y) = u(y) - u(x^0) \leq C\kappa r$ and $u > 0$ in $B_{c(\kappa)r}(y)$ for small constants $c(\kappa)$. This combined with (4.53) gives

$$\int_{B_r(x^0)} G_y(x) d\lambda \geq -C\kappa r - CF_0r^d + c(1 - \kappa) \int_{\partial B_r(x^0)} u d\mathcal{H}^{d-1} \geq c_0r, \quad (c_0 > 0), \quad (4.54)$$

provided that κ and r small enough. Note that we have

$$\int_{B_r(x^0)} G_y(x) d\lambda = \int_{B_r(x^0) \cap \{u=0\}} G_y(x) d\lambda \leq \sup_{B_r(x^0) \cap \{u=0\}} G_y(x) \int_{B_r(x^0)} d\lambda \leq C(\kappa)r^{2-d} \int_{B_r(x^0)} d\lambda. \quad (4.55)$$

Here we used the fact that $\text{dist}(y, B_r(x^0) \cap \{u = 0\}) \geq c(\kappa)r$. Combining (4.54) and (4.55) together gives the left hand side of (4.49).

We now prove (4.50). Firstly, the right-hand side of (4.50) is obvious since $\int_{\partial B_{r,x^0}^+(u)} d\lambda \leq \int_{B_r(x^0)} d\lambda$. Moreover,

$$\begin{aligned} \int_{B_r(x^0)} d\lambda &= \int_{B_r(x^0) \cap \partial\{u>0\}} d\lambda + \int_{B_r(x^0) \cap \{u=0\}} d\lambda \\ &\leq \int_{B_r(x^0) \cap \partial\{u>0\}} d\lambda + \int_{B_r(x^0) \cap \{u=0\}} f(0) dx \\ &\leq \int_{B_r(x^0) \cap \partial\{u>0\}} d\lambda + C_d F_0 r^d, \end{aligned}$$

which implies that

$$\int_{B_r(x^0) \cap \partial\{u>0\}} d\lambda \geq \int_{B_r(x^0)} d\lambda - C_d F_0 r^d \geq cr^{d-1} - C_d F_0 r^d \geq cr^{d-1},$$

for sufficiently small r , which gives the left-hand side of (4.50). \square

The next theorem easily follows easily from Proposition 4.2, precisely as in [1].

Theorem 4.3 (Representation theorem). *Let u be a minimizer. Then:*

- (1) $\mathcal{H}^{d-1}(D \cap \partial\{u > 0\}) < +\infty$ for every $D \subset\subset \Omega$.
- (2) There exists a Borel function q_u , such that

$$\Delta u + f(u)\chi_{\{u>0\}} = q_u \mathcal{H}^{d-1} \llcorner \partial\{u > 0\}.$$

That is, for any $\xi \in C_0^\infty(\Omega)$,

$$-\int_{\Omega} \nabla u \cdot \nabla \xi dx + \int_{\Omega^+(u)} f(u)\xi dx = \int_{\partial\Omega^+(u)} q_u \xi d\mathcal{H}^{d-1}.$$

- (3) For any compact subset D of Ω , and for any $B_r(x^0) \subset D$ with $x^0 \in \partial\Omega^+(u)$, there exists $0 < c \leq C < \infty$, independent of r and x^0 , so that

$$0 < c \leq q_u(x) \leq C < +\infty \quad \text{and} \quad cr^{d-1} \leq \mathcal{H}^{d-1}(B_r(x^0) \cap \partial\{u > 0\}) \leq Cr^{d-1}.$$

As a direct corollary of Theorem 4.3 and [18, Theorem 1, Section 5.11], we have

Corollary 4.4 (Finite perimeter of $\Omega^+(u)$). *Let $u \in \mathcal{K}$ be a minimizer of J , then the set $\Omega^+(u)$ has locally finite perimeter in Ω .*

In the rest of this section, we shall deal with the blow-up sequences

$$u_n(x) = \frac{u(x^0 + r_n x)}{r_n} \quad \text{for } x^0 \in \partial\Omega^+(u).$$

Since $|\nabla u_n| \leq C$ and $u_n(0) = 0$, we have by a diagonal argument and the theorem of Ascoli-Arzelà that there exists a non-relabel subsequence and a function $u_0 \in W_{\text{loc}}^{1,\infty}(\mathbb{R}^d)$, so that

$$u_n \rightarrow u_0 \quad \text{in} \quad C_{\text{loc}}^\alpha(\mathbb{R}^d) \quad \text{for some} \quad \alpha \in (0, 1), \quad (4.56)$$

and

$$\nabla u_n \xrightarrow{*} \nabla u_0 \quad \text{in} \quad L_{\text{loc}}^\infty(\mathbb{R}^d). \quad (4.57)$$

The function u_0 is usually called a *blow-up limit*.

Proposition 4.5 (Properties of blow-up limits). *Let $\{u_n\}$ be the blow-up sequence and let u_0 be given as in (4.56) and (4.57), respectively. Then*

- (1) $\partial\Omega^+(u_n) \rightarrow \partial\Omega^+(u_0)$ locally in the Hausdorff distance of \mathbb{R}^d .
- (2) $\chi_{\{u_n > 0\}} \rightarrow \chi_{\{u_0 > 0\}}$ in $L_{\text{loc}}^1(\mathbb{R}^d)$.
- (3) $\nabla u_n \rightarrow \nabla u_0$ a.e. in \mathbb{R}^d .
- (4) $u_n \rightarrow u_0$ strongly in $H_{\text{loc}}^1(\mathbb{R}^d)$.
- (5) u_0 is a global minimizer of the functional $J_0(v)$ in B_r for any $r > 0$. Namely,

$$J_0(u_0) = \min_v J_0(v) \quad \text{for any } v \in H^1(B_r) \text{ with } v = u_0 \text{ on } \partial B_r, \quad (4.58)$$

where $J_0(v) := \int_{B_r} |\nabla v|^2 + Q(x^0)\chi_{\{v > 0\}} dx$.

- (6) u_0 satisfies the problem $\Delta u_0 = 0$ in $B_1^+(u_0)$ and $|\nabla u_0| = Q(x^0)$ on $B_1 \cap \partial\{u_0 > 0\}$ and the non-degeneracy property: there exists some constants $0 < c_0 < 1$ so that

$$c_0 < \frac{|B_1 \cap \{u_0 > 0\}|}{|B_1|} < 1 - c_0. \quad (4.59)$$

Proof. The argument (1)-(3) are standard. We only remark that these three properties strongly rely on Corollary 3.4 in our semilinear case. For the fourth property, it suffices to show that there exists a non-relabel subsequence $\{u_n\}$ such that $\nabla u_n \rightarrow \nabla u_0$ strongly in $L_{\text{loc}}^2(\mathbb{R}^d)$, namely, for any $R > 0$ the inequality

$$\limsup_{n \rightarrow \infty} \int_{B_R} |\nabla u_n|^2 \xi dx \leq \int_{B_R} |\nabla u_0|^2 \xi dx \quad \text{for every } \xi \in C_0^\infty(B_R).$$

For each n , u_n is a solution to the problem

$$\Delta u_n + r_n f(r_n u_n) = 0 \quad \text{in} \quad \Omega^+(u_n), \quad |\nabla u_n| = Q(x^0 + r_n x) \quad \text{on} \quad \partial\Omega^+(u_n). \quad (4.60)$$

Since u_n converges to u_0 locally uniformly, it follows from (4.60) that u_0 is harmonic in $\{u_0 > 0\}$. Also, using the uniform convergence, the continuity of u_0 and its harmonicity in $\{u_0 > 0\}$ we obtain

$$\begin{aligned} \int_{\mathbb{R}^d} |\nabla u_n|^2 \xi dx &= - \int_{\mathbb{R}^d} u_n (\nabla u_n \cdot \nabla \xi - r_n f(r_n u_n) \xi) dx \\ &\rightarrow - \int_{\mathbb{R}^d} u_0 \nabla u_0 \cdot \nabla \xi dx = \int_{\mathbb{R}^d} |\nabla u_0|^2 \xi dx \quad \text{as } n \rightarrow \infty. \end{aligned}$$

It therefore follows that u_n converges to u_0 strongly in $W_{\text{loc}}^{1,2}(\mathbb{R}^d)$ as $n \rightarrow \infty$.

We now prove (5). Consider a function $\eta_\varepsilon(x) := \min\{\frac{\text{dist}(x, \mathbb{R}^d \setminus B_r)}{\varepsilon}, 1\}$, it is easy to see that $\eta_\varepsilon \in C_0^{0,1}(B_r)$ and $0 \leq \eta_\varepsilon \leq 1$. Define $v_n := v + (1 - \eta_\varepsilon)(u_n - u_0)$ and it follows immediately that $v_n = u_n$ outside B_r and $\{v_n > 0\} \subset \{v > 0\} \cap \{\eta_\varepsilon < 1\}$. Thus,

$$\begin{aligned} & \int_{B_r} |\nabla u_n|^2 + F(r_n u_n) + Q^2(x^0 + r_n x) \chi_{\{u_n > 0\}} dx \\ & \leq \int_{B_r} |\nabla v_n|^2 + F(r_n v_n) + Q^2(x^0 + r_n x) \chi_{\{v_n > 0\}} dx \\ & \leq \int_{B_r} |\nabla v_n|^2 + F(r_n v_n) + Q^2(x^0 + r_n x) \chi_{\{v_n > 0\}} dx + \int_{B_r} Q^2(x^0 + r_n x) \chi_{\{\eta_\varepsilon < 1\}} dx. \end{aligned}$$

Passing to the limit as $n \rightarrow \infty$ gives

$$\int_{B_r} |\nabla u_0|^2 + Q^2(x^0) \chi_{\{u_0 > 0\}} dx \leq \int_{B_r} |\nabla v|^2 + Q^2(x^0) \chi_{\{v > 0\}} + Q^2(x^0) \int_{B_r} \chi_{\{\eta_\varepsilon < 1\}} dx, \quad (4.61)$$

where we have used the statements (2) and (3). Taking $\varepsilon \rightarrow 0$ in (4.61) gives the desired result. The fact that $\Delta u_0 = 0$ follows from the uniform convergence of u_n to u_0 and the first equation in (4.60). The second equation in (4.60) follows from the fact that u_0 is a global minimizer of $J_0(v)$. Finally for the last property, we point out it directly follows from (5) and [1, Lemma 3.7]. \square

Remark 4.6. *In conclusion, we compare the classical homogeneous problems with our semilinear non-homogeneous problems as follows:*

TABLE 1. Homogeneous v.s. Nonhomogeneous

<i>Problem</i>	<i>Equations for Blow-up sequences</i>	<i>Equations for Blow-ups</i>
<i>Problem (1.7)</i>	$\Delta u_n + r_n f(u_n) = 0$	$\Delta u_0 = 0$
<i>Problem (1.2)</i>	$\Delta u_n = 0$	$\Delta u_0 = 0$

One of the significant differences we read from the second column of Table 1 between the two problems is that the blow-up sequence satisfies different equations. However, in view of the last column of Table 1, we see that the blow-up limits of the two problems are both harmonic functions. We stress that this is the main reason we can establish the regularity of the regular set $\partial_{\text{reg}} \Omega^+(u)$ and determine the dimension of the singular set. We will discuss these in the coming four sections precisely.

5. WEISS-TYPE MONOTONICITY FORMULAS

In this section, we establish a Weiss-type monotonicity formula, which will be a useful tool in the study of homogeneity of blow-ups u_0 . The Weiss boundary-adjusted energy developed by Weiss in [35] originally considered the problem (1.1), who proved that blow-up limits of (1.1) are homogeneous functions of degree 1. We borrow this idea and adapt it to our settings. However, the strategy here is slightly different, since our functional J involves a non-linear term F . Our final results reveal that the functional J still possesses monotonicity after subtracting the boundary-adjusted term. Before establishing that, we need to introduce some notation. For every $B_r(x^0) \subset \Omega$

and $u \in H^1(B_r(x^0))$, we define

$$W(u, r; Q) = \frac{1}{r^d} \int_{B_r(x^0)} (|\nabla u|^2 + F(u) + Q^2(x)\chi_{\{u>0\}}) dx - \frac{1}{r^{d+1}} \int_{\partial B_r(x^0)} u^2 d\mathcal{H}^{d-1}, \quad (5.62)$$

and

$$\begin{aligned} W(u_{x^0, r}) &:= W(u_{x^0, r}, 1, Q(x^0 + rx)) \\ &= \int_{B_1} |\nabla u_{x^0, r}|^2 + F(ru_{x^0, r}) + Q^2(x^0 + rx)\chi_{\{u_{x^0, r}>0\}} dx - \int_{\partial B_1} u_{x^0, r}^2 d\mathcal{H}^{d-1}. \end{aligned} \quad (5.63)$$

Remark 5.1. *It is easy to verify the following fact: If $u \in H^1(B_1)$ is a minimizer of J for all $v \in H^1(B_1)$ with $v = u$ on ∂B_1 . Then u is also a minimizer of $W = W(\cdot, r; Q)$ in the sense that $W(u) \leq W(v)$ for every $v \in H^1(B_1)$ with $v = u$ on ∂B_1 .*

Remark 5.2. *Notice that the first integral in (5.62) is exactly our standard energy $J(u; B_r)$, while the second integral in (5.62) is the noted Weiss boundary-adjusted energy. We pause here for a second to display our pattern for constructing the formula (5.62). Firstly, we focus on the integral involving the gradient of u and the non-linear term F . The term $|\nabla u|^2 + F(u)$ is invoked in the first integral because we require a minimizer to satisfy the equation. Moreover, the order of $|\nabla u|^2 + F(u)$ is in fact r^{d-1} , due to Proposition 3.1. Indeed, if u grows like Cr , then ∇u would be bounded and consequently, $\int_{B_r(x^0)} |\nabla u|^2 dx$ should grow like Cr^d , and so we divide $J(u)$ by r^d . Secondly, our final goal is to neutralize the order r^d caused by the first integral using the boundary adjusted term so that $W(\cdot)$ exhibits some boundedness property. Therefore, using the fact again u grows like Cr , it is handy that $\int_{\partial B_r(x^0)} u^2 d\mathcal{H}^{d-1}$ grows like Cr^{d+1} and this is the reason we divide this boundary integral by $\frac{1}{r^{d+1}}$. The matched order between two integrals is crucial so that W is preserved under the scaling $u_{x^0, r}(x) = u(x^0 + rx)/r$ in the sense that $W(u, r, Q) = W(u_{x^0, r})$. This allows us to passing the limit $\lim_{r \rightarrow 0^+} W(\cdot, \cdot, r)$ and obtains the wanted homogeneity for blow up limits.*

The main body in this section is to prove the following proposition.

Proposition 5.3 (Weiss-type monotonicity formula). *Let $B_r(x^0) \subset \Omega$ with $r \in (0, \delta_0)$, where $\delta_0 = \text{dist}(x^0, \partial\Omega)$. Let $u \in \mathcal{K}$ be a minimizer in $B_{\delta_0}(x^0)$. Then*

$$\frac{d}{dr} W(u, r; Q) = \frac{2}{r^d} \int_{\partial B_r(x^0)} \left(\nabla u \cdot \nu - \frac{u}{r} \right)^2 d\mathcal{H}^{d-1} + \mathcal{F}_1(u, r; F) + \mathcal{Q}_1(r; Q), \quad (5.64)$$

where

$$\mathcal{F}_1(u, r; F) := \frac{1}{r^{d+1}} \int_{B_r(x^0)} u F'(u) dx, \quad (5.65)$$

and

$$\mathcal{Q}_1(r; Q) := -\frac{1}{r^{d+1}} \int_{B_r} Q^2(x) x \cdot d\mu(x) - \frac{d}{r^{d+1}} \int_{B_r} Q^2(x)\chi_{\{u>0\}} dx + \frac{1}{r^d} \int_{\partial B_r} Q^2(x)\chi_{\{u>0\}} d\mathcal{H}^{d-1}. \quad (5.66)$$

Here we employed the notation

$$d\mu = \nabla \chi_{\{u>0\}}. \quad (5.67)$$

Remark 5.4. Since $\Omega^+(u)$ is locally finite perimeter in Ω , we have that $\chi_{\{u>0\}}$ is a function of bounded variation and thus (5.67) is well-defined.

Remark 5.5. If in particular $Q(x) \equiv Q(x^0) > 0$, then $\mathcal{Q}_r(r; Q) = 0$. This can be easily obtained by (5.66) and the following simple fact

$$\begin{aligned} \frac{1}{2}r^{d+1}\mathcal{Q}_1(r; Q) &= -\frac{1}{2} \int_{B_r(x^0)} x \cdot d\mu(x) - \frac{d}{2} \int_{B_r(x^0)} \chi_{\{u>0\}} dx + \frac{r}{2} \int_{\partial B_r(x^0)} \chi_{\{u>0\}} d\mathcal{H}^{d-1} \\ &= 0. \end{aligned} \quad (5.68)$$

To prove Proposition 5.3, we need some fundamental tools, the first is a domain variation formula.

Lemma 5.6 (Domain variation formula). *Let $Q(x) \in C^0(\Omega)$ be any continuous function, let $u \in \mathcal{K}$ be a local minimizer of J in Ω and let $\xi \in C_0^\infty(\Omega; \mathbb{R}^d)$. Then we have*

$$\int_{\Omega} 2\nabla u D\xi(\nabla u)^T - (|\nabla u|^2 + F(u)) \operatorname{div} \xi \, dx + \int_{\Omega} Q^2(x)\xi(x) \cdot d\mu = 0, \quad (5.69)$$

where $d\mu$ is defined in (5.67).

Proof. Let $L := |D\xi|$ be the Lipschitz constant of ξ . For $t > 0$, we define $\Psi_t(x) = x + t\xi(x)$ and $u_t(x) = u(x + t\xi(x))$. One can show that, for $0 < t < \frac{1}{L}$, $\Psi_t: \Omega \rightarrow \Omega$ is bijective, $\Psi_t^{-1} \in C^\infty(\Omega; \mathbb{R}^d)$, and moreover $\Psi_t^{-1}(y) - y \in C_0^\infty(\Omega; \mathbb{R}^d)$ with $\operatorname{supp}(\Psi_t^{-1}(y) - y) \subset \operatorname{supp}(\Psi_t)$. A direct calculation gives $D\Psi_t = \operatorname{Id} + tD\xi$ and $|\det D\Psi_t(x)| = 1 + t \operatorname{div} \xi + o(t)$. By a change of variables,

$$\begin{aligned} \int_{\Omega} |\nabla u_t(x)|^2 \, dx &= \int_{\Omega} |\nabla u_t(\Psi_t^{-1}(y))|^2 |\det D\Psi_t^{-1}(y)| \, dy \\ &= \int_{\Omega} (|\nabla u(y)|^2 + 2t\nabla u D\xi(\nabla u)^T + o(t))(1 - t \operatorname{div} \xi + o(t)) \, dy \\ &= \int_{\Omega} |\nabla u(y)|^2 \, dy + t \int_{\Omega} 2\nabla u D\xi(\nabla u)^T \, dy - t \int_{\Omega} |\nabla u(y)|^2 \operatorname{div} \xi \, dy + o(t). \end{aligned} \quad (5.70)$$

Similarly, one has

$$\begin{aligned} \int_{\Omega} F(u_t(x)) \, dx &= \int_{\Omega} F(u_t(\Psi_t^{-1}(y))) |\det D\Psi_t^{-1}(y)| \, dy \\ &= \int_{\Omega} F(u(y))(1 - t \operatorname{div} \xi + o(t)) \, dy \\ &= \int_{\Omega} F(u(y)) \, dy - t \int_{\Omega} F(u(y)) \operatorname{div} \xi \, dy + o(t). \end{aligned} \quad (5.71)$$

By differentiating (5.70) and (5.71) with respect to t we obtain

$$\left. \frac{d}{dt} \left(\int_{\Omega} |\nabla u_t|^2 + F(u_t) \, dx \right) \right|_{t=0} = \int_{\Omega} 2\nabla u D\xi(\nabla u)^T - (|\nabla u|^2 + F(u)) \operatorname{div} \xi \, dy. \quad (5.72)$$

We also differentiate the last term in our functional

$$\left. \frac{d}{dt} \left(\int_{\Omega} Q^2(x)\chi_{\{u_t>0\}}(x) \, dx \right) \right|_{t=0} = \frac{d}{dt} \int_{\Omega} Q^2(x)\chi_{\{u>0\}}(\Psi_t(x)) \, dx = \int_{\Omega} Q^2(x)\xi \cdot d\mu(x). \quad (5.73)$$

Combining (5.72), (5.73) and the fact that u is a minimizer, we have

$$0 = \frac{d}{dt} \Big|_{t=0} J(u_t) = \int_{\Omega} 2\nabla u D\xi(\nabla u)^T - (|\nabla u|^2 + F(u)) \operatorname{div} \xi \, dx + \int_{\Omega} Q^2(x) \xi \cdot d\mu.$$

□

Next we introduce an energy identity comes from the equation satisfied by minimizers.

Lemma 5.7 (Energy identity). *Let $u \in \mathcal{K}$ be a local minimizer of J in Ω , then for a.e. $r \in (0, \delta_0)$ with $\delta_0 := \operatorname{dist}(x^0, \partial\Omega)$,*

$$\int_{B_r(x^0)} \left[|\nabla u|^2 + \frac{uF'(u)}{2} \right] dx = \int_{\partial B_r(x^0)} u \nabla u \cdot \nu \, d\mathcal{H}^{d-1}. \quad (5.74)$$

Proof. Without loss of generality, we might assume that $x^0 = 0$. For any $0 < \varepsilon < r$, we define $\phi_{\varepsilon,r} \in C_0^\infty(B_r)$ such that $\phi_{\varepsilon,r}(x) = 1$ in $B_{r-\varepsilon}$ and $\nabla \phi_{\varepsilon,r}(x) = -\frac{1}{\varepsilon} \frac{x}{|x|} \chi_{\{B_r \setminus B_{r-\varepsilon}\}}$. It is easy to see that $u\phi_{\varepsilon,r} \in H_0^1(B_r^+(u))$. Inserting $u\phi_{\varepsilon,r}$ as a test function in (2.18) yields

$$\begin{aligned} - \int_{B_r^+(u)} \frac{F'(u)}{2} (\phi_{\varepsilon,r} u) \, dx &= \int_{B_r^+(u)} \nabla u \cdot \nabla (\phi_{\varepsilon,r} u) \, dx \\ &= \int_{B_r} |\nabla u|^2 \phi_{\varepsilon,r} \, dx - \frac{1}{\varepsilon} \int_{B_r \setminus \overline{B_{r-\varepsilon}}} \nabla u \cdot \frac{x}{|x|} u \, dx. \end{aligned}$$

Letting $\varepsilon \rightarrow 0$, we obtain that for a.e. $r \in (0, \delta_0)$,

$$\int_{B_r} \left[|\nabla u|^2 + \frac{uF'(u)}{2} \right] dx = \int_{\partial B_r} u \nabla u \cdot \nu \, d\mathcal{H}^{d-1}.$$

□

In [7], Caffarelli and et. al. developed a Pohožaev-type identity for vectorial Bernoulli problem, we borrow their idea and adapt it to our settings, which will be applied to prove the monotonicity formula.

Lemma 5.8 (Pohožaev-type identity). *Let $u \in \mathcal{K}$ be a local minimizer of J in Ω , let $B_r(x^0) \subset \Omega$ with $r \in (0, \delta_0)$ where $\delta_0 := \operatorname{dist}(x^0, \partial\Omega)$. Assume that $Q(x)$ is a continuous function, then we have the following identity*

$$\begin{aligned} d \int_{B_r(x^0)} (|\nabla u|^2 + F(u)) \, dx &= 2 \int_{B_r(x^0)} |\nabla u|^2 \, dx + r \int_{\partial B_r(x^0)} (|\nabla u|^2 - 2(\nabla u \cdot \nu)^2) \, d\mathcal{H}^{d-1} \\ &\quad + r \int_{\partial B_r(x^0)} F(u) \, d\mathcal{H}^{d-1} + \int_{B_r(x^0)} Q^2(x) (x - x^0) \cdot d\mu(x), \end{aligned} \quad (5.75)$$

where the vector Radon measure μ is defined in (5.67).

Proof. Assume $x^0 = 0$ and define $\xi(x) := x\phi_{\varepsilon,r}$ where $\phi_{\varepsilon,r}$ is the test function given in Lemma 5.7. Introducing $\xi(x)$ as a test function into the domain variation formula (5.69) gives

$$\begin{aligned} & d \int_{\Omega} (|\nabla u|^2 + F(u))\phi_{\varepsilon,r} \, dx \\ &= \int_{\Omega} 2|\nabla u|^2\phi_{\varepsilon,r} + 2(\nabla u \cdot x)(\nabla\phi_{\varepsilon,r} \cdot \nabla u) \, dx - \int_{\Omega} (|\nabla u|^2 + F(u))(x \cdot \nabla\phi_{\varepsilon,r}(x)) \, dx \\ & \quad + \int_{\Omega} Q^2(x)(x\phi_{\varepsilon,r}) \cdot d\mu(x). \end{aligned} \quad (5.76)$$

Passing to the limit as $\varepsilon \rightarrow 0$ in (5.76) gives the desired result. \square

We next show that the formula (5.64) is a direct application of Lemma 5.7 and Lemma 5.8.

Proof of the formula (5.64). Assume $x^0 = 0$, then for a.e. $r \in (0, \delta_0)$, a straightforward calculation yields

$$\frac{d}{dr} \left(r^{-d-1} \int_{\partial B_r} u^2 \, d\mathcal{H}^{d-1} \right) = -2r^{-d-2} \int_{\partial B_r} u^2 \, d\mathcal{H}^{d-1} + 2r^{-d-1} \int_{\partial B_r} u \nabla u \cdot \nu \, d\mathcal{H}^{d-1}. \quad (5.77)$$

We further compute

$$\begin{aligned} \frac{1}{2}r^{d+1} \frac{d}{dr} W(u, r; Q) &= -\frac{d}{2} \int_{B_r} |\nabla u|^2 + F(u) \, dx - \frac{d}{2} \int_{B_r} Q^2(x)\chi_{\{u>0\}} \, dx \\ & \quad + \frac{r}{2} \int_{\partial B_r} |\nabla u|^2 + F(u) + Q^2(x)\chi_{\{u>0\}} \, d\mathcal{H}^{d-1} \\ & \quad + \frac{1}{r} \int_{\partial B_r} u^2 \, d\mathcal{H}^{d-1} - \int_{\partial B_r} u \nabla u \cdot \nu \, d\mathcal{H}^{d-1}. \end{aligned}$$

Using the identity (5.75) for the first integral on the right-hand side, we get

$$\begin{aligned} \frac{1}{2}r^{d+1} \frac{d}{dr} W(u, r; Q) &= -\frac{1}{2} \left(2 \int_{B_r} |\nabla u|^2 + r \int_{\partial B_r} |\nabla u|^2 - 2(\nabla u \cdot \nu)^2 \, d\mathcal{H}^{d-1} \right. \\ & \quad \left. + r \int_{\partial B_r} F(u) \, d\mathcal{H}^{d-1} + \int_{B_r} Q^2(x)x \cdot d\mu(x) \right) \\ & \quad - \frac{d}{2} \int_{B_r} Q^2(x)\chi_{\{u>0\}} \, dx + \frac{r}{2} \int_{\partial B_r} (|\nabla u|^2 + F(u)) \, d\mathcal{H}^{d-1} \\ & \quad + \frac{r}{2} \int_{\partial B_r} Q^2(x)\chi_{\{u>0\}} \, d\mathcal{H}^{d-1} + \frac{1}{r} \int_{\partial B_r} u^2 \, d\mathcal{H}^{d-1} \\ & \quad - \int_{\partial B_r} u \nabla u \cdot \nu \, d\mathcal{H}^{d-1}. \end{aligned}$$

Tidying the results, we obtain

$$\begin{aligned} \frac{1}{2}r^{d+1}\frac{d}{dr}W(u,r;Q) &= -\int_{B_r}|\nabla u|^2 dx + r\int_{\partial B_r}(\nabla u \cdot \nu)^2 d\mathcal{H}^{d-1} + \frac{1}{r}\int_{\partial B_r}u^2 d\mathcal{H}^{d-1} \\ &\quad - \int_{\partial B_r}u\nabla u \cdot \nu d\mathcal{H}^{d-1} - \frac{1}{2}\int_{B_r}Q^2(x)x \cdot d\mu(x) \\ &\quad - \frac{d}{2}\int_{B_r}Q^2(x)\chi_{\{u>0\}} dx + \frac{r}{2}\int_{\partial B_r}Q^2(x)\chi_{\{u>0\}} d\mathcal{H}^{d-1}, \end{aligned} \quad (5.78)$$

where μ is the vector Radon measure defined in (5.67). It follows from Lemma 5.7 and applying (5.74) for the first integral on the right-hand side of (5.78), we obtain for a.e. $r \in (0, \delta_0)$,

$$\begin{aligned} \frac{1}{2}r^{d+1}\frac{d}{dr}W(u,r;Q) &= r\int_{\partial B_r}(\nabla u \cdot \nu)^2 d\mathcal{H}^{d-1} + \frac{1}{r}\int_{\partial B_r}u^2 d\mathcal{H}^{d-1} \\ &\quad - 2\int_{\partial B_r}u\nabla u \cdot \nu d\mathcal{H}^{d-1} + \int_{B_r}\frac{uF'(u)}{2} dx \\ &\quad - \frac{1}{2}\int_{B_r}Q^2(x)x \cdot d\mu(x) - \frac{d}{2}\int_{B_r}Q^2(x)\chi_{\{u>0\}} dx \\ &\quad + \frac{r}{2}\int_{\partial B_r}Q^2(x)\chi_{\{u>0\}} d\mathcal{H}^{d-1}. \end{aligned} \quad (5.79)$$

Notice that

$$r\int_{\partial B_r}(\nabla u \cdot \nu)^2 d\mathcal{H}^{d-1} - 2\int_{\partial B_r}u\nabla u \cdot \nu d\mathcal{H}^{d-1} + \frac{1}{r}\int_{\partial B_r}u^2 d\mathcal{H}^{d-1} = r\int_{\partial B_r}\left(\nabla u \cdot \nu - \frac{u}{r}\right)^2 d\mathcal{H}^{d-1}. \quad (5.80)$$

The desired result follows from (5.79) and (5.80). \square

We next introduce another monotonicity formula, which is different from (5.64) and is obtained by comparing $u_{x^0,r}$ with its one-homogeneous extensions in B_1 .

Proposition 5.9. *Let $B_r(x^0) \subset \Omega$ with $r \in (0, \delta_0)$, where $\delta_0 = \text{dist}(x^0, \partial\Omega)$. Let $u \in \mathcal{K}$ be a local minimizer in $B_{\delta_0}(x^0)$ and let $z_{x^0,r}: B_1 \rightarrow \mathbb{R}$ be the one-homogeneous extension of $u_{x^0,r}$ defined by*

$$z_{x^0,r} := |x|u_{x^0,r}\left(\frac{x}{|x|}\right). \quad (5.81)$$

Then

$$\begin{aligned} \frac{d}{dr}W(u_{x^0,r}) &= \frac{d}{r}(W(z_{x^0,r}) - W(u_{x^0,r})) + \frac{1}{r}\int_{\partial B_1}|x \cdot \nabla u_{x^0,r} - u_{x^0,r}|^2 d\mathcal{H}^{d-1} \\ &\quad + \mathcal{F}_2(u_{x^0,r}; F) + \mathcal{Q}_2(u_{x^0,r}; Q), \end{aligned} \quad (5.82)$$

where

$$\mathcal{F}_2(u_{x^0,r}; F) = \frac{1}{r}\int_{\partial B_1}F(ru_{x^0,r}) d\mathcal{H}^{d-1} - \frac{d}{r}\int_{B_1}F(rz_{x^0,r}) dx, \quad (5.83)$$

and

$$\mathcal{Q}_2(u_{x^0,r}; Q) = \frac{1}{r}\int_{\partial B_1}Q^2(x^0 + rx)\chi_{\{u_{x^0,r}>0\}} d\mathcal{H}^{d-1} - \frac{d}{r}\int_{B_1}Q^2(x^0 + rx)\chi_{\{z_r>0\}} dx. \quad (5.84)$$

Remark 5.10. In [27], Reifenberg exhibited the idea that a purely variational inequality, known as an epiperimetric inequality, will encode the local behavior of the free boundary. Since then, extensive literature has been dedicated to prove different kinds of epiperimetric inequalities. (We refer to [29] for epiperimetric inequality for problem (1.1)). The original idea of Reifenberg is: if $\mathcal{E}: H^1(B_1) \rightarrow \mathbb{R}$ is a given energy which satisfies the following properties:

- (1) $\mathcal{E}(u_r) \leq \mathcal{E}(v_r)$ for every $v_r \in H^1(B_1)$ with $v_r = u_r$ on ∂B_1 ;
- (2) $\frac{d}{dr}\mathcal{E}(u_r) \geq \frac{C_1}{r} \int_{\partial B_1} |x \cdot \nabla u_r - u_r|^2 d\mathcal{H}^{d-1}$ for every $r \in (0, 1)$ and for some positive constants $C_1 > 0$;
- (3) $\frac{d}{dr}\mathcal{E}(u_r) \geq \frac{C_2}{r} (\mathcal{E}(z_r) - \mathcal{E}(u_r))$ for every $r \in (0, 1)$ and for some positive constants $C_2 > 0$. Here z_r is the one-homogeneous extension of u_r defined in (5.81).
- (4) $\mathcal{E}(h_r) \leq (1 - \varepsilon)\mathcal{E}(u_r)$ for all $h_r = u_r = z_r$ on ∂B_1 .

Then the uniqueness of blow-up limit is a direct corollary of (1)-(4). As an application of the uniqueness of blow-up limit, one can determine the regularity of the regular set $\partial_{\text{reg}}\Omega^+(u)$, even the regularity of the singular set $\partial_{\text{sing}}\Omega^+(u)$. In our context, the energy to which will apply (1)-(4) is

$$\mathcal{E}(u) - \frac{\omega_d}{2} = \left(\int_{B_1} |\nabla u|^2 + F(u) + \chi_{\{u>0\}} dx - \int_{\partial B_1} u^2 d\mathcal{H}^{d-1} \right) - \frac{\omega_d}{2}.$$

This is the core reason we put forward the formula (5.82). Observe that (1)-(3) is a direct result of the formula (5.82) (Please see Remark 5.12 below). However, (4) is unknown and remains to be an open problem. In this paper, we will use an approach to show the uniqueness of blow-up limit (see Section) heavily dependent on the geometry (flatness) of the free boundary. However, we believe that the regularity of problem (1.7) can also be obtained following Reifenberg's idea and Proposition 5.9 will help us to prove (2) and (3).

Proof. Assume $x^0 = 0$. It follows from the definition of z_r that for each $r \in (0, \delta_0)$ one can write $z_r: B_1 \rightarrow \mathbb{R}$ in polar coordinates $\rho \in (0, 1]$, $\theta \in \mathbb{S}^{d-1}$ as $z_r(\rho, \theta) = \rho z_r(1, \theta)$. Therefore, we have the following identity.

$$\begin{aligned} \int_{B_1} |\nabla z_r|^2 dx - \int_{\partial B_1} z_r^2 d\mathcal{H}^{d-1} &= \int_0^1 \rho^{d-1} d\rho \int_{\mathbb{S}^{d-1}} (z_r^2(1, \theta) + |\nabla_{\theta} z_r(1, \theta)|^2) d\theta - \int_{\mathbb{S}^{d-1}} z_r^2 d\theta \\ &= \frac{1}{d} \int_{\mathbb{S}^{d-1}} |\nabla_{\theta} z_r(1, \theta)|^2 d\theta - \frac{d-1}{d} \int_{\mathbb{S}^{d-1}} z_r^2(1, \theta) d\theta \\ &= \frac{1}{d} \int_{\partial B_1} (|\nabla u_r|^2 - (x \cdot \nabla u_r)^2) d\mathcal{H}^{d-1} - \frac{d-1}{d} \int_{\partial B_1} u_r^2 d\mathcal{H}^{d-1}, \end{aligned}$$

where we have used $|\nabla u|^2 = (\partial_\rho u)^2 + \rho^{-2}|\nabla_\theta u|^2$ if $u = u(\rho, \theta)$. Thus, let $\mathcal{F}_2(u_r; F)$ and $\mathcal{Q}_2(u_r; Q)$ be defined in (5.83) and (5.84), and it follows from (5.63) that

$$\begin{aligned} & \frac{d}{dr}(W(z_r) - W(u_r)) + \frac{1}{r} \int_{\partial B_1} |x \cdot \nabla u_r - u_r|^2 d\mathcal{H}^{d-1} + \mathcal{F}_2(u_r; F) + \mathcal{Q}_2(u_r; F) \\ &= -\frac{d}{r} \int_{B_1} (|\nabla u_r|^2 + F(ru_r) + Q^2(rx)\chi_{\{u_r > 0\}}) dx \\ &+ \frac{1}{r} \int_{\partial B_1} (|\nabla u_r|^2 + F(ru_r) + Q^2(rx)\chi_{\{u_r > 0\}}) d\mathcal{H}^{d-1} \\ &+ \frac{2}{r} \int_{\partial B_1} u_r^2 d\mathcal{H}^{d-1} - \frac{2}{r} \int_{\partial B_1} u_r (\nabla u_r \cdot x) d\mathcal{H}^{d-1}. \end{aligned}$$

On the other hand, by a direct calculation, we have

$$\begin{aligned} & \frac{d}{dr} \left(r^{-d} \int_{B_r} |\nabla u|^2 + F(u) + Q^2(x)\chi_{\{u > 0\}} \right) \\ &= -\frac{d}{r} \int_{B_1} (|\nabla u_r|^2 + F(ru_r) + Q^2(rx)\chi_{\{u_r > 0\}}) dx \\ &+ \frac{1}{r} \int_{\partial B_1} (|\nabla u_r|^2 + F(ru_r) + Q^2(rx)\chi_{\{u_r > 0\}}) d\mathcal{H}^{d-1}. \end{aligned}$$

This proves the result by employing the equality (5.77). \square

In the end of this section, we collect some properties of the functions $\mathcal{F}_i(u, r; F)$ and $\mathcal{Q}_i(u, r; Q)$ for $i = 1, 2$.

Proposition 5.11. *Let $\mathcal{F}_i(u, r; F)$ and $\mathcal{Q}_i(u, r; Q)$ for $i = 1, 2$ be defined as in Proposition 5.3 and Proposition 5.9, respectively. Then the following does hold:*

- (1) *If $u \in \mathcal{K}$ is a minimizer of J in $B_r(x^0)$ and let $F(t)$ satisfies (1.8) with $F'(0) = 0$, then $\mathcal{F}_i(u, r; F) \geq 0$ for a.e. $r \in (0, \delta_0)$.*
- (2) *If $u \in \mathcal{K}$ is a minimizer of J in $B_r(x^0)$ and let $F(t)$ satisfies (1.8), then*

$$\mathcal{F}_i(u, r; F) \geq -C \quad \text{for } i = 1, 2, \quad (5.85)$$

where C is a constant depending on d, M_0 and F_0 but independent of r and x^0 .

- (3) *If $Q(x) \in C^0(B_r(x^0))$, then*

$$\mathcal{Q}_i(r; Q) \geq -C \frac{Q_{\max}}{r} \text{osc}_{B_r(x^0)} Q \quad \text{for a.e. } r \in (0, \delta_0). \quad (5.86)$$

Here C is a constant depending only on the continuity norm of Q and $\text{osc}_{B_r(x^0)}(Q) := \max_{B_r} Q - \min_{B_r(x^0)} Q$.

- (4) *If $Q(x) \in C^\beta(B_r(x^0))$, then*

$$\mathcal{Q}_i(r; Q) \geq -C \frac{Q_{\max}}{r^{1-\beta}} [Q]_{\beta; B_r(x^0)} \quad \text{for a.e. } r \in (0, \delta_0), \quad (5.87)$$

Here C is a constant depending only on $[Q]_{\beta; B_r(x^0)} := \sup_{x \neq y} \frac{|Q(x) - Q(y)|}{|x - y|^\beta}$.

Proof. Assume $x^0 = 0$. (1). In view of (1.8), $F'(t)$ is an increasing function for $t \in \mathbb{R}$, therefore $F'(0) = 0$ implies that $F'(t) \geq F'(0) = 0$ for all $t \geq 0$. It follows from Proposition 2.2 that $u \geq 0$ in $B_r(x^0)$ and the definition of $\mathcal{F}_1(u, r; F)$ in (5.65) that $\mathcal{F}_1(u, r; F) \geq 0$. In order to prove $\mathcal{F}_2(u, r; F) \geq 0$, it suffices to prove that

$$\frac{d}{r} \int_{B_1} F(rz_r) dx = \frac{d}{r} \int_0^1 \rho^{d-1} d\rho \int_{\mathbb{S}^{d-1}} F(r\rho z_r(1, \theta)) d\theta \leq \frac{1}{r} \int_{\partial B_1} F(ru_r) d\mathcal{H}^{d-1}. \quad (5.88)$$

In fact, it follows from $F'(t) \geq 0$ for all $t \geq 0$ that

$$\begin{aligned} \int_0^1 \rho^{d-1} d\rho \int_{\mathbb{S}^{d-1}} F(r\rho z_r(1, \theta)) d\theta &\leq \int_0^1 \rho^{d-1} d\rho \int_{\mathbb{S}^{d-1}} F(rz_r(1, \theta)) d\theta \\ &= \frac{1}{d} \int_{\mathbb{S}^{d-1}} F(rz_r(1, \theta)) d\theta = \frac{1}{d} \int_{\partial B_1} F(ru_r) d\mathcal{H}^{d-1}, \end{aligned}$$

where we have used the fact that $0 \leq r\rho z_1(1, \theta) \leq rz_r(1, \theta)$ for $\rho \in (0, 1]$ in the first inequality and $u_r = z_r \in H^1(\partial B_1)$ in the last inequality. This gives (5.88).

(2). Since u is a minimizer, we have from Proposition 3.1 that $|\nabla u| \leq C$ for some constant depending on d and F_0 . This implies that $|u(x)| \leq C|x - x^0| \leq Cr$ in $B_r(x^0)$. On the other hand, in via of (1.8) and a direct calculation, we obtain

$$\max_{u \geq 0} |F'(u)| \leq 2(|f(0)| + \max_{t \geq 0} |f'(t)||u|) \leq 2 \left(\frac{F_0}{2} + \frac{F_0}{2} M_0 \right) \leq C(F_0, M_0). \quad (5.89)$$

Therefore, it follows from (5.65) and (5.89) that

$$|\mathcal{F}_1(u, r; F)| \leq \frac{1}{r^{d+1}} \int_{B_r} |uF'(u)| dx \leq \frac{C}{r^d} \max_{u \geq 0} |F'(u)| \int_{B_r} dx \leq C(d, F_0, M_0). \quad (5.90)$$

This proves (5.85) for $i = 1$. For $\mathcal{F}_2(u_r; F)$, we compute

$$\begin{aligned} \mathcal{F}_2(u_r; F) &= \frac{1}{r} \int_{\partial B_1} F(0) + F'(r\xi_r)ru_r d\mathcal{H}^{d-1} - \frac{d}{r} \int_0^1 \rho^{d-1} d\rho \int_{\mathbb{S}^{d-1}} F(0) + F'(r\zeta_r)r\rho z_r(1, \theta) d\theta \\ &= \int_{\partial B_1} F'(r\xi_r)u_r d\mathcal{H}^{d-1} - d \int_0^1 \rho^d d\rho \int_{\mathbb{S}^{d-1}} F'(r\zeta_r)z_r(1, \theta) d\theta \\ &= \int_{\partial B_1} F'(r\xi_r)u_r d\mathcal{H}^{d-1} - \frac{d}{d+1} \int_{\partial B_1} F'(r\zeta_r)u_r d\mathcal{H}^{d-1} \\ &= \frac{1}{r^d} \int_{\partial B_r} F'(\xi)u d\mathcal{H}^{d-1} - \frac{d}{d+1} \frac{1}{r^d} \int_{\partial B_r} F'(\zeta)u d\mathcal{H}^{d-1} \end{aligned}$$

Then it follows from (5.89) and a similar calculation as in (5.90) that (5.85) holds for $i = 2$.

(3). Recalling (5.66) and compute

$$\begin{aligned}
\mathcal{Q}_1(r; Q) &= -\frac{1}{r^{d+1}} \int_{B_r} (Q^2(x) - Q^2(0))x \cdot d\mu(x) - \frac{d}{r^{d+1}} \int_{B_r} (Q^2(x) - Q^2(0))\chi_{\{u>0\}} dx \\
&\quad + \frac{1}{r^d} \int_{\partial B_r} (Q^2(x) - Q^2(0))\chi_{\{u>0\}} d\mathcal{H}^{d-1} \\
&\geq -C \frac{Q_{\max} \text{osc}_{B_r} Q}{r^d} \int_{B_r} d|\mu|(x) - C \frac{dQ_{\max} \text{osc}_{B_r} Q}{r^{d+1}} \int_{B_r} \chi_{\{u>0\}} dx \\
&\quad - C \frac{Q_{\max} \text{osc}_{B_r} Q}{r^d} \int_{\partial B_r} \chi_{\{u>0\}} d\mathcal{H}^{d-1} \\
&\geq -C \frac{Q_{\max}}{r} \text{osc}_{B_r} Q,
\end{aligned} \tag{5.91}$$

where we have used Remark 5.5 in the first identity. As for \mathcal{Q}_2 , we first notice that if Q is a constant, then $\mathcal{Q}_2 \equiv 0$, indeed,

$$\int_{B_1} \chi_{\{z_r>0\}} = \int_0^1 s^{d-1} ds \int_{\mathbb{S}^{d-1}} \chi_{\{u_r>0\}} d\mathcal{H}^{d-1} = \frac{1}{d} \int_{\partial B_1} \chi_{\{u_r>0\}} d\mathcal{H}^{d-1}.$$

Therefore, we can rewrite \mathcal{Q}_2 in the following form

$$\mathcal{Q}_2(u_r; Q) = \frac{1}{r} \int_{\partial B_1} \chi_{\{u_r>0\}} (Q^2(rx) - Q^2(0)) d\mathcal{H}^{d-1} - \frac{d}{r} \int_{B_1} \chi_{\{z_r>0\}} (Q^2(rx) - Q^2(0)) dx.$$

Then we can follow a similar argument as in (5.91) to conclude the proof.

(4). The desired result (5.87) follows by a similar calculation as in (3), so we omit it. \square

Remark 5.12. *If in particular $f(0) = 0$ and $Q \equiv \text{const}$, it follows from (5.82), (5.83) and the first property in Proposition 5.11 that*

$$\frac{d}{dr} W(u_{x^0, r}) \geq \frac{1}{r} \int_{\partial B_1} |x \cdot \nabla u_{x^0, r} - u_{x^0, r}|^2 d\mathcal{H}^{d-1} \quad \text{for a.e. } r \in (0, \delta_0),$$

and

$$\frac{d}{dr} W(u_{x^0, r}) \geq \frac{d}{r} (W(z_{x^0, r}) - W(u_{x^0, r})) \quad \text{for a.e. } r \in (0, \delta_0).$$

Notice that these two inequalities were exactly the second and the third property we have mentioned in Remark 5.10.

6. STRUCTURE OF THE FREE BOUNDARY

This section dedicates to prove Theorem 1.2 using those technical tools obtained in the previous sections. The first one is a direct application of the Weiss-type monotonicity formula we obtained in Proposition 5.3 and Proposition 5.9, which tells us that blow-up limits of minimizers are homogeneous functions of degree one.

Proposition 6.1 (Homogeneity of blow-up limits). *Let u be a minimizer of J in Ω with $0 \in \Omega$ and $u(0) = 0$, let $F(t)$ satisfy (1.8). Assume that Q is Dini-continuous at the origin, namely,*

$$\lim_{r \rightarrow 0^+} \int_r^1 \frac{1}{s} \operatorname{osc}_{B_s} Q \, ds < \infty. \quad (6.92)$$

Let $B_{r_n} \subset \Omega$ with $r_n \rightarrow 0$ as $n \rightarrow \infty$. Let u_0 be a blow-up limit of u at the origin with respect to the sequence r_n . Then

- (1) The limit $\lim_{r \rightarrow 0^+} W(u, r, Q)$ exists and is finite for a.e. $r \in (0, \delta_0)$;
- (2) u_0 is a homogeneous function of degree one.

Proof. Define $\rho(r) := \int_0^r \frac{1}{s} \operatorname{osc}_{B_s} Q \, ds$. Then it follows from the assumption (6.92) that $\rho(r)$ is absolutely continuous with respect to $r \in (0, \delta_0)$ and thus $\rho'(r)$ and $\lim_{r \rightarrow 0^+} \rho(r)$ exists for a.e. $r \in (0, \delta_0)$. It follows from (5.64) that $\frac{d}{dr} W(u, r; Q) \geq \mathcal{F}_i(u, r; F) + \mathcal{Q}_i(r; Q)$ for $i = 1$ and $i = 2$ with $\mathcal{Q}_2(r; Q) = 0$. It follows from the second and the third property of Proposition (5.11) that $\frac{d}{dr} W(u, r; Q) \geq -C - \frac{Q_{\max}}{r} \operatorname{osc}_{B_r} Q$. Thus the function $G(u, r; Q) := W(u, r; Q) + Cr + CQ_{\max}\rho(r)$ satisfies $\frac{d}{dr} G(u, r; Q) \geq 0$ for a.e. $r \in (0, \delta_0)$ and therefore the limit $\lim_{r \rightarrow 0^+} G(u, r; Q)$ exists. Since $\lim_{r \rightarrow 0^+} Cr = 0$, due to the fact that C is independent of r and $\lim_{r \rightarrow 0^+} \rho(r) = 0$ we obtain that the limit $\lim_{r \rightarrow 0^+} W(u, r; Q) \in [-\infty, +\infty)$ exists. Since $u(0) = 0$ we may see by regularity results that $W(u, 0^+; Q) > -\infty$. This proves (1).

(2). To prove the second statement, we use the formula (5.64) and one can also use the formula (5.82) to obtain the same result without effort. We integrate the inequality (5.64) and we have

$$\begin{aligned} & \int_{r_n \varrho}^{r_n \sigma} \frac{1}{r} \int_{\partial B_1} |x \cdot \nabla u_r - u_r|^2 \, d\mathcal{H}^{d-1} \, dr \\ & \leq W(u, r_n \sigma, Q) - W(u, r_n \varrho, Q) + \int_{r_n \varrho}^{r_n \sigma} C \, dr + CQ_{\max}(\rho(r_n \sigma) - \rho(r_n \varrho)), \end{aligned} \quad (6.93)$$

where we have used (5.85), (5.86) and (6.92). It follows by rescaling (6.93) that

$$\begin{aligned} 0 & \leq \int_{B_\sigma \setminus B_\varrho} |z|^{-d-2} (\nabla u_n \cdot z - u_n) \, dz \\ & \leq W(u, r_n \sigma, Q) - W(u, r_n \varrho, Q) + Cr_n(\sigma - \varrho) + CQ_{\max}(\rho(\sigma r_n) - \rho(\varrho r_n)). \end{aligned}$$

Passing to the limit as $n \rightarrow \infty$ we have the desired homogeneity. \square

Remark 6.2. *Although in the above Proposition, we mainly focused on the homogeneity of free boundary point $0 \in \partial\Omega^+(u)$, one can also consider the case $0 \in \Omega^+(u)$. In fact, if $0 \in \Omega$ with $u(0) > 0$, then it follows from the regularity results $|\nabla u|^2 \leq C$ in Ω that $W(u, r, Q) \leq C_1 - \frac{C_2}{r^2}$ for any $0 < r < \delta_0$. This implies that $W(u, 0^+, Q) = -\infty$ for all such points. The homogeneity of u_0 at these points can be verified same as in the second part of the proof in Proposition 6.1.*

Remark 6.3. *In a significant work [33] on two-dimensional gravity water waves Vărvărucă and Weiss on two-dimensional gravity water waves with vorticity. They put forward a monotonicity formula to investigate the local behavior of "stagnation points", at which points u is a $\frac{3}{2}$ -homogeneous*

function. We point out that one can apply Proposition 6.1 to deduce the homogeneity for those points away from the stagnation points.

In the rest of this section, we will focus on the limit

$$\vartheta := \lim_{r \rightarrow 0^+} \frac{|B_r(x^0) \cap \{u > 0\}|}{|B_r(x^0)|} \quad \text{for every } x^0 \in \partial\Omega^+(u). \quad (6.94)$$

It follows from Proposition 3.6 that $\vartheta \in (0, 1)$ and we will finally prove that $\vartheta \in [\frac{1}{2}, 1)$ for every free boundary point and that the regular part of the free boundary $\partial_{\text{reg}}\Omega^+(u)$ is fully characterized by this limit. Most of the ideas come from [25] and we begin with the following observation.

Proposition 6.4. *Let $u \in \mathcal{K}$ be a minimizer of J in Ω with $0 \in \Omega$ and $u(0) = 0$, let $F(t)$ satisfy (1.8) and let $Q(x)$ be a Hölder continuous function at the origin which satisfies (1.9), then the limit ϑ defined in (6.94) exists and*

$$\vartheta = \frac{|B_1 \cap \{u_0 > 0\}|}{|B_1|}. \quad (6.95)$$

Proof. Let $\{r_n\}_{n=1}^\infty > 0$ be a vanishing sequence and let u_n be a blow-up sequence at with respect to B_{r_n} . It follows from Proposition 4.5 (3), (4) and Proposition 6.1 (2) that

$$\begin{aligned} \lim_{n \rightarrow \infty} W(u_n; Q(r_n x)) &= \int_{B_1} |\nabla u_0|^2 dx - \int_{\partial B_1} u_0^2 d\mathcal{H}^{d-1} + Q^2(0) \lim_{r \rightarrow 0^+} \frac{\int_{B_r} \chi_{\{u > 0\}} dx}{r^d} \\ &= Q^2(0) \omega_d \lim_{r \rightarrow 0^+} \frac{|B_r \cap \{u > 0\}|}{|B_r|} = Q^2(0) \omega_d \vartheta, \end{aligned}$$

The limit here exists because $\lim_{r \rightarrow 0^+} W(u, r; Q)$ exists (by Proposition 6.1 (1)). On the other hand, since $\chi_{\{u_n > 0\}} \rightarrow \chi_{\{u_0 > 0\}}$ in $L^1_{\text{loc}}(\mathbb{R}^d)$ (by Proposition 4.5 (2)), we have that

$$\lim_{n \rightarrow \infty} \int_{B_1} \chi_{\{u_n > 0\}} Q^2(r_n x) dx = Q^2(0) \int_{B_1} \chi_{\{u_0 > 0\}} dx$$

Thus

$$Q^2(0) \omega_d \frac{|B_1 \cap \{u_0 > 0\}|}{|B_1|} = \lim_{n \rightarrow \infty} W(u_n; Q(r_n x)) = Q^2(0) \omega_d \vartheta,$$

which is exactly (6.95). \square

Remark 6.5. *In view of (6.95) and (4.59) that $\vartheta \in (0, 1)$ and thus u_0 is a non-constantly-vanishing function in B_1 . That is, $u_0 \not\equiv 0$ in $B_1(0)$.*

At this stage, we point out Theorem 1.2 is a direct corollary of the following proposition.

Proposition 6.6. *Let $u \in \mathcal{K}$ be a minimizer of J in Ω with $0 \in \Omega$ and $u(0) = 0$, let $F(t)$ satisfy (1.8) and let $Q(x)$ be a Hölder continuous function at the origin which satisfies (1.9) with $Q(0) = 1$. Then for ϑ be defined in (6.94), we have*

- (1) $\vartheta \in [\frac{1}{2}, 1)$ for every $x^0 \in \partial\Omega^+(u)$.
- (2) $\partial_{\text{reg}}\Omega^+(u) = (\Omega^+(u))^{(1/2)}$.

Proof. The upper bound of ϑ follows directly from (6.95) and (4.59). Assume that $x^0 \in \partial\Omega^+(u)$ and that u_n is the blow-up sequence converges to a blow-up limit $u_0: \mathbb{R}^d \rightarrow \mathbb{R}$. It follows once again from Proposition 4.5 (6) and Proposition 6.1 (2) that u_0 is one-homogeneous and harmonic in $B_1^+(u_0)$. This implies that u_0 is a solution to the equation (in polar coordinates) $-\Delta_{\mathbb{S}} u_0 = (d-1)u_0$ on $\partial B_1^+(u_0)$, where $\Delta_{\mathbb{S}}$ is defined in (1.11). It follows from Remark 4.8 in [25] that $\mathcal{H}^{d-1}(\partial B_1^+(u_0)) \geq \frac{d\omega_d}{2}$. Using the homogeneity of u_0 , we immediately have $|B_1 \cap \{u_0 > 0\}| \geq \frac{\omega_d}{2}$. Now, the convergence $\lim_{n \rightarrow \infty} \int_{B_1} \chi_{\{u_n > 0\}} dx = \int_{B_1} \chi_{\{u_0 > 0\}} dx$ by Proposition 4.5 (2) implies that

$$\vartheta = \lim_{n \rightarrow \infty} \frac{|B_1 \cap \{u_n > 0\}|}{|B_1|} = \frac{|B_1 \cap \{u_0 > 0\}|}{|B_1|} \geq \frac{1}{2},$$

which concludes (1). In the case of equality $\gamma = \frac{1}{2}$, we have by Remark 4.8 in [26] that $u_0|_{\partial B_1}$ is precisely the first eigenvalue on \mathbb{S}_+^{d-1} , whose one-homogeneous extension is precisely the one-plane solution $(x \cdot \nu)^+$ for some $\nu \in \partial B_1$. This implies that if u is a minimizer of J in Ω , then $\partial_{\text{reg}} \Omega^+(u) = (\Omega^+(u))^{(1/2)}$. \square

7. REGULARITY OF THE FREE BOUNDARY NEAR REGULAR POINTS

In this section, we focus on the regularity of regular free boundary points, and the structure of singular free boundary points will be considered in the forthcoming section. We will first prove the uniqueness of blow-up limit and the polynomial convergence rate of a blow-up sequence. Then regularity of free boundaries is a direct consequence of these two facts. Such techniques are prevalent in recent research and are robust because there are quantities of arguments, for instance, via an epiperimetric inequality, to verify the uniqueness of blow-up limit and the rate of convergence for blow-up sequences. This method was mainly borrowed from [Chapter 8, [31]]. In this section we assume without loss of generality that $0 \in \Omega$ and that $Q(0) = 1$. We begin by showing that minimizers of J in Ω are viscosity solutions in the sense of Definition 1.3.

Proposition 7.1 (Minimizers are viscosity solutions). *Let $u \in \mathcal{K}$ be a minimizer of J in Ω . Then u is a viscosity solution of (1.7) in the sense of definition 1.3.*

Proof. Step I. Assume that $x^0 \in \Omega^+(u)$ and that $\phi \in C^\infty(\Omega)$ touching u from below at x^0 . Then $u \in C^{2,\alpha}$ smooth at x^0 , $(u - \phi)(x^0) = 0$ and $(u - \phi)(x) \geq 0$ in a small neighborhood of x^0 . This implies that the function $(u - \phi)$ attains a local minimum at x^0 and so $\Delta(u - \phi)(x^0) \geq 0$ by the second derive test of calculus. Therefore,

$$\Delta\phi(x^0) + f(\phi(x^0)) = \Delta\phi(x^0) + f(u(x^0)) \leq \Delta u(x^0) + f(u(x^0)) = 0,$$

where we have used the fact that $\Delta u(x^0) + f(u(x^0)) = 0$ for $x^0 \in \Omega^+(u)$. It is easy to check the case in a similar way when ϕ touches u from above at x^0 .

Step II. It remains to check that u satisfies the free boundary condition $|\nabla u| = Q(x^0)$ in the viscosity sense. To this end, suppose that the function ϕ touches u from below at $x^0 \in \partial\Omega^+(u)$ and assume without loss of generality that $x^0 = 0$. Consider the following two blow-up sequences

$u_n(x) := \frac{u(r_n x)}{r_n}$ and $\phi_n(x) := \frac{\phi(r_n x)}{r_n}$ where $r_n \rightarrow 0^+$ as $n \rightarrow \infty$. Since u is a minimizer, we have that the convergence is uniform in B_1 . Since ϕ is a smooth function in Ω , the limit $\phi_0 := \lim_{n \rightarrow \infty} \phi_n(x)$ exists and we have $\phi_0 = \xi_0 \cdot x$, where the vector $\xi_0 \in \mathbb{R}^d$ is precisely the gradient $\nabla\phi(0)$. Without loss of generality we may assume that $\xi_0 = A e_d$ for some non-negative constant $A \geq 0$. Thus,

$$|\nabla\phi(0)| = |\nabla\phi_0(0)| = A \quad \text{and} \quad \phi_0(x) = Ax_d. \quad (7.96)$$

Moreover, we can assume that $A > 0$ since otherwise the inequality $|\nabla\phi| \leq Q(0) = 1$ holds trivially. It follows from Proposition 4.5 (6), Proposition 6.1 (2) and Remark 6.5 that $u_0 \in H_{\text{loc}}^1(\mathbb{R}^d)$ is a non-trivial, continuous and one-homogeneous function, harmonic in $B_1^+(u_0)$. By the uniform convergence we have $u_0 \geq \phi_0$ and this implies $\{u_0 > 0\} \supset \{\phi_0 > 0\} = \{x_d > 0\}$. Thus it follows from Lemma 5.31 [28] that $\{u_0 > 0\} = \{x_d > 0\}$ because the case $\{u_0 > 0\} = \{x_d \neq 0\}$ is ruled out (by (4.59) or Proposition 6.6). Moreover, u_0 is a global minimizer of $J_0(v)$ where $J_0(v)$ is defined in (4.58) and hence satisfies the optimality condition $|\nabla u_0| = Q(0) = 1$. In conclusion, $u_0(x) = x_d^+$ for $x \in \mathbb{R}^d$. Finally, it follows from $u_0 \geq \phi_0$ and (7.96) that $A \leq 1$, and we have from (7.96) that $|\nabla\phi(0)| \leq 1$, as desired. The case when ϕ touches u from above is similar and we omit it. \square

Remark 7.2. *Since minimizers are continuous in Ω , we define*

$$\tilde{f}(x) := -(f \circ u)(x), \quad \text{for every } x \in \Omega. \quad (7.97)$$

It follows from a direct calculation that

$$|\tilde{f}(x)| = |f(u(x))| \leq |f(0)| + \max_{s \geq 0} |f'(s)| \|u\|_{L^\infty(\Omega)} \leq C(F_0, M_0),$$

where $M_0 := \sup_\Omega \Psi$ and Ψ is given in (1.6). Thus, $\tilde{f} \in C^0(\Omega) \cap L^\infty(\Omega)$. Moreover, we infer from Proposition 7.1 that u is a viscosity solution to

$$\Delta u = \tilde{f}(x) \quad \text{in} \quad \Omega^+(u), \quad |\nabla u| = Q(x) \quad \text{on} \quad \partial\Omega^+(u). \quad (7.98)$$

In what follows, we will apply De Silva's viscosity approach to prove the regularity of the regular set $\partial_{\text{reg}}\Omega^+(u)$ of the free boundary. The first observation is that the free boundaries are ε -flat near points in $\partial_{\text{reg}}\Omega^+(u)$.

Lemma 7.3 (ε -flat for regular free boundary points). *Let $u \in \mathcal{K}$ be a minimizer of J in Ω . Assume that $0 \in \partial_{\text{reg}}\Omega^+(u)$, then there exists $\varepsilon_1 > 0$ such that the rescaling u_r is ε_1 -flat in B_1 in the direction $\nu \in \partial B_1$ for every $r \in (0, 1)$. That is,*

$$(x \cdot \nu - \varepsilon_1)^+ \leq u_r(x) \leq (x \cdot \nu + \varepsilon_1)^+ \quad \text{for every } x \in B_1, \quad \text{and some } \nu \in \partial B_1.$$

Proof. Recall the definition $\partial_{\text{reg}}\Omega^+(u)$, there is a vanishing sequence $r_n \rightarrow 0^+$ such that the blow-up sequence $u_n(x) = \frac{u(r_n x)}{r_n}$ converges uniformly in B_1 to a function $u_0: \mathbb{R}^d \rightarrow \mathbb{R}$ of the form $u_0(x) = (x \cdot \nu)^+$ for some $\nu \in \mathbb{R}^d$. Then there exists $\varepsilon > 0$ such that for n large enough,

$$\|u_n - u_0\|_{L^\infty(B_1)} < \varepsilon,$$

which implies that

$$u_n > 0 \quad \text{in} \quad \{x \cdot \nu > \varepsilon\} \quad \text{and} \quad u_n = 0 \quad \text{in} \quad \{x \cdot \nu < -\varepsilon\}.$$

This gives that u_n is 2ε -flat in B_1 . Namely,

$$(x \cdot \nu - 2\varepsilon)^+ \leq u_n(x) \leq (x \cdot \nu + 2\varepsilon)^+ \quad \text{in} \quad B_1, \quad (7.99)$$

for some $\nu \in \partial B_1$. Let now $r \in (0, 1)$ be arbitrary, then there must exist $n \in \mathbb{N}$ such that $r_{n+1} \leq r \leq r_n$. Thus, there is $\varrho \in (0, 1]$ such that $r = \varrho r_n$. Now since u_n satisfies (7.99), we get that $u_r = u_{\varrho r_n}$ satisfies

$$\left(x \cdot \nu - \frac{2\varepsilon}{\varrho}\right)^+ \leq u_r(x) \leq \left(x \cdot \nu + \frac{2\varepsilon}{\varrho}\right)^+ \quad \text{in} \quad B_1.$$

Set $\varepsilon_1 = \frac{2\varepsilon}{\varrho} > 0$ we obtain for every $r \in (0, 1)$, u_r satisfies

$$(x \cdot \nu - \varepsilon_1)^+ \leq u_r \leq (x \cdot \nu + \varepsilon_1)^+ \quad \text{in} \quad B_1,$$

for some $\nu \in \partial B_1$, as desired. \square

With Lemma 7.3 at hand we obtain the following *improvement of flatness* [Lemma 4.1 in [16]] for viscosity solutions.

Lemma 7.4 (Improvement of flatness). *Let $u \in \mathcal{K}$ be a minimizer of J in Ω . Assume that $0 \in \partial_{\text{reg}} \Omega^+(u)$. Then there are universal constants $\kappa \in (0, 1)$, $\sigma \in (0, 1)$, $C_0 > 0$ and $\varepsilon_0 > 0$ such that if u_r is ε -flat in B_1 in the direction $\nu \in \partial B_1$ for every $r \in (0, 1)$ and $\varepsilon \in (0, \varepsilon_0]$, then $u_{\kappa r}$ is $\sigma\varepsilon$ -flat in B_1 in the direction $\tilde{\nu} \in \partial B_1$ with $|\tilde{\nu} - \nu| \leq C_0\varepsilon$.*

With Lemma 7.4 at hand, we are able to prove the core Lemma in this section:

Lemma 7.5 (Uniqueness of the blow-up). *Let $u \in \mathcal{K}$ be a minimizer of J in Ω . Assume that $0 \in \partial_{\text{reg}} \Omega^+(u)$. Then there is a unique vector $\nu_0 \in \partial B_1 \subset \mathbb{R}^d$ such that*

$$\|u_r - u_0\|_{L^\infty(B_1)} \leq C_1 r^\gamma \quad \text{for every} \quad r \leq \frac{1}{2}, \quad (7.100)$$

where

$$u_0(x) := (\nu_0 \cdot x)^+ \quad \text{for every} \quad x \in \mathbb{R}^d. \quad (7.101)$$

Here C_1, γ are positive constants depending on κ, σ, C_0 and ε_0 (Recall Lemma 7.4).

Proof. Repeated use of Lemma 7.4 via n times gives $u_{\frac{\kappa^n}{2}}$ is $\varepsilon_0 \sigma^n$ -flat in the direction $\nu_n \in \partial B_1$ such that

$$|\nu_n - \nu_{n+1}| \leq C_0 \varepsilon_0 \sigma^n \quad \text{for every} \quad n \in \mathbb{N}.$$

This implies that $\{\nu_n\}$ is a Cauchy sequence so that there is a vector ν_0 such that

$$\nu_0 := \lim_{n \rightarrow \infty} \nu_n \quad \text{and} \quad |\nu_0 - \nu_n| \leq \frac{C_0 \varepsilon_0}{1 - \sigma} \sigma^n.$$

This implies

$$|(x \cdot \nu_0)_+ - u_{\kappa^n/2}| \leq |(x \cdot \nu_0)_+ - (x \cdot \nu_n + \varepsilon_0 \sigma^n)| \leq \left(1 + \frac{C_0}{1 - \sigma}\right) \sigma^n \quad \text{in } B_1. \quad (7.102)$$

Thus we set

$$u_0(x) = (x \cdot \nu_0)^+.$$

Now we verify (7.100). Let $r \in (0, \frac{1}{2}]$ and let $n \in \mathbb{N}$ be such that $\frac{\kappa^{n+1}}{2} < r \leq \frac{\kappa^n}{2}$. It follows that there must exist $\varrho \in (\kappa, 1]$ such that $r = \varrho r_n$. Since $u_{\kappa^n/2}$ satisfies

$$(x \cdot \nu_n - \varepsilon_0 \sigma^n)^+ \leq u_{\kappa^n/2} \leq (x \cdot \nu_n + \varepsilon_0 \sigma^n)^+ \quad \text{in } B_1,$$

it is handy that u_r satisfies

$$\left(x \cdot \nu_n - \frac{\varepsilon_0}{\varrho} \sigma^n\right)^+ \leq u_r \leq \left(x \cdot \nu_n + \frac{\varepsilon_0}{\varrho} \sigma^n\right)^+ \quad \text{in } B_1.$$

Thus, $\|u_n - u_r\|_{L^\infty(B_1)} \leq \frac{\varepsilon_0}{\varrho} \sigma^n \leq \frac{\varepsilon_0}{\kappa} \sigma^n$. It follows from (7.102) that $\|u_r - u_0\|_{L^\infty(B_1)} \leq (1 + \frac{C_0}{1 - \sigma} + \frac{1}{\kappa}) \varepsilon_0 \sigma^n$. Choose γ so that $\kappa^\gamma = \sigma$, and we obtain

$$\|u_r - u_0\|_{L^\infty(B_1)} \leq (2/\kappa)^\gamma \left(1 + \frac{C_0}{1 - \sigma} + \frac{1}{\kappa}\right) \varepsilon_0 r^\gamma.$$

This concludes the proof by letting $C_1 = \left(1 + \frac{C_0}{1 - \sigma} + \frac{1}{\kappa}\right)$. \square

We now introduce the following notion: For any $\varepsilon > 0$, $x^0 \in \Omega$ and $\nu \in \mathbb{R}^d$, we define $\mathcal{C}_\varepsilon^\pm(x^0, \nu)$ to be the cones

$$\mathcal{C}_\varepsilon^\pm(x^0, \nu) := \{x \in \mathbb{R}^d : \pm \nu \cdot (x - x^0) > \varepsilon |x - x^0|\}.$$

In fact, as an application of the uniqueness of blow-ups, we have

Corollary 7.6 ($\partial_{\text{reg}} \Omega^+(u)$ is Lipschitz continuity). *Let $u \in \mathcal{K}$ be a minimizer of J in Ω . Assume that $0 \in \partial_{\text{reg}} \Omega^+(u)$. Then $\partial \Omega^+(u)$ is a Lipschitz graph in a small neighborhood of 0 .*

Proof. The idea of this proof is originally from Proposition 8.6 in [31], we adapt it to our settings.

Step I. In this first step, we show that if u minimizes J in B_1 with $u(0) = 0$ and $0 \in \partial_{\text{reg}} \Omega^+(u)$. Then for every $\varepsilon > 0$ there exists $R > 0$ such that

$$\begin{cases} u > 0 & \text{in } \mathcal{C}_\varepsilon^+(0, \nu_0) \cap B_R, \\ u = 0 & \text{in } \mathcal{C}_\varepsilon^-(0, \nu_0) \cap B_R, \end{cases} \quad (7.103)$$

where $\nu_0 \in \partial B_1$. To this end, let γ and C_1 be defined as in the Lemma 7.5. Then it follows from (7.100) that $u_r \geq (x \cdot \nu_0 - C_1 r^\gamma)^+$ for every $x \in B_1$. Choose C be any positive constant such that $C \geq C_1$, we obtain

$$\{x \in B_1 : x \cdot \nu_0 > C r^\gamma\} \subset \Omega^+(u_r). \quad (7.104)$$

On the other hand, observe that u_r is also a minimizer of J in B_1 , therefore u_r is non-degenerate in B_1 so that

$$\Omega^+(u_r) \cap \{x \in B_1 : x \cdot \nu_0 < -Cr^\gamma\} = \emptyset \quad (7.105)$$

Estimates (7.104) and (7.105) imply (7.103) by taking $R > 0$ such that $CR^\gamma \leq \varepsilon$.

Step II. In this step, we prove that the free boundary near the origin can be expressed by the graph of a function g , precisely, there exists a $\delta > 0$ such that

$$(B'_\delta \times (-\delta, \delta)) \cap \partial\Omega^+(u) = \{(x', t) \in B'_\delta \times (-\delta, \delta) : t = g(x')\}.$$

Without loss of generality, we assume that $\nu_0 = e_d$ and therefore $u_0(x) = x_d^+$ is the blow-up limit in 0. Thus $\Omega^+(u_0) = \{x_d > 0\}$. Let now $\varepsilon \in (0, 1)$ and R be as in (7.103), it follows that

$$\begin{cases} u > 0 & \text{in } \mathcal{C}_\varepsilon^+(0, e_d) \cap B_R, \\ u = 0 & \text{in } \mathcal{C}_\varepsilon^-(0, e_d) \cap B_R, \end{cases}$$

where $\mathcal{C}_\varepsilon^\pm$ is the cone $\{x \in \mathbb{R}^d : \pm x_d > \varepsilon|x|\}$. Let $x'_0 \in B'_\delta$ where $\delta \leq R\sqrt{1 - \varepsilon^2}$, it follows that the segment $\{(x'_0, t) : t > \varepsilon R\}$ is contained in $\{u > 0\}$ and the segment $\{(x'_0, t) : t < -\varepsilon R\}$ is contained in $\{u = 0\}$. Therefore, $g(x'_0) := \inf\{t : u(x'_0, t) > 0\}$ is well defined and $x^0 := (x'_0, g(x'_0)) \in \partial\Omega^+(u) \cap B_R$. Moreover, one has

$$-\varepsilon|x'_0| \leq g(x'_0) \leq \varepsilon|x'_0|,$$

which implies that $|x^0| \leq |x'_0|\sqrt{1 + \varepsilon^2} \leq \sqrt{2}\delta$. In order to finish the claim, it suffices to show that if $\delta > 0$ is small enough, then

$$\begin{cases} u > 0 & \text{in } \mathcal{C}_{2\varepsilon}^+(x^0, e_d) \cap B_R(x^0), \\ u = 0 & \text{in } \mathcal{C}_{2\varepsilon}^-(x^0, e_d) \cap B_R(x^0). \end{cases} \quad (7.106)$$

Indeed, as a direct consequence of (7.106), one can easily deduce that

$$\begin{aligned} (B'_\delta \times (-\delta, \delta)) \cap \{u > 0\} &= \{(x', t) \in B'_\delta \times (-\delta, \delta) : t > g(x')\}, \\ (B'_\delta \times (-\delta, \delta)) \cap \{u = 0\} &= \{(x', t) \in B'_\delta \times (-\delta, \delta) : t \leq g(x')\}. \end{aligned}$$

We now prove (7.106) to conclude this step. Applying (7.106) for the free boundary point x^0 , we have

$$\begin{cases} u > 0 & \text{in } \mathcal{C}_\varepsilon^+(x^0, \nu_{x^0}) \cap B_R(x^0), \\ u = 0 & \text{in } \mathcal{C}_\varepsilon^-(x^0, \nu_{x^0}) \cap B_R(x^0). \end{cases}$$

In view of (7.106), we only need to show $\mathcal{C}_{2\varepsilon}^\pm(x^0, e_d) \subset \mathcal{C}_\varepsilon^\pm(x^0, \nu_{x^0})$. Since u is a Lipschitz function with Lipschitz coefficient L in B_1 and $0, x^0 \in \partial\Omega^+(u) \cap B_R$, let $\alpha := \frac{\gamma}{1+\gamma}$ and $r := |x^0|^{1-\alpha}$,

$$|u_r - u_{x^0, r}| = \frac{1}{r}|u(rx) - u(x^0 + rx)| \leq \frac{L}{r}|x^0| = L|x^0|^\alpha,$$

for every $x \in B_1$. This implies $\|u_r - u_{x^0, r}\|_{L^\infty(B_1)} \leq L|x^0|^\alpha$. On the other hand, if we denote u_0 and u_{x^0} the unique blow-up limit of u at 0 and x^0 respectively. It follows from (7.106) that

$$\|u_{x^0} - u_{x^0, r}\|_{L^\infty(B_1)} \leq C_1 r^\gamma \quad \text{and} \quad \|u_r - u_0\|_{L^\infty(B_1)} \leq C_1 r^\gamma.$$

Choose R such that $(2R)^{1-\alpha} \leq \frac{1}{2}$, we have by $r^\gamma = |x^0|^\alpha$ that

$$\|u_{x^0} - u_0\|_{L^\infty(B_1)} \leq (L + 2C_1)|x^0|^\alpha.$$

This implies $|\nu_0 - e_d| \leq C|x^0|^\alpha$ by a direct calculation. Let $x \in \mathcal{C}_{2\varepsilon}^\pm(x^0, e_d)$. Then,

$$\begin{aligned} \nu_{x^0} \cdot (x - x^0) &= e_d \cdot (x - x^0) + (\nu_{x^0} - e_d) \cdot (x - x^0) \\ &> 2\varepsilon|x - x^0| - C(\sqrt{2}\delta)^\alpha|x - x^0| > \varepsilon|x - x^0|, \end{aligned}$$

where we choose δ such that $C(\sqrt{2}\delta)^\alpha \leq \varepsilon$. This finishes $\mathcal{C}_{2\varepsilon}^\pm(x^0, e_d) \subset \mathcal{C}_\varepsilon^\pm(x^0, \nu_{x^0})$ and thus (7.106) is concluded.

Step III. We now prove that $g: B'_\delta \rightarrow \mathbb{R}$ is Lipschitz continuous on B'_δ . Let $x'_1, x'_2 \in B'_\delta$ where $x^1 = (x'_1, g(x'_1))$ and $x^2 = (x'_2, g(x'_2))$. Since $x^1 \notin \mathcal{C}_{2\varepsilon}^+(x^2, e_d)$, we have that

$$g(x'_1) - g(x'_2) = (x^1 - x^2) \cdot e_d \leq 2\varepsilon|x'_1 - x'_2| + 2\varepsilon|g(x'_1) - g(x'_2)|.$$

A same inequality can be derived since $x^2 \notin \mathcal{C}_{2\varepsilon}^+(x^1, e_d)$. Two estimates give $(1 - 2\varepsilon)|g(x'_1) - g(x'_2)| \leq 2\varepsilon|x'_1 - x'_2|$. Choosing $\varepsilon \leq \frac{1}{4}$, we obtain

$$|g(x'_1) - g(x'_2)| \leq 4\varepsilon|x'_1 - x'_2|,$$

which concludes the proof of the Lipschitz continuity of g . \square

We are now in a position to prove the regularity of the free boundary near regular points.

Proposition 7.7 (Regularity of $\partial_{\text{reg}}\Omega^+(u)$). *Let $u \in \mathcal{K}$ be a minimizer of J in $\Omega \subset \mathbb{R}^d$. Assume that $0 \in \partial_{\text{reg}}\Omega^+(u)$, then there exists a small neighborhood, the size of which depends on d , $[Q]_{C^\beta; \Omega}$, F_0 and M_0 so that $\partial_{\text{reg}}\Omega^+(u)$ is $C^{1, \gamma}$ regularity in that neighborhood for $\gamma \in (0, \beta)$.*

Proof. Consider the blow-up sequence $u_k(x) := u_{\delta_k}(x) = \frac{u(\delta_k x)}{\delta_k}$ with $\delta_k \rightarrow 0$ as $k \rightarrow \infty$. It is easy to see that for every k , u_k is a solution to

$$\Delta u_k = \tilde{f}_k(x) \quad \text{in} \quad B_1^+(u_k), \quad |\nabla u_k| = Q_k(x) \quad \text{on} \quad \partial B_1^+(u_k), \quad (7.107)$$

where $\tilde{f}_k(x) = \delta_k \tilde{f}(\delta_k x)$ and $Q_k(x) = Q(\delta_k x)$. It follows that there is a positive constant $\bar{\varepsilon}$ such that the following does hold:

$$|\tilde{f}_k(x)| = |\delta_k \tilde{f}(\delta_k x)| = \delta_k \|\tilde{f}\|_{L^\infty} \leq \bar{\varepsilon},$$

and

$$|Q_k(x) - 1| = |Q(\delta_k x) - Q(0)| \leq \delta_k^\beta [Q]_\beta \leq \bar{\varepsilon}.$$

Thus, using Proposition 4.5, up to extracting a subsequence: $u_k \rightarrow u_0$ locally and uniformly in \mathbb{R}^d and $\partial\Omega^+(u_k) \rightarrow \partial\Omega^+(u_0)$ locally in the Hausdorff distance in \mathbb{R}^d for a globally defined function

$u_0: \mathbb{R}^d \rightarrow \mathbb{R}$. Moreover, every blow-up limit u_0 is a solution to $\Delta u_0 = 0$ in $B_1^+(u_0)$ and $|\nabla u_0| = 1$ on $\partial B_1^+(u_0)$. It follows from Corollary 7.6 that $\partial\Omega^+(u)$ is a Lipschitz graph in a small neighborhood of 0. We also see from (1)-(2) above that $\partial\Omega^+(u_0)$ is Lipschitz continuous. Thus, u_0 is a so-called one-plane solution, namely (up to rotations), $u_0(x) = x_d^+$. Consequently, we conclude that for all k large enough, u_k is $\bar{\varepsilon}$ -flat in B_1 , i.e.

$$(x_d - \bar{\varepsilon})^+ \leq u_k(x) \leq (x_d + \bar{\varepsilon})^+, \quad \text{in } B_1.$$

Thus, u_k satisfies the flatness condition required by Theorem 1.1 in [16]. Moreover, \tilde{f}_k and Q_k also satisfy the requirements of Theorem 1.1 in [16]. Combining these facts, we conclude from Theorem 1.1 in [16] that $\partial\Omega^+(u)$ is $C^{1,\gamma}$ in $B_{1/2}$. This finishes the proof. \square

8. DIMENSION OF THE SINGULAR SET

This concluding section dedicates to investigate the structure of the singular set. We will prove Theorem 1.3 for our semilinear case. Our results discover that the non-linear right-hand side in our settings does not affect the structure of singularities and our argument follows by "dimension reduction" process. We begin by giving the definition of d^* .

Definition 8.1 (Definition of d^* , compare to [35]). *We define d^* to be the smallest dimension d so that there exists a function $z: \mathbb{R}^d \rightarrow \mathbb{R}$ such that*

- (1) z is a non-negative one-homogeneous absolute minimum of J_0 given in (4.58);
- (2) The free boundary $\partial\{z > 0\}$ is not a $(d-1)$ -dimensional C^1 -regular surface in \mathbb{R}^d .

In [13], it was proved that $\partial_{\text{sing}}\Omega^+(u) = \emptyset$ when $d = 2$. We present here a different proof, which depends mainly on the homogeneity of minimizers and this is completely different from [13].

Proposition 8.1. *Let $u \in \mathcal{K}$ be a minimizer of J in $\Omega \subset \mathbb{R}^2$, then $\partial\Omega^+(u) = \partial_{\text{reg}}\Omega^+(u)$.*

Proof. Let $u \in \mathcal{K}$ be a minimizer of J in $\Omega \subset \mathbb{R}^2$. Then it follows from Proposition 4.5 (4) that $u_0: \mathbb{R}^2 \rightarrow \mathbb{R}$ is a global minimizer of $J_0(v)$ (recall (4.58) in \mathbb{R}^2). We write u_0 in polar coordinates $u_0(\rho, \theta) = \rho c(\theta)$. Since c is continuous, we have that $\{c > 0\}$ is open and so it is a countable union of disjoint arcs. Based on the fact that local minimizers of $J_0(v)$ cannot have isolated zeros, we have that $\{c > 0\} \neq \mathbb{S}^1$. It follows from Proposition 4.5 and Proposition 6.1 (2) that up to a rotation, the trace c is a solution to

$$c''(\theta) + c(\theta) = 0 \quad \text{in } (0, \pi), \quad c(0) = c(\pi) = 0.$$

It follows that $c(\theta)$ is a multiple of $\sin \theta$ on $(0, \pi)$. Thus, $\{c > 0\}$ is a union of disjoint arcs, each one of length π . Since $0 \in \partial\Omega^+(u_0)$, thanks (4.59), one has $|B_1 \cap \{u_0 > 0\}| < |B_1| = \pi$. The homogeneity of u_0 then implies that $\mathcal{H}^1(\{c > 0\}) < 2\pi$. Therefore, $\{c > 0\}$ is an arc of length π and u_0 is of the form $u_0(x) = a(x \cdot \nu)^+$ for some $a > 0$. Thanks to Proposition 4.5 (5), we obtain $a = Q(0)$. Consequently, $u_0(x) = Q(0)(x \cdot \nu)^+$ and it follows from the definition of $\partial_{\text{reg}}\Omega^+(u)$ that $\partial\Omega^+(u) = \partial_{\text{reg}}\Omega^+(u)$. Thus by Definition 8.1, $d^* \geq 3$. \square

When $d < d^*$, we have the following fact

Proposition 8.2. *Let $u \in \mathcal{K}$ be a minimizer of J in $\Omega \subset \mathbb{R}^d$ with $2 \leq d < d^*$, then $\partial_{\text{sing}}\Omega^+(u) = \emptyset$ and $\partial\Omega^+(u) = \partial_{\text{reg}}\Omega^+(u)$.*

Proof. Let $x^0 \in \partial\Omega^+(u)$ be an arbitrary free boundary point and let $r_n \rightarrow 0$ be a infinitesimal sequence. Then it follows that the blow-up sequence u_n converges uniformly to a blow-up limit u_0 . Thanks to Proposition 4.5 (4), u_0 is a global minimizer of J_0 in \mathbb{R}^d where J_0 is defined in (4.58). It then follows from the definition of d^* that $\partial_{\text{sing}}\Omega^+(u_0) = \emptyset$. By the definition of $\partial_{\text{reg}}\Omega^+(u)$, we obtain that every blow-up u_{00} of u_0 is of the form $Q(0)(x \cdot \nu)^+$ for some $\nu \in \partial B_1$. In particular, it holds for every blow-up limit in 0. Since (by Proposition 6.1 (2)) u_0 is one-homogeneous function, we have

$$u_{00}(x) = \lim_{n \rightarrow \infty} \frac{u_0(r_n x)}{r_n} = \lim_{n \rightarrow \infty} u_0(x) = u_0(x),$$

which implies that the blow-up limit u_{00} of u_0 is indeed u_0 itself, and so

$$u_0(x) = Q(0)(x \cdot \nu)^+, \quad \text{for some } \nu \in B_1.$$

Thus, for n large enough, we have

$$\|u_n(x) - Q(0)(x \cdot \nu)^+\|_{L^\infty(B_1)} \leq \varepsilon, \quad \text{for some } \nu \in B_1.$$

Set $y = x^0 + r_n x$ and we have from the definition of u_n that

$$\left\| \frac{u(y)}{r_n} - \frac{Q(0)}{r_n}((y - x^0) \cdot \nu)^+ \right\|_{L^\infty(B_{r_n}(x^0))} \leq \varepsilon, \quad \text{for some } \nu \in B_1.$$

This implies that there exists $r \in (0, R)$ such that

$$\|u(y) - Q(0)((y - x^0) \cdot \nu)^+\|_{L^\infty(B_r(x^0))} \leq \varepsilon r, \quad \text{for some } \nu \in B_1. \quad (8.108)$$

In via of (8.108), there must exist $\bar{\varepsilon} > 0$ such that u is $\bar{\varepsilon}$ -flat in B_1 . Moreover, since u minimizes J in B_1 , it follows from Proposition 7.1 that u is a viscosity solution in $B_1 \subset \mathbb{R}^d$. Thanks to Proposition 7.7, we know that $\partial\Omega^+(u)$ is $C^{1,\gamma}$ in a small neighborhood of $x^0 \in \partial\Omega^+(u)$. Thus, $x^0 \in \partial_{\text{reg}}\Omega^+(u)$. Since x^0 is arbitrary, we conclude that $\partial_{\text{sing}}\Omega^+(u) = \emptyset$ and this concludes the proof. \square

We now investigate the case $d = d^*$. Let $u \in \mathcal{K}$ be a minimizer of J in $\Omega \subset \mathbb{R}^{d^*}$ (Proposition 8.6 below). To this end, we begin by studying the singular set of a one-homogeneous function in \mathbb{R}^d . In [35], Weiss showed that the singular set of one-homogeneous function is a cone in \mathbb{R}^d in homogeneous case. We follow his approach and extend the same result to our case.

Lemma 8.3. *Let $u \in \mathcal{K}$ be a minimizer of J in $\Omega \subset \mathbb{R}^d$. Then the graph of any blow-up limit u_0 is a minimal cone with vertex at $0 \in \mathbb{R}^d$.*

Proof. It suffices to prove the following two facts:

- (1) Every blow-up limit u_0 of u is a non-negative one-homogeneous global minimizer of J_0 in \mathbb{R}^d .

(2) For any open and bounded set $D \subset \mathbb{R}^d$ satisfying $\bar{D} \cap (\partial\Omega^+(u_0) \setminus \partial_{\text{red}}\Omega^+(u_0)) = \emptyset$,

$$\text{Per}(\Omega^+(u_0), D) \leq \text{Per}(F, D),$$

for any $F \subset \Omega^+(u_0)$ so that $\partial F \cap D$ is a C^2 surface and $\text{supp}(\chi_F - \chi_{\{u_0 > 0\}}) \subset D$. Here $\text{Per}(E)$ denotes the perimeter of a given set $E \subset \mathbb{R}^d$.

Notice that (1) is a direct consequence of Proposition 4.5 (5) and Proposition 6.1 (2). To prove the above mentioned second property, since $Q(x) \in C^{0,\beta}(\Omega)$, it follows from Theorem 5.5 in [1] that u_0 is a weak solution. Then a same argument as in Theorem 2.8 in [35] gives the desired result. \square

We now consider a blow-up limit u_{00} of u_0 at $x^0 \neq 0 \in \partial\Omega^+(u_0)$, the following fact was first proved by Weiss in [35] for minimizers of J_0 , we adapt it to our settings.

Lemma 8.4. *Let $u \in \mathcal{K}$ be a minimizer of J in $\Omega \subset \mathbb{R}^d$ and let u_0 be any blow-up limits of u at any free boundary points. Then the graph of u_{00} is a minimal cone with vertex at 0 and u_{00} is constant in the direction x^0 . That is, $u_{00}(x) = u_{00}(x + tx^0)$ for $x \in \mathbb{R}^d$ and $t \in \mathbb{R}$. Moreover, $\{t \in \mathbb{R} : tx^0\} \subset \partial_{\text{sing}}\Omega^+(u)$.*

Proof. The proof is similar to Lemma 3.1 in [35]. Notice that u_0 is a global minimizer of J_0 in \mathbb{R}^d , therefore one has

$$\begin{aligned} u_{00}(x + tx^0) &= \lim_{n \rightarrow \infty} \frac{u_0(x^0 + r_n(x + tx^0))}{r_n} \\ &= \lim_{n \rightarrow \infty} \left(\frac{1 + tr_n}{r_n} \right) u_0 \left(x^0 + \frac{r_n}{1 + tr_n} x \right), \end{aligned}$$

where we have used the homogeneity of u_0 in the second identity. We now claim that

$$\lim_{n \rightarrow \infty} \left(\frac{1 + tr_n}{r_n} \right) u_0 \left(x^0 + \frac{r_n}{1 + tr_n} x \right) = \lim_{n \rightarrow \infty} \frac{u_0(x^0 + r_n x)}{r_n}. \quad (8.109)$$

Since u_0 is a global minimizer of J_0 , thanks to Corollary 3.3 in [1], u_0 is a Lipschitz function with Lipschitz coefficient $L_0 := \|\nabla u_0\|_{L^\infty(B_1)}$. Then a straightforward estimate gives

$$\begin{aligned} &\left| \frac{1 + tr_n}{r_n} u_0 \left(x^0 + \frac{r_n}{1 + tr_n} x \right) - \frac{1}{r_n} u_0(x^0 + r_n x) \right| \\ &\leq t \left| u_0 \left(x^0 + \frac{r_n}{1 + tr_n} x \right) - u_0(x^0) \right| + \frac{1}{r_n} \left| u_0 \left(x^0 + \frac{r_n}{1 + tr_n} x \right) - u_0(x^0 + r_n x) \right| \\ &\leq \frac{2tr_n L_0 |x|}{1 + tr_n} \rightarrow 0 \quad \text{as } n \rightarrow \infty. \end{aligned}$$

Then the desired result follows from (8.109). \square

In via of a same argument as in Lemma 3.2 in [35], one has the following property for u_{00} given in Lemma 8.4.

Lemma 8.5. *The graph of $u_{00}|_{\mathbb{R}^{d-1}} = \bar{u}$ is a minimal cone and $0 \in \partial_{\text{sing}}\Omega^+(\bar{u})$.*

As a direct corollary of Lemma 8.4 and Lemma 8.5, one has

Proposition 8.6. *Let $u \in \mathcal{K}$ be minimizer of J in $\Omega \subset \mathbb{R}^{d^*}$. Then $\partial_{\text{sing}}\Omega^+(u) \setminus \{0\} = \emptyset$. In particular, this means $\dim_{\mathcal{H}} \partial_{\text{sing}}\Omega^+(u) = 0$.*

Proof. Step I. Suppose there is a sequence of singular free boundary points $x_n \in \partial_{\text{sing}}\Omega^+(u)$ converging to some point x^0 as $n \rightarrow \infty$, and we claim $x^0 \in \partial_{\text{sing}}\Omega^+(u)$. Let us consider the blow-up sequence u_n at x^0 and we denote by u_0 the blow-up limit of u_n as $n \rightarrow \infty$. On the one hand, the uniform convergence of u_n implies that $u_0(x^0) = 0$. On the other hand, since for each n , u_n minimizes J , we may infer from Corollary 3.5 that for every $r > 0$ small

$$\begin{aligned} \|u_0\|_{L^\infty(B_r(x^0))} &\geq \lim_{n \rightarrow \infty} (\|u_n\|_{L^\infty(B_r(x^0))} - \|u_n - u_0\|_{L^\infty(B_r(x^0))}) \\ &\geq \liminf_{n \rightarrow \infty} \|u_n\|_{L^\infty(B_{r/2}(x^0))} \geq \eta r, \end{aligned}$$

which gives $x^0 \in \partial\Omega^+(u_0)$. If $x^0 \in \partial_{\text{reg}}\Omega^+(u_0)$, so that every blow-up limit u_{00} of u_0 at x^0 is of the form $Q(0)((x - x^0) \cdot \nu)^+$ for some $\nu \in \partial B_1$. Moreover, there is a sequence $r_n \rightarrow 0$ so that

$$\lim_{n \rightarrow \infty} \left\| \frac{u_0(x^0 + r_n(x - x^0))}{r_n} - Q(0)((x - x^0) \cdot \nu)^+ \right\|_{L^\infty(B_1)} = 0.$$

Let $y := x^0 + r_n(x - x^0)$, one has

$$0 = \lim_{n \rightarrow \infty} \frac{1}{r_n} \|u_0(y) - Q_0((y - x^0) \cdot \nu)^+\|_{L^\infty(B_{r_n}(x^0))}.$$

In particular, there exists $r \in (0, R)$ so that

$$\|u_0(\cdot) - Q_0((\cdot - x^0) \cdot \nu)^+\|_{L^\infty(B_r(x^0))} \leq \frac{\varepsilon r}{3}.$$

Since u_0 is a continuous function, we have for sufficiently large n ,

$$\|u_0(\cdot) - Q(0)((\cdot - x_n) \cdot \nu)^+\|_{L^\infty(B_r(x_n))} \leq \frac{2\varepsilon r}{3}$$

Since u_n converges to u_0 uniformly, we get for sufficiently large n ,

$$\|u_n(\cdot) - Q(0)((\cdot - x_n) \cdot \nu)^+\|_{L^\infty(B_r(x_n))} \leq \varepsilon r.$$

This implies that there exists $\bar{\varepsilon} > 0$ so that u_n is $\bar{\varepsilon}$ -flat in $B_r(x_n)$. Moreover, u_n is a viscosity solution. Thanks to the regularity results we have $\partial\Omega^+(u_n)$ is $C^{1,\gamma}$ in a small neighborhood of x_n and this implies that $x_n \in \partial_{\text{reg}}\Omega^+(u_n)$, which yields a contradiction with the initial assumption.

Step II. The minimality of u and Proposition 6.1 (2) imply that $u_0: \mathbb{R}^{d^*} \rightarrow \mathbb{R}$ is a non-negative global minimizer of J_0 in \mathbb{R}^{d^*} . Then it follows from Lemma 8.4, Lemma 8.5 and similar argument in [35] that

$$\partial_{\text{sing}}\Omega^+(u_0) \setminus \{0\} = \emptyset. \tag{8.110}$$

Let $x_n \in \partial_{\text{sing}}\Omega^+(u)$ and let $r_n := |x_n - x^0|$. Notice now that for every $n > 0$ the point $\xi_n := \frac{x_n - x^0}{r_n} \in \partial B_1$ is a singular point for u_n . Up to extracting a subsequence, we may assume that ξ_n converges to a point $\xi_0 \in \partial B_1$. It follows from Step I that $\xi_0 \in \partial_{\text{sing}}\Omega^+(u_0)$. This yields a contradiction to (8.110). \square

Proposition 8.7. *Let $u \in \mathcal{K}$ be a minimizer of J in $\Omega \subset \mathbb{R}^d$ for $d > d^*$, then*

$$\dim_{\mathcal{H}}(\partial_{\text{sing}}\Omega^+(u)) \leq d - d^*. \quad (8.111)$$

Proof. Let $d > d^*$ and let $s > 0$ be fixed. We will prove

$$\mathcal{H}^{d-d^*+s}(\partial_{\text{sing}}\Omega^+(u)) = 0. \quad (8.112)$$

Suppose that this is not the case and $\mathcal{H}^{d-d^*+s}(\partial_{\text{sing}}\Omega^+(u)) > 0$. Then it follows that there is a point $x^0 \in \partial_{\text{sing}}\Omega^+(u)$ and a sequence $r_n \rightarrow 0$ such that

$$\mathcal{H}^{d-d^*+s}(\partial_{\text{sing}}\Omega^+(u) \cap B_{r_n}(x^0)) \geq \varepsilon r_n^{d-d^*+s}.$$

Choosing u_n a blow-up sequence, we get

$$\mathcal{H}^{d-d^*+s}(\partial_{\text{sing}}\Omega^+(u_n) \cap B_1) \geq \varepsilon.$$

Up to extracting a subsequence, u_n converges to a blow-up limit u_0 and u_0 is a one-homogeneous global minimizer of J_0 in \mathbb{R}^d . Consequently,

$$\mathcal{H}^{d-d^*+s}(\partial_{\text{sing}}\Omega^+(u_0) \cap B_1) \geq \varepsilon,$$

which yields a contradiction to Federer dimension reduction Theorem [19] for one-homogeneous global minimizer of J_0 in \mathbb{R}^d . \square

REFERENCES

- [1] H. W. Alt, L. A. Caffarelli, Existence and regularity for a minimum problem with free boundary, *J. Reine Angew. Math.*, **325**, 105-144, (1981).
- [2] H. W. Alt, L. A. Caffarelli, A. Friedman, A free boundary problem for quasilinear elliptic equations, *Ann. Scuola Norm. Sup. Pisa Cl. Sci.*, **11**, 1-44, (1984).
- [3] L. A. Caffarelli, A Harnack inequality approach to the regularity of free boundaries. I. Lipschitz free boundaries are $C^{1,\alpha}$, *Rev. Mat. Iberoamericana.*, **3**, 139-162, (1987).
- [4] L. A. Caffarelli, A Harnack inequality approach to the regularity of free boundaries. II. Flat free boundaries are Lipschitz, *Comm. Pure Appl. Math.*, **42**, 55-78, (1989).
- [5] L. A. Caffarelli, A Harnack inequality to the regularity of free boundaries. Part III: existence theory, compactness, and dependence on X , *Ann. Scuola Norm. Sup. Pisa*, **15**(4), 583-602, (1988).
- [6] H. W. Alt, L. A. Caffarelli, A. Friedman, Variational problems with two phases and their free boundaries, *Trans. Amer. Math. Soc.*, **282**(2), 431-461, (1984).
- [7] L. A. Caffarelli, H. Shahgholian, K. Yeressian, A minimization problem with free boundary related to a cooperative system, *Duke. Math. J.*, **167**(10), 1825-1882, (2018).
- [8] M. G. Crandall, H. Ishii, P. L. Lions, User's guide to viscosity solutions of second order partial differential equations, *Bull. Amer. Math. Soc.*, **27**, 1-67, (1992).
- [9] L. A. Caffarelli, D. S. Jerison, C. E. Kenig, Global energy minimizers for free boundary problems and full regularity in three diemnsions, *Contemp. Math. Amer. Math. Soc. Providence.*, **350**, 83-97, (2004).
- [10] A. Constantin, W. Strauss, Exact steady periodic water waves with vorticity, *Comm. Pure Appl. Math.*, **57**(4), 481-527, (2004).
- [11] J. F. Cheng, L. L. Du, Y. F. Wang, Two-dimensional impinging jets in hydrodynamic rotational flows, *Ann. Inst. H. Poincaré Anal. Non Linéaire.*, **34**, 1355-1386, (2017).

- [12] J. F. Cheng, L. L. Du, Q. Zhang, Two-dimensional cavity flow in an infinitely long channel with non-zero vorticity, *J. Differential Equations.*, **263**, 4126-4155, (2017).
- [13] J. F. Cheng, L. L. Du, A free boundary problem for semilinear elliptic equation and its applications, *arXiv:2006.02269v1*.
- [14] D. Danielli, A Petrosyan, A minimum problem with free boundary for a degenerate quasilinear operator, *Calc. Var. PDE.*, **23**(1), 97-124, (2005).
- [15] D. De Silva, D. Jerison, A singular energy minimizing free boundary, *J. Reine Angew. Math.*, **635**, 1-21, (2009).
- [16] D. De Silva, Free boundary regularity for a problem with right hand side, *Interfaces Free Bound.*, **13**, 223-238, (2011).
- [17] L. L. Du, C. L. Yang, The free boundary of steady axisymmetric inviscid flow with vorticity II: near the non-degenerate points, *arXiv:2310.09477*.
- [18] L. C. Evans, Measure Theory and Fine Properties of Functions, *Boca Raton*, CRC Press, 2nd revised, edition (2015).
- [19] H. Federer, Geometric Measure Theory, *Springer Berlin, Heidelberg*, (1996).
- [20] B. Gustafsson, H. Shahgholian, Existence and geometric properties of solutions of a free boundary problem in potential theory, *J. Reine Angew. Math.*, **473**, 137-179, (1996).
- [21] D. Gilbarg, N. S. Trudinger, Elliptic Partial Differential Equations of Second Order, *Springer-Verlag*, Berlin, (1998).
- [22] D. S. Jerison, O. Savin, Some remarks on stability of cones for the one-phase free boundary problem, *Geom. Funct. Anal.*, **25**, 1240-1257, (2015).
- [23] R. Leitão, E V. Teixeira, Regularity and geometric estimates for minima of discontinuous functionals, *Rev. Mat. Iberoam.*, **31**, 69-108, (2015).
- [24] C. B. Morrey, Multiple Integrals in The Calculus of Variations, *Classics in Mathematics.*, Springer, Berlin, (2008).
- [25] D. Mazzoleni, S. Terracini, B. Velichkov, Regularity of the optimal sets for some spectral functionals, *Geom. Funct. Anal.*, **27**, 373-426 (2017).
- [26] D. Mazzoleni, S. Terracini, B. Velichkov, Regularity of the free boundary for the vectorial Bernoulli problem, *Anal. PDE.*, **13**, 741-764, (2020).
- [27] E. R. Reifenberg, An epiperimetric inequality related to the analytic of minimal surfaces, *Ann. of Math.*, **80**, 1-14, (1964).
- [28] E. Russ, B. Tery, B. Velichkov, Existence and regularity of optimal shapes for elliptic operators with drift, *Calc. Var. PDE.*, **58**, 199, (2019).
- [29] L. Spolaor, B. Velichkov, An epiperimetric inequality for the regularity of some free boundary problems: the 2-dimensional case, *Comm. Pure. Appl. Math.*, **72**(2), 375-421, (2019).
- [30] N. S. Trudinger, On Harnack type inequalities and their application to quasilinear elliptic equations, *Comm. Pure Appl. Math.*, **20**, 721-747, (1967).
- [31] B. Velichkov, Regularity of the One-phase Free Boundaries, *Lect. Notes*(2023). <https://doi.org/10.1007/978-3-031-13238-4>.
- [32] E. Vărvărucă, On some properties of the traveling water waves with vorticity, *Siam. J. Math. Anal.*, **39**(5), 1686-1692, (2008).
- [33] E. Vărvărucă, G. Weiss, The stokes conjecture for waves with vorticity, *Ann. Inst. H. Poincaré.*, **29**, 861-885, (2012).
- [34] G. Weiss, G. H. Zhang, A free boundary approach to two-dimensional steady capillary gravity water waves, *Arch. Ration. Mech. Anal.*, **203**, 747-768, (2012).
- [35] G. Weiss, Partial regularity for a minimum problem with free boundary, *J. Geom. Anal.*, **9**(2), 317-326, (1999).

COLLEGE OF MATHEMATICS AND STATISTICS, SHENZHEN UNIVERSITY, SHENZHEN 518061, P. R. CHINA.

Email address: `dulili@szu.edu.cn`

DEPARTMENT OF MATHEMATICS, SICHUAN UNIVERSITY, CHENGDU 610064, P. R. CHINA.

Email address: `yang_chunlei@stu.scu.edu.cn`