# Maximum k- vs. $\ell$ -colourings of graphs<sup>\*</sup>

Tamio-Vesa Nakajima University of Oxford tamio-vesa.nakajima@cs.ox.ac.uk Stanislav Živný University of Oxford standa.zivny@cs.ox.ac.uk

13th February 2024

#### Abstract

We present polynomial-time SDP-based algorithms for the following problem: For fixed  $k \leq \ell$ , given a real number  $\varepsilon > 0$  and a graph G that admits a k-colouring with a  $\rho$ -fraction of the edges coloured properly, it returns an  $\ell$ -colouring of G with an  $(\alpha \rho - \varepsilon)$ -fraction of the edges coloured properly in polynomial time in G and  $1/\varepsilon$ . Our algorithms are based on the algorithms of Frieze and Jerrum [Algorithmica'97] and of Karger, Motwani and Sudan [JACM'98].

For k = 2,  $\ell = 3$ , our algorithm achieves an approximation ratio  $\alpha = 1$ , which is the best possible. When k is fixed and  $\ell$  grows large, our algorithm achieves an approximation ratio of  $\alpha = 1 - o(1/\ell)$ . When k,  $\ell$  are both large, our algorithm achieves an approximation ratio of  $\alpha = 1 - 1/\ell + 2 \ln \ell/k\ell - o(\ln \ell/k\ell) - O(1/k^2)$ ; if we fix  $d = \ell - k$  and allow k,  $\ell$  to grow large, this is  $\alpha = 1 - 1/\ell + 2 \ln \ell/k\ell - o(\ln \ell/k\ell)$ .

By extending the results of Khot, Kindler, Mossel and O'Donnell [SICOMP'07] to the promise setting, we show that for large k and  $\ell$ , assuming Khot's Unique Games Conjecture (UGC), it is NP-hard to achieve an approximation ratio  $\alpha$  greater than  $1 - 1/\ell + 2 \ln \ell/k\ell + o(\ln \ell/k\ell)$ , provided that  $\ell$  is bounded by a function that is  $o(\exp(\sqrt[3]{k}))$ . For the case where  $d = \ell - k$  is fixed, this bound matches the performance of our algorithm up to  $o(\ln \ell/k\ell)$ . Furthermore, by extending the results of Guruswami and Sinop [ToC'13] to the promise setting, we prove that it is NP-hard to achieve an approximation ratio greater than  $1 - 1/\ell + 8 \ln \ell/k\ell + o(\ln \ell/k\ell)$ , provided again that  $\ell$  is bounded as before (but this time without assuming the UGC).

## 1 Introduction

The three most studied objectives in approximation algorithms are to maximise the number of satisfied constraints, to minimise the number of unsatisfied constraints, and to find a solution that satisfies a  $(1 - f(\varepsilon))$ fraction of the constraints given an instance in which a  $(1 - \varepsilon)$ -fraction of the constraints is satisfiable, where f is some function satisfying  $f \rightarrow 0$  as  $\varepsilon \rightarrow 0$  and not depending on the input size.<sup>1</sup> All three objectives are examples of a *quantitative* approximation. Another approach to approximation is a *qualitative* approximation, which insists on satisfying *all* constraints but possibly in a weaker form. A canonical example of this is the *approximate graph colouring* problem [G]76]: Given a *k*-colourable graph, find an  $\ell$ -colouring, where  $k \leq \ell$ . In this work, we shall combine the two approaches. In particular, we are interested in the following type of problems: Given a graph in which a large fraction of edges can be properly *k*-coloured, can we

<sup>\*</sup>This work was supported by UKRI EP/X024431/1 and by a Clarendon Fund Scholarship. For the purpose of Open Access, the authors have applied a CC BY public copyright licence to any Author Accepted Manuscript version arising from this submission. All data is provided in full in the results section of this paper.

<sup>&</sup>lt;sup>1</sup>This notion of tractability, coined *robust solvability*, was introduced by Zwick [Zwi98].

find an  $\ell$ -colouring of it with a good fraction of the edges properly coloured? Our main result is an efficient algorithm for this problem and showing its optimality in many cases.

Given a graph G = (V, E) and  $k \in \mathbb{N}$ , a k-colouring of G is an assignment  $c : V \to \{1, \dots, k\}$  of colours to the vertices of G. The value  $\rho_k(c)$  of a k-colouring c is the fraction of properly coloured edges:

$$\rho_k(c) = \frac{|\{(u,v) \in E \mid c(u) \neq c(v)\}|}{|E|}$$

A *k*-colouring *c* is called *proper* if  $\rho_k(c) = 1$ ; i.e., if no edge is monochromatic under *c*. We denote by  $\rho_k(G)$  the largest value of  $\rho_k(c)$  over all *k*-colourings *c* of *G*:

$$\rho_k(G) = \max_{c:V \to \{1,\dots,k\}} \rho_k(c) \, .$$

Testing whether  $\rho_k(G) = 1$  is the same as determining whether G admits a proper k-colouring; this problem is NP-hard for  $k \ge 3$ , as shown by Karp [Kar72], and solvable in polynomial time for k = 1, 2.

Given a graph G, say with  $\rho = \rho_2(G) < 1$  (since the case  $\rho_2(G) = 1$  is solvable exactly efficiently), the celebrated result of Goemans and Williamson uses a semidefinite programming (SDP) relaxation, equivalent to an eigenvalue minimisation problem proposed earlier by Delorme and Poljak [DP93a, DP93b], to design a polynomial-time randomised algorithm that finds a 2-colouring c of G with  $\rho_2(c) \ge \alpha_{GW}\rho$  [GW95], where  $\alpha_{GW} \approx 0.87856$ . Their algorithm was later derandomised by Mahajan and Ramesh [MR99]. On the hardness side, the work of Håstad [Hås01] and Trevisan, Sorkin, Sudan, and Williamson [TSSW00] showed that obtaining a 2-colouring c with  $\rho_2(c) \ge \alpha \rho$  is NP-hard for any  $\alpha \ge 16/17 + \varepsilon$  for an arbitrarily small  $\varepsilon > 0$ . Note that  $16/17 \approx 0.94117$  and thus there is a gap between  $\alpha_{GW}$  and 16/17. However, under Khot's influential *Unique Games Conjecture* (UGC) [Kho02], Khot, Kindler, Mossel and O'Donnell showed that finding a 2-colouring c with  $\rho_2(c) \ge \alpha \rho$  is NP-hard for any  $\alpha \ge \alpha_{GW} + \varepsilon$  [KKM007],<sup>2</sup> thus showing that the algorithm of Goemans and Williamson [GW95] is optimal (up to an arbitrarily small additive constant). In fact, the Goemans-Williamson algorithm finds, given a graph G with  $\rho_2(G) = 1 - \varepsilon$ , a 2-colouring c of G with  $\rho_2(c) = 1 - O(\sqrt{\varepsilon})$  [GW95]. Moreover, the dependence on  $\varepsilon$  is UGC-optimal [KKM007].

What about colourings with more than two colours? Building on the work of Goemans and Williamson [GW95], Frieze and Jerrum [FJ97] provided an SDP-based algorithm for approximating  $\rho_k(G)$  for every G and constant  $k \ge 2$ . Asymptotic optimality of this algorithm for large k (up to an arbitrarily small additive constant) was shown by Khot, Kindler, Mossel, and O'Donnell under the UGC [KKM007], as we will discuss in more detail later. All the results mentioned so far are concerned with quantitative approximation. We now turn to qualitative approximation.

Let *G* be a graph that can be properly *k*-coloured; i.e.,  $\rho_k(G) = 1$ . Is it possible to find efficiently a proper  $\ell$ -colouring of *G* for some constant  $k \leq \ell$ ? Garey and Johnson conjectured that this problem is NP-hard as long as  $k \geq 3$  [G]76]. For k = 3, NP-hardness is known for  $\ell = 3$  [Kar72],  $\ell = 4$  [KLS00, GK04], and  $\ell = 5$  [BBKO21]; the case of  $\ell \geq 6$  is open. For  $k \geq 4$ , better bounds are known [KOWŽ23]. However, NP-hardness has been shown for all constant  $3 \leq k \leq \ell$  under stronger assumptions. Namely, under a non-standard variant of the Unique Games Conjecture by Dinur, Mossel, and Regev [DMR09], under the *d*-to-1 conjecture of Khot [Kh002] (for any fixed *d*) by Guruswami and Sandeep [GS20], and under the rich 2-to-1 conjecture of Braverman, Khot, and Minzer [BKM21] by Braverman, Khot, Lifshitz, and Minzer in [BKLM21].

We now combine the quantitative and qualitative approaches. Given a graph G of value  $\rho_k(G)$ , what is the largest  $0 < \alpha \le 1$  so that  $\rho_\ell(G)$  can be  $\alpha$ -approximated? It is not hard to show that, for  $3 \le k \le \ell$ ,

<sup>&</sup>lt;sup>2</sup>The results in [KKM007] was initially conditional on the "majority is stablest" conjecture, later proved by Mossel, O'Donnell, and Oleszkiewicz [M0010].

$k^{\ell}$	3	4	5	6	7	8	9	10	11	12	13	14	15
3	.836	.904	.938	.957	.969	.976	.982	.985	.988	.990	.992	.993	.994
4		.858	.899	.924	.940	.952	.960	.967	.972	.975	.979	.981	.983
5			.877	.904	.923	.936	.946	.954	.960	.964	.968	.972	.974
6				.892	.911	.926	.936	.945	.952	.957	.961	.965	.968
7					.903	.918	.930	.938	.945	.951	.956	.960	.963
8						.913	.924	.934	.941	.947	.952	.956	.960
9							.920	.930	.937	.944	.949	.953	.957
10								.927	.935	.941	.946	.951	.954
11									.932	.939	.944	.949	.953
12										.937	.942	.947	.951
13											.941	.946	.950
14												.944	.949
15													.948

Table 1: Approximate values of  $\alpha_{k\ell}$ . As all values are between 0 and 1, we omit the leading 0.

a 1-approximation is at least as hard approximate graph colouring, cf. Appendix A. For  $\alpha < 1$ , not much is known other than what follows immediately from the already mentioned previous work: The algorithm from [FJ97] gives an  $\alpha$ -approximation with  $\alpha \ge 1-1/\ell + (1+\varepsilon(\ell))(2\ln\ell/\ell^2)$ , where  $\varepsilon(\ell) \to 0$  as  $\ell \to \infty$ , and for  $\ell = k \ge 2$  this algorithm is UGC-optimal (up to an arbitrarily small additive constant) [KKM007]. However, the situation is unclear for general k and  $\ell$  and in fact already for  $k = 2 \le \ell$  and  $\alpha = 1$ . If  $\rho_2(G) = 1$  then G is bipartite and the problem is easy but what if  $\rho_2(G) < 1$ ?

**Contributions** We initiate a systematic investigation of promise maximum colouring, i.e., k-vs.  $\ell$ -colourings. As our first result, we extend the algorithm of Frieze and Jerrum [FJ97] to work for k-vs.  $\ell$ -colourings. We analyse the power of the algorithm in two cases: for k = 2,  $\ell = 3$ , and for  $k \leq \ell$  as  $k, \ell \rightarrow \infty$ . For the case k = 2 one gets the best possible scenario: There is an efficient approximation algorithm with approximation factor  $\alpha = 1$  already for 3-colourings (and thus also for  $\ell$ -colourings for any  $\ell \geq 3$ ). In particular, we will show the following result.

**Theorem 1.** There is a randomised algorithm which, given a graph G that admits a k-colouring of value  $\rho$  and a real number  $\varepsilon > 0$ , finds an  $\ell$ -colouring of expected value  $\alpha_{k\ell} \rho - \varepsilon$  in polynomial time in G and  $\log(1/\varepsilon)$ . In particular,

- 1. For  $k = 2, \ell = 3$ , we have  $\alpha_{2,3} = 1$ .
- 2. We have that  $\alpha_{k\ell} \ge 1 \frac{1}{\ell} + \frac{2\ln\ell}{k\ell} o\left(\frac{\ln\ell}{k\ell}\right) O\left(\frac{1}{k^2}\right)$ .
- 3. Furthermore,  $\alpha_{k\ell} > 1 1/\ell$ , hence the algorithm is better than random guessing for  $\rho$  near 1.<sup>3</sup>

For illustration, we tabulated numerical approximate values for  $\alpha_{k\ell}$  in Table 1 on page 3.<sup>4</sup> Our algorithm solves the Frieze-Jerrum SDP for *k*-colourings, then rounds like Frieze and Jerrum do for  $\ell$ -colourings [FJ97]. For 2- vs. 3-colourings our analysis begins much like, but is simpler than that of Frieze and Jerrum, because we only care about the case where we get 3 colours; in particular, we establish a closed form for the probability that a pair of vectors is cut properly using the results of Cheng [Che68, Che69]. For *k*- vs.  $\ell$ -colourings, we largely follow the analysis from [FJ97].

We will also show how to derandomise our algorithm.

<sup>&</sup>lt;sup>3</sup>The  $o(\cdot)$ ,  $O(\cdot)$  notation hides only terms and factors dependant on  $k, \ell$ , not on  $\varepsilon$ .

<sup>&</sup>lt;sup>4</sup>The exact definition of  $\alpha_{k\ell}$  is given in Definition 7. The probabilities  $P_{\ell}(a)$  that appear in that definition were computed using the methods and R library from [AG18].

**Theorem 2.** Fix  $2 \le k \le l$  and let  $\alpha_{kl}$  be as in Theorem 1. There is a deterministic algorithm which, given a graph G that admits a k-colouring of value  $\rho$  and a real number  $\varepsilon > 0$ , finds an l-colouring of value  $\alpha_{kl} \rho - \varepsilon$  in polynomial time in G and  $1/\varepsilon$ .

The proof of Theorem 2 uses the method of conditional expectation; however our method is conceptually simpler (if less efficient) than previous ones: to compute our conditional probabilities, we approximate the normal distribution with a normalised binomial distribution. Moreover, our method explicitly works not only for the rounding scheme in our algorithm but also in the rounding scheme of Frieze and Jerrum [FJ97] (which we use), in contrast to previous work that did not explicitly handle this rounding scheme [MR99, EIO02, BK05, GOWZ10]; cf. Section 4 and the discussion therein for details. Theorem 2 implies the following corollary for k = 2,  $\ell = 3$ , which eliminates any dependence on  $\varepsilon$ .

**Corollary 3.** There is a deterministic algorithm which, given a graph G that admits a 2-colouring of value  $\rho$ , finds a 3-colouring of value  $\rho$  in polynomial time in G.

*Proof.* By Theorems 1 and 2, there exists an algorithm which, if given a graph *G* that admits a 2-colouring of value  $\rho$ , will find a 3-colouring of value  $\rho - \varepsilon$  in polynomial time in *G* and  $1/\varepsilon$ . Note that the value of any colouring is of the form k/m, where *m* is the number of edges of *G*; in particular suppose  $\rho = k/m$ . Set  $\varepsilon = 1/2m$  and note that the algorithm returns a 3-colouring of value at least k/m - 1/2m; since the value of the returned colouring is of the form k'/m, it must be at least k/m as required.

The algorithm from Theorem 2 has good performance when both k and  $\ell$  grow large. What if k is fixed and only  $\ell$  grows large? We give an algorithm that has good performance in this case as well. The idea is based on an algorithm of Karger, Motwani and Sudan for approximate graph colouring [KMS98, Section 6], but rather than cutting by  $\Theta(\log(n))$  random hyperplanes we cut with  $\lfloor \log_2(\ell) \rfloor$  hyperplanes.

**Theorem 4.** Let k > 2 be fixed<sup>5</sup> and  $\ell \ge k$  be large. There is a deterministic algorithm which, given a graph G that admits a k-colouring of value  $\rho$  and a real number  $\varepsilon > 0$ , finds an  $\ell$ -colouring of G of value  $\alpha'_{k\ell}\rho - \varepsilon$  in polynomial time in G and  $1/\varepsilon$ . In particular, for a fixed k there exists a constant  $u_k > 1$  such that

$$\alpha_{k\ell}' \geq 1 - O\left(1/\ell^{u_k}\right),\,$$

which is  $1 - o(1/\ell)$  when  $\ell$  grows large.<sup>6</sup>

Of course, by running the algorithms of Theorem 2 and Theorem 4 in parallel and then taking the better of the two results, we can get an algorithm that is at least as good as either of them.

Using the framework of Khot, Kindler, Mossel, and O'Donnel [KKMO07], we will show that, under the UGC, it is NP-hard to beat the approximation guarantee of our algorithm from Theorem 2 by more than a constant that grows small for any large k,  $\ell$  with  $k \leq \ell$  and  $\ell$  bounded by a function that is  $o(e^{\sqrt[3]{k}})$ . We combine this with the methods of Guruswami and Sinop [GS13]<sup>7</sup> to also find some weaker (non-tight) unconditional results.<sup>8</sup> We will present a unified version of these proofs, using ideas from the work of Dinur, Mossel and Regev [DMR06] (on which [GS13] also draws). The way the unification of these two proofs works out is also similar to the work of Guruswami and Sandeep [GS20].

<sup>&</sup>lt;sup>5</sup>The case  $k = 2, \ell \ge 2$  is covered by Theorem 2.

<sup>&</sup>lt;sup>6</sup>For example, for k = 3 we have  $u_k = 2$  and the approximation ratio is  $1 - O(1/\ell^2)$ . This is significantly better, for large  $\ell$ , than the random guessing algorithm which has performance  $1 - 1/\ell$ .

<sup>&</sup>lt;sup>7</sup>The results of [GS13] were, at the time, conditional on the 2-to-1 conjecture of Khot [Kho02]; however this has recently been proved by Khot, Minzer and Safra [KMS23].

<sup>&</sup>lt;sup>8</sup>We thank Venkat Guruswami for bringing [GS13] to our attention.

**Theorem 5.** Fix some function  $M(k) = o(e^{\sqrt[3]{k}})$ . Let  $2 \le k \le \ell$  be such that  $\ell \le M(k)$ . For any small enough  $\varepsilon > 0$ , consider the problem of deciding whether a given graph G admits a k-colouring of value  $1 - \varepsilon$ , or not even an  $\ell$ -colouring of value  $\beta + \varepsilon$ . We have the following.

• Assuming the UGC, the problem is NP-hard for  $\beta = \beta_{k\ell} = 1 - \frac{1}{\ell} + \frac{2\ln\ell}{k\ell} + o\left(\frac{\ln\ell}{k\ell}\right)$ .

• Unconditionally, the problem is NP-hard for  $\beta = \beta'_{k\ell} = 1 - \frac{1}{\ell} + \frac{8 \ln \ell}{k\ell} + o\left(\frac{\ln \ell}{k\ell}\right)$ .

Both of these results only hold when  $\beta_{k\ell}, \beta'_{k\ell} \in (0, 1)$ .

The NP-hardness bound in Theorem 5 is limited due to the fact that, for a fixed k, we cannot have  $\ell$  arbitrarily large. This is intrinsic to the expression above: for a large  $\ell$  and a fixed k we have  $\beta_{k\ell} > 1$ . Moreover, any NP-hardness bound for a fixed k and a large  $\ell$  must take into account the algorithm with approximation ratio  $1 - o(1/\ell)$  we gave in Theorem 4.

**Evaluation of performance bounds** For  $k = \ell$  we recover the positive results of [FJ97] (and indeed our algorithm is the same as that of [FJ97] for  $k = \ell$ ) and the negative result of [KKMO07]. In detail, we have that our algorithm from Theorem 1 has performance  $1 - 1/\ell + 2 \ln \ell/\ell^2 - o(\ln \ell/\ell^2)$  and that it is NP-hard to do, under the UGC, any better than  $1 - 1/\ell + 2 \ln \ell/\ell\ell + o(\ln \ell/\ell^2)$ .<sup>10</sup> We also recover the unconditional (in light of [KMS23]) result of [GS13], i.e. that it is NP-hard to do any better than  $1 - 1/\ell + 8 \ln \ell/\ell\ell + o(\ln \ell/\ell^2)$ . For k = 2,  $\ell = 3$ , we get an approximation ratio of 1. This is the best possible result.<sup>11</sup>

For the fixed-gap case, i.e.  $\ell = k + d$  for some fixed  $d \ge 0$ , we get the same type of result as for  $k = \ell$ : performance  $1 - 1/\ell + 2 \ln \ell/k\ell - o(\ln \ell/k\ell)$  and NP-hardness, under the UGC, of  $1 - 1/\ell + 2 \ln \ell/k\ell + o(\ln \ell/k\ell)$ , since in this case the  $1/k^2$  term is strictly dominated by  $\ln \ell/k\ell$ .<sup>12</sup> Furthermore, unconditionally we find that an approximation ratio of  $1 - 1/\ell + 8 \ln \ell/k\ell + o(\ln \ell/k\ell)$  is NP-hard to achieve. This unconditional result is not yet tight, since already the second order term is different.

For fixed k and large  $\ell$ , the algorithm from Theorem 4 has performance  $1 - o(1/\ell)$ . This algorithm cannot be improved by more than  $o(1/\ell)$ , since no algorithm can have approximation ratio greater than 1. We believe that the algorithm from Theorem 2 is at least as strong as the algorithm from Theorem 4 even for fixed k and large  $\ell$ .

We note that the fact that our NP-hardness bound in Theorem 5 only works for  $\ell$  bounded by some function of k mirrors the current state-of-the-art for approximate graph colouring: distinguishing proper k-vs.  $\ell$ -colourings is only known to be NP-hard whenever  $\ell \leq \binom{k}{\lfloor k/2 \rfloor} - 1$  [KOWŽ23].

**Related work** Graph colouring is a canonical example a *Constraint Satisfaction Problem* (CSP) [FV98, KSTW00]. Robust solvability of CSPs was studied, among others, by Charikar, Makarychev, and Makarychev [CMM09], Guruswami and Zhou [GZ12], and Barto and Kozik [BK16]. Raghavandra showed UGC-optimality of the basic SDP programming relaxation for all CSPs [Rag08]. The notion of an almost *k*-colouring (a large fraction of the graph being properly *k*-coloured) was recently studied by Hecht, Minzer, and Safra [HMS23],

<sup>&</sup>lt;sup>9</sup>The constants hidden in the expression defining  $\beta_{k\ell}$  depend on  $M(k), k, \ell$ , but not on  $\varepsilon$ .

<sup>&</sup>lt;sup>10</sup>The negative result of [KKMO07] is slightly more specific as their asymptotic error term is  $O(\ln \ln \ell/\ell^2)$ . A careful inspection of our analysis shows that our error term is  $O(1/\ell^2 + \ln \ell \ln \ln k/\ell k \ln k)$ , which for  $\ell = k$  is precisely their  $O(\ln \ln \ell/\ell^2)$ .

<sup>&</sup>lt;sup>11</sup>The graph on 2p + q vertices consisting of p disjoint edges and q disjoint loops has a k-colouring of value p/(p+q), and no  $\ell$ -colouring of value greater than p/(p+q). Thus it is not possible in general to find an  $\ell$ -colouring of value greater than p/(p+q) for a graph with a k-colouring of value p/(p+q).

<sup>&</sup>lt;sup>12</sup>As an example, this implies that for large k we can do k- vs. ( $\ell = k + 10$ )-colourings with approximation ratio  $1 - 1/\ell + 1.999 \ln \ell/k\ell$ , but it is NP-hard under the UGC to do it with approximation ratio  $1 - 1/\ell + 2.001 \ln \ell/k\ell$ .

who showed that finding an almost *k*-colouring of a graph that admits an almost 3-colouring is NP-hard for every constant *k*. Austrin, O'Donnell, Tan and Wright showed NP-hardness of distinguishing whether  $\rho_3(G) = 1$  or  $\rho_3(G) < \frac{16}{17} + \varepsilon$  [AOTW14].

Approximate graph colouring is an example of a *Promise Constraint Satisfaction Problem* (PCSP) [AGH17, BG21, BBKO21]. Robust solvability of PCSPs has recently been investigated by Brakensiek, Guruswami, and Sandeep [BGS23]. Bhangale, Khot, and Minzer have recently studied approximability of certain Boolean PCSPs [BKM22, BKM23a, BKM23b].

We finish with a remark about our result on 2- vs. 3-colourings in the context of PCSPs: our tractability result cannot be explained by a finite sandwich (in the sense of [BG19]), cf. Appendix B.

## 2 Preliminaries

For any positive integer n let  $[n] = \{1, ..., n\}$ . For any predicate  $\phi$ , we let  $[\phi] = 1$  if  $\phi$  is true, and 0 otherwise. We shall use semidefinite programming and refer the reader to [GM12] for a reference.

For an event  $\phi$  we let  $\Pr[\phi]$  be the probability that  $\phi$  is true. For a random variable X, we let  $\mathbb{E}[X]$  denote its expected value. Note that  $\mathbb{E}[[\phi]] = \Pr[\phi]$ .

For any two distributions  $\mathcal{D}$ ,  $\mathcal{D}'$  with domains A, A', we let  $\mathcal{D} \times \mathcal{D}'$  denote the product distribution, whose domain is  $A \times A'$ . For any distribution  $\mathcal{D}$  over  $\mathbb{R}$  and  $a, b \in \mathbb{R}$ , the distribution  $a\mathcal{D} + b$  is the distribution of aX + b when  $X \sim \mathcal{D}$ . We use the standard probability theory abbreviations i.i.d. (independent and identically distributed) and p.m.f. (probability mass function).

We introduce a few classic distributions we will need. The uniform distribution  $\mathcal{U}(D)$  over a finite set D is the distribution with p.m.f.  $f: D \to [0, 1]$  given by f(x) = 1/|D|. Note that  $\mathcal{U}(D^n)$  is the same as  $\mathcal{U}(D)^n$ , a fact which we will use implicitly. We let NBin(n) denote a normalised binomial distribution: it is the distribution of  $X_1 + \cdots + X_n$ , where  $X_i \sim \mathcal{U}(\{-1/\sqrt{n}, 1/\sqrt{n}\})$ . The domain of this distribution is  $\{(-n+2k)/\sqrt{n} \mid 0 \le k \le n\}$ , the probability mass function is  $(-n+2k)/\sqrt{n} \mapsto \binom{n}{k}/2^n$ , the expectation is 0, and the variance is 1. If  $\mu, \sigma \in \mathbb{R}$ , then we let  $\mathcal{N}(\mu, \sigma^2)$  denote the normal distribution with mean  $\mu$  and variance  $\sigma^2$ . Fixing d, if  $\mu \in \mathbb{R}^d, \Sigma \in \mathbb{R}^{d \times d}$ , then we let  $\mathcal{N}(\mu, \Sigma)$  denote the multivariate normal distribution with mean  $\mu$  and covariance matrix  $\Sigma$ . We let  $\mathbf{I}_d$  denote the  $d \times d$  identity matrix. Observe that if  $\mathbf{x} \sim \mathcal{N}(\mu, \Sigma)$ , where  $\mathbf{x} \in \mathbb{R}^d$ , then for any matrix  $\mathbf{A} \in \mathbb{R}^{d' \times d}$  we have that  $\mathbf{Ax} \sim \mathcal{N}(\mathbf{A}\mu, \mathbf{A\Sigma}\mathbf{A}^T)$ . Furthermore if  $\mathbf{x} \sim \mathcal{N}(\mu, \Sigma)$  with  $\Sigma$  positive semidefinite, then by finding the Cholesky decomposition  $\Sigma = \mathbf{A}\mathbf{A}^T$ , where  $\mathbf{A} \in \mathbb{R}^{d \times d}$ , we find that  $\mathbf{x}$  is identically distributed to  $\mathbf{Ax}' + \mu$ , where  $\mathbf{x}' \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_d)$ .

## 3 Main result

In this section, we will prove our main result, restated here.

**Theorem 1.** There is a randomised algorithm which, given a graph G that admits a k-colouring of value  $\rho$  and a real number  $\varepsilon > 0$ , finds an  $\ell$ -colouring of expected value  $\alpha_{k\ell}\rho - \varepsilon$  in polynomial time in G and  $\log(1/\varepsilon)$ . In particular,

1. For  $k = 2, \ell = 3$ , we have  $\alpha_{2,3} = 1$ .

2. We have that 
$$\alpha_{k\ell} \ge 1 - \frac{1}{\ell} + \frac{2\ln\ell}{k\ell} - o\left(\frac{\ln\ell}{k\ell}\right) - O\left(\frac{1}{k^2}\right)$$
.

3. Furthermore,  $\alpha_{k\ell} > 1 - 1/\ell$ , hence the algorithm is better than random guessing for  $\rho$  near 1.<sup>13</sup>

<sup>&</sup>lt;sup>13</sup>The  $o(\cdot)$ ,  $O(\cdot)$  notation hides only terms and factors dependant on  $k, \ell$ , not on  $\varepsilon$ .

In order to prove Theorem 1, we first introduce an auxiliary notion, which already appears in [F]97].

**Definition 6.** Fix  $a, b \in \mathbb{R}$  such that  $a^2 + b^2 = 1, b \ge 0$ , and  $\ell \in \mathbb{N}$ . Suppose that  $x_1, \ldots, x_\ell, y_1, \ldots, y_\ell \sim \mathcal{N}(0, 1)$ ; i.e., they are i.i.d. standard normal variables. We let  $P_\ell(a)$  denote the probability that

$$x_1 \ge x_2 \land \dots \land x_1 \ge x_\ell \land ax_1 + by_1 \ge ax_2 + by_2 \land \dots \land ax_1 + by_1 \ge ax_\ell + by_\ell.$$

We then write  $N_{\ell}(a) = \ell P_{\ell}(a)$ . This is just the probability that  $\arg \max_{c} x_{c} = \arg \max_{c} ax_{c} + by_{c}$ .

The following quantity is similar to  $\alpha_k$  from [F]97].

Definition 7. Let

$$\alpha_{k\ell} = \min_{-1/(k-1) \le a < 1} \frac{k(1 - \ell P_{\ell}(a))}{(k-1)(1-a)}$$

Observe that for a = 1 the ratio would be 0/0, hence for  $-1/(k-1) \le a \le 1$  it holds that

$$\alpha_{k\ell} \frac{k-1}{k} (1-a) \le 1 - \ell P_{\ell}(a). \tag{1}$$

The proof of Theorem 1 is split into the following four propositions.

**Proposition 8.** There is a randomised algorithm which, given a graph G that admits a k-colouring of value  $\rho$ , finds an  $\ell$ -colouring of expected value  $\alpha_{k\ell}\rho - \varepsilon$  in polynomial time in G and  $\log(1/\varepsilon)$  for an arbitrarily small  $\varepsilon > 0$ .

**Proposition 9.**  $\alpha_{2,3} = 1$ .

**Proposition 10.**  $\alpha_{k\ell} \ge 1 - \frac{1}{\ell} + \frac{2\ln\ell}{k\ell} - o\left(\frac{\ln\ell}{k\ell}\right) - O\left(\frac{1}{k^2}\right).$ 

**Proposition 11.**  $\alpha_{k\ell} > 1 - 1/\ell$ .

#### 3.1 **Proof of Proposition 8**

Our algorithm solves the SDP of [FJ97] for *k*-colourings, then rounds the solution of the SDP like [FJ97] but for  $\ell$ -colourings. Henceforth fix  $2 \le k \le \ell$ , and  $\varepsilon > 0$ . The following lemma also appears, essentially, as [FJ97, Lemma 3] and the preceding definitions; we include it for completeness.

**Lemma 12.** For any  $n \ge k$ , there exist vectors  $\mathbf{e}_1, \ldots, \mathbf{e}_k \in \mathbb{R}^n$  such that  $\mathbf{e}_i \cdot \mathbf{e}_i = 1$  and  $\mathbf{e}_i \cdot \mathbf{e}_j = -1/(k-1)$  for  $i \ne j$ .

*Proof.* Let  $\mathbf{e}_i = \frac{1}{k(k-1)} (1, \dots, 1, 1-k, 1, \dots, 1, 0, \dots, 0)^T$ , where there are k nonzero values, and where the value 1 - k appears at the *i*-th position. These vectors satisfy the required conditions.

*Proof of Proposition 8.* Suppose we are given a graph G = (V, E), which we are promised has a *k*-colouring of value  $\rho$ . Suppose V = [n] and |E| = m. If  $n \le k$  then simply iterating over all possible *k*-colourings and finding the best one is sufficient, thus assume  $n \ge k$ .

By relabelling the promised colouring to  $\mathbf{e}_1, \ldots, \mathbf{e}_k \in \mathbb{R}^n$ , we find that there exist variables  $\mathbf{a}_i^* \in {\mathbf{e}_1, \ldots, \mathbf{e}_k} \subseteq \mathbb{R}^n$  for  $i \in [n]$  such that  $\frac{1}{m} \sum_{(i,j) \in E} [\mathbf{a}_i^* \neq \mathbf{a}_j^*] \ge \rho$ . We find that  $[\mathbf{a}_i^* \neq \mathbf{a}_j^*] = \frac{k-1}{k} (1 - \mathbf{a}_i^* \cdot \mathbf{a}_j^*)$ , when

 $\mathbf{a}_{i}^{*}, \mathbf{a}_{j}^{*} \in {\mathbf{e}_{1}, \dots, \mathbf{e}_{k}}$ . We now relax as Frieze and Jerrum [FJ97], and Goemans and Williamson before them [GW95], to a semidefinite program; namely, we solve the following program:

$$\max \quad \frac{1}{m} \sum_{(i,j) \in E} \frac{k-1}{k} (1 - \mathbf{a}_i \cdot \mathbf{a}_j)$$
  
s.t. 
$$\mathbf{a}_i \cdot \mathbf{a}_i = 1,$$
  
$$\mathbf{a}_i \cdot \mathbf{a}_j \ge -\frac{1}{k-1}, i \neq j,$$
  
$$\mathbf{a}_i \in \mathbb{R}^n.$$
 (2)

The semidefinite program (2) can be solved with an additive error of at most  $\varepsilon/\alpha_{k\ell}$  in time polynomial with respect to n, m and  $\log(\alpha_{k\ell}/\varepsilon) = \log(1/\varepsilon) + O(1)$ . By the discussion in the previous paragraph, we see that the SDP must have value at least  $\rho$ , due to the potential solution  $\mathbf{a}_1^*, \ldots, \mathbf{a}_n^*$ . Thus, by solving the program we now have a collection of n unit vectors  $\mathbf{a}_1, \ldots, \mathbf{a}_n \in \mathbb{R}^n$  with pairwise inner product at least -1/(k-1) such that

$$\frac{1}{m}\sum_{(i,j)\in E}\frac{k-1}{k}(1-\mathbf{a}_i\cdot\mathbf{a}_j)\geq \rho-\varepsilon/\alpha_{k\ell}.$$

Our algorithm now *randomly rounds* as Frieze and Jerrum does [FJ97], for  $\ell$ -colourings. Namely, we take  $\ell$  standard normal variables  $\mathbf{x}_1, \ldots, \mathbf{x}_{\ell} \in \mathbb{R}^n$ ; for each vertex  $i \in V$  we compute  $c = \arg \max_j \mathbf{a}_i \cdot \mathbf{x}_j$ , and then assign vertex i colour c (breaking possible ties arbitrarily).

Now, let us compute the expected value of the resulting rounding. Consider an edge  $(i, j) \in E$ ; in terms of  $\frac{k-1}{k}(1 - \mathbf{a}_i \cdot \mathbf{a}_j)$ , what is the probability that (i, j) is properly coloured? This is the same as the probability that arg max<sub>c</sub>  $\mathbf{a}_i \cdot \mathbf{x}_c \neq \arg \max_c \mathbf{a}_j \cdot \mathbf{x}_c$ , which, by symmetry, is equal to

$$1 - \ell \Pr_{\mathbf{x}_1, \dots, \mathbf{x}_{\ell}} \left[ \bigwedge_{c=2}^{\ell} \mathbf{a}_i \cdot \mathbf{x}_1 \ge \mathbf{a}_i \cdot \mathbf{x}_c, \bigwedge_{c=2}^{\ell} \mathbf{a}_j \cdot \mathbf{x}_1 \ge \mathbf{a}_j \cdot \mathbf{x}_c \right].$$
(3)

Since  $\mathbf{x}_1, \ldots, \mathbf{x}_\ell$  are drawn from a rotationally symmetric distribution, we can rotate everything to be in a 2dimensional plane without affecting the probability in (3). Furthermore, rotate so that  $\mathbf{a}_i$  is moved to (1, 0), and  $\mathbf{a}_j$  is at (a, b), where  $a = \mathbf{a}_i \cdot \mathbf{a}_j$  and  $b = \sqrt{1 - a^2}$  (note that this rotation is possible since it preserves the angle between  $\mathbf{a}_i$  and  $\mathbf{a}_j$ , and their lengths). Since the vectors  $\mathbf{x}_1, \ldots, \mathbf{x}_\ell$  are (after the rotation) bivariate standard normal variables, we can see them as pairs  $(x_1, y_1), \ldots, (x_\ell, y_\ell)$ , where  $x_1, \ldots, x_\ell, y_1, \ldots, y_\ell \sim$  $\mathcal{N}(0, 1)$  are i.i.d. standard normal variables. Then, we can rewrite (3) as

$$1 - \ell \Pr_{\substack{x_1, \dots, x_\ell \\ y_1, \dots, y_\ell}} \left[ \bigwedge_{c=2}^{\ell} x_1 \ge x_c, \bigwedge_{c=2}^{\ell} ax_1 + by_1 \ge ax_c + by_c \right] = 1 - \ell P_\ell(a) = 1 - \ell P_\ell(\mathbf{a}_i \cdot \mathbf{a}_j).$$
(4)

Since  $-1/(k-1) \leq \mathbf{a}_i \cdot \mathbf{a}_j \leq 1$ , by (1), we have that

$$\alpha_{k\ell}\frac{k-1}{k}(1-\mathbf{a}_i\cdot\mathbf{a}_j)\leq 1-\ell P_\ell(\mathbf{a}_i\cdot\mathbf{a}_j).$$

Hence, by linearity of expectation the expected value of the  $\ell$ -colouring we return is, as required, at least

$$\alpha_{k\ell} \frac{1}{m} \sum_{(i,j)\in E} \frac{k-1}{k} (1 - \mathbf{a}_i \cdot \mathbf{a}_j) \ge \alpha_{k\ell} \rho - \varepsilon.$$

## 3.2 **Proof of Proposition 9**

In this section we prove Proposition 9, i.e. that  $\alpha_{2,3} = 1$ . For this case, we can find  $P_3(a)$  explicitly.

Lemma 13. 
$$P_3(a) = \frac{1}{9} + \frac{\arcsin a + \arcsin \frac{a}{2}}{4\pi} + \frac{(\arcsin a)^2 - (\arcsin \frac{a}{2})^2}{4\pi^2}$$

To prove Lemma 13, we will use the following result of Cheng.

**Theorem 14** ([Che68][Che69, Equation (2.18)]). Suppose  $\mathbf{u} = (u_1, u_2, u_3, u_4) \sim \mathcal{N}(\mathbf{0}, \boldsymbol{\Sigma})$  are drawn from a quadrivariate normal distribution with mean zero and covariance matrix

$$\boldsymbol{\Sigma} = \begin{pmatrix} 1 & a & b & ab \\ a & 1 & ab & b \\ b & ab & 1 & a \\ ab & b & a & 1 \end{pmatrix},$$

where  $a, b \in [-1, 1]$ . Then  $\Pr_{\mathbf{u}}[u_1 \ge 0, u_2 \ge 0, u_3 \ge 0, u_4 \ge 0]$  is

$$\frac{1}{16} + \frac{\arcsin a + \arcsin b + \arcsin ab}{4\pi} + \frac{(\arcsin a)^2 + (\arcsin b)^2 - (\arcsin ab)^2}{4\pi^2}$$

Proof of Lemma 13. Define the following normally distributed random variables:

$$u_1 = (x_1 - x_2)/\sqrt{2}, \qquad u_3 = (ax_1 + by_1 - ax_2 - by_2)/\sqrt{2}, u_2 = (x_1 - x_3)/\sqrt{2}, \qquad u_4 = (ax_1 + by_1 - ax_3 - by_3)/\sqrt{2}.$$

By simple computation and since  $a^2 + b^2 = 1$ , we have that  $\mathbf{u} = (u_1, u_2, u_3, u_4) \sim \mathcal{N}(\mathbf{0}, \boldsymbol{\Sigma})$ , where

$$\boldsymbol{\Sigma} = \begin{pmatrix} 1 & \frac{1}{2} & a & \frac{a}{2} \\ \frac{1}{2} & 1 & \frac{a}{2} & a \\ a & \frac{a}{2} & 1 & \frac{1}{2} \\ \frac{a}{2} & a & \frac{1}{2} & 1 \end{pmatrix}.$$

The probability we want is just  $\Pr_{\mathbf{u}}[u_1 \ge 0, u_2 \ge 0, u_3 \ge 0, u_4 \ge 0]$ . Apply Theorem 14 with  $a = \frac{1}{2}$  and b = a to find that the probability we want is

$$\frac{1}{16} + \frac{\arcsin\frac{1}{2} + \arcsin a + \arcsin\frac{a}{2}}{4\pi} + \frac{(\arcsin\frac{1}{2})^2 + (\arcsin a)^2 - (\arcsin\frac{a}{2})^2}{4\pi^2} = \frac{1}{9} + \frac{\arcsin a + \arcsin \frac{a}{2}}{4\pi} + \frac{(\arcsin a)^2 - (\arcsin\frac{a}{2})^2}{4\pi^2},$$
required.

as required.

The bound we want on  $P_3$  is the following.

Lemma 15. For  $-1 \le a \le 1$ ,  $1 - 3P_3(a) \ge \frac{1-a}{2}$ .

The functions involved are shown in Figure 1.



Figure 1: Plot of expressions from Lemma 15.

Proof. Define

$$f(a) = 1 - 3P_3(a) - \frac{1 - a}{2} = \frac{1}{6} + \frac{a}{2} - \frac{3}{4\pi} \left( \arcsin a + \arcsin \frac{a}{2} \right) - \frac{3}{4\pi^2} \left( (\arcsin a)^2 - \left( \arcsin \frac{a}{2} \right)^2 \right).$$

We want to show that  $f(x) \ge 0$  for  $x \in [-1, -1]$ . First we show that  $f(x) \ge 0$  for  $x \in [-1, 0]$ . Numerically, we can find that

$$\max_{-1 < x < 1} f^{\prime\prime\prime}(x) \approx -0.1454 < 0,$$

at  $x \approx -0.5681$ . Thus f''(x) is decreasing, and f'(x) is concave. Thus, by Jensen's inequality, for  $x \in (-1, 0)$ ,

$$f'(x) \ge -x \lim_{t \to -1^+} f'(t) + (1+x)f'(0),$$

(as f' is not defined at -1). But  $f'(0) \approx 0.1419 > 0$ , and  $\lim_{t \to -1^+} f'(t) \approx 0.1642 > 0$ , so f'(x) > 0 for  $x \in (-1, 0)$ . It follows that f is increasing on [-1, 0], which is sufficient to show that  $f(x) \ge 0$  for  $x \in [-1, 0]$ , as f(-1) = 0.

Now, we consider  $x \in [0, 1]$ . Observe again that we know that f''(x) is decreasing. But since  $f''(0) \approx -0.1139 < 0$ , it follows that f''(x) < 0 for  $x \in [0, 1)$ ; so f is concave on [0, 1]. Again applying Jensen's inequality, we find that for  $x \in [0, 1]$ ,  $f(x) \ge xf(0) + (1 - x)f(1)$ . As f(0) = 1/6 > 0 and f(1) = 0, we get  $f(x) \ge 0$  for  $x \in [0, 1]$ . Thus our conclusion follows in all cases.

*Proof of Proposition 9.* By Lemma 15 we have that

$$\frac{2(1-3P_3(a))}{(2-1)(1-a)} = \frac{1-3P_3(a)}{\frac{1}{2}(1-a)} \ge 1$$

for all  $a \in [-1, 1)$ . Hence

$$\alpha_{2,3} = \min_{a \in [-1,1)} \frac{2(1-3P_3(a))}{(2-1)(1-a)} \ge 1$$

Furthermore, the minimised quantity is 1 when a = -1, hence  $a_{2,3} = 1$ .

#### 3.3 **Proof of Proposition 10**

Our proof of Proposition 10, restated below, very closely follows [FJ97, Corollary 6, Corollary 7].

**Proposition 10.** 
$$\alpha_{k\ell} \ge 1 - \frac{1}{\ell} + \frac{2\ln\ell}{k\ell} - o\left(\frac{\ln\ell}{k\ell}\right) - O\left(\frac{1}{k^2}\right)$$

The following result follows from the analysis in [FJ97, Lemma 5, Corollary 6, Corollary 7].

**Theorem 16.** The Taylor series for  $N_{\ell}(x)$ , given by

$$N_{\ell}(x) = \sum_{i=0}^{\infty} c_i x^i$$

*converges for*  $-1 \le x \le 1$ . *Every*  $c_i \ge 0$ . *Furthermore*  $c_0 = 1/\ell$ ,  $c_1 \sim 2 \ln \ell/(\ell - 1)$ , and  $\sum_{i=0}^{\infty} c_{2i} = 1/2$ .

The following fact was observed in [FJ97]; we include a proof for completeness.

**Lemma 17.** For  $0 \le a \le 1$ , we have  $\frac{k-1}{k}(1-a) \le 1 - N_{\ell}(a)$ .

*Proof.* We first wish to find  $N_{\ell}(0)$ ,  $N_{\ell}(1)$ .  $P_{\ell}(0)$  is just the probability that  $x_1 \ge x_i$  and  $y_1 \ge y_i$  for  $x_1, \ldots, x_{\ell}, y_1, \ldots, y_{\ell} \sim \mathcal{N}(0, 1)$ . By symmetry these events occur with probability  $1/\ell$  each, and thus overall they occur with probability  $1/\ell^2$ . On the other hand,  $P_{\ell}(1)$  is just the probability that  $x_1 \ge x_i$  for  $x_1, \ldots, x_{\ell} \sim \mathcal{N}(0, 1)$ . By symmetry this is  $1/\ell$ .

Observe that since every term in the Taylor series of  $N_{\ell}$  is nonnegative, the function is convex on [0, 1], hence  $1 - N_{\ell}(a)$  is concave. Since furthermore  $(k - 1)(1 - 0)/k = 1 - 1/k \le 1 - 1/\ell = 1 - N_{\ell}(0)$  and  $(k - 1)(1 - 1)/k = 0 \le 0 = 1 - N_{\ell}(1)$ , by Jensen's inequality we have, for 0 < a < 1, that

$$\frac{k-1}{k}(1-a) < 1 - \ell N_{\ell}(a).$$

*Proof of Proposition 10.* First observe that we only need to prove this result for large enough k; for all small k we can just force the bound to hold by increasing the  $o(\cdot)$  term arbitrarily. Thus we will prove that the bound holds only for large enough k. We will try to find some  $R \in [1/2, 1]$  such that

$$R\frac{k-1}{k}(1-a) \le 1 - \ell P_{\ell}(a) = 1 - N_{\ell}(a),$$
(5)

for  $-1/(k-1) \le a \le 1$ . We will then conclude  $\alpha_{k\ell} \ge R$ . (The stipulation that  $R \ge 1/2$  will be necessary later; it is justified by the fact that at the end we will find such an R.)

By Lemma 17, (5) is true for any  $0 \le R \le 1$  and  $0 \le a \le 1$ . In other words, we need only to care about  $-1/(k-1) \le a \le 0$ ; thus assume that this is the case.

Now, for  $-1/(k-1) \le a \le 0$ , we have  $a^{2i} \le a^2$  and  $a^{2i+1} \le 0$ ; since we know the first two coefficients of the Taylor series of  $N_{\ell}$ , and the sum of the even coefficients, by ignoring the higher-order odd terms and summing together the even terms we can deduce therefore that

$$N_{\ell}(a) \le \frac{1}{\ell} + (1 + \varepsilon(\ell)) \frac{2\ln\ell}{\ell} a + \frac{a^2}{2},\tag{6}$$

where  $\lim_{\ell \to \infty} \varepsilon(\ell) = 0$ . (This is because we know that the first-order coefficient is of order  $2 \ln \ell / (\ell - 1) \sim 2 \ln \ell / \ell$ .) We suppress the  $\ell$  in  $\varepsilon(\ell)$  henceforth.

By substituting (6) into (5) and factoring out (-a), we get the following sufficient condition on R

$$1 - \frac{1}{\ell} + (-a) \underbrace{\left( (1+\varepsilon) \frac{2\ln\ell}{\ell} + \frac{a}{2} \right)}_{A} \ge \frac{k-1}{k} (1-a)R.$$

Observe that the left-hand side is a linear function of A with nonnegative slope; thus by substituting A with its minimum value we get another sufficient condition on R. Observe that the value of a that minimises A is a = -1/(k - 1), i.e. the minimum value. Hence the following holding for all  $-1/(k - 1) \le a \le 0$  is a sufficient condition for R:

$$1 - \frac{1}{\ell} + (-a)\left((1+\varepsilon)\frac{2\ln\ell}{\ell} - \frac{1}{2(k-1)}\right) \ge \frac{k-1}{k}(1-a)R.$$

Now subtract -a(k-1)R/k to get that the following must hold

$$1 - \frac{1}{\ell} + (-a) \underbrace{\left( (1+\varepsilon) \frac{2\ln\ell}{\ell} - \frac{1}{2(k-1)} - \frac{k-1}{k} R \right)}_{B} \ge \frac{k-1}{k} R.$$

$$\tag{7}$$

*B* is negative for large enough *k*, as  $\ell \ge k$  and  $R \ge 1/2$ . Hence to minimise the left-hand side of (7) we must take a = -1/(k-1) again. Thus it is a sufficient condition on *R* that

$$1 - \frac{1}{\ell} + (1 + \varepsilon) \frac{2\ln \ell}{\ell(k-1)} - \frac{1}{2(k-1)^2} - \frac{R}{k} \ge \frac{k-1}{k} R.$$

Add R/k and reverse the bound to find the sufficient condition

$$R \le 1 - \frac{1}{\ell} + (1 + \varepsilon) \frac{2 \ln \ell}{\ell (k - 1)} - \frac{1}{2(k - 1)^2}$$

Rearrange again to find the sufficient condition

$$R \leq 1 - \frac{1}{\ell} + \frac{2\ln\ell}{\ell(k-1)} + \frac{2\varepsilon(\ell)\ln\ell}{\ell(k-1)} - \frac{1}{2(k-1)^2} = 1 - \frac{1}{\ell} + \frac{2\ln\ell}{\ell k} + \underbrace{\left(\frac{2\ln\ell}{\ell k(k-1)} + \frac{2\varepsilon(\ell)\ln\ell}{\ell(k-1)} - \frac{1}{2(k-1)^2}\right)}_{C}$$

Observe that  $C = O(\ln \ell/k^2 \ell) + o(\ln \ell/k \ell) + O(1/k^2) = o(\ln \ell/\ell k) + O(1/k^2)$ . Hence, since this condition is a sufficient condition on *R*, we find that for large enough *k*,

$$\alpha_{k\ell} \ge 1 - \frac{1}{\ell} + \frac{2\ln\ell}{k\ell} - o\left(\frac{\ln\ell}{k\ell}\right) - O\left(\frac{1}{k^2}\right).$$

### 3.4 **Proof of Proposition 11**

Now, we prove Proposition 11.

**Proposition 11.**  $\alpha_{k\ell} > 1 - 1/\ell$ .

This proposition serves a role analogous to [FJ97, Corollary 6] (which is equivalent to the case  $k = \ell$ ). We believe that our proof of this fact is simpler; also the direct generalisation of the proof in [FJ97] does not seem to work for k much smaller than  $\ell$ . We will first need a technical lemma.

**Lemma 18.** For  $-1 \le a \le 0$ , we have  $N_{\ell}(a) \le 1/\ell$ , with equality only at a = 0.

*Proof.* First, recall that  $N_{\ell}(0) = 1/\ell$ . Note that  $N_{\ell}(a)$  is the probability that  $\arg \max_i x_i = \arg \max_i ax_i + \sqrt{1-a^2} y_i$ , where  $x_1, \ldots, x_{\ell}, y_1, \ldots, y_{\ell} \sim \mathcal{N}(0, 1)$  are i.i.d. variables. Now, suppose without loss of generality that the values of  $(x_i)_{i \in [\ell]}$  are fixed, and in particular  $x_1 > \cdots > x_{\ell}$  (the inequalities are strict with probability 1). Letting  $A = -a/\sqrt{1-a^2} \ge 0$ , we have that  $N_{\ell}(a)$  is just  $\Pr[\arg \max_i y_i - Ax_i = 1]$ . Now, fix  $y_1$ , and note that conditional on this, the probability above becomes

$$\prod_{i=2}^{\ell} \Pr[y_1 - Ax_1 > y_i - Ax_i] = \prod_{i=2}^{\ell} \Pr[y_i < y_1 - A(x_1 - x_i)]$$

by independence. As  $x_1 - x_i > 0$ , term-by-term this probability is maximised at A = a = 0 (and only there). Since all the probabilities are nonzero, we get that the only *a* that minimises this expression is a = 0. Hence, after integrating over all possible choices of  $y_1$ , we get that  $N_{\ell}(a) \le N_{\ell}(0) = 1/\ell$ , with equality only at a = 0.

*Proof of Proposition 11.* We wish to prove that, for  $-1/(k-1) \le a < 1$ ,

$$\frac{k-1}{k}\frac{\ell-1}{\ell}(1-a) < 1 - N_{\ell}(a).$$

For  $a \in [0, 1)$ , this follows immediately by Lemma 17, so assume a < 0. For such a, we know that  $N_{\ell}(a) < 1/\ell$ . Furthermore, note that  $0 \le (k - 1)(1 - a)/k \le 1$  for our choice of a, so

$$\frac{k-1}{k}\frac{\ell-1}{\ell}(1-a) \le \frac{\ell-1}{\ell} = 1 - \frac{1}{\ell} < 1 - N_{\ell}(a).$$

## 4 Derandomisation

In this section, we will show how to derandomise our algorithm and thus establish Theorem 2.<sup>14</sup>

We start with briefly comparing our approach to previous work. There is a line of work that uses pseudorandom generators for derandomisation, such as the work of Engebretsen, Indyk, and O'Donnell [EIO02], and Gopalan, O'Donnell, Wu, and Zuckerman [GOWZ10]. In [EIO02], the loss due to derandomisation has to be constant for the method to be polynomial, whereas we want this to be of the order of 1/m, where mis the number of edges of the input graph. The result of [GOWZ10] does not immediately give a polynomialtime algorithm in our setting due to their required seed length. The method of conditional expectations, which we use, was also used by Mahajan and Ramesh [MR99]. In comparison to [MR99], our approach is simple, easy to generalise, and also explicitly works for our rounding scheme (and also for [FJ97], which we use). The approach of Bhargava and Kosaraju [BK05] derandomises conditional probabilities by an approximation of normal distributions via polynomials; we approximate simply just with a scaled bionomial distribution. Finally, there is also the approach of Sivakumar [Siv02], which goes through complexity theory. We believe that our approach is more straightforward overall.

Our goal will be the following general derandomisation theorem.

<sup>&</sup>lt;sup>14</sup>Throughout we will ignore issues of real precision.

**Theorem 19.** Fix a constant d. There exists an algorithm that does the following. Suppose we are given  $n, m \in \mathbb{N}$ ,  $\mathbf{a}_{ij} \in \mathbb{R}^n$  and  $b_{ij}, \varepsilon \in \mathbb{R}$  for all  $i \in [m]$ ,  $j \in [d]$ . Suppose  $\mathbf{x} = (x_1, \ldots, x_d) \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_d)$  and that

$$\sum_{i=1}^{m} \Pr_{\mathbf{x}} \left[ \bigwedge_{j=1}^{d} \mathbf{a}_{ij} \cdot \mathbf{x} > b_{ij} \right] \ge \alpha$$

for some  $\alpha \in \mathbb{R}$ . Then the algorithm computes some particular  $\mathbf{x}^* = (x_1^*, \dots, x_d^*) \in \mathbb{R}^d$  such that

$$\sum_{i=1}^{m} \left[ \bigwedge_{j=1}^{d} \mathbf{a}_{ij} \cdot \mathbf{x}^* > b_{ij} \right] \geq \alpha - \varepsilon,$$

in polynomial time with respect to  $n, m, 1/\varepsilon$ .

To facilitate the proof of Theorem 19, we will need a multidimensional version of the Berry-Esseen theorem. We will use the following version with explicit constants, due to Raič [Rai19].

**Theorem 20** ([Rail9, Theorem 1.1]). Suppose  $\mathbf{t}_1, \ldots, \mathbf{t}_N \in \mathbb{R}^d$  are independent random variables with mean zero, such that the sum of their covariance matrices is  $\mathbf{I}_d$ . Let  $\mathbf{s} = \mathbf{t}_1 + \cdots + \mathbf{t}_N$ . Suppose  $\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_d)$ , and let  $S \subseteq \mathbb{R}^d$  be convex and measurable. Then

$$\left|\Pr_{\mathbf{s}}[\mathbf{s} \in S] - \Pr_{\mathbf{x}}[\mathbf{x} \in S]\right| \le \left(42\sqrt[4]{d} + 16\right) \sum_{i=1}^{N} \mathbf{E}\left[||\mathbf{t}_{i}||^{3}\right].$$

The following is an easy and well-known corollary of Theorem 20: We can approximate a multivariate normal distribution with binomial distributions. For completeness, we provide a proof.

**Corollary 21.** Let  $d \in \mathbb{N}$  be a constant and take  $\varepsilon \in (0, 1)$ . Take

$$N = P_{\varepsilon} \ge \left(\frac{42d^{7/4} + 16d^{3/2}}{\varepsilon}\right)^2 = \frac{\xi_d}{\varepsilon^2},\tag{8}$$

where  $\xi_d = O(d^{7/2})$  depends only on d. Suppose  $s_1, \ldots, s_N \sim \operatorname{NBin}(N)$  are *i.i.d.*, and let  $\mathbf{s} = (s_1, \ldots, s_d)$ . Let  $\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_d)$ . Then for all convex measurable sets  $S \subseteq \mathbb{R}^d$  we have

$$\left|\Pr_{\mathbf{s}}[\mathbf{s}\in S] - \Pr_{\mathbf{x}}[\mathbf{x}\in S]\right| \le \varepsilon.$$

*Proof.* Note that each component of **s** is i.i.d. and distributed as the sum of N independent trials that take value  $\pm 1/\sqrt{N}$  equiprobably. In other words, we can see **s** as the sum  $\mathbf{t}_1 + \cdots + \mathbf{t}_N$ , where  $\mathbf{t}_1, \ldots, \mathbf{t}_N \sim \mathcal{U}(\{-1/\sqrt{N}, 1/\sqrt{N}\}^d)$  are i.i.d. Observe that the covariance matrix of  $\mathbf{t}_i$  is  $\mathbf{I}_d/N$ , so the sum of these covariance matrices for all *i* is  $\mathbf{I}_d$ . Furthermore

$$||\mathbf{t}_i|| = \sqrt{\left(\pm \frac{1}{\sqrt{N}}\right)^2 + \dots + \left(\pm \frac{1}{\sqrt{N}}\right)^2} = \sqrt{\frac{d}{N}}$$

with probability 1. Now, apply Theorem 20 to  $\mathbf{s} = \sum_{i=1}^{N} \mathbf{t}_i$ . We find that

$$\left|\Pr_{\mathbf{s}}[\mathbf{s} \in S] - \Pr_{\mathbf{x}}[\mathbf{x} \in S]\right| \le \left(42\sqrt[4]{d} + 16\right) \sum_{i=1}^{N} \mathbb{E}\left[||\mathbf{t}_{i}||^{3}\right] = \left(42\sqrt[4]{d} + 16\right) N\sqrt{\frac{d}{N}^{3}} = \frac{42d^{7/4} + 16d^{3/2}}{\sqrt{N}}.$$

Substituting (8), it follows that  $|\Pr_{\mathbf{s}}[\mathbf{s} \in S] - \Pr_{\mathbf{x}}[\mathbf{x} \in S]| \le \varepsilon$ , as required.

**Theorem 22.** Fix a constant d, and take  $\mathbf{a}_1, \ldots, \mathbf{a}_d \in \mathbb{R}^n, b_1, \ldots, b_d \in \mathbb{R}, c_1, \ldots, c_d \in \mathbb{R}, \varepsilon \in \mathbb{R}$ . Consider the function

$$p(t) = \Pr_{\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_n)} \left[ \bigwedge_{i=1}^d \mathbf{a}_i \cdot \mathbf{x} + c_i t > b_i \right].$$

There exists a step function  $\hat{p}$  with  $poly(1/\varepsilon)$  steps, where the steps and the values at those steps are computable in polynomial time with respect to  $1/\varepsilon$  and n, such that  $|\hat{p}(t) - p(t)| \le \varepsilon$  for all  $t \in \mathbb{R}$ .

*Proof.* Observe that the tuple  $(\mathbf{a}_1 \cdot \mathbf{x}, \dots, \mathbf{a}_d \cdot \mathbf{x})$  (interpreted as a column vector) is a *d*-variate normally distributed vector; namely, if we let

$$\mathbf{A} = \begin{pmatrix} \mathbf{a}_1^T \\ \vdots \\ \mathbf{a}_d^T \end{pmatrix}$$

be a block matrix whose rows are  $\mathbf{a}_1^T, \ldots, \mathbf{a}_d^T$ , then  $(\mathbf{a}_1 \cdot \mathbf{x}, \ldots, \mathbf{a}_d \cdot \mathbf{x}) = \mathbf{A}\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{A}\mathbf{I}_D\mathbf{A}^T) = \mathcal{N}(\mathbf{0}, \mathbf{A}\mathbf{A}^T)$ .

We can compute the covariance matrix, namely  $\mathbf{A}\mathbf{A}^T$ , in polynomial time with respect to *n*. Now, by computing the Cholesky decomposition of this positive semidefinite matrix, we can find  $\mathbf{A}' \in \mathbb{R}^{d \times d}$  such that  $\mathbf{A}'\mathbf{A}'^T = \mathbf{A}\mathbf{A}^T$ . Thus  $(\mathbf{a}_1 \cdot \mathbf{x}, \dots, \mathbf{a}_d \cdot \mathbf{x}) \sim \mathcal{N}(\mathbf{0}, \mathbf{A}'\mathbf{A}'^T)$ . Letting  $\mathbf{a}_1'^T, \dots, \mathbf{a}_d'^T$  be the rows of  $\mathbf{A}'$ , we find that  $(\mathbf{a}_1 \cdot \mathbf{x}, \dots, \mathbf{a}_d \cdot \mathbf{x})$  is identically distributed to  $(\mathbf{a}_1' \cdot \mathbf{x}', \dots, \mathbf{a}_d' \cdot \mathbf{x}')$ , when  $\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_n)$  and  $\mathbf{x}' \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_d)$ , since both follow the distribution  $\mathcal{N}(\mathbf{0}, \mathbf{A}'\mathbf{A}'^T)$ . Thus,

$$p(t) = \Pr_{\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_d)} \left[ \bigwedge_{i=1}^d \mathbf{a}'_i \cdot \mathbf{x} + c_i t > b_i \right].$$

In other words, we have reduced the dimensionality of our problem from n to d, a constant.

Note that the set defined by  $\bigwedge_{i=1}^{d} \mathbf{a}'_i \cdot \mathbf{x} > b_i + t$  is necessarily convex and measurable, being the intersection of finitely many half-spaces. Thus we can apply Corollary 21. Let  $N = P_{\varepsilon}$  and  $s_1, \ldots, s_N \sim \text{NBin}(N)$ , and suppose  $\mathbf{s} = (s_1, \ldots, s_d)$ . Then, we know that

$$\left| p(t) - \Pr_{\mathbf{s}} \left[ \bigwedge_{i=1}^{d} \mathbf{a}'_{i} \cdot \mathbf{s} + c_{i}t > b_{i} \right] \right| \leq \varepsilon.$$

This suggests using the following definition:

$$\widehat{p}(t) = \Pr_{\mathbf{s}} \left[ \bigwedge_{i=1}^{d} \mathbf{a}'_{i} \cdot \mathbf{s} + c_{i}t > b_{i} \right],$$

as this must satisfy the condition  $|\hat{p}(t) - p(t)| \leq \varepsilon$ . It remains to show that  $\hat{p}$  is a step function, and that these steps can be efficiently computed. Intuitively, this is the case since the probability distribution we define  $\hat{p}$  over is discrete. More precisely, letting

$$D = \{-N/\sqrt{N}, (-N+2)/\sqrt{N}, \dots, (N-2)/\sqrt{N}, N/\sqrt{N}\}^{d}$$

be the domain of s, and letting q be the p.m.f. of s (note that it can be efficiently computed, since s is essentially distributed according to a product distribution of normalised binomials), we note that

$$\widehat{p}(t) = \Pr_{\mathbf{s}}\left[\bigwedge_{i=1}^{d} \mathbf{a}'_{i} \cdot \mathbf{s} + c_{i}t > b_{i}\right] = \sum_{\mathbf{u}\in D} q(\mathbf{u})\left[\bigwedge_{i=1}^{d} \mathbf{a}'_{i} \cdot \mathbf{u} + c_{i}t > b_{i}\right] = \sum_{\mathbf{u}\in D} q(\mathbf{u})\left[c_{i}t > \max_{1 \le i \le d} b_{i} - \mathbf{a}'_{i} \cdot \mathbf{u}\right]$$

Now observe that, for each  $\mathbf{u} \in D$ , the function

$$t \mapsto q(\mathbf{u}) \left[ c_i t > \max_{1 \le i \le d} b_i - \mathbf{a}'_i \cdot \mathbf{u} \right]$$

is a step function with at most one step: if  $c_i = 0$  then the function is a constant (whose value is easy to compute); otherwise the step is at  $(1/c_i) \max_{1 \le i \le d} b_i - \mathbf{a}'_i \cdot \mathbf{u}$ , where the step being increasing or decreasing is determined by the sign of  $c_i$  (and again the values of the function are easy to compute). It therefore follows that  $\hat{p}$  is a step function that has at most  $|D| = P_{\varepsilon}^d = \operatorname{poly}(1/\varepsilon)$  steps, and that each of the values that the function takes can be computed in polynomial time with respect to  $1/\varepsilon$  and n.

*Proof of Theorem 19.* We give a recursive algorithm. If n = 0 then there is nothing to output, so assume  $n \ge 1$ . Let  $\mathbf{x} = (t, \mathbf{x}')$ , and  $\mathbf{a}_{ij} = (c_{ij}, \mathbf{a}'_{ij})$  — in other words, separate out the first variable. We are given

$$\sum_{i=1}^{m} \Pr_{\mathbf{x}',t} \left[ \bigwedge_{j=1}^{d} \mathbf{a}'_{ij} \cdot \mathbf{x}' + c_{ij}t > b_{ij} \right] = \sum_{i=1}^{m} \Pr_{\mathbf{x}} \left[ \bigwedge_{j=1}^{d} \mathbf{a}_{ij} \cdot \mathbf{x} > b_{ij} \right] \ge \alpha,$$

when  $\mathbf{x} = (\mathbf{x}', t) \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_d)$ . But then this must be true for some particular value of t, say  $t^*$ , i.e.

$$\sum_{i=1}^{m} \Pr_{\mathbf{x}'} \left[ \bigwedge_{j=1}^{d} \mathbf{a}'_{ij} \cdot \mathbf{x}' + c_{ij}t^* > b_{ij} \right] \ge \alpha.$$
(9)

Apply Theorem 22 to each of the probabilities above viewed as functions of *t*, with  $\varepsilon' = \varepsilon/2nm$ ; we thus build step functions  $p_1, \ldots, p_m$  in polynomial time with respect to *n* and  $1/\varepsilon' = 2nm/\varepsilon$ , such that

$$\left| p_i(t) - \Pr_{\mathbf{x}'} \left[ \bigwedge_{j=1}^d \mathbf{a}'_{ij} \cdot \mathbf{x}' + c_{ij}t > b_{ij} \right] \right| \le \frac{\varepsilon}{2nm}$$

for all *t*. Add these equations for i = 1, ..., m to find

$$\left|\sum_{i=1}^{m} p_i(t) - \sum_{i=1}^{m} \Pr_{\mathbf{x}'} \left[ \bigwedge_{j=1}^{d} \mathbf{a}'_{ij} \cdot \mathbf{x}' + c_{ij}t > b_{ij} \right] \right| \le \frac{\varepsilon}{2n}.$$
 (10)

Observe that  $\sum_{i=1}^{m} p_i$  is a step function with polynomially many steps with respect to n, m, whose values are also computable in polynomial time. Thus it is easy to find some value  $\hat{t}$  that maximises the expression  $\sum_{i=1}^{m} p_i(\hat{t})$ . By (9) and (10) we have that  $\sum_{i=1}^{m} p_i(\hat{t}) \ge \sum_{i=1}^{m} p_i(t^*) \ge \alpha - \frac{\varepsilon}{2n}$ , and by (10) again we find that

$$\sum_{i=1}^{m} \Pr_{\mathbf{x}'} \left[ \bigwedge_{j=1}^{d} \mathbf{a}'_{ij} \cdot \mathbf{x}' + c_{ij}\hat{t} > b_{ij} \right] \ge \alpha - \frac{\varepsilon}{n}$$

Equivalently,

$$\sum_{i=1}^{m} \Pr_{\mathbf{x}'} \left[ \bigwedge_{j=1}^{d} \mathbf{a}'_{ij} \cdot \mathbf{x}' > b_{ij} - c_{ij} \hat{t} \right] \geq \alpha - \frac{\varepsilon}{n},$$

and we can recursively find optimal values for the remaining random variables in  $\mathbf{x}'$ . Observe that our recursive depth is *n*, that at each level we use polynomial time with respect to *n*, *m*,  $1/\varepsilon$ , and that, finally, at each

step we lose  $\varepsilon/n$  from the sum of our probabilities. These facts together imply the correctness of our general derandomisation procedure.

We note in passing that the total time complexity of our method is exponential in d; however this does not matter, as we consider d a constant.

This is enough to derandomise our algorithm.

**Theorem 2.** Fix  $2 \le k \le l$  and let  $\alpha_{kl}$  be as in Theorem 1. There is a deterministic algorithm which, given a graph G that admits a k-colouring of value  $\rho$  and a real number  $\varepsilon > 0$ , finds an l-colouring of value  $\alpha_{kl} \rho - \varepsilon$  in polynomial time in G and  $1/\varepsilon$ .

*Proof.* Let G = (V, E), where V = [n] and m = |E|. Assume that  $n \ge k$  (otherwise simply check all possible colourings). By the analysis of our randomised algorithm from Theorem 1, using SDP we can find, in polynomial time with respect to G and  $\log(1/\varepsilon) + O(1)$ , a set of vectors  $\mathbf{a}_1, \ldots, \mathbf{a}_n$  such that  $\mathbf{a}_i \cdot \mathbf{a}_i = 1$ ,  $\mathbf{a}_i \cdot \mathbf{a}_i \ge -1/(k-1)$  for  $i \ne j$  and, if  $\mathbf{x}_1, \ldots, \mathbf{x}_k \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_n)$  are normally distributed variables, then

$$\frac{1}{m}\sum_{(i,j)\in E} \Pr_{\mathbf{x}_1,\ldots,\mathbf{x}_{\ell}} \left[ \arg\max_{c} \mathbf{x}_{c} \cdot \mathbf{a}_{i} \neq \arg\max_{c} \mathbf{x}_{c} \cdot \mathbf{a}_{j} \right] \geq \alpha_{k\ell} \rho - \frac{\varepsilon}{2}.$$

Now, let  $\mathbf{x} = (\mathbf{x}_1, \dots, \mathbf{x}_{\ell}) \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_{\ell n})$ , and define  $\mathbf{a}_{ic}$  such that  $\mathbf{x} \cdot \mathbf{a}_{ic} = \mathbf{x}_c \cdot \mathbf{a}_i$ ; in other words, pad out  $\mathbf{a}_i$  with  $(\ell - 1)n$  zeroes. We first claim that the event

$$\arg\max \mathbf{x}_c \cdot \mathbf{a}_i \neq \arg\max \mathbf{x}_c \cdot \mathbf{a}_j$$

can be seen as the disjoint union of  $\ell(\ell - 1)$  intersections of  $2(\ell - 1)$  hyperplanes in the space of **x**. To express it in this way, first fix the value of the respective sides to  $c_0 \neq c_1$ , where  $c_0, c_1 \in [\ell]$ , in  $\ell(\ell - 1)$  ways. Observe that the event that arg max<sub>c</sub>  $\mathbf{x}_c \cdot \mathbf{a}_i = c_0$  is the same as

$$\bigwedge_{c\neq c_0} \mathbf{x}_{c_0} \cdot \mathbf{a}_i > \mathbf{x}_c \cdot \mathbf{a}_i.$$

Now, using the notation from before, this is equivalent to

$$\bigwedge_{c\neq c_0} \mathbf{x} \cdot (\mathbf{a}_{ic_0} - \mathbf{a}_{ic}) > 0.$$

It follows that

$$\begin{aligned} \alpha_{k\ell}\rho - \frac{\varepsilon}{2} &\leq \frac{1}{m} \sum_{(i,j)\in E} \Pr_{\mathbf{x}_{1},...,\mathbf{x}_{\ell}} \left[ \arg\max_{c} \mathbf{x}_{c} \cdot \mathbf{a}_{i} \neq \arg\max_{c} \mathbf{x}_{c} \cdot \mathbf{a}_{j} \right] \\ &= \frac{1}{m} \sum_{(i,j)\in E} \sum_{c_{0}\neq c_{1}} \Pr_{\mathbf{x}_{1},...,\mathbf{x}_{\ell}} \left[ \bigwedge_{c\neq c_{0}} \mathbf{x} \cdot (\mathbf{a}_{ic_{0}} - \mathbf{a}_{ic}) > 0 \land \bigwedge_{c\neq c_{1}} \mathbf{x} \cdot (\mathbf{a}_{jc_{1}} - \mathbf{a}_{jc}) > 0 \right]. \end{aligned}$$

By Theorem 19 for  $d = 2(\ell - 1)$ , in polynomial time with respect to  $n, m, 1/\varepsilon$ , we can find particular values **x**<sup>\*</sup> such that

$$\frac{1}{m}\sum_{(i,j)\in E}\sum_{c_0\neq c_1}\left[\bigwedge_{c\neq c_0}\mathbf{x}^*\cdot(\mathbf{a}_{ic_0}-\mathbf{a}_{ic})>0\wedge\bigwedge_{c\neq c_1}\mathbf{x}^*\cdot(\mathbf{a}_{jc_1}-\mathbf{a}_{jc})>0\right]\geq \alpha_{k\ell}\rho-\frac{\varepsilon}{2}-\frac{\varepsilon}{2}=\alpha_{k\ell}\rho-\varepsilon.$$

Defining  $(\mathbf{x}_1^*, \ldots, \mathbf{x}_{\ell}^*) = \mathbf{x}^*$ , this is equivalent to

$$\frac{1}{m} \sum_{(i,j)\in E} [\arg\max_{c} \mathbf{x}_{c}^{*} \cdot \mathbf{a}_{i} \neq \arg\max_{c} \mathbf{x}_{c}^{*} \cdot \mathbf{a}_{j}] \geq \alpha_{k\ell} \rho - \varepsilon.$$

In other words, if we set the colour of vertex *i* to  $\arg \max_{c} \mathbf{x}_{c}^{*} \cdot \mathbf{a}_{i}$ , then the resulting  $\ell$ -colouring will have value  $\alpha_{k\ell} \rho - \varepsilon$ , as required.

## 5 Algorithm for fixed k and large $\ell$

We show the following theorem.

**Theorem 4.** Let k > 2 be fixed<sup>15</sup> and  $\ell \ge k$  be large. There is a deterministic algorithm which, given a graph G that admits a k-colouring of value  $\rho$  and a real number  $\varepsilon > 0$ , finds an  $\ell$ -colouring of G of value  $\alpha'_{k\ell}\rho - \varepsilon$  in polynomial time in G and  $1/\varepsilon$ . In particular, for a fixed k there exists a constant  $u_k > 1$  such that

$$\alpha_{k\ell}' \geq 1 - O\left(1/\ell^{u_k}\right),\,$$

which is  $1 - o(1/\ell)$  when  $\ell$  grows large.<sup>16</sup>

Note that Theorem 4 together with Theorem 5 do not contradict the UGC and P  $\neq$  NP, since Theorem 5 only works for bounded  $\ell$ .

*Proof.* As always, we will only care about when  $\ell$  grows large. We will first give a randomised algorithm, and then derandomise it. We solve the same semi-definite program as in Theorem 1 (which is also the same as in [FJ97, KMS98]), i.e.

$$\max \quad \frac{1}{m} \sum_{(i,j) \in E} \frac{k-1}{k} (1 - \mathbf{a}_i \cdot \mathbf{a}_j)$$
  
s.t. 
$$\mathbf{a}_i \cdot \mathbf{a}_i = 1,$$
$$\mathbf{a}_i \cdot \mathbf{a}_j \ge -\frac{1}{k-1}, i \neq j,$$
$$\mathbf{a}_i \in \mathbb{R}^n.$$

As in Theorem 1, the value is at least  $\rho$ . We now randomly round in the following way: Sample  $t = \lfloor \log_2(\ell) \rfloor$ random hyperplanes that pass through the origin  $H_1, \ldots, H_t$  in n dimensions; then, to colour node i check on which side of  $H_1, \ldots, H_t$  the vector  $\mathbf{a}_i$  is, and depending on this assign a unique colour. Note that we use  $2^t \leq \ell$  colours in this way. Fix an edge (i, j) and consider  $a = \mathbf{a}_i \cdot \mathbf{a}_j$ ; what is the probability that the colours assigned to i and j are different? Note that the probability that  $\mathbf{a}_i, \mathbf{a}_j$  are separated by one hyperplane among  $H_1, \ldots, H_t$  is just  $\frac{1}{\pi} \arccos a$  (this observation is originally from [GW95]). So the probability that  $\mathbf{a}_i, \mathbf{a}_j$  will be separated by at least one hyperplane is

$$1 - \left(1 - \frac{1}{\pi} \arccos a\right)^t.$$

<sup>&</sup>lt;sup>15</sup>The case  $k = 2, \ell \ge 2$  is covered by Theorem 2.

<sup>&</sup>lt;sup>16</sup>For example, for k = 3 we have  $u_k = 2$  and the approximation ratio is  $1 - O(1/\ell^2)$ . This is significantly better, for large  $\ell$ , than the random guessing algorithm which has performance  $1 - 1/\ell$ .

Now, the approximation ratio is given by

$$\alpha'_{k\ell} = \min_{-1/(k-1) \le a < 1} \frac{k(1 - (1 - \arccos a/\pi)^t)}{(k-1)(1-a)}.$$

We first deal with *a* around a neighbourhood of 1, similarly to Lemma 17. We claim that there exists some  $0 < a_k < 1$  such that

$$\frac{\arccos a}{\pi} \ge \frac{(k-1)(1-a)}{k}$$

for all  $a \in (a_k, 1]$ . Indeed, consider  $\arccos a/\pi - (k-1)(1-a)/k$ . The derivative tends to  $-\infty$  as  $a \to 1$  (from below), so for some neighbourhood of 1 the derivative is negative. Suppose  $(a_k, 1]$  is this neighbourhood. Thus the function is decreasing on this interval. Since the function is equal to 0 at 1, our conclusion follows.

Now, observe that  $\arccos a/\pi \in [0, 1]$ , hence

$$1 - \left(1 - \frac{\arccos a}{\pi}\right)^t \ge 1 - \left(1 - \frac{\arccos a}{\pi}\right) = \frac{\arccos a}{\pi} \ge \frac{(k-1)(1-a)}{k}$$

when  $a \in (a_k, 1]$ . Hence the expression minimised in the definition of  $\alpha_{k\ell}$  is at least 1 whenever  $a \in (a_k, 1]$ , and thus does not affect the value of  $\alpha'_{k\ell}$ . We now focus on the case  $-1/(k-1) < a \le a_k < 1$ .

Define

$$f(a) = \frac{1 - (1 - \arccos a/\pi)^t}{1 - a}.$$

Observe that

$$(a-1)^{2} f'(a) = 1 - \frac{t}{\pi} \sqrt{\frac{1-a}{1+a}} \left(1 - \frac{\arccos(a)}{\pi}\right)^{t-1} - \left(1 - \frac{\arccos(a)}{\pi}\right)^{t}.$$

Note that for large enough t (i.e. large enough  $\ell$ ), we have that f'(a) > 0 for  $-1/(k-1) \le a \le a_k < 1$ . (The size required of  $\ell$  depends on  $a_k$  and hence on k.) Hence,

$$\alpha_{k\ell}' = \frac{k}{k-1} \frac{1 - \left(1 - \frac{1}{\pi} \arccos\left(-\frac{1}{k-1}\right)\right)^t}{\left(1 + \frac{1}{k-1}\right)} = 1 - \left(1 - \frac{1}{\pi} \arccos\left(-\frac{1}{k-1}\right)\right)^t.$$

(This value is indeed less than 1, so  $a \in (a_k, 1]$  did not matter.) We now observe that for any fixed k > 2,

$$X_k \coloneqq 1 - \frac{1}{\pi} \arccos\left(-\frac{1}{k-1}\right) \in \left(0, \frac{1}{2}\right).$$

Define  $u_k = -\log_2(X_k) > 1$ . Hence

$$\alpha'_{k\ell} = 1 - X_k^t = 1 - 2^{\log_2 X_k \lfloor \log_2 \ell \rfloor}.$$

Observe that

$$\log_2 X_k \lfloor \log_2 \ell \rfloor \le \log_2 X_k (-1 + \log_2 \ell) = -\log_2 X_k + \log_2 X_k \log_2 \ell = -\log_2 X_k - u_k \log_2 \ell.$$

Thus

$$\alpha'_{k\ell} \ge 1 - 1/(X_k \ell^{u_k}) = 1 - O(1/\ell^{u_k})$$

We now turn to derandomising this algorithm. It is sufficient to show that the event that two vectors  $\mathbf{a}_i$ ,  $\mathbf{a}_j$  are properly cut by one of the  $H_1, \ldots, H_t$  hyperplanes is the disjoint union of the intersection of constantly many half-spaces in some multivariate normal probability distribution. Then, the derandomisation works precisely as for Theorem 1, using Theorem 19. First, we must express our hyperplanes  $H_1, \ldots, H_t$  in terms of normal variables. As was first observed by [FJ97], a uniformly random hyperplane  $H_i$  can be sampled by taking the set of points at equal distance between two vectors  $\mathbf{x}_i$ ,  $\mathbf{y}_i \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_n)$ , and the points to one side or the other of the hyperplane are those points closer (in terms of inner product) to  $\mathbf{x}_i$  or  $\mathbf{y}_i$  respectively. We label the two sides of  $H_i$  with 0 and 1, with side 0 containing  $\mathbf{x}_i$  and side 1 containing  $\mathbf{y}_i$ , and take the convention that if a vector is on  $H_i$  then it is on side 0. Then the event that  $\mathbf{a}_i$  is on side 0 of  $H_j$  is

$$\mathbf{a}_i \cdot \mathbf{x}_j \geq \mathbf{a}_i \cdot \mathbf{y}_j.$$

Call this event E(i, j, 0), and the complementary event E(i, j, 1). If as before we write

$$\mathbf{x} = (\mathbf{x}_1, \ldots, \mathbf{x}_n, \mathbf{y}_1, \ldots, \mathbf{y}_n) \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_{2tn}),$$

then each event E(i, j, x) is equivalent to **x** belonging to a half-space. Now, the event that vertex *i* is assigned colour *c* (call it E(i, c)) is equivalent to a conjunction of *t* of these events, one for each hyperplane  $H_1, \ldots, H_t$ . Furthermore, the event that  $\mathbf{a}_i, \mathbf{a}_j$  are properly cut is equivalent to the disjoint union of at most  $2^t \times 2^t$  of conjunctions of these events, namely

$$\bigvee_{c\neq c'} E(i,c) \wedge E(j,c').$$

Hence we can derandomise as before, and our conclusion follows.

## 6 Hardness

In this section we use the approach of Khot, Kindler, Mossel and O'Donnell [KKMO07] and of Guruswami and Sinop [GS13] respectively to prove the following hardness result.

**Theorem 5.** Fix some function  $M(k) = o(e^{\sqrt[3]{k}})$ . Let  $2 \le k \le \ell$  be such that  $\ell \le M(k)$ . For any small enough  $\varepsilon > 0$ , consider the problem of deciding whether a given graph G admits a k-colouring of value  $1 - \varepsilon$ , or not even an  $\ell$ -colouring of value  $\beta + \varepsilon$ . We have the following.

• Assuming the UGC, the problem is NP-hard for 
$$\beta = \beta_{k\ell} = 1 - \frac{1}{\ell} + \frac{2\ln\ell}{k\ell} + o\left(\frac{\ln\ell}{k\ell}\right)$$

• Unconditionally, the problem is NP-hard for 
$$\beta = \beta'_{k\ell} = 1 - \frac{1}{\ell} + \frac{8 \ln \ell}{k\ell} + o\left(\frac{\ln \ell}{k\ell}\right)$$
.

Both of these results only hold when  $\beta_{k\ell}, \beta'_{k\ell} \in (0, 1)$ .<sup>17</sup>

The conditional bound in the first bullet point matches the bound in Theorem 1 (up to the asymptotic error terms), but only when  $\ell$  is bounded by some M(k) that is strictly smaller, asymptotically, than the superpolynomial function  $e^{\sqrt[3]{k}}$ . Throughout this entire section, fix the function M(k) and  $2 \le k \le \ell$ .

In the rest of this introduction to Section 6, we give a brief overview of the proof of Theorem 5 and how it differs from existing work. All definitions not stated here explicitly can be found, together with all details, in later subsections. We first recall the definition of label cover.

<sup>&</sup>lt;sup>17</sup>The constants hidden in the expression defining  $\beta_{k\ell}$  depend on  $\mathcal{M}(k), k, \ell$ , but not on  $\varepsilon$ .

**Definition 23.** An instance of *label cover with p-to-1 constraints with domain size r* is a tuple  $I = (V = V_A \cup V_B, E, \pi)$ , where  $(V_A \cup V_B, E)$  is a bipartite graph, and for each edge  $(a, b) \in E$  we have a constraint<sup>18</sup>  $\pi_{a,b} : [pr] \rightarrow [r]$  that is *p*-to-1; i.e., for every  $x \in [r]$  there are precisely *p* values  $y \in [pr]$  such that  $x = \pi_{a,b}(y)$ . We call the instance left-regular if every vertex  $a \in V_A$  has the same degree. A solution to this instance is a mapping *c* that takes  $V_A$  to [r] and  $V_B$  to [pr]. The value of the solution is the proportion of edges  $(a, b) \in E$  with  $c(a) = \pi_{a,b}(c(b))$ . The value of the instance is the maximum value of any solution.

The problem of  $(1 - \eta, \eta)$ -approximating a label cover with *p*-to-1 constraints with domain size *r* is the following: Given an instance *I* of label cover with *p*-to-1 constraints and domain size *r*, decide if its value is at least  $1 - \eta$  or at most  $\eta$ .

The following two are not the original forms of the Unique Games Conjecture or the 2-to-1 Theorem, but they are equivalent to them due to the reductions in [KR08] — the original forms considered weighted non-left-regular label cover instances.

**Conjecture 24** (Unique Games Conjecture (UGC) [Kho02]). For every small  $\eta > 0$ , there exists an  $r \in \mathbb{N}$  such that it is NP-hard to  $(1 - \eta, \eta)$ -approximate a left-regular label cover with 1-to-1 constraints and domain size r.

**Theorem 25** (2-to-1 Theorem [KMS23]). For every small  $\eta > 0$ , there exists an  $r \in \mathbb{N}$  such that it is NP-hard to  $(1 - \eta, \eta)$ -approximate a left-regular label cover with 2-to-1 constraints and domain size r.

Theorem 5 follows from Propositions 26 and 27 stated below. Proposition 26 serves the same role as [KKMO07, Proposition 12] and [GS13, Proposition 3.21]; we reprove it here since the promise version does not immediately follow from the non-promise version, and since Lemma 43 (needed in the proof of Proposition 26) fixes a small bug in the published proofs. On the other hand, Proposition 27 is a standard PCP construction, analogous to [KKMO07, Section 11.4] and [GS13, Section 3.4]; we prove it here in a unified way (covering simultaneously 2-to-1 and 1-to-1 constraints) rather than repeating most of the proof twice. The way the unification works is similar to the proof in [GS20].

**Proposition 26.** Fix  $p \in \{1, 2\}$  and  $2 \le k \le \ell$  such that  $\ell \le M(k)$ . Suppose T is a symmetric Markov operator on  $[k^p]$  with spectral radius 0 < c/(k-1) < 1, where  $c \le 4$ . Then there exists  $\tau > 0$  and  $d \in \mathbb{N}$  such that for any  $f : [k^p]^r \to \Delta_{\ell}$  with  $\operatorname{Inf}_i^{\le d}(f) \le \tau$  for all  $i \in [r]$ , we have

$$\langle f, T^{\otimes n} f \rangle \ge \frac{1}{\ell} - \frac{2c \ln \ell}{k\ell} - o\left(\frac{\ln \ell}{k\ell}\right).$$

**Proposition 27.** Let  $p \in \{1, 2\}$  and  $2 \le k \le \ell$ . Assume that there is a colourful symmetric Markov operator T on  $[k^p]$  and  $\tau > 0$ ,  $d \in \mathbb{N}$  such that for any  $f : [k^p]^r \to \Delta_\ell$  with  $\operatorname{Inf}_i^{\le d}(f) \le \tau$  for all  $i \in [n]$ , we have that  $\langle f, T^{\otimes r} f \rangle \ge 1 - \beta$  for some  $\beta \in (0, 1)$ . Assume further that all the values of T are nonnegative integer multiples of a rational number L. Then, assuming that  $(1 - \eta, \eta)$ -approximating a label cover instance with p-to-1 constraints is NP-hard, for any small enough  $\varepsilon$ , it is NP-hard to decide whether a given graph G has a k-colouring of value  $1 - \varepsilon$ , or not even an  $\ell$ -colouring of value  $\beta + \varepsilon$ .

<sup>&</sup>lt;sup>18</sup>We note that alternatively we could have defined our constraints as *p*-to-1 relations  $\pi \subseteq [pr] \times [r]$  i.e. relations where for every  $x \in [r]$  there exist exactly *p* values  $y \in [pr]$  such that  $(y, x) \in \pi$ , and furthermore every  $y \in [pr]$  corresponds to exactly one  $x \in [r]$  such that  $(y, x) \in \pi$ . To translate from this view to ours, map every  $x \in [pr]$  to that  $y \in [r]$  such that  $(x, y) \in \pi$ ; to translate from our view to this one, take the graph of the function we use as a constraint.

The Markov operators mentioned above will be given in the following theorems. The notion of a "colourful" Markov chain merely unifies two properties that we are interested in for our PCP construction. This property appears without a name also in [GS20, Lemma 10].<sup>19</sup>

**Theorem 28.** Fix  $k \ge 2$ . The Bonami-Beckner operator  $T_{-1/(k-1)}$  on [k] is a symmetric Markov operator that is colourful with spectral radius 1/(k-1). Furthermore, all the values in the matrix are nonnegative integer multiples of 1/(k-1).

**Theorem 29** ([GS13, Lemma 3.8]). Fix  $k \ge 6$ . There exists a colourful symmetric Markov operator on  $[k^2]$ whose spectral radius is at most 4/(k-1). Furthermore, all the values in the matrix are nonnegative integer multiples of 1/(k-1)(k-2)(k-3).

The proof of Theorem 5 thus follows by combining all the facts listed above, together with the observation that we can take k to be large, since the fact that  $\ell$  is bounded by a function of k means that the asymptotic term can be increased for small k to make the theorem hold for small k.

#### 6.1 Fourier-analytic notions

We closely follow the exposition of Fourier analysis on discrete domains from [DMR06]. We also include some results from [GS13]. We diverge from these only in that they number their colours  $0, \ldots, k-1$ , whereas we number them  $1, \ldots, k$ ; also we simplify the notation for our Fourier coefficients.

We will be looking in general at functions of the form  $[D]^r \to [\ell]$  or  $[D]^r \to [0,1]$ . Note that  $[\ell]$  can be naturally embedded in the set of probability distributions over  $[\ell]$ , which can be seen as the set  $\Delta_{\ell} = \{(x_1, \dots, x_{\ell}) \mid \sum_i x_i = 1, x_i \ge 0\}$ . Such functions form a vector space under point-wise addition and multiplication, and they have a natural inner product, namely

$$\langle f, g \rangle = \mathbb{E}_{\mathbf{x} \sim \mathcal{U}([D]^r)} \left[ f(\mathbf{x}) \cdot g(\mathbf{x}) \right].$$

The inner product induces a norm  $|| \cdot ||_2$ . In general we will assume that a variable **x** that is mentioned in an expectation will be taken uniformly at random from an appropriate set. Observe that if  $f(\mathbf{x}) =$  $(f^{1}(\mathbf{x}), ..., f^{\ell}(\mathbf{x}))$  and  $g(\mathbf{x}) = (g^{1}(\mathbf{x}), ..., g^{\ell}(\mathbf{x}))$ , then

$$\langle f,g\rangle = \sum_{i=1}^{\ell} \langle f^i,g^i\rangle.$$

So in particular  $||f||_2^2 = \sum_{i=1}^{\ell} ||f_i||_2^2$ . For any two functions  $f : A \to \mathbb{R}$ ,  $g : B \to \mathbb{R}$  we define  $f \otimes g : A \times B \to R$  by  $(f \otimes g)(a, b) =$ f(a)g(b); thus  $\otimes$  is a tensor product. For every [D], fix some orthonormal (under  $\langle \cdot, \cdot \rangle^{20}$ ) basis of the set of functions  $[D] \to \mathbb{R}$ , namely some functions  $\alpha_1, \ldots, \alpha_D$ , such that  $\alpha_1(1) = \ldots = \alpha_1(D) = 1.^{21}$  For  $\mathbf{x} \in [D]^r$  we define  $\alpha_{\mathbf{x}} : [D]^r \to \mathbb{R}$  by

$$\alpha_{\mathbf{x}} = \alpha_{x_1} \otimes \cdots \otimes \alpha_{x_r}$$

<sup>&</sup>lt;sup>19</sup>[G\$13, Lemma 3.8] constructs a Markov chain that is colourful in our terminology, and implicitly uses this fact in their PCP construction. However their lemma states merely that the Markov chain has diagonal elements equal to zero, which is not by itself enough to make the PCP reduction work.

<sup>&</sup>lt;sup>20</sup>The only difference between orthonormality of function  $[D] \to \mathbb{R}$  under  $\langle \cdot, \cdot \rangle$  and of  $\mathbb{R}^D$  under the normal inner product is a matter of normalisation.

<sup>&</sup>lt;sup>21</sup>In Lemma 38, and only there, we will need a particular choice of  $\alpha_1, \ldots, \alpha_D$ . However, the lemma does not mention any quantity dependant on this choice in its statement, so the choice is "contained" within that lemma.

or equivalently

$$\alpha_{\mathbf{x}}(y_1,\ldots,y_r) = \prod_{i=1}^r \alpha_{x_i}(y_i).$$

It can be seen that for  $\mathbf{x} \neq \mathbf{y}$  we have that  $\alpha_{\mathbf{x}} \perp \alpha_{\mathbf{y}}$ . Furthermore, since there are  $[D]^r$  such function (i.e. the same as the dimension of the set of functions from  $[D]^r$  to  $\mathbb{R}$ ), for any  $f : [D]^r \to \mathbb{R}$  we have

$$f = \sum_{\mathbf{x} \in [D]^r} \langle f, \alpha_{\mathbf{x}} \rangle \alpha_{\mathbf{x}} = \sum_{\mathbf{x} \in [D]^r} \hat{f}(\mathbf{x}) \alpha_{\mathbf{x}},$$

where  $\hat{f}(\mathbf{x}) = \langle f, \alpha_{\mathbf{x}} \rangle$ . This  $\hat{f}(\mathbf{x})$  is known as a Fourier coefficient. Clearly  $\overline{af + bg} = a\hat{f} + b\hat{g}$ , hence  $\hat{i}$  is linear. We have a version of Parseval's identity now, since  $\{\alpha_{\mathbf{x}}\}_{\mathbf{x}}$  forms a basis:

$$||f||_2^2 = \sum_{\mathbf{x} \in [D]^r} \hat{f}^2(\mathbf{x})$$

Indeed, in general we have that

$$\langle f, g \rangle = \sum_{\mathbf{x} \in [D]^r} \hat{f}(\mathbf{x}) \hat{g}(\mathbf{x}).$$

We also generalise our notion of Fourier coefficient to functions  $f : [D]^r \to \mathbb{R}^{\ell}$ . If  $f(\mathbf{x}) = (f_1(\mathbf{x}), \dots, f_{\ell}(\mathbf{x}))$ , then

$$\hat{f}(\mathbf{x}) = (\hat{f}_1(\mathbf{x}), \dots, \hat{f}_\ell(\mathbf{x}))$$

With these, Parseval's inequality generalises to

$$||f||_{2}^{2} = \sum_{i=1}^{\ell} ||f_{i}||_{2}^{2} = \sum_{i=1}^{\ell} \sum_{\mathbf{x} \in [D]^{r}} \hat{f}_{i}^{2}(\mathbf{x}) = \sum_{\mathbf{x} \in [D]^{r}} |\hat{f}(\mathbf{x})|^{2}.$$

We now introduce the notion of *low-degree influence*. First for  $\mathbf{x} \in [D]^r$ , we let  $|\mathbf{x}|$  to be the number of coordinates  $i \in [r]$  such that  $\mathbf{x}_i \neq 1$ . With this in hand, for  $f : [D]^r \to \mathbb{R}$ ,  $i \in [r]$ ,  $d \leq r$ , we define

$$\operatorname{Inf}_{i}^{\leq d}(f) = \sum_{\substack{\mathbf{x} \in [D]^{r} \\ \mathbf{x}_{i} \neq 1 \\ |\mathbf{x}| \leq d}} \hat{f}^{2}(\mathbf{x}).$$

This definition does not depend on the basis  $\alpha_1, \ldots, \alpha_D$  taken initially (only that  $\alpha_1(x) = 1$ ); for details see [DMR06, Definition 2.5].

We now generalise this definition to functions of the form  $f : [D]^r \to \mathbb{R}^{\ell}$ ; for such a function suppose  $f(\mathbf{x}) = (f_1(\mathbf{x}), \dots, f_d(\mathbf{x}))$ . Then, for  $i \in [r], d \leq r$ , we define

$$\operatorname{Inf}_{i}^{\leq d}(f) = \sum_{j=1}^{\ell} \operatorname{Inf}_{i}^{\leq d}(f_{j})$$

Observe that

$$\operatorname{Inf}_{i}^{\leq d}(f) = \sum_{j=1}^{\ell} \sum_{\substack{\mathbf{x} \in [D]^{r} \\ \mathbf{x}_{i} \neq 1 \\ |\mathbf{x}| \leq d}} \hat{f_{j}}^{2}(\mathbf{x}) = \sum_{\substack{\mathbf{x} \in [D]^{r} \\ \mathbf{x}_{i} \neq 1 \\ |\mathbf{x}| \leq d}} \sum_{j=1}^{\ell} \hat{f_{j}}^{2}(\mathbf{x}) = \sum_{\substack{\mathbf{x} \in [D]^{r} \\ \mathbf{x}_{i} \neq 1 \\ |\mathbf{x}| \leq d}} |\hat{f}(\mathbf{x})|^{2}$$

Next, we deduce some classic inequalities for sums of low-level influences. Consider  $f : [D]^r \to \Delta_{\ell}$ . Note that

$$||f||_{2}^{2} = \mathbb{E}[|f(\mathbf{x})|^{2}] \le 1,$$

since for any  $\mathbf{y} \in \Delta_{\ell}$  we have  $|\mathbf{y}|^2 \leq 1$ . Thus, by Parseval's identity, we have

$$1 \ge ||f||_2^2 = \sum_{\mathbf{x} \in [D]} |\hat{f}(\mathbf{x})|^2$$

Observe that the formula giving  $\text{Inf}_i^{\leq d}(f)$  as a sum of square lengths of Fourier coefficients contains each term in the sum above at most *d* times; since the sum is at most 1, we derive that  $\text{Inf}_i^{\leq d}(f) \leq d$ .

**Minor operations** Consider any vector  $\mathbf{x} \in [D]^r$  and a function  $\pi : [s] \to [r]$ . Then we define  $\mathbf{x}^{\pi} \in [D]^s$  by

$$\mathbf{x}^{\pi} = (\mathbf{x}_{\pi(1)}, \ldots, \mathbf{x}_{\pi(s)}).$$

Furthermore, consider any function  $f : [D]^r \to \mathbb{R}^d$ , and let  $\pi : [r] \to [s]$ . Then define  $f^{\pi} : [D]^s \to \mathbb{R}^d$  by

$$f^{\pi}(\mathbf{x}) = f(\mathbf{x}^{\pi}) = f(x_{\pi(1)}, \dots, x_{\pi(r)}).$$

Observe that when the function  $\pi : [r] \to [r]$  is a bijection, and for function  $f, g : [D]^r \to \mathbb{R}^d$ , we have that

$$\langle f^{\pi}, g^{\pi} \rangle = \mathbb{E}_{\mathbf{x}}[\langle f^{\pi}(\mathbf{x}), g^{\pi}(\mathbf{x}) \rangle] = \mathbb{E}_{\mathbf{x}}[\langle f(\mathbf{x}^{\pi}), g(\mathbf{x}^{\pi}) \rangle] = \mathbb{E}_{\mathbf{x}}[\langle f(\mathbf{x}), g(\mathbf{x}) \rangle] = \langle f, g \rangle.$$

Next, observe that for such  $\pi$ ,

$$\alpha_{\mathbf{x}}^{\pi}(y_1,\ldots,y_r) = \alpha_{\mathbf{x}}(y_{\pi(1)},\ldots,y_{\pi(r)}) = \prod_{i=1}^r \alpha_{x_i}(y_{\pi(i)}) = \prod_{i=1}^r \alpha_{x_{\pi^{-1}(i)}}(y_i) = \alpha_{\mathbf{x}^{\pi^{-1}}}(y_1,\ldots,y_r).$$

Hence  $\alpha_{\mathbf{x}}^{\pi} = \alpha_{\mathbf{x}^{\pi^{-1}}}$ . Furthermore this implies that for bijective  $\pi$ ,

$$\widehat{f^{\pi}}(\mathbf{x}) = \langle f^{\pi}, \alpha_{\mathbf{x}} \rangle = \langle f, \alpha_{\mathbf{x}}^{\pi^{-1}} \rangle = \langle f, \alpha_{\mathbf{x}^{\pi}} \rangle = \widehat{f}(\mathbf{x}^{\pi}).$$

**Coordinate regrouping lemmas** We reuse and slightly modify the notation from [DMR06, Definition 2.6], which reappears in [GS13, Definition 3.22]. The notation there does not include the (2) we use — this will be important for us since what we do is in greater generality.

We will implicitly use the fact that  $[k^2] \cong [k]^2$ ; fix some arbitrary bijection between the two. For  $\mathbf{x} \in [k]^{2r}$ , we define  $\overline{\mathbf{x}}^{(2)} \in [k^2]^r \cong ([k]^2)^r$  by

$$\overline{\mathbf{x}}^{(2)} = ((x_1, x_2), \dots, (x_{2r-1}, x_{2r})).$$

Conversely, for  $\mathbf{x} \in [k^2]^r \cong ([k]^2)^r$ , where  $\mathbf{x} = ((x_1, x'_1), \dots, (x_n, x'_n))$ , we have

$$\underline{\mathbf{x}}_{(2)} = (x_1, x'_1, \dots, x_n, x'_n).$$

From the definitions,  $\underline{\mathbf{x}}_{(2)}^{(2)} = \mathbf{x}$ . Now, we define these transformations on the inputs for functions, as follows. For  $f : [k]^{2r} \to \mathbb{R}^d$ ,  $g : [k^2]^r \to \mathbb{R}^d$ , we have

$$\overline{f}^{(2)}(\mathbf{x}) = f(\underline{\mathbf{x}}_{(2)}),$$
$$\underline{g}_{(2)}(\mathbf{x}) = g(\overline{\mathbf{x}}^{(2)}).$$

Again,  $\overline{\underline{f}}_{(2)}^{(2)} = f$ . Next, we recall the following result.

**Lemma 30** ([DMR06, Claim 2.7]). For  $f : [k]^{2r} \to \mathbb{R}$ ,  $i \in [r]$ , and  $1 \le d \le r$ , we have

$$\operatorname{Inf}_{i}^{\leq d}(\overline{f}^{(2)}) \leq \operatorname{Inf}_{2i-1}^{\leq d}(f) + \operatorname{Inf}_{2i}^{\leq d}(f).$$

The following is an obvious corollary.

**Corollary 31.** For  $f : [k]^{2r} \to \mathbb{R}^{\ell}$ ,  $i \in [r]$ , and  $1 \le d \le r$ , we have

$$\operatorname{Inf}_{i}^{\leq d}(\overline{f}^{(2)}) \leq \operatorname{Inf}_{2i-1}^{\leq d}(f) + \operatorname{Inf}_{2i}^{\leq d}(f).$$

*Proof.* Apply the previous lemma to each term in the sum that defines  $Inf_i^{\leq d}$  for functions to  $\mathbb{R}^{\ell}$ .

Now, we define  $\overline{\mathbf{x}}^{(1)} = \underline{\mathbf{x}}_{(1)} = \mathbf{x}$  and likewise  $\overline{f}^{(1)} = \underline{f}_{(1)} = f$ , and thus immediately

$$\operatorname{Inf}_{i}^{\leq d}(\overline{f}^{(1)}) \leq \operatorname{Inf}_{i}^{\leq d}(f).$$

Therefore we find that

**Corollary 32.** Fix  $p \in \{1, 2\}$ . For  $f : [k]^{pr} \to \mathbb{R}^{\ell}$ ,  $i \in [r]$  and  $d \in [r]$ , we have

$$\operatorname{Inf}_{i}^{\leq d}(\overline{f}^{(p)}) \leq \sum_{j=p(i-1)+1}^{pi} \operatorname{Inf}_{j}^{\leq d}(f).$$

This setup will allow us to painlessly unify the proofs of [KKMO07] and [GS13]. We note that Corollary 32 is a special case (i.e. for  $p \in \{1, 2\}$ ) of [GS20, Lemma 6].

## 6.2 Markov operators

We recount some definitions for Markov operators, following [DMR06, GS13], as well as an important result from [GS13].

A Markov operator T on [D] is a  $D \times D$  stochastic matrix; we say that the operator is symmetric if T is symmetric. For such an operator, we say that the spectral radius r(T) is the second-largest absolute value of any eigenvalue (such an operator has an eigenvalue equal to 1, and all its eigenvalues must be at most 1 in absolute value). We will usually let  $T(x \leftrightarrow y)$  denote the probability that x goes to y i.e. the element at position (x, y) in T.

Such an operator operates on the space of functions  $f : [D] \to \mathbb{R}^d$ , in the following way. For  $x \in [D]$ , we let T(x) be the distribution associated with row (or equivalently column) x in T. Then we have

$$(Tf)(x) = \mathop{\mathbb{E}}_{y \sim T(x)} [f(y)].$$

Note that *T* acts linearly on *f*, since T(af + bg) = a(Tf) + b(Tg) by linearity of expectation. Observe that if  $f(x) = (f^1(x), \dots, f^d(x))$ , then

$$(Tf)(x) = ((Tf^{1})(x), \dots, (Tf^{d})(x)).$$

For any two operators T, T' on [D], [D'] respectively, we define the operator  $T \otimes T'$  on  $[D] \times [D']$ as being the matrix which, at position ((x, x'), (y, y')) for  $x, y \in [D], x', y' \in [D']$  has value  $T(x \leftrightarrow y)T(x' \leftrightarrow y')$ . (In other words,  $T \otimes T'$  is the Kronecker product of T and T'.)

Furthermore, define  $T^{\otimes r} = T \otimes \ldots \otimes T$  where *T* is multiplied *r* times. Observe that  $T^{\otimes r}$  acts on a function  $f : [D]^r \to \mathbb{R}^d$  in the following way. Let  $T^{\otimes r}(\mathbf{x})$  be the product distribution over  $[D]^r$  which gives **y** probability  $\prod_{i=1}^r T(x_i \leftrightarrow y_i)$ . Then

$$(Tf)(\mathbf{x}) = \underset{\mathbf{y} \sim T^{\otimes r}(\mathbf{x})}{\mathbb{E}}[f(\mathbf{y})].$$

Furthermore, we see immediately that  $(T \otimes T')(f \otimes f') = (Tf) \otimes (Tg)$ . Thus, in particular,

$$T^{\otimes r}\alpha_{\mathbf{x}} = \bigotimes_{i=1}^r T\alpha_{x_i}.$$

We observe that for a symmetric Markov operator *T*, we have  $\langle Tf, g \rangle = \langle f, Tg \rangle$ .

The following unifies concepts from [KKMO07, GS13]. The idea also appears without a name in [GS20].

**Definition 33.** Consider a symmetric Markov chain T on  $[k^p] \cong [k]^p$ . We say that the Markov chain is colourful if, for any  $x_1, \ldots, x_p, y_1, \ldots, y_p \in [k]$  such that  $T((x_1, \ldots, x_p) \leftrightarrow (y_1, \ldots, y_p)) > 0$ , we have  $\{x_1, \ldots, x_p\} \cap \{y_1, \ldots, y_p\} = \emptyset$ .

We now introduce two very important colourful operators. The first is (a special case of) the *Bonami-Beckner operator*.

**Definition 34.** Fix  $D \ge 2$ . For each  $-1/(D-1) \le \rho \le 1$  we define the Bonami-Beckner operator by

$$T_{\rho} = \rho \mathbf{I}_D + \frac{1-\rho}{D} \mathbf{J}_D.$$

The matrix  $\mathbf{J}_D$  is the all-ones matrix of size  $D \times D$ . This operator is clearly symmetric and doubly stochastic i.e. it is a symmetric Markov operator. Its eigenvalues are 1 and (k - 1) copies of  $\rho$ . Furthermore, any vector  $\mathbf{x}$  whose sum is zero (i.e. is perpendicular to the all-ones vector) is an eigenvector of  $T_\rho$  with eigenvalue  $\rho$ , as

$$T_{\rho}\mathbf{x} = \rho\mathbf{x} + \frac{1-\rho}{D}\mathbf{J}_{D}\mathbf{x} = \rho\mathbf{x}.$$

**Theorem 28.** Fix  $k \ge 2$ . The Bonami-Beckner operator  $T_{-1/(k-1)}$  on [k] is a symmetric Markov operator that is colourful with spectral radius 1/(k-1). Furthermore, all the values in the matrix are nonnegative integer multiples of 1/(k-1).

*Proof.* The operator is given by the symmetric matrix whose diagonal elements are zero (this is sufficient for colourfulness), and whose off-diagonal elements are 1/(k-1). The eigenvalues of this operator are 1 and (k-1) copies of -1/(k-1), hence the spectral radius is |-1/(k-1)| = 1/(k-1).

**Theorem 29** ([GS13, Lemma 3.8]). Fix  $k \ge 6$ . There exists a colourful symmetric Markov operator on  $[k^2]$  whose spectral radius is at most 4/(k-1). Furthermore, all the values in the matrix are nonnegative integer multiples of 1/(k-1)(k-2)(k-3).

(We note that [GS13, Lemma 3.8] does not explicitly state the fact that the operator is colourful, or that its elements are multiples of 1/(k-1)(k-2)(k-3), but these are easy to observe.)

Given these definitions, we also define the notion of noise stability.

**Definition 35.** Let  $f : [D]^r \to \mathbb{R}$ . Then we define, for  $-1/(D-1) \le \rho \le 1$ , the noise stability of f as

$$\mathbb{S}_{\rho}(f) = \langle f, T_{\rho}^{\otimes r} f \rangle$$

Equivalently,

$$\mathbb{S}_{\rho}(f) = \sum_{\mathbf{x} \in [D]^r} \rho^{|\mathbf{x}|} \hat{f}^2(\mathbf{x}),$$

regardless of the choice of  $\alpha_1, \ldots, \alpha_D$  provided  $\alpha_1(x) = 1$ .

#### MOO theorem, bounds 6.3

We introduce some simple definitions from [KKMO07]. (In fact, the exact definition of the quantity  $\Lambda_{\rho}(\mu)$ is not needed, only the bound in Theorem 39.)

**Definition 36.** Fix 
$$-1 \le \rho \le 1$$
 and  $0 \le \mu \le 1$ . Let  $u$  be some value such that if  $x \sim \mathcal{N}(0, 1)$ , then  
 $\Pr[x \le u] = \mu$ . Let  $\Sigma = \begin{pmatrix} 1 & \rho \\ \rho & 1 \end{pmatrix}$ , and suppose  $(x, y) \sim \mathcal{N}(\mathbf{0}, \Sigma)$ . Then, we define  
 $\Lambda_{\rho}(\mu) = \Pr[x \le u, y \le u].$ 

We will need the following theorem to prove the main technical lemma. It follows from what is called the MOO Theorem (i.e. Mossell, O'Donnell, Oleszkiewicz) in [KKMO07], proved originally by Mossel, O'Donnel and Oleszkiewicz [MOO10], together with [KKMO07, Proposition 13].

**Theorem 37.** For any  $k \ge 2, 0 \le \rho < 1$  and  $\varepsilon > 0$ , there exists  $\tau > 0, d \in \mathbb{N}$  such that the following holds. Suppose  $f: [k]^r \to [0,1]$  is such that  $\operatorname{Inf}_i^{\leq d}(f) \leq \tau$  for all  $i \in [n]$ , and let  $\mu = \mathbb{E}[f]$ . Then

$$\mathbb{S}_{\rho}(f) \leq \Lambda_{\rho}(\mu) + \varepsilon.$$

In order to link this theorem with our setting, we will need the following relation between  $\langle f, T^{\otimes r} f \rangle$  for some symmetric Markov operator T with spectral radius  $\rho$  and  $\mathbb{S}_{\rho}(f)$ . The following bound generalises the first step in proving [KKMO07, Proposition 12], following the first step in the proof of [GS13, Propositio 3.21].

**Lemma 38.** Let T be a symmetric Markov operator on [D] with spectral radius  $0 < \rho < 1$ . For any f:  $[D]^r \to [0,1]$ , where  $\mu = \mathbb{E}[f]$ , we have that

$$\langle f, T^{\otimes r} f \rangle \ge \mu^2 - \mathbb{S}_{\rho}(f).$$

*Proof.* Suppose that  $\alpha_1, \ldots, \alpha_D : [D] \to \mathbb{R}$  are orthonormal (with respect to  $\langle \cdot, \cdot \rangle$ ) eigenvectors of T, seen as functions, whose eigenvalues are  $\lambda_1, \ldots, \lambda_D$ . Suppose  $\alpha_1 = 1$  is the constant one function, with eigenvalue 1. Then we find that  $\alpha_2, \ldots, \alpha_D$  are all perpendicular to the constant-ones function, and have eigenvalue at most  $\rho$  in absolute value. Furthermore, we find that  $\alpha_1, \ldots, \alpha_D$  are also eigenvectors of  $T_{\rho}$ , with eigenvalues 1,  $\rho$ , ...,  $\rho$ . Recall that  $T^{\otimes r} \alpha_{\mathbf{x}} = \bigotimes_{i=1}^{r} T \alpha_{x_i}$ ; now  $T \alpha_{x_i} = \lambda_{x_i} \alpha_{x_i}$ , so

$$T^{\otimes r} \alpha_{\mathbf{x}} = \bigotimes_{i=1}^r \lambda_{x_i} \alpha_{x_i} = \left(\prod_{i=1}^r \lambda_{x_i}\right) \left(\bigotimes_{i=1}^r \alpha_{x_i}\right) = \left(\prod_{i=1}^r \lambda_{x_i}\right) \alpha_{\mathbf{x}}.$$

So, recalling that T is symmetric and hence  $T^{\otimes r}$  is also thus,

$$\begin{aligned} \langle f, T^{\otimes r} f \rangle &= \sum_{\mathbf{x} \in [D]^r} \hat{f}(\mathbf{x}) \widehat{T^{\otimes r} f}(\mathbf{x}) = \sum_{\mathbf{x} \in [D]^r} \hat{f}(\mathbf{x}) \langle T^{\otimes r} f, \alpha_{\mathbf{x}} \rangle \\ &= \sum_{\mathbf{x} \in [D]^r} \hat{f}(\mathbf{x}) \langle f, T^{\otimes r} \alpha_{\mathbf{x}} \rangle = \sum_{\mathbf{x} \in [D]^r} \hat{f}(\mathbf{x}) \left\langle f, \left(\prod_{i=1}^r \lambda_{x_i}\right) \alpha_{\mathbf{x}} \right) = \sum_{\mathbf{x} \in [D]^r} \left(\prod_{i=1}^r \lambda_{x_i}\right) \hat{f}(\mathbf{x}) \left\langle f, \alpha_{\mathbf{x}} \right\rangle \\ &= \sum_{\mathbf{x} \in [D]^r} \left(\prod_{i=1}^r \lambda_{x_i}\right) \hat{f}(\mathbf{x}) \left\langle f, \alpha_{\mathbf{x}} \right\rangle \end{aligned}$$

Observe now that  $\prod_{i=1}^{r} \lambda_1 = 1$  and that  $\prod_{i=1}^{r} \lambda_{x_i} \ge -\rho^{|\mathbf{x}|}$ . Noting that  $\hat{f}(\mathbf{1}) = \langle \mathbf{1}, f \rangle = \mathbb{E}_{\mathbf{x}} f(\mathbf{x}) = \mu$ , we have that

$$\langle f, T^{\otimes r} f \rangle \ge 2\mu^2 - \sum_{\mathbf{x} \in [D]^r} \rho^{|\mathbf{x}|} \hat{f}^2(\mathbf{x}) = 2\mu^2 - \mathbb{S}_{\rho}(f).$$

We will also need an estimate for  $\Lambda_{\rho}(\mu)$ . The following appears in [KKMO07].

**Theorem 39** ([KKMO07, Proposition 11]). For all small enough  $\mu$  and  $0 < \rho \le 1/\ln^3(1/\mu)$ , we have

$$\Lambda_{\rho}(\mu) \leq \mu \left( \mu + 2\rho\mu \ln(1/\mu) \left( 1 + O\left( \frac{\ln \ln(1/\mu) + \ln \ln(1/\rho)}{\ln(1/\mu)} \right) \right) \right)$$

We will also need an estimate for when  $\mu$  is not small. A fact similar to the following appears in the proof of [KKM007, Proposition 12]; we offer a derivation from the literature for completeness.

**Proposition 40.** Suppose  $\rho \ge 0$  is small and  $0 < \mu < 1$ . Then

$$\Lambda_{\rho}(\mu) \leq \mu^2 + 3\rho.$$

We use the following bound due to Willink [Wil05]. Their results are expressed using the following functions:

$$L(u, v, \rho) = \Pr[x \ge u, y \ge v]$$
  

$$\Phi(u) = \Pr[x \le u]$$
  

$$\Phi(u, v, \rho) = \Pr[x \le u, y \le u],$$

where  $(x, y) \sim \mathcal{N}(\mathbf{0}, \mathbf{\Sigma}), \mathbf{\Sigma} = \begin{pmatrix} 1 & \rho \\ \rho & 1 \end{pmatrix}$ . Observe that (-x, -y) and (x, y) have the same distribution; furthermore  $\Pr[x \ge u, y \ge v] = \Pr[-x \le -u, -y \le -v]$ . Hence we see that if u is selected so that  $\Phi(u) = \Pr[x \le u] = \mu$ , then  $\Lambda_{\rho}(\mu) = \Pr[x \le u, y \le u] = \Pr[-x \ge -u, -y \ge -u] = L(-u, -u, \rho)$ .

**Theorem 41** ([Wil05, Equation (1.2)]). Define  $\theta = \sqrt{\frac{1-\rho}{1+\rho}}$ . For  $h > 0, \rho \ge 0$ , we have

$$L(b, h, \rho) \le \Phi(-b)\Phi(-\theta h)(1+\rho).$$

Also recall the following intuitive fact.

**Theorem 42** ([Wil05, Equation (1.1)]).  $L(b, k, \rho) = 1 - \Phi(b) - \Phi(k) + \Phi(b, k, \rho)$ .

*Proof of Proposition 40.* First suppose  $\mu < 1/2$ . Take u < 0 such that  $\Phi(u) = \mu$ . Note that as  $\Lambda_{\rho}(\mu) = L(-u, -u, \rho)$ , we have

$$\Lambda_{\rho}(\mu) \leq \Phi(u)\Phi(\theta u)(1+\rho)$$

Let  $\phi(x) = \Phi'(x)$  be the density function of the normal distribution. It is well known that  $\Phi(u) \le -\phi(u)/u$  for u < 0. So, as  $\max_x \phi(x) = 1/\sqrt{2\pi} \le 1$ , we have that  $-1/u \ge -\phi(u)/u \ge \Phi(u) = \mu$ , hence  $-u \le 1/\mu$ . Observe now that

$$\Phi(\theta u) \leq \Phi(u) - (1 - \theta)u \leq \mu + \frac{1 - \theta}{\mu}$$

as  $\Phi$  is Lipschitz with constant  $1/\sqrt{2\pi} \leq 1$ . So,

$$\Lambda_{\rho}(\mu) \leq \Phi(u)\Phi(\theta u)(1+\rho) \leq \mu\left(\mu + \frac{1-\theta}{\mu}\right)(1+\rho).$$

Now note that  $\theta = \sqrt{(1-\rho)/(1+\rho)} \ge 1-\rho$  for  $0 \le \rho \le 1$ , hence

$$\Lambda_{\rho}(\mu) \leq (1+\rho)(\mu^{2}+\rho) = \mu^{2} + \rho(\mu^{2}+1) + \rho^{2} \leq \mu^{2} + 3\rho,$$

for small enough  $\rho$ .

Now we deal with the case  $\mu > 1/2$  i.e. u > 0. By Theorem 42, and since  $\Phi(-u) = 1 - \Phi(u) = 1 - \mu$ , we have that

$$\Lambda_{\rho}(\mu) = L(-u, -u, \rho) = 1 - 2\Phi(-u) + \Phi(-u, -u, \rho) = 1 - 2(1 - \mu) + \Lambda_{\rho}(1 - \mu).$$

Now apply the result we have proved above to  $1 - \mu < 1/2$ , to find that

$$\Lambda_{\rho}(\mu) \leq 1 - 2(1-\mu) + (1-\mu)^2 + 3\rho = \mu^2 + 3\rho$$

The case  $\mu = 1/2$  follows by continuity.

#### 6.4 **Proof of Proposition 26**

In this section, we prove Proposition 26, which we restate here.

**Proposition 26.** Fix  $p \in \{1, 2\}$  and  $2 \le k \le \ell$  such that  $\ell \le M(k)$ . Suppose T is a symmetric Markov operator on  $[k^p]$  with spectral radius 0 < c/(k-1) < 1, where  $c \le 4$ . Then there exists  $\tau > 0$  and  $d \in \mathbb{N}$  such that for any  $f : [k^p]^r \to \Delta_\ell$  with  $\operatorname{Inf}_i^{\le d}(f) \le \tau$  for all  $i \in [r]$ , we have

$$\langle f, T^{\otimes n} f \rangle \geq \frac{1}{\ell} - \frac{2c \ln \ell}{k\ell} - o\left(\frac{\ln \ell}{k\ell}\right).$$

We first prove the following technical lemma.<sup>22</sup>

**Lemma 43.** Let  $F_T(x) = x^2(1 + T \ln x)$ , where  $F_T(0) = 0$ . Fix  $\ell$  and take some T > 0 smaller than an absolute constant, such that  $\ell < e^{1/T}$ . Suppose  $x_1 + \cdots + x_{\ell} = 1$ ,  $x_i \ge 0$ . Then,  $\sum_{i=1}^{\ell} F_T(x) \ge 1/\ell - T \ln \ell/\ell - 4\ell e^{-1/T}$ .

*Proof.* Note that  $F'_T(x) = 2Tx \ln x + Tx + 2x$  and  $F''_T(x) = 2T \ln(x) + 3T + 2$ . The second derivative is negative for  $x < e^{-1/T-3/2}$ , positive for  $x > e^{-1/T-3/2}$  and zero for  $x = e^{-1/T-3/2}$ . The first derivative is zero at  $e^{-1/T-1/2}$ , negative for smaller  $x \ge 0$ , and positive for greater x. Hence the function decreases below  $e^{-1/T-1/2}$ , then increases above it, and is convex whenever  $x > e^{-1/T-3/2}$ . These functions can be seen for T = 10 in Figure 2.

We wish first to prove that the function is Lipschitz continuous on [0, 1]. Observe that the derivative is minimised at  $e^{-1/T-3/2}$ , and is maximised at 1. At  $e^{-1/T-3/2}$  the derivative is

$$2Te^{-1/T-3/2}\left(-\frac{1}{T}-\frac{3}{2}\right)+Te^{-1/T-3/2}+2T,$$

<sup>&</sup>lt;sup>22</sup>A variant of this lemma exists implicitly within [KKMO07], in the proof of [KKMO07, Proposition 12], but this assumes that  $F_T$  is convex on an interval of the form (0, c) where *c* does not depend on *T*. This seems to not be the case, so we give a different proof here. The proof in [GS13] claims their bound follows precisely as in [KKMO07], so it also implicitly makes this claim about  $F_T$ 



Figure 2: Plot of  $F_T(x)$ ,  $F'_T(x)$ ,  $F''_T(x)$  for T = 10.

which for small T is at least -1; furthermore at 1 the derivative is just T + 2, which for small T is at most 3. So we find that  $F_T$  is Lipschitz continuous with constant 3 for all small enough T. Split  $[\ell]$  into two sets A, B: let  $i \in A$  if  $x_i > e^{-1/T - 3/2}$ , and let  $i \in B$  otherwise. Since  $\ell e^{-1/T - 3/2} < 1$ ,

we have that  $|A| \ge 1$ . Now consider two cases.

**Summing over** A Observe that  $\sum_{i \in A} x_i = 1 - \sum_{i \in B} x_i$ . Note that since  $F_T$  is convex above  $e^{-1/T - 3/2}$ , the minimum value of  $\sum_{i \in A} F_T(x_i)$  when  $\sum_{i \in A} x_i$  is fixed is attained when all  $x_i$  for  $i \in A$  are equal. Hence

$$\sum_{i \in A} F_T(x_i) \ge |A| F_T\left(\frac{1 - \sum_{i \in B} x_i}{|A|}\right) \ge |A| F_T(1/|A|) - 3\frac{\sum_{i \in B} x_i}{|A|} \ge |A| F_T(1/|A|) - 3\ell e^{-1/T - 3/2}$$

by Lipschitz continuity, and since  $x_i \leq e^{-1/T-3/2}$  when  $i \in B$ . Observe that this quantity is just

$$\frac{1 - T \ln |A|}{|A|} - 3\ell e^{-1/T - 3/2} \ge \frac{1 - T \ln |A|}{|A|} - 3\ell e^{-1/T}.$$

Now consider the function

$$a \mapsto \frac{1 - T \ln a}{a}$$

The first derivative of this is

$$\frac{T\ln a - T - 1}{a^2}$$

For  $a \le \ell \le e^{1/T}$ , we have that this derivative is negative. Thus the function is minimised when *a* is as large as possible i.e.  $|A| = \ell$ . Hence, we find that.

$$\sum_{i\in\mathcal{A}}F_T(x_i)\geq \frac{1}{\ell}-\frac{T\ln\ell}{\ell}-3\ell e^{-1/T}$$

**Summing over** *B* For  $x_i \le e^{-1/T-3/2}$  we have that that  $F_T(x_i)$  is decreasing, with its minimum at  $e^{-1/T-3/2}$ . Note that this minimum is

$$F_T(e^{-1/T-3/2}) = -\frac{3Te^{-2/T-3}}{2} \ge -e^{-1/T},$$

for small enough T > 0. Hence, summing over B, we find that

$$\sum_{i \in B} F_T(x_i) \ge -\ell e^{-1/T}$$

Thus we conclude that

$$\sum_{i=1}^{\ell} F_T(x_i) \ge \frac{1}{\ell} - \frac{T \ln \ell}{\ell} - 4\ell e^{-1/T}.$$

The proof of Proposition 26 given below follows the proof of [KKMO07, Proposition 12] and of [GS13, Proposition 3.21].

We also use the following notation from [KKMO07]: let  $[x]^+ = \max(x, 0)$ .

*Proof of Proposition 26.* Observe that, as in the proof of Proposition 10, we need only prove this result for large enough k. Thus assume k is large. Fix  $\tau$ , d to be those numbers given by Theorem 37 for k,  $\rho = c/(k-1)$ ,  $\varepsilon = 1/\ell^3$ .

We will actually prove that

$$\langle f, T^{\otimes r} f \rangle \ge \frac{1}{\ell} - \frac{2c \ln \ell}{(k-1)\ell} - C \frac{2c \ln \ell}{\ell} \frac{\ln \ln k}{(k-1)\ln k} - 4\ell e^{-k/3c} - D\ell e^{-\sqrt[3]{k-1}} - \frac{1}{\ell^2}$$
(11)

for some absolute constants C, D > 0. Since  $\ell \leq M(k) = o(e^{\sqrt[3]{k}}), 2c \ln \ell/(k-1)\ell = 2c \ln \ell/k\ell + 2c \ln \ell/k(k-1)\ell = 2c \ln \ell/k\ell + o(\ln \ell/k\ell)$ , and also  $1/\ell^2 \leq 1/k\ell = o(\ln \ell/k\ell)$ , our conclusion that

$$\langle f, T^{\otimes r} f \rangle \ge \frac{1}{\ell} - \frac{2c \ln \ell}{k\ell} - o\left(\frac{\ln \ell}{k\ell}\right).$$

follows immediately.

Define  $f^1, \ldots, f^{\ell} : [k]^r \to \Delta_{\ell}$  by  $f(\mathbf{x}) = (f^1(\mathbf{x}), \ldots, f^{\ell}(\mathbf{x}))$ , and define  $\mu_i = \mathbb{E}_{\mathbf{x}}[f^i(\mathbf{x})]$ . Since  $\sum_i f^i(\mathbf{x}) = 1$ , by linearity we have that  $\sum_i \mu_i = 1$ . By Lemma 38,

$$\langle f^i, T^{\otimes r} f^i \rangle \ge \mu_i^2 - \mathbb{S}_{c/(k-1)}(f^i).$$

Furthermore, since the codomain of  $f^i$  contains only nonnegative numbers we have that  $\langle f^i, T^{\otimes r} f^i \rangle \ge 0$ . Hence

$$\langle f^i, T^{\otimes r} f^i \rangle \ge \left[ 2\mu_i^2 - \mathbb{S}_{c/(k-1)}(f^i) \right]^+,$$

and summing over  $\ell$ , we have

$$\langle f, T^{\otimes r} f \rangle = \sum_{i=1}^{\ell} \langle f^i, T^{\otimes r} f^i \rangle \ge \sum_{i=1}^{\ell} \left[ 2\mu_i^2 - \mathbb{S}_{c/(k-1)}(f^i) \right]^+.$$
(12)

Our goal will now be to prove the inequality

$$\sum_{i=1}^{\ell} \left[ 2\mu_i^2 - \mathbb{S}_{c/(k-1)}(f^i) \right]^+ \ge \left( \sum_{i=1}^{\ell} \mu_i^2 - \frac{2c\mu_i^2 \ln(1/\mu_i)}{k-1} \left( 1 + C\frac{\ln\ln k}{\ln k} \right) \right) - D\ell e^{-\sqrt[3]{k-1}} - \frac{1}{\ell^2}$$
(13)

for some absolute constants C, D > 0. We take the convention that  $\mu_i^2 \ln(1/\mu_i) = 0$  when  $\mu_i = 0$ .

We now split the integers  $[\ell]$  into two sets A, B. Let  $i \in A$  if  $c/(k-1) \leq 1/\ln^3(1/\mu_i)$  i.e.  $e^{-\sqrt[3]{(k-1)/c}} \leq \mu_i$ , and let  $i \in B$  otherwise. We will prove (13) first on A, then on B, then sum. We will fix C when looking at A, then fix D depending on C when looking at B.

Summing over A. We wish to prove that

$$\sum_{i \in \mathcal{A}} \left[ 2\mu_i^2 - \mathbb{S}_{c/(k-1)}(f^i) \right]^+ \ge \left( \sum_{i \in \mathcal{A}} \mu_i^2 - \frac{2c\mu_i^2 \ln(1/\mu_i)}{k-1} \left( 1 + C\frac{\ln\ln k}{\ln k} \right) \right) - \frac{1}{\ell^2}$$

for some value of C that does not depend on c,  $\mu_i$ , k. First, by Theorem 37 with  $\varepsilon = 1/\ell^3$ , we find that

$$\mathbb{S}_{c/(k-1)}(f^i) \leq \Lambda_{c/(k-1)}(\mu_i) + \frac{1}{\ell^3}$$

If any  $\mu_i = 1$ , then the bound we want holds immediately (for large enough k), as then  $\Lambda_{c/(k-1)}(\mu_i) = 1$ , and all other  $\mu_j = 0$ . Thus suppose  $\mu_i < 1$ . Note that by Proposition 40, each term for  $i \in A$  within the sum from the right-hand side of (12) contributes (for large k) at least  $\mu_i^2 - 3c/(k-1) - 1/\ell^3$  to the sum. If there exists some  $\mu_i > (1/k)^{1/10}$ , say, then these values are large enough (for large k) to make (11) hold automatically (since for large enough k we have that  $(1/k)^{1/5}$  is larger than O(1/k)). Thus we can assume that  $\mu_i \leq (1/k)^{1/10}$  for all  $i \in A$  i.e. all  $\mu_i$  are small for  $i \in A$ .

Since  $c/(k-1) \le 1/\ln^3(1/\mu_i)$  and all  $\mu_i$  are small when  $i \in A$ , we apply Theorem 39 to find that

$$\begin{split} \mathbb{S}_{c/(k-1)}(f^{i}) &\leq \Lambda_{c/(k-1)}(\mu_{i}) + \frac{1}{\ell^{3}} \\ &\leq \mu_{i} \left( \mu_{i} + \frac{2c\mu_{i}\ln(1/\mu_{i})}{k-1} \left( 1 + O\left(\frac{\ln\ln(1/\mu_{i}) + \ln\ln(k-1)/c}{\ln(1/\mu_{i})}\right) \right) \right) + \frac{1}{\ell^{3}}. \end{split}$$

We observe that as  $e^{-\sqrt[3]{(k-1)/c}} \le \mu_i \le (1/k)^{1/10}$  for  $i \in A$  there exists some constant C such that for large k this quantity is bounded by

$$\mu_i \left( \mu_i + \frac{2c\mu_i \ln(1/\mu_i)}{k-1} \left( 1 + C\frac{\ln\ln k}{\ln k} \right) \right) + \frac{1}{\ell^3}.$$

Hence by rearranging the sum, we get that

$$\begin{split} \sum_{i \in \mathcal{A}} \left[ 2\mu_i^2 - \mathbb{S}_{c/(k-1)}(f^i) \right]^+ &\geq \sum_{i \in \mathcal{A}} \mu_i^2 - \frac{2c\mu_i^2 \ln(1/\mu_i)}{k-1} \left( 1 + C\frac{\ln\ln k}{\ln k} \right) - \frac{1}{\ell^3} \\ &\geq \left( \sum_{i \in \mathcal{A}} \mu_i^2 - \frac{2c\mu_i^2 \ln(1/\mu_i)}{k-1} \left( 1 + C\frac{\ln\ln k}{\ln k} \right) \right) - \frac{1}{\ell^2}. \end{split}$$

**Summing over** *B***.** We wish to prove that

$$\sum_{i\in B} \left[2\mu_i^2 - \mathbb{S}_{c/(k-1)}(f^i)\right]^+ \ge \underbrace{\left(\sum_{i\in B} \mu_i^2 - \frac{2c\mu_i^2\ln(1/\mu_i)}{k-1}\left(1 + C\frac{\ln\ln k}{\ln k}\right)\right)}_{\mathcal{S}} - D\ell e^{-\sqrt[3]{k-1}}.$$

Note that the left-hand side is nonnegative; furthermore, every term in *S* is, for large *k*, at most some universal constant times  $e^{-2\sqrt[3]{(k-1)/c}}$ . By assumption  $c \le 4$ , so  $e^{-2\sqrt[3]{(k-1)/c}} \le e^{-\sqrt[3]{k-1}}$ . Thus setting *D* large enough makes the inequality true.

Now, by adding the bound when summing over A and B, we get that (13) is true. Combined with (13) with (12), what we must now show to prove (11) is:

$$\sum_{i=1}^{\ell} \mu_i^2 - \frac{2c\mu_i^2 \ln(1/\mu_i)}{k-1} \left( 1 + C\frac{\ln\ln k}{\ln k} \right) \ge \frac{1}{\ell} - \frac{2c\ln\ell}{(k-1)\ell} - C\frac{2c\ln\ell}{\ell} \frac{\ln\ln k}{(k-1)\ln k} - 4\ell e^{-k/3\epsilon}.$$
 (14)

Recall the function  $F_T(x)$  from Lemma 43. We observe that (14) is equivalent to

$$\sum_{i=1}^{\ell} F_T(\mu_i) \ge \frac{1}{\ell} - \frac{2c \ln \ell}{(k-1)\ell} - C \frac{2c \ln \ell}{\ell} \frac{\ln \ln k}{(k-1) \ln k}$$

where

$$T = \frac{2c}{k-1} \left( 1 + C \frac{\ln \ln k}{\ln k} \right) \ge 0.$$

For large k, we have that  $T \leq 3c/k$ . Hence for large k we have T arbitrarily small; furthermore, by assumption  $\ell < M(k) = o(e^{\sqrt[3]{k}}) = o(e^{k/3c}) = o(e^{1/T})$ . Thus, for large k, we have that  $\ell < e^{1/T}$ . Thus, applying Lemma 43, we have simply that

$$\sum_{i=1}^{\ell} F_T(\mu_i) \ge \frac{1}{\ell} - \frac{\ln \ell}{\ell} \left( \frac{2c}{k-1} \left( 1 + C \frac{\ln \ln k}{\ln k} \right) \right) - 4\ell e^{-k/3c},$$

which by rearranging yields (14).

#### 6.5 **Proof of Proposition 27**

We will prove the following hardness fact.

**Proposition 27.** Let  $p \in \{1, 2\}$  and  $2 \le k \le \ell$ . Assume that there is a colourful symmetric Markov operator T on  $[k^p]$  and  $\tau > 0$ ,  $d \in \mathbb{N}$  such that for any  $f : [k^p]^r \to \Delta_\ell$  with  $\operatorname{Inf}_i^{\le d}(f) \le \tau$  for all  $i \in [n]$ , we have that  $\langle f, T^{\otimes r} f \rangle \ge 1 - \beta$  for some  $\beta \in (0, 1)$ . Assume further that all the values of T are nonnegative integer multiples of a rational number L. Then, assuming that  $(1 - \eta, \eta)$ -approximating a label cover instance with p-to-1 constraints is NP-hard, for any small enough  $\varepsilon$ , it is NP-hard to decide whether a given graph G has a k-colouring of value  $1 - \varepsilon$ , or not even an  $\ell$ -colouring of value  $\beta + \varepsilon$ .

The construction here is very standard, and essentially identical to that in [KKMO07, Section 11.4] or [GS13, Section 3.4]. We will express our results in a more algebraic way, though, rather than using the language of PCP verifiers.

*Proof of Proposition 27.* Fix  $p \in \{1, 2\}$ . Henceforth let  $\overline{\cdot}, \underline{\cdot}$  be  $\overline{\cdot}^{(p)}$  and  $\underline{\cdot}_{(p)}$  respectively. We will consider some value  $\eta$  that depends on  $p, \varepsilon$ ; at the end of the proof we will fix  $\eta$  small enough for everything to follow. By assumption, there exists an  $r \in \mathbb{N}$  such that, given a left-regular label cover instance  $I = (V = V_A \cup V_B, E, \pi_*)$ , with *p*-to-1 constraints, it is NP-hard to decide whether there exists a solution with value at least  $1 - \eta$ , or all solutions have value at most  $\eta$ .

We observe that a constraint is a *p*-to-1 function from [pr] to [r]; such a function can be written as a composite between a permutation  $\pi : [pr] \rightarrow [pr]$  and the function  $\sigma : [pr] \rightarrow [r]$ , given by  $\sigma(1) = \ldots = \sigma(p) = 1, \sigma(p+1) = \ldots = \sigma(2p) = 2$  i.e.  $\sigma(x) = \lceil x/p \rceil$ . Thus we assume that the constraint that corresponds to the edge (a, b) is given by  $\sigma \circ \pi_{a,b}$ .

We will reduce this instance in polynomial time to an instance of maximum k- vs.  $\ell$ -colouring, namely a graph G = (V', E'), and then prove the completeness and soundness of the reduction.

**Reduction** For every variable  $b \in V_B$ , introduce a set of variables in our graph in the following way. For every vector  $\mathbf{x} \in [k]^{pr}$ , we introduce a vertex  $v_b(\mathbf{x})$ . Thus our graph G will have the vertex set

$$V' = \{v_b(\mathbf{x}) \mid b \in V_B, \mathbf{x} \in [k]^{pr}\},\$$

with  $k^{pr}|V_B|$  vertices. As for the edges, consider every pair of edges  $(a, b), (a, b') \in E$ . For such a pair, for every  $\mathbf{x}, \mathbf{y} \in [k]^{pr}$ , we add in an edge between  $(\mathbf{x}^{\pi_{a,b}})$  and  $v_{b'}(\mathbf{x}^{\pi_{a,b'}})$  exactly  $L^r T^{\otimes r}(\overline{\mathbf{x}} \leftrightarrow \overline{\mathbf{y}}) \leq L^r$  times. This is well defined since  $\overline{\mathbf{x}}, \overline{\mathbf{y}} \in [k^p]^r$ , and  $T^{\otimes r}$  can be seen as a Markov chain over  $[k^p]^r$ . Furthermore each probability in T is a nonnegative integer multiple of L, hence we never add non-integer or negative numbers of edges between vertices.

Throughout the following, we will let  $f_a(\mathbf{x})$  denote the colour of  $v_a(\mathbf{x})$ . Observe that this reduction works in polynomial time, since *L*, *k*, *p*, *r* are constants.

**Completeness** Suppose there exists a solution to *I*, say  $c : V \to [r]$ , that satisfies a  $1 - \eta$  fraction of constraints. Consider the following *k*-colouring of *G*: let  $f_v(x_1, \ldots, x_r) = x_{c(v)}$ . Now, we must compute the value of this colouring. Suppose val(*a*) is the proportion of edges incident to  $a \in V_A$  solved by *c*. Since the instance *I* is regular on  $V_A$ , we have that the value of the instance is  $\mathbb{E}[val(a)] \ge 1 - \eta$ , where *a* is drawn uniformly at random from  $V_A$ .

What fraction of the edges in E' were constructed due to a pair of edges  $(a, b), (a, b') \in E$  which are both satisfied by c? Since the degree of all vertices in  $V_A$  is equal, and thus the same number of edges is added for each a, this is equivalent to asking "what is the probability that, if we select  $a \in V_A$  and  $b, b' \in V_B$ incident to a uniformly and independently at random, then the edges (a, b), (a, b') are solved by c". But observe that this probability, for fixed a, is at least  $1-2(1-\operatorname{val}(a)) = 2\operatorname{val}(a)-1$ . By linearity of expectation, the required probability is thus at least  $\mathbb{E}[2\operatorname{val}(a) - 1] \ge 2(1 - \eta) - 1 = 1 - 2\eta$ .

Now, note that every edge  $(v_b(\mathbf{x}^{\pi_{a,b}}), v_{b'}(\mathbf{y}^{\pi_{a,b'}})) \in E'$  where (a, b), (a, b') are solved by c will also be properly coloured by c'. To see why, note that  $f_b(\mathbf{x}^{\pi_{a,b}}) = f_b(x_{\pi_{a,b}(1)}, \dots, x_{\pi_{a,b}(pr)}) = x_{\pi_{a,b}(c(b))}$ , and likewise  $f_{b'}(\mathbf{y}^{\pi_{a,b'}}) = y_{\pi_{a,b'}(c(b'))}$ . Defining  $i = \pi_{a,b}(c(b)), j = \pi_{a,b'}(c(b'))$ , these are  $x_i$  and  $y_j$ . But since the edges (a, b), (a, b') are solved, we have  $\sigma(i) = c(a) = \sigma(j)$ , or equivalently

$$i, j \in \{pc(a), pc(a) + 1, \dots, pc(a) + p - 1\}.$$

Since we have added in the edge  $(v_b(\mathbf{x}^{\pi_{a,b}}), v_{b'}(\mathbf{y}^{\pi_{a,b'}}))$ , it follows that the transition between  $\overline{\mathbf{x}}$  and  $\overline{\mathbf{y}}$  in  $T^{\otimes r}$  has nonzero probability. Thus, the transition between  $(x_{pc(a)}, \ldots, x_{pc(a)+p-1})$  and  $(y_{pc(a)}, \ldots, y_{pc(a)+p-1})$  has nonzero probability in T; by the colourfulness of T,

$$\{x_{pc(a)}, \ldots, x_{pc(a)+p-1}\} \cap \{y_{pc(a)}, \ldots, y_{pc(a)+p-1}\} = \emptyset.$$

Thus  $x_i \neq y_i$ . Hence the edge is properly coloured, as

$$f_b(\mathbf{x}^{\pi_{a,b}}) = x_{\pi_{a,b}(c(b))} = x_i \neq y_j = y_{\pi_{a,b'}(c(b'))} = f_{b'}(\mathbf{y}^{\pi_{a,b'}}).$$

Thus the resulting graph has a k-colouring of value  $1 - 2\eta$ ; taking  $2\eta < \varepsilon$  thus implies completeness.

**Soundness** Suppose that the graph G has an  $\ell$ -colouring of value at least  $\beta + \varepsilon$ ; call it c. As opposed to the completeness case, let val(a) denote the proportion of edges in G added due to edges  $(a, b), (a, b') \in E$  that are properly coloured. Having fixed such an a, note that every choice of  $\mathbf{x} \in [k^p]^r$  induces the same number of edges, and for this  $\mathbf{x}$  every choice of  $\mathbf{y}$  induces a number of edges proportional to  $T^{\otimes}(\mathbf{x} \leftrightarrow \mathbf{y})$ . Hence

$$\operatorname{val}(a) = \underset{\overline{\mathbf{x}} \in [k^{p}]^{r}}{\mathbb{E}} [f_{a}(\mathbf{x}^{\pi_{a,b}}) \neq f_{b}(\mathbf{y}^{\pi_{a,b}})].$$

Observe that since all vertices in  $V_A$  have the same degree, similarly the same number of edges of the form  $(a, b), (a, b') \in E$  exist for every  $a \in V_A$ ; hence the value of c can be expressed as  $\mathbb{E}[val(a)] \ge \beta + \varepsilon$ , where a is drawn uniformly at random from  $V_A$ . Now apply Markov's inequality to 1 - val(a) to find that the probability that  $1 - val(a) \ge 1 - \beta$ , i.e.,  $val(a) \le \beta$  is at most  $(1 - \beta - \varepsilon)/(1 - \beta) = 1 - \varepsilon/(1 - \beta) \le 1 - \varepsilon$ . Hence the probability that  $val(a) > \beta$  is at least  $\varepsilon$ . Let  $S \subseteq V_A$  be the set of  $a \in V_A$  for which  $val(a) \ge \beta$ ; we have that  $|S| \ge \varepsilon |V_A|$ .

Our goal will be to assign, to each vertex in  $S \cup V_B$ , at most  $C = \max(\lceil 2pd/\tau \rceil, 1)$  possible values; if  $c(a) \subseteq [r]$  for  $a \in S$  and  $c(b) \subseteq [pr]$  for  $b \in V_B$  is the set of values, we will want

$$c(a) \cap \sigma(\pi_{a,b}(c(b))) \neq \emptyset \tag{15}$$

for at least a  $\tau/2p$  fraction of the edges  $(a, b) \in E$  for any fixed  $a \in S$ . Then, by randomly selecting a value for v from c(v), we find that each of these  $\tau/2p$  fraction of edges of the form  $(a, b) \in E$  for any fixed  $a \in S$  are satisfied with probability at least  $1/C^2$ . Since I is regular on  $V_A$  and  $|S| \ge \varepsilon |V_A|$ , this implies that this solution (if extended arbitrarily to  $V_A \setminus S$ ) has value at least  $\varepsilon \tau/2pC^2$ . Taking  $\eta$  small enough so that  $\varepsilon \tau/2pC^2 \ge \eta$  is then enough to prove soundness.

Note that  $[f_b(\mathbf{x}) \neq f_{b'}(\mathbf{y})] = 1 - f_b(\mathbf{x}) \cdot f_{b'}(\mathbf{y})$ , if we see the codomain  $[\ell]$  of  $f_b$  as being embedded within  $\Delta_{\ell}$ ; fixing some  $a \in S$ , we then observe that

$$\beta < \operatorname{val}(a) = \underbrace{\mathbb{E}}_{\substack{(a,b),(a,b') \in E \\ \mathbf{x} \in [k^p]^r \\ \overline{\mathbf{y}} \sim T^{\otimes r}(\overline{\mathbf{x}})}} \begin{bmatrix} 1 - f_b(\mathbf{x}^{\pi_{a,b}}) \cdot f_{b'}(\mathbf{y}^{\pi_{a,b'}}) \end{bmatrix} = 1 - \underbrace{\mathbb{E}}_{\substack{\overline{\mathbf{x}} \in [k^p]^r \\ \overline{\mathbf{y}} \sim T^{\otimes r}(\overline{\mathbf{x}})}} \begin{bmatrix} \mathbb{E}\\ (a,b),(a,b') \in E \end{bmatrix} \begin{bmatrix} f_b(\mathbf{x}^{\pi_{a,b}}) \cdot f_{b'}(\mathbf{y}^{\pi_{a,b'}}) \end{bmatrix} \end{bmatrix}$$

Observe that the inner product above is between two independent variables, hence the expression is equal to

$$1 - \mathop{\mathbb{E}}_{\substack{\overline{\mathbf{x}} \in [k^p]^r \\ \overline{\mathbf{y}} \sim T^{\otimes r}(\overline{\mathbf{x}})}} \left[ \mathop{\mathbb{E}}_{(a,b) \in E} \left[ f_b(\mathbf{x}^{\pi_{a,b}}) \right] \cdot \mathop{\mathbb{E}}_{(a,b) \in E} \left[ f_b(\mathbf{y}^{\pi_{a,b}}) \right] \right].$$

Now, define  $g_a(\mathbf{x}) = \mathbb{E}_{(a,b)\in E} \left[ f_b(\mathbf{x}^{\pi_{a,b}}) \right]$  i.e.  $g_a = \mathbb{E}_{(a,b)\in E} \left[ f_b^{\pi_{a,b}} \right]$ . Hence, by substituting  $\mathbf{x}, \mathbf{y}$  with  $\underline{\mathbf{x}}, \underline{\mathbf{y}}$ , we get that the expression is

$$1 - \mathop{\mathbb{E}}_{\substack{\overline{\mathbf{x}} \in [k^{p}]^{r} \\ \overline{\mathbf{y}} \sim T^{\otimes r}(\overline{\mathbf{x}})}} \left[ g_{a}(\mathbf{x}) \cdot g_{a}(\mathbf{y}) \right] = 1 - \mathop{\mathbb{E}}_{\substack{\mathbf{x} \in [k^{p}]^{r} \\ \mathbf{y} \sim T^{\otimes r}(\mathbf{x})}} \left[ \overline{g_{a}}(\mathbf{x}) \cdot \overline{g_{a}}(\mathbf{y}) \right].$$

Now, separating the choice of  $\mathbf{x}$  from  $\mathbf{y}$ , and by linearity, this is

$$1 - \mathbb{E}_{\mathbf{x} \in [k^p]^r} \left[ \overline{g_a}(\mathbf{x}) \cdot \mathbb{E}_{\mathbf{y} \sim T^{\otimes r}(\mathbf{x})} \overline{g_a}(\mathbf{y}) \right] = 1 - \mathbb{E}_{\mathbf{x} \in [k^p]^r} \left[ \overline{g_a}(\mathbf{x}) \cdot \left( (T^{\otimes r} \overline{g_a})(\mathbf{x}) \right) \right] = 1 - \langle \overline{g_a}, T^{\otimes r} \overline{g_a} \rangle$$

Hence,  $\langle \overline{g_a}, T^{\otimes r} \overline{g_a} \rangle < 1 - \beta$  for  $a \in S$ . By assumption we must have  $\operatorname{Inf}_i^{\leq d}(\overline{g_a}) > \tau$  for at least one  $i \in [pr]$ . By Corollary 32, for at least one  $i \in [pr]$  we must have  $\operatorname{Inf}_i^{\leq d}(g_a) > \tau/p$ . Let  $c(a) = \{i\}$ ; thus  $|c(a)| \leq C$ . Thus we have labelled S. For  $b \in V_B$ , define

$$c(b) = \{i \in [r] \mid \operatorname{Inf}_{i}^{\leq d}(f_{b}) \geq \tau/2p\}.$$

Note that  $\sum_{i=1}^{r} \text{Inf}_{i}^{\leq d}(f_{b}) \leq d$ , and furthermore  $\text{Inf}_{i}^{\leq d}(f_{b}) \geq 0$  as  $f_{b}$  takes nonnegative values, so  $|c(b)| \leq 2pd/\tau \leq C$ .

Now, we must prove that this mapping *c* satisfies (15) for any fixed  $a \in S$ , and for a  $\tau/2p$  fraction of edges of the form  $(a, b) \in E$ . Fixing  $a \in S$ , letting  $g = g_a$ ,  $\{i\} = c(a)$ , we observe first that

$$\hat{g} = \widehat{\mathbb{E}[f_b^{\pi_{a,b}}]} = \mathbb{E}_{(a,b)\in E}[\widehat{f_b^{\pi_{a,b}}}],$$

by linearity of  $\hat{\cdot}$  and hence  $\hat{g}(\mathbf{x}) = \mathbb{E}_{(a,b)\in E}[\widehat{f_b^{\pi_{a,b}}}(\mathbf{x})]$ . Thus

$$\begin{aligned} \tau/p < \mathrm{Inf}_{i}^{\leq d}(g) &= \sum_{\substack{\mathbf{x} \in [k]^{r} \\ x_{i} \neq 1 \\ |\mathbf{x}| \leq d}} \left| \hat{g}(\mathbf{x}) \right|^{2} = \sum_{\substack{\mathbf{x} \in [k]^{r} \\ x_{i} \neq 1 \\ |\mathbf{x}| \leq d}} \left| \left| \mathbb{E}_{(a,b) \in E} \left[ \widehat{f_{b}^{\pi_{a,b}}}(\mathbf{x}) \right] \right|^{2} \leq \sum_{\substack{\mathbf{x} \in [k]^{r} \\ x_{i} \neq 1 \\ |\mathbf{x}| \leq d}} \mathbb{E} \left[ \left| \widehat{f_{b}^{\pi_{a,b}}}(\mathbf{x}) \right|^{2} \right] \\ &= \mathbb{E}_{(a,b) \in E} \left[ \sum_{\substack{\mathbf{x} \in [k]^{r} \\ x_{i} \neq 1 \\ |\mathbf{x}| \leq d}} \left| \widehat{f_{b}^{\pi_{a,b}}}(\mathbf{x}) \right|^{2} \right] = \mathbb{E}_{(a,b) \in E} \left[ \mathrm{Inf}_{i}^{\leq d}(f_{b}^{\pi_{a,b}}) \right], \end{aligned}$$

where all the inequalities follow by linearity or convexity. We now apply Markov's inequality to  $\max(\inf_{i=1}^{\leq d} (f_b^{\pi_{a,b}})) - \inf_{i=1}^{\leq d} (f_b^{\pi_{a,b}})$ . For a  $\tau/2p$  fraction of the edges  $(a, b) \in E$  we have

$$\mathrm{Inf}_i^{\leq d}(f_b^{\pi_{a,b}}) \geq \tau/2p.$$

Now note that for such *b*,

$$\tau/2p \leq \mathrm{Inf}_{i}^{\leq d}(f_{b}^{\pi_{a,b}}) = \sum_{\substack{\mathbf{x}\in[k]^{r}\\x_{i}\neq 1\\|\mathbf{x}|\leq d}} \left|\widehat{f_{b}^{\pi_{a,b}}}(\mathbf{x})\right|^{2} = \sum_{\substack{\mathbf{x}\in[k]^{r}\\x_{i}\neq 1\\|\mathbf{x}|\leq d}} \left|\widehat{f_{b}}(\mathbf{x}^{\pi_{a,b}})\right|^{2} = \sum_{\substack{\mathbf{y}\in[k]^{r}\\y_{\pi_{a,b}^{-1}(i)}\neq 1\\|\mathbf{y}|\leq d}} \left|\widehat{f_{b}}(\mathbf{y})\right|^{2} = \mathrm{Inf}_{\pi_{a,b}^{-1}(i)}(f_{b}).$$

where  $\mathbf{y} = \mathbf{x}^{\pi_{a,b}}$ . Hence  $\pi_{a,b}^{-1}(i) \in c(b)$ , so in particular (15) holds for this *a* and for at least a  $\tau/2p$  fraction of the edges (a, b). This concludes the proof.

## References

- [AG18] Dario Azzimonti and David Ginsbourger. Estimating orthant probabilities of high-dimensional gaussian vectors with an application to set estimation. J. Comput. Graph. Stat., 27(2):255–267, 2018. doi:10.1080/10618600.2017.1360781.
- [AGH17] Per Austrin, Venkatesan Guruswami, and Johan Håstad. (2+ε)-Sat is NP-hard. SIAM J. Comput., 46(5):1554–1573, 2017. doi:10.1137/15M1006507.
- [AOTW14] Per Austrin, Ryan O'Donnell, Li-Yang Tan, and John Wright. New np-hardness results for 3-coloring and 2-to-1 label cover. *ACM Trans. Comput. Theory*, 6(1):2:1–2:20, 2014. doi:10.1145/2537800.
- [BBKO21] Libor Barto, Jakub Bulín, Andrei A. Krokhin, and Jakub Opršal. Algebraic approach to promise constraint satisfaction. J. ACM, 68(4):28:1–28:66, 2021. arXiv:1811.00970, doi:10.1145/3457606.
- [BG19] Joshua Brakensiek and Venkatesan Guruswami. An algorithmic blend of LPs and ring equations for promise CSPs. In Proc. 30th Annual ACM-SIAM Symposium on Discrete Algorithms (SODA'19), pages 436–455, 2019. arXiv:1807.05194, doi:10.1137/1.9781611975482.28.

- [BG21] Joshua Brakensiek and Venkatesan Guruswami. Promise Constraint Satisfaction: Algebraic Structure and a Symmetric Boolean Dichotomy. *SIAMJ. Comput.*, 50(6):1663–1700, 2021. arXiv:1704.01937, doi:10.1137/19M128212X.
- [BGS23] Joshua Brakensiek, Venkatesan Guruswami, and Sai Sandeep. SDPs and Robust Satisfiability of Promise CSP. In Proc. Annual 55th ACM Symposium on Theory of Computing (STOC'23), pages 609–622. ACM, 2023. arXiv:2211.08373, doi:10.1145/3564246.3585180.
- [BK05] Ankur Bhargava and S. Rao Kosaraju. Derandomization of dimensionality reduction and SDP based algorithms. In *Proc. 8th International Workshop on Algorithms and Data Structures (WADS'05)*, pages 396–408. Springer Berlin Heidelberg, 2005. doi:10.1007/11534273\\_35.
- [BK16] Libor Barto and Marcin Kozik. Robustly solvable constraint satisfaction problems. *SIAM J. Comput.*, 45(4):1646–1669, 2016. doi:10.1137/130915479.
- [BKLM21] Mark Braverman, Subhash Khot, Noam Lifshitz, and Dor Minzer. An Invariance Principle for the Multi-slice, with Applications. In Proc. 62nd IEEE Annual Symposium on Foundations of Computer Science (FOCS'21), pages 228–236. IEEE, 2021. arXiv:2110.10725, doi:10.1109/F0CS52979.2021.00030.
- [BKM21] Mark Braverman, Subhash Khot, and Dor Minzer. On rich 2-to-1 games. In Proc. 12th Innovations in Theoretical Computer Science Conference (ITCS'21), volume 185 of LIPIcs, pages 27:1–27:20. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2021. doi:10.4230/LIPIcs.ITCS.2021.27.
- [BKM22] Amey Bhangale, Subhash Khot, and Don Minzer. On Approximability of Satisfiable k-CSPs: I. In Proc. 54th Annual ACM Symposium on Theory of Computing (STOC'22), pages 976–988. ACM, 2022. doi:10.1145/3519935.3520028.
- [BKM23a] Amey Bhangale, Subhash Khot, and Don Minzer. On Approximability of Satisfiable k-CSPs: II. In Proc. 55th Annual ACM Symposium on Theory of Computing (STOC'23), pages 632–642. ACM, 2023. doi:10.1145/3564246.3585120.
- [BKM23b] Amey Bhangale, Subhash Khot, and Don Minzer. On Approximability of Satisfiable k-CSPs: III. In Proc. 55th Annual ACM Symposium on Theory of Computing (STOC'23), pages 643–655. ACM, 2023. doi:10.1145/3564246.3585121.
- [Che68] M. Cheng. The clipping loss in correlation detectors for arbitrary input signal-to-noise ratios. *IEEE Trans. Inf. Theory*, 14(3):382–389, 1968. doi:10.1109/TIT.1968.1054159.
- [Che69] M. C. Cheng. The orthant probabilities of four gaussian variates. *Ann. Math. Stat.*, 40(1):152–161, 1969. doi:10.1214/aoms/1177697812.
- [CMM09] Moses Charikar, Konstantin Makarychev, and Yury Makarychev. Near-optimal algorithms for maximum constraint satisfaction problems. ACM Trans. Algorithms, 5(3):32:1–32:14, 2009. doi:10.1145/1541885.1541893.
- [DMR06] Irit Dinur, Elchanan Mossel, and Oded Regev. Conditional hardness for approximate coloring. In Proceedings of the Thirty-Eighth Annual ACM Symposium on Theory of Computing, STOC '06, pages 344–353. Association for Computing Machinery, 2006. doi:10.1145/1132516.1132567.
- [DMR09] Irit Dinur, Elchanan Mossel, and Oded Regev. Conditional Hardness for Approximate Coloring. *SIAM J. Comput.*, 39(3):843–873, 2009. doi:10.1137/07068062X.
- [DP93a] Charles Delorme and Svatopluk Poljak. Combinatorial properties and the complexity of a max-cut approximation. *Eur. J. Comb.*, 14(4):313–333, 1993. doi:10.1006/eujc.1993.1035.
- [DP93b] Charles Delorme and Svatopluk Poljak. Laplacian eigenvalues and the maximum cut problem. *Math. Program.*, 62:557–574, 1993. doi:10.1007/BF01585184.
- [EIO02] Lars Engebretsen, Piotr Indyk, and Ryan O'Donnell. Derandomized dimensionality reduction with applications. In Proc. 13th Annual ACM-SIAM Symposium on Discrete Algorithms (SODA'02), pages 705-712. ACM/SIAM, 2002. URL: http://dl.acm.org/citation.cfm?id=545381.545476.

- [FJ97] Alan M. Frieze and Mark Jerrum. Improved Approximation Algorithms for MAX k-CUT and MAX BISECTION. *Algorithmica*, 18(1):67–81, 1997. doi:10.1007/BF02523688.
- [FV98] Tomás Feder and Moshe Y. Vardi. The Computational Structure of Monotone Monadic SNP and Constraint Satisfaction: A Study through Datalog and Group Theory. SIAM J. Comput., 28(1):57–104, 1998. doi:10.1137/S0097539794266766.
- [GJ76] M. R. Garey and D. S. Johnson. The complexity of near-optimal graph coloring. *J. ACM*, 23(1):43–49, 1976. doi:10.1145/321921.321926.
- [GK04] Venkatesan Guruswami and Sanjeev Khanna. On the hardness of 4-coloring a 3-colorable graph. *SIAM J. Discret. Math*, 18(1):30–40, 2004. doi:10.1137/S0895480100376794.
- [GM12] Bernd Gärtner and Jiří Matoušek. *Approximation algorithms and semidefinite programming*. Springer Science & Business Media, 2012.
- [GOWZ10] Parikshit Gopalan, Ryan O'Donnell, Yi Wu, and David Zuckerman. Fooling Functions of Halfspaces under Product Distributions. In Proc. 25th Annual IEEE Conference on Computational Complexity (CCC'2010), pages 223–234. IEEE Computer Society, 2010. doi:10.1109/CCC.2010.29.
- [GS13] Venkatesan Guruswami and Ali Kemal Sinop. Improved inapproximability results for maximum kcolorable subgraph. *Theory Comput.*, 9:413–435, 2013. doi:10.4086/T0C.2013.V009A011.
- [GS20] Venkatesan Guruswami and Sai Sandeep. d-To-1 Hardness of Coloring 3-Colorable Graphs with O(1) Colors. In Proc. 47th International Colloquium on Automata, Languages, and Programming (IC-ALP'20), volume 168 of LIPIcs, pages 62:1–62:12. Schloss Dagstuhl – Leibniz-Zentrum für Informatik, 2020. doi:10.4230/LIPIcs.ICALP.2020.62.
- [GW95] Michel X. Goemans and David P. Williamson. Improved approximation algorithms for maximum cut and satisfiability problems using semidefinite programming. J. ACM, 42(6):1115–1145, 1995. doi:10.1145/227683.227684.
- [GZ12] Venkatesan Guruswami and Yuan Zhou. Tight bounds on the approximability of almost-satisfiable horn SAT and exact hitting set. *Theory Comput.*, 8(1):239–267, 2012. doi:10.4086/toc.2012.v008a011.
- [Hås01] Johan Håstad. Some optimal inapproximability results. J. ACM, 48(4):798-859, jul 2001. doi:10.1145/502090.502098.
- [HMS23] Yahli Hecht, Dor Minzer, and Muli Safra. NP-Hardness of Almost Coloring Almost 3-Colorable Graphs. In Proc. Approximation, Randomization, and Combinatorial Optimization. Algorithms and Techniques (APPROX/RANDOM'23), volume 275 of LIPIcs, pages 51:1–51:12. Schloss Dagstuhl -Leibniz-Zentrum für Informatik, 2023. doi:10.4230/LIPIcs.APPROX/RANDOM.2023.51.
- [Kar72] Richard M. Karp. Reducibility among combinatorial problems. In Complexity of Computer Computations: Proceedings of a symposium on the Complexity of Computer Computations, pages 85–103. Springer US, 1972. doi:10.1007/978-1-4684-2001-2\_9.
- [Kho02] Subhash Khot. On the power of unique 2-prover 1-round games. In Proc. 34th Annual ACM Symposium on Theory of Computing (STOC'02), pages 767–775. ACM, 2002. doi:10.1145/509907.510017.
- [KKMO07] Subhash Khot, Guy Kindler, Elchanan Mossel, and Ryan O'Donnell. Optimal Inapproximability Results for MAX-CUT and Other 2-Variable CSPs? SIAM J. Comput., 37(1):319–357, 2007. doi:10.1137/S0097539705447372.
- [KKR17] Vladimir Kolmogorov, Andrei A. Krokhin, and Michal Rolínek. The Complexity of General-Valued CSPs. SIAM J. Comput., 46(3):1087–1110, 2017. arXiv:1502.07327, doi:10.1137/16M1091836.
- [KLS00] Sanjeev Khanna, Nathan Linial, and Shmuel Safra. On the hardness of approximating the chromatic number. *Comb.*, 20(3):393–415, 2000. doi:10.1007/s004930070013.

- [KMS98] David R. Karger, Rajeev Motwani, and Madhu Sudan. Approximate graph coloring by semidefinite programming. J. ACM, 45(2):246–265, 1998. doi:10.1145/274787.274791.
- [KMS23] Subhash Khot, Dor Minzer, and Muli Safra. Pseudorandom sets in Grassmann graph have near-perfect expansion. *Ann. Math.*, 198(1):1 92, 2023. doi:10.4007/annals.2023.198.1.1.
- [KO15] Marcin Kozik and Joanna Ochremiak. Algebraic properties of valued constraint satisfaction problem. In Proc. 42nd International Colloquium on Automata, Languages, and Programming (ICALP'22), pages 846–858. Springer Berlin Heidelberg, 2015. arXiv:1403.0476, doi:10.1007/978-3-662-47672-7\\_69.
- [KOWŽ23] Andrei A. Krokhin, Jakub Opršal, Marcin Wrochna, and Stanislav Živný. Topology and adjunction in promise constraint satisfaction. SIAM J. Computing, 52(1):37–79, 2023. arXiv:2003.11351, doi:10.1137/20M1378223.
- [KR08] Subhash Khot and Oded Regev. Vertex cover might be hard to approximate to within  $2 \varepsilon$ . J. Comput. Syst. Sci., 74(3):335–349, 2008. doi:10.1016/j.jcss.2007.06.019.
- [KSTW00] Sanjeev Khanna, Madhu Sudan, Luca Trevisan, and David P. Williamson. The approximability of constraint satisfaction problems. SIAM J. Comput., 30(6):1863–1920, 2000. doi:10.1137/S0097539799349948.
- [MOO10] Elchanan Mossel, Ryan O'Donnell, and Krzysztof Oleszkiewicz. Noise stability of functions with low influences: invariance and optimality. *Ann. of Math. (2)*, 171(1):295–341, 2010. doi:10.4007/annals.2010.171.295.
- [MR99] Sanjeev Mahajan and H. Ramesh. Derandomizing approximation algorithms based on semidefinite programming. SIAM J. Comput., 28(5):1641–1663, 1999. doi:10.1137/S0097539796309326.
- [Rag08] Prasad Raghavendra. Optimal algorithms and inapproximability results for every CSP? In Proc. 40th Annual ACM Symposium on Theory of Computing (STOC'08), pages 245–254, 2008. doi:10.1145/1374376.1374414.
- [Rai19] Martin Raič. A multivariate Berry–Esseen theorem with explicit constants. *Bernoulli*, 25(4A):2824 2853, 2019. arXiv:1802.06475, doi:10.3150/18-BEJ1072.
- [Siv02] D. Sivakumar. Algorithmic derandomization via complexity theory. In *Proc. 34th Annual ACM Symposium on Theory of Computing (STOC'02)*, page 619–626. ACM, 2002. doi:10.1145/509907.509996.
- [TSSW00] Luca Trevisan, Gregory B. Sorkin, Madhu Sudan, and David P. Williamson. Gadgets, approximation, and linear programming. SIAM J. Comput., 29(6):2074–2097, 2000. doi:10.1137/S0097539797328847.
- [VŽ21] Caterina Viola and Stanislav Živný. The combined basic LP and affine IP relaxation for promise vcsps on infinite domains. *ACM Trans. Algorithms*, 17(3):21:1–21:23, 2021. arXiv:2007.01779, doi:10.1145/3458041.
- [Wil05] R. Willink. Bounds on the bivariate normal distribution function. *Communications in Statistics Theory and Methods*, 33(10):2281–2297, 2005. doi:10.1081/STA-200031505.
- [Zwi98] Uri Zwick. Finding almost-satisfying assignments. In Proc. 30th Annual ACM Symposium on Theory of Computing (STOC'98), pages 551–560. ACM, 1998. doi:10.1145/276698.276869.

## A AGC-hardness of 1-approximation

We show a reduction from AGC to 1-approximation of  $\rho_k(G)$  via  $\rho_\ell(G)$ .

**Proposition 44.** Fix  $3 \le k \le \ell$  and some rational  $\rho \in (0, 1]$ . There is a log-space reduction from the problem of distinguishing  $\rho_k(G) = 1$  vs.  $\rho_\ell(G) < 1$  to the problem of distinguishing  $\rho_k(G) \le \rho$  vs.  $\rho_\ell(G) > \rho$ .

*Proof.* Let  $\rho = p/q$ , p > 0. Suppose we are given a graph G; we are then asked to decide if it is k-colourable or not even  $\ell$ -colourable. Let G have m edges and let 1 denote the graph with one vertex and an edge from that vertex to itself. (This notation is justified, since this graph is a unit with respect to the direct product of graphs.) Let + denote disjoint union of graphs. We also allow multiplication of a graph by a scalar in the obvious way. (For example, 3G = G + G + G.) Then our reduction takes the graph G to the graph pG + (q - p)m1.

Note first that the reduction can be done in logarithmic space. For completeness, note that if G is kcolourable, then pG + (q - p)m1 has a k-colouring of value  $\rho$ , namely the one that colours each of the p disjoint copies of G as in the k-colouring of G. This colouring correctly colours pm of the pm + (q - p)m =qm edges i.e. it has value  $\rho = pm/qm = p/q$ . For soundness, suppose that pG + (q - p)m1 has an  $\ell$ colouring of value  $\rho$ . This colouring must correctly colour a p/q fraction of the edges of pG + (q - p)m1. Since this graph has qm edges, it must correctly colour pm edges. But the only edges that could possibly be correctly coloured are the ones in pG (since the remaining edges in (q - p)m1 are all loops). Furthermore, there are pm edges in pG, thus all the edges in pG must be correctly coloured. But this implies that G has an  $\ell$ -colouring, as required.

## **B** Nonsolvability by finite sandwiches

The problem studied in this paper is an example of a promise Max-CSP, which is a special case of promise valued CSP [VŽ21]. A graph (in particular  $K_2$  and  $K_3$ ) is seen as a {0, 1}-valued structure. We denote by  $\rightarrow$  a fractional homomorphism between two valued structures; a fractional polymorphism from G to H is a fractional homomorphism from  $G^n$  to H for some  $n \in \mathbb{N}$  [VŽ21].

**Proposition 45.** Assume  $P \neq NP$ . Let  $\Gamma$  be a valued structure such that  $K_2 \rightarrow \Gamma \rightarrow K_3$ . Then, VCSP( $\Gamma$ ) is NP-hard.

*Proof.* It is sufficient to show that the template  $(K_2, K_3)$  admits no cyclic fractional polymorphisms of arity at least 2 as in this case  $\Gamma$  has no such fractional polymorphisms, which implies that VCSP( $\Gamma$ ) is NP-hard by [KO15], cf. also [KKR17]. To see why  $(K_2, K_3)$  has no cyclic fractional polymorphisms of arity at least 2, we split into two cases. First, consider the case of even arity. In this case, consider the following matrix:

$$\begin{pmatrix} 0 & 1 & \cdots & 0 & 1 \\ 1 & 0 & \cdots & 1 & 0 \end{pmatrix}.$$

Any cyclic fractional symmetric polymorphism outputs a constant tuple (i.e. one with value 0) with probability 1; whereas all the columns of the matrix are within  $K_2$ , and thus the value of the outputted tuple should have expected value at least 1.<sup>23</sup> Thus no cyclic fractional polymorphisms of even arity exists. Now, consider odd arity not less than 3. In this case, consider the following matrix:

$$\begin{pmatrix} 0 & 1 & \cdots & 0 & 1 & 1 \\ 1 & 0 & \cdots & 1 & 1 & 0 \end{pmatrix}.$$

In the arity 3 case, for example, the matrix is

$$\begin{pmatrix} 0 & 1 & 1 \\ 1 & 1 & 0 \end{pmatrix}.$$

As before, the fractional polymorphism outputs a tuple with value 0, whereas it should output a tuple with value at least 2/3 (in any case, a positive value). Thus no such polymorphism exists.

<sup>&</sup>lt;sup>23</sup>The definition of a fractional polymorphism in [KO15, KKR17] requires that the expected value should be at most 1; this is because [KO15, KKR17] deal with minimisation problems.

**Remark 46.** To the best knowledge of the authors, our result is the first example of a promise  $\{0, 1\}$ -valued CSP [VŽ21] whose tractability does not come from a finite sandwich. There are examples of *non*- $\{0, 1\}$ -valued PVCSPs whose tractability does not come from a finite sandwich but these examples actually encode solving a Max-CSP approximately. For example, if  $\Gamma$  denotes a set of rational-valued functions let  $\rho\Gamma$  be the same set but with all functions scaled by  $\rho$ . Then, a  $\rho$ -approximation of any Max-CSP( $\Gamma$ ) is encoded by PVCSP( $\rho\Gamma, \Gamma$ ). We believe that these examples do not essentially use the gap between two combinatorially different structures, whereas ( $K_2, K_3$ ) *does* — and the reason why it must use this combinatorial structure is because the PCSP template ( $K_2, K_3$ ) is  $\{0, 1\}$ -valued.