

Proper and Improper Variants of Mathias and Silver Forcing

Liu Shixiao

Abstract

In this paper we answer several questions in [15] regarding density variants of Mathias and Silver forcing. These questions include whether each of the forcings is proper, adds dominating reals, or adds Cohen reals. We also generalize one of the proofs to Mathias forcings parametrized by lower semi-continuous submeasures satisfying certain properties.

1 Introduction

The aim of this paper is to establish several new results on variants of Mathias and Silver forcing introduced by Laguzzi, Mildenerger, and Stuber-Rousselle in [15]. These variants of Mathias and Silver forcing differ from previously considered ones as they are parametrized neither by filters nor by co-ideals, but by families of sets that satisfy certain conditions on their upper or lower density.

One of the reasons for analyzing such variants of Mathias and Silver forcing is to gain a better understanding of the cardinals $\text{cov}^*(\mathcal{Z}_0)$ and $\text{non}^*(\mathcal{Z}_0)$, which are cardinal characteristics associated to the ideal \mathcal{Z}_0 of sets of asymptotic density 0. Recall that:

Definition 1. Let $A \subseteq \omega$. The upper density and lower density of A are defined by

$$d^-(A) = \liminf_{n \rightarrow \infty} \frac{|A \cap n|}{n},$$

$$d^+(A) = \limsup_{n \rightarrow \infty} \frac{|A \cap n|}{n}.$$

In the case of $d^-(A) = d^+(A) = x$, we also say the density of A is x and write $d(A) = x$. And we define

$$\mathcal{Z}_0 = \{A \subseteq \omega : d(A) = 0\}$$

It is well-known (see [8] or [19]) that \mathcal{Z}_0 is an analytic P-ideal. Also recall that:

Definition 2. A function $\varphi : \mathcal{P}(\omega) \rightarrow [0, \infty]$ is called a *submeasure on ω* if

1. $\varphi(\emptyset) = 0$;
2. $\varphi(X) \leq \varphi(Y)$, for all $X \subseteq Y \subseteq \omega$;
3. $\varphi(X \cup Y) \leq \varphi(X) + \varphi(Y)$, for all $X, Y \in \mathcal{P}(\omega)$;
4. $\varphi(\{n\}) < \infty$, for all $n \in \omega$.

We say a submeasure φ on ω is *lower semi-continuous*, if $\varphi(X) = \lim_{n \rightarrow \infty} \varphi(X \cap n)$ for every $X \in \mathcal{P}(\omega)$. For a lower semi-continuous submeasure φ on ω , we define

$$\text{Exh}(\varphi) = \{X \subseteq \omega : \lim_{m \rightarrow \infty} \varphi(X \setminus m) = 0\}$$

It is not hard to see that $\text{Exh}(\varphi)$ is an $F_{\sigma\delta}$ P-ideal for every lower semi-continuous submeasure φ on ω . Indeed, analytic P-ideals are always of this form; by a theorem of Solecki [19], every analytic P-ideal \mathcal{I} is $F_{\sigma\delta}$ and there is a lower semi-continuous submeasure φ on ω such that $\mathcal{I} = \text{Exh}(\varphi)$. Cardinal invariants associated with such analytic ideals as well as properties of their quotients have been well-studied. For example, in [8], Farah studied gaps in quotients of the form $\mathcal{P}(\omega)/\mathcal{I}$, where \mathcal{I} is an analytic ideal, and in [9] he proved that $\mathcal{P}(\omega) / \mathcal{Z}_0$ is forcing equivalent to $(\mathcal{P}(\omega)/\text{FIN}) * \mathcal{B}$, where \mathcal{B} is a specific measure algebra. Brendle and Shelah [5] defined and investigated certain cardinal invariants for analytic ideals. In [12], Hernández-Hernández and Hrušák focused on four cardinal invariants associated to any tall analytic P-ideal, establishing various connections with the classical cardinal invariants.

Definition 3. A non-principal ideal \mathcal{I} on ω is said to be *tall* if for all $A \in [\omega]^{\aleph_0}$ there is $B \in [A]^{\aleph_0}$ such that $B \in \mathcal{I}$. For a non-principal tall ideal \mathcal{I} on ω , define the cardinals

$$\begin{aligned} \text{add}^*(\mathcal{I}) &= \min\{|\mathcal{F}| : \mathcal{F} \subseteq \mathcal{I} \wedge \forall X \in \mathcal{I} \exists A \in \mathcal{F} (A \not\subseteq^* X)\} \\ \text{cov}^*(\mathcal{I}) &= \min\{|\mathcal{F}| : \mathcal{F} \subseteq \mathcal{I} \wedge \forall X \in [\omega]^{\aleph_0} \exists A \in \mathcal{F} (|A \cap X| = \aleph_0)\} \\ \text{cof}^*(\mathcal{I}) &= \min\{|\mathcal{F}| : \mathcal{F} \subseteq \mathcal{I} \wedge \forall X \in \mathcal{I} \exists A \in \mathcal{F} (X \subseteq^* A)\} \\ \text{non}^*(\mathcal{I}) &= \min\{|\mathcal{F}| : \mathcal{F} \subseteq [\omega]^{\aleph_0} \wedge \forall X \in \mathcal{I} \exists A \in \mathcal{F} (|A \cap X| < \aleph_0)\}. \end{aligned}$$

The cardinals $\text{cov}^*(\mathcal{I})$ and $\text{non}^*(\mathcal{I})$ are duals. Similarly, $\text{add}^*(\mathcal{I})$ and $\text{cof}^*(\mathcal{I})$ are duals. See [2] for more about duality of cardinal invariants. For definitions of the classical cardinal invariants, such as $\mathfrak{b}, \mathfrak{d}, \mathfrak{s}, \mathfrak{r}$, the reader can refer to Blass [2] or to Bartoszyński and Judah [1]. In [12], Hernández-Hernández and Hrušák pointed out that $\text{cov}^*(\mathcal{Z}_0)$ and $\text{non}^*(\mathcal{Z}_0)$ play a basic role in the general theory these invariants. They established (see Theorems 3.10 and 3.12 of [12]) that $\min\{\text{cov}(\mathcal{N}), \mathfrak{b}\} \leq \text{cov}^*(\mathcal{Z}_0) \leq \text{non}(\mathcal{M})$, $\text{cov}(\mathcal{M}) \leq \text{non}^*(\mathcal{Z}_0) \leq \max\{\mathfrak{d}, \text{non}(\mathcal{N})\}$, $\text{cov}^*(\mathcal{Z}_0) \leq \max\{\mathfrak{b}, \text{non}(\mathcal{N})\}$, and that $\text{non}^*(\mathcal{Z}_0) \leq \min\{\mathfrak{d}, \text{cov}(\mathcal{N})\}$. They raised the question of whether $\text{cov}^*(\mathcal{Z}_0) \leq \mathfrak{d}$. Raghavan and Shelah [18] answered their question by proving the dual inequalities $\text{cov}^*(\mathcal{Z}_0) \leq \mathfrak{d}$ and $\mathfrak{b} \leq \text{non}^*(\mathcal{Z}_0)$ in ZFC. Raghavan [17] further improved these results by showing in ZFC that $\min\{\mathfrak{d}, \mathfrak{r}\} \leq \text{non}^*(\mathcal{Z}_0)$ and that $\text{cov}^*(\mathcal{Z}_0) \leq \max\{\mathfrak{b}, \mathfrak{s}(\mathfrak{pr})\}$. Here, $\mathfrak{s}(\mathfrak{pr})$ is defined to be the minimal cardinality of a family \mathcal{F} of partitions of ω into infinitely many pieces such that for every $X \in [\omega]^{\aleph_0}$, there is a partition in \mathcal{F} such that all pieces have infinite intersection with X . It is not known whether $\mathfrak{s}(\mathfrak{pr}) = \mathfrak{s}$.

The cardinal $\text{cov}^*(\mathcal{I})$ is closely connected with the question of which forcing notions can diagonalize \mathcal{I} . For any ideal $\mathcal{I} \in V$, a forcing $\mathbb{P} \in V$ is said to *diagonalize* \mathcal{I} if there is a \mathbb{P} -name \dot{X} such that $\Vdash_{\mathbb{P}} \dot{X} \in [\omega]^{\aleph_0}$ and for each $I \in V \cap \mathcal{I}$, $\Vdash_{\mathbb{P}} |\dot{X} \cap I| < \aleph_0$. Note that if \mathbb{P} diagonalizes \mathcal{I} , then \mathbb{P} tends to increase $\text{cov}^*(\mathcal{I})$. Furthermore, if \mathcal{I} is *Katětov below* \mathcal{J} , which means that $\exists f \in \omega^\omega \forall I \in \mathcal{I} (f^{-1}(I) \in \mathcal{J})$, then any \mathbb{P} that diagonalizes \mathcal{J} will also diagonalize \mathcal{I} . The question of whether an ideal can be diagonalized without adding certain types of reals has a long history in set theory of the reals. For instance, Laflamme [14] showed that every F_σ ideal can be diagonalized by an ω^ω -bounding forcing, and Canjar [6] showed that certain ultrafilters can be diagonalized without adding dominating reals. The result of [18] shows that any proper forcing which diagonalizes \mathcal{Z}_0 must add an unbounded real, and it is proved in [17] that any Suslin c.c.c. forcing which diagonalizes \mathcal{Z}_0 must add a dominating real. Raghavan [17] also asked whether \mathcal{Z}_0 can be diagonalized by any proper forcing without adding dominating reals; this question still remains open.

Brendle, Guzmán, Hrušák and Raghavan [3] have recently shown that the ideal generated by a m.a.d. family is always *nearly* Katětov below \mathcal{Z}_0 , which means that the reduction holds after removing countably many members of the m.a.d. family. It is a long standing open problem whether $\mathfrak{d} = \aleph_1$ implies $\mathfrak{a} = \aleph_1$. In [4], Brendle and Raghavan raised the question of whether $\mathfrak{b} = \mathfrak{s} = \aleph_1$ already implies $\mathfrak{a} = \aleph_1$, and their question remains open as well. The results of [3] show that an understanding of what forcing notions are capable of diagonalizing \mathcal{Z}_0 is useful for analyzing \mathfrak{a} . By [4], for any collection of closed sets whose union is a m.a.d. family, there is a forcing that does not add dominating reals and forces that the reinterpreted union is not maximal in the extension.

Laguzzi, Mildner, and Stuber-Rousselle [15] recently introduced new variants of Mathias and Silver forcing parametrized by upper or lower density. Their variants of Mathias forcing diagonalize \mathcal{Z}_0 .

Definition 4. The Mathias forcing \mathbb{M} consists of tuples (s, A) , where $s \subset \mathbb{N}$ is finite, $A \subseteq \mathbb{N}$ is infinite, and $\max s \leq \min A$. For two tuples, $(s_1, A_1) \leq (s_2, A_2)$ iff

- $s_1 \supseteq s_2$
- $A_1 \subseteq A_2$
- If $n \in s_1 \setminus s_2$, then $n \in A_2$

The density variants of Mathias forcing are denoted with superscripts $^-$ $^+$ and subscripts \geq_ε $>_0$, where the density requirements are imposed on the infinite set A . For instance, $\mathbb{M}_{\geq_\varepsilon}^-$ requires $d^-(A) \geq \varepsilon$ for all conditions (s, A) , and $\mathbb{M}_{>_0}^+$ requires $d^+(A) > 0$.

Definition 5. The Silver forcing \mathbb{V} consists of partial functions $f : \omega \rightarrow 2$ such that $\omega \setminus \text{dom}(f)$ is infinite. For two conditions, $f_1 \leq f_2$ iff $f_1 \supseteq f_2$. The density variants of Silver forcing are denoted with superscripts $^-$ $^+$ and subscripts \geq_ε $>_0$, where the density requirements are imposed on $\omega \setminus \text{dom}(f)$. For instance, $\mathbb{V}_{\geq_\varepsilon}^-$ requires $d^-(\omega \setminus \text{dom}(f)) \geq \varepsilon$ for all conditions f , and $\mathbb{V}_{>_0}^+$ requires $d^+(\omega \setminus \text{dom}(f)) > 0$.

Properties of the classical Mathias and Silver models are well-known. See [1]. Nevertheless new properties of these models are still being discovered, for instance [7] showed there are no P-points in the Silver model. While many variants of Mathias and Silver forcing have been considered previously, for example [10] and [13], the ones considered by Laguzzi, Mildner, and Stuber-Rousselle in [15] are quite different because they are determined by neither a filter nor a co-ideal. Their results are summarized in the table below. Note that the paper also claims to have proven the properness of $\mathbb{M}_{\geq_\varepsilon}^-$, but we will soon show an error in their proof.

	$\mathbb{M}_{>_0}^-$	$\mathbb{M}_{>_0}^+$	$\mathbb{M}_{\geq_\varepsilon}^-$	$\mathbb{M}_{\geq_\varepsilon}^+$	$\mathbb{V}_{>_0}^-$	$\mathbb{V}_{>_0}^+$	$\mathbb{V}_{\geq_\varepsilon}^-$	$\mathbb{V}_{\geq_\varepsilon}^+$
proper		✓		✓	✗			✓
dominating		✓		✓	✓			✗
Cohen	✓	✓	✓	✓	✓			✗

Several questions were left open in [15], indicated by the blank boxes in the table above. In section 2 of this paper, we study the Mathias forcings $\mathbb{M}_{\geq_\varepsilon}^-$ and $\mathbb{M}_{>_0}^-$. In particular, Question 3.13 of [15] asks whether $\mathbb{M}_{\geq_\varepsilon}^-$ adds dominating reals. They pointed out that since $\mathbb{M}_{\geq_\varepsilon}^-$ diagonalizes \mathcal{Z}_0 , a negative answer to their question would give a positive answer to Question 38 of [17]. While

a general criterion for whether a Mathias-Prikry type forcing adds a dominating real is given in [13], this is not applicable to $\mathbb{M}_{\geq \varepsilon}^-$ because it is not parametrized by a filter. We will give a positive answer to Question 3.13 of [15] by showing that $\mathbb{M}_{\geq \varepsilon}^-$ always adds a dominating real. In addition to this, we will introduce a generalization of $\mathbb{M}_{\geq \varepsilon}^-$ for a sequence of lower semi-continuous submeasures. In section 3 we study the Silver forcings $\mathbb{V}_{\geq \varepsilon}^-$ and $\mathbb{V}_{>0}^+$. We will show that they both collapse the continuum to ω . With all these results, we complete the above table by the end of this paper. Some of the results will rely on the Darboux property of upper and lower density, which can be found in [16].

2 Mathias Forcings with Density

In this section we deal with properties of two Mathias forcings that remain unproven in [15]. Let's first show a counterexample to Corollary 3.5 of [15]. The corollary claims that, for any set A with $d^-(A) = \epsilon$, the set $\mathcal{F}(A) = \{B \subseteq A : d^-(B) = \epsilon\}$ is a filter. This erroneous corollary was used in the paper to prove the properness of $\mathbb{M}_{\geq \varepsilon}^-$.

Example 6. Divide \mathbb{N} into intervals $[a_i, a_{i+1})$ such that $a_0 = 0$ and $a_{i+1} = 2^{a_i}$. We take A, B_1, B_2 such that

$$\begin{aligned} A \cap [a_i, a_{i+1}) &= \begin{cases} [a_i, a_{i+1}) & \text{if } i \text{ is even} \\ \{2n : n \in \mathbb{N}\} \cap [a_i, a_{i+1}) & \text{if } i \text{ is odd} \end{cases} \\ B_1 \cap [a_i, a_{i+1}) &= \begin{cases} \{2n : n \in \mathbb{N}\} \cap [a_i, a_{i+1}) & \text{if } i \text{ is even} \\ \{2n : n \in \mathbb{N}\} \cap [a_i, a_{i+1}) & \text{if } i \text{ is odd} \end{cases} \\ B_2 \cap [a_i, a_{i+1}) &= \begin{cases} \{2n+1 : n \in \mathbb{N}\} \cap [a_i, a_{i+1}) & \text{if } i \text{ is even} \\ \{2n : n \in \mathbb{N}\} \cap [a_i, a_{i+1}) & \text{if } i \text{ is odd} \end{cases} \end{aligned}$$

Then we have $d^-(A) = d^-(B_1) = d^-(B_2) = 1/2$ and $B_1, B_2 \subseteq A$, but $d^-(B_1 \cap B_2) = 0$. Hence $\{B \subseteq A : d^-(B) = 1/2\}$ is not a filter.

Furthermore, consider the equivalence relation $=^*$ on 2^ω , namely $f =^* g$ iff they are equal modulo a finite set. Pick one representative f_α from each equivalence class. The set $\{f_\alpha : \alpha \in I\}$ has size 2^{\aleph_0} . For each f_α , we take $B_\alpha \subseteq \mathbb{N}$ such that

$$B_\alpha \cap [a_i, a_{i+1}) = \begin{cases} \{2n+1 : n \in \mathbb{N}\} \cap [a_i, a_{i+1}) & \text{if } i \text{ is even and } f_\alpha(i/2) = 1 \\ \{2n : n \in \mathbb{N}\} \cap [a_i, a_{i+1}) & \text{if } i \text{ is odd, or } i \text{ is even but } f_\alpha(i/2) = 0 \end{cases}$$

By definition, each B_α has density $1/2$ and $B_\alpha \subseteq A$. Moreover, for $f_\alpha \neq f_\beta$, the intersection $B_\alpha \cap B_\beta$ will be empty on infinitely many intervals $[a_i, a_{i+1})$. Since the size of these intervals grow exponentially, we have that $d^-(B_\alpha \cap B_\beta) = 0$. Therefore, the set $\{(\emptyset, B_\alpha) : \alpha \in I\}$ is an antichain in $\mathbb{M}_{\geq 1/2}^-$ of size 2^{\aleph_0} , and it lies below the condition (\emptyset, A) with $d^-(A) = 1/2$.

Since the original proof is based on an incorrect corollary, we need to give a new proof that $\mathbb{M}_{\geq \varepsilon}^-$ is proper. Indeed, we prove it for the generalized forcing notion.

Definition 7. For each $m \in \mathbb{N}$, let $\varphi_m : \mathcal{P}(\mathbb{N}) \rightarrow [0, \infty]$ be a lower semi-continuous submeasure on \mathbb{N} , such that $\lim_{m \rightarrow \infty} \varphi_m(\{n\}) = 0$ for all $n \in \mathbb{N}$. Then we can define the submeasure $\varphi^-(A) :=$

$\liminf_{m \rightarrow \infty} \varphi_m(A)$ and the corresponding Mathias forcing

$$\mathbb{M}_{\geq \varepsilon}^{\varphi^-} := \{(s, A) : \max(s) < \min(A) \text{ and } \varphi^-(A) \geq \varepsilon\}$$

where the ordering is defined as usual.

It's clear that, when we take $\varphi_m(A) = \frac{|A \cap m|}{m}$, the above definition gives us $\mathbb{M}_{\geq \varepsilon}^-$, so it indeed generalizes Mathias forcing with lower density. Other examples satisfying the conditions include weighted density $\varphi_m(A) = \frac{\sum_{i \in A \cap m} f(i)}{\sum_{i < m} f(i)}$, where f is the weight function (see also Erdős-Ulam submeasure, for instance in [8]), and the filter of cofinite sets (see Example 11 below).

Fact 8. If $A =^* B$, then $\varphi^-(A) = \varphi^-(B)$.

Proof. Suppose $A \cup A_0 = B \cup B_0$ where A_0, B_0 are both finite. Then by subadditivity we have $\lim_{m \rightarrow \infty} \varphi_m(A_0) = \lim_{m \rightarrow \infty} \varphi_m(B_0) = 0$. Hence

$$\varphi^-(A) = \liminf_{m \rightarrow \infty} \varphi_m(A) \leq \liminf_{m \rightarrow \infty} \varphi_m(B \cup B_0) \leq \liminf_{m \rightarrow \infty} \varphi_m(B) + \liminf_{m \rightarrow \infty} \varphi_m(B_0) = \varphi^-(B)$$

and vice versa. \square

Theorem 9. $\mathbb{M}_{\geq \varepsilon}^{\varphi^-}$ is proper for any $\varepsilon \geq 0$.

Proof. Take κ large enough and $\mathcal{M} < H_\kappa$ countable with $\mathbb{M}_{\geq \varepsilon}^{\varphi^-} \in \mathcal{M}$. List all ordinal names $\dot{\alpha}_0, \dot{\alpha}_1, \dots$ in \mathcal{M} . Fix $(s, A_0) \in \mathbb{M}_{\geq \varepsilon}^{\varphi^-} \cap \mathcal{M}$ and we find an extension $(s, B) \leq (s, A_0)$ and sets Γ_n of ordinals in \mathcal{M} such that $(s, B) \Vdash \dot{\alpha}_n \in \Gamma_n$ for all $n \in \mathbb{N}$.

The construction is done by induction. In each stage $i + 1$, we construct finite sets B_{i+1} and $\Gamma_{n,i+1}$, and in the end we let $B = \bigcup_{i \geq 1} B_i$ and $\Gamma_n = \bigcup_{i \geq 1} \Gamma_{n,i}$. We also construct auxiliary sequences $\{A_i\}, \{N_i\}, \{M_i\}$, where $A_0 \supseteq A_1 \supseteq A_2 \supseteq \dots$ is a shrinking sequence of infinite subsets of \mathbb{N} , $0 = N_0 < N_1 < N_2 < \dots$ are the boundaries of A_{i+1} at each stage where we freeze anything below N_i and only shrink beyond N_i , and $0 = M_0 < M_1 < M_2 < \dots$ are the boundaries of m at each stage such that $\varphi_m(A_{i+1})$ is close enough to ε for all $m \geq M_{i+1}$. Each stage $i + 1$ will be a recursive construction on its own, where we find the sequence $A_i = A_{i+1,0} \supseteq A_{i+1,1} \supseteq \dots \supseteq A_{i+1,l} = A_{i+1}$ such that $A_{i+1,m} \cap N_i = A_i \cap N_i$ for all m , and therefore $A_i \cap N_i = A_{i+1} \cap N_i$.

- Stage 0:

A_0 is given by the condition (s, A_0) we fix. Start with $M_0 = N_0 = 0$.

- Stage $i + 1$:

There are finitely many possible ways to extend s with elements in $\bigcup_{k=0}^i B_k$, since each B_k is finite. Enumerate all such extensions as $s_i^0, s_i^1, \dots, s_i^p$. Start with $A_{i+1,0} = A_i$. Let $\{(n_m, j_m) : 0 \leq m < (p+1)(i+2)\}$ enumerate all pairs (n, j) such that $0 \leq n \leq i+1$ and $0 \leq j \leq p$. In each step m , ask (in \mathcal{M}) whether:

$$\exists C \subseteq (A_{i+1,m} \setminus N_i) \exists \beta \in \text{Ord} (s_i^{j_m}, C) \Vdash \dot{\alpha}_{n_m} = \check{\beta}$$

1. If the answer is “no”, let $A_{i+1,m+1} = A_{i+1,m}$.

2. If the answer is “yes”, choose a pair of witnesses $C_{i+1,m}$ and $\beta_{i+1,m}$ for C and β , and let $A_{i+1,m+1} = (A_i \cap N_i) \cup C_{i+1,m}$. Note that $(s_i^j, C_{i+1,m})$ is a condition and therefore $\varphi^-(A_{i+1,m+1}) \geq \varepsilon$.

In the end we let $A_{i+1} = A_{i+1,(p+1)(i+2)-1}$ and $\Gamma_{n,i+1} = \{\beta_{i+1,m} : n_m = n\}$. Clearly $A_{i+1} \cap N_i = A_i \cap N_i$ and each $\Gamma_{n,i+1}$ is a finite set of ordinals in \mathcal{M} .

For the boundary at this stage, since $\varphi^-(A_{i+1}) \geq \varepsilon$, we can find $M_{i+1} > M_i$ such that $\varphi_m(A_{i+1}) \geq \varepsilon - \frac{1}{2^{i+1}}$ for all $m \geq M_{i+1}$. By lower semi-continuity, for each m , there is $L_{i+1,m}$ such that $|\varphi_m(A_{i+1}) - \varphi_m(A_{i+1} \cap n)| \leq \frac{1}{2^{i+1}}$ for all $n \geq L_{i+1,m}$. Let $N_{i+1} = N_i + \max\{L_{i+1,m} : m < M_{i+1}\}$.

For the last action at this stage, let $B_{i+1} = A_i \cap [N_i, N_{i+1})$. Note that B_{i+1} is not a truncate of A_{i+1} , but rather, a truncate of A_i from the previous stage.

Finally, let $B = \bigcup_{i \geq 1} B_i$ and $\Gamma_n = \bigcup_{i \geq 1} \Gamma_{n,i}$ as we promised. Notice that all B_{i+1} are disjoint. Also, $A_i \cap N_i = A_{i+1} \cap N_i$ for all i , hence we have $B \cap N_{i+1} = A_i \cap N_{i+1}$. Now we prove that B and Γ_n we constructed do satisfy the requirements.

- $B \subseteq A$.

Each B_{i+1} is a truncate of A_i , and $A_i \subseteq \dots \subseteq A_0 = A$. In fact, $B \subseteq A_i$ for all i .

- $\varphi_m(B \cap n) \geq \varepsilon - \frac{1}{2^{i-1}}$ holds if for some $i \geq 1$ we have $M_i \leq m < M_{i+1}$ and $n \geq N_{i+1}$, and hence $\varphi^-(B) \geq \varepsilon$.

Since $m \geq M_i$, we have $\varphi_m(A_i) \geq \varepsilon - \frac{1}{2^i}$. Since $m < M_{i+1}$ and therefore $N_{i+1} > L_{i+1,m}$, we have $|\varphi_m(A_i) - \varphi_m(A_i \cap N_{i+1})| \leq \frac{1}{2^i}$. Thus $\varphi_m(B \cap n) \geq \varphi_m(B \cap N_{i+1}) = \varphi_m(A_i \cap N_{i+1}) \geq \varphi_m(A_i) - |\varphi_m(A_i) - \varphi_m(A_i \cap N_{i+1})| \geq \varepsilon - \frac{1}{2^{i-1}}$.

- $(s, B) \Vdash \dot{\alpha}_n \in \Gamma_n$ for all n .

Take an arbitrary extension $(t, D) \leq (s, B)$ such that $(t, D) \Vdash \dot{\alpha}_n = \gamma$ for some $\gamma \in \text{Ord}$ and some $n \in \mathbb{N}$. Then t extends s with elements from $\bigcup_{k=0}^i B_k$ for some i . Without loss of generality suppose $i \geq n$. So by our construction, $t = s_i^j$ for some j . In the corresponding step m of stage $i+1$, we asked in \mathcal{M} whether

$$\exists C \subseteq (A_{i+1,m} \setminus N_i) \exists \beta \in \text{Ord} (s_i^j, C) \Vdash \dot{\alpha}_n = \check{\beta}$$

By elementarity of $\mathcal{M} < H_\kappa$, we get the same answer as asking in H_κ , and the answer must be “yes”, because $(s_i^j, D) \Vdash \dot{\alpha}_n = \gamma$ and $D \subseteq B \subseteq A_{i+1,m}$, and thus $(s_i^j, D \cap (A_{i+1,m} \setminus N_i)) \Vdash \dot{\alpha}_n = \gamma$. Since the answer is “yes”, we must’ve set $A_{i+1,m+1} = (A_i \cap N_i) \cup C_{i+1,m}$ at this step and thus $D \subseteq B \subseteq A_{i+1} \subseteq^* C_{i+1,m}$. Therefore $(s_i^j, D \cap C_{i+1,m})$ is a common extension of $(s_i^j, C_{i+1,m})$ and (s_i^j, D) . Now $(s_i^j, C_{i+1,m}) \Vdash \dot{\alpha}_n = \beta_{i+1,m}$. On the other hand, $(s_i^j, D) \Vdash \dot{\alpha}_n = \gamma$. Hence $\beta_{i+1,m} = \gamma$. That is to say, $\gamma \in \Gamma_{n,i+1}$, and hence $(t, D) \Vdash \dot{\alpha}_n \in \Gamma_n$. By the arbitrariness of (t, D) , we see that $(s, B) \Vdash \dot{\alpha}_n \in \Gamma_n$.

□

Corollary 10. $\mathbb{M}_{\geq \varepsilon}^-$ is proper for any $0 \leq \varepsilon \leq 1$.

□

Now the remaining question is whether $\mathbb{M}_{\geq \varepsilon}^{\varphi^-}$ adds a dominating real. The answer is: it depends. We give one example of such forcing not adding dominating reals and one that does.

Example 11. Consider $\mathbb{M}_{\geq \varepsilon}^{\varphi^-}$ where

$$\varphi_m(A) = \begin{cases} 1 & m \in A \\ 0 & m \notin A \end{cases}$$

and $0 < \varepsilon \leq 1$. Then $\varphi^-(A) \geq \varepsilon$ iff A is cofinite. Hence Cohen forcing densely embeds in $\mathbb{M}_{\geq \varepsilon}^{\varphi^-}$ by the map $s \mapsto (s^{-1}(1), \omega \setminus \text{dom}(s))$. Therefore it does not add dominating reals.

Next, we will show that $\mathbb{M}_{\geq \varepsilon}^-$ adds a dominating real. The proof relies on the following (weak) Darboux property of lower density:

Lemma 12. For any $A \subseteq N$ with $d^-(A) = \delta > \varepsilon$, there is $A' \subset A$ such that $d^-(A') = \varepsilon$.

Proof. See [16] Corollary 1&2. □

Theorem 13. $\mathbb{M}_{\geq \varepsilon}^-$ adds a dominating real for any $0 < \varepsilon \leq 1$.

Proof. Let G be the generic filter and $g = \bigcup \{s : \exists A (s, A) \in G\}$. By the Darboux property, we can fix a maximal antichain $M \subseteq \mathbb{M}_{\geq \varepsilon}^-$ such that $d^-(A) = \varepsilon$ for all $(s, A) \in M$. Suppose $M \cap G = \{(s_0, A_0)\}$. Enumerate A_0 as $a_0 < a_1 < a_2 < \dots$ and let $A_n = \{a_i : i = 2^k n, k \in \mathbb{N}^+\}$ for $n \geq 1$. Clearly $A_0 \supset A_1 \supset A_2 \supset \dots$ and $d^+(A_n) = \frac{1}{2^n} d^+(A_0) \leq \frac{1}{2^n}$. Also, $d^-(A_0 \setminus A_n) = \frac{2^n - 1}{2^n} \varepsilon$. Therefore $A \cap A_n \neq \emptyset$ for all $A \subseteq A_0$ with $d^-(A) = \varepsilon$ and for all n .

In $V[G]$, define function $F : \omega \rightarrow \omega$ by

$$F(n) = \min\{k : k \in g \cap A_n\}$$

Such k exists because $D_n = \{(s, A) : \exists k \in s \cap A_n\}$ is dense below (s_0, A_0) . We show that F is a dominating real.

Fix $f \in (\omega^\omega)^V$ and $(s, A) \in \mathbb{M}_{\geq \varepsilon}^-$ such that $(s, A) \leq (s_0, A_0)$. Let

$$A_f = \bigcup_{n \in \omega} \{k \in A_n : k \leq f(n)\}$$

We claim that $d^-(A \setminus A_f) = \varepsilon$. Indeed, since each summand of A_f is finite, for all $m \in \mathbb{N}$ we have

$$\begin{aligned} d^+(A_f) &= d^+\left(\bigcup_{n=1}^{\infty} \{k \in A_n : k \leq f(n)\}\right) \\ &= d^+\left(\bigcup_{n=m}^{\infty} \{k \in A_n : k \leq f(n)\}\right) \\ &\leq d^+\left(\bigcup_{n=m}^{\infty} A_n\right) \\ &= d^+(A_m) \leq \frac{1}{2^m} \end{aligned}$$

and therefore $d(A_f) = 0$. Thus $(s, A \setminus A_f)$ is a condition. Then, since the infinite part $A \setminus A_f$ has excluded any $k \in A_n$ with $k \leq f(n)$, we have

$$(s, A \setminus A_f) \Vdash \forall n (\dot{F}(n) \geq \min(A \setminus A_f) \rightarrow f(n) < \dot{F}(n)).$$

Note that $F(n) \in A_n$ and $\min(A_n) \geq 2^n$, and therefore this translates to

$$(s, A \setminus A_f) \Vdash \forall n > \log_2 \min(A \setminus A_f) (f(n) < \dot{F}(n)).$$

Hence $D_f = \{(s, A) : (s, A) \Vdash \exists m \forall n > m (f(n) < \dot{F}(n))\}$ is dense below (s_0, A_0) for all ground model $f \in \omega^\omega$. \square

Note that for the submeasure φ^- in Example 11, if we attempt to apply the same proof, A_f will be infinite and therefore $(s, A \setminus A_f)$ is no longer a condition.

We switch to another density Mathias forcing. Instead of $d^-(A) \geq \varepsilon$, the condition is now $d^-(A) > 0$. This forcing notion was also investigated by Matthew Harrison-Trainor, Liu Lu and Patrick Lutz from a recursion-theoretic perspective in [11]. The following proof is inspired by Proposition 4.5 of their paper, where they constructed $A_X \subseteq N$ for a fixed set $X \subseteq N$, such that any subset $B \subseteq A_X$ with $d^-(B) > 0$ computes X uniformly.

Theorem 14. $\mathbb{M}_{>0}^-$ collapses the continuum to ω .

Proof. Fix a bijection $\langle \cdot, \cdot, \cdot \rangle : \mathbb{N} \times \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ such that $\langle 0, 0, 0 \rangle = 0$. In the following we construct disjoint intervals $I_{a,b,c}$ for each $n = \langle a, b, c \rangle$. Later, for some fixed condition (s, A) and set $S \in (2^\omega)^V$, we will code S with a one-step extension $(s, A_0) \leq (s, A)$. The b -th digit of S will be stored infinitely many times in $A_0 \cap I_{a,b,c}$ for the smallest a such that $d^-(A) > 1/(a+2)$ and for all $c \in \mathbb{N}$.

We build $I_{a,b,c}$ inductively. Let $I_{0,0,0} = [0, 1)$. Suppose $I_{a,b,c} = [i_n, i_{n+1})$ is constructed for $\langle a, b, c \rangle = n$, then let $I_{a,b,c} = [i_{n+1}, (2a+4)^{2c+4} i_{n+1})$ for $\langle a, b, c \rangle = n+1$. We further divide each $I_{a,b,c}$ into intervals $I_{a,b,c}^k = [(2a+4)^k \cdot i_{n+1}, (2a+4)^{k+1} \cdot i_{n+1})$ for $0 \leq k \leq 2c+3$. Later in the coding step $A_0 \subseteq A$, depending on whether the b -th digit of S is 1 or 0, we will either store the information in odd intervals $I_{a,b,c}^1, I_{a,b,c}^3, \dots, I_{a,b,c}^{2c+1}$, or in even intervals $I_{a,b,c}^2, I_{a,b,c}^4, \dots, I_{a,b,c}^{2c+2}$. The first interval $I_{a,b,c}^0$ and last interval $I_{a,b,c}^{2c+3}$ will act as buffer areas; they aren't necessary but having them makes the proof easier.

Claim. Fix a_0 such that $d^-(A) > 1/(a_0+2)$. Take $A_0 \subseteq A$ satisfying

- For $a = a_0$ and $b, c \in \mathbb{N}$, either

$$I_{a,b,c} \cap (A \setminus A_0) = I_{a,b,c}^2 \cup I_{a,b,c}^4 \cup \dots \cup I_{a,b,c}^{2c+2}$$

or

$$I_{a,b,c} \cap (A \setminus A_0) = I_{a,b,c}^1 \cup I_{a,b,c}^3 \cup \dots \cup I_{a,b,c}^{2c+1};$$

- For $a \neq a_0$ and $b, c \in \mathbb{N}$, $I_{a,b,c} \cap (A \setminus A_0) = \emptyset$.

Then $d^-(A_0) > 0$.

Proof of claim. Since $d^-(A) > 1/(a_0 + 2)$, there is n such that $\frac{|x \cap A|}{x} > \frac{1}{a_0 + 2}$ for all $x > n$. Consider any interval $I_{a,b,c}^k = [i, (2a+4)i)$ such that $i \gg n$. It is of one of the following three cases:

1. $a = a_0$ and the previous interval $I_{a_0,b,c}^{k-1} = [\frac{i}{2a_0+4}, i)$ has empty intersection with $A \setminus A_0$:

By counting we have $|I_{a_0,b,c}^{k-1} \cap A_0| = |I_{a_0,b,c}^{k-1} \cap A| \geq |i \cap A| - \frac{i}{2a_0+4} > \frac{i}{2a_0+4}$. Thus

$$\frac{|x \cap A_0|}{x} \geq \frac{|I_{a_0,b,c}^{k-1} \cap A_0|}{x} > \frac{1}{(2a_0+4)^2} \text{ for all } x \in I_{a_0,b,c}^k.$$

2. $a = a_0$ and the previous interval $I_{a_0,b,c}^{k-1} \subset A \setminus A_0$:

In this case we must have $k \geq 2$, and $I_{a_0,b,c}^{k-2}$ has empty intersection with $A \setminus A_0$. Therefore

$$\frac{|x \cap A_0|}{x} \geq \frac{|I_{a_0,b,c}^{k-2} \cap A_0|}{x} > \frac{1}{(2a_0+4)^3} \text{ for all } x \in I_{a_0,b,c}^k.$$

3. $a = a_0$ and $k = 0$, or $a \neq a_0$;

Notice that we only remove elements from $I_{a,b,c} \cap A$ when $a = a_0$. Also we have a buffer area $I_{a_0,b,c}^{2c+3}$ at the end of each $I_{a_0,b,c}$ which we never touch. Whether $[\frac{i}{2a_0+4}, i)$ is the previous interval or not, we always have $[\frac{i}{2a_0+4}, i) \cap (A \setminus A_0) = \emptyset$. By the same argument as in 1.,

$$\text{we have } \frac{|x \cap A_0|}{x} > \frac{1}{(2a_0+4)^2} \text{ for all } x \in I_{a,b,c}^k.$$

Hence, we have $d^-(A_0) \geq \frac{1}{(2a_0+4)^3} > 0$. □

Now let's define the coding. Let G be the generic filter and $g = \bigcup \{s : (s, A) \in G\}$. Define $F : \mathbb{N} \times \mathbb{N} \rightarrow 2^\omega$ in $V[G]$ as follows. Fix $a, b, n \in \mathbb{N}$. Let

$$l(a, b) = \min\{l \in \mathbb{N} : \exists c \exists k (l \in g \cap I_{a,b,c}^k \text{ and } 1 \leq k \leq 2c+2)\}.$$

To see such l exists, notice that missing all the consecutive intervals $I_{a,b,c}^k$ for $1 \leq k \leq 2c+2$ forces $\frac{|x \cap A|}{x}$ to temporarily drop below $\frac{1}{(2a+4)^{2c+2}}$, and therefore any set A with $d^-(A) > 0$ cannot miss all these intervals for fixed a, b . Now define $F(a) \in 2^\omega$ by

$$F(a)(b) = \begin{cases} 1 & \text{if } l(a, b) \in I_{a,b,c}^k \text{ where } k \text{ is odd} \\ 0 & \text{if } l(a, b) \in I_{a,b,c}^k \text{ where } k \text{ is even} \end{cases}$$

and \dot{F} be a name for F . It remains to show $D_S = \{(s, A) : \exists a (s, A) \Vdash \dot{F}(a) = S\}$ is dense for all $S \in (2^\omega)^V$.

Fix (s, A) and $S \in (2^\omega)^V$. Choose the smallest natural number a_0 such that $d^-(A) > 1/(a_0+2)$. Now, for all $b, c \in \mathbb{N}$, we remove from A all the even intervals $I_{a_0,b,c}^k$ if $S(b) = 1$, and all the odd intervals $I_{a_0,b,c}^k$ if $S(b) = 0$. That is, take $A_0 \subseteq A$ such that

$$A \setminus A_0 = \bigcup \left\{ I_{a_0,b,c}^k : S(b) \equiv k+1 \pmod{2}, 1 \leq k \leq 2c+2 \right\}$$

By the above claim, (s, A_0) is a condition. Now, (s, A_0) already decides whether $l(a_0, b)$ lies in an odd interval (or even interval) $I_{a_0,b,c}^k$, because either all the even intervals (or all the odd intervals)

have empty intersection with A_0 , depending on the value of $S(b)$. Thus $(s, A_0) \Vdash \dot{F}(a_0) = S$. Therefore, D_S is dense for all ground model $S \in 2^\omega$. \square

3 Silver Forcings with Density

In this section we investigate the Silver forcings that are yet to be categorized in [15]. The first density Silver forcing we investigate is $\mathbb{V}_{\geq \varepsilon}^-$. Unlike the Mathias case, this forcing collapses the continuum. The intuition is, each condition f is allowed to contain infinite positive information, and hence we can code any real in the ground model via one-step extension, by simply adding it to the domain of f . The tough part is how to recover this information from the generic. Namely, we need to find which bits of g were used to code f back in the one-step extension. We start with a lemma.

Lemma 15. Fix A such that $d^-(A) = \varepsilon$. Then there exists an infinite sequence $N_0 < N_1 < N_2 < \dots$ such that for every $B \subseteq A$ with $d^-(B) = \varepsilon$, we always have

$$\lim_{i \rightarrow \infty} \frac{|(A \setminus B) \cap [N_i, N_{i+1})|}{N_{i+1} - N_i} = 0$$

Moreover, $\{N_i\}_{i \in \omega}$ can grow arbitrarily fast.

Proof. Take $\{N_i\}_{i \in \omega}$ such that $\frac{|A \cap N_i|}{N_i} \leq \varepsilon + \frac{1}{i}$ and $N_{i+1} > 2N_i$ for all i . Such sequence exists since $d^-(A) = \varepsilon$. Suppose we have some $B \subseteq A$ with $d^-(B) = \varepsilon$ such that

$$\frac{|(A \setminus B) \cap [N_i, N_{i+1})|}{N_{i+1} - N_i} \geq \delta$$

for infinitely many $i \in \mathbb{N}$ and some fixed $\delta > 0$. Then for these $i \in \mathbb{N}$, we have

$$\frac{|B \cap N_{i+1}|}{N_{i+1}} \leq \frac{|A \cap N_{i+1}| - |(A \setminus B) \cap [N_i, N_{i+1})|}{N_{i+1}} \leq \varepsilon + \frac{1}{i+1} - \frac{\delta}{2}$$

and therefore $d^-(B) \leq \varepsilon - \frac{\delta}{2}$. Contradiction. The moreover part of the lemma follows from the proof itself since we never imposed a cap on the growth of N_{i+1} . \square

Theorem 16. For any $0 < \varepsilon \leq 1$, forcing with $\mathbb{V}_{\geq \varepsilon}^-$ collapses the continuum to ω .

Proof. Let G be the generic filter and $g = \bigcup G$ is a total function $g : \omega \rightarrow 2$. Fix a maximal antichain $M \subseteq \mathbb{V}_{\geq \varepsilon}^-$ such that $d^-(\omega \setminus \text{dom}(f)) = \varepsilon$ for all $f \in M$. Suppose $M \cap G = \{f_0\}$. Now that $d^-(\omega \setminus \text{dom}(f_0)) = \varepsilon$, we take the sequence $\{N_i\}_{i \in \omega}$ satisfying Lemma 15. Also fix a bijection $\langle \cdot, \cdot \rangle : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$. Let I_i^j be the interval $[N_{\langle i, j \rangle}, N_{\langle i, j \rangle + 1})$.

We'd like to design a coding F with the help of g , such that $\{F(n)\}_{n \in \omega}$ enumerates $(2^\omega)^V$. For each interval I_i^j , we list the first $\lfloor \frac{\varepsilon \cdot |I_i^j|}{2^{i+1}} \rfloor$ elements of $I_i^j \setminus \text{dom}(f_0)$, and check the value of g at these slots. We say that I_i^j is positive, if there are more 1's in g than 0's among the slots we check; otherwise it's negative. Notice that, since $d^-(\omega \setminus \text{dom}(f_0)) = \varepsilon$, there will be enough elements in $I_i^j \setminus \text{dom}(f_0)$ for $k = \langle i, j \rangle$ large enough.

Fix $n \in \mathbb{N}$. We construct $F(n)$ together with a sequence $\{m_{i,n}\}_{i \in \omega}$. The idea is to use an interval $I_i^{m_{i,n}}$ to code $F(n)(i)$, then several intervals afterwards to code the position of the next

interval $I_{i+1}^{m_{i+1,n}}$. Start with $m_{0,n} = n$. Suppose $m_{i,n}$ is constructed. We let $F(n)(i) = 1$ if $I_i^{m_{i,n}}$ is positive, and $F(n)(i) = 0$ if $I_i^{m_{i,n}}$ is negative. Then let $m_{i+1,n}$ be the largest number $j \geq 0$ such that $I_i^{m_{i,n}+1}, \dots, I_i^{m_{i,n}+j}$ are all positive ($j = 0$ if the first interval $I_i^{m_{i,n}+1}$ is already negative). Such finite j exists by the usual genericity argument: for any $f \leq f_0$ and fixed i , by the above lemma, $\lim_{j \rightarrow \infty} \frac{|(\text{dom}(f) \setminus \text{dom}(f_0)) \cap I_i^j|}{|I_i^j|} = 0$ and hence f only decides whether I_i^j is positive/negative for finitely many j 's. Then we can extend $f' \leq f$ to ensure at least one of the intervals $I_i^{m_{i,n}+j}$ is negative, and therefore j is finite.

The above definition gives us a map $F : \omega \rightarrow 2^\omega$ in $V[G]$. Let \dot{F} be its name in V , and $\dot{m}_{i,n}$ be a name for each $m_{i,n}$. For every $A \in (2^\omega)^V$, define the set $D_A = \{f \in \mathbb{V}_{\geq \varepsilon}^- : \exists n \ f \Vdash \dot{F}(n) = A\}$. It suffices to prove D_A is dense below f_0 .

Fix $f \leq f_0$ and $A \in (2^\omega)^V$. By Lemma 15, for each $i \in \mathbb{N}$ we can find n_i large enough such that

$$|(\text{dom}(f) \setminus \text{dom}(f_0)) \cap I_i^j| < \frac{\varepsilon \cdot |I_i^j|}{2^{i+3}}$$

for all $j \geq n_i$. Now we take the extension $f' \leq f$ such that

$$f'(k) = \begin{cases} 1 & k \notin \text{dom}(f) \text{ and } k \text{ is among the first } \lfloor \frac{\varepsilon \cdot |I_i^{n_i}|}{2^{i+1}} \rfloor \text{ elements of } I_i^{n_i} \setminus \text{dom}(f_0) \text{ for } A(i) = 1 \\ 0 & k \notin \text{dom}(f) \text{ and } k \text{ is among the first } \lfloor \frac{\varepsilon \cdot |I_i^{n_i}|}{2^{i+1}} \rfloor \text{ elements of } I_i^{n_i} \setminus \text{dom}(f_0) \text{ for } A(i) = 0 \\ 1 & k \notin \text{dom}(f) \text{ and } k \text{ is among the first } \lfloor \frac{\varepsilon \cdot |I_i^{n_i+j}|}{2^{i+1}} \rfloor \text{ elements of } I_i^{n_i+j} \setminus \text{dom}(f_0) \text{ for } 1 \leq j \leq n_{i+1} \\ 0 & k \notin \text{dom}(f) \text{ and } k \text{ is among the first } \lfloor \frac{\varepsilon \cdot |I_i^{n_i+n_{i+1}+1}|}{2^{i+1}} \rfloor \text{ elements of } I_i^{n_i+n_{i+1}+1} \setminus \text{dom}(f_0) \end{cases}$$

and $f'(k)$ is not defined except for $k \in \text{dom}(f)$ or when k satisfies one of the above four conditions. Notice that for each i , we extended the domain on the intervals I_i^j for only finitely many j 's, and in each I_i^j the extended part has density at most $\frac{\varepsilon}{2^{i+1}}$. Thus $d(\text{dom}(f') \setminus \text{dom}(f)) = 0$. So f' is a condition.

Now we claim $f' \Vdash \dot{F}(n_0) = A$. This is because f' already has enough information to decide whether each relevant interval is positive/negative, since $(\text{dom}(f') \setminus \text{dom}(f)) \cap I_i^j$ has occupied at least $3/4$ of the slots that we check when deciding whether I_i^j is positive. Now by definition we have $m_{0,n_0} = n_0$, and the intervals $I_i^{n_i+1}, \dots, I_i^{n_i+n_{i+1}+1}$ indicate $m_{i+1,n_0} = n_{i+1}$. Therefore by induction $f' \Vdash \dot{m}_{i,n_0} = n_i$ for all i . Finally, the intervals $I_i^{n_i}$ ensure $F(n_0)(i) = A(i)$. Therefore $f' \Vdash \dot{F}(n_0)(i) = A(i)$. We conclude that D_A is dense below f_0 , and thus F enumerates $(2^\omega)^V$. \square

The last density forcing we investigate is $\mathbb{V}_{>0}^+$. Since $d^+(A) > 0$ is equivalent to saying A is not a density zero set, this forcing naturally splits into a two-step iteration.

Lemma 17. $\mathbb{V}_{>0}^+$ is forcing equivalent with the two-step iteration $\mathcal{P}(\omega)/\mathcal{Z}_0 * \mathbb{V}_{\text{ground}}(\dot{\mathcal{F}})$, where $\mathcal{Z}_0 = \{A \subset \omega : d(A) = 0\}$, $\dot{\mathcal{F}}$ is the name for $\mathcal{P}(\omega)/\mathcal{Z}_0$ -generic filter, and $\mathbb{V}_{\text{ground}}(\dot{\mathcal{F}})$ consists of partial functions $f : \omega \rightarrow 2$ in the ground model such that $\omega \setminus \text{dom}(f) \in \dot{\mathcal{F}}$.

Proof. Define the map

$$\begin{aligned} j : \mathbb{V}_{>0}^+ &\rightarrow \mathcal{P}(\omega)/\mathcal{Z}_0 * \mathbb{V}_{\text{ground}}(\dot{\mathcal{F}}) \\ f &\mapsto \langle [\omega \setminus \text{dom}(f)], \dot{f} \rangle \end{aligned}$$

It's clear that $f \leq g$ implies $j(f) \leq j(g)$. Suppose $f \perp g$. Then either $f(n) \neq g(n)$ for some n , or $d(\omega \setminus (\text{dom}(f) \cup \text{dom}(g))) = 0$. In both cases $j(f) \perp j(g)$. To show that j is a dense embedding, fix $\langle [A], \dot{f} \rangle \in \mathcal{P}(\omega)/\mathcal{Z}_0 * \mathbb{V}_{\text{ground}}(\dot{\mathcal{F}})$. So $[A] \Vdash \omega \setminus \text{dom}(\dot{f}) \in \dot{\mathcal{F}}$. Hence $A \subseteq \omega \setminus \text{dom}(f)$ modulo a density zero set. Let $f' = f \cup \{(n, 0) : n \notin A \wedge n \notin \text{dom}(f)\}$. Then $j(f') = \langle [A], \dot{f}' \rangle \leq \langle [A], \dot{f} \rangle$. \square

This intermediate model $V[\mathcal{F}]$ will help us show that $\mathbb{V}_{>0}^+$ collapses the continuum.

Theorem 18. Forcing with $\mathbb{V}_{>0}^+$ collapses the continuum to ω .

Proof. Let \mathcal{F} be the generic filter in $\mathcal{P}(\omega)/\mathcal{Z}_0$. In $V[\mathcal{F}]$, we inductively construct a partition $\mathbb{N} = \bigcup_{n \in \omega} A_n$ satisfying the following:

- Each A_n is in the ground model and $\omega \setminus \bigcup_{i=0}^n A_i \in \mathcal{F}$ for all n .
- For each $F \in \mathcal{F}$, there is n such that $A_n \cap F$ is infinite.

Start with $A_0 = \emptyset$. Clearly $\omega \setminus A_0 \in \mathcal{F}$. Suppose A_i is constructed for $i \leq n$. Enumerate the elements of $\omega \setminus \bigcup_{i=0}^n A_i$ as $b_n^0 < b_n^1 < b_n^2 < \dots$ and consider the disjoint union $\omega \setminus \bigcup_{i=0}^n A_i = B_n^1 \cup B_n^2$ where $B_n^1 = \{b_n^i : i \text{ is odd}\}$ and $B_n^2 = \{b_n^i : i \text{ is even}\}$. Both B_n^1 and B_n^2 are in the ground model, since by induction hypothesis A_i is in the ground model for $i \leq n$. Notice that \mathcal{F} is an ultrafilter over the ground model, and by induction hypothesis $\omega \setminus \bigcup_{i=0}^n A_i \in \mathcal{F}$; therefore exactly one of B_n^1, B_n^2 is in \mathcal{F} . Let A_{n+1} be the one *not* in \mathcal{F} . Then $\omega \setminus \bigcup_{i=0}^{n+1} A_i \in \mathcal{F}$.

It's clear that $\mathbb{N} = \bigcup_{n \in \omega} A_n$ and $d(A_n) = \frac{1}{2^n}$ for $n \geq 1$. Now suppose $A_n \cap F$ is finite for all n . Then

$$d^+(F) = d^+(\bigcup_{i \in \omega} (A_i \cap F)) = d^+(\bigcup_{i \geq n} (A_i \cap F)) \leq d^+(\bigcup_{i \geq n} A_i) = \frac{1}{2^{n-1}}$$

for all $n \geq 1$. Hence $d(F) = 0$ and thus $F \notin \mathcal{F}$. Finally, let $a_n^0 < a_n^1 < a_n^2 < \dots$ enumerate each A_n , and define $\iota_n(i) = a_n^i$. Each ι_n is also in the ground model.

In V , we construct a sufficiently surjective function $\sigma : 2^\omega \rightarrow 2^\omega$. That is, for every $x \in 2^\omega$ and for every partial function $f : \omega \rightarrow 2$ such that $\omega \setminus \text{dom}(f)$ is infinite, there is $y \in 2^\omega$ such that $f \subset y$ and $\sigma(y) = x$. This is done by transfinite induction: let $\{\langle x_\alpha, f_\alpha \rangle : \alpha < 2^{\aleph_0}\}$ enumerate all such pairs of x and f . At each stage, we choose $y_\alpha \supset f_\alpha$ which is not in the domain of σ yet and let $\sigma(y_\alpha) = x_\alpha$. This is doable because $|\omega \setminus \text{dom}(f_\alpha)| = \aleph_0$, and at the current stage, σ is only defined at $< 2^{\aleph_0}$ many points. Then σ is obtained by taking union at limit stages.

Let G be the generic filter in $\mathbb{V}_{\text{ground}}(\mathcal{F})$ and $g = \bigcup G$ be the generic real. In $V[\mathcal{F}][G]$, define

$$\begin{aligned} \eta : \omega &\rightarrow (2^\omega)^V \\ n &\mapsto \sigma(g \circ \iota_n) \end{aligned}$$

To see that this is well-defined, *i.e.* $g \circ \iota_n \in (2^\omega)^V$, notice that each A_n is in the ground model and $\omega \setminus A_n \in \mathcal{F}$. Hence, given any $f \in \mathbb{V}_{\text{ground}}(\mathcal{F})$ we can extend it to $f' \leq f$ such that $A_n \subseteq \text{dom}(f')$. Therefore $f' \Vdash g \circ \iota_n \in (2^\omega)^V$.

Now we claim that η is surjective. Fix $x \in (2^\omega)^V$ and take an arbitrary $f \in \mathbb{V}_{\text{ground}}(\mathcal{F})$. Then $A_n \cap (\omega \setminus \text{dom}(f))$ is infinite for some n , or equivalently, $\omega \setminus \text{dom}(f \circ \iota_n)$ is infinite. So there is $y \supset f \circ \iota_n$ such that $\sigma(y) = x$. Now let $f' = f \cup (y \circ \iota_n^{-1})$. Then $\text{dom}(f') = \text{dom}(f) \cup A_n$ and therefore it is a condition, and it follows from the definition of η that $f' \Vdash \dot{\eta}(n) = x$. \square

To conclude, we've completed the following table from [15]. The shaded boxes are the results proven in this paper. Note that if a forcing notion collapses the continuum to ω , then it fails to be proper and also adds both dominating and Cohen reals.

	$\mathbb{M}_{>0}^-$	$\mathbb{M}_{>0}^+$	$\mathbb{M}_{\geq \epsilon}^-$	$\mathbb{M}_{\geq \epsilon}^+$	$\mathbb{V}_{>0}^-$	$\mathbb{V}_{>0}^+$	$\mathbb{V}_{\geq \epsilon}^-$	$\mathbb{V}_{\geq \epsilon}^+$
proper	✗	✓	✓	✓	✗	✗	✗	✓
dominating	✓	✓	✓	✓	✓	✓	✓	✗
Cohen	✓	✓	✓	✓	✓	✓	✓	✗

Question 19. How about the corresponding forcings with subscript $> \epsilon$ for $\epsilon \neq 0$? Are they proper? If so, do they add dominating and Cohen reals?

Acknowledgements. Thanks to Dilip Raghavan for advice on almost everything in this paper. Thanks to Zhang Zhentao for early discussion on properties of upper and lower density. Thanks to Will Johnson for a counterexample on upper density which is not included in this paper; it saved the author from trying to prove a lemma that turns out to be incorrect.

References

- [1] Tomek Bartoszyński and Haim Judah. *Set theory*. A K Peters, Ltd., Wellesley, MA, 1995. On the structure of the real line.
- [2] Andreas Blass. Combinatorial cardinal characteristics of the continuum. In *Handbook of set theory*. Vols. 1, 2, 3, pages 395–489. Springer, Dordrecht, 2010.
- [3] Jörg Brendle, Osvaldo Guzmán, Michael Hrušák, and Dilip Raghavan. Combinatorial properties of MAD families. *arXiv e-prints*, pages 1–43 arXiv:2206.14936, June 2022.
- [4] Jörg Brendle and Dilip Raghavan. Bounding, splitting, and almost disjointness. *Ann. Pure Appl. Logic*, 165(2):631–651, 2014.
- [5] Jörg Brendle and Saharon Shelah. Ultrafilters on ω —their ideals and their cardinal characteristics. *Trans. Amer. Math. Soc.*, 351(7):2643–2674, 1999.
- [6] R. Michael Canjar. Mathias forcing which does not add dominating reals. *Proc. Amer. Math. Soc.*, 104(4):1239–1248, 1988.
- [7] David Chodounský and Osvaldo Guzmán. There are no P-points in Silver extensions. *Israel J. Math.*, 232(2):759–773, 2019.
- [8] I. Farah. Analytic quotients: theory of liftings for quotients over analytic ideals on the integers. *Mem. Amer. Math. Soc.*, 148(702):xvi+177, 2000.
- [9] Ilijas Farah. Analytic Hausdorff gaps. II. The density zero ideal. *Israel J. Math.*, 154:235–246, 2006.
- [10] Serge Grigorieff. Combinatorics on ideals and forcing. *Ann. Math. Logic*, 3(4):363–394, 1971.
- [11] Matthew Harrison-Trainor, Lu Liu, and Patrick Lutz. Coding information into all infinite subsets of a dense set, 2023. arXiv:2306.01226.

- [12] Fernando Hernández-Hernández and Michael Hrušák. Cardinal invariants of analytic P -ideals. *Canad. J. Math.*, 59(3):575–595, 2007.
- [13] Michael Hrušák and Hiroaki Minami. Mathias-Prikry and Laver-Prikry type forcing. *Ann. Pure Appl. Logic*, 165(3):880–894, 2014.
- [14] Claude Laflamme. Zapping small filters. *Proc. Amer. Math. Soc.*, 114(2):535–544, 1992.
- [15] G. Laguzzi, H. Mildenberger, and B. Stuber-Rousselle. Mathias and Silver forcing parametrized by density. *Arch. Math. Logic*, pages 00881–7, 26 pp., 2023.
- [16] Paolo Leonetti and Salvatore Tringali. Upper and lower densities have the strong Darboux property. *J. Number Theory*, 174:445–455, 2017.
- [17] Dilip Raghavan. The density zero ideal and the splitting number. *Ann. Pure Appl. Logic*, 171(7):102807, 15 pp., 2020.
- [18] Dilip Raghavan and Saharon Shelah. Two inequalities between cardinal invariants. *Fund. Math.*, 237(2):187–200, 2017.
- [19] S. Solecki. Analytic ideals and their applications. *Ann. Pure Appl. Logic*, 99(1-3):51–72, 1999.