

Generalized regular Turán numbers

Dániel Gerbner*, Hilal Hama Karim†

Abstract

We combine two generalizations of ordinary Turán problems. Given graphs H and F and a positive integer n , we study $\text{rex}(n, H, F)$, which is the largest number of copies of H in F -free regular n -vertex graphs.

1 Introduction

One of the fundamental theorems in extremal graph theory is due to Turán [18]. It states that among n -vertex K_{k+1} -free graphs, the most edges are contained in the complete k -partite graph with each part of order $\lfloor n/k \rfloor$ or $\lceil n/k \rceil$. This graph is called the *Turán graph* and is denoted by $T(n, k)$. More generally, given a graph F , we denote by $\text{ex}(n, F)$ the largest number of edges in an n -vertex F -free graph. This topic has attracted a lot of attention, see [8] for a survey.

A natural generalization is the study of the largest number of copies of another graph H instead of the number of edges in n -vertex F -free graphs. Let $\mathcal{N}(H, G)$ denote the number of not necessarily induced copies of H in G , and let $\text{ex}(n, H, F) = \max\{\mathcal{N}(H, G) : G \text{ is an } n\text{-vertex } F\text{-free graph}\}$. After several sporadic results, the systematic study of these so-called generalized Turán problems was initiated by Alon and Shikhelman [1].

Another natural generalization was recently considered in [11, 5, 4, 12, 17]. Here we study $\text{rex}(n, F)$, which is the largest number of edges in an n -vertex *regular* F -free graph.

In this paper we combine the above generalizations. Let $\text{rex}(n, H, F) := \max\{\mathcal{N}(H, G) : G \text{ is an } F\text{-free regular } n\text{-vertex graph}\}$. Our goal is to show some examples where $\text{rex}(n, H, F)$ behaves similarly to $\text{ex}(n, H, F)$ and also show some examples where they differ significantly.

Alon and Shikhelman [1] proved that $\text{ex}(n, H, F) = \Theta(n^{|V(H)|})$ if and only if F is not a subgraph of any blow-up of H . We extend this theorem to the regular setting.

Theorem 1.1. *For any graph F and H , we have that $\text{rex}(n, H, F) = \Theta(n^{|V(H)|})$ if and only if F is not a subgraph of a blow-up of H .*

*Alfréd Rényi Institute of Mathematics, Hun-Ren, E-mail: gerbner@renyi.hu.

†Department of Computer Science and Information Theory, Faculty of Electrical Engineering and Informatics, Budapest University of Technology and Economics, Műegyetem rkp. 3., H-1111 Budapest, Hungary. E-mail: hilal.hamakarim@edu.bme.hu

Another result of Alon and Shikhelman [1] is that $\text{ex}(n, K_3, F) = O(n)$ if and only if F is an extended friendship graph. In the extended friendship graph, every cycle is a triangle and there is a vertex v such that every pair of triangles intersect in v . We extend this theorem as well to our setting.

Theorem 1.2. $\text{rex}(n, K_3, F) = O(n)$ if and only if F is an extended friendship graph.

Let us turn to problems where adding the regularity changes the situation. It is well-known and easy to see that for any forest F , any graph with minimum degree at least $|V(F)|$ contains F . This implies that $\text{rex}(n, F) \leq (|V(F)|-1)n$. Let H be a connected graph, then the vertices of H have an ordering such that each but the first vertex has a neighbor that is earlier in the ordering. The copies of H in an F -free r -regular graph can be counted by picking the vertices in the above order. The first vertex can be picked n ways, and then each other vertex can be picked at most r ways among the neighbors of at least one of the vertices picked earlier. This shows that $\text{rex}(n, H, F) = O(n)$. On the other hand, $\text{ex}(n, P_\ell, P_k) = \Theta(n^{\lfloor \ell/2 \rfloor})$ by a theorem of Györi, Salia, Tompkins and Zamora [16].

Another example where the order of magnitude of $\text{rex}(n, H, F)$ is much smaller than that of $\text{ex}(n, H, F)$ is given by even cycles. When C_{2k} is forbidden, the regularity does not have to be constant, but it is $O(n^{1/k})$ by a theorem of Bondy and Simonovits [3]. Therefore, $\text{rex}(n, C_\ell, C_{2k}) = O(n^{1+\frac{\ell-1}{k}})$, while we have $\text{ex}(n, C_\ell, C_{2k}) = \Theta(n^{\lfloor \ell/2 \rfloor})$ if $3 \leq \ell \neq 2k$ [13].

Note that we have $\text{ex}(n, C_\ell, C_{2k+1}) = \Theta(n^\ell)$ if ℓ is even or $\ell > 2k + 1$, as shown by the blow-up of C_ℓ . Interestingly, in the remaining case $3 < \ell < 2k + 1$ is odd, we have $\text{ex}(n, C_\ell, C_{2k+1}) = \Theta(n^{\lfloor \ell/2 \rfloor})$ [13], while the above argument does not give any non-trivial bound. It is a natural question to ask whether $\text{rex}(n, C_\ell, C_{2k+1})$ is significantly smaller in this case. We can answer this question in the negative.

Proposition 1.3. *If $3 < \ell < 2k + 1$ is odd, then $\text{rex}(n, C_\ell, C_{2k+1}) = \Theta(n^{\lfloor \ell/2 \rfloor})$.*

So far we considered only the order of magnitude of $\text{rex}(n, H, F)$. Let us turn to exact and asymptotic results. As shown in [5, 4], for $k \geq 3$ we have $\text{rex}(n, K_{k+1}) = (1 + o(1))|E(T(n, k))|$ (obviously we have $\text{rex}(n, K_{k+1}) = |E(T(n, k))|$ if k divides n). The exact value of $\text{rex}(n, K_{k+1})$ was determined for all sufficiently large n in [12]. Let $T^*(n, k)$ denote an arbitrary n -vertex K_{k+1} -free regular graph with $\text{rex}(n, K_{k+1})$ edges. Forbidding K_3 is very different from forbidding larger cliques in the regular Turán problem. If n is even, then $T(n, 2)$ is the regular n -vertex triangle-free graph with the most edges. If n is odd, then a regular n -vertex triangle-free graph with the most edges is obtained by deleting some edges of an n -vertex blow-up of C_5 , as shown in [5, 4].

Given H with $\chi(H) \leq k$, there has been a lot of research on whether $\text{ex}(n, H, K_{k+1}) = \mathcal{N}(H, T(n, k))$ for sufficiently large n , see e.g. [15, 10, 9]. There have been two types of counterexamples found (where even $\text{ex}(n, H, K_{k+1}) = (1 + o(1))\mathcal{N}(H, T(n, k))$ does not hold). If H is a very unbalanced bipartite graph, then an unbalanced complete k -partite graph may contain more copies of H than the Turán graph. For some graphs H , there are n -vertex K_{k+1} -free graphs that contain more copies of H than any n -vertex complete k -partite graph, see [14]. In each of the known constructions, most of the vertices of H would belong

to two different classes of k -partite graphs, but they can belong to the same class of the blow-up of another graph. Then that class has many vertices.

Both counterexamples are very far from being regular. This suggests that maybe there are no regular counterexamples at all.

Conjecture 1.4. *Let $\chi(H) \leq k \geq 3$. Then $\text{rex}(n, H, K_{k+1}) = (1 + o(1))\mathcal{N}(H, T(n, k))$. Moreover, if n is sufficiently large and is divisible by k , then $\text{rex}(n, H, K_{k+1}) = \mathcal{N}(H, T(n, k))$.*

Proposition 1.5. *Conjecture 1.4 holds for complete k -partite graphs H .*

Proposition 1.6. *Let H be a bipartite graph. If n is even and sufficiently large, then $\text{rex}(n, H, K_3) = \mathcal{N}(H, T(n, 2))$. If H is a tree and n is odd and sufficiently large, then $\text{rex}(n, H, K_3) = \mathcal{N}(H, G^*)$, where G^* is a regular graph obtained by deleting some edges of an n -vertex blow-up of C_5 .*

Finally, we determine the exact value for $\text{rex}(n, K_3, P_k)$, when n is large enough and P_k is a path on k vertices, with $k \geq 7$. To ease the notation and describe the extremal graphs, we define some graphs first. Let \mathcal{G}_{k-1} denote the graphs obtained from K_{k-1} by removing the edges of a triangle-free 2-regular subgraph, i.e., the union of vertex-disjoint cycles of length more than 3 such that the total length of the cycles is $k-1$. In the case k is even, let $G_{k-2} := K_{k-2} - M$, a clique on $k-2$ vertices in which a perfect matching is removed. Note that each of the above graphs is $(k-4)$ -regular and P_k -free. If k is odd, let $G'_{k-1} := K_{k-1} - M$, a clique on $k-1$ vertices in which a perfect matching is removed. It is easy to see that $\mathcal{N}(K_3, G_{k-1}) = 8\binom{k/2-1}{3} + 3 - k/2$ for any graph $G_{k-1} \in \mathcal{G}_{k-1}$, $\mathcal{N}(K_3, G_{k-2}) = 8\binom{k/2-1}{3}$ and $\mathcal{N}(K_3, G'_{k-1}) = (k-1)(k-3)(k-5)/6 = 8\binom{(k-1)/2}{3}$. We denote by $H + F$ the disjoint union of two graphs H and F , and by mF we mean m disjoint copies of the graph F .

Theorem 1.7. *Let P_k be a path on k vertices and n be large enough. Then:*

1. *If $(k-1)|n$, then $\text{rex}(n, K_3, P) = \frac{n}{k-1} \binom{k-1}{3}$, and the unique extremal graph is $\frac{n}{k-1}K_{k-1}$.*
2. *Assume that $(k-1) \nmid n$, $k \geq 7$ and either $k-2$ divides n or k is odd. Let $n = a(k-2) + b$ with $b < k-2$. Then we have $\text{rex}(n, K_3, P_k) = (a-b) \binom{k-2}{3} + 8b \binom{k-1}{3}$, and the unique extremal graph is $(a-b)(K_{k-2}) + bG'_{k-1}$.*
3. *If $k \geq 7$ is even, and n is neither divisible by $k-1$ nor by $k-2$. Let $n = a(k-3) + b$, with $b < k-3$. Then*

$$\text{rex}(n, K_3, P_k) = (a - \ell - \lfloor b/2 \rfloor) \binom{k-3}{3} + \ell \mathcal{N}(K_3, G_{k-2}) + \lfloor b/2 \rfloor \mathcal{N}(K_3, G_{k-1}),$$

and the extremal graphs are formed by adding $\lfloor b/2 \rfloor$ graphs from \mathcal{G}_{k-1} to $(a - \ell - \lfloor b/2 \rfloor)K_{k-3} + \ell G_{k-2}$, where $\ell = 0$ if b is even and $\ell = 1$ otherwise.

2 Tools

We will use the following well-known theorem of Erdős and Sachs [7].

Theorem 2.1 ([7]). *For every r and g , there exists an r -regular graph of girth at least g .*

We will use the following simple corollaries of the above theorem.

Lemma 2.2. *(i) For any r and k , if n is sufficiently large and nr is even, then there is an n -vertex r -regular graph with girth at least k .*

(ii) For any r , k and i , if n is sufficiently large and $nr - i$ is even, then there is an n -vertex graph with girth at least k that contains i vertices of degree $r - 1$ and each other vertex has degree r . Moreover, we can have that the vertices of degree $r - 1$ are at distance at least $k - 1$.

Proof. Let us start by proving **(i)**. We know such a graph G_1 exists on m vertices for some m . If r is even, we take $r/2$ vertex-disjoint copies of G_1 and remove an edge from each. We add a new vertex and connect it to the endpoints of the removed edges. The resulting graph G_2 satisfies the desired properties on $\frac{rm}{2} + 1$ vertices. For each $n \geq rm^2$, we can write n as $a(\frac{rm}{2} + 1) + bm$, thus we can create an n -vertex graph by taking vertex-disjoint copies of G_1 and G_2 .

If r is odd, we take r vertex-disjoint copies of G_1 and remove an edge from each. We add two new vertices u, v and connect u to one of the endpoints of each removed edge and v to the other endpoint. The resulting graph G'_2 satisfies the desired properties on $rm + 2$ vertices. For each even $n \geq r^2m$, we can write n as $a(rm + 2) + bm$, thus we can create an n -vertex graph by taking vertex-disjoint copies of G_1 and G'_2 .

We continue with the proof of **(ii)**. If i is even, we take a graph guaranteed by **(i)** and remove $i/2$ independent edges such that the endpoints of these edges are at distance at least $k - 1$. If n is sufficiently large, we can greedily find such edges. Indeed, we take an edge u_1v_1 , then at most $2r - 2$ other vertices are adjacent to u or v , and at most $2(r - 1)^j$ vertices are at distance j from u or v . Altogether there are at most $2(r - 1)^k$ vertices at distance at most $k - 1$ from u or v . We take a vertex u_2 different from those at most $2(r - 1)^k$ vertices and an arbitrary neighbor v_2 of u_2 . Repeating this, we can find $i/2$ edges if we can pick a vertex $u_{i/2}$ that is not among the $i - 2$ vertices picked earlier and the at most $(i - 2)(r - 1)^k$ vertices at distance at most $k - 1$ from the vertices picked earlier. In other words, we can pick the desired edges if $n > i - 2 + (i - 2)(r - 1)^k$. Note that the distance of u_i and v_i is at least $k - 1$ after removing the edge u_iv_i because of the girth condition.

If i is odd, observe that both n and r are odd. Let G_3 be an $(r - 1)$ -regular m -vertex graph for some odd m . Let G_4 be an r -regular graph on m' vertices for some m' sufficiently large. We take $(m - i)/2$ copies of G_4 and remove an edge from each. This way we obtain $m - i$ vertices of degree $r - 1$, we connect each of them to a different vertex of G_3 . The resulting graph has exactly i vertices of degree $r - 1$ and each other vertex has degree r .

In each of the above constructions, we removed an edge uv from some copy of a graph of girth at least k , then we added some edges incident to u and v and outside vertices. After

removing uv , the distance of u and v becomes at least $k - 1$, thus this way we do not create cycles of length less than k . ■

Corollary 2.3. *For any sequence (a_n) of positive integers with $a_n = \omega(1)$, we can take n vertex graphs G_n such that they satisfy the assumptions of Lemma 2.2 with $r \leq a_n$ and $r = \omega(1)$.*

We will use the following simple corollary of Hall's theorem.

Observation 2.4. *For every $k \leq n$, we have a k -regular bipartite graph with both parts of order n .*

We will use a theorem of Andrásfai, Erdős, and Sós [2], stating that a non-bipartite triangle-free graph on n vertices contains a vertex of degree at most $2n/5$.

We will use a stability result on $\text{rex}(2n + 1, K_3)$, which may be interesting on its own. When we talk about V_{i+j} in the statement or the proof, then $+$ is meant modulo 5.

Lemma 2.5. *Let G be a d -regular n -vertex triangle-free graph with n odd. Let $d \geq 2n/5 - o(n)$. Then $V(G)$ contains disjoint sets V_1, \dots, V_5 such that $|V_i| = n/5 - o(n)$ and from V_i there is no edge to V_i, V_{i+2} and V_{i+3} , and $n/5 - o(n)$ edges go to V_{i+1} and V_{i+4} . In particular G is obtained by deleting some edges of an n -vertex blow-up of C_5 .*

Proof. Observe that G cannot be bipartite, thus $d \leq 2n/5$ by the result of Andrásfai, Erdős, and Sós [2]. Let C_{2k+1} be a shortest odd cycle in G and C be a copy of C_{2k+1} . Then every vertex outside C is adjacent to at most two vertices of C . This implies that there are at most $2(n - 2k - 1)$ edges between C and the other vertices. On the other hand, there are at least $(2k + 1)d - (2k + 1) \geq 2(2k + 1)n/5 - o(n)$ edges between C and the other vertices by our assumption on the degrees of the vertices of C (which is d). Here we use that there are $2k + 1$ edges inside C , since it is the shortest odd cycle and so does not have any chords.

This shows that $k \leq 2$. Since G is triangle-free, we have $k = 2$. Furthermore, $n - o(n)$ vertices outside C have two neighbors in C , otherwise there are at most $2(n - 2k - 1) - \Omega(n) < (2k + 1)d - (2k + 1)$ edges between C and the other vertices. Let v_1, \dots, v_5 be the vertices of C in the cyclic order. Observe that no vertex can be adjacent to both v_i and v_{i+1} , thus $n - o(n)$ vertices are each, for some i , adjacent to v_i and v_{i+2} . We place those vertices to V_{i+1} . Let $U = V(G) \setminus (V_1 \cup V_2 \cup V_3 \cup V_4 \cup V_5)$, then $|U| = o(n)$.

Let $u \in V_i$. As u has a common neighbor with every vertex of V_i, V_{i+2} and V_{i+3} , there are no neighbors of u in $V_i \cup V_{i+2} \cup V_{i+3}$, thus all the neighbors of u are in V_{i+1} and V_{i+4} and U . In particular, $|V_{i+1}| + |V_{i+4}| \geq 2n/5 - o(n)$. This holds for every non-adjacent pair of classes. If $|V_i| \leq n/5 - \alpha n$, then $|V_{i+2}|, |V_{i+3}| \geq n/5 + \alpha n - o(n)$. Then $|V_{i+1}| + |V_{i+4}| \leq 2n/5 - \alpha n - o(n)$, thus $\alpha = o(1)$, completing the proof. ■

3 Proofs

Let us prove Theorem 1.1. Recall that it states that $\text{rex}(n, H, F) = \Theta(n^{|V(H)|})$ if and only if F is not a subgraph of a blow-up of H .

Proof of Theorem 1.1. If F is a subgraph of a blow-up of H , then obviously $rex(n, H, F) \leq \text{ex}(n, H, F) = o(n^{|V(H)|})$, where we use the result of Alon and Shikhelman [1] mentioned in the introduction. Assume now that F is not a subgraph of any blow-up of H . If H is the empty graph, the statement follows. Observe that otherwise we also have that F has chromatic number at least 3. We can also assume that there are no isolated vertices in F .

In the analogous statement for $\text{ex}(n, H, F)$, this is the trivial direction, as the blow-up $H(m)$ with $m = \lfloor n/|V(H)| \rfloor$ is F -free and contains $\Omega(n^{|V(H)|})$ copies of H . However, we have two problems here: the first is that $H(m)$ is not regular if H is not regular, and the second is that we may need to add some vertices of degree 0 to obtain an n -vertex graph.

Let Δ be the largest degree in H . Let H_i be a graph with girth more than $3|V(F)|$ that has a set S of i vertices of degree 2Δ and all the other vertices of degree $2\Delta + 1$, such that the vertices of S are of distance at least $|V(F)|$. Such a graph exists by Lemma 2.2 where the number of vertices is large enough compared to Δ and $|V(F)|$, but constant compared to n .

For each i and each vertex v of H of degree i , we take a copy of $H_{2\Delta+1-i}$ and join v to the $2\Delta + 1 - i$ vertices of this copy of degree 2Δ . This way we obtain a $(2\Delta + 1)$ -regular graph H' on constant many vertices.

Claim 3.1. *The blow-up $H'(m)$ is F -free for any m .*

Proof of Claim. We can assume that m is large enough. Let us assume that there is a copy of F in $H'(m)$, that we will denote with F^* . Let H^* denote an arbitrary copy of an H_i for some i in H' . Let F_0 denote a connected component of the intersection of F^* with the blow-up of H^* . Observe that F_0 is bipartite and intersects the blow-up of at most one vertex v of S . Indeed, otherwise F_0 would contain an odd cycle of length more than $|V(F)|$ or two vertices of distance more than $|V(F)|$. Let u be the vertex of H that is joined to v in H' and u' be an arbitrary neighbor of u in H .

Now we can delete F_0 and embed it to the complete bipartite graph between the blow-ups of u and u' , using only vertices that were not in F^* . We repeat this for every subgraph of F outside $H(m)$. At the end, we obtain a copy of F in $H(m)$, a contradiction. ■

Let us return to the proof of the theorem. We are done if $|V(H')|$ divides n , as we can pick m to be $n/|V(H')|$. To prove the theorem for every n , we do the following. Let H'' denote the vertex-disjoint union of $H'(2m)$ and $C_{2|V(F)|+1}((2\Delta + 1)m)$. Note that H'' is F -free, since every subgraph of $C_{2|V(F)|+1}((2\Delta + 1)m)$ on at most $|V(F)|$ vertices is bipartite. If there is a copy of F in H'' , then the components that are in $C_{2|V(F)|+1}((2\Delta + 1)m)$ could be easily replaced by copies in $H'(2m)$, to find a copy of F in $H'(2m)$, a contradiction.

Clearly, H'' is $2(2\Delta + 1)m$ -regular for any m , and the number of vertices have the same parity as m . Let us pick the largest m such that $n - |V(H'')|$ is even. Observe that $n - |V(H'')|$ is a constant. Now we modify the $C_{2|V(F)|+1}((2\Delta + 1)m)$ subgraph. Note that this is similar to the way the odd cycles were modified in [5].

Let $A_1, \dots, A_{2|V(F)|+1}$ be the blown-up parts of the cycle in this order. We take a pair of neighboring parts, say A_i and A_{i+1} , and add $b = (n - |V(H'')|)/2$ vertices to each of A_i and A_{i+1} . We add them in such a way that we still have a complete bipartite graph between any

pair of consecutive blown-up parts A_j, A_{j+1} , i.e., we connect the new vertices of A_i to each vertex of A_{i-1} and A_{i+1} , and connect the new vertices of A_{i+1} to each vertex of A_i and A_{i+2} . Then we remove the edges of a spanning bipartite graph B between A_{i-1} and A_i such that each vertex of A_{i-1} has degree b and each vertex of A_i has degree $\lfloor \frac{(2\Delta+1)mb}{(2\Delta+1)m+b} \rfloor$ or $\lceil \frac{(2\Delta+1)mb}{(2\Delta+1)m+b} \rceil$ in B . We remove the edges of a copy of B between A_{i+1} and A_{i+2} as well such that the vertices of degree b are in A_{i+2} .

At this point the vertices outside A_i and A_{i+1} have degree $2(2\Delta + 1)m$. The part A_i consists of a set A'_i of vertices of degree $2(2\Delta + 1)m + b - \lfloor \frac{(2\Delta+1)mb}{(2\Delta+1)m+b} \rfloor$ and a set A''_i of vertices of degree $2(2\Delta + 1)m + b - \lceil \frac{(2\Delta+1)mb}{(2\Delta+1)m+b} \rceil$. Similarly, A_{i+1} consists of a set A'_{i+1} of vertices of degree $2(2\Delta + 1)m + b - \lfloor \frac{(2\Delta+1)mb}{(2\Delta+1)m+b} \rfloor$ and a set A''_{i+1} of vertices of degree $2(2\Delta + 1)m + b - \lceil \frac{(2\Delta+1)mb}{(2\Delta+1)m+b} \rceil$. Observe that by the analogous construction, we have that $|A'_i| = |A'_{i+1}|$. We pick a perfect matching M' between A'_i and A'_{i+1} , and extend it to a perfect matching M between A_i and A_{i+1} . We delete the edges of M .

Then the resulting graph between A_i and A_{i+1} is $2(2\Delta + 1)m + b - 1$ -regular, thus we can delete matchings between A_i and A_{i+1} till we obtain a $\left((2\Delta + 1)m + \lfloor \frac{(2\Delta+1)mb}{(2\Delta+1)m+b} \rfloor \right)$ -regular graph between A_i and A_{i+1} . After that, we add the edges of M that are not in M' . Observe that vertices of A'_i have $\left((2\Delta + 1)m + \lfloor \frac{(2\Delta+1)mb}{(2\Delta+1)m+b} \rfloor \right)$ neighbors in A_{i+1} and $(2\Delta + 1)m - \lfloor \frac{(2\Delta+1)mb}{(2\Delta+1)m+b} \rfloor$ neighbors in A_{i-1} . Vertices of A''_i have one more neighbor in A_{i+1} and one less neighbor in A_{i-1} . The same holds for vertices in A_{i+1} . Let G denote the resulting n -vertex graph. Then G is $2(2\Delta + 1)m$ -regular and contains at least $m^{|V(H)|} = \Theta(n^{|V(H)|})$ copies of H , completing the proof. \blacksquare

Let us continue with the proof of Theorem 1.2. Recall that it states that $\text{rex}(n, K_3, F) = O(n)$ if and only if F is an extended friendship graph.

Proof of Theorem 1.2. If F is an extended friendship graph, then $\text{rex}(n, K_3, F) \leq \text{ex}(n, K_3, F) = O(n)$.

Assume that F is not an extended friendship graph. Then it either contains $2K_3$ or a longer cycle C_k with $k \geq 4$. In the first case, we take the K_3 -free graph on $n - 1$ vertices with regularity $r = \Omega(n)$ due to Caro and Tuza [5]. In particular, it contains an induced copy K of $K_{r/2, r/2}$. We remove a matching from K , and add a new vertex v , connected to the vertices of K . The resulting graph is r -regular, and contains $r^2/4$ triangles that all contain v , completing our proof.

Let us assume now that F contains C_k . Let n be sufficiently large. We pick $r = \omega(1)$ such that r is small enough to have a $2r$ -regular graph of girth more than $3k^2$ on m vertices whenever $m \geq 11n/36$. We take an r -regular m -vertex graph G_0 of girth at least $3k^2$ where $m = \lfloor n/4 \rfloor$. We consider G_0^k as an auxiliary graph. Recall that the k th power G_0^k of a graph G_0 is obtained by joining vertices of distance at most k .

It is easy to see that G_0^k is r' -reg, where $r' = r + r(r - 1) + \dots + r(r - 1)^{k-1}$. We take a proper $r' + 1$ -edge-coloring of G_0^k . Since G_0 is a subgraph of G_0^k , we obtain a proper edge-coloring of G_0 . For each color i , we partition the edges of color i to some number of r -sets

and a set of order at most r . For each such set, we add a new vertex and connect it to the at most $2r$ vertices that are incident to those at most r edges. This way we obtain G_1 .

The vertices of G_0 have degree r in G_0 , thus they are incident to edges of r colors, hence their degree is $2r$ in G_1 . The newly added vertices have degree $2r$, except $r' + 1$ vertices, that are connected to the endpoints of less than r edges. Let us assume that the sum of degrees in G_1 is $2r|V(G_1)| - \ell$. Note that ℓ is even since each vertex has an even degree. There are rm edges from $V(G_0)$ to $V(G_1) \setminus V(G_0)$ and at least $2r(|V(G_1) \setminus V(G_0)| - r' - 1)$ edges from $V(G_1) \setminus V(G_0)$ to $V(G_0)$, thus $|V(G_1) \setminus V(G_0)| \leq \frac{m}{2} + r' + 1$, hence $V(G_1) \leq 3n/8 + O(1) \leq 4n/9$.

Now we make G_1 regular. We take another copy of G_0 and remove $\ell/2$ edges. The resulting graph Q_1 is of girth more than k with a set S of ℓ vertices of degree $2r - 1$ and all the other vertices of degree r , such that the vertices of S are at distance at least $|V(F)|$. We join each vertex v of G_1 to $2r - d(v)$ vertices of degree $2r - 1$ in this new graph. The resulting graph G_2 is $2r$ -regular on at most $25n/36$ vertices.

Finally, we add a $2r$ -regular graph of girth more than k on $n - |V(G_2)|$ vertices. This exists by the choice of r . ■

Let us continue with the proof of Proposition 1.3. Recall that it states that if $3 < \ell < 2k + 1$ is odd, then $\text{rex}(n, C_\ell, C_{2k+1}) = \Theta(n^{\lfloor \ell/2 \rfloor})$.

Proof of Proposition 1.3. The upper bound is shown by $\text{ex}(n, C_\ell, C_{2k+1}) = \Theta(n^{\lfloor \ell/2 \rfloor})$ in [13].

Let us turn to the lower bound. We start with an unbalanced blow-up of C_ℓ , where we blow up $(\ell - 1)/2$ independent vertices to m -sets, and keep the other vertices (note that this construction shows the analogous bound for $\text{ex}(n, C_\ell, C_{2k+1})$, but it is far from regular). Let H denote this graph, then the largest degree is $2m$ in H . We add sets A_1, A_2, \dots, A_{2k} of new vertices of order $2m - 2$. We take two blown-up parts A and A' of order m of H . We take all the possible edges between A and A_1 , then an m -regular graph between A_i and A_{i+1} for each $i \leq 2k - 1$ (this exists because of Observation 2.4), and then take all the possible edges between A_{2k} and A' . It is easy to see that each vertex of A, A' and each A_i has degree $2m$ and no C_{2k+1} is created this way. We repeat this by taking $4k(m - 1)$ new vertices as long as there are at least two blown up classes of order m .

We are left with several vertices of degree $2m$, at most one blown up class with m vertices of degree 2 and exactly two adjacent vertices u, v of degree $m + 1$. In the case there are only the vertices u, v of degree less than $2m$, we take sets B, B' of order $m - 1$ and B_1, \dots, B_{2k+1} of order $2m - 1$. We take all the edges between u and B and between B and B_1 . Then we take an $(m + 1)$ -regular graph between B_{2i+1} and B_{2i+2} , and an $(m - 1)$ -regular graph between B_{2i} and B_{2i+1} . Finally, we take all the edges between B_{2k+1} and B' and between B' and v .

In the case we have a blown-up class A left, we take sets A_1, A_2, \dots, A_{2k} as above. Then we add all the edges between A_{2k} and a new m -set C . We connect each vertex of C to u and a new vertex u' . Then we take sets B, B', B_i as above and add edges as above, with u replaced by u' . It is easy to see that in the resulting graph H' each vertex has degree $2m$ and there is no C_{2k+1} . Each time we added a bipartite graph, thus there is no odd cycle without

the edges of H . However, we made sure that vertices of H are at distance at least $2k$ (and adjacent vertices of H are at distance more than $2k$) in the newly added parts of the graph.

It is left to add $n - |V(H')|$ vertices without ruining these properties. Observe that we added at most $4\ell km$ vertices to H . We pick m to be the largest odd number below $\lfloor n/10k\ell \rfloor$, thus H' has at most $n/2$ vertices. If $n - |V(H')|$ is even, we can pick a bipartite $2m$ -regular graph on those vertices, completing the proof. If $n - |V(H')|$ is odd, we additionally pick a copy of $C_{(2k+3)m}$, and then pick a bipartite $2m$ -regular graph on the remaining vertices, completing the proof. \blacksquare

Let us continue with the proof of Proposition 1.5. Recall that it states that if H is a complete k -partite graph and $k \geq 3$, then $\text{rex}(n, H, K_{k+1}) = (1+o(1))\mathcal{N}(H, T(n, k))$, without the error term if k divides n .

Proof. Let $H = K_{s_1, \dots, s_k}$ with $s_1 \leq s_2 \leq \dots \leq s_k$. For simplicity, we will deal with labeled copies of H , clearly the same n -vertex K_{k+1} -free regular graph maximizes (asymptotically) the number of labeled copies of H as the number of copies. We will show that for $k \geq 2$, any n -vertex K_{k+1} -free graph contains at most $k! \prod_{i=1}^k \binom{n/k}{s_i}$ copies of H . Clearly $T(n, k)$ satisfies this with equality if k divides n , and $T^*(n, k)$ gives the correct asymptotics, thus this upper bound completes the proof for other values of n as well.

We apply induction on k and on $\sum_{i=1}^k s_i$. The base case $k = 2$ follows from Proposition 1.6 (note that the case n is even is equivalent to the even case of Proposition 1.6, while the case of odd n is much weaker. One could prove this weaker statement by following the proof of the even case of Proposition 1.6 word by word). The other base case $\sum_{i=1}^k s_i = k$ follows from Zykov's theorem [19], which states that $\text{ex}(n, K_r, K_{k+1}) = \mathcal{N}(H, T(n, k))$.

Let G be an n -vertex K_{k+1} -free r -regular graph, then $r \leq (k-1)n/k$ by Turán's theorem. We consider two cases. Assume first that $s_1 = 1$ and let H' be the graph we obtain by deleting the first class from H . Then we first pick a vertex v of G corresponding to the single vertex in the first class, at most n ways. Then we pick a labeled copy of H' in the neighborhood of v , at most $(k-1)! \prod_{i=2}^k \binom{r/(k-1)}{s_i}$ by the induction on k . This way we picked the labeled copies of H at most $n(k-1)! \prod_{i=2}^k \binom{r/(k-1)}{s_i} \leq nk! / k \prod_{i=2}^k \binom{(k-1)n/k}{s_i} = k! \frac{n}{k} \prod_{i=2}^k \binom{n/k}{s_i} = k! \binom{n/k}{s_1} \prod_{i=2}^k \binom{n/k}{s_i}$.

Assume now that $s_1 > 1$ and let H'' be the graph we obtain by deleting a K_k from H . We first pick an unlabeled copy K of K_k , then a labeled copy of H'' from the remaining vertices, and then add the labels to the vertices of K . By Zykov's theorem, the number of unlabeled copies of K_k is maximized by the Turán graph, thus it is at most n^k/k^k . The number of labeled copies of H'' is at most $k! \prod_{i=1}^k \binom{(n-k)/k}{s_i-1}$ by induction on $\sum_{i=1}^k s_i$. Afterwards, we add the vertices of K to the vertices of H'' . Observe that each vertex of K has a copy of K_{k-1} in their neighborhood in H'' . As G is K_{k+1} -free, the vertices of a K_{k-1} cannot be adjacent to two adjacent vertices. This implies that each copy of K_{k-1} has at most one common neighbor in K . Therefore, each vertex of K can belong to at most one of the classes of H . This means that the number of labels the vertices of K can receive is at most $\prod_{i=1}^k s_i$, hence the number of labeled copies of H is at most $\frac{n^k}{k^k} k! \prod_{i=1}^k s_i \prod_{i=1}^k \binom{(n-k)/k}{s_i-1} = k! \prod_{i=1}^k \binom{n/k}{s_i}$. This completes the proof. \blacksquare

Let us continue with Proposition 1.6. Recall that it extends the above proposition to the case $k = 2$ if n is even. If n is odd, then it deals with the case H is a tree and claims that the extremal graph is obtained by deleting some edges of a blowup of C_5 .

Proof of Proposition 1.6. Let n be even, consider a component H_0 of H and an ordering of the vertices of H_0 such that each but the first vertex has an earlier neighbor. Such an ordering obviously exists by first picking an arbitrary vertex and then each time picking a neighbor of a vertex already picked.

Let G be an n -vertex r -regular triangle-free graph, then $r \leq n/2$. Moreover, either G is bipartite (with both parts of order $n/2$, thus G is contained in $T(n, 2)$, completing the proof), or $r \leq 2n/5$. Assume that $r \leq 2n/5$. There are at most n ways to pick the first vertex and at most $2n/5$ ways to pick each subsequent vertices. In $T(n, 2)$ there are n and $(1 + o(1))n/2$ ways to do this. The copies of H_0 may be counted multiple times, but the number of times is a fixed constant c depending only on the automorphisms of H and not the host graph. Therefore, G contains at most $2 \left(\frac{2n}{5}\right)^{|V(H_0)|} / c$, while $T(n, 2)$ contains $(2 + o(1)) \left(\frac{n}{2}\right)^{|V(H_0)|} / c$ copies of H_0 . Then we pick the other components of H . Similarly, there are more ways to pick each component in $T(n, 2)$ than in G if n is large enough. Therefore, $T(n, 2)$ contains more copies of H for sufficiently large n , completing the proof.

Assume now that n is odd and H is a forest. Let G be an n -vertex r -regular triangle-free graph, then $r \leq 2n/5$. Moreover, by Lemma 2.5, either G is obtained by deleting some edges of an n -vertex blow-up of C_5 , or $r \leq 2n/5 - \varepsilon n$ for some $\varepsilon > 0$. In the first case, we are done. In the second case, we can proceed similarly to the argument in the case where n is even. G contains at most $2 \left(\frac{2n}{5} - \varepsilon n\right)^{|V(H_0)|} / c$ copies of H_0 , while G^* contains $2 \left(\frac{2n}{5}\right)^{|V(H_0)|} / c$ copies of H_0 . The same holds for other components, thus G^* contains more copies of H for sufficiently large n , completing the proof in this case. \blacksquare

Before the proof of Theorem 1.7, let us mention some results that will be used. By a theorem of Erdős and Gallai [6], if a connected graph has at least k vertices and minimum degree $\lfloor k/2 \rfloor$, then it contains a P_k . Gerbner, Patkós, Tuza and Vizer [12] gave the exact value of $\text{regex}(n, T)$ for any tree T and large n . Note that $\text{regex}(n, F) = \max\{d : \text{there is an } n\text{-vertex, } d\text{-regular, } F\text{-free graph } G\}$

Theorem 3.2 ([12]). *Let T be a tree on t vertices and $n > n_0(T)$. Then*

$$\text{regex}(n, T) = \begin{cases} t - 2 & \text{if } t - 1 \text{ divides } n \text{ or } T \text{ is a star and } t \text{ or } n \text{ is even,} \\ t - 3 & \text{if the above does not hold, and either } t \text{ is odd, or } t - 2 \text{ divides } n, \\ & \text{or } T \text{ is a star or } T \text{ is an almost-star and } n \text{ is even.} \\ t - 4 & \text{otherwise.} \end{cases}$$

A tree is an *almost-star* if in its proper 2-coloring, one of the classes consists of at most two vertices (thus a path on at least 7 vertices is not an almost star).

Now we are ready to present the proof of Theorem 1.7, which determines $\text{rex}(n, K_3, P_k)$ if $k \geq 7$ and n is sufficiently large.

Proof of Theorem 1.7. Let G be an n -vertex r -regular P_k -free graph containing the maximum number of triangles. Observe that each vertex of G is in at most $\binom{r}{2}$ triangles, and hence $\mathcal{N}(K_3, G) \leq \frac{n}{3} \binom{r}{2}$, with equality only when each vertex is in a clique K_{r+1} . Note that in each of the constructions described in the introduction, the r -regular n -vertex graph contains $n/(r+1) - O(1)$ copies of K_{r+1} , hence contains $\frac{n}{3} \binom{r}{2} - O(1)$ triangles. This implies that, for large n , a graph with smaller regularity cannot contain more triangles than our construction. Therefore, r is at least the regularity of the claimed unique construction. Using Theorem 3.2, in each of the cases we know that r is at most the regularity of the claimed unique construction, and it is left to show that no other r -regular graph can contain at least as many triangles as our construction, in each case.

As G is P_k -free, we have $r \leq k-2$ by Theorem 3.2. Therefore, $\mathcal{N}(K_3, G) \leq \frac{n}{3} \binom{k-2}{2} = \frac{n}{k-1} \binom{k-1}{3}$, with equality only when $(k-1)|n$ and G is $n/k-1$ disjoint copies of K_{k-1} , proving the first case. If n is not divisible by $(k-1)$ and $r = k-2$, then by Theorem 3.2 we have that P_k is a star, i.e., $k \leq 3$. Consequently, $r < k-2$.

In the second case, we have k is odd and assume $r = k-3$. We can write n as $a(k-2) + b$, where $0 \leq b \leq k-3$. Since $k \geq 7$, we have $k-3 > k/2$, and hence, by the result of Erdős and Gallai, each component has at most $k-1$ vertices. Thus, each component is either a K_{k-2} or G'_{k-1} , for these are the only $(k-3)$ -regular graphs on at most $k-1$ vertices. This means $G = xK_{k-2} + yG'_{k-1}$, which gives $x(k-2) + y(k-1) = n = a(k-2) + b$, implying

$$y = a'(k-2) + b, \text{ where } a' = a - x - y.$$

If $y > b$, then $a' \geq 1$, and hence, $y \geq (k-2) + b$, then we can replace $k-2$ copies of G'_{k-1} by $k-1$ copies of K_{k-2} , increasing the number of triangles as $\mathcal{N}(K_3, K_{k-2}) > \mathcal{N}(K_3, G'_{k-1})$, contradicting the choice of G . Also, if $y < b$, then as $k-2 > 0$, we must have $a' < 0$, which implies $y < 0$, a contradiction. Therefore, $y = b$ and $x = a - b$, which proves the second case.

Note that if k is even, then K_{k-1} does not contain a perfect matching, and hence, if $(k-2) \nmid n$, then $r < k-3$, leading to the third case.

Finally, assume $r = k-4$ and $k \geq 7$ is even. We can write n as $a(k-3) + b$, where $0 \leq b \leq k-4$. First recall that each graph in \mathcal{G}_{k-1} contains the same number of triangles. By the same reasoning of the previous case, we may assume that G consists of $xK_{k-3} + yG_{k-2}$ and z copies of graphs from \mathcal{G}_{k-1} . This gives $x(k-3) + y(k-2) + z(k-1) = n = a(k-3) + b$, implying

$$2z \geq a'(k-3) - y + b, \text{ where } a' = a - x - y - z.$$

Note that $\mathcal{N}(K_3, K_{k-3}) + \mathcal{N}(K_3, G_{k-1}) > 2\mathcal{N}(K_3, G_{k-2})$, for any graph $G_{k-1} \in \mathcal{G}_{k-1}$, and hence, whenever there are two copies of G_{k-2} in G , we can replace them by a copy of K_{k-3} and a copy of G_{k-1} , increasing the number of triangles. Therefore, we have that y is either 0 or 1. If $z > \lfloor b/2 \rfloor$, then we have $a' \geq 1$, which means $z \geq (k-3)/2 - y/2 + b/2$. If b is even we may assume $y = 0$, and hence, in both cases of b being odd or even, we still have $z \geq \lceil (k-3)/2 \rceil + \lfloor b/2 \rfloor$. We can then replace $\lceil (k-3)/2 \rceil$ copies of G_{k-1} by $\lfloor (k-1)/2 \rfloor$ copies of K_{k-3} and a copy of G_{k-2} , increasing the number of triangles, which contradicts the

extremality of G . Again, due to compatibility of the number of vertices, z cannot be less than $\lfloor b/2 \rfloor$. This completes the proof. ■

Funding: Research supported by the National Research, Development and Innovation Office - NKFIH under the grants FK 132060 and KKP-133819.

References

- [1] N. Alon, C. Shikhelman. Many T copies in H -free graphs. *Journal of Combinatorial Theory, Series B*, **121**, 146–172, 2016.
- [2] B. Andrásfai, P. Erdős and V. T. Sós. On the connection between chromatic number, maximal clique and minimum degree of a graph, *Discrete Math.* **8**, 205–218, 1974.
- [3] J. A. Bondy and M. Simonovits. Cycles of even length in graphs. *Journal of Combinatorial Theory, Series B*, **16**(2), 97–105, 1974.
- [4] S. Cambie, R. de Joannis de Verclos, R. J. Kang. Regular Turán numbers and some Gan–Loh–Sudakov-type problems. *Journal of Graph Theory*, **102**(1), 67–85, 2023.
- [5] Y. Caro and Zs. Tuza. Regular Turán numbers. *The Australasian Journal of Combinatorics*, **78**, 133–144, 2020.
- [6] P. Erdős and T. Gallai. On maximal paths and circuits of graphs. *Acta Math. Acad. Sci. Hungar.*, **10**, 337–356, 1959.
- [7] P. Erdős, H. Sachs. Reguläre Graphen gegebener Tailenweite mit minimaler Knotenzahl (in German). *Wiss. Z. Martin-Luther-Univ. Halle-Wittenberg Math.-Natur. Reihe* **12**, 251–257, 1963.
- [8] Z. Füredi, M. Simonovits, (2013). The history of degenerate (bipartite) extremal graph problems. In Erdős centennial (pp. 169–264). Berlin, Heidelberg: Springer Berlin Heidelberg.
- [9] D. Gerbner. On Turán-good graphs, *Discrete Mathematics*, **344**(8), 112445, 2021.
- [10] D. Gerbner, C. Palmer. Some exact results for generalized Turán problems. *European Journal of Combinatorics*, **103**, 103519, 2022.
- [11] D. Gerbner, B. Patkós, Zs. Tuza, M. Vizer. Singular Turán numbers and WORM-colorings, *Disc. Math. Graph Theory*, **42**(4), 1061–1074, 2022.
- [12] D. Gerbner, B. Patkós, Z. Tuza, M. Vizer. Some exact results for regular Turán problems, *European Journal of Combinatorics*, available online, 2023.

- [13] L. Gishboliner and A. Shapira. A generalized Turán problem and its applications. In Proceedings of the 50th Annual ACM SIGACT Symposium on Theory of Computing, pages 760–772. ACM, 2018.
- [14] A. Grzesik, E Győri, N. Salia, C. Tompkins. Subgraph Densities in K_r -Free Graphs. *The Electronic Journal of Combinatorics*, P1.51, 2023.
- [15] E. Győri, J. Pach, M. Simonovits. On the maximal number of certain subgraphs in K_r -free graphs. *Graphs and Combinatorics*, **7**(1), 31–37, 1991.
- [16] E. Győri, N. Salia, C. Tompkins, O. Zamora. The maximum number of P_l copies in P_k -free graphs. *Discrete Mathematics & Theoretical Computer Science*, **21**, #14 2019.
- [17] M. Tait, C. Timmons. Regular Turán numbers of complete bipartite graphs. *Discrete Mathematics*, **344**(10), 112531, 2021.
- [18] P. Turán. On an extremal problem in graph theory. *Mat. Fiz. Lapok* **48**, 436–452, 1941.
- [19] A. A. Zykov. On some properties of linear complexes. *Matematicheskii Sbornik*, **66**(2), 163–188, 1949.