### CENTERS OF QUANTUM-WAJSBERG ALGEBRAS

### LAVINIA CORINA CIUNGU

Department of Mathematics
St Francis College
179 Livingston Street, Brooklyn, New York, NY 11201, USA
lciungu@sfc.edu

ABSTRACT. We define the Wajsberg-center and the OML-center of a quantum-Wajsberg algebra, and study their structures. We prove that the Wajsberg-center is a Wajsberg subalgebra of a quantum-Wajsberg algebra, and that it is a distributive sublattice of its corresponding poset. If the quantum-Wajsberg algebra is quasi-linear, we show that the Wajsberg-center is a linearly ordered Wajsberg algebra. We also show that the lattice subreduct of the Wajsberg-center is a Kleene algebra. Furthermore, we prove that the OML-center is an orthomodular lattice, and that the orthomodular lattices form a subvariety of the variety of quantum-Wajsberg algebras.

 $\bf Keywords:$  quantum-Wajsberg algebra, Wajsberg-center, OML-center, distributive lattice, orthomodular lattice, Kleene algebra

**AMS** classification (2020): 06F35, 03G25, 06A06, 81P10, 06C15

#### 1. Introduction

In the last decades, the study of algebraic structures related to the logical foundations of quantum mechanics became a central topic of research. Generally known as quantum structures, these algebras serve as algebraic semantics for the classical and non-classical logics, as well as for the quantum logics. As algebraic structures connected with quantum logics we mention the following algebras: bounded involutive lattices, De Morgan algebras, ortholattices, orthomodular lattices, MV algebras, quantum MV algebras.

The quantum-MV algebras (or QMV algebras) were introduced by R. Giuntini in [7] as non-lattice generalizations of MV algebras ([3]) and as non-idempotent generalizations of orthomodular lattices ([1, 26]). These structures were intensively studied by R. Giuntini ([8, 9, 10, 11, 12]), A. Dvurečenskij and S. Pulmannová ([5]), R. Giuntini and S. Pulmannová ([13]) and by A. Iorgulescu in [20, 21, 22, 23, 24, 25]. An extensive study on the orthomodular structures as quantum logics can be found in [31]. Many algebraic semantics for the classical and non-classical logics studied so far (pseudo-effect algebras, residuated lattices, pseudo-MV/BL/MTL algebras, bounded non-commutative  $R\ell$ -monoids, pseudo-hoops, pseudo-BCK/BCI algebras), as well as their commutative versions, are quantum-B algebras.

Quantum-B algebras, defined and investigated by W. Rump and Y.C. Yang ([33, 32]), arise from the concept of quantales which was introduced in 1984 as a framework for quantum mechanics with a view toward non-commutative logic ([29]). Interesting results on quantum-B algebras have been presented in [34, 35, 15, 16].

We redefined in [4] the quantum-MV algebras starting from involutive BE algebras and we introduced and studied the notion of quantum-Wajsberg algebras (QW algebras, for short). We proved that any Wajsberg algebra is a quantum-Wajsberg algebra, and the commutative quantum-Wajsberg algebras are Wajsberg algebras. It was also shown that the Wajsberg algebras are both quantum-Wajsberg algebras and commutative quantum-B algebras.

In this paper, we define the Wajsberg-center or the commutative center of a quantum-Wajsberg algebra X as the set of those elements of X that commute with all other elements of X. We study certain properties of the Wajsberg-center, and we prove that the Wajsberg-center is a Wajsberg subalgebra of X, and it is also a distributive sublattice of its corresponding poset. If the quantum-Wajsberg algebra is quasi-linear, we show that the Wajsberg-center is a linearly ordered Wajsberg algebra. We also prove that the lattice subreduct of the Wajsberg-center is a Kleene algebra. Furthermore, we define the OML-center of a quantum-Wajsberg algebra, and study its properties. We prove that the OML-center is an orthomodular lattice, and that the orthomodular lattices form a subvariety of the variety of quantum-Wajsberg algebras. Additionally, we prove new properties of quantum-Wajsberg algebras.

#### 2. Preliminaries

In this section, we recall some basic notions and results regarding BCK algebras, Wajsberg algebras, BE algebras and quantum-Wajsberg algebras that will be used in the paper. Additionally, we prove new properties of quantum-Wajsberg algebras. For more details regarding the quantum-Wajsberg algebras we refer the reader to [4].

Starting from the systems of positive implicational calculus, weak systems of positive implicational calculus and BCI and BCK systems, in 1966 Y. Imai and K. Isèki introduced the BCK algebras ([17]). BCK algebras are also used in a dual form, with an implication  $\rightarrow$  and with one constant element 1, that is the greatest element ([28]). A (dual) BCK algebra is an algebra  $(X, \rightarrow, 1)$  of type (2,0) satisfying the following conditions, for all  $x, y, z \in X$ :  $(BCK_1)$   $(x \rightarrow y) \rightarrow ((y \rightarrow z) \rightarrow (x \rightarrow z)) = 1$ ;  $(BCK_2)$   $1 \rightarrow x = x$ ;  $(BCK_3)$   $x \rightarrow 1 = 1$ ;  $(BCK_4)$   $x \rightarrow y = 1$  and  $y \rightarrow x = 1$  imply x = y. In this paper, we use the dual BCK algebras. If  $(X, \rightarrow, 1)$  is a BCK algebra, for  $x, y \in X$  we define the relation  $\leq$  by  $x \leq y$  if and only if  $x \rightarrow y = 1$ , and  $\leq$  is a partial order on X.

Wajsberg algebras were introduced in 1984 by Font, Rodriguez and Torrens in [6] as algebraic model of  $\aleph_0$ -valued Łukasiewicz logic. A Wajsberg algebra is an algebra  $(X, \to, ^*, 1)$  of type (2,1,0) satisfying the following conditions for all  $x,y,z\in X$ :  $(W_1)$   $1\to x=x$ ;  $(W_2)$   $(y\to z)\to ((z\to x)\to (y\to x))=1$ ;  $(W_3)$   $(x\to y)\to y=(y\to x)\to x$ ;  $(W_4)$   $(x^*\to y^*)\to (y\to x)=1$ . Wajsberg algebras are bounded with  $0=1^*$ , and they are involutive. It was proved in [6] that Wajsberg algebras are termwise equivalent to MV algebras.

BE algebras were introduced in [27] as algebras  $(X, \to, 1)$  of type (2,0) satisfying the following conditions, for all  $x, y, z \in X$ :  $(BE_1)$   $x \to x = 1$ ;  $(BE_2)$   $x \to 1 = 1$ ;  $(BE_3)$   $1 \to x = x$ ;  $(BE_4)$   $x \to (y \to z) = y \to (x \to z)$ . A relation  $\leq$  is defined on X by  $x \leq y$  iff  $x \to y = 1$ . A BE algebra X is bounded if there exists  $0 \in X$  such that  $0 \leq x$ , for all  $x \in X$ . In a bounded BE algebra  $(X, \to, 0, 1)$  we define  $x^* = x \to 0$ , for all  $x \in X$ . A bounded BE algebra X is called involutive if  $x^{**} = x$ , for any  $x \in X$ .

A BE algebra X is called *commutative* if  $(x \to y) \to y = (y \to x) \to x$ , for all  $x, y \in X$ . A

bounded BE algebra X is called *involutive* if  $x^{**} = x$ , for any  $x \in X$ .

Obviously, any BCK algebra is a BE algebra, but the exact connection between BE algebras and BCK algebras is made in the papers [18, 19]: a BCK algebra is a BE algebra satisfying  $(BCK_4)$  (antisymmetry) and  $(BCK_1)$ .

A suplement algebra (S-algebra, for short) is an algebra  $(X, \oplus, *, 0, 1)$  of type (2, 1, 0, 0) satisfying the following axioms for all  $x, y, z \in X$ :  $(S_1)$   $x \oplus y = y \oplus x$ ;  $(S_2)$   $x \oplus (y \oplus z) = (x \oplus y) \oplus z$ ;  $(S_3)$   $x \oplus x^* = 1$ ;  $(S_4)$   $x \oplus 0 = x$ ;  $(S_5)$   $x^{**} = x$ ;  $(S_6)$   $0^* = 1$ ;  $(S_7)$   $x \oplus 1 = 1$  ([14]).

The following additional operations can be defined in a supplement algebra:

$$x \odot y = (x^* \oplus y^*)^*, \ x \cap_S y = (x \oplus y^*) \odot y, \ x \cup_S y = (x \odot y^*) \oplus y.$$

A quantum-MV algebra (QMV algebra, for short) is an S-algebra  $(X, \oplus, ^*, 0, 1)$  satisfying the following axiom for all  $x, y, z \in X$  ([8]):

 $(QMV) \ x \oplus ((x^* \cap_S y) \cap_S (z \cap_S x^*)) = (x \oplus y) \cap_S (x \oplus z).$ 

**Lemma 2.1.** Let  $(X, \to, 1)$  be a BE algebra. The following hold for all  $x, y, z \in X$ :

- (1)  $x \rightarrow (y \rightarrow x) = 1;$
- $(2) \ x \leq (x \to y) \to y.$

If X is bounded, then:

- (3)  $x \rightarrow y^* = y \rightarrow x^*$ ;
- $(4) \ x \leq x^{**}$ .

If X is involutive, then:

- (5)  $x^* \to y = y^* \to x$ ;
- (6)  $x^* \to y^* = y \to x$ ;
- (7)  $(x \to y)^* \to z = x \to (y^* \to z);$
- (8)  $x \to (y \to z) = (x \to y^*)^* \to z;$
- (9)  $(x^* \to y)^* \to (x^* \to y) = (x^* \to x)^* \to (y^* \to y).$

*Proof.* (1)-(6) See [4].

- (7) Applying  $(BE_4)$  we get:  $(x \to y)^* \to z = z^* \to (x \to y) = x \to (z^* \to y) = x \to (y^* \to z)$ .
- (8) Using  $(BE_4)$ , we have:  $x \to (y \to z) = x \to (z^* \to y^*) = z^* \to (x \to y^*) = (x \to y^*)^* \to z$ .
- (9) Applying twice (7), we get:  $(x^* \to y)^* \to (x^* \to y) = x^* \to (y^* \to (x^* \to y)) = x^* \to (x^* \to (y^* \to y)) = (x^* \to x)^* \to (y^* \to y).$

In a BE algebra X, we define the additional operation  $x \cup y = (x \to y) \to y$ . If X is involutive, we define the operations  $x \cap y = ((x^* \to y^*) \to y^*)^* = (x^* \cup y^*)^*$ ,  $x \odot y = (x \to y^*)^* = (y \to x^*)^*$ , and the relation  $\leq_Q$  by  $x \leq_Q y$  iff  $x = x \cap y$ .

**Proposition 2.2.** Let X be an involutive BE algebra. Then the following hold for all  $x, y, z \in X$ :

- (1)  $x \leq_Q y$  implies  $x \leq y$ ,  $x = y \cap x$  and  $y = x \cup y$ ;
- $(2) \leq_Q is \ reflexive \ and \ antisymmetric;$
- $(3) (x \cap y) \to z = (y \to x) \to (y \to z);$
- $(4) (x \cap y)^* \to (y \to x)^* = y \uplus (y \to x)^*;$
- $(5) (x \cap (y \cap z))^* = ((z \to x) \cap (z \to y)) \to z^*;$
- (6)  $x, y \leq_Q z$  and  $z \to x = z \to y$  imply x = y; (cancellation law)
- (7)  $x \cap y = y \odot (y \rightarrow x)$ .

*Proof.* (1) - (3) See [4].

(4) We have:

$$(x \cap y)^* \to (y \to x)^* = ((x^* \to y^*) \to y^*) \to (y \to x)^*$$

$$= ((y \to x) \to y^*) \to (y \to x)^*$$

$$= (y \to (y \to x)^*) \to (y \to x)^* = y \uplus (y \to x)^*.$$

(5) Aplying (3), we get:

$$\begin{array}{c} ((z \to x) \cap (z \to y)) \to z^* = ((x^* \to z^*) \cap (y^* \to z^*)) \to z^* \\ = ((y^* \to z^*) \to (x^* \to z^*)) \to ((y^* \to z^*) \to z^*) \\ = (x^* \to ((y^* \to z^*) \to z^*)) \to ((y^* \to z^*) \to z^*) \\ = (x^* \to (y^* \uplus z^*)) \to (y^* \uplus z^*) \\ = (x^* \to (y \cap x)^*) \to (y \cap z)^* \\ = (x \cap (y \cap z))^*. \end{array}$$

(6) Since  $x, y \leq_Q z$  and  $z \to x = z \to y$ , we have:

$$x = x \cap z = ((x^* \to z^*) \to z^*)^* = ((z \to x) \to z^*)^* = ((z \to y) \to z^*)^* = ((y^* \to z^*) \to z^*)^* = y \cap z = y.$$

(7) We have 
$$y \odot (y \to x) = (y \to (y \to x)^*)^* = ((y \to x) \to y^*)^* = ((x^* \to y^*) \to y^*)^* = x \cap y$$
.

A (left-)quantum-Wajsberg algebra (QW algebra, for short)  $(X, \to, ^*, 1)$  is an involutive BE algebra  $(X, \to, ^*, 1)$  satisfying the following condition for all  $x, y, z \in X$ :

$$(\mathrm{QW}) \ x \to ((x \cap y) \cap (z \cap x)) = (x \to y) \cap (x \to z).$$

Condition (QW) is equivalent to the following conditions:

$$(QW_1) \ x \to (x \cap y) = x \to y;$$

$$(QW_2)$$
  $x \to (y \cap (z \cap x)) = (x \to y) \cap (x \to z).$ 

**Definition 2.3.** ([20]) A (left-)m-BE algebra is an algebra  $(X, \odot, *, 1)$  of type (2, 1, 0) satisfying the following properties, for all  $x, y, z \in X$ : (PU)  $1 \odot x = x = x \odot 1$ ; (Pcomm)  $x \odot y = y \odot x$ ; (Pass)  $x \odot (y \odot z) = (x \odot y) \odot z$ ; (m-L)  $x \odot 0 = 0$ ; (m-Re)  $x \odot x^* = 0$ , where  $0 := 1^*$ .

Note that, according to [25, Cor. 17.1.3], the involutive (left-)BE algebras  $(X, \to, ^*, 1)$  are term-equivalent to involutive (left-)m-BE algebras  $(X, \odot, ^*, 1)$ , by the mutually inverse transformations ([20, 25]):

$$\Phi: x \odot y := (x \to y^*)^* \quad \text{and} \quad \Psi: x \to y := (x \odot y^*)^*.$$

**Definition 2.4.** ([24, Def. 3.10]) A (left-)quantum-MV algebra, or a (left-)QMV algebra for short, is an involutive (left-)m-BE algebra  $(X, \odot, ^*, 1)$  verifying the following axiom: for all  $x, y, z \in X$ ,

$$(\operatorname{Pqmv}) \ x \odot ((x^* \uplus y) \uplus (z \uplus x^*)) = (x \odot y) \uplus (x \odot z).$$

**Proposition 2.5.** The (left-)quantum-Wajsberg algebras are term-equivalent to (left-)quantum-MV algebras.

*Proof.* We prove that the axioms (Pqmv) and (QW) are equivalent. Using the transformation  $\Phi$ , from (Pqmv) we get:

$$x\odot((x^*\uplus y)\uplus(z\uplus x^*))=(x\to((x^*\uplus y)\uplus(z\uplus x^*))^*)^*=(x\to((x\Cap y^*)\Cap(z^*\Cap x)))^*\text{ and }(x\odot y)\uplus(x\odot z)=(x\to y^*)^*\uplus(x\to z^*)^*=((x\to y^*)\Cap(x\to z^*))^*,$$
 hence (Pqmv) becomes:

$$(x \to ((x \cap y^*) \cap (z^* \cap x)))^* = ((x \to y^*) \cap (x \to z^*))^*,$$

for all  $x, y, z \in X$ . Replacing y by  $y^*$  and z by  $z^*$ , we get axiom (QW). Similarly axiom (QW) implies axiom (Pqmv).

In what follows, by quantum-MV algebras and quantum-Wajsberg algebras we understand the left-quantum-MV algebras and left-quantum-Wajsberg algebras, respectively.

**Proposition 2.6.** ([4]) Let X be a quantum-Wajsberg algebra. The following hold for all  $x, y, z \in X$ :

```
(1) x \to (y \cap x) = x \to y and (x \to y) \to (y \cap x) = x;
```

- (2)  $x \leq_Q x^* \to y \text{ and } x \leq_Q y \to x;$
- (3)  $x \le y$  iff  $y \cap x = x$ ;
- $(4) (x \rightarrow y) \uplus (y \rightarrow x) = 1.$

If  $x \leq_Q y$ , then:

- $(5) \ y = y \uplus x;$
- (6)  $y^* \leq_Q x^*$ ;
- (7)  $y \to z \leq_Q x \to z \text{ and } z \to x \leq_Q z \to y;$
- (8)  $x \cap z \leq_Q y \cap z \text{ and } x \cup z \leq_Q y \cup z;$
- (9)  $x \odot z \leq_Q y \odot z$ .

**Proposition 2.7.** Let X be a quantum-Wajsberg algebra. The following hold, for all  $x, y, z \in X$ :

- $(1) (x \cap y) \cap (y \cap z) = (x \cap y) \cap z;$
- $(2) \leq_Q is transitive;$
- $(3) (z \cap x) \to (y \cap x) = (z \cap x) \to y;$
- (4)  $x \leq_Q y$  and  $y \leq x$  imply x = y;
- (5)  $x \leq_Q y \text{ implies } x \cap (y \cap z) = x \cap z;$
- (6)  $z \cap ((y^* \rightarrow z) \cap (x^* \rightarrow y)) = z \cap (x^* \rightarrow y);$
- $(7) \ x \uplus (x \to y)^* = x;$
- (8)  $x = y \rightarrow x \text{ iff } y = x \rightarrow y;$
- $(9) \ x \cap y, y \cap x \leq_Q x \to y.$

*Proof.* (1) - (3) See [4].

- (4) By Proposition 2.6(3),  $y \le x$  implies  $x \cap y = y$ . Since  $x \le_Q y$ , we have  $x \cap y = x$ , hence x = y.
- (5) Using (1),  $(x \cap y) \cap (y \cap z) = (x \cap y) \cap z$ . Since  $x \leq_Q y$  implies  $x \cap y = x$ , we get  $x \cap (y \cap z) = x \cap z$ .
- (6) It follows by (5), since  $z \leq_Q y^* \to z$ ;
- (7) By Proposition 2.6(2),(5), we have  $x^* \leq_Q x \to y$ , so that  $(x \to y)^* \leq_Q x$  and  $x \uplus (x \to y)^* = x$ .
- (8) Suppose  $x = y \to x$ , so that  $y^* \cap x^* = (y \cup x)^* = ((y \to x) \to x)^* = (x \to x)^* = 1^* = 0$ . Using  $(QW_1)$ , we get  $y = (y^*)^* = y^* \to 0 = y^* \to (y^* \cap x^*) = y^* \to x^* = x \to y$ . The converse follows similarly.
- (9) Using Proposition 2.6(2), we have  $y \leq_Q x \to y$ , so that  $(x \to y)^* \leq_Q y^* \leq_Q (x^* \to y^*) \to y^* = (x \cap y)^*$ . Hence  $x \cap y \leq_Q x \to y$ . Similarly  $(x \to y)^* \leq (x \to y) \to x^* = (y^* \to x^*) \to x^* = (y \cap x)^*$ . Thus  $y \cap x \leq_Q x \to y$ .

By Propositions 2.2(2), 2.7(2), in a quantum-Wajsberg algebra X,  $\leq_Q$  is a partial order on X.

A quantum-Wajsberg algebra X is called *commutative* if  $x \cup y = y \cup x$ , or equivalently  $x \cap y = y \cap x$  for all  $x, y \in X$ .

Since:

- commutative BE algebras are commutative BCK algebras ([36]]),
- bounded commutative BCK are term-equivalent to MV algebras ([30]) and
- Wajsberg algebras are term-equivalent to MV algebras ([6]),

it follows that bounded commutative BE algebras are bounded commutative BCK algebras, hence are term-equivalent to MV algebras, hence to Wajsberg algebras.

Hence the commutative quantum-Wajsberg algebras are the Wajsberg algebras.

It was proved in [4] that a quantum-Wajsberg algebra is a bounded commutative BCK algebra, that is a Wajsberg algebra, if and only if the relations  $\leq$  and  $\leq_Q$  coincide.

**Proposition 2.8.** ([4]) Let  $(X, \to, 0, 1)$  be a bounded commutative BCK algebra. The following hold for all  $x, y, z \in X$ :

- $\begin{array}{l} (1)\ x \leq_Q y\ and\ x \leq_Q z\ imply\ x \leq_Q y \cap z; \\ (2)\ y \leq_Q x\ and\ z \leq_Q x\ imply\ y \uplus z \leq_Q x; \end{array}$
- (3)  $x \leq_Q y$  implies  $x \cup z \leq_Q y \cup z$  and  $x \cap z \leq_Q y \cap z$ .

### 3. The Wajsberg-center of quantum-Wajsberg algebras

In this section, we investigate the commutativity property of quantum-Wajsberg algebras. We define the Wajsberg-center or the commutative center of a quantum-Wajsberg algebra X as the set of those elements of X that commute with all other elements of X. We study certain properties of the Wajsberg-center, and prove that the Wajsberg-center is a Wajsberg subalgebra of X. In what follows,  $(X, \rightarrow, ^*, 1)$  will be a quantum-Wajsberg algebra, unless otherwise stated.

**Definition 3.1.** We say that the elements  $x, y \in X$  commute, denoted by  $x \mathcal{C} y$ , if  $x \cap y = y \cap x$ .

**Definition 3.2.** The commutative center of X is the set  $\mathcal{Z}(X) = \{x \in X \mid x\mathcal{C}y, \text{ for all } x \in$  $y \in X$  }.

Obviously  $0, 1 \in \mathcal{Z}(X)$ .

**Lemma 3.3.** If xCy, then  $x \cup y = y \cup x$ .

*Proof.* Applying twice Proposition 2.6(1), we have:

$$x \uplus y = (x \to y) \to y = (x \to y) \to ((y \to x) \to (x \cap y))$$

$$= (y \to x) \to ((x \to y) \to (x \cap y))$$

$$= (y \to x) \to ((x \to y) \to (y \cap x)) = (y \to x) \to x = y \uplus x.$$

**Lemma 3.4.** Let  $x, y \in X$ . The following are equivalent:

- (a) xCy;
- (b)  $(x \to y) \to (x \cap y) = x$ .

*Proof.*  $(a) \Rightarrow (b)$  By Proposition 2.6(1), we get  $x = (x \rightarrow y) \rightarrow (y \cap x) = (x \rightarrow y) \rightarrow (x \cap y)$ .  $(b) \Rightarrow (a)$  Suppose  $(x \rightarrow y) \rightarrow (x \cap y) = x$ , and applying Proposition 2.6(1), we have:  $(x \to y) \to (x \cap y) = (x \to y) \to (y \cap x) = x$ . Since by Proposition 2.7(9),  $x \cap y$ ,  $y \cap x \leq_Q x \to y$ , by cancellation law (Proposition 2.2(6)), we get  $x \cap y = y \cap x$ . Hence xCy.

**Proposition 3.5.** The following hold:

- (1) the relation C is reflexive and symmetric;
- (2) if  $x \leq_Q y$  or  $y \leq_Q x$ , then xCy;
- (3) xCy implies  $x^*Cy^*$ ;
- $(4) (x \cap y)^* \mathcal{C}(x \to y)^*.$

*Proof.* (2) If  $x \leq_Q y$ , then  $x = x \cap y$  and, by Proposition 2.2(1) we have  $x = y \cap x$ . Hence  $x \cap y = y \cap x$ , that is  $x \in \mathcal{C}y$ , and similarly  $y \leq_Q x$  implies  $x \in \mathcal{C}y$ .

- (3) Using Lemma 3.3, we have:  $x^* \cap y^* = (x \cup y)^* = (y \cup x)^* = y^* \cap x^*$ , hence  $x^* \mathcal{C} y^*$ .
- (4) Since  $x \cap y \leq_Q y \leq_Q x \to y$ , we get  $(x \to y)^* \leq_Q (x \cap y)^*$ . Applying (2), it follows that  $(x \cap y)^* \mathcal{C}(x \to y)^*$ .

Corollary 3.6.  $\mathcal{Z}(X)$  is closed under \*.

*Proof.* Let  $x \in \mathcal{Z}(X)$ , that is  $x\mathcal{C}z$  for all  $z \in X$ . We also have  $x\mathcal{C}z^*$ , and applying Lemma 3.3 we have  $x \uplus z^* = z^* \uplus x$ . It follows that  $x^* \cap z = (x \uplus z^*)^* = (z^* \uplus x)^* = z \cap x^*$ . Hence  $x^* \in \mathcal{Z}(X)$ .

**Proposition 3.7.** If  $x, y, z \in X$  such that xCy and xCz, then  $(x \cap y) \cap z = y \cap (x \cap z)$ .

*Proof.* From  $x \cap y = y \cap x$ ,  $x \cap z = z \cap x$ , and applying Proposition 2.7(1),(3), we get:

$$(x \cap y) \cap z = (y \cap x) \cap z = (y \cap x) \cap (x \cap z)$$

$$= (((y \cap x)^* \to (x \cap z)^*) \to (x \cap z)^*)^*$$

$$= (((y \cap x)^* \to (z \cap x)^*) \to (z \cap x)^*)^*$$

$$= (((z \cap x) \to (y \cap x)) \to (z \cap x)^*)^*$$

$$= (((z \cap x) \to y) \to (z \cap x)^*)^*$$

$$= ((y^* \to (z \cap x)^*) \to (z \cap x)^*)^*$$

$$= y \cap (z \cap x) = y \cap (x \cap z).$$

**Corollary 3.8.** If xCy, yCz and xCz, then  $(x \cap y) \cap z = z \cap (x \cap y)$ .

*Proof.* By hypothesis and using Proposition 3.7, we get:

$$(x \cap y) \cap z = y \cap (x \cap z) = y \cap (z \cap x) = z \cap (y \cap x) = z \cap (x \cap y).$$

Corollary 3.9.  $\mathcal{Z}(X)$  is closed under  $\cap$ .

*Proof.* Let  $x, y \in \mathcal{Z}(X)$  and let  $z \in X$ . It follows that  $x\mathcal{C}y, y\mathcal{C}z, x\mathcal{C}z$ , and by Corollary 3.8 we get  $(x \cap y) \cap z = z \cap (x \cap y)$ . Hence  $x \cap y \in \mathcal{Z}(X)$ , that is  $\mathcal{Z}(X)$  is closed under  $\cap$ .

**Proposition 3.10.** Let  $x, y, z \in X$  such that yCz. Then  $x \to (y \cap z) \leq_Q (x \to y) \cap (x \to z)$ .

*Proof.* From  $z \cap y \leq_Q y$ , we get  $x \to (z \cap y) \leq_Q x \to y$ , so that  $(x \to y)^* \leq_Q (x \to (z \cap y))^*$  and  $(x \to (z \cap y))^* \to (x \to z)^* \leq_Q (x \to y)^* \to (x \to z)^*$ . It follows that:

 $((x \rightarrow (z \mathbin{\widehat{\boxtimes}} y))^* \rightarrow (x \rightarrow z)^*) \mathbin{\widehat{\boxtimes}} ((x \rightarrow y)^* \rightarrow (x \rightarrow z)^*) = (x \rightarrow (z \mathbin{\widehat{\boxtimes}} y))^* \rightarrow (x \rightarrow z)^*.$ 

Similarly, from  $y \cap z \leq_Q z$  we have  $x \to (y \cap z) \leq_Q x \to z$ , hence  $(x \to (y \cap z)) \cap (x \to z) = x \to (y \cap z)$ . Applying Proposition 2.2(5), and taking into consideration that yCz, we have:

```
 \begin{array}{l} (x \to (y \cap z)) \cap ((x \to y) \cap (x \to z)) = \\ = ((((x \to z) \to (x \to (y \cap z))) \cap ((x \to z) \to (x \to y))) \to (x \to z)^*)^* \\ = ((((x \to (y \cap z))^* \to (x \to z)^*) \cap ((x \to y)^* \to (x \to z)^*)) \to (x \to z)^*)^* \\ = ((((x \to (z \cap y))^* \to (x \to z)^*) \cap ((x \to y)^* \to (x \to z)^*)) \to (x \to z)^*)^* \\ = (((x \to (z \cap y))^* \to (x \to z)^*) \to (x \to z)^*)^* \end{array}
```

$$= (((x \to (y \cap z))^* \to (x \to z)^*) \to (x \to z)^*)^*$$

$$= (x \to (y \cap z)) \cap (x \to z) = x \to (y \cap z).$$
Hence  $x \to (y \cap z) \leq_Q (x \to y) \cap (x \to z).$ 

**Lemma 3.11.** If xCy, xCz, yCz, then  $y \cap (z \cap x) \leq_Q y \cap z$ .

*Proof.* From  $z \cap x = x \cap z \leq_Q z$ , by Proposition 2.6(8) we get  $(z \cap x) \cap y \leq_Q z \cap y = y \cap z$ . Using Corollary 3.8, we get  $y \cap (z \cap x) \leq_Q y \cap z$ .

**Proposition 3.12.** *If* xCy, xCz, yCz, then  $(x \to y) \cap (x \to z) \leq x \to (y \cap z)$ .

*Proof.* Applying Proposition 2.2(3) and Lemma 3.11, we get:

$$\begin{split} ((x \to y) & \cap (x \to z)) \to (x \to (y \cap z)) = \\ & = (x \to (y \cap z))^* \to ((x \to y) \cap (x \to z))^* \\ & = (x \to (y \cap z))^* \to ((x \to y)^* \cup (x \to z)^*) \\ & = (x \to (y \cap z))^* \to (((x \to y)^* \to (x \to z)^*) \to (x \to z)^*) \\ & = ((x \to y)^* \to (x \to z)^*) \to ((x \to (y \cap z))^* \to (x \to z)^*) \\ & = ((x \to z) \to (x \to y)) \to ((x \to z) \to (x \to (y \cap z))) \\ & = ((z \cap x) \to y) \to ((z \cap x) \to (y \cap z)) \\ & = (y \cap (z \cap x)) \to (y \cap z) = 1. \end{split}$$

It follows that  $(x \to y) \cap (x \to z) \le x \to (y \cap z)$ .

**Proposition 3.13.** If xCy, xCz, yCz, then  $x \to (y \cap z) = (x \to y) \cap (x \to z)$ .

*Proof.* It follows by Propositions 3.10, 3.12, 2.7(4).

Corollary 3.14. If  $y \in \mathcal{Z}(X)$  and  $x, z \in X$ , then  $(z \cap x)^* \to ((z \to x)^* \cap y) = ((z \cap x)^* \to (z \to x)^*) \cap ((z \cap x)^* \to y)$ .

*Proof.* It follows by Propositions 3.5(4) and 3.13, since  $y\mathcal{C}(z \cap x)^*$  and  $y\mathcal{C}(z \to x)^*$ .

Corollary 3.15. If  $x, y, z \in \mathcal{Z}(X)$ , then  $(z \cap x)^* \to y = (y^* \to z) \cap (x^* \to y)$ .

*Proof.* Since  $y \in \mathcal{Z}(X)$  implies  $y^* \in \mathcal{Z}(X)$ , applying Proposition 3.13, we get:  $(z \cap x)^* \to y = y^* \to (z \cap x) = (y^* \to z) \cap (y^* \to x) = (y^* \to z) \cap (x^* \to y)$ .

**Proposition 3.16.** If  $x, y \in \mathcal{Z}(X)$ , then  $x^* \to y \in \mathcal{Z}(X)$ .

*Proof.* Let  $x, y \in \mathcal{Z}(X)$  and let  $z \in X$ . Then  $x\mathcal{C}z$  and  $y\mathcal{C}(z \to x)^*$ . Applying Lemma 3.4, we get:

$$z = (z \to x) \to (z \cap x) = ((z \to x)^*)^* \to (z \cap x)$$
 and  $(z \to x)^* = ((z \to x)^* \to y) \to ((z \to x)^* \cap y)$ ,

respectively. It follows that:

$$z = ((z \to x)^*)^* \to (z \cap x)$$

$$= (((z \to x)^* \to y) \to ((z \to x)^* \cap y))^* \to (z \cap x)$$

$$= ((z \to x)^* \to y) \to (((z \to x)^* \cap y)^* \to (z \cap x)) \text{ (by Lemma 2.1(7))}$$

$$= ((z \to x)^* \to y) \to (((z \cap x)^* \to ((z \to x)^* \cap y))$$

$$= ((z \to x)^* \to y) \to (((z \cap x)^* \to (z \to x)^*) \cap ((z \cap x)^* \to y)) \text{ (by Corollary 3.14)}$$

$$= ((z \to x)^* \to y) \to (((x \cap z)^* \to (z \to x)^*) \cap ((z \cap x)^* \to y))$$

$$= ((z \to x)^* \to y) \to ((z \cup (z \to x)^*) \cap ((z \cap x)^* \to y)) \text{ (by Proposition 2.2(4))}$$

$$= ((z \to x)^* \to y) \to (z \cap ((z \cap x)^* \to y)) \text{ (by Proposition 2.7(7))}$$

$$= ((z \to x)^* \to y) \to (z \cap ((y^* \to z) \cap (x^* \to y)) \text{ (by Corrolary 3.15)}$$

$$= ((z \to x)^* \to y) \to (z \cap (x^* \to y)) \text{ (by Proposition 2.7(6))}$$

$$= (z \to (x^* \to y)) \to (z \cap (x^* \to y)) \text{ (by Lemma 2.1(7))}.$$

Using Lemma 3.4, we conclude that  $x^* \to y \in \mathcal{Z}(X)$ .

Corollary 3.17. If  $x, y \in \mathcal{Z}(X)$ , then  $x \to y \in \mathcal{Z}(X)$ .

*Proof.* Since  $x \in \mathcal{Z}(X)$ , by Corollary 3.6 we get  $x^* \in \mathcal{Z}(X)$ . Applying Proposition 3.16,  $x^*, y \in \mathcal{Z}(X)$  implies  $x \to y \in \mathcal{Z}(X)$ .

**Theorem 3.18.**  $(\mathcal{Z}(X), \rightarrow, 0, 1)$  is a Wajsberg subalgebra of X.

*Proof.* Since by Corollary 3.17,  $x, y \in \mathcal{Z}(X)$  implies  $x \to y \in \mathcal{Z}(X)$ , it follows that  $\mathcal{Z}(X)$  is closed under  $\to$ . Moreover  $0, 1 \in \mathcal{Z}(X)$ , hence it is a quantum-Wajsberg subalgebra of X. Since  $x, y \in \mathcal{Z}(X)$  implies  $x \cap y = y \cap x$ ,  $(\mathcal{Z}(X), \to, 0, 1)$  is a commutative quantum-Wajsberg algebra, that is a bounded commutative BCK subalgebra of X. Hence it is a Wajsberg subalgebra of X.

Corollary 3.19. A quantum-Wajsberg algebra X is a Wajsberg algebra if and only if  $\mathcal{Z}(X) = X$ .

Taking into consideration the above results, the commutative center  $\mathcal{Z}(X)$  will be also called the *Wajsberg-center* of X. Similarly as in [7] for the case of QMV algebras, we define the notion of a quasi-linear quantum-Wajsberg algebra.

**Definition 3.20.** A QW algebra X is said to be *quasi-linear* if, for all  $x, y \in X$ ,  $x \nleq_Q y$  implies y < x.

**Proposition 3.21.** If X is a quasi-linear QW algebra, then  $\mathcal{Z}(X)$  is a linearly ordered Wajsberg algebra.

*Proof.* According to [4], a quantum-Wajsberg algebra is a Wajsberg algebra if and only if the relations  $\leq$  and  $\leq_Q$  coincide. Since  $\mathcal{Z}(X)$  is a quasi-linear Wajsberg algebra,  $x \nleq y$  implies y < x, that is  $\mathcal{Z}(X)$  is linearly ordered.

### 4. The lattice structure of Wajsberg-centers

We study certain lattice properties of the Wajsberg-center of a quantum-Wajsberg algebra, and prove that the Wajsberg-center of a quantum-Wajsberg algebra X is a distributive sublattice of the poset  $(X, \leq_Q, 0, 1)$ . If the quantum-Wajsberg algebra is quasi-linear, we prove that the Wajsberg-center is a linearly ordered Wajsberg algebra. Finally, we show that the lattice subreduct of the Wajsberg-center is a Kleene algebra. In what follows,  $(X, \rightarrow, ^*, 1)$  will be a quantum-Wajsberg algebra, unless otherwise stated.

**Proposition 4.1.** The following hold for all  $x, y, z \in \mathcal{Z}(X)$ :

- (1)  $x \to (y \cap z) = (x \to y) \cap (x \to z)$  (distributivity of  $\to$  over  $\cap$ );
- (2)  $x \odot (y \cup z) = (x \odot y) \cup (x \odot z)$  (distributivity of  $\odot$  over  $\cup$ );
- (3)  $x \cap (y \cup z) = (x \cap y) \cup (x \cap z)$  (distributivity of  $\cap$  over  $\cup$ );
- $(4) \ x \uplus (y \cap z) = (x \uplus y) \cap (x \uplus z) \ (distributivity \ of \uplus \ over \cap).$

*Proof.* (1) It follows by Proposition 3.13.

(2) Applying (1), we get:

$$\begin{array}{l} x\odot(y\uplus z)=(x\to (y\uplus z)^*)^*=(x\to (y^*\Cap z^*))^*\\ =((x\to y^*)\Cap (x\to z^*))^*=(x\to y^*)^*\uplus (x\to z^*)^*\\ =(x\odot y)\uplus (x\odot z). \end{array}$$

(3) By commutativity we have  $y, z \leq_Q y \uplus z$ , so that  $(y \uplus z) \to x \leq_Q y \to x, z \to x$ . Applying Propositions 2.6(9) and 2.2(7), we get:

$$\begin{array}{l} y\odot ((y \uplus z) \to x) \leq_Q y\odot (y \to x) = x \Cap y \text{ and } \\ z\odot ((y \uplus z) \to x) \leq_Q z\odot (z \to x) = x \Cap z. \end{array}$$

Using Proposition 2.2(7) and (2), we have:

$$x \mathbin{\cap} (y \mathbin{\cup} z) = (y \mathbin{\cup} z) \odot ((y \mathbin{\cup} z) \to x) = (y \odot ((y \mathbin{\cup} z) \to x)) \mathbin{\cup} (z \odot ((y \mathbin{\cup} z) \to x)) \\ \leq_Q (x \mathbin{\cap} y) \mathbin{\cup} (x \mathbin{\cap} z).$$

On the other hand,  $x \cap y \leq_Q y$ ,  $x \cap z \leq_Q z$  imply  $(x \cap y) \cup (x \cap z) \leq_Q y \cup z$ , and  $x \cap y \leq_Q x$ ,  $x \cap z \leq_Q x$  imply  $(x \cap y) \cup (x \cap z) \leq_Q x$ . Hence by Proposition 2.8,  $(x \cap y) \cup (x \cap z) \leq_Q x \cap (y \cup z)$ . Since  $\mathcal{Z}(X)$  is a commutative bounded BCK algebra, the relation  $\leq_Q$  is antisymmetric, and we conclude that  $x \cap (y \cup z) = (x \cap y) \cup (x \cap z)$ .

(4) Applying (3), we have:

$$\begin{array}{l} x \uplus (y \Cap z) = (x^* \Cap (y \Cap z)^*)^* = (x^* \Cap (y^* \uplus z^*))^* \\ = ((x^* \Cap y^*) \uplus (x^* \Cap z^*))^* = ((x \uplus y)^* \uplus (x \uplus z)^*)^* \\ = (x \uplus y) \Cap (x \uplus z). \end{array}$$

**Lemma 4.2.** The following hold for all  $x, y \in \mathcal{Z}(X)$ :

- (1)  $x \cup y$  is the least upper bound (l.u.b.) of  $\{x, y\}$ ;
- (2)  $x \cap y$  is the greatest lower bound (g.l.b.) of  $\{x, y\}$ .

Proof. (1) By Corollaries 3.9 and 3.6,  $\mathcal{Z}(X)$  is closed under  $\mathbb{Q}$  and  $\mathbb{Q}$ . Since  $x \uplus y = (x^* \cap y^*)^*$  for all  $x,y \in \mathcal{Z}(X)$ , it follows that  $\mathcal{Z}(X)$  is also closed under  $\mathbb{Q}$ . Since by commutativity  $x,y \leq_Q x \uplus y$ , it follows that  $x \uplus y$  is an upper bound of  $\{x,y\}$ . Let z be another upper bound of  $\{x,y\}$ , so that  $x,y \leq_Q z$ , that is  $x=x \cap z$  and  $y=y \cap z$ . Using Proposition 4.1(3), we have  $(x \uplus y) \cap z = z \cap (x \uplus y) = (z \cap x) \cup (z \cap y) = (x \cap z) \cup (y \cap z) = x \cup y$ . Hence  $x \cup y \leq_Q z$ , so that  $x \cup y$  is the l.u.b. of  $\{x,y\}$ .

(2) By commutativity we also have  $x \cap y \leq_Q x, y$ , thus  $x \cap y$  is a lower bound of  $\{x,y\}$ . Let z be another lower bound of  $\{x,y\}$ , so that  $z \leq_Q x, y$ , that is  $z = z \cap x$  and  $z = z \cap y$ . Using Proposition 3.7 and Corollary 3.8, we have:  $z \cap (x \cap y) = (x \cap y) \cap z = y \cap (x \cap z) = y \cap (z \cap x) = y \cap z = z \cap y = z$ , that is  $z \leq_Q x \cap y$ . It follows that  $x \cap y$  is the g.l.b. of  $\{x,y\}$ .

**Theorem 4.3.**  $(\mathcal{Z}(X), \cap, \cup, 0, 1)$  is a distributive sublattice of the poset  $(X, \leq_Q, 0, 1)$ .

*Proof.* It follows by Lemma 4.2, Theorem 3.18 and Proposition 4.1.

**Proposition 4.4.** The following hold for all  $x, y, z \in \mathcal{Z}(X)$ :

- (1)  $x \to (y \cup z) = (x \to y) \cup (x \to z)$  (distributivity of  $\to$  over  $\cup$ );
- (2)  $x \odot (y \cap z) = (x \odot y) \cap (x \cup z)$  (distributivity of  $\odot$  over  $\cap$ );
- (3)  $(y \cup z) \rightarrow x = (y \rightarrow x) \cap (z \rightarrow x);$
- $(4) (y \cap z) \to x = (y \to x) \cup (z \to x).$

*Proof.* (1) Since by commutativity  $y \uplus z \geq_Q y, z$ , we have  $x \to (y \uplus z) \geq_Q x \to y, x \to z$ , so that  $x \to (y \uplus z)$  is an upper bound of  $\{x \to y, x \to z\}$ . Let u be another upper bound of

 $\{x \to y, x \to z\}, \text{ that is } u \geq_Q x \to y, x \to z. \text{ It follows that } u \odot x \geq_Q (x \to y) \odot x = x \cap y \text{ and } u \odot x \geq_Q (x \to z) \odot x = x \cap z. \text{ Hence, by Proposition 4.1(3)}, \\ x \cap (y \cup z) = (x \cap y) \cup (x \cap z) \leq_Q u \odot x. \text{ Using } (QW_1), \text{ we get } x \to (y \cup z) = x \to (x \cap (y \cup z)) \leq_Q x \to (u \odot x) = x \to (u \to x^*)^* = (u \to x^*) \to x^* = u \cup x^* = u \text{ (since } u \geq_Q x \to y \geq_Q x^*). \text{ Thus } x \to (y \cup z) \text{ is the least upper bound of } \{x \to y, x \to z\}, \text{ and so } x \to (y \cup z) = (x \to y) \cup (x \to z).$ 

(2) Using (1), we have:

$$x \odot (y \cap z) = (x \to (y \cap z)^*)^* = (x \to (y^* \cup z^*))^* = ((x \to y^*) \cup (x \to z^*))^* = ((x \odot y)^* \cup (x \odot z)^*)^* = (x \odot y) \cap (x \odot z).$$

(3) Applying Proposition 4.1(1), we have:

$$(y \uplus z) \to x = x^* \to (y \uplus z)^* = x^* \to (y^* \cap z^*)$$
$$= (x^* \to y^*) \cap (x^* \to z^*) = (y \to x) \cap (z \to x).$$

(4) By (1), we get:

**Proposition 4.5.** The following hold for all  $x, y, z \in \mathcal{Z}(X)$ :

- $(1) (x \uplus y) \to (x \uplus z) \ge_Q x \uplus (y \to z);$
- $(2) (x \cap y) \to (x \cap z) \ge_Q x \cap (y \to z).$

*Proof.* (1) Applying Proposition 4.4, since  $y \to x \ge_Q x$  we get:

$$\begin{split} (x \uplus y) \to (x \uplus z) &= (x \to (x \uplus z)) \Cap (y \to (x \uplus z)) \\ &= ((x \to x) \uplus (x \to z)) \Cap ((y \to x) \uplus (y \to z)) \\ &= (1 \uplus (x \to z)) \Cap ((y \to x) \uplus (y \to z)) \\ &= 1 \Cap ((y \to x) \uplus (y \to z)) \\ &= (y \to x) \uplus (y \to z) \geq_Q x \uplus (y \to z). \end{split}$$

(2) Similarly, using Proposition 4.1 we have:

$$\begin{split} (x \cap y) &\to (x \cap z) = (x \to (x \cap z)) \uplus (y \to (x \cap z)) \\ &= ((x \to x) \cap (x \to z)) \uplus ((y \to x) \cap (y \to z)) \\ &= (1 \cap (x \to z)) \uplus ((y \to x) \cap (y \to z)) \\ &= (x \to z) \uplus ((y \to x) \cap (y \to z)) \\ &\geq_Q (y \to x) \cap (y \to z) \geq_Q x \cap (y \to z). \end{split}$$

**Proposition 4.6.** The following hold for all  $x, y \in \mathcal{Z}(X)$ :

- (1)  $(x^* \odot y) \cap (x \odot y^*) = 0$ ;
- $(2) (x \cap x^*) \odot (y \cap y^*) = 0;$
- $(3) x \cap x^* \leq_Q y \cup y^*.$

*Proof.* (1) Applying Proposition 2.6(4), we get:

$$(x^* \odot y) \cap (x \odot y^*) = (x^* \to y^*)^* \cap (x \to y)^* = (y \to x)^* \cap (x \to y)^* = ((y \to x) \cup (x \to y))^* = 1^* = 0.$$

(2) By distributivity of  $\odot$  over  $\cap$  and using (1), we have:

$$\begin{array}{l} (x \mathbin{\cap} x^*) \odot (y \mathbin{\cap} y^*) = ((x \mathbin{\cap} x^*) \odot y) \mathbin{\cap} ((x \mathbin{\cap} x^*) \odot y^*) \\ = (x \odot y) \mathbin{\cap} (x^* \odot y) \mathbin{\cap} (x \odot y^*) \mathbin{\cap} (x^* \odot y^*) \\ = (x \odot y) \mathbin{\cap} 0 \mathbin{\cap} (x^* \odot y^*) = 0. \end{array}$$

(3) Using (2), we get:

$$(x \cap x^*) \to (y \cup y^*) = ((x \cap x^*) \odot (y \cup y^*)^*)^* = ((x \cap x^*) \odot (y \cap y^*))^* = 0^* = 1.$$
 Since  $\leq_Q$  and  $\leq$  coincide in  $\mathcal{Z}(X)$ , it follows that  $x \cap x^* \leq_Q y \cup y^*$ .

**Definition 4.7.** A Kleene algebra is a structure  $(L, \wedge, \vee, *, 0, 1)$ , where  $(L, \wedge, \vee, 0, 1)$  is a bounded distributive lattice and \* is a unary operation satisfying the following conditions for all  $x, y \in L$ :

- $(K_1) (x^*)^* = x;$
- $(K_2) (x \vee y)^* = x^* \wedge y^*;$
- $(K_3)$   $x \wedge x^* \leq y \vee y^*$ .

**Theorem 4.8.**  $(\mathcal{Z}(X), \cap, \cup, *, 0, 1)$  is a Kleene algebra.

*Proof.* It follows from Theorem 4.3 and Proposition 4.6(3).

#### 5. The OML-center of quantum-Wajsberg algebras

Given a quantum-Wajsberg algebra X, we define the OML-center  $\mathcal{O}(X)$  of X, we study its properties, and show that  $\mathcal{O}(X)$  is a subalgebra of X. We prove that  $\mathcal{O}(X)$  is an orthomodular lattice, and the orthomodular lattices form a subvariety of the variety of quantum-Wajsberg algebras. In what follows,  $(X, \to, ^*, 1)$  will be a quantum-Wajsberg algebra, unless otherwise stated.

Denote  $\mathcal{O}(X) = \{x \in X \mid x = x^* \to x\}$ . Obviously  $0, 1 \in \mathcal{O}(X)$ .

**Lemma 5.1.**  $\mathcal{O}(X)$  is closed under \* and  $\rightarrow$ .

Proof. If  $x \in \mathcal{O}(X)$ , then  $x = x^* \to x$ , and by Proposition 2.7(8), we get  $x^* = x \to x^* = (x^*)^* \to x^*$ , hence  $x^* \in \mathcal{O}(X)$ . Let  $x, y \in \mathcal{O}(X)$ , that is  $x = x^* \to x$  and  $y = y^* \to y$ . By Lemma 2.1(9), we have  $(x^* \to y)^* \to (x^* \to y) = (x^* \to x)^* \to (y^* \to y) = x^* \to y$ , thus  $x^* \to y \in \mathcal{O}(X)$ . Finally, from  $x^*, y \in \mathcal{O}(X)$ , we get  $x \to y \in \mathcal{O}(X)$ . Hence  $\mathcal{O}(X)$  is closed under \* and  $\to$ .

Corollary 5.2. The following hold:

- (1)  $\mathcal{O}(X) = \{x \in X \mid x^* = x \to x^*\};$
- (2)  $(\mathcal{O}(X), \rightarrow, 0, 1)$  is a subalgebra of  $(X, \rightarrow, 0, 1)$ ;
- (3)  $\mathcal{O}(X)$  is closed under  $\cap$ ,  $\cup$  and  $\odot$ .

**Proposition 5.3.**  $\mathcal{O}(X) = \{x \in X \mid x^* \cup x = 1\} = \{x \in X \mid x^* \cap x = 0\}.$ 

Proof. If  $x \in \mathcal{O}(X)$ , then  $x = x^* \to x$ , so that  $x^* \uplus x = (x^* \to x) \to x = x \to x = 1$ . Conversely, if  $x^* \uplus x = 1$ , then  $(x^* \to x) \to x = 1$ , that is  $x^* \to x \le x$ . Since by Proposition 2.6(2),  $x \le_Q x^* \to x$ , using Proposition 2.7(4) we get  $x = x^* \to x$ , that is  $x \in \mathcal{O}(X)$ . Similarly  $\mathcal{O}(X) = \{x \in X \mid x^* \cap x = 0\}$ .

**Proposition 5.4.** The following hold for all  $x \in \mathcal{O}(X)$  and  $y \in X$ :

- (1)  $x \to (x \to y) = x \to y$ ;
- $(2) (x \rightarrow y) \rightarrow x = x;$
- $(3) (y \to x)^* \to x = y \to x;$
- (4)  $(y \to x)^* \to (y \to x) = y \to (y \to x)$ .

*Proof.* (1) Using Lemma 2.1(7), we get:  $x \to (x \to y) = (x \to y)^* \to x^* = x \to (y^* \to x^*) = y^* \to (x \to x^*) = y^* \to x^* = x \to y$ .

- (2) It follows by (1), applying Proposition 2.7(8).
- (3) By Lemma 2.1(7),  $(y \to x)^* \to x = y \to (x^* \to x) = y \to x$ .
- (4) Replacing y by  $y^*$  in Lemma 2.1(9) and taking into consideration that  $x^* \to x = x$ , we get  $(x^* \to y^*)^* \to (x^* \to y^*) = x^* \to (y \to y^*)$ , so that  $(y \to x)^* \to (y \to x) = y \to (x^* \to y^*)$ . Hence  $(y \to x)^* \to (y \to x) = y \to (y \to x)$ .

For any  $x, y \in \mathcal{O}(X)$ , define the operations:  $x \uplus_L y = x^* \to y$ ,  $x \cap_L y = x \odot y$  and the relation  $x \leq_L y$  iff  $x^* \to y = y$ . One can easily check that  $x \uplus_L y = (x^* \cap_L y^*)^*$  and  $x \cap_L y = (x^* \uplus_L y^*)^*$ .

# **Proposition 5.5.** The following hold for all $x, y \in \mathcal{O}(X)$ :

- $(1) \leq_L = \leq_{Q|\mathcal{O}(X)};$
- (2)  $x \uplus y \leq_Q x \uplus_L y \text{ and } x \cap_L y \leq_Q x \cap y;$
- (3)  $(x \cup_L y) \rightarrow x^* = x^*$  and  $(x \cap_L y)^* \rightarrow x = x$ ;
- (4)  $(x \uplus_L y)^* \to y = x \uplus_L y \text{ and } (x \cap_L y) \to y^* = (x \cap_L y)^*;$
- (5)  $(\mathcal{O}(X), \cap_L, \cup_L, 0, 1)$  is a bounded lattice.

*Proof.* (1) Let  $x, y \in \mathcal{O}(X)$  such that  $x \leq_Q y$ . It follows that  $y^* \leq_Q x^*$  and  $x^* \to y \leq_Q y^* \to y = y$ . On the other hand,  $y \leq_Q x^* \to y$ , hence  $x^* \to y = y$ , that is  $x \leq_L y$ . Conversely, if  $x \leq_L y$  we have  $x \leq_Q x^* \to y = y$ . Thus  $\leq_L = \leq_{Q|\mathcal{O}(X)}$ .

- (2) Since  $x^* \leq_Q x \to y$ , we have  $(x \to y) \to y \leq_Q x^* \to y$ , that is  $x \uplus y \leq_Q x \uplus_L y$ . Similarly  $x \leq_Q x^* \to y^*$  implies  $(x^* \to y^*) \to y^* \leq_Q x \to y^*$ . Hence  $(x \to y^*)^* \leq_Q x \cap y$ , so that  $x \odot y \leq_Q x \cap y$ , that is  $x \cap_L y \leq_Q x \cap y$ .
- (3) It follows from Proposition 5.4(2), replacing x by  $x^*$  and y by  $y^*$ , respectively.
- (4) Since  $y \in \mathcal{O}(X)$ , by Proposition 5.4(3) we have  $(x \to y)^* \to y = x \to y$ . Replacing x by  $x^*$  we get  $(x \uplus_L y)^* \to y = x \uplus_L y$ , and replacing y by  $y^*$  we have  $(x \cap_L y) \to y^* = (x \cap_L y)^*$ .
- (5) Clearly  $\bigcup_L$  and  $\bigcap_L$  are commutative and idempotent. Moreover, using Lemma 2.1(7) we can easily check that  $\bigcup_L$  and  $\bigcap_L$  are associative. Finally, applying Proposition 5.4(2), we have:

$$x \uplus_L (x \cap_L y) = x^* \to (x \to y^*)^* = (x \to y^*) \to x = x, x \cap_L (x \uplus_L y) = (x \to (x^* \to y)^*)^* = ((x^* \to y) \to x^*)^* = (x^*)^* = x,$$

for all  $x, y \in \mathcal{O}(X)$ , hence  $\bigcup_L$  and  $\bigcap_L$  satisfy the absorption laws. Thus  $(\mathcal{O}(X), \bigcap_L, \bigcup_L, 0, 1)$  is a bounded lattice.

**Corollary 5.6.** The following hold for all  $x, y \in \mathcal{O}(X)$ :

- (1)  $x \leq_Q y$  iff  $y = y \cup_L x$ ;
- $(2) x \uplus y = (x \to y)^* \uplus_L y.$

Proof. (1) 
$$x \leq_Q y$$
 iff  $x \leq_L y$  iff  $y = x^* \to y = y^* \to x = y \uplus_L x$ .  
(2)  $x \uplus y = (x \to y) \to y = ((x \to y)^*)^* \to y = (x \to y)^* \uplus_L y$ .

In what follows, if  $x, y \in \mathcal{O}(X)$ , we will use  $x \leq_Q y$  instead of  $x \leq_L y$ .

**Proposition 5.7.** For any  $x, y \in \mathcal{O}(X)$ ,  $x \cup_L y$  and  $x \cap_L y$  are the l.u.b. and g.l.b. of  $\{x, y\}$ , respectively.

*Proof.* Obviously  $x, y \leq_Q x^* \to y$ , so that  $x \uplus_L y$  is an upper bound of  $\{x, y\}$ . Let  $z \in \mathcal{O}(X)$  be another upper bound of  $\{x, y\}$  in  $\mathcal{O}(X)$ , that is  $x, y \leq_Q z$ . It follows that  $z^* \leq_Q x^*$ , so that  $x^* \to y \leq_Q x^* \to z \leq_Q z^* \to z = z$ . Hence  $x \uplus_L y \leq_Q z$ , that is  $x \uplus_L y$  is the l.u.b.

of  $\{x,y\}$ . Similarly  $x\odot y\leq_Q x,y$ , thus  $x\cap_L y$  is a lower bound of  $\{x,y\}$ . Let  $z\in\mathcal{O}(X)$  be another lower bound of  $\{x,y\}$  in  $\mathcal{O}(X)$ , so that  $z\leq_Q x,y$ . We get  $x^*,y^*\leq_Q z^*$ , so that  $z^*$  is an upper bound of  $\{x^*,y^*\}$ , hence  $x^*\uplus_L y^*\leq_Q z^*$ , that is  $x\to y^*\leq_Q z^*$ . It follows that  $z\leq_Q (x\to y^*)^*=x\odot y=x\cap_L y$ , thus  $x\cap_L y$  is the g.l.b. of  $\{x,y\}$ .

**Definition 5.8.** ([2]) An algebra  $(X, \wedge, \vee, ', 0, 1)$  with two binary, one unary and two nullary operations is an *ortholattice* if it satisfies the following axioms for all  $x, y, z \in X$ :

- $(Q_1)$   $(X, \wedge, \vee, 0, 1)$  is a bounded lattice;
- $(Q_2) \ x \wedge x' = 0 \text{ and } x \vee x' = 1;$
- $(Q_3)^{'}(x \wedge y)' = x' \vee y' \text{ and } (x \vee y)' = x' \wedge y';$
- $(Q_4) (x')' = x.$

An orthomodular lattice is an ortholattice satisfying the following axiom:

 $(Q_5)$   $x \leq y$  implies  $x \vee (x' \wedge y) = y$  (where  $x \leq y$  iff  $x = x \wedge y$ ).

**Theorem 5.9.**  $(\mathcal{O}(X), \cap_L, \cup_L, *, 0, 1)$  is an orthomodular lattice called the *orthomodular center* or *OML-center of* X.

*Proof.* Let X be a quantum-Wajsberg algebra. Using Propositions 5.5, 5.7, 5.3 we can easily check that  $(\mathcal{O}(X), \cap_L, \uplus_L, *, 0, 1)$  is an ortholattice. We show that axiom  $(Q_5)$  is also satisfied. Let  $x, y \in \mathcal{O}(X)$  such that  $x \leq_Q y$ , and we have:  $x \uplus_L (x^* \cap_L y) = x \uplus_L (x^* \circ y) = x \uplus_L (x^* \to y^*)^* = x \uplus_L (y \to x)^* = x^* \to (y \to x)^* = (y \to x) \to x = y \uplus x = y$ , since  $x \leq_Q y$ .

**Theorem 5.10.** If  $(X, \wedge, \vee, ', 0, 1)$  is an orthomodular lattice, then  $(X, \rightarrow, 0, 1)$  is a quantum-Wajsberg algebra, where  $x \rightarrow y = x' \vee y$  for all  $x, y \in X$ .

*Proof.* According to [5, Thm. 2.3.9], every orthomodular lattice  $(X, \wedge, \vee, ', 0, 1)$  determines a QMV algebra by taking  $\oplus$  as the supremum  $\vee$  and  $^*$  as the orthocomplement ', and conversely, if an ortholattice X determines a QMV algebra  $(X, \oplus, ^*, 0, 1)$  taking  $\oplus = \vee$  and  $^* = '$ , then X is orthomodular. By [4, Thm. 5.3], any quantum-MV algebra  $(X, \oplus, ^*, 0, 1)$  is a quantum-Wajsberg algebra  $(X, \to, 0, 1)$ , where  $x \to y = x^* \oplus y$ . It follows that every orthomodular lattice  $(X, \wedge, \vee, ', 0, 1)$  determines a quantum-Wajsberg algebra  $(X, \to, 0, 1)$  with  $x \to y = x^* \oplus y = x^* \vee y$  for all  $x, y \in X$ .

Corollary 5.11.  $(X, \cap_L, \cup_L, *, 0, 1)$  is an orthomodular lattice if and only if  $\mathcal{O}(X) = X$ .

Similarly as [5, Cor. 2.3.13] for the case of QMV algebras, we have the following result.

**Corollary 5.12.** The orthomodular lattices form a subvariety of the variety of quantum-Wajsberg algebras. This subvariety satisfies the condition  $x = x^* \to x$ , or equivalently,  $x^* \uplus x = 1$ , or equivalently,  $x^* \cap x = 0$ .

*Proof.* The equivalence of conditions  $x = x^* \to x$ ,  $x^* \uplus x = 1$  and  $x^* \cap x = 0$  follows from Proposition 5.3. If a quantum-Wajsberg algebra  $(X, \to, 0, 1)$  is an orthomodular lattice with  $x \lor y = x^* \to y$ , than  $x^* \to x = x \lor x = x$ . Conversely, if X satisfies condition  $x^* \to x = x$  for any  $x \in X$ , then  $\mathcal{O}(X) = X$ , hence X is an orthomodular lattice.

**Example 5.13.** Let  $X = \{0, a, b, c, d, 1\}$  and let  $(X, \rightarrow, 0, 1)$  be the involutive BE algebra with  $\rightarrow$  and the corresponding operation  $\cap$  given in the following tables:

$\rightarrow$	0	a	b	c	d	1	$ \  \   \bigcap$	0	a	b	c	d	1
0									0				
a	c	1	1	c	1	1	a	0	a	b	0	d	a
b	d	1	1	1	d	1	b	0	a	b	c	0	b .
c	a	a	1	1	1	1	c	0	0	b	c	d	c
d	b	1	b	1	1	1	d	0	a	0	c	d	d
1	0	a	b	c	d	1	1	0	a	b	c	d	1

Then X is a quantum-Wajsberg algebra and  $\mathcal{Z}(X) = \{0,1\}$ ,  $\mathcal{O}(X) = X$ . Therefore  $(X, \cap_L, \cup_L, *, 0, 1)$  is an orthomodular lattice with  $\cup_L$  and  $\cap_L$  given below.

$\uplus_L$	0	a	b	c	d	1	$\Cap_L$	0	a	b	c	d	1
0							0	0	0	0	0	0	0
a	a	a	1	1	1	1	a	0	a	0	0	0	a
b	b	1	b	1	1	1	b	0	0	b	0	0	b .
c	c	1	1	c	1	1	c	0	0	0	c	0	c
d	d	1	1	1	d	1	d	0	0	0	0	d	d
1	1	1	1	1	1	1	1	0	a	b	c	d	1

As we can see in this example, in general,  $\bigcup_L \neq \bigcup$  and  $\bigcap_L \neq \bigcap$ .

Remark 5.14. In general, the lattice  $(\mathcal{O}(X), \cap_L, \cup_L, 0, 1)$  is not distributive. Indeed, in Example 5.13 we have  $a \cup_L (b \cap_L c) = a \neq 1 = (a \cup_L b) \cap_L (a \cup_L c)$ .

# 6. Concluding remarks and future work

In this paper, we continued the study of quantum-Wajsberg algebras ([4]). We defined the Wajsberg-center and the OML-center of a quantum-Wajsberg algebra  $(X, \to, ^*, 1)$ , proving that the Wajsberg-center is a Wajsberg subalgebra of X, and that it is a distributive sublattice of the poset  $(X, \leq_Q, 0, 1)$  (where  $0 = 1^*$ ). We introduced the notion of quasi-linear quantum-Wajsberg algebras, and we proved that the Wajsberg-center of a quasi-linear quantum-Wajsberg algebra is a linearly ordered Wajsberg algebra. We also proved that the OML-center is an orthomodular lattice, and that the orthomodular lattices form a subvariety of the variety of quantum-Wajsberg algebras.

There are several ways this work can be continued, as follows:

- Introduce and study certain generalizations of quantum-Wajsberg algebras, such as implicative-orthomodular, pre-Wajsberg and meta-Wajsberg algebras.
- Define the implicative-orthomodular lattices as a special subclass of quantum-Wajsberg algebras, and study their properties.
  - Prove an analogue of Foulis-Holland theorem for implicative-orthomodular lattices.
- Study the Baer \*-semigroup associated to an implicative-orthomodular lattice X and its relationship with the Sasaki projections defined on X.
  - Investigate the central lifting property for implicative-orthomodular lattices.

Another direction of research could be the solving of the following open problem.

**Open problem.** Is the variety of quasi-linear quantum-Wajsberg algebras axiomatizable (in the sense of [9])?

## Declaration of interest

The author has no relevant financial or non-financial interests to disclose.

#### Acknowledgement

The author is very grateful to the anonymous referees for their useful remarks and suggestions on the subject that helped improving the presentation.

#### References

- [1] L. Beran, Orthomodular Lattices: Algebraic Approach. Mathematics and its Applications, Springer, Netherland, 1985.
- [2] S. Burris, H. P. Sankappanavar, A course in Universal Algebra, Springer-Verlag, New York, 1981.
- [3] C.C. Chang, Algebraic analysis of many valued logics, Trans. Amer. Math. Soc. 88(1958), 467–490.
- [4] L.C. Ciungu, Quantum-Wajsberg algebras, arxiv.org:2303.16481v2.
- [5] A. Dvurečenskij, S. Pulmannová, New trends in Quantum Structures, Kluwer Academic Publishers, Dordrecht, Ister Science, Bratislava, 2000.
- [6] J.M. Font, A.J. Rodriguez, A. Torrens, Wajsberg algebras, Stochastica 8(1)(1984), 5–31.
- [7] R. Giuntini, Quasilinear QMV algebras, Inter. J. Theor. Phys. 34(1995), 1397–1407.
- [8] R. Giuntini, Quantum MV-algebras, Studia Logica 56(1996), 393-417.
- [9] R. Giuntini, Axiomatizing Quantum MV-algebras, Mathware and Soft Comput. 4(1997), 23–39.
- [10] R. Giuntini, Quantum MV-algebras and commutativity, Inter. J. Theor. Phys. 37(1998), 65–74.
- [11] R. Giuntini, An independent axiomatization of quantum MV-algebras. In: C. Carola, A. Rossi (eds.), The Foundations of Quantum Mechanics, World Scientific, Singapore, 2000, pp. 233–249.
- [12] R. Giuntini, Weakly linear quantum MV-alqebras, Algebra Universalis 53(2005), 45-72.
- [13] R. Giuntini, S. Pulmannová, *Ideals and congruences in effect algebras and QMV-algebras*, Comm. Algebra **28**(2000), 1567–1592.
- [14] S. Gudder, Total extension of effect algebras, Found. Phys. Letters 8(1995), 243–252.
- [15] S.W. Han, X.T. Xu, F. Qin, The unitality of quantum B-algebras, Int. J. Theor. Phys. 57(2018), 1582–1590.
- [16] S. Han, R. Wang, X. Xu, The injective hull of quantum B-algebras, Fuzzy Sets Syst. 369(2019), 114–121.
- [17] Y. Imai, K. Iséki, On axiom systems of propositional calculi. XIV. Proc. Japan Acad. 42, 19–22, (1966)
- [18] A. Iorgulescu, New generalizations of BCI, BCK and Hilbert algebras Part I, J. of Multiple-Valued Logic and Soft Computing 27(4)(2016), 353–406.
- [19] A. Iorgulescu, New generalizations of BCI, BCK and Hilbert algebras Part II, J. of Multiple-Valued Logic and Soft Computing 27(4)(2016), 407–456.
- [20] A. Iorgulescu, Algebras of logic vs. algebras, In Adrian Rezus, editor, Contemporary Logic and Computing, Vol. 1 of Landscapes in Logic, pages 15–258, College Publications, 2020.
- [21] A. Iorgulescu, On quantum-MV algebras Part I: The orthomodular algebras, Sci. Ann. Comput. Sci. 31(2)(2021), 163–221.
- [22] A. Iorgulescu, On quantum-MV algebras Part II: Orthomodular lattices, softlattices and widelattices, Trans. Fuzzy Sets Syst. 1(1)(2022), 1–41.
- [23] A. Iorgulescu, On quantum-MV algebras Part III: The properties (m-Pabs-i) and (WNMm), Sci. Math. Jpn. **35**(e-2022), Article 4 Vol. 86, No. 1, 2023, 49–81.

- [24] A. Iorgulescu, M. Kinyon, Putting quantum-MV algebras on the map, Sci. Math. Jpn. 34(e-2021), Article 10 - Vol. 85, No. 2, 2022, 89–115.
- [25] A. Iorgulescu, BCK algebras versus m-BCK algebras. Foundations, Studies in Logic, Vol. 96, 2022.
- [26] G. Kalmbach, Orthomodular Lattices, Academic Press, London, New York, 1983.
- [27] H.S. Kim, Y.H. Kim, On BE-algebras, Sci. Math. Jpn. 66(2007), 113-116.
- [28] K.H. Kim, Y.H. Yon, Dual BCK-algebra and MV-algebra, Sci. Math. Jpn. 66(2007), 247–254.
- [29] C.J. Mulvey, &, In: Second Topology Conference, Taormina, April 47, 1984, Rend. Circ. Mat. Palermo Suppl. 12(1986), 99–104.
- [30] D. Mundici, MV-algebras are categorically equivalent to bounded commutative BCK-algebras, Math. Japonica, 6(1986), 889–894.
- [31] P. Pták, S. Pulmannová, Orthomodular Structures as Quantum Logics, Veda and Kluwer Acad. Publ., Bratislava and Dordrecht, 1991.
- [32] W. Rump, Quantum B-algebras, Cen. Eur. J. Math. 11(2013), 1881–1899.
- [33] W. Rump, Y.C. Yang, Non-commutative logic algebras and algebraic quantales, Ann. Pure Appl. Logic 165(2014), 759–785.
- [34] W. Rump, The completion of a quantum B-algebra, Cah. Topol. Géom. Différ. Catég. 57(2016), 203–228.
- [35] W. Rump, Quantum B-algebras: their omnipresence in algebraic logic and beyond, Soft Comput. 21(2017), 2521–2529.
- [36] A. Walendziak, On commutative BE-algebras, Sci. Math. Jpn. 69(2009), 281-284.