

Superdiffusive magnetization transport in the XX spin chain with non-local dephasing

Marko Žnidarič

*Department of Physics, Faculty of Mathematics and Physics,
University of Ljubljana, Jadranska 19, SI-1000 Ljubljana, Slovenia*

(Dated: November 14, 2023)

We study recently discussed XX spin chain with non-local dephasing [1] in a steady-state boundary-driven setting, confirming superdiffusive magnetization transport in the thermodynamic limit. The emergence of superdiffusion is rather interesting as the Lindblad operators causing it are a coherent sum of two terms, each of which would separately cause diffusion. One therefore has a quantum phenomenon where a coherent sum of two diffusive terms results in superdiffusion. We also study perturbations of the superdiffusive model, finding that breaking the exact form of dissipators, as well as adding interactions to the XX chain, results in superdiffusion changing into diffusion.

I. INTRODUCTION

Transport is one of the simplest nonequilibrium properties which though is not necessarily easy to address, particularly in one-dimensional quantum lattice models. Namely, in one-dimensional system one can have an interesting interplay between integrability on one hand, which in itself favors ballistic transport where disturbances spread linearly in time, and chaos on the other hand, where one expects diffusion with its square-root growth of disturbances [2]. Different transport types can be distinguished by a dynamical scaling exponent z that tells us how fast the size of a disturbance, say a width σ of a packet, spreads in time, $\sigma \sim t^{1/z}$.

Specific cases with ballistic $z = 1$ as well as diffusive $z = 2$ were known for a long time. It was also known that in quadratic systems, i.e. systems that are non-interacting in a single-particle basis, one can have an intermediate superdiffusive transport with $1 < z < 2$ if one allows for an inhomogeneous Hamiltonian, for instance, a site-dependent potential [3]. An example of such a superdiffusive system is the Fibonacci model [5–7], or a random dimer model [8]. Significant progress has been made in the last decade also for interacting models [2], with realization that one can have superdiffusion also in a homogeneous interacting system. This was first observed numerically in the isotropic Heisenberg spin chain at infinite temperature [9] where $z = 3/2$. By now we have a fairly thorough understanding of why and when such an “interacting” superdiffusion occurs [10–18], see also Ref. [19] for a review, with a microscopic framework being provided by a generalized hydrodynamics [20, 21]. It is limited to a zero-magnetization sector in integrable models with a continuous non-Abelian symmetry. Integrability is required in order to have ballistically propagating quasiparticles, while the non-Abelian symmetry ensures appropriate properties of those quasiparticles (scaling of their velocity and magnetization they carry with their size). Intriguing was also observation [22, 23] of not just the scaling exponent $z = 3/2$ but also of the associated Kardar-Parisi-Zhang (KPZ) [24] scaling functions in a fully coherent (noiseless) quantum system. Superdiffusion in the Heisenberg spin chain has been also observed

experimentally [25–28].

Very recently a surprisingly simple new way of obtaining superdiffusion in quantum lattice models has been revealed [1], namely, rather than using symmetry one can use a multi-site dephasing dissipation to induce superdiffusion in an otherwise free fermionic model (equivalent to the XX spin chain). The fact that the dephasing dissipators acts on multiple sites is crucial; for local dephasing one instead gets diffusion [29]. Superdiffusion comes due to the dephasing strength being zero at some momentum, resulting in a diverging scattering length for those ballistic plane-wave quasiparticles of the XX chain, causing the dynamical scaling exponent $z = 3/2$. One can also get other values of z [30] if one has a higher order zero in the momentum-space dephasing strength, or if the free-fermion dispersion relation has a zero in the velocity. The mechanism of this newly discovered superdiffusion is different than in previously mentioned interacting integrable models as well as in free inhomogeneous systems; the model is translationally invariant and the phenomenon is not limited to one spatial dimension. Ref. [1] presented theoretical arguments explaining superdiffusion and verified its prediction by a direct numerical simulation of time evolution of a fully polarized domain wall. The largest size $L = 256$ was not large enough to really be in the asymptotic regime of long times, however a hydrodynamic approximation with a Wigner function that though can be simulated in the asymptotic regime did agree with the exact numerics.

In the present paper we use a boundary-driven Lindblad setting that allows us to (i) probe much larger systems upto $L = 6000$, thereby confirming asymptotic superdiffusion in an exact lattice model, (ii) probe the role of weak interactions and weak breaking of dephasing dissipation, both resulting in diffusion, and (iii) verify that the superdiffusion in question is a genuine bulk thermodynamic property and is e.g. not particular to a specific initial state (a fully polarized domain wall can be a non-generic initial state in some situations, like e.g., in the XXZ spin chain, with a non-generic transport type being specific to that state).

II. NONEQUILIBRIUM STEADY-STATE SETTING

We will use spin language rather than fermions [1] and study a chain of spin 1/2 particles with the bulk described by the XX spin chain, written in terms of Pauli operators ($\sigma_j^x, \sigma_j^y, \sigma_j^z$ and $\sigma_j^\pm = (\sigma_j^x \pm i\sigma_j^y)/2$) as

$$H = \sum_{j=1}^{L-1} \sigma_j^x \sigma_{j+1}^x + \sigma_j^y \sigma_{j+1}^y. \quad (1)$$

In fermionic language it describes a system of L spinless non-interacting fermions. On top of the Hamiltonian part we also have a bulk dissipation, such that the evolution equation of the density operator $\rho(t)$ is the Lindblad master equation [32, 33],

$$\frac{d\rho}{dt} = i[\rho, H] + \gamma \sum_{j=2}^{L-1} \mathcal{L}_j^{(\text{deph})}(\rho) + \mathcal{L}^{(\text{bath})}(\rho). \quad (2)$$

There are two dissipative parts. The one with $\mathcal{L}_j^{(\text{deph})}$ of strength γ (set to $\gamma = 1$) describes bulk non-local dephasing and is the term responsible for interesting superdiffusive transport. The bath part $\mathcal{L}^{(\text{bath})}$ will act only on the boundary and is there solely to efficiently probe transport properties.

Dephasing superoperator $\mathcal{L}_j^{(\text{deph})}$ will act on few sites surrounding the site j , in our case on 3 neighboring sites $j-1, j$ and $j+1$, and is described by a single Lindblad operator L_j of form $L_j = l_j^\dagger l_j$,

$$\mathcal{L}_j^{(\text{deph})}(\rho) = 2L_j\rho L_j^\dagger - \rho L_j^\dagger L_j - L_j^\dagger L_j \rho, \quad L_j = l_j^\dagger l_j. \quad (3)$$

We will use different forms of l_j , resulting in either superdiffusion or diffusion. Just as an example, taking

$$l_j = \frac{1}{\sqrt{2}}(\sigma_{j-1}^- + Z_{j-1}^{[2]}\sigma_{j+1}^-), \quad (4)$$

where $Z_j^{[r]}$ is a product of σ_k^z on r consecutive sites, starting with the j -th,

$$Z_j^{[r]} = \prod_{k=j}^{j+r-1} \sigma_k^z, \quad (5)$$

will result in superdiffusion. Such dissipation is called a non-local dephasing in analogy with the standard (local) dephasing, where one takes $L_j = \sigma_j^+ \sigma_j^- = (\mathbb{1} + \sigma_j^z)/2$, as it can be thought of as a dephasing acting on quasiparticles delocalized over few sites [1]. It is instructive to write out the Lindblad operator $L_j = l_j^\dagger l_j$; for the above choice (4) we get

$$L_j = \frac{2 + \sigma_{j-1}^z + \sigma_{j+1}^z}{4} + \frac{\sigma_{j-1}^+ Z_{j-1}^{[2]} \sigma_{j+1}^- - \sigma_{j-1}^- Z_{j-1}^{[2]} \sigma_{j+1}^+}{2}. \quad (6)$$

We can see that for real spins, living on sites j , the Lindblad operator is a coherent sum of dephasing and of next-nearest-neighbor hopping. Each of these terms individually is expected to lead to diffusion, for dephasing see Ref. [29], for hopping Ref. [34], but both together, as we shall see, cause superdiffusion. Interestingly, at first sight a benign looking phase term $Z_{j-1}^{[2]} = \sigma_{j-1}^z \sigma_j^z$ is absolutely crucial – leaving it out in l_j (4) will lead to diffusion.

To study transport we couple the first and the last spin to magnetization baths described phenomenologically by the following 4 Lindblad operators,

$$\begin{aligned} \mathcal{L}^{(\text{bath})}(\rho) &= \sum_{k=1}^4 2L'_k \rho L_k'^\dagger - \rho L_k'^\dagger L'_k - L_k'^\dagger L'_k \rho \\ L'_1 &= \sqrt{\Gamma(1+\mu)} \sigma_1^+, \quad L'_2 = \sqrt{\Gamma(1-\mu)} \sigma_1^- \\ L'_3 &= \sqrt{\Gamma(1-\mu)} \sigma_L^+, \quad L'_4 = \sqrt{\Gamma(1+\mu)} \sigma_L^-. \end{aligned} \quad (7)$$

This standard setup [35] can be thought of as an infinite-temperature magnetization driving. The coupling strength Γ is set to $\Gamma = 1$, while the driving parameter μ determines magnetization that the bath is trying to impose on the two boundary spins (μ on σ_1^z and $-\mu$ on σ_L^z). We use small $\mu = 0.1$ throughout the paper, meaning that we are in a linear response regime where all observables relevant for magnetization transport are proportional to μ .

Lindblad equation (2) has a single steady-state solution ρ_∞ . For $\mu = 0$, i.e., no magnetization bias, the steady-state is a trivial infinite-temperature state $\rho_\infty \sim \mathbb{1}$, for nonzero μ though it is a true nonequilibrium steady state (NESS) with nontrivial magnetization profile and a site-independent NESS magnetization current J . An example




Figure 1. NESS magnetization profile for non-local dephasing described by eq.(4), and $\gamma = \Gamma = 1$, $\mu = 0.1$. x is a scaled coordinate along the chain. In the thermodynamic limit magnetization smoothly varies from $+\mu$ at the left edge to $-\mu$ at the right edge.

of a NESS magnetization profile $\text{tr}(\rho_\infty \sigma_j^z)$ for dissipation Eq. (4) is shown in Fig. 1. We can see that the profile is not linear, as one would expect for an ordinary diffusion, suggesting superdiffusion.

The type of transport is most easily inferred from the scaling of the NESS current J with system size L , keeping

driving μ constant [2]. Namely, the current will in general have a power law dependence,

$$J \sim \frac{1}{L^{z-1}}, \quad (8)$$

with a dynamical scaling exponent z characterizing transport type. For ballistic transport one has $z = 1$, an example is the XX chain without dephasing [36], diffusion is characterized by $z = 2$, an example being the XX chain with local dephasing [29], while $1 < z < 2$ indicates superdiffusion.

III. CURRENT SCALING

Let us study the scaling of J with L more in detail. First, because we have a nontrivial dephasing in the bulk, the local current operator is not simply equal to the standard $j_k = 2(\sigma_k^x \sigma_{k+1}^y - \sigma_k^y \sigma_{k+1}^x)$. Writing the continuity equation for the expectation value $z_j = \text{tr}(\rho_\infty \sigma_j^z)$,

$$\frac{dz_k}{dt} = J_{k-1} - J_k, \quad (9)$$

defines the local current operator J_k , which has in the bulk an additional term due to nonzero $\langle \sigma_k^z, \sum_j \mathcal{L}_j^{(\text{deph})}(\rho) \rangle$. For instance, for the l_j in Eq.(4) we get

$$J_k = j_k + \frac{\gamma}{2}(\sigma_{k-1}^z + \sigma_k^z - \sigma_{k+1}^z - \sigma_{k+2}^z). \quad (10)$$

Due to a 3-site action of $\mathcal{L}_j^{(\text{deph})}$ the additional term involves 4 sites surrounding the bond $k - (k + 1)$ across which the current J_k flows. Note that in all cases studied the total magnetization is conserved, $\sum_j \mathcal{L}_j^{(\text{deph})}(\sum_k \sigma_k^z) = 0$.

Crucial for the correct assessment of transport is being able to obtain results for sufficiently large system sizes L . It is not uncommon to make bold incorrect claims of superdiffusion simply due to an insufficiently large L [37]. We will use two different numerical methods to obtain NESS ρ_∞ , and in turn the NESS current $J = \text{tr}(J_k \rho_\infty)$. One is time-evolved-block decimation (TEBD) method [38, 39], where the expansion coefficients of ρ in the Pauli basis are written in terms of a product of matrices – a so-called matrix product operator ansatz. Time evolution by Lindblad equation is then split into small Trotter-Suzuki time steps so that the elementary operation involves two nearest-neighbor spins. Because the dephasing in our case acts on three consecutive sites we write the chain of L spins as a ladder of $L/2$ rungs, so that all operations are indeed nearest-neighbor ones but acting on rungs instead of spins. The price one has to pay is that the local operator space dimension is 4^2 instead of 4. Details of our TEBD implementation for the Lindblad equation can be found in Ref. [40]. The method works for any Hamiltonian, not just for the non-interacting XX

chain, with the efficiency boiling down to the size of matrices required for a given numerical precision.

The second method works when equations for all 2-point observables (2-point in the fermionic language) form a closed set of equations. That is, instead of having to solve a system of size 4^L , one has to deal with a system of L^2 linear equations. The method can be applied to Lindblad operators that are Hermitian and quadratic in fermionic operators, and only for the XX chain Hamiltonian. Namely, for such class of systems one has a set of hierarchical equations, first observed for the XX chain with dephasing [29] and then generalized [34, 41], see also Refs. [42–44]. k -point observables form a closed set of linear equations with an inhomogeneous term coming from lower orders. Equations can therefore be solved order by order, starting with 2-point expectation values. In our spin language those 2-point observables are energy-density like ($r \geq 2$),

$$A_j^{(r)} = \sigma_j^x Z_{j+1}^{[r-2]} \sigma_{j+r-1}^x + \sigma_j^y Z_{j+1}^{[r-2]} \sigma_{j+r-1}^y, \quad (11)$$

while $A_j^{(1)} = -\sigma_j^z$, and current like ($r \geq 2$),

$$B_j^{(r)} = \sigma_j^x Z_{j+1}^{[r-2]} \sigma_{j+r-1}^y - \sigma_j^y Z_{j+1}^{[r-2]} \sigma_{j+r-1}^x. \quad (12)$$

There are in total L^2 such observables. If we put their NESS expectation value in a vector \mathbf{y} , we have to solve a system of linear equations

$$M\mathbf{y} = \mu\mathbf{m}, \quad (13)$$

where a sparse matrix M depends on the dephasing strength γ and the bath coupling strength Γ , while a constant source vector \mathbf{m} comes solely due to bath driving. Both our superdiffusive examples do possess such hierarchical structure of NESS correlations and so Eq.(13) can be used to study large systems. Because of dealing with a finite system, where one has to correctly write equations also at the boundary, construction of M is a bit messy and we give details in Appendix A.

A. Superdiffusion with $z = 3/2$

Let us start with the dissipator already mentioned in Eq.(10), namely $L_j = l_j^\dagger l_j$ with

$$l_j = \frac{1}{\sqrt{2}} (\sigma_{j-1}^- + \sigma_{j-1}^z \sigma_j^z \sigma_{j+1}^-). \quad (14)$$

For such l_j one has a closed set of L^2 linear equations for 2-point observables (11,12), and we study current in the NESS in systems with up to $L = 6000$ spins. Results are shown in Fig. 2, where we can see that the dynamical scaling exponent is indeed $z = \frac{3}{2}$. This confirms theoretical prediction based on Ref. [1] with numerically exact lattice simulation. We can also see in the inset that the convergence of z is rather slow as $\frac{3}{2} - z(L) \sim 1/\sqrt{L}$, which




Figure 2. Scaling of the NESS current J with system's length L for hierarchy-preserving dephasing in Eq.(4), blue circles, showing superdiffusion with $z = 3/2$. Red squares are for dissipation Eq.(15) showing diffusion with $z = 2$. The inset shows convergence of $z - 1$ with system size, with the full curve suggesting a slow $\sim 1/\sqrt{L}$ convergence.

is in line with a slow convergence with time observed in Ref. [1] of a hydrodynamic Wigner function approximation for a unitary evolution of a domain wall initial state.

Considering we are working in spin language it is natural to ask, what would happen if we would use a simpler-looking l_j without the product of two σ_k^z (that are remnants of the Jordan-Wigner transformation). To that end we take

$$l_j = \frac{1}{\sqrt{2}} (\sigma_{j-1}^- + \sigma_{j+1}^-). \quad (15)$$

Current for such l_j stays the same (10). At first sight the difference between Eq.(4) and Eq.(15) is minuscule – sometimes such phase factors are simply neglected when doing Jordan-Wigner transformations as they are believed not to be important. In our case they are crucial. We use full TEBD to calculate the NESS J (full red squares in Fig. 2), as well as an approximation using just 2-point observables. Namely, for such l_j (15) 2-point observables do not form a closed set anymore; one has equations of form $M\mathbf{y} + N\mathbf{w} = \mu\mathbf{m}$, where \mathbf{w} are expectation values of higher point observables. This comes about because L_j is a sum of terms that are quadratic in fermions as well as terms that are quartic. One can therefore get 2-point observables (quadratic) from 4-point ones, for instance $\mathcal{L}_2^{(\text{deph})}(\sigma_1^x \sigma_2^x \sigma_3^z)$ will result also in $\sigma_2^x \sigma_3^x$. Nevertheless, the approximation we make is simply dropping all higher point expectations, i.e. $\mathbf{w} = 0$, and solving resulting equations for 2-point expectation values, see Appendix A 3. Those results are shown with empty red squares in Fig. 2. We can see that the approximation with only 2-point observables works surprisingly good. Both TEBD and 2-point approximation show clear diffusion. Therefore, as soon as one breaks a closed hierarchy of correlations one gets diffusion.




Figure 3. Scaling of the NESS current J with L . Blue circles are for dissipator in Eq. (16) that respects hierarchical structure of correlations, showing superdiffusion with $z = 5/3$, while the red squares are for Eq.(19) without the phase factors and which shows diffusion with $z = 2$. The inset shows convergence of $z - 1$ with system size, with the full curve suggesting $\sim 1/\sqrt{L}$ asymptotics.

B. Superdiffusion with $z = 5/3$

It was predicted in Ref. [1] that the exponent is not always $z = 3/2$. It depends on the order of a zero at k_0 in the momentum-space dephasing strength, as well as on special points k_0 where the velocity of free quasiparticles might be zero. An example of such higher order zero is dephasing dissipators $L_j = l_j^\dagger l_j$ with

$$l_j = \frac{1}{\sqrt{6}} (\sigma_{j-1}^- - 2Z_{j-1}^{[1]}\sigma_j^- + Z_{j-1}^{[2]}\sigma_{j+1}^-), \quad (16)$$

which is a particular case of a more general

$$l_j = \frac{1}{\sqrt{2+a^2}} (\sigma_{j-1}^- - aZ_{j-1}^{[1]}\sigma_j^- + Z_{j-1}^{[2]}\sigma_{j+1}^-), \quad (17)$$

studied in Ref. [1]. For $-2 < a < 2$ one expects $z = \frac{3}{2}$ while the chosen $a = 2$ is marginal with a 2nd order zero and prediction [1] that the dynamical scaling exponent is $z = 5/3$. We again calculate the NESS and the scaling of current, which in this case (16) is

$$J_k = j_k + \frac{\gamma}{18} (\sigma_{k-1}^z + 9\sigma_k^z - 9\sigma_{k+1}^z - \sigma_{k+2}^z) + \frac{\gamma}{9} (A_{k-1}^{(2)} - A_{k+1}^{(2)}) + \frac{\gamma}{9} (A_{k-1}^{(3)} - A_k^{(3)}). \quad (18)$$

The dissipator (16) preserves the 2-point expectations and we can study large systems (Appendix A 2). In Fig. 3 we show results, demonstrating clear convergence to theoretical prediction. Let us also not that the additional terms in the current expression in Eq.(16) as well as in Eq.(4) are all differences of operators on neighboring sites. Because the steady state expectations are continuous in the spatial index k (see Fig.1) they all scale




Figure 4. Interactions cause superdiffusion to go into diffusion. Frame (a) shows dephasing in Eq. (4), and (b) dephasing in Eq. (16). Blue circles are the non-interacting case, i.e., the same data as in Figs. 2 and 3, while red triangles are for the XXZ chain (21) with interaction strength $\Delta = 0.2$.

as $\sim 1/L$, and therefore in the thermodynamic limit for superdiffusion one has $J_k \approx j_k$.

Next, we check what happens if we remove the phase factors in the above l_j , that is, we take

$$l_j = \frac{1}{\sqrt{6}} (\sigma_{j-1}^- - 2\sigma_j^- + \sigma_{j+1}^-). \quad (19)$$

The corresponding current operator is

$$J_k = j_k + \frac{\gamma}{18} (\sigma_{k-1}^z + 9\sigma_k^z - 9\sigma_{k+1}^z - \sigma_{k+2}^z) + \frac{\gamma}{9} (A_{k-1}^{(3)} - A_k^{(3)}). \quad (20)$$

Results are shown in Fig. 3, with full red squares for TEBD simulations, and empty red squares using a 2-point correlation function approximation (Appendix A 4), similarly as in the previous subsection for $z = 3/2$. Again, we can see that already this subtle change leads to diffusion.

C. Interactions

We have seen that while one does get superdiffusion for a whole class of dissipators parameterized by a (17), superdiffusion goes away if we remove product of σ_j^z in the definition of l_j . In this subsection we test what happens in we keep the form of l_j but add interactions to the Hamiltonian. To this end we study the XXZ chain,

$$H = \sum_{j=1}^{L-1} \sigma_j^x \sigma_{j+1}^x + \sigma_j^y \sigma_{j+1}^y + \Delta \sigma_j^z \sigma_{j+1}^z, \quad (21)$$

where Δ represents interaction. We use TEBD to get the NESS with which we can go upto $L = 1024$ with rather modest matrix sizes. As we can see in Fig. 4 we obtain diffusion for both $z = 3/2$ dephasing in Eq. (4), and for $z = 5/3$ dephasing in Eq. (16) already for relatively small interaction $\Delta = 0.1$.

While one might jump to a conclusion that this is expected and that superdiffusion is only a property of the

XX chain and the specific form of non-local dephasing, upon reflection things are not that clear. Namely, one can heuristically understand the emergent superdiffusion in the XX chain in the following way [1]: looking at l_j (17) in momentum space, one finds that such non-local l_j results in a momentum dephasing strength that depends on the momentum k . This dephasing strength can in particular have a zero at some k_0 , resulting in a diverging scattering length of free (quasi)particles at that k_0 . Superdiffusion therefore emerges from a measure zero of non-dephasing ballistic quasiparticles. Following this explanation one could argue that because the XXZ chain is integrable, and as such also harbors ballistic quasiparticles, the same phenomenon should be possible. The important difference compared to the XX chain is that the transformation to quasiparticles is not a simple Fourier transformation and perhaps one would have to construct a dephasing that would be zero for those non-plane wave quasiparticles. On the other hand, for the XXZ model one does not have a closed hierarchy of correlations [45], which seem to be important to get superdiffusion – breaking that, as we have seen, immediately leads to diffusion, even in the XX model. Therefore more studies are needed to clarify the generality or speciality of the proposed superdiffusion scenario due to non-local dephasing.

IV. CONCLUSION

We have demonstrated superdiffusive magnetization transport in the XX spin chain in the presence of non-local dephasing dissipation in the bulk. The effect is simple and interesting: one can view it as the emergence of superdiffusion out of a coherent sum of two diffusive contributions in Lindblad operators. It is different than other known cases of superdiffusion, for instance the one in integrable models with a non-Abelian symmetry, or in inhomogeneous non-interacting systems. We have not touched upon superdiffusive classical systems, however, what we can say is that it seems to be qualitatively different than the superdiffusion observed in the stochastic momentum exchange model [46, 47] where the effect changes with dimension.

While a number of questions has been answered, many remain, and some new arose. For instance, during TEBD simulations we have observed that the required size of matrices can be very small. In other words, the operator Schmidt spectrum of the NESS seems to decay quickly. That seems to be the case for both superdiffusive, and to a lesser extent also for diffusive cases studied. One question is can any of the superdiffusive NESSs be written in a matrix product operator form with a low-rank matrices? The fact that the numerical rank is small is perhaps related to two known similar cases of low-rank NESS: for the XX chain and our boundary driving and without dephasing the ballistic NESS requires matrices of size 4 (independent of L) [36], while in the presence of local dephasing the same holds in the leading order in

elements of $R^{(11)}$ are $R_{j,j}^{(11)} = (1, \frac{3}{2}, 2, \dots, 2, \frac{3}{2}, 1)$, and $R_{j,j+1}^{(11c)} = R_{j+1,j}^{(11c)} = \frac{1}{2}$.

The 2-diagonal, $r = 2$, is given by $\tilde{\mathbf{c}}_2 = R^{(24)}\mathbf{c}_4 + \mathbf{c}_2 + R^{(22c)}\mathbf{c}_2^*$, with $(L-2)$ -dimensional $R_{j,j}^{(24)} = R_{k+2,k}^{(24)} = \frac{1}{2}$, and $R_{j,j}^{(22c)} = (-\frac{1}{2}, -\frac{1}{2}, 0, \dots, 0, -\frac{1}{2}, -\frac{1}{2})$.

Matrix elements of \tilde{C} below the diagonal are determined from those above by hermiticity, $\tilde{C}^\dagger = \tilde{C}$.

2. Dephasing with $z = 5/3$

Here l_j is given by Eq.(16), that is

$$l_j = \frac{1}{\sqrt{6}}(\sigma_{j-1}^- - 2\sigma_{j-1}^z\sigma_j^- + Z_{j-1}^{[2]}\sigma_{j+1}^-). \quad (\text{A7})$$

Matrix elements of \tilde{C} on all r -diagonals with $|r| \geq 3$ can be again expressed as $\tilde{C} = LC + CL$, with nonzero elements of L being $L_{k,k} = \frac{1}{6}(1, 5, 6, \dots, 6, 5, 1)$, $L_{k,k+1} = L_{k+1,k} = -\frac{1}{3}(1, 2, \dots, 2, 1)$, $L_{k,k+2} = L_{k+2,k} = \frac{1}{6}$.

The main diagonal is instead equal to $\tilde{\mathbf{c}}_0 = R^{(00)}\mathbf{c}_0 + R^{(01)}(\mathbf{c}_1 + \mathbf{c}_1^*)/2 + R^{(02)}(\mathbf{c}_2 + \mathbf{c}_2^*)/2$, with nonzero elements of $R^{(00)}$ being $R_{j,j}^{(00)} = -\frac{1}{18}(5, 13, 18, \dots, 18, 13, 5)$, $R_{j+1,j}^{(00)} = R_{j,j+1}^{(00)} = \frac{2}{9}(1, 2, \dots, 2, 1)$, and $R_{j+2,j}^{(00)} = R_{j,j+2}^{(00)} = \frac{1}{18}$. Nonzero elements of L -dimensional $R^{(01)}$ are $R_{j,j}^{(01)} = \frac{2}{9}(2, 1, \dots, 1, -1, 1)$, $R_{j,j+1}^{(01)} = -\frac{2}{9}$, $R_{j+1,j}^{(01)} = \frac{2}{9}(-1, 1, \dots, 1, 2)$, and $R_{j+2,j}^{(01)} = -\frac{2}{9}$. Nonzero elements of L -dimensional $R^{(02)}$ are $R_{j,j}^{(02)} = -\frac{2}{9}$, $R_{j+1,j}^{(02)} = \frac{4}{9}$, and $R_{j+2,j}^{(02)} = -\frac{2}{9}$.

The 1-diagonal is $\tilde{\mathbf{c}}_1 = R^{(10)}\mathbf{c}_0 + R^{(11r)}\mathbf{c}_1 + R^{(11c)}\mathbf{c}_1^* + R^{(12r)}\mathbf{c}_2 + R^{(12c)}\mathbf{c}_2^* + R^{(13)}\mathbf{c}_3$, with nonzero $R_{j,j}^{(13)} = R_{k+2,k}^{(13)} = \frac{1}{6}$ of a $(L-1)$ -dimensional $R^{(13)}$. Nonzero elements of $(L-1)$ -dimensional $R^{(12r)}$ are $R_{j,j}^{(12r)} = -\frac{1}{9}(5, \dots, 5, 2, 5)$ and $R_{j+1,j}^{(12r)} = -\frac{1}{9}(2, 5, \dots, 5)$, while nonzero elements of $R^{(12c)}$ are $R_{j,j}^{(12c)} = R_{j+1,j}^{(12c)} = \frac{1}{9}$. Nonzero elements of $(L-1)$ -dimensional $R^{(11r)}$ are $R_{j,j}^{(11r)} = \frac{1}{18}(14, 25, 28, \dots, 28, 25, 14)$, $R_{j,j+1}^{(11r)} = R_{j+1,j}^{(11r)} = \frac{2}{9}$, while of $R^{(11c)}$ are $R_{j,j}^{(11c)} = -\frac{2}{9}(1, 2, \dots, 2, 1)$ and $R_{j,j+1}^{(11c)} = R_{j+1,j}^{(11c)} = -\frac{1}{18}$. Finally, nonzero elements of L -dimensional $R^{(10)}$ are $R_{j,j}^{(10)} = -\frac{1}{9}(2, 1, \dots, 1)$, $R_{j,j}^{(10)} = -\frac{1}{9}(2, 1, \dots, 1, -1, 1)$, $R_{j,j+1}^{(10)} = -\frac{1}{9}(-1, 1, \dots, 1, 2)$ and $R_{j,j+2}^{(10)} = R_{j+1,j}^{(10)} = \frac{1}{9}$ (in $R^{(10)}\mathbf{c}_0$ only the first $(L-1)$ components go into $\tilde{\mathbf{c}}_1$).

The 2-diagonal is $\tilde{\mathbf{c}}_2 = R^{(20)}\mathbf{c}_0 + R^{(21r)}\mathbf{c}_1 + R^{(21c)}\mathbf{c}_1^* + R^{(22r)}\mathbf{c}_2 - \frac{1}{18}\mathbf{c}_2^* + R^{(23)}\mathbf{c}_3 + R^{(24)}\mathbf{c}_4$. Nonzero elements of $(L-2)$ -dimensional $R^{(22r)}$ are $R_{j,j}^{(22r)} = \frac{2}{9}(5, 8, \dots, 8, 5)$. Nonzero elements of $(L-1)$ -dimensional $R^{(21r)}$ are $R_{j,j}^{(21r)} = -\frac{1}{9}(5, 5, \dots, 5, 2, 5)$ and $R_{j,j+1}^{(21r)} = -\frac{1}{9}(2, 5, \dots, 5)$, while $R_{j,j}^{(21c)} = R_{k,k+1}^{(21c)} = \frac{1}{9}$ (only the

first $(L-2)$ components of $R^{(21r)}\mathbf{c}_1 + R^{(21c)}\mathbf{c}_1^*$ matter). Nonzero elements of $(L-2)$ -dimensional $R^{(24)}$ are $R_{j,j}^{(24)} = R_{k+2,k}^{(24)} = \frac{1}{6}$, while of the same-sized $R^{(23)}$ are $R_{j,j}^{(23)} = -\frac{1}{3}(2, 2, \dots, 2, 1, 2)$ and $R_{j+1,j}^{(23)} = -\frac{1}{3}(1, 2, \dots, 2)$. Finally, nonzero elements of L -dimensional $R^{(20)}$ are $R_{j,j}^{(20)} = -\frac{2}{9}$ and $R_{j,j+1}^{(20)} = R_{j+1,j}^{(20)} = \frac{1}{9}$ (middle $(L-2)$ components of $R^{(20)}\mathbf{c}_0$ is what goes into $\tilde{\mathbf{c}}_2$).

3. Perturbed $z = 3/2$ dephasing

The operators l_j are (15)

$$l_j = \frac{1}{\sqrt{2}}(\sigma_{j-1}^- + \sigma_{j+1}^-). \quad (\text{A8})$$

Remember that in this case one does not have a closed set of equations for 2-point observables. Nevertheless, as explained, dropping higher order expectation values in equations for 2-point functions gives a good approximation (red squares in Fig.2). Elements of $\tilde{C}_{j,j+r}$ with $|r| \geq 3$ are equal to corresponding matrix elements of $LC + CL$ with diagonal $L_{k,k} = \frac{1}{2}(1, 2, 3, \dots, 3, 2, 1)$.

The main diagonal is instead $\tilde{\mathbf{c}}_0 = R^{(0)}\mathbf{c}_0$, with nonzero $R_{j,j}^{(0)} = -\frac{1}{2}(1, 1, 2, \dots, 2, 1, 1)$, $R_{j,j+2}^{(0)} = R_{j+2,j}^{(0)} = \frac{1}{2}$. The 1-diagonal is $\tilde{\mathbf{c}}_1 = R^{(1)}\mathbf{c}_1$ with $R_{j,j}^{(1)} = \frac{1}{2}(2, 3, 4, \dots, 4, 3, 2)$. The 2-diagonal is given by $\tilde{\mathbf{c}}_2 = R^{(2)}\mathbf{c}_2 - \frac{1}{2}\mathbf{c}_2^*$, with $R_{j,j}^{(2)} = \frac{1}{2}(3, 4, \dots, 4, 3)$.

4. Perturbed $z = 5/3$ dephasing

In this case the operators l_j are (19)

$$l_j = \frac{1}{\sqrt{6}}(\sigma_{j-1}^- - 2\sigma_j^- + \sigma_{j+1}^-). \quad (\text{A9})$$

Neglecting higher order correlations, elements of $\tilde{C}_{j,j+r}$ with $|r| \geq 3$ are equal to the matrix elements of $LC + CL$, with $L_{k,k} = \frac{1}{18}(3, 16, 19, \dots, 19, 16, 3)$, $L_{k,k+1} = L_{k+1,k} = \frac{1}{18}(5, 10, \dots, 10, 5)$, $L_{k,k+2} = L_{k+2,k} = \frac{1}{9}$.

Elements $\tilde{C}_{j,j+r}$ with $r = 0, 1, 2$ must instead be given separately. The main diagonal is $\tilde{\mathbf{c}}_0 = R^{(0)}\mathbf{c}_0 + R^{(02)}\mathbf{c}'_2$, where $\mathbf{c}'_2 = (a_1^{(1)}, (\mathbf{c}_2 + \mathbf{c}_2^*)/2, a_{L-1}^{(1)})$, and nonzero elements $R_{j,j}^{(0)} = -\frac{1}{18}(5, 13, 18, \dots, 18, 13, 5)$, $R_{j,j+1}^{(0)} = R_{j+1,j}^{(0)} = \frac{2}{9}(1, 2, \dots, 2, 1)$, $R_{j,j+2}^{(0)} = R_{j+2,j}^{(0)} = \frac{1}{18}$, as well as $R_{j,j}^{(02)} = \frac{1}{9}(-3, 4, \dots, 4, -3)$, $R_{j,j+1}^{(02)} = -\frac{1}{9}(2, \dots, 2, -3)$ and $R_{j+1,j}^{(02)} = -\frac{1}{9}(-3, 2, \dots, 2)$.

The 1-diagonal is $\tilde{\mathbf{c}}_1 = R^{(12)}\mathbf{c}_2 + R^{(13)}\mathbf{c}_3 + R^{(11r)}\mathbf{c}_1 + R^{(11c)}\mathbf{c}_1^*$. Nonzero elements of $(L-1)$ -dimensional $R^{(13)}$ are $R_{j,j}^{(13)} = R_{k+2,k}^{(13)} = \frac{1}{9}$. Nonzero elements of $(L-1)$ -dimensional $R^{(12)}$ are $R_{j,j}^{(12)} = \frac{1}{9}(4, \dots, 4, \frac{3}{2}, 4)$ and

$R_{j+1,j}^{(12)} = \frac{1}{9}(\frac{3}{2}, 4, \dots, 4)$. Nonzero elements of $(L-1)$ -dimensional $R^{(11r)}$ are $R_{j,j+1}^{(11r)} = R_{j+1,j}^{(11r)} = -\frac{2}{9}$ and $R_{j,j}^{(11r)} = \frac{1}{18}(14, 25, 28, \dots, 28, 25, 14)$, while for $R^{(11c)}$ we have $R_{j,j}^{(11c)} = -\frac{2}{9}(1, 2, \dots, 2, 1)$ and $R_{j,j+1}^{(11c)} = R_{j+1,j}^{(11c)} = \frac{1}{9}$.

The 2-diagonal is $\tilde{\mathbf{c}}_2 = R^{(20)}\mathbf{c}_0 + R^{(21)}\mathbf{c}_1 + R^{(22r)}\mathbf{c}_2 - \frac{1}{18}\mathbf{c}_2^* + R^{(23)}\mathbf{c}_3 + R^{(24)}\mathbf{c}_4$. Nonzero elements of $(L-2)$ -dimensional $R^{(24)}$ are $R_{j,j}^{(24)} = R_{j+2,j}^{(24)} = \frac{1}{9}$. Nonzero

elements of $(L-2)$ -dimensional $R^{(23)}$ are $R_{j,j}^{(23)} = \frac{5}{9}(1, \dots, 1, \frac{1}{2}, 1)$ and $R_{j+1,j}^{(23)} = \frac{5}{9}(\frac{1}{2}, 1, \dots, 1)$. Nonzero elements of $(L-2)$ -dimensional $R^{(22r)}$ are $R_{j,j}^{(22r)} = (\frac{7}{6}, \frac{17}{9}, \dots, \frac{17}{9}, \frac{7}{6})$. Nonzero elements of $R^{(21)}$ are $R_{j,j}^{(21)} = \frac{4}{9}(1, \dots, 1, \frac{3}{8}, 1)$ and $R_{j,j+1}^{(21)} = \frac{4}{9}(\frac{3}{8}, 1, \dots, 1)$. Finally, nonzero elements of $R^{(20)}$ are $R_{j,j}^{(20)} = -\frac{2}{9}$ and $R_{j,j+1}^{(20)} = R_{j+1,j}^{(20)} = \frac{1}{9}$.

-
- [1] Y. Wang, C. Fang, and J. Ren, *Superdiffusive transport in quasi-particle dephasing models*, arXiv:2310.03069 (2023).
- [2] B. Bertini et al., *Finite-temperature transport in one-dimensional quantum lattice models*, Rev. Mod. Phys. **93**, 025003 (2021).
- [3] In inhomogeneous quadratic systems one can in fact construct an arbitrary transport type, see e.g. Ref. [4].
- [4] G. Mantica, *Quantum intermittency in almost-periodic lattice systems derived from their spectral properties*, Physica D **103**, 576 (1997).
- [5] M. Kohmoto, L. P. Kadanoff, and C. Tang, *Localization problem in one dimension: mapping and escape*, Phys. Rev. Lett. **50**, 1870 (1983).
- [6] S. Ostlund, R. Pandit, D. Rand, H. J. Schellnhuber, and E. D. Siggia, *One-dimensional Schrödinger equation with an almost periodic potential*, Phys. Rev. Lett. **50**, 1873 (1983).
- [7] H. Hiramoto and S. Abe, *Dynamics of an electron in quasiperiodic systems. I. Fibonacci model*, J. Phys. Soc. Japan **57**, 230 (1988).
- [8] D. H. Dunlap, H.-L. Wu, and P. W. Phillips, *Absence of localization in a random-dimer model*, Phys. Rev. Lett. **65**, 88 (1990).
- [9] M. Žnidarič, *Spin transport in a one-dimensional anisotropic Heisenberg model*, Phys. Rev. Lett. **106**, 220601 (2011).
- [10] S. Gopalakrishnan and R. Vasseur, *Kinetic theory of spin diffusion and superdiffusion in XXZ spin chains*, Phys. Rev. Lett. **122**, 127202 (2019).
- [11] M. Dupont and J. E. Moore, *Universal spin dynamics in infinite-temperature one-dimensional quantum magnets*, Phys. Rev. B **101**, 121106(R) (2020).
- [12] V. B. Bulchandani, *Kardar-Parisi-Zhang universality from soft gauge modes*, Phys. Rev. B **101**, 041411(R) (2020).
- [13] J. De Nardis, S. Gopalakrishnan, E. Ilievski, and R. Vasseur, *Superdiffusion from emergent classical solitons in quantum spin chains*, Phys. Rev. Lett. **125**, 070601 (2020).
- [14] E. Ilievski, J. De Nardis, S. Gopalakrishnan, R. Vasseur, and B. Ware, *Superuniversality of superdiffusion*, Phys. Rev. X **11**, 031023 (2021).
- [15] P. Glorioso et al., *Hydrodynamics in lattice models with continuous non-Abelian symmetries*, SciPost Phys. **10**, 015 (2021).
- [16] P. W. Claeys, A. Lamacraft, and J. Herzog-Arbeitman, *Absence of superdiffusion in certain random spin models*, Phys. Rev. Lett. **128**, 246603 (2022).
- [17] Ž. Krajinik and T. Prosen, *Kardar-Parisi-Zhang physics in integrable rotationally symmetric dynamics on discrete space-time lattice*, J. Stat. Phys. **179**, 110 (2020).
- [18] M. Dupont, N. E. Sherman, and J. E. Moore, *Spatiotemporal crossover between low- and high-temperature dynamical regimes in the quantum Heisenberg magnet*, Phys. Rev. Lett. **127**, 107201 (2021).
- [19] V. B. Bulchandani, S. Gopalakrishnan and E. Ilievski, *Superdiffusion in spin chains*, J. Stat. Mech. **2021**, 084001 (2021).
- [20] O. A. Castro-Alvaredo, B. Doyon, and T. Yoshimura, *Emergent hydrodynamics in integrable quantum systems out of equilibrium*, Phys. Rev. X **6**, 041065 (2016).
- [21] B. Bertini, M. Collura, J. De Nardis, and M. Fagotti, *Transport in out-of-equilibrium XXZ chains: Exact profiles of charges and currents*, Phys. Rev. Lett. **117**, 207201 (2016).
- [22] M. Ljubotina, M. Žnidarič, and T. Prosen, *Kardar-Parisi-Zhang physics in the quantum Heisenberg magnet*, Phys. Rev. Lett. **122**, 210602 (2019).
- [23] F. Weiner, P. Schmittecker, S. Bera, and F. Evers, *High-temperature spin dynamics in the Heisenberg chain: Magnon propagation and emerging Kardar-Parisi-Zhang scaling in the zero-magnetization limit*, Phys. Rev. B **101**, 045115 (2020).
- [24] M. Kardar, G. Parisi, Y.-C. Zhang, *Dynamic scaling of growing interfaces*, Phys. Rev. Lett. **56**, 889 (1986).
- [25] P. N. Jepsen et al., *Spin transport in a tunable Heisenberg model realized with ultracold atoms*, Nature **588**, 403 (2020).
- [26] A. Scheie et al., *Detection of Kardar-Parisi-Zhang hydrodynamics in a quantum Heisenberg spin-1/2 chain*, Nature Physics **17**, 726 (2021).
- [27] D. Wei et al., *Quantum gas microscopy of Kardar-Parisi-Zhang superdiffusion*, Science **376**, 716 (2022).
- [28] N. Keenan et al., *Evidence of Kardar-Parisi-Zhang scaling on a digital quantum simulator*, npj Quantum Inf. **9**, 72 (2023).
- [29] M. Žnidarič, *Exact solution for a diffusive nonequilibrium steady state of an open quantum chain*, J. Stat. Mech. **2010**, L05002 (2010).
- [30] Predicted exponents do not necessarily follow Fibonacci sequence [31].
- [31] V. Popkov, A. Schadschneider, J. Schmidt, and G. M. Schütz, *Fibonacci family of dynamical universality classes*, PNAS **112**, 12645 (2015).
- [32] V. Gorini, A. Kossakowski, and E. C. G. Sudarshan, *Completely positive dynamical semigroups of N-level systems*, J. Math. Phys. **17**, 821 (1976).

- [33] G. Lindblad, *On the generators of quantum dynamical semigroups*, Commun. Math. Phys. **48**, 119 (1976).
- [34] V. Eisler, *Crossover between ballistic and diffusive transport: the quantum exclusion process*, J. Stat. Mech. **2011**, P06007 (2011).
- [35] G. T. Landi, D. Poletti, and G. Schaller, *Nonequilibrium boundary-driven quantum systems: Models, methods and properties*, Rev. Mod. Phys. **94**, 045006 (2022).
- [36] M. Žnidarič, *A matrix product solution for a nonequilibrium steady state of an XX chain*, J. Phys. A **43**, 415004 (2010).
- [37] M. Žnidarič, *Comment on “Nonequilibrium steady state phases of the interacting Aubry-André-Harper model”*, Phys. Rev. B **103**, 237101 (2021).
- [38] U. Schollwöck, *The density-matrix renormalization group in the age of matrix product states*, Annals of Physics **326**, 96 (2011).
- [39] M. Zwolak and G. Vidal, *Mixed-state dynamics in one-dimensional quantum lattice systems: A time-dependent superoperator renormalization algorithm*, Phys. Rev. Lett. **93**, 207205 (2004).
- [40] M. Žnidarič, *Dephasing-induced diffusive transport in the anisotropic Heisenberg model*, New J. Phys. **12**, 043001 (2010).
- [41] B. Žunkovič, *Closed hierarchy of correlations in Markovian open quantum systems*, New J. Phys. **16**, 013042 (2014).
- [42] K. Temme, M. M. Wolf, and F. Verstraete, *Stochastic exclusion processes versus coherent transport*, New J. Phys. **14**, 075004 (2012).
- [43] B. Horstmann, J. I. Cirac, and G. Giedke, *Noise-driven dynamics and phase transitions in fermionic systems*, Phys. Rev. A **87**, 012108 (2013).
- [44] X. Turkeshi and M. Schiro, *Diffusion and thermalization in a boundary-driven dephasing model*, Phys. Rev. B **104**, 144301 (2021).
- [45] It is worth mentioning that the closeness of correlations is not always limited solely to noninteracting models, see Ref. [41].
- [46] G. Basile, C. Bernardin, and S. Olla, *Momentum conserving model with anomalous thermal conductivity in low dimensional systems*, Phys. Rev. Lett. **96**, 204303 (2006).
- [47] S. Lepri, C. Mejia-Monasterio, and A. Politi, *A stochastic model of anomalous heat transport: analytical solution of the steady state*, J. Phys. A **42**, 025001 (2009).
- [48] M. Žnidarič, *Solvable quantum nonequilibrium model exhibiting a phase transition and a matrix product representation*, Phys. Rev. E **83**, 011108 (2011).
- [49] M. Horvat and M. Žnidarič, *Eur. Phys. J. B* **86**, 67 (2013).
- [50] M. de Leeuw, C. Paletta, and B. Pozsgay, *Constructing integrable Lindblad superoperators*, Phys. Rev. Lett. **126**, 240403 (2021).