A SIGNED *e*-EXPANSION OF THE CHROMATIC QUASISYMMETRIC FUNCTION

FOSTER TOM

ABSTRACT. We prove a new signed elementary symmetric function expansion of the chromatic quasisymmetric function of any natural unit interval graph. We then use a signreversing involution to prove a new combinatorial formula for K-chains, which are graphs formed by joining cliques at single vertices. This formula immediately implies *e*-positivity and *e*-unimodality for K-chains. We also prove a version of our signed *e*-expansion for arbitrary graphs.

1. INTRODUCTION

In 1995, Stanley [49] introduced the chromatic symmetric function $X_G(\mathbf{x})$, a generalization of the chromatic polynomial of a graph G, and famously conjectured with Stembridge [49, 53] that $X_G(\mathbf{x})$ is *e*-positive whenever G is the incomparability graph of a $(\mathbf{3} + \mathbf{1})$ -free partially ordered set. Guay-Paquet [32] showed that it suffices to prove *e*-positivity for the smaller class of natural unit interval graphs. Shareshian and Wachs [45] defined a further generalization called the chromatic quasisymmetric function $X_G(\mathbf{x};q)$, showed that if G is a natural unit interval graph, then $X_G(\mathbf{x};q)$ is in fact symmetric; and conjectured that it is *e*-positive and *e*-unimodal for such graphs. This notorious problem has received considerable attention. Many authors proved *e*-positivity for certain families of graphs [7, 13, 15, 17, 22, 27, 28, 29, 36, 39, 40, 54, 55, 56], studied related positivity problems [9, 10, 20, 21, 23, 24, 25, 26, 30, 32, 42, 47], and explored alternative generalizations of the chromatic symmetric function [1, 6, 8, 18, 19, 31, 51]. The Stanley–Stembridge conjecture is also closely related to positivity of immanants of Jacobi–Trudi matrices [53], the cohomology of regular semisimple Hessenberg varieties [2, 11, 14, 33, 35, 46], and characters of Hecke algebras evaluated at Kazhdan–Lusztig basis elements [16, 48].

²⁰²⁰ Mathematics Subject Classification. Primary 05E05; Secondary 05E10, 05C15.

Key words and phrases. chromatic quasisymmetric function, elementary symmetric function, natural unit interval graph, proper colouring, Shareshian–Wachs conjecture, Stanley–Stembridge conjecture.

Department of Mathematics, MIT.

In this paper, we prove a new signed elementary symmetric function expansion of the chromatic quasisymmetric function of any natural unit interval graph, in terms of objects called *forest triples*. We then use a sign-reversing involution on forest triples to prove a new combinatorial e-expansion for K-chains, which are graphs formed by joining a sequence of cliques at single vertices.

In Section 2, we introduce chromatic quasisymmetric functions and natural unit interval graphs. In Section 3, we prove our signed formula (Theorem 3.4)

(1.1)
$$X_G(\boldsymbol{x};q) = \sum_{\mathcal{F} \in \mathrm{FT}(G)} \mathrm{sign}(\mathcal{F}) q^{\mathrm{weight}(\mathcal{F})} e_{\mathrm{type}(\mathcal{F})}.$$

In Section 4, we use a sign-reversing involution on forest triples to prove the explicit combinatorial e-expansion (Corollary 4.18) for a generalization of K-chains, namely

(1.2)
$$X_{K_{\gamma}^{\epsilon}}(\boldsymbol{x};q) = [\gamma_1 - 2]_q! \cdots [\gamma_{\ell} - 2]_q! \sum_{\alpha \in A_{\gamma}^{\epsilon}} [\alpha_1]_q \prod_{i=2}^{\ell+1} q^{m_i} [|\alpha_i - (\gamma_{i-1} - \epsilon_{i-1} - 1)|]_q e_{\operatorname{sort}(\alpha)}.$$

This formula implies that $X_{K_{\gamma}^{\epsilon}}(\boldsymbol{x};q)$ is *e*-positive and *e*-unimodal (Corollary 4.20). In Section 5, we propose some new directions of study for the Stanley–Stembridge conjecture and we prove a version of our forest triple formula for arbitrary graphs (Theorem 5.10).

2. Background

Let G = ([n], E) be a graph with vertex set $[n] = \{1, \ldots, n\}$. A colouring of G is a function $\kappa : [n] \to \mathbb{N} = \{1, 2, 3, \ldots\}$ and we say that κ is proper if $\kappa(i) \neq \kappa(j)$ whenever $\{i, j\} \in E$, in other words, adjacent vertices are assigned different positive integers, which we think of colours. We denote by $\operatorname{asc}(\kappa)$ the number of ascents of κ , which are pairs of vertices (i, j) with i < j and $\kappa(i) < \kappa(j)$. The chromatic quasisymmetric function of G is the formal power series in an infinite alphabet of variables $\boldsymbol{x} = (x_1, x_2, x_3, \ldots)$ and an additional variable q given by [45, Definition 1.2]

(2.1)
$$X_G(\boldsymbol{x};q) = \sum_{\substack{\kappa:[n] \to \mathbb{N} \\ \kappa \text{ proper}}} q^{\operatorname{asc}(\kappa)} x_{\kappa(1)} x_{\kappa(2)} \cdots x_{\kappa(n)}.$$

Unlike the chromatic symmetric function $X_G(\mathbf{x}) = X_G(\mathbf{x}; 1)$, which is always symmetric in the x_i variables, $X_G(\mathbf{x}; q)$ is not symmetric in general. Nevertheless, if G is a natural unit interval graph, meaning that

(2.2) for all $1 \le i < j < k \le n$, if $\{i, k\} \in E$, then $\{i, j\} \in E$ and $\{j, k\} \in E$,

then $X_G(\boldsymbol{x};q)$ is symmetric in the x_i variables [45, Theorem 4.5] and palindromic in q with center of symmetry $\frac{|E|}{2}$ [45, Corollary 4.6], meaning that $X_G(\boldsymbol{x};q) = q^{|E|}X_G(\boldsymbol{x};q^{-1})$. In this

FIGURE 1. The bowtie graph G and the chromatic quasisymmetric function $X_G(\boldsymbol{x};q)$

G =



case, we can consider the expansion

(2.3)
$$X_G(\boldsymbol{x};q) = \sum_{\mu} c_{\mu}(q) e_{\mu}$$

in terms of the elementary symmetric functions e_{μ} , which are indexed by integer partitions $\mu = \mu_1 \cdots \mu_\ell$ and defined by

(2.4)
$$e_{\mu} = e_{\mu_1} \cdots e_{\mu_{\ell}}, \text{ where } e_k = \sum_{i_1 < \cdots < i_k} x_{i_1} \cdots x_{i_k}.$$

We say that $X_G(\boldsymbol{x}; q)$ is *e-positive* if every polynomial $c_{\mu}(q)$ has positive coefficients and we say that $X_G(\boldsymbol{x}; q)$ is *e-unimodal* if every polynomial $c_{\mu}(q)$ is *unimodal*, meaning that the coefficients form a weakly increasing sequence followed by a weakly decreasing sequence.

Example 2.1. The bowtie graph G and the chromatic quasisymmetric function $X_G(\boldsymbol{x}; q)$ are shown in Figure 1. We see that G is a natural unit interval graph because it satisfies (2.2) and we see from (2.5) that $X_G(\boldsymbol{x}; q)$ is indeed palindromic in q with center of symmetry $\frac{|E(G)|}{2} = \frac{6}{2} = 3$, and it is *e*-positive and *e*-unimodal.

Example 2.2. For the complete graph K_n on n vertices, a proper colouring κ must use n distinct colours and can be associated with a permutation $\sigma \in S_n$ by setting $\sigma(i) = j$ if $\kappa(i)$ is the *j*-th smallest colour used. Therefore, we have

(2.6)
$$X_{K_n}(\boldsymbol{x};q) = \left(\sum_{\sigma \in S_n} q^{|\{(i,j): i < j, \sigma_i < \sigma_j\}|}\right) e_n.$$

Then defining the q-integer and the q-factorial

(2.7)
$$[k]_q = 1 + q + q^2 + \dots + q^{k-1} = \frac{q^k - 1}{q - 1} \text{ and } [n]_q! = [n]_q [n - 1]_q \cdots [2]_q [1]_q,$$

we have by a standard induction argument that the sum in (2.6) is $[n]_q!$, and so

(2.8)
$$X_{K_n} = [n]_q! e_n.$$

Therefore $X_{K_n}(\boldsymbol{x};q)$ is *e*-positive and *e*-unimodal.

Stanley was interested in characterizing *e*-positivity of $X_G(\mathbf{x})$ in order to generalize positivity results of immanants of Jacobi–Trudi matrices [53]. The celebrated Stanley–Stembridge conjecture [49, Conjecture 5.1], [53, Conjecture 5.5] asserts that we have *e*-positivity for incomparability graphs of $(\mathbf{3} + \mathbf{1})$ -free partially ordered sets and Guay-Paquet [32, Theorem 5.1] showed that it suffices to prove this for the smaller class of natural unit interval graphs. The following quasisymmetric refinement by Shareshian and Wachs has seen connections to Hessenberg varieties and Hecke algebras.

Conjecture 2.3. [45, Conjecture 1.3] If G is a natural unit interval graph, then $X_G(\boldsymbol{x};q)$ is *e*-positive and *e*-unimodal.

Remark 2.4. An abstract graph is called a *unit interval graph* if its vertices can be labelled 1 through n so that (2.2) holds. The name comes from an equivalent characterization as a graph whose vertices are unit intervals of the real line and whose edges join intersecting intervals. It is convenient to consider *natural unit interval graphs*, where the adjective "natural" means that such a labelling is given. The chromatic quasisymmetric function does not depend on the choice of labelling, as long as (2.2) is satisfied [45, Proposition 4.2].

Remark 2.5. Let G = ([n], E) be a natural unit interval graph and consider the graph $G' = ([n], \{\{n+1-j, n+1-i\} : \{i, j\} \in E\})$ given by reversing the vertex labels. If we restrict to n variables and associate a proper colouring $\kappa : [n] \to [n]$ of G to $\kappa' : [n] \to [n]$ by setting $\kappa'(i) = n+1-\kappa(n+1-i)$, it follows that $X_G(x_1, \ldots, x_n; q) = X_{G'}(x_n, \ldots, x_1; q)$. Then we have $X_G(\boldsymbol{x}; q) = X_{G'}(\boldsymbol{x}; q)$ by symmetry in the x_i variables. This proves palindromicity in q and shows that we could have equivalently defined $X_G(\boldsymbol{x}; q)$ using descents instead of ascents. Similarly, we can switch the roles of i and (n+1-i) in everything that follows, for example, we can use increasing trees instead of decreasing trees.

3. SIGNED FORMULA

In this section, we prove a signed combinatorial *e*-expansion of the chromatic quasisymmetric function of any natural unit interval graph G = ([n], E). This formula has the advantage of allowing us to calculate a particular term without calculating the entire chromatic quasisymmetric function. We will walk through an example in Figure 3.

Definition 3.1. A subtree T of G is *decreasing* if for every vertex v of T, the unique path in T from the largest vertex of T to v is decreasing. Equivalently, every vertex of T except the largest has exactly one larger neighbour.

Given a decreasing subtree T of G, we define a permutation of V(T), denoted list(T), by reading the vertices of T starting from the smallest, and at each step, reading the smallest unread vertex of T that is adjacent to a read vertex of T. An example is given in Figure 2.

Remark 3.2. These trees and permutations also arise in work of Abreu and Nigro [1], Athanasiadis [9], and D'Adderio, Riccardi, and Siconolfi [21] in their study of chromatic quasisymmetric functions.

FIGURE 2. A decreasing tree T and the permutation list(T)

$$T = \underbrace{\begin{array}{c}1 \\ \bullet \\ \end{array}}_{2} \underbrace{\begin{array}{c}5 \\ \bullet \\ \end{array}}_{2} \underbrace{\begin{array}{c}6 \\ \bullet \\ \end{array}}_{1} \underbrace{\begin{array}{c}7 \\ \bullet \\ \end{array}}_{8} \underbrace{\begin{array}{c}8 \\ \end{array}}_{1} \underbrace{\left(T\right)}_{1} = 15632478$$

Definition 3.3. A tree triple of G is an object $\mathcal{T} = (T, \alpha, r)$ consisting of the following data.

- \circ T is a decreasing subtree of G.
- $\alpha = \alpha_1 \cdots \alpha_\ell$ is an integer composition with size $|\alpha| = \sum_{i=1}^\ell \alpha_i = |V(T)|$. r is a positive integer with $1 \le r \le \alpha_1$, the first part of α .

A forest triple of G is a sequence of tree triples $\mathcal{F} = (\mathcal{T}_i = (\mathcal{T}_i, \alpha^{(i)}, r_i))_{i=1}^m$ such that each vertex of G is in exactly one tree T_i and we have

(3.1)
$$\min(V(T_1)) < \min(V(T_2)) < \dots < \min(V(T_m)).$$

The type of a forest triple \mathcal{F} is the integer partition

(3.2)
$$\operatorname{type}(\mathcal{F}) = \operatorname{sort}(\alpha^{(1)} \cdots \alpha^{(m)})$$

obtained by concatenating the compositions and sorting in decreasing order. The sign of \mathcal{F} is the integer

(3.3)
$$\operatorname{sign}(\mathcal{F}) = (-1)^{\sum_{i=1}^{m} (\ell(\alpha^{(i)}) - 1)} = (-1)^{\ell(\operatorname{type}(\mathcal{F})) - m}$$

Letting $\sigma = \text{list}(T_1) \cdots \text{list}(T_m)$ be the concatenation of the permutations of the $V(T_i)$, we define the *weight* of \mathcal{F} to be the integer

(3.4)
$$\operatorname{weight}(\mathcal{F}) = \operatorname{inv}_G(\sigma) + \sum_{i=1}^m (r_i - 1), \text{ where}$$
$$\operatorname{inv}_G(\sigma) = |\{(i, j) : i < j, \sigma_i > \sigma_j, \{\sigma_j, \sigma_i\} \in E(G)\}$$

is the number of *G*-inversions of the permutation $\sigma \in S_n$. We denote by FT(G) the set of forest triples of G and by $FT_{\mu}(G)$ the set of forest triples of G of type μ .

We now state our signed combinatorial formula.

Theorem 3.4. The chromatic quasisymmetric function $X_G(\boldsymbol{x};q)$ of a natural unit interval graph G satisfies

(3.5)
$$X_G(\boldsymbol{x};q) = \sum_{\mathcal{F} \in \mathrm{FT}(G)} \mathrm{sign}(\mathcal{F}) q^{\mathrm{weight}(\mathcal{F})} e_{\mathrm{type}(\mathcal{F})} = \sum_{\mu} \left(\sum_{\mathcal{F} \in \mathrm{FT}_{\mu}(G)} \mathrm{sign}(\mathcal{F}) q^{\mathrm{weight}(\mathcal{F})} \right) e_{\mu}.$$



Example 3.5. Let us calculate the coefficient of e_{32} in the chromatic quasisymmetric function $X_G(\boldsymbol{x};q)$ for the bowtie graph G in Figure 3. Forest triples $\mathcal{F} \in \mathrm{FT}_{32}(G)$ come in one of two flavours.

We could have a single tree triple $\mathcal{T} = (T, \alpha, r)$, where T is a decreasing spanning tree of G and α is either 32 or 23. The four possibilities of T are given in Figure 3 along with $\operatorname{inv}_G(\sigma)$. We can independently choose either $\alpha = 32$ and $r = 1, 2, \text{ or } 3, \text{ or } \alpha = 23$ and r = 1or 2, and we have $\operatorname{sign}(\mathcal{F}) = (-1)^{2-1} = -1$, so these forest triples contribute

(3.6)
$$-(1+2q+q^2)(1+q+q^2+1+q)e_{32} = -(1+q)^2(2+2q+q^2)e_{32}.$$

Alternatively, \mathcal{F} can consist of two tree triples $\mathcal{T} = (T, \alpha, r)$ and $\mathcal{T}' = (T', \alpha', r')$ with trees of sizes 3 and 2 and compositions the single parts 3 and 2. The four possibilities of (T, T')are given in Figure 3 along with $\operatorname{inv}_G(\sigma)$. We can independently choose r and r', and we have $\operatorname{sign}(\mathcal{F}) = (-1)^{(1-1)+(1-1)} = +1$, so these forest triples contribute

(3.7)
$$+(2+2q)(1+q+q^2)(1+q)e_{32} = +(1+q)^2(2+2q+2q^2)e_{32}.$$

Therefore, by summing (3.6) and (3.7), the coefficient of e_{32} in $X_G(\boldsymbol{x};q)$ is $q^2(1+q)^2$, as we saw in (2.5).

Example 3.6. Forest triples of G of type n consist of a single tree triple $\mathcal{T} = (T, \alpha, r)$, where T is a decreasing spanning tree of G, α consists of the single part n, and $1 \leq r \leq n$. Therefore, the coefficient of e_n in $X_G(\boldsymbol{x}; q)$ is

(3.8)
$$c_n(q) = [n]_q \sum_{\substack{T \text{ dec. sp. tree}}} q^{\text{inv}_G(\text{list}(T))}.$$

We will show in Lemma 3.13 that (3.8) is equal to $[n]_q[b_2]_q[b_3]_q \cdots [b_n]_q$, where b_i is the number of smaller neighbours of vertex *i* in *G*. For example, the coefficient of e_5 in $X_G(\boldsymbol{x};q)$ for the bowtie graph *G* is $[5]_q[2]_q[2]_q$, as we saw in (2.5). This recovers the result [45, Corollary 7.2].

Before we prove Theorem 3.4, we will use the example of paths to show how finding a sign-reversing involution φ on the set FT(G) of forest triples of G can be used to prove that $X_G(\boldsymbol{x};q)$ is e-positive and give an explicit formula. The following result is well-known [45, Section 5], [46, Proposition 5.3], but this proof technique is new.

Proposition 3.7. The path $P_n = ([n], \{\{i, i+1\} : 1 \le i \le n-1\})$ has chromatic quasisymmetric function given by

(3.9)
$$X_{P_n}(\boldsymbol{x};q) = \sum_{\alpha \models n} [\alpha_1]_q ([\alpha_2]_q - 1) \cdots ([\alpha_m]_q - 1) e_{\operatorname{sort}(\alpha)}$$
$$= \sum_{\alpha \models n} q^{m-1} [\alpha_1]_q [\alpha_2 - 1]_q \cdots [\alpha_m - 1]_q e_{\operatorname{sort}(\alpha)},$$

where we sum over integer compositions $\alpha = \alpha_1 \cdots \alpha_m$ of size *n*. In particular, $X_{P_n}(\boldsymbol{x};q)$ is *e*-positive.

Example 3.8. The chromatic quasisymmetric function $X_{P_6}(\boldsymbol{x};q)$ with the compositions contributing to (3.9), is given by

Proof of Proposition 3.7. Because decreasing subtrees of P_n are simply paths from some vertex i to some vertex j > i, which we will denote $P_{i \to j}$, a forest triple $\mathcal{F} \in FT(P_n)$ must be of the form

(3.11)
$$\mathcal{F} = (\mathcal{T}_1 = (P_{1 \to i_2 - 1}, \alpha^{(1)}, r_1), \mathcal{T}_2 = (P_{i_2 \to i_3 - 1}, \alpha^{(2)}, r_2), \dots, \mathcal{T}_m = (P_{i_m \to n}, \alpha^{(m)}, r_m)).$$

Also note that we always have $\sigma = 123 \cdots n$, so $\operatorname{inv}_{P_n}(\sigma) = 0$ and $\operatorname{weight}(\mathcal{F}) = \sum_{i=1}^m (r_i - 1)$. We will now define a forest triple $\varphi(\mathcal{F})$ as follows. Let $1 \leq j \leq m$ be maximal, if one exists, such that either $\ell(\alpha^{(j)}) \geq 2$, or we have $j \geq 2$, $\ell(\alpha^{(j)}) = 1$, and $r_j = 1$. If $\ell(\alpha^{(j)}) \geq 2$, then letting $\alpha^{(j)} \setminus \alpha_{\ell}^{(j)}$ denote the composition $\alpha^{(j)}$ with its last part $\alpha_{\ell}^{(j)}$ removed, we define $\varphi(\mathcal{F})$ by replacing \mathcal{T}_j by the two tree triples

(3.12)
$$S_1 = (P_{i_j \to i_{j+1}-1-\alpha_\ell^{(j)}}, \alpha^{(j)} \setminus \alpha_\ell^{(j)}, r_j) \text{ and } S_2 = (P_{i_{j+1}-\alpha_\ell^{(j)} \to i_{j+1}-1}, \alpha_\ell^{(j)}, 1).$$

On the other hand, if $j \ge 2$, $\ell(\alpha^{(j)}) = 1$, and $r_j = 1$, then we define $\varphi(\mathcal{F})$ by replacing \mathcal{T}_{j-1} and \mathcal{T}_j by the tree triple

(3.13)
$$\mathcal{T} = (P_{i_{j-1} \to i_{j+1}-1}, \alpha^{(j-1)} \cdot \alpha^{(j)}, r_{j-1}).$$

If no such j exists, we set $\varphi(\mathcal{F}) = \mathcal{F}$. Informally, we look at our tree triples from right to left, identifying the first one where either the composition has at least two parts, in which case we break the path into two, or where we can join with the previous tree triple to undo this process.

We now claim that φ is an *involution*, meaning that $\varphi(\varphi(\mathcal{F})) = \mathcal{F}$. This is clear if \mathcal{F} is a *fixed point*, meaning that $\varphi(\mathcal{F}) = \mathcal{F}$, otherwise by maximality of j, we have $\ell(\alpha^{(k)}) = 1$ and $r_k \geq 2$ for $j + 1 \leq k \leq m$, so applying φ again will replace the desired tree triples. We also see that the map φ preserves type because the parts of the compositions do not change and it preserves weight because the r_i that appear or disappear are always equal to one. If \mathcal{F} is a fixed point we have $\operatorname{sign}(\mathcal{F}) = +1$ because every composition must have length one, and if \mathcal{F} is not a fixed point then φ is *sign-reversing*, meaning that $\operatorname{sign}(\varphi(\mathcal{F})) = -\operatorname{sign}(\mathcal{F})$, because the total number of tree triples changes by one.

In summary, this means that the map φ is a type-and-weight-preserving bijection between the negatively-signed non-fixed forest triples and the positively-signed non-fixed forest triples of P_n . Therefore, (3.5) reduces to a sum over only the fixed forest triples. Such an \mathcal{F} can be associated with a composition $\alpha \models n$ by reading the tree lengths from left to right, we will have type(\mathcal{F}) = sort(α), and we must have $r_i \ge 2$ for every $i \ge 2$, so we have as desired

(3.14)
$$X_{P_n}(\boldsymbol{x};q) = \sum_{\mathcal{F} \text{ fixed}} q^{\text{weight}(\mathcal{F})} e_{\text{type}(\mathcal{F})} = \sum_{\alpha \models n} \sum_{\substack{1 \le r_i \le \alpha_i \\ r_i \ge 2 \text{ if } i \ge 2}} q^{\sum_{i=1}^m (r_i-1)} e_{\text{sort}(\alpha)}$$
$$= \sum_{\alpha \models n} [\alpha_1]_q ([\alpha_2]_q - 1) \cdots ([\alpha_m]_q - 1) e_{\text{sort}(\alpha)}.$$

Example 3.9. Figure 4 shows all of the forest triples of P_6 of type 222 paired under our signreversing involution φ . The compositions are not written but they are 2, 22, or 222. We have indicated whether each r is 1 or 2 by circling the r-th smallest vertex of the corresponding tree. Informally, we can join two paths if the path on the right has its leftmost vertex circled. The coefficient of e_{222} in $X_{P_6}(\boldsymbol{x}; q)$ is $(q^2 + q^3)$, corresponding to the two fixed points.

We now prepare to prove Theorem 3.4. We begin by studying the permutation list(T). We will then use Alexandersson and Sulzgruber's combinatorial *e*-expansion of shifted *LLT polynomials*, which were originally introduced by Lascoux, Leclerc, and Thibon [38] and shown by Carlsson and Mellit to be related to chromatic quasisymmetric functions via an operation called *plethystic substitution*. This relationship was also studied by Alexandersson and Panova [4].

Definition 3.10. Let $\sigma = \sigma_1 \cdots \sigma_h$ be a permutation of a subset $A \subseteq [n]$. A *descent* of σ is a pair of adjacent entries (σ_i, σ_{i+1}) with $\sigma_i > \sigma_{i+1}$. A *left-to-right (LR) maximum* of σ is an entry σ_i such that $\sigma_i > \sigma_i$ for every i < j. In other words, reading the entries of σ from left



to right, when we read the entry σ_j , it is the maximum entry read by that point. We say that σ is a *tree list* of G if it satisfies the following two conditions.

- (1) (Descent condition) For every descent $\sigma_i > \sigma_{i+1}$ of σ , we have $\{\sigma_{i+1}, \sigma_i\} \in E(G)$.
- (2) (LR maxima condition) Let $a_1 < \cdots < a_s$ be the sequence of LR maxima of σ . Then for every $1 \le i \le s 1$, we have $\{a_i, a_{i+1}\} \in E(G)$.

Lemma 3.11. The function list is a bijection between decreasing subtrees of G and tree lists of G.

Proof. Let T be a decreasing subtree of G and let $\sigma = \text{list}(T)$ have sequence of LR maxima $a_1 < \cdots < a_s$. If σ_j is not an LR maximum, then we define its *parent* to be the rightmost entry σ_i with $\sigma_i > \sigma_j$ and i < j. We claim that the unique larger neighbour of σ_j in T is its parent σ_i . To have read σ_j after the larger entry σ_i , it must be adjacent to some $\sigma_{i'}$ for $i \leq i' < j$, which if j = i + 1 can only be σ_i , while if $j \geq i + 2$ can again only be σ_i because every $\sigma_{i'}$ for i < i' < j already has unique larger neighbour its parent by induction on j - i. We also claim that for every $1 \leq i \leq s - 1$, the unique larger neighbour of a_i in T is a_{i+1} . When we read a_{i+1} , it must be adjacent to a vertex read already, but the non-LR maxima already have unique larger neighbour their parent and the LR maxima $a_{i'}$ for $1 \leq i' < i$ already have unique larger neighbour $a_{i'+1}$ by induction on i, so the only possibility is a_i .

Now σ is a tree list of G because if $\sigma_i > \sigma_{i+1}$, then we have shown that $\{\sigma_{i+1}, \sigma_i\} \in E(T)$ so $\{\sigma_{i+1}, \sigma_i\} \in E(G)$, and for every $1 \le i \le s - 1$, we have shown that $\{a_i, a_{i+1}\} \in E(T)$ so $\{a_i, a_{i+1}\} \in E(G)$. Conversely, given a tree list σ , we can recover T because we have identified the unique larger neighbour of every vertex. The LR maxima condition ensures that the edges $\{a_i, a_{i+1}\}$ are present in G and thus available for T, and if σ_j has parent σ_i , then we have $\sigma_{i+1} \le \sigma_j < \sigma_i$, so the descent condition ensures that $\{\sigma_{i+1}, \sigma_i\} \in E(G)$ and (2.2) ensures that the edge $\{\sigma_j, \sigma_i\}$ is present in G and thus available for T.

Remark 3.12. Tree lists $\sigma \in S_n$ of G are also in bijection with acyclic orientations of G with unique sink 1, given by orienting each edge toward the earlier entry of σ .

Lemma 3.13. Let $A \subseteq [n]$ be nonempty and for a vertex $i \in A$, let $b_i(A)$ denote the number of vertices $j \in A$ with j < i and $\{j, i\} \in E(G)$. Then we have

(3.15)
$$\sum_{\sigma \in S_A: \ \sigma \text{ tree list of } G} q^{\operatorname{inv}_G(\sigma)} = \prod_{i \in A: \ i > \min(A)} [b_i(A)]_q.$$

Proof. We use induction on A, noting that the statement holds if |A| = 1 or if $b_{\max(A)}(A) = 0$, so assume otherwise and let $A' = A \setminus \{\max(A)\}$. A tree list $\sigma \in S_A$ arises from a tree list $\sigma' \in S_{A'}$ precisely by inserting $\max(A)$ either at the end of σ' or directly before one of its neighbours other than the one occurring first in σ' . If $\max(A)$ is inserted at the end, it creates no new *G*-inversions, while if it is inserted directly before the *j*-th neighbour from the right, where $1 \leq j \leq b_{\max(A)}(A) - 1$, then it creates *j* new *G*-inversions, so we multiply by $[b_{\max(A)}(A)]_q$ when we go from A' to A.

We now introduce LLT polynomials and plethystic substitution. All of the properties we will need can be blackboxed.

Definition 3.14. Let G = ([n], E) be a natural unit interval graph. The *LLT polynomial* is

(3.16)
$$\operatorname{LLT}_{G}(\boldsymbol{x};q) = \sum_{\substack{\kappa:[n] \to \mathbb{N} \\ \text{not necessarily proper}}} q^{\operatorname{asc}(\kappa)} x_{\kappa(1)} x_{\kappa(2)} \cdots x_{\kappa(n)}.$$

Note that we are summing over arbitrary colourings of G, not just the proper ones. Now these power series are related by the following operation.

Definition 3.15. Let $a(q) = a_0 + a_1q + \cdots + a_jq^j$ be a polynomial. The *plethystic substitution* $f \mapsto f[a(q)\boldsymbol{x}]$ is defined by setting $p_k[a(q)\boldsymbol{x}] = (a_0 + a_1q^k + \cdots + a_jq^{jk})p_k$ for every k, where $p_k = \sum_i x_i^k$ is the *power sum symmetric function*, and extending to an algebra homomorphism on symmetric functions.

Proposition 3.16. [12, Proposition 3.5] The chromatic quasisymmetric function $X_G(\boldsymbol{x};q)$ is related to the LLT polynomial $\text{LLT}_G(\boldsymbol{x};q)$ via the equation

(3.17)
$$X_G(\boldsymbol{x};q) = \frac{\mathrm{LLT}_G[(q-1)\boldsymbol{x}]}{(q-1)^n}.$$

Now Alexandersson conjectured [3, Conjecture 10] and proved with Sulzgruber [5, Corollary 2.10] the following combinatorial e-expansion of $LLT_G(\mathbf{x}; q+1)$. They formulated it in terms of the highest reachable vertex, but this is equivalent by Remark 2.5.

Definition 3.17. Let $\theta \subseteq E(G)$. For $u \in [n]$, the lowest reachable vertex $\operatorname{lrv}_{\theta}(u)$ is the smallest $v \in [n]$ such that the subgraph $([n], \theta)$ has a decreasing path from u to v. Defining $u \sim_{\theta} u'$ if $\operatorname{lrv}_{\theta}(u) = \operatorname{lrv}_{\theta}(u')$, we let $\pi(\theta) = \{[u_1]_{\theta}, \ldots, [u_m]_{\theta}\}$ be the equivalence classes of \sim_{θ} and we define $\lambda(\theta)$ to be the partition of n determined by their sizes.

Theorem 3.18. Let G be a natural unit interval graph. The shifted LLT polynomial $LLT_G(\boldsymbol{x}; q+1)$ satisfies

(3.18)
$$\operatorname{LLT}_{G}(\boldsymbol{x}; q+1) = \sum_{\boldsymbol{\theta} \subseteq E(G)} q^{|\boldsymbol{\theta}|} e_{\lambda(\boldsymbol{\theta})}.$$

Therefore, we get an expression for $X_G(\boldsymbol{x};q)$ by replacing q by (q-1) and applying plethystic substitution to the right hand side of (3.18). We will use the following identities.

Lemma 3.19.

- (1) We have $e_n[(q-1)\boldsymbol{x}] = \sum_{k=0}^n e_k[q\boldsymbol{x}]e_{n-k}[-\boldsymbol{x}].$ (2) We have $e_n[-\boldsymbol{x}] = (-1)^n h_n$, where $h_n = \sum_{i_1 \leq \cdots \leq i_n} x_{i_1} \cdots x_{i_n}$ is the complete homogeneous symmetric function.

(3) We have
$$h_n = \sum_{\alpha \models n} (-1)^{n-\ell(\alpha)} e_{\operatorname{sort}(\alpha)}$$
.

(4) We have

(3.19)
$$\frac{e_n[(q-1)\boldsymbol{x}]}{q-1} = \sum_{\alpha \models n} (-1)^{\ell(\alpha)-1} [\alpha_1]_q e_{\operatorname{sort}(\alpha)}.$$

Proof. The first three parts are [34, Equation 1.63], [50, Page 448], and [41, Theorem 2.18]. To prove the fourth, we use the others and we associate a composition of (n-k) to a composition of n by prepending a k to find that

$$(3.20) \qquad \frac{e_n[(q-1)\boldsymbol{x}]}{q-1} = \frac{1}{q-1} \sum_{k=0}^n e_k[q\boldsymbol{x}] e_{n-k}[-\boldsymbol{x}] = \frac{1}{q-1} \sum_{k=0}^n q^k e_k(-1)^{n-k} h_{n-k}$$
$$= \frac{1}{q-1} \sum_{k=0}^n q^k e_k(-1)^{n-k} \left(\sum_{\alpha \vDash n-k} (-1)^{n-k-\ell(\alpha)} e_{\operatorname{sort}(\alpha)} \right)$$
$$= \frac{1}{q-1} \left(\sum_{k=1}^n \sum_{\alpha \vDash n-k} q^k(-1)^{\ell(\alpha)} e_k e_{\operatorname{sort}(\alpha)} - \sum_{\alpha \vDash n} (-1)^{\ell(\alpha)-1} e_{\operatorname{sort}(\alpha)} \right)$$
$$= \sum_{\alpha \vDash n} (-1)^{\ell(\alpha)-1} \frac{q^{\alpha_1}-1}{q-1} e_{\operatorname{sort}(\alpha)} = \sum_{\alpha \vDash n} (-1)^{\ell(\alpha)-1} [\alpha_1]_q e_{\operatorname{sort}(\alpha)}.$$

Now we are ready to prove Theorem 3.4.

Proof of Theorem 3.4. We consider which $\theta \subseteq E(G)$ have $\pi(\theta) = \mathcal{A}$ for a fixed set partition $\mathcal{A} = \{A_1, \ldots, A_m\}$ of [n] with $\min(A_1) < \cdots < \min(A_m)$. For each of the (n-m) vertices i that are not the minimum of their block, say A_j , θ must have at least one of the $b_i(A_j)$ edges of $B(i) = \{\{i', i\} \in E(G) : i', i \in A_i, i' < i\}$ joining i to a smaller vertex in the same block. Additionally, θ may optionally have any number of edges in

(3.21)
$$I(\mathcal{A}) = \{\{u, v\} \in E(G) : u < v, u \in A_i, v \in A_j, i > j\}$$

that join a smaller vertex in a later block to a larger vertex in an earlier block, because these edges do not affect $\pi(\theta)$. Therefore, using Lemma 3.11 and Lemma 3.13, we have

$$(3.22) \qquad \sum_{\substack{\theta \subseteq E(G)\\\pi(\theta)=\mathcal{A}}} \frac{(q-1)^{|\theta|}}{(q-1)^{n-m}} = \left(\prod_{j=1}^{m} \sum_{\substack{i \in A_j\\i > \min(A_j)}} \frac{1}{q-1} \sum_{\substack{\emptyset \neq S_i \subseteq B(i)}} (q-1)^{|S_i|} \right) \sum_{S \subseteq I(\mathcal{A})} (q-1)^{|S|}$$
$$= q^{|I(\mathcal{A})|} \prod_{j=1}^{m} \prod_{\substack{i \in A_j\\i > \min(A_j)}} [b_i(A_j)]_q = q^{|I(\mathcal{A})|} \prod_{j=1}^{m} \sum_{\substack{\sigma^{(j)} \in S_{A_j}\\\sigma^{(j)} \text{ tree list of } G}} q^{\operatorname{inv}_G(\sigma^{(j)})}$$
$$= \sum_{\substack{T_1, \dots, T_m \text{ dec. trees}\\V(T_j) = A_j}} q^{\operatorname{inv}_G(\operatorname{list}(T_1) \cdots \operatorname{list}(T_m))},$$

where the factor of $q^{|I(\mathcal{A})|}$ accounts for G-inversions of $\sigma = \text{list}(T_1) \cdots \text{list}(T_m)$ occurring between vertices in different blocks. Now using Proposition 3.16 and Theorem 3.18, and separating $\theta \subseteq E(G)$ by the set partition $\pi(\theta) = \mathcal{A} = \{A_1, \ldots, A_m\}$, we have

$$\begin{split} X_{G}(\boldsymbol{x};q) &= \sum_{\theta \subseteq E(G)} \frac{(q-1)^{|\theta|} e_{\lambda(\theta)}[(q-1)\boldsymbol{x}]}{(q-1)^{n}} = \sum_{\mathcal{A}} \left(\sum_{\substack{\theta \subseteq E(G) \\ \pi(\theta) = \mathcal{A}}} \frac{(q-1)^{|\theta|}}{(q-1)^{n-m}} \right) \prod_{j=1}^{m} \frac{e_{|A_{j}|}[(q-1)\boldsymbol{x}]}{q-1} \\ &= \sum_{\mathcal{A}} \sum_{\substack{T_{1}, \dots, T_{m} \text{ dec. trees} \\ V(T_{j}) = A_{j}}} q^{\mathrm{inv}_{G}(\sigma)} \prod_{j=1}^{m} \sum_{\alpha^{(j)} \models |A_{j}|} (-1)^{\ell(\alpha^{(j)})-1} [\alpha_{1}^{(j)}]_{q} e_{\mathrm{sort}(\alpha^{(j)})} \\ &= \sum_{\substack{T_{1}, \dots, T_{m} \text{ dec. trees} \\ V(T_{1}) \sqcup \cdots \sqcup V(T_{m}) = [n] \\ \alpha^{(1)} \models |V(T_{1})|, \dots, \alpha^{(m)} \models |V(T_{m})| \\ 1 \leq r_{1} \leq \alpha_{1}^{(1)}, \dots, 1 \leq r_{m} \leq \alpha_{1}^{(m)}} \\ &= \sum_{\mathcal{F} \in \mathrm{FT}(G)} \mathrm{sign}(\mathcal{F}) q^{\mathrm{weight}(\mathcal{F})} e_{\mathrm{type}(\mathcal{F})}. \end{split}$$

4. Complete chains

In this section, we prove a combinatorial e-expansion of the chromatic quasisymmetric function of any K-chain. We begin by setting some notation.

Definition 4.1. We define the sum of graphs $G_1 = ([n_1], E_1)$ and $G_2 = ([n_2], E_2)$ to be (4.1) $G_1 + G_2 = ([n_1 + n_2 - 1], E_1 \cup \{\{i + n_1 - 1, j + n_1 - 1\} : \{i, j\} \in E_2\}).$

Informally, vertex n_1 of G_1 and vertex 1 of G_2 are glued together. For a composition $\gamma = \gamma_1 \cdots \gamma_\ell$ with all parts at least 2, we define the graph

(4.2)
$$K_{\gamma} = K_{\gamma_1} + \dots + K_{\gamma_\ell},$$

where K_a is the complete graph on [a]. Graphs of the form K_{γ} are called *K*-chains. Note that *K*-chains are natural unit interval graphs. The bowtie graph from Figure 1 is K_{33} .

This notation is due to Gebhard and Sagan, who proved [31, Corollary 7.7] that $X_{K_{\gamma}}(\boldsymbol{x})$ is *e*-positive for every γ . We will prove that the chromatic *quasisymmetric* function $X_{K_{\gamma}}(\boldsymbol{x};q)$ is *e*-positive and *e*-unimodal by finding a sign-reversing involution on forest triples. We will in fact find an involution whose fixed points have all compositions of length one, and which preserves $\alpha_1^{(1)}(\mathcal{F})$ and $r_1(\mathcal{F})$, where $\mathcal{T}_1 = (T_1, \alpha^{(1)}(\mathcal{F}), r_1(\mathcal{F}))$ is the tree triple of \mathcal{F} with $1 \in V(T_1)$, so we make the following definitions. Let G be a natural unit interval graph.

Definition 4.2. A tree triple $\mathcal{T} = (T, \alpha, r)$ of G is *breakable* if $\ell(\alpha) \geq 2$. A forest triple \mathcal{F} of G is an *atom* if all of its tree triples are not breakable.

Note that if $\mathcal{T} = (T, \alpha, r)$ is not breakable, then the composition α consists of the single part |V(T)|. In particular, if \mathcal{F} is an atom, then $\operatorname{sign}(\mathcal{F}) = (-1)^{\sum_{i=1}^{m} (1-1)} = +1$.

Definition 4.3. A forest triple \mathcal{F} of G is simple if $r_1(\mathcal{F}) = 1$. Let $\mathrm{FT}_{\mathrm{simple}}(G)$ denote the set of simple forest triples of G. A nice involution for G is a function $\varphi : \mathrm{FT}_{\mathrm{simple}}(G) \to \mathrm{FT}_{\mathrm{simple}}(G)$ with the following properties.

- (1) φ is an *involution*, meaning that $\varphi(\varphi(\mathcal{F})) = \mathcal{F}$ for all $\mathcal{F} \in \mathrm{FT}_{\mathrm{simple}}(G)$. In other words, φ is its own inverse, so in particular it is a bijection.
- (2) φ preserves type, weight, and $\alpha_1^{(1)}$, which is the first part of the composition associated to the tree containing the vertex 1.
- (3) If \mathcal{F} is a fixed point of φ , meaning that $\varphi(\mathcal{F}) = \mathcal{F}$, then \mathcal{F} is an atom.
- (4) φ is sign-reversing, meaning that if $\varphi(\mathcal{F}) \neq \mathcal{F}$, then $\operatorname{sign}(\varphi(\mathcal{F})) = -\operatorname{sign}(\mathcal{F})$.

Note that a forest triple $\mathcal{F}' \in FT(G)$ arises from some simple forest triple \mathcal{F} by choosing some $1 \leq r_1(\mathcal{F}') \leq \alpha_1^{(1)}(\mathcal{F})$. Therefore, if we can find a nice involution φ for G, then we have

(4.3)
$$X_G(\boldsymbol{x};q) = \sum_{\mathcal{F} \in \mathrm{FT}_{\mathrm{simple}}(G), \varphi(\mathcal{F}) = \mathcal{F}} [\alpha_1^{(1)}(\mathcal{F})]_q \ q^{\mathrm{weight}(\mathcal{F})} e_{\mathrm{type}(\mathcal{F})},$$

thus proving that $X_G(\boldsymbol{x}; q)$ is *e*-positive. The following Lemma will help us describe forest triples of a sum of graphs.

Lemma 4.4. Let T be a decreasing subtree of G that contains a cut vertex c. Then the tree list $\sigma = \text{list}(T)$ contains all of the vertices of T at most c in some order, followed by all of the vertices of T strictly greater than c in some order.

Proof. If not, then there is some i with $\sigma_i > c$ and $\sigma_{i+1} \leq c$. By the LR maxima condition, the cut vertex c must appear before σ_i , but this means that $\sigma_{i+1} < c$ and σ does not satisfy the descent condition.

We will want a standard way of taking a permutation σ and writing a new permutation whose first entry is min(σ). The following choice is not the only possibility, but it will work.

Definition 4.5. Let $\sigma = \sigma_1 \cdots \sigma_h$ be a permutation of some set $A = \{a_1 < \cdots < a_h\} \subseteq [n]$. Define indstart (σ) to be the integer $1 \leq r \leq h$ such that $\sigma_1 = a_r$ and define the permutation

(4.4)
$$\operatorname{startmin}(\sigma) = w_1 w_2 \cdots w_{r-1}(\sigma),$$

where w_i is the transposition that switches the positions of the entries a_i and a_{i+1} in σ . Note that startmin(σ) starts with its minimum element a_1 , and if the entries in A are all adjacent in G, then we have

(4.5)
$$\operatorname{inv}_{G}(\sigma) = \operatorname{inv}_{G}(\operatorname{startmin}(\sigma)) + (\operatorname{indstart}(\sigma) - 1)$$

because each of the (indstart(σ) - 1) transpositions that are applied decreases the number of *G*-inversions by exactly one. Conversely, given a permutation $\sigma = \sigma_1 \cdots \sigma_h$ of *A* with $\sigma_1 = a_1$ and an integer $1 \le r \le h$, define the permutation

(4.6)
$$\operatorname{startr}(\sigma, r) = w_{r-1} \cdots w_2 w_1(\sigma).$$

Note that $\operatorname{startr}(\sigma, r)$ starts with the element a_r and $\operatorname{startmin}(\operatorname{startr}(\sigma, \operatorname{indstart}(\sigma))) = \sigma$.

Example 4.6. For $\sigma = 61743$, we have $indstart(\sigma) = 4$ and $startmin(\sigma) = 13764$.

We also define useful breaking and joining maps on tree triples. For convenience, we may refer to a tree T and the permutation $\sigma = \text{list}(T)$ interchangeably.

Definition 4.7. Let $\mathcal{T} = (T, \alpha, r)$ be a breakable tree triple of G with all vertices of T adjacent in G, let $\alpha \setminus \alpha_{\ell}$ be the composition α with its last part α_{ℓ} removed, and divide $\sigma = \operatorname{list}(T)$ into its head and tail,

(4.7)
$$\sigma^{\text{head}} = \sigma_1 \cdots \sigma_{|V(T)|-\alpha_\ell} \text{ and } \sigma^{\text{tail}} = \sigma_{|V(T)|-\alpha_\ell+1} \cdots \sigma_{|V(T)|}.$$

Then we define the pair of tree triples $easybreak(\mathcal{T}) = (\mathcal{S}_1, \mathcal{S}_2)$, where

(4.8)
$$\mathcal{S}_1 = (\sigma^{\text{head}}, \alpha \setminus \alpha_\ell, r) \text{ and } \mathcal{S}_2 = (\text{startmin}(\sigma^{\text{tail}}), \alpha_\ell, \text{indstart}(\sigma^{\text{tail}})).$$

Conversely, given tree triples $S_1 = (S_1, \alpha^{(1)}, r_1)$ and $S_2 = (S_2, \alpha^{(2)}, r_2)$ with $\ell(\alpha^{(2)}) = 1$, min $(V(S_1)) < \min(V(S_2))$, and $V(S_1) \sqcup V(S_2)$ all adjacent in G, we define the tree triple

(4.9)
$$\operatorname{easyjoin}(\mathcal{S}_1, \mathcal{S}_2) = \mathcal{T} = (\operatorname{list}(S_1) \cdot \operatorname{startr}(\operatorname{list}(S_2), r_2), \alpha^{(1)} \cdot \alpha^{(2)}, r_1).$$

Note that easybreak $(\mathcal{T}) = (\mathcal{S}_1, \mathcal{S}_2)$ if and only if easyjoin $(\mathcal{S}_1, \mathcal{S}_2) = \mathcal{T}$, and by (4.5), we have

(4.10)
$$\operatorname{inv}_G(\operatorname{list}(T)) = \operatorname{inv}_G(\operatorname{list}(S_1) \cdot \operatorname{list}(S_2)) + (r_2 - 1)$$

FIGURE 5. Some simple forest triples \mathcal{F} of K_6 and $\varphi(\mathcal{F})$



\mathcal{F}	$arphi(\mathcal{F})$	${\cal F}$	$arphi(\mathcal{F})$
(123456, 33, 1)	(123, 3, 1), (456, 3, 1)	(136254, 222, 1)	(1362, 22, 1), (45, 2, 2)
(123546, 33, 1)	(123, 3, 1), (456, 3, 2)	(13, 2, 1), (2645, 22, 2)	(13, 2, 1), (26, 2, 2), (45, 2, 1)
(123654, 33, 1)	(123, 3, 1), (465, 3, 3)	(15, 2, 1), (2463, 22, 1)	(15, 2, 1), (24, 2, 1), (36, 2, 2)

We now show how we can use the easybreak and easyjoin maps to find a nice involution for the complete graph K_n . We have already seen in Example 2.2 that $X_{K_n}(\boldsymbol{x};q) = [n]_q!e_n$, but this proof will be instructive for when we prove the main result of this section. Some examples are given in Figure 5.

Proposition 4.8. There is a nice involution φ for the complete graph K_n . Furthermore, the fixed points of φ are exactly

(4.11) $\operatorname{Fix}(\varphi) = \{ \mathcal{F} \in \operatorname{FT}_{\operatorname{simple}}(K_n) : \mathcal{F} \text{ is an atom, } \mathcal{F} \text{ consists of a single tree triple} \}.$

In particular, by Lemma 3.13, the chromatic quasisymmetric function is

(4.12)
$$X_{K_n}(\boldsymbol{x};q) = \sum_{\mathcal{F} \in \operatorname{Fix}(\varphi)} [\alpha_1^{(1)}(\mathcal{F})]_q \ q^{\operatorname{weight}(\mathcal{F})} e_{\operatorname{type}(\mathcal{F})} = [n]_q \sum_{\sigma \in S_n: \ \sigma_1 = 1} q^{\operatorname{inv}_{K_n}(\sigma)} e_n = [n]_q! e_n.$$

Proof. We build our nice involution φ using the easybreak and easyjoin maps. Let

(4.13)
$$\mathcal{F} = (\mathcal{T}_1 = (T_1, \alpha^{(1)}, 1), \dots, \mathcal{T}_m = (T_m, \alpha^{(m)}, r_m)) \in \mathrm{FT}_{\mathrm{simple}}(K_n).$$

If \mathcal{T}_m is breakable, with easybreak $(\mathcal{T}) = (\mathcal{S}_1, \mathcal{S}_2)$, then we define $\varphi(\mathcal{F})$ by replacing \mathcal{T}_m by \mathcal{S}_1 and \mathcal{S}_2 , that is, $\varphi(\mathcal{F}) = (\mathcal{T}_1, \ldots, \mathcal{T}_{m-1}, \mathcal{S}_1, \mathcal{S}_2)$. Note that by our definition in (4.7), the tree triples are in the correct order and weight is preserved by (4.10). We have that φ reverses sign and preserves type and $\alpha_1^{(1)}$.

If \mathcal{T}_m is not breakable and m = 1, then $\mathcal{F} \in \operatorname{Fix}(\varphi)$ and we perforce define $\varphi(\mathcal{F}) = \mathcal{F}$. If \mathcal{T}_m is not breakable and $m \geq 2$, then we define $\varphi(\mathcal{F})$ by replacing \mathcal{T}_{m-1} and \mathcal{T}_m by $\mathcal{T} = \operatorname{easyjoin}(\mathcal{T}_{m-1}, \mathcal{T}_m)$, that is, $\varphi(\mathcal{F}) = (\mathcal{T}_1, \ldots, \mathcal{T}_{m-2}, \mathcal{T})$. Note that by our definition in (4.7), the tree triples are in the correct order and weight is preserved by (4.10). We have that φ reverses sign and preserves type and $\alpha_1^{(1)}$. By construction, the map φ is an involution with fixed points exactly $\operatorname{Fix}(\varphi)$, as desired.



\mathcal{F}	$\operatorname{seg}_{[6]}(\mathcal{F})$	$ \mathcal{T}_i _{\geq 6}$	$ \mathcal{F} _{\geq 6}$
(123456789, 9, 1)	1	(6789, 4, 1)	(1234, 4, 1)
(123654789, 72, 1)	1	(6789, 22, 1)	(1234, 22, 1)
(1362548, 52, 1), (79, 2, 2)	2	(68, 2, 1)	(13, 2, 1), (24, 2, 2)
(143, 3, 1), (265978, 33, 3)	2	(6978, 13, 1)	(1423, 13, 1)
(164598, 231, 1), (23, 2, 2), (7, 1, 1)	3	(698, 21, 1)	(143, 21, 1), (2, 1, 1)

We now define a notion of restricting a simple forest triple \mathcal{F} of $G_1 + G_2$ to one of G_2 and we define an invariant that describes how many "segments" of trees \mathcal{F} has present in G_1 . Some examples are given in Figure 6.

Definition 4.9. Let $G_1 = ([n_1], E_1)$ and $G_2 = ([n_2], E_2)$ be natural unit interval graphs and let $\mathcal{F} = (\mathcal{T}_1 = (T_1, \alpha^{(1)}, r_1 = 1), \dots, \mathcal{T}_m = (T_m, \alpha^{(m)}, r_m))$ be a simple forest triple of $G = G_1 + G_2$. Let *i* be such that $n_1 \in V(T_i)$, let $\sigma = \text{list}(T_i)$, let $k = |V(T_i) \cap [n_1]|$, and noting that $k \leq |V(T_i)| = |\alpha^{(i)}|$, we define $\text{seg}_{[n_1]}(\mathcal{T}_i)$ to be the number *j* such that

(4.14)
$$\alpha_1^{(i)} + \dots + \alpha_{j-1}^{(i)} < k \le \alpha_1^{(i)} + \dots + \alpha_j^{(i)}.$$

Letting $i' \ge i$ be maximal with $V(T_{i'}) \cap [n_1] \ne \emptyset$, we say that the tree triples $\mathcal{T}_1, \ldots, \mathcal{T}_{i'}$ are present in $[n_1]$, we define $\mathcal{F}|_{[n_1]} = (\mathcal{T}_1, \ldots, \mathcal{T}_{i'})$, and we define the integer

(4.15)
$$\sup_{[n_1]}(\mathcal{F}) = \sup_{[n_1]}(\mathcal{T}_i) + \sum_{1 \le t \le i', \ t \ne i} \ell(\alpha^{(t)})$$

We now define the tree triple of G

(4.16)
$$\mathcal{T}_i|_{\geq n_1} = (n_1 \ \sigma_{k+1} \ \cdots \ \sigma_{\ell(\sigma)}, (\alpha_1^{(i)} + \cdots + \alpha_j^{(i)} - k + 1)\alpha_{j+1}^{(i)} \cdots \alpha_{\ell}^{(i)}, 1)$$

and the simple forest triple of G_2

(4.17)
$$\mathcal{F}|_{\geq n_1} = (\mathcal{T}_i|_{\geq n_1} - (n_1 - 1), \mathcal{T}_{i'+1} - (n_1 - 1), \dots, \mathcal{T}_m - (n_1 - 1))$$

where $\mathcal{T} - x$ denotes the tree triple obtained by subtracting x from each entry of list(T).

Note that by Lemma 4.4, we have $\sigma_t \leq n_1$ for $1 \leq t \leq k$ and $\sigma_t > n_1$ for t > k, so the permutation $n_1 \sigma_{k+1} \cdots \sigma_{\ell(\sigma)}$ starts with its minimum element and its descents and successive LR maxima appear in σ , so it is indeed a tree list of G and $\mathcal{F}|_{\geq n_1} \in \mathrm{FT}_{\mathrm{simple}}(G_2)$.

We now prove the key result that will let us deduce *e*-positivity for *K*-chains by inductively adding complete graphs one at a time. The argument is identical if we add an *almost-complete* graph, where a single edge is missing, so we prove both of these cases together. For $\epsilon \in \{0, 1\}$, let K_a^{ϵ} denote the complete graph on [a] if $\epsilon = 0$ and the almost-complete graph on [a], with the edge (1, a) removed, if $\epsilon = 1$. Note that every σ with $\sigma_1 = \min(\sigma)$ is a tree list for K_a , and is a tree list for $K_a^{(1)}$ unless $\sigma_1 = 1$ and $\sigma_2 = a$.

Theorem 4.10. Let φ' be a nice involution for a natural unit interval graph G'. Then there is a nice involution φ for $G = K_a^{\epsilon} + G'$. Furthermore, the fixed points of φ are exactly

(4.18) Fix(
$$\varphi$$
) = { $\mathcal{F} \in \operatorname{FT}_{\operatorname{simple}}(G)$: \mathcal{F} is an atom, $\mathcal{F}|_{\geq a}$ is fixed under φ' , $\operatorname{seg}_{[a]}(\mathcal{F}) \leq 2$,
if $1 \notin V(T_i)$, then $r_i \geq a - \epsilon$, if $1 \in V(T_i)$, then $|V(T_i)| \geq a$ },

where $\mathcal{T}_i = (T_i, \alpha^{(i)}, r_i)$ is the tree triple of \mathcal{F} with $a \in V(T_i)$.

Proof. We perforce define $\varphi(\mathcal{F}) = \mathcal{F}$ for $\mathcal{F} \in \operatorname{Fix}(\varphi)$, so we now build a sign-reversing involution with no fixed points on $\operatorname{FT}_{\operatorname{simple}}(G) \setminus \operatorname{Fix}(\varphi)$ using several cases. Let

(4.19)
$$\mathcal{F} = (\mathcal{T}_1 = (T_1, \alpha^{(1)}, r_1), \dots, \mathcal{T}_m = (T_m, \alpha^{(m)}, r_m)) \in \mathrm{FT}_{\mathrm{simple}}(G) \setminus \mathrm{Fix}(\varphi),$$

let *i* be such that $a \in V(T_i)$, and let $i' \geq i$ be maximal with $V(T_{i'}) \cap [a] \neq \emptyset$. Illustrative examples for $G = K_{64} = K_6 + K_4$ will be provided in the figures.

Case 1: $\mathcal{F}|_{\geq a}$ is not fixed under φ' . We first define a sign-reversing involution $\varphi|_{\geq a}$ with no fixed points on those \mathcal{F} for which $\varphi'(\mathcal{F}|_{\geq a}) \neq \mathcal{F}|_{\geq a}$. The idea will be to restrict \mathcal{F} to G', apply the map φ' , and reattach to the rest of \mathcal{F} . Let $\sigma = \operatorname{list}(T_i), k = |V(T_i) \cap [a]|$, and $j = \operatorname{seg}_{[a]}(\mathcal{T}_i)$. Apply the map φ' to get $\varphi'(\mathcal{F}|_{\geq a}) = \mathcal{F}' = (\mathcal{T}'_1 = (\mathcal{T}'_1, \alpha^{(1)'}, 1), \mathcal{T}'_2, \ldots, \mathcal{T}'_{m'})$ and let $\sigma' = \operatorname{list}(T'_1)$. We now define the tree triple of G

(4.20)
$$\mathcal{T}_i \vee \mathcal{T}'_1 = (\sigma_1 \cdots \sigma_k \ (\sigma'_2 + a - 1) \cdots (\sigma'_{|V(\mathcal{T}'_1)|} + a - 1), \alpha_1^{(i)} \cdots \alpha_j^{(i)} \alpha_2^{(1)'} \cdots \alpha_\ell^{(1)'}, r_i)$$

and we define $\varphi|_{>a}(\mathcal{F}) = \mathcal{F} \vee \mathcal{F}'$, where

$$(4.21) \qquad \mathcal{F} \lor \mathcal{F}' = (\mathcal{T}_1, \dots, \mathcal{T}_{i-1}, \mathcal{T}_i \lor \mathcal{T}_1', \mathcal{T}_{i+1}, \dots, \mathcal{T}_{i'}, \mathcal{T}_2' + (a-1), \dots, \mathcal{T}_{m'}' + (a-1)).$$

The permutation $\sigma_1 \cdots \sigma_k \ (\sigma'_2 + a - 1) \cdots (\sigma'_{|V(T'_1)|} + a - 1)$ starts with its minimum element and its descents and successive LR maxima are those of σ or translated ones in σ' so it is a tree list of G. Because φ' preserves the first part of the first composition, we have

$$(4.22) \ \alpha_1^{(i)} + \dots + \alpha_j^{(i)} + \alpha_2^{(1)'} + \dots + \alpha_\ell^{(1)'} = k - 1 + \alpha_1^{(1)} (\mathcal{F}|_{\geq a}) + \alpha_2^{(1)'} + \dots + \alpha_\ell^{(1)'} = k - 1 + |V(T_1')|$$

and we have $r_i \leq \alpha_1^{(i)}$, so $\mathcal{T}_i \vee \mathcal{T}'_1$ is indeed a tree triple of G and $\varphi|_{\geq a}(\mathcal{F})$ is a simple forest triple of G. Informally, $\mathcal{F} \vee \mathcal{F}'$ replaces the part of \mathcal{F} in G' by \mathcal{F}' . By construction, we have $\mathcal{F} \vee (\mathcal{F}|_{\geq a}) = \mathcal{F}$ and we have $(\mathcal{F} \vee \mathcal{F}')|_{\geq a} = \mathcal{F}'$, which by hypothesis is not fixed under φ' . The map $\varphi|_{\geq a}$ reverses sign, preserves type, weight, and $\alpha_1^{(1)}$, has no fixed points, and is an involution because

(4.23)
$$\varphi|_{\geq a}(\varphi|_{\geq a}(\mathcal{F})) = \mathcal{F} \lor (\varphi'(\varphi'(\mathcal{F}|_{\geq a}))) = \mathcal{F} \lor (\mathcal{F}|_{\geq a}) = \mathcal{F}.$$

FIGURE 7. Some simple forest triples \mathcal{F} of K_{64} and $\varphi|_{\geq 6}(\mathcal{F})$

$K_{64} = 1 + 1 + 1 + 1 + 1 + 1 + 1 + 1 + 1 + 1$	

${\cal F}$	$\mathcal{F} _{\geq 6}$	\mathcal{F}'	$arphi _{\geq 6}(\mathcal{F}) = \mathcal{F} \lor \mathcal{F}'$
(123654789, 72, 1)	(1234, 22, 1)	(12, 2, 1), (34, 2, 1)	(1236547, 7, 1), (89, 2, 1)
(136254897, 432, 1)	(1342, 22, 1)	(13, 2, 1), (24, 2, 2)	(1362548, 43, 1), (79, 2, 2)
(154623978, 522, 1)	(1423, 22, 1)	(14, 2, 1), (23, 2, 1)	(1546239, 52, 1), (78, 2, 1)
(163452987, 2322, 1)	(1432, 22, 1)	(14, 2, 1), (23, 2, 2)	(1634529, 232, 1), (78, 2, 2)

Case 2: $\mathcal{F}|_{\geq a}$ is fixed under φ' and $\operatorname{seg}_{[a]}(\mathcal{F}) \leq 2$. First note that if $\operatorname{seg}_{[a]}(\mathcal{F}) = 1$, then $i = 1, \mathcal{F}$ is an atom, $[a] \subseteq V(T_1)$, and $\mathcal{F} \in \operatorname{Fix}(\varphi)$, so we must have $\operatorname{seg}_{[a]}(\mathcal{F}) = 2$. We now define a bijection φ_2 between those \mathcal{F} with $\operatorname{seg}_{[a]}(\mathcal{T}_i) = 2$, for which the only tree triple present in [a] is of the form $\mathcal{T}_i = \mathcal{T}_1 = (T_1, \alpha^{(1)} = \alpha_1^{(1)} \alpha_2^{(1)}, 1)$, and those with $\operatorname{seg}_{[a]}(\mathcal{T}_i) = 1$, which have two tree triples present in [a], which are both not breakable. Then we get a sign-reversing involution by either applying φ_2 or its inverse.

More specifically, for each k we define a bijection

$$\begin{array}{ll} (4.24) & \operatorname{break}_k : \{T \ \operatorname{dec.} \ \operatorname{subtree} \ \operatorname{of} \ G : \ [a] \subseteq V(T), \ k < |V(T)| < k + a\} \\ & \to \{(S_1, S_2, r_2) : \ S_1, S_2 \ \operatorname{dec.} \ \operatorname{subtrees} \ \operatorname{of} \ G, \ [a] \subseteq V(S_1) \sqcup V(S_2), \ 1 \in V(S_1), \\ & |V(S_1)| < a, \ 1 \leq r_2 \leq |V(S_2)| = k, \ \operatorname{if} \ a \in V(S_2), \ \operatorname{then} \ r_2 \leq a - \epsilon - 1\}, \end{array}$$

breaking a tree T into S_1 and S_2 , with $1 \in V(S_1)$ and $|V(S_2)| = k$, such that

(4.25)
$$\operatorname{inv}_G(\operatorname{list}(T)) = \operatorname{inv}_G(\operatorname{list}(S_1) \cdot \operatorname{list}(S_2)) + (r_2 - 1).$$

Then we define $\varphi_2(\mathcal{F})$ by taking $\operatorname{break}_{\alpha_2^{(1)}}(T_1) = (S_1, S_2, r_2)$ and replacing \mathcal{T}_1 by the tree triples

(4.26)
$$S_1 = (S_1, \alpha_1^{(1)}, 1) \text{ and } S_2 = (S_2, \alpha_2^{(1)}, r_2).$$

By construction, φ_2 will reverse sign, will preserve type and $\alpha_1^{(1)}$, will preserve weight by (4.25), and will have no fixed points. Let T be a decreasing subtree of G with $[a] \subseteq V(T)$ and k < |V(T)| < k + a, let $\sigma = \text{list}(T)$, and let $2 \leq j \leq |V(T)|$ be such that $\sigma_j = a$. Note that if j = 2, then we must have $\epsilon = 0$. We will have three cases, depending on the position j of the entry a in σ .

FIGURE 8. Some simple forest triples \mathcal{F} of K_{64} and $\varphi_2(\mathcal{F})$



\mathcal{F}	$\varphi_2(\mathcal{F})$	${\cal F}$	$\varphi_2(\mathcal{F})$
(123456789, 36 , 1)	(123, 3, 1), (456789, 6, 1)	(123456789, 54 , 1)	(12345, 5, 1), (6789, 4, 1)
(132465789, 36 , 1)	(132, 3, 1), (465789, 6, 2)	(156324789, 54, 1)	(15324, 5, 1), (6789, 4, 4)
(123654789, 36 , 1)	(123, 3, 1), (465789, 6, 3)	(162345789, 54, 1)	(16789, 5, 1), (2345, 4, 1)
(146532789, 36 , 1)	(145, 3, 1), (263789, 6, 4)	(163425789, 54, 1)	(16789, 5, 1), (2435, 4, 2)
(165234789, 36 , 1)	(152, 3, 1), (346789, 6, 5)	(165432789, 54, 1)	(16789, 5, 1), (2543, 4, 4)

If
$$j \ge |V(T)| - k + 1$$
, then as in Definition 4.7, we divide σ into its head and tail,

(4.27)
$$\sigma^{\text{head}} = \sigma_1 \cdots \sigma_{|V(T)|-k} \text{ and } \sigma^{\text{tail}} = \sigma_{|V(T)|-k+1} \cdots \sigma_{|V(T)|},$$

and we define (S_1, S_2, r_2) by

(4.28)
$$\operatorname{list}(S_1) = \sigma^{\operatorname{head}}, \operatorname{list}(S_2) = \operatorname{startmin}(\sigma^{\operatorname{tail}}), \text{ and } r_2 = \operatorname{indstart}(\sigma^{\operatorname{tail}}).$$

Because |V(T)| - k < a and by Lemma 4.4, σ^{head} has all entries in [a - 1] so is a tree list of G, we have $\min(\sigma^{\text{tail}}) \leq \sigma_1^{\text{tail}} \leq a$, so the only transpositions applied to σ^{tail} switch entries at most a, which are all adjacent, and its descents and successive LR maxima appear in the tree list σ , so startmin(σ^{tail}) is a tree list of G, and (4.25) follows from (4.5). Also note that $a \in V(S_2)$ and we have

$$(4.29) 1 \le r_2 \le |V(S_2) \cap [a]| = a - (|V(T)| - k) \le a - 1$$

with equality only if k = |V(T)| - 1 and $\sigma_2 = a$, meaning that $\epsilon = 0$, so $r_2 \le a - \epsilon - 1$.

If $a - k + 1 \le j \le |V(T)| - k$, then the entry *a* would appear in σ^{head} and σ^{tail} would have entries less than *a* and greater than *a* and therefore would have no hope of satisfying the LR maximum condition. Therefore, we move the entry *a* from position *j* to the start of the tail, at position |V(T)| - k + 1, and define

(4.30)
$$\sigma^{\text{head}'} = \sigma_1 \cdots \sigma_{j-1} \sigma_{j+1} \cdots \sigma_{|V(T)|-k+1} \text{ and } \sigma^{\text{tail}'} = a \sigma_{|V(T)|-k+2} \cdots \sigma_{\ell(\sigma)},$$

and we define (S_1, S_2, r_2) by

(4.31)
$$\operatorname{list}(S_1) = \sigma^{\operatorname{head}'}, \operatorname{list}(S_2) = \operatorname{startmin}(\sigma^{\operatorname{tail}'}), \text{ and } r_2 = a - j + 1.$$

Informally, r_2 records the original position of the entry a. Again, $\sigma^{\text{head}'}$ has all entries in [a-1] and the descents and successive LR maxima of startmin $(\sigma^{\text{tail}'})$ are either in $\{2, \ldots, a\}$

or appear in σ so these are tree lists of G. Moving the entry a from position j to position (|V(T)| - k + 1) created (|V(T)| - k + 1 - j) new G-inversions, and then we applied

(4.32)
$$\operatorname{indstart}(\sigma^{\operatorname{tail}'}) - 1 = a - \ell(\sigma^{\operatorname{head}'}) - 1 = a - (|V(T)| - k) - 1$$

transpositions, so (4.25) holds. Also note that $a \in V(S_2)$ and we have

(4.33)
$$a - (|V(T)| - k) + 1 \le r_2 \le a - \epsilon - 1$$

because $j \ge 2$ with equality only if $\epsilon = 0$.

If $j \leq a - k$, then we remove k entries from the inside of σ and define

(4.34)
$$\sigma^{\text{in}} = \sigma_{a-k+1} \cdots \sigma_a \text{ and } \sigma^{\text{out}} = \sigma_1 \cdots \sigma_{a-k} \sigma_{a+1} \cdots \sigma_{|V(T)|},$$

and we define (S_1, S_2, r_2) by

(4.35)
$$\operatorname{list}(S_1) = \sigma^{\operatorname{out}}, \ \operatorname{list}(S_2) = \operatorname{startmin}(\sigma^{\operatorname{in}}), \ \operatorname{and} \ r_2 = \operatorname{indstart}(\sigma^{\operatorname{in}}).$$

Because $j \leq a - k$, the entry a appears in σ^{out} so its descents and successive LR maxima appear in σ and $\operatorname{startmin}(\sigma^{\text{in}})$ has all entries in [a - 1] so these are tree lists of G, and we have $r_2 \leq \ell(\sigma^{\text{in}}) = k$. Moving the string σ^{in} does not affect the number of G-inversions because its entries are less than a, whereas every $\sigma_t > a$ for t > a by Lemma 4.4, so again (4.25) follows from (4.5). Also note that the entries σ_1 and σ_2 are unaffected and $a \in V(S_1)$.

It remains to show that the map break_k is a bijection. The idea is that we can determine which case of the map was applied by whether $a \in V(S_2)$ and from r_2 , using (4.29) and (4.33), so we can recover list(T). If $a \in V(S_2)$ and $r_2 \leq a - (|V(S_1) \sqcup V(S_2)| - k)$, then we must have been in the first case where $j \geq |V(T)| - k + 1$ and the inverse map is given by

(4.36)
$$\operatorname{list}(T) = \operatorname{list}(S_1) \cdot \operatorname{startr}(\operatorname{list}(S_2), r_2).$$

Note that because $r_2 \leq a - \epsilon - 1$, the r_2 -th smallest entry of S_2 can only be an a if $\epsilon = 0$, so $\operatorname{list}(T)$ is indeed a tree list of G. If $a \in V(S_2)$ and $r_2 \geq a - (|V(S_1) \sqcup V(S_2)| - k) + 1$, then we must have been in the second case where $a - k + 1 \leq j \leq |V(T)| - k$ and the inverse map is given by concatenating the permutations $\tau^{(1)} = \operatorname{list}(S_1)$ and $\tau^{(2)} = \operatorname{startr}(\operatorname{list}(S_2, a - |V(S_1)|))$ and then returning the entry a from the start of $\tau^{(2)}$ to position $j = a - r_2 + 1$, that is,

(4.37)
$$\operatorname{list}(T) = \tau_1^{(1)} \cdots \tau_{a-r_2}^{(1)} a \ \tau_{a-r_2+2}^{(1)} \cdots \tau_{|V(S_1)|}^{(1)} \ \tau_2^{(2)} \cdots \tau_{k-1}^{(2)}.$$

Note that because $r_2 \leq a - \epsilon - 1$, the entry *a* can only move to position j = 2 if $\epsilon = 0$, so list(T) is indeed a tree list of *G*. Finally, if $a \in V(S_1)$, then we must have been in the third case where $j \leq a - k$ and the inverse map is given by inserting $\tau^{(2)} = \text{startr}(\text{list}(S_2, r_2))$ into $\tau^{(1)} = \text{list}(S_1)$, specifically,

(4.38)
$$\operatorname{list}(T) = \tau_1^{(1)} \cdots \tau_{a-|V(S_1)|}^{(1)} \tau_1^{(2)} \cdots \tau_k^{(2)} \tau_{a-|V(S_1)|+1}^{(1)} \cdots \tau_{|V(S_1)|}^{(1)}$$

These constructions are inverse to each other, so we indeed have a sign-reversing involution.

$K_{64} = \underbrace{\begin{smallmatrix} 2 \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ $	
${\cal F}$	$arphi_{\subseteq [5]}(\mathcal{F})$
(136789, 6, 1), (2, 1, 1), (45, 2, 1)	(136789, 6, 1), (245, 12, 1)
(1, 1, 1), (23, 2, 2), (456789, 33, 2)	(132, 12, 1), (456789, 33, 2)
(1, 1, 1), (256789, 123, 1), (34, 2, 1)	(153, 12, 1), (246789, 123, 1)
(1, 1, 1), (264789, 6, 4), (35, 2, 2)	(145, 12, 1), (263789, 6, 4)
(1, 1, 1), (2463789, 7, 5), (5, 1, 1)	(14, 11, 1), (2365789, 7, 5)

FIGURE 9. Some simple forest triples \mathcal{F} of K_{64} and $\varphi_{\subset [5]}(\mathcal{F})$

Case 3: $\mathcal{F}|_{\geq a}$ is fixed under φ' , $\operatorname{seg}_{[a]}(\mathcal{F}) \geq 3$, and $\operatorname{seg}_{[a]}(\mathcal{T}_i) \leq \operatorname{seg}_{[a]}(\mathcal{F}) - 2$. We now define a sign-reversing involution $\varphi_{\subseteq [a-1]}$ on these \mathcal{F} by applying the easybreak and easyjoin maps used in the proof of Proposition 4.8, but now we carefully work around \mathcal{T}_i . Note that the tree triples present in [a] other than \mathcal{T}_i have all vertices in [a-1]. Because

(4.39)
$$\operatorname{seg}_{[a]}(\mathcal{F}) - \operatorname{seg}_{[a]}(\mathcal{T}_i) = \sum_{1 \le t \le i', t \ne i} \ell(\alpha^{(t)}) \ge 2,$$

either some such tree triple is breakable or there are at least two such tree triples.

First suppose that $\sum_{i+1 \leq t \leq i'} \ell(\alpha^{(t)}) \geq 2$, meaning that $i' \geq i+1$ and either $\mathcal{T}_{i'}$ is breakable, or $\mathcal{T}_{i'}$ is not breakable and in fact $i' \geq i+2$. Informally, this means there are at least two segments of trees occurring after T_i . If $\mathcal{T}_{i'}$ is breakable, then we define $\varphi_{\subset [a-1]}(\mathcal{F})$ by replacing $\mathcal{T}_{i'}$ by the two tree triples of easybreak $(\mathcal{T}_{i'})$, and if $\mathcal{T}_{i'}$ is not breakable, then we define $\varphi_{\subset [a-1]}(\mathcal{F})$ by replacing $\mathcal{T}_{i'-1}$ and $\mathcal{T}_{i'}$ by easyjoin $(\mathcal{T}_{i'-1}, \mathcal{T}_{i'})$. The map $\varphi_{\subset [a-1]}$ is a sign-reversing involution on these \mathcal{F} , it preserves type and $\alpha_1^{(1)}$, and it preserves weight by (4.10) and because it preserves the order of the trees.

Similarly, suppose that i' = i, and either \mathcal{T}_{i-1} is not breakable, or \mathcal{T}_{i-1} is breakable with easybreak $(\mathcal{T}_{i-1}) = (\mathcal{S}_1, \mathcal{S}_2 = (S_2, \beta^{(2)}, r'))$, and $\min(V(S_2)) < \min(V(T_i))$. Then we again define $\varphi_{\subseteq [a-1]}(\mathcal{F})$ by either replacing \mathcal{T}_{i-1} by the two tree triples of easybreak (\mathcal{T}_{i-1}) if \mathcal{T}_{i-1} is breakable, or by replacing \mathcal{T}_{i-2} and \mathcal{T}_{i-1} by easyjoin $(\mathcal{T}_{i-2}, \mathcal{T}_{i-1})$ otherwise. The map $\varphi_{\subset [a-1]}$ is a sign-reversing involution on these \mathcal{F} , it preserves type and $\alpha_1^{(1)}$, and it preserves weight by (4.10) and because it preserves the order of the trees.

It remains to find a bijection $\varphi_{\subseteq[a-1]}$ between the case where i' = i + 1 and \mathcal{T}_{i+1} is not breakable, and the case where i' = i and \mathcal{T}_{i-1} is breakable with easybreak $(\mathcal{T}_{i-1}) = (\mathcal{S}_1, \mathcal{S}_2 = (S_2, \beta^{(2)}, r'))$, but this time $\min(V(T_i)) < \min(V(S_2))$. Then we get a sign-reversing involution by either applying $\varphi_{\subseteq[a-1]}$ or its inverse. Suppose that \mathcal{F} is in the former case.

We would like to join \mathcal{T}_{i+1} to \mathcal{T}_{i-1} , but the problem is that when we compute weight, the entries of T_{i+1} would now be read before the entries of T_i . The idea is to first move \mathcal{T}_{i+1} "through" \mathcal{T}_i , preserving the relative order of entries x with $\min(V(T_i)) < x < a$, before joining to \mathcal{T}_{i-1} . Let $k = |V(T_i) \cap [a]|, \sigma = \operatorname{list}(T_i), \sigma' = \operatorname{startr}(\operatorname{list}(T_{i+1}), r_{i+1})$, and let $2 \leq j \leq k$ be such that $\sigma_j = a$. We now define

(4.40)
$$\tau = \sigma_2 \cdots \sigma_{j-1} \sigma_{j+1} \cdots \sigma_k \sigma'_1 \cdots \sigma'_{\ell(\sigma')},$$

the permutation consisting of the entries x in $V(T_i) \sqcup V(T_{i+1})$ with $\min(V(T_i)) < x < a$, which in particular are all adjacent. We use the first $|V(T_{i+1})|$ entries of τ to join with \mathcal{T}_{i-1} to make the tree triple

(4.41)
$$\mathcal{T} = (\text{list}(T_{i-1}) \cdot (\tau_1 \cdots \tau_{|V(T_{i+1})|}), \alpha^{(i-1)} \cdot \alpha^{(i+1)}, r_{i-1}),$$

and we use the remaining entries of τ to modify \mathcal{T}_i to make the tree triple

(4.42) $\mathcal{T}'_i = (\sigma_1 \ \tau_{|V(T_{i+1})|+1} \ \cdots \ \tau_{|V(T_{i+1})|+j-2} \ a \ \tau_{|V(T_{i+1})|+j-1} \ \cdots \ \tau_{\ell(\tau)} \ \sigma_{k+1} \ \cdots \ \sigma_{\ell(\sigma)}, \alpha^{(i)}, r_i).$ Then we define $\varphi_{\subseteq [a-1]}(\mathcal{F})$ by replacing by tree triples $\mathcal{T}_{i-1}, \ \mathcal{T}_i$, and \mathcal{T}_{i+1} by \mathcal{T} and \mathcal{T}'_i . The entries that moved are in [a-1] so are adjacent in G and these are indeed tree triples. Note that entries of τ occurring after a, which make a G-inversion with a and not σ_1 , are exchanged with entries occurring before σ_1 , which make a G-inversion with σ_1 and not a. Therefore, the map $\varphi_{\subseteq [a-1]}$ preserves weight, and by construction it also reverses sign and preserves type and $\alpha_1^{(1)}$. We have indeed produced a forest triple for which $i' = i, \ \mathcal{T}_{i-1}$ is breakable with easybreak $(\mathcal{T}_{i-1}) = (\mathcal{S}_1, \mathcal{S}_2 = (S_2, \beta^{(2)}, r'))$, and $\min(V(T_i)) < \min(V(S_2))$.

To define the inverse map, we break \mathcal{T}_{i-1} as easybreak $(\mathcal{T}_{i-1}) = (\mathcal{S}_1, \mathcal{S}_2 = (S_2, \beta^{(2)}, r'))$ and move \mathcal{S}_2 back "through" \mathcal{T}_i . Let $k = |V(T_i) \cap [a]|, \sigma = \text{list}(T_i), \sigma' = \text{startr}(\text{list}(S_2), r')$, and let $2 \leq j \leq k$ be such that $\sigma_j = a$. We now define

(4.43)
$$\tau = \sigma'_1 \cdots \sigma'_{\ell(\sigma')} \sigma_2 \cdots \sigma_{j-1} \sigma_{j+1} \cdots \sigma_k,$$

the permutation consisting of the entries x in $V(S_2) \sqcup V(T_i)$ with $\min(V(T_i)) < x < a$. We use the first (k-2) entries of τ to modify \mathcal{T}_i to make the tree triple

(4.44)
$$\mathcal{T}'_{i} = (\sigma_{1} \ \tau_{1} \ \cdots \ \tau_{j-2} \ a \ \tau_{j-1} \ \cdots \ \tau_{k-2} \ \sigma_{k+1} \ \cdots \ \sigma_{\ell(\sigma)}, \alpha^{(i)}, r_{i}),$$

and we use the remaining entries of τ to make the tree triple

(4.45)
$$\mathcal{S}'_2 = (\operatorname{startmin}(\tau_{k-1} \cdots \tau_{\ell(\tau)}), \beta^{(2)}, \operatorname{indstart}(\tau_{k-1} \cdots \tau_{\ell(\tau)})).$$

Then the inverse map is given by replacing the tree triples \mathcal{T}_{i-1} and \mathcal{T}_i by $\mathcal{S}_1, \mathcal{T}'_i$, and \mathcal{S}'_2 .



(1, 1, 1), (23465789, 35, 1)

(1, 1, 1), (23456789, 35, 2)

(1, 1, 1), (25634789, 341, 3)

FIGURE 10. Some simple forest triples \mathcal{F} of K_{64} and $\varphi_{>3}(\mathcal{F})$

Case 4: $\mathcal{F}|_{\geq a}$ is fixed under φ' , $\operatorname{seg}_{[a]}(\mathcal{F}) \geq 3$, and $\operatorname{seg}_{[a]}(\mathcal{T}_i) - \operatorname{seg}_{[a]}(\mathcal{F}) = 0$ or 1. We define a bijection $\varphi_{\geq 3}$ between such \mathcal{F} with $\operatorname{seg}_{[a]}(\mathcal{T}_i) = \operatorname{seg}_{[a]}(\mathcal{F})$, meaning that the only tree triple present in [a] is $\mathcal{T}_1 = (T_1, \alpha^{(1)}, 1)$, where $[a] \subseteq V(T_1)$, $\ell(\alpha^{(1)}) \geq 3$, and $\alpha_1^{(1)} + \alpha_2^{(1)} < a$; and those with $\operatorname{seg}_{[a]}(\mathcal{T}_i) = \operatorname{seg}_{[a]}(\mathcal{F}) - 1$, meaning there is exactly one other tree triple present in [a] and it is not breakable. Then we get a sign-reversing involution by either applying $\varphi_{\geq 3}$ or its inverse. Let $\sigma = \operatorname{list}(T_1)$ and let j be such that $\sigma_j = a$. Note that $2 \leq j \leq a$ by Lemma 4.4 and we can have j = 2 only if $\epsilon = 0$. We will have two cases, depending on the position of the entry a in σ .

If
$$j \leq a - \alpha_2^{(1)}$$
, then we remove $\alpha_2^{(1)}$ entries from the inside of σ and define

(4.46)
$$\sigma^{\text{in}} = \sigma_{a-\alpha_2^{(1)}+1} \cdots \sigma_a \text{ and } \sigma^{\text{out}} = \sigma_1 \cdots \sigma_{a-\alpha_2^{(1)}} \sigma_{a+1} \cdots \sigma_{|V(T)|},$$

and we define $\varphi_{\geq 3}(\mathcal{F})$ by replacing \mathcal{T}_1 by the tree triples

(132456789, 135, 1)

(123465789, 135, 1)

145623789, 1341, 1

(4.47)
$$S_1 = (\sigma^{\text{out}}, \alpha_1^{(1)} \alpha_3^{(1)} \cdots \alpha_\ell^{(1)}, 1) \text{ and } S_2 = (\operatorname{startmin}(\sigma^{\text{in}}), \alpha_2^{(1)}, \operatorname{indstart}(\sigma^{\text{in}})).$$

Because $j \leq a - \alpha_2^{(1)}$, the entry *a* appears in σ^{out} so its descents and successive LR maxima appear in σ , and startmin(σ^{in}) has all entries in [a - 1] so these are tree lists of *G*, and we have $r_2 \leq \ell(\sigma^{\text{in}}) = \alpha_2^{(1)}$, so \mathcal{S}_2 is indeed a tree triple of *G*. Moving the string σ^{in} does not affect the number of *G*-inversions because its entries are less than *a*, whereas every $\sigma_t > a$ for t > a by Lemma 4.4. The map $\varphi_{\geq 3}$ reverses sign, preserves type, $\alpha_1^{(1)}$, and weight. Also note that the vertices 1 and *a* are in the same tree.

If $a - \alpha_2^{(1)} + 1 \le j \le a$, which in particular means that $j \ne 2$ because $\alpha_1^{(1)} + \alpha_2^{(1)} < a$, then we separate σ into its head and tail,

(4.48)
$$\sigma^{\text{head}} = \sigma_1 \cdots \sigma_{\alpha_1^{(1)}} \text{ and } \sigma^{\text{tail}} = \sigma_{\alpha_1^{(1)}+1} \cdots \sigma_{|V(T)|}.$$

As before, we apply transpositions to σ^{tail} to put its minimum element in front. We will now do something a little unusual. We move the entry *a* to position $(a - \text{indstart}(\sigma^{\text{tail}}) + 1)$ so that it makes $(\text{indstart}(\sigma^{\text{tail}}) - 1)$ *G*-inversions to account for these transpositions. We then use the new value of r = a - j + 1 to account for the original (a - j) *G*-inversions made by the entry *a*. Specifically, let $r' = \text{indstart}(\sigma^{\text{tail}})$ and let τ be startmin (σ^{tail}) with the entry *a* removed, and define $\varphi_{>3}(\mathcal{F})$ by replacing \mathcal{T}_1 by the tree triples

(4.49)
$$S_1 = (\sigma^{\text{head}}, \alpha_1^{(1)}, 1)$$
 and $S_2 = (\tau_1 \cdots \tau_{a-r'} \ a \ \tau_{a-r'+1} \cdots \tau_{\ell(\tau)}, \alpha_2^{(1)} \cdots \alpha_{\ell}^{(1)}, a-j+1).$

The entries of σ^{head} and the entries of τ that moved are all in [a-1] so are adjacent and these are tree lists of G. We also have $a - j + 1 \leq \alpha_1^{(2)}$, so S_2 is indeed a tree triple of G. The map $\varphi_{\geq 3}$ reverses sign and preserves type, $\alpha_1^{(1)}$, and weight. Also note that the vertices 1 and a are in different trees.

It remains to show that the map $\varphi_{\geq 3}$ is a bijection. Suppose that $\operatorname{seg}_{[a]}(\mathcal{T}_i) = \operatorname{seg}_{[a]}(\mathcal{F}) - 1$. The idea is that we can determine which case of the map was applied by whether the vertices 1 and a are in the same tree in \mathcal{F} , so we can recover the original tree triple.

If the vertices 1 and a are in the same tree $T_i = T_1$, then we must have been in the first case where $j \leq a - \alpha_2^{(1)}$. Let $\sigma = \text{list}(T_1)$, $\sigma' = \text{startr}(\text{list}(T_2), r_2)$, $k = |V(T_2) \cap [a]|$, and then the inverse map is given by inserting σ' back into σ , specifically, by replacing \mathcal{T}_1 and \mathcal{T}_2 by

(4.50)
$$\mathcal{T} = (\sigma_1 \cdots \sigma_k \sigma'_1 \cdots \sigma'_{|V(T_2)|} \sigma_{k+1} \cdots \sigma_{|V(T_1)|}, \alpha_1^{(1)} \alpha_1^{(2)} \alpha_2^{(1)} \cdots \alpha_\ell^{(1)}, 1)$$

If the vertices 1 and a are in different trees, so that $1 \in V(T_1)$ and $a \in V(T_2)$, then we must have been in the second case where $a - \alpha_2^{(1)} + 1 \leq j \leq a$. Let $\sigma' = \text{list}(T_1)$, let $\sigma = \text{list}(T_2)$, let $k = |V(T_2) \cap [a]|$, let j' be such that $\sigma_{j'} = a$, and let τ be startr(list(T_2), k - j' + 1) with the entry a removed. The position of the entry a tells us the original first entry of the tail and the value of r_2 tells us the original position of the entry a. The inverse map is given by replacing \mathcal{T}_1 and \mathcal{T}_2 by

(4.51)
$$\mathcal{T} = (\sigma'_1 \cdots \sigma'_{|V(T_2)|} \tau_1 \cdots \tau_{k-r_2-1} a \tau_{k-r_2} \cdots \tau_{\ell(\tau)}, \alpha_1^{(1)} \alpha_1^{(2)} \cdots \alpha_{\ell}^{(2)}, 1).$$

These constructions are inverse to each other, so we indeed have a sign-reversing involution in this case.

By combining our sign-reversing involutions in these four cases, we build a nice involution φ for G with the prescribed fixed points, thus completing the proof.

Now we can repeatedly apply Theorem 4.10 to obtain a combinatorial description of fixed forest triples for any *almost-K-chain*, which is a graph of the form

(4.52)
$$K_{\gamma}^{\epsilon} = K_{\gamma_1}^{\epsilon_1} + \dots + K_{\gamma_{\ell}}^{\epsilon_{\ell}},$$

where $\gamma = \gamma_1 \cdots \gamma_\ell$ is a composition with all parts at least 2 and $\epsilon = \epsilon_1 \cdots \epsilon_\ell$ is a sequence of 0's and 1's. Aliniaeifard, Wang, and van Willigenburg proved that for such graphs, the chromatic symmetric function $X_{K_{\gamma}^{\epsilon}}(\boldsymbol{x})$ is *e*-positive [7, Proposition 5.4].

Definition 4.11. Let K^{ϵ}_{γ} be an almost-K-chain and for $0 \leq t \leq \ell(\gamma)$, let

$$(4.53) c_t = \gamma_1 + \dots + \gamma_t - t + 1,$$

so that $c_0 = 1$, $c_{\ell(\gamma)} = n$, and $c_1, \ldots, c_{\ell(\gamma)-1}$ are the cut vertices of K^{ϵ}_{γ} . Let $\operatorname{Fix}(K^{\epsilon}_{\gamma})$ denote the set of forest triples $\mathcal{F} \in \operatorname{FT}(K^{\epsilon}_{\gamma})$ that are atoms and such that for every $1 \leq t \leq \ell(\gamma)$, letting $\mathcal{T} = (T, \alpha, r)$ denote the tree triple of \mathcal{F} with $c_t \in V(T)$, we have the following.

- (C1) There is at most one other tree triple $\mathcal{T}' = (T', \alpha', r')$ in \mathcal{F} such that there exists a vertex $v \in V(T')$ with $c_{t-1} \leq v \leq c_t$.
- (C2) If $c_{t-1} \notin V(T)$, we have $r \ge \gamma_t \epsilon_t$.
- (C3) If $c_{t-1} \in V(T)$, we have $|\{v \in V(T) : v \ge c_{t-1}\}| \ge \gamma_t$.

Let $\operatorname{Fix}_{\operatorname{simple}}(K^{\epsilon}_{\gamma})$ denote the set of simple forest triples $\mathcal{F} \in \operatorname{Fix}(K^{\epsilon}_{\gamma})$.

Corollary 4.12. The chromatic quasisymmetric function of an almost-K-chain K^{ϵ}_{γ} is

$$(4.54) \quad X_{K^{\epsilon}_{\gamma}}(\boldsymbol{x};q) = \sum_{\mathcal{F} \in \mathrm{Fix}_{\mathrm{simple}}(K^{\epsilon}_{\gamma})} [\alpha_{1}^{(1)}(\mathcal{F})]_{q} \ q^{\mathrm{weight}(\mathcal{F})} e_{\mathrm{type}(\mathcal{F})} = \sum_{\mathcal{F} \in \mathrm{Fix}(K^{\epsilon}_{\gamma})} q^{\mathrm{weight}(\mathcal{F})} e_{\mathrm{type}(\mathcal{F})}.$$

In particular, $X_{K^{\epsilon}_{\gamma}}(\boldsymbol{x};q)$ is *e*-positive.

Proof. We use induction on $\ell(\gamma)$ to show that there is a nice involution whose fixed points are exactly $\operatorname{Fix_{simple}}(K_{\gamma}^{\epsilon})$. The case where $\ell(\gamma) = 1$ is done in Proposition 4.8, so suppose that $\ell(\gamma) \geq 2$ and let $G' = K_{\gamma_2}^{\epsilon_2} + \cdots + K_{\gamma_{\ell}}^{\epsilon_{\ell}}$. By Theorem 4.10 and our induction hypothesis, there is a nice involution φ for $K_{\gamma}^{\epsilon} = K_{\gamma_1}^{\epsilon_1} + G'$ with fixed points exactly

(4.55) Fix(
$$\varphi$$
) = { $\mathcal{F} \in \operatorname{FT}_{\operatorname{simple}}(K_{\gamma}^{\epsilon})$: \mathcal{F} is an atom, $\mathcal{F}|_{\geq \gamma_1} \in \operatorname{Fix}_{\operatorname{simple}}(G')$, $\operatorname{seg}_{[\gamma_1]}(\mathcal{F}) \leq 2$,
if $1 \notin V(T_i)$, then $r_i \geq \gamma_1 - \epsilon_1$, if $1 \in V(T_i)$, then $|V(T_i)| \geq \gamma_1$ },

where $\mathcal{T}_i = (T_i, \alpha^{(i)}, r_i)$ is the tree triple of \mathcal{F} with $\gamma_1 \in V(T_i)$. Note that by induction, the condition $\mathcal{F}|_{\geq \gamma_1} \in \operatorname{Fix}_{\operatorname{simple}}(G')$ means that for the restricted tree triple $\mathcal{T}_i|_{\geq \gamma_1} = (T', \alpha', 1)$, if $\gamma_2 \in V(T')$, then we must have $|V(T')| \geq \gamma_2$, which means that for the tree triple \mathcal{T}_i , if $c_1 \in V(T_i)$, we must have $|\{v \in V(T_i) : v \geq c_1\}| \geq \gamma_2$, because it is only these vertices of T_i that appear in the restriction to G'. The other conditions of $\operatorname{Fix}(K_{\gamma}^{\epsilon})$ are exactly those in (4.55), so we have $\operatorname{Fix}(\varphi) = \operatorname{Fix}_{\operatorname{simple}}(K_{\gamma}^{\epsilon})$, as desired. \Box

Now by enumerating these fixed points, we can simplify (4.54) to a more explicit expression. We first state a Lemma about *q*-counting subsets.

Lemma 4.13. For integers $0 \le k \le n$, we have

(4.56)
$$\sum_{S \subseteq [n]: |S|=k} q^{|\{(i,j): i>j, i \notin S, j \in S\}|} = \binom{n}{k}_q,$$

where the *q*-binomial coefficient is $\binom{n}{k}_q = \frac{[n]_q!}{[k]_q![n-k]_q!}$.

Proof. As with ordinary binomial coefficients, the result follows from induction on n, the identity $\binom{n}{k}_q = q^k \binom{n-1}{k}_q + \binom{n-1}{k-1}_q$, and from considering the cases of $n \notin S$ and $n \in S$. \Box

Before we prove our explicit formula for $X_{K_{\gamma}^{\epsilon}}(\boldsymbol{x};q)$, we will first consider the simpler case of a K-chain with two cliques because this will demonstrate the main ideas.

Corollary 4.14. Consider the *K*-chain K_{ab} with two cliques of sizes *a* and *b*, which has n = a + b - 1 vertices. Then the chromatic quasisymmetric function of K_{ab} is

(4.57)
$$X_{K_{ab}}(\boldsymbol{x};q) = [a-1]_q! [b-1]_q! \sum_{k=\max\{a,b\}}^{n} q^{n-k} [2k-n]_q e_{k(n-k)}.$$

Proof. We count fixed forest triples $\mathcal{F} \in \operatorname{Fix}(K_{ab})$ by weight. Let $\mathcal{T} = (T, \alpha, r)$ be the tree triple of \mathcal{F} with $n \in V(T)$ and let k = |V(T)|. We must have $a \in V(T)$, otherwise $V(T) \subseteq \{a + 1, \ldots, n\}$ and $r \leq k \leq b - 1$, while (C2) requires $r \geq b$. Now by (C3), we must have $\{a, \ldots, n\} \subseteq V(T)$ and $k \geq b$. If $1 \notin V(T)$, (C2) implies that $k \geq r \geq a$, while if $1 \in V(T)$, (C3) again implies that $k \geq a$, so we must have $k \geq \max\{a, b\}$. By (C1), there is at most one other tree triple $\mathcal{T}' = (T', \alpha', r')$ in \mathcal{F} , and we have |V(T')| = n - k and $V(T') \subseteq [a-1]$. We now consider the possibilities of \mathcal{F} with $1 \in V(T')$ and with $1 \in V(T)$.

If $1 \in V(T')$, then we must have $r \geq a$ by (C2). By using Lemma 3.13 to account for *G*-inversions within the trees *T* and *T'* and using Lemma 4.13 to account for *G*-inversions between trees *T'* and *T*, the *q*-weighted sum over these fixed points is

(4.58)
$$\sum_{\substack{\mathcal{F}\in \mathrm{Fix}(K_{ab})\\|V(T)|=k\\1\in V(T')}} q^{\mathrm{weight}(\mathcal{F})} = \binom{a-2}{k-b}_q [n-k-1]_q! [k-b]_q! [b-1]_q! [n-k]_q ([k]_q - [a-1]_q) = q^{a-1} [a-2]_q! [b-1]_q! [n-k]_q [k-a+1]_q.$$

If $1 \in V(T)$, then we can have any $1 \leq r \leq k$. We have (n-k) *G*-inversions between vertex $n \in V(T)$ and the vertices in T'. By Lemma 3.13 and Lemma 4.13, we have

(4.59)
$$\sum_{\substack{\mathcal{F}\in \operatorname{Fix}(K_{ab})\\|V(T)|=k\\1\in V(T)}} q^{\operatorname{weight}(\mathcal{F})} = q^{n-k} \binom{a-2}{n-k}_{q} [n-k-1]_{q}! [k-b]_{q}! [b-1]_{q}! [n-k]_{q} [k]_{q}$$
$$= q^{n-k} [a-2]_{q}! [b-1]_{q}! [k-b]_{q} [k]_{q}.$$

FIGURE 11. The K-chain K_{66} and the chromatic quasisymmetric function $X_{K_{66}}(\boldsymbol{x};q)$



 $(4.60) \ X_{K_{66}}(\boldsymbol{x};q) = [5]_q! [5]_q! (q^5 e_{65} + q^4 [3]_q e_{74} + q^3 [5]_q e_{83} + q^2 [7]_q e_{92} + q [9]_q e_{(10)1} + [11]_q e_{(11)}).$

Now by summing (4.58) and (4.59) over all $\max\{a, b\} \le k \le n$, we have as desired

$$X_{K_{ab}}(\boldsymbol{x};q) = [a-2]_q![b-1]_q! \sum_{k=\max\{a,b\}}^n (q^{a-1}[n-k]_q[k-a+1]_q + q^{n-k}[k-b]_q[k]_q)e_{k(n-k)}$$
$$= [a-1]_q![b-1]_q! \sum_{k=\max\{a,b\}}^n q^{n-k}[2k-n]_qe_{k(n-k)}.$$

We now state the general formula for any almost-K-chain.

Definition 4.15. Let $\gamma = \gamma_1 \cdots \gamma_\ell$ be a composition with all parts at least 2 and let $\epsilon = \epsilon_1 \cdots \epsilon_\ell$ be a list of 0's and 1's. We define A^{ϵ}_{γ} to be the set of weak compositions $\alpha = \alpha_1 \cdots \alpha_{\ell+1}$ of length $\ell + 1$ and size $|\gamma| - \ell + 1$ such that $\alpha_1 \ge 1$ and for each $2 \le i \le \ell + 1$ we have either

(4.61)
$$\alpha_i < \gamma_{i-1} - \epsilon_{i-1} - 1 \text{ and } \alpha_i + \dots + \alpha_{\ell+1} < \gamma_i + \dots + \gamma_\ell - (\ell - i), \text{ or}$$

(4.62)
$$\alpha_i \ge \gamma_{i-1} - \epsilon_{i-1} \text{ and } \alpha_i + \dots + \alpha_{\ell+1} \ge \gamma_i + \dots + \gamma_\ell - (\ell - i).$$

Remark 4.16. Note that for $i = \ell + 1$, the conditions are that $\alpha_{\ell+1} = 0$ or $\alpha_{\ell+1} \ge \gamma_{\ell} - \epsilon_{\ell}$. In particular, if $\epsilon_{\ell} = 0$, we must have $\alpha_{\ell+1} = 0$, otherwise these conditions will force $\alpha_{\ell+1} \ge \gamma_{\ell}$ and $\alpha_i \ge \gamma_{i-1} - 1$ for every $2 \le i \le \ell$, but then $|\alpha| \ge |\gamma| - \ell + 2$.

Example 4.17. If $\gamma = abc$ and $\epsilon = 000$, then A_{abc}^{000} is the set of weak compositions $\alpha = \alpha_1 \alpha_2 \alpha_3 \alpha_4$ such that $\alpha_1 + \alpha_2 + \alpha_3 + \alpha_4 = a + b + c - 2$, $\alpha_1 \ge 1$, $\alpha_4 = 0$ by Remark 4.16, and

(4.63)
$$\alpha_2 < a - 1 \text{ and } \alpha_2 + \alpha_3 < b + c - 1, \text{ or } \alpha_2 \ge a \text{ and } \alpha_2 + \alpha_3 \ge b + c - 1,$$

(4.64) $\alpha_3 < b-1 \text{ and } \alpha_3 < c, \text{ or } \alpha_3 \ge b \text{ and } \alpha_3 \ge c.$

Corollary 4.18. The chromatic quasisymmetric function of an almost-K-chain K^{ϵ}_{γ} is

(4.65)
$$X_{K_{\gamma}^{\epsilon}}(\boldsymbol{x};q) = [\gamma_{1}-2]_{q}! \cdots [\gamma_{\ell}-2]_{q}! \sum_{\alpha \in A_{\gamma}^{\epsilon}} [\alpha_{1}]_{q} \prod_{i=2}^{\ell+1} q^{m_{i}} [|\alpha_{i}-(\gamma_{i-1}-1-\epsilon_{i-1})|]_{q} e_{\operatorname{sort}(\alpha)},$$

where $\ell = \ell(\gamma)$ and $m_i = \min\{\alpha_i, \gamma_{i-1} - 1 - \epsilon_{i-1}\}.$

Proof. The idea is to associate a weak composition $\alpha \in A^{\epsilon}_{\gamma}$ to each fixed point $\mathcal{F} \in \operatorname{Fix}(K^{\epsilon}_{\gamma})$ and then to enumerate $q^{\text{weight}(\mathcal{F})}$ over all \mathcal{F} corresponding to a particular α . Given a fixed forest triple \mathcal{F} , we define $\alpha = \alpha_1 \cdots \alpha_{\ell+1}$ as follows. For $1 \leq i \leq \ell+1$, let T_i be the tree with $c_{i-1} \in V(T_i)$. Note that different indices i may refer to the same tree because for example it is possible that $c_{i-2} \in V(T_i)$ as well. Let $\alpha_1 = |V(T_1)| \ge 1$ be the size of the tree containing vertex 1. Then for $2 \leq i \leq \ell + 1$, if $c_{i-2} \notin V(T_i)$, we let $\alpha_i = |V(T_i)|$ be the size of the tree T_i , while if $c_{i-2} \in V(T_i)$, then there is at most one other tree T'_i with $C_i \cap V(T'_i) \neq \emptyset$; we let $\alpha_i = |V(T'_i)|$ be the size of this other tree T'_i if there is one, otherwise we set $\alpha_i = 0$. Note that all trees have been considered, so $|\alpha| = n = |\gamma| - \ell + 1$.

If $c_{i-2} \notin V(T_i)$, meaning that $\alpha_i = |V(T_i)|$, then we must have $\alpha_i \geq \gamma_{i-1} - \epsilon_{i-1}$ by (C2) and the trees T_j for $j \ge i$ must use all the vertices at least c_{i-1} , so we must have

(4.66)
$$\alpha_i + \dots + \alpha_{\ell+1} \ge \gamma_i + \dots + \gamma_\ell - (\ell - i).$$

If $c_{i-2} \in V(T_i)$, meaning that $\alpha_i = |V(T'_i)|$ is the size of the other tree T'_i present in C_{i-1} , if any, then we must have $V(T_{i'}) \subseteq \{c_{i-2}+1,\ldots,c_{i-1}-1\}$ so $\alpha_i < \gamma_{i-1}-1$, the tree T_i must contain at least γ_{i-2} vertices at least c_{i-2} by (C3), and therefore the tree T'_i along with the trees to the right of T_i use the remaining vertices at least c_{i-2} , so we must have

(4.67)
$$\alpha_i + \dots + \alpha_{\ell+1} < \gamma_i + \dots + \gamma_\ell - (\ell - i)$$

Also note that if $\epsilon_{i-1} = 0$ and $\alpha_i = \gamma_{i-1} - \epsilon_{i-1} - 1$, then the summand in (4.65) is zero so we can equivalently replace the condition $\alpha_i < \gamma_{i-1} - 1$ by the condition $\alpha_i < \gamma_{i-1} - \epsilon_{i-1} - 1$, and therefore the weak compositions that arise are exactly those $\alpha \in A^{\epsilon}_{\gamma}$. We now enumerate $q^{\operatorname{weight}(\mathcal{F})}$ over all \mathcal{F} associated to a particular $\alpha \in A^{\epsilon}_{\gamma}$. The factor $[\alpha_1]_q$ accounts for the possible choices of $1 \le r_1 \le \alpha_1$. Now for $2 \le i \le \ell + 1$, we consider the contributions given by the choices for r_i and the inversion-weighted choices of how to join vertices in the clique C_{i-1} .

If $c_{i-2} \notin V(T_i)$, then we must have $r_i \geq \gamma_{i-1} - \epsilon_{i-1}$. Let $k = \alpha_1 + \cdots + \alpha_{i-1} - c_{i-2} + 1$ be the number of vertices of T_i in C_{i-1} . Then by Lemma 3.13 and Lemma 4.13, the contribution given by the possible choices in C_{i-1} is

(4.68)
$$\binom{\gamma_{i-1}-2}{k-1}_{q} [k-1]_{q}! [\gamma_{i-1}-k-1]_{q}! ([\alpha_{i}]_{q}-[\gamma_{i-1}-1-\epsilon_{i-1}]_{q})$$
$$= q^{\gamma_{i-1}-1-\epsilon_{i-1}} [\gamma_{i-1}-2]_{q}! [\alpha_{i}-(\gamma_{i-1}-1-\epsilon_{i-1})]_{q}.$$

If $c_{i-2} \in V(T_i)$, then we can have any $1 \leq r_i \leq \alpha_i$. We have α_i G-inversions between vertex $c_{i-1} \in V(T_i)$ and the vertices in $T_{i'}$. By Lemma 3.13 and Lemma 4.13, the contribution given by the possible choices in C_{i-1} is

(4.69)
$$q^{\alpha_{i}} \binom{\gamma_{i-1}-2}{\alpha_{i}}_{q} [\alpha_{i}-1]_{q}! [\gamma_{i-1}-2-\alpha_{i}]_{q}! [\gamma_{i-1}-1-\epsilon_{i-1}-\alpha_{i}]_{q} [\alpha_{i}]_{q} = q^{\alpha_{i}} [\gamma_{i-1}-2]_{q}! [(\gamma_{i-1}-1-\epsilon_{i-1})-\alpha_{i}]_{q}.$$

Note that if $\alpha_i = 0$, meaning there is no tree T'_i present, the expression in (4.69) is still valid. Therefore, in either case, we can use the absolute value to express the contribution as

(4.70)
$$q^{m_i}[\gamma_{i-1}-2]_q![|\alpha_i-(\gamma_{i-1}-1-\epsilon_{i-1})|]_q$$

Multiplying these contributions over all $2 \le i \le \ell + 1$, we have as desired

$$(4.71) \qquad X_{K_{\gamma}^{\epsilon}}(\boldsymbol{x};q) = \sum_{\alpha \in A_{\gamma}^{\epsilon}} [\alpha_{1}]_{q} \prod_{i=2}^{\ell+1} q^{m_{i}} [\gamma_{i-1} - 2]_{q}! [|\alpha_{i} - (\gamma_{i-1} - 1 - \epsilon_{i-1})|]_{q} e_{\operatorname{sort}(\alpha)}$$
$$= [\gamma_{1} - 2]_{q}! \cdots [\gamma_{\ell} - 2]_{q}! \sum_{\alpha \in A_{\gamma}^{\epsilon}} [\alpha_{1}]_{q} \prod_{i=2}^{\ell+1} q^{m_{i}} [|\alpha_{i} - (\gamma_{i-1} - 1 - \epsilon_{i-1})|]_{q} e_{\operatorname{sort}(\alpha)}.$$

Our explicit formula also allows us to confirm e-unimodality for almost-K-chains. We make the following elementary observation.

Observation 4.19. Let $a(q) = \sum_{k=i}^{j} a_k q^k \in \mathbb{N}[q]$ and $b(q) = \sum_{k=i'}^{j'} b_k q^k \in \mathbb{N}[q]$ be positive palindromic unimodal polynomials with centers of symmetry $\frac{i+j}{2}$ and $\frac{i'+j'}{2}$.

- (O1) The product a(q)b(q) is a positive palindromic unimodal polynomial with center of symmetry $\frac{i+j}{2} + \frac{i'+j'}{2}$.
- (O2) If $\frac{i+j}{2} = \frac{i'+j'}{2}$, then the sum a(q) + b(q) is a positive palindromic unimodal polynomial with the same center of symmetry.

Corollary 4.20. The chromatic quasisymmetric function of an almost-K-chain K_{γ}^{ϵ} is *e*-positive and *e*-unimodal.

Proof. By (4.65), the coefficient of e_{μ} in $X_{K_{\gamma}^{\epsilon}}(\boldsymbol{x};q)$ is

$$(4.72) \quad c_{\mu}(q) = [\gamma_1 - 2]_q! \cdots [\gamma_{\ell} - 2]_q! \sum_{\alpha \in A_{\gamma}^{\epsilon}: \text{ sort}(\alpha) = \mu} [\alpha_1]_q \prod_{i=2}^{\ell+1} q^{m_i} [|\alpha_i - (\gamma_{i-1} - 1 - \epsilon_{i-1})|]_q.$$

Note that regardless of whether $\alpha_i < \gamma_{i-1} - 1 - \epsilon_{i-1}$ or $\alpha_i \ge \gamma_{i-1} - 1 - \epsilon_{i-1}$, the factor $q^{m_i}[|\alpha_i - (\gamma_{i-1} - 1 - \epsilon_{i-1})|]_q$ is a positive palindromic unimodal polynomial with center of symmetry $\frac{\alpha_i + \gamma_{i-1} - 2 - \epsilon_{i-1}}{2}$. Therefore, by (O1), each summand in (4.72) is a positive palindromic unimodal polynomial with center of symmetry

(4.73)
$$\frac{\alpha_1 - 1}{2} + \sum_{i=2}^{\ell+1} \frac{\alpha_i + \gamma_{i-1} - 2 - \epsilon_{i-1}}{2} = \frac{|\alpha| + |\gamma| - 2\ell - 1 - |\epsilon|}{2} = \frac{2|\gamma| - 3\ell - |\epsilon|}{2},$$

which does not depend on α , and therefore we can apply (O2) to see that the sum is a positive palindromic unimodal polynomial with the same center of symmetry. Then we apply (O1) again to see that $c_{\mu}(q)$ is positive and unimodal.

5. Further directions

We conclude by proposing some further avenues of study for the Stanley–Stembridge conjecture. First, we can continue to seek sign-reversing involutions on forest triples to prove *e*-positivity and *e*-unimodality for other graphs. The author checked by computer that every natural unit interval graph with at most 9 vertices has a nice involution.

Problem 5.1. Let G be a natural unit interval graph. Find a nice involution for G to prove that $X_G(\boldsymbol{x};q)$ is *e*-positive. Furthermore, by enumerating the fixed points, find an explicit *e*-expansion to prove that $X_G(\boldsymbol{x};q)$ is *e*-unimodal.

For example, Dahlberg [22, Theorem 4.5] proved that for the class of triangular ladders

(5.1)
$$P_{n,2} = ([n], \{\{i, j\} : 1 \le j - 2 \le i \le j \le n\})$$

the chromatic symmetric function $X_{P_{n,2}}(\boldsymbol{x})$ is *e*-positive. We could try to strengthen this result by finding a nice involution to prove a quasisymmetric version. More specifically, because triangular ladders are built by successively joining a new vertex to two vertices, we could take an inductive approach.

Problem 5.2. Let G = ([n], E) be a natural unit interval graph with $\{1, 2\} \in E$ and let φ be a nice involution for G. Find a nice involution for the natural unit interval graph

(5.2)
$$([n+1], \{\{1,2\}, \{1,3\}\}) \cup \{\{i+1, j+1\}: \{i,j\} \in E\})$$

Furthermore, by enumerating the fixed points, prove that $X_{P_{n,2}}(\boldsymbol{x};q)$ is *e*-unimodal.

Much of the progress toward the Stanley-Stembridge conjecture considers a particular natural unit interval graph G and shows that every coefficient $c_{\mu}(q)$ of the *e*-expansion $X_G(\boldsymbol{x};q) = \sum_{\mu} c_{\mu}(q)e_{\mu}$ is a positive polynomial. We could also consider a dual approach where we fix a particular partition μ and show positivity for every natural unit interval graph G. Upcoming work of Sagan and the author [43] proves that $c_{\mu}(q)$ is positive whenever $\mu_1 \leq 3$. Hwang [37, Theorem 5.13] proved that $c_{\mu}(q)$ is positive whenever μ is a *hook*, meaning that $\mu_2 \leq 1$. In the q = 1 case, Abreu and Nigro [2, Corollary 1.10] proved that $c_{\mu}(1)$ is positive whenever $\mu = (n-k)k$ has two parts. We could try to strengthen this result by using forest triples to prove a quasisymmetric version.

Proposition 5.3. Let G = ([n], E) be a natural unit interval graph, let $s(G) = [b_2]_q \cdots [b_n]_q$, using the notation of Example 3.6, and for $1 \le k \le n-1$, define

(5.3)
$$s_k(G) = \sum_{\substack{T_1, T_2 \text{ dec. trees, } V(T_1) \sqcup V(T_2) = [n]\\ 1 \in V(T_1), \ |V(T_2)| = k}} q^{\text{inv}_G(\text{list}(T_1) \cdot \text{list}(T_2))}.$$

Then for $1 \leq k < \frac{n}{2}$, the coefficient of $e_{(n-k)k}$ in $X_G(\boldsymbol{x};q)$ is

(5.4)
$$c_{(n-k)k}(q) = [n-k]_q[k]_q s_k(G) + [n-k]_q[k]_q s_{n-k}(G) - ([n-k]_q + [k]_q)s(G)$$

(5.5)
$$= [n-k]_q([k]_q s_k(G) - s(G)) + [k]_q([n-k]_q s_{n-k}(G) - s(G)).$$

If $k = \frac{n}{2}$, then both terms in (5.5) are equal and the coefficient is given by just one of them.

FIGURE 12. The triangular ladder $P_{8,2}$



Proof. Forest triples $\mathcal{F} \in \mathrm{FT}_{(n-k)k}(G)$ come in one of two flavours. We could have a single tree triple $\mathcal{T} = (T, \alpha, r)$, where T is a decreasing spanning tree, contributing a factor of s(G) by Lemma 3.13, and α is either (n-k)k or k(n-k). Alternatively, \mathcal{F} can consist of two tree triples $\mathcal{T} = (T, \alpha, r)$ and $\mathcal{T}' = (T', \alpha', r')$ with trees of sizes (n-k) and k and compositions the single parts (n-k) and k. Now the result follows from Theorem 3.4.

Therefore positivity of $c_{(n-k)k}(q)$ would follow from the following. The author checked by computer that this holds for every natural unit interval graph with at most 10 vertices.

Problem 5.4. Let G = ([n], E) be a natural unit interval graph. Prove that for every $1 \le k \le n-1$, the polynomial $[k]_q s_k(G) - s(G)$ has positive coefficients.

Another property that is actively studied in relation to unimodality is *log-concavity* [52].

Definition 5.5. A polynomial $a(q) = \sum_{k=i}^{j} a_k q^k \in \mathbb{N}[q]$ is *log-concave* if for every k, we have $a_k^2 \ge a_{k-1}a_{k+1}$.

Note that if a(q) is a positive log-concave polynomial, then it must be unimodal because we cannot have $a_{k-1} > a_k$ and $a_k < a_{k+1}$. Therefore, log-concavity is a stronger property than unimodality. Thank you to Bruce Sagan for suggesting the following Conjecture. The author checked by computer that this holds for every natural unit interval graph with at most 10 vertices.

Conjecture 5.6. Let G be a natural unit interval graph. Then the chromatic quasisymmetric function $X_G(\mathbf{x};q)$ is *e-log-concave*, meaning that for every μ , the coefficient $c_{\mu}(q)$ of e_{μ} is a log-concave polynomial.

Recall that if G is not a natural unit interval graph, then $X_G(\mathbf{x}; q)$ is not generally symmetric in the x_i variables, so it may not have an *e*-expansion. However, when q = 1, the chromatic symmetric function $X_G(\mathbf{x}) = X_G(\mathbf{x}; 1)$ is always symmetric and we can use a variant of forest triples, with a slightly different condition on our trees, to calculate a signed *e*-expansion. We conclude by describing this variant. Let G = ([n], E) be an arbitrary graph with a fixed total ordering \leq of the edge set E.

Definition 5.7. A no-broken-circuit (NBC) tree of G is a subtree T of G such that E(T) does not contain a subset of the form $C \setminus \{\max(C)\}$, called a broken circuit, where C is a set of edges that forms a cycle and $\max(C)$ is the largest edge of C under \leq .

FIGURE 13. The claw graph G and the chromatic symmetric function $X_G(\mathbf{x})$



Remark 5.8. If G is a natural unit interval graph, we can define $\leq lexicographically$, meaning that for edges $\{i, j\}$ and $\{i', j'\}$ with i < j and i' < j', we define $\{i, j\} < \{i', j'\}$ if either i < i', or i = i' and j < j'. Then NBC trees are precisely decreasing trees. If $i \in V(T)$ has two larger neighbours j < k, then $\{j, k\} \in E(G)$ by (2.2) and E(T) contains the broken circuit $\{\{i, j\}, \{i, k\}, \{j, k\}\} \setminus \{\{j, k\}\}$. Conversely, if E(T) contains a broken circuit $C \setminus \{\max(C)\}$, then the smallest vertex in the edges of C has two larger neighbours in T.

Definition 5.9. A tree triple of G is an object $\mathcal{T} = (T, \alpha, r)$ consisting of the following data.

- \circ T is an NBC tree of G.
- $\alpha = \alpha_1 \cdots \alpha_\ell$ is an integer composition with size $|\alpha| = |V(T)|$.
- $\circ r$ is a positive integer with $1 \leq r \leq \alpha_1$.

A forest triple of G is a sequence of tree triples $\mathcal{F} = (\mathcal{T}_i = (T_i, \alpha^{(i)}, r_i))_{i=1}^m$ such that each vertex of G is in exactly one tree T_i and we have

(5.6)
$$\min(V(T_1)) < \min(V(T_2)) < \dots < \min(V(T_m)).$$

We also define the type and sign of \mathcal{F} , and the sets FT(G) and $FT_{\mu}(G)$, as in Definition 3.3.

Theorem 5.10. The chromatic symmetric function $X_G(\mathbf{x})$ of an arbitrary graph G satisfies

(5.7)
$$X_G(\boldsymbol{x}) = \sum_{\mathcal{F} \in \mathrm{FT}(G)} \mathrm{sign}(\mathcal{F}) e_{\mathrm{type}(\mathcal{F})} = \sum_{\mu} \left(\sum_{\mathcal{F} \in \mathrm{FT}_{\mu}(G)} \mathrm{sign}(\mathcal{F}) \right) e_{\mu}.$$

In particular, the *e*-expansion does not depend on the choice of total ordering \lt .

Example 5.11. Let us calculate the coefficient of e_{22} in $X_G(\boldsymbol{x})$ for the *claw graph* G in Figure 13. Because G has no cycles, every subtree is an NBC tree. Note that we cannot break G into two trees of size 2, because both trees would need to include vertex 3. Therefore, the only possibility is for \mathcal{F} to consist of a single tree triple $\mathcal{T} = (T, \alpha, r)$, where T is all of G, $\alpha = 22$, and $1 \leq r \leq 2$. We have $\operatorname{sign}(\mathcal{F}) = (-1)^{2-1} = -1$, so these forest triples give us the term $-2e_{22}$. Note that $X_G(\boldsymbol{x})$ is not *e*-positive.

To prove Theorem 5.10, we use a power sum expansion of $X_G(\boldsymbol{x})$ and then we perform a change-of-basis from power sum to elementary symmetric functions. The technique of changing bases is also employed in upcoming work of Sagan and the author [43]. Stanley originally proved the following result using Möbius inversion, and more recently a signreversing involution proof was given by Sagan and Vatter [44, Section 3].

Theorem 5.12. [49, Theorem 2.9] The chromatic symmetric function $X_G(\boldsymbol{x})$ is given by

(5.9)
$$X_G(\boldsymbol{x}) = \sum_{\substack{T_1, \dots, T_m \text{ NBC trees}\\V(T_1) \sqcup \cdots \sqcup V(T_m) = [n]}} (-1)^{n-m} p_{|V(T_1)|} \cdots p_{|V(T_m)|}.$$

Mendes and Remmel proved the following change-of-basis formula by showing that both sides satisfy the same recurrence.

Lemma 5.13. [41, Theorem 2.22] We have the identity

(5.10)
$$p_n = \sum_{\alpha \vDash n} (-1)^{n-\ell(\alpha)} \alpha_1 e_{\operatorname{sort}(\alpha)}.$$

Proof of Theorem 5.10. By Theorem 5.12 and Lemma 5.13, we have

$$(5.11) X_G(\boldsymbol{x}) = \sum_{\substack{T_1, \dots, T_m \text{ NBC trees}\\V(T_1) \sqcup \dots \sqcup V(T_m) = [n]}} (-1)^{n-m} p_{|V(T_1)|} \cdots p_{|V(T_m)|} \\ = \sum_{\substack{T_1, \dots, T_m \text{ NBC trees}\\V(T_1) \sqcup \dots \sqcup V(T_m) = [n]}} (-1)^{n-m} \prod_{i=1}^m \sum_{\alpha^{(i)} \models |V(T_i)|} (-1)^{|V(T_i)| - \ell(\alpha^{(i)})} \alpha_1^{(i)} e_{\operatorname{sort}(\alpha^{(i)})} \\ = \sum_{\substack{T_1, \dots, T_m \text{ NBC trees}\\V(T_1) \sqcup \dots \sqcup V(T_m) = [n]\\\alpha^{(1)} \models |V(T_1)|, \dots, \alpha^{(m)} \models |V(T_m)|} \\ 1 \le r_1 \le \alpha_1^{(1)}, \dots, 1 \le r_m \le \alpha_1^{(m)}} \\ = \sum_{\mathcal{F} \in \operatorname{FT}(G)} \operatorname{sign}(\mathcal{F}) e_{\operatorname{type}(\mathcal{F})}. \end{aligned}$$

Therefore, even for arbitrary graphs G, we can study *e*-positivity of the chromatic symmetric function $X_G(\boldsymbol{x})$ by fixing a total ordering \lt and seeking a sign-reversing involution on forest triples of G.

6. Acknowledgments

The author would like to thank Per Alexandersson, Alexander Postnikov, Bruce Sagan, Richard Stanley, Aarush Vailaya, and Stephanie van Willigenburg for helpful discussions.

References

- [1] A. Abreu and A. Nigro. "A symmetric function of increasing forests". In: Forum of Mathematics, Sigma 9 (2021).
- [2] A. Abreu and A. Nigro. "Splitting the cohomology of Hessenberg varieties and *e*-positivity of chromatic symmetric functions". In: (2023). arXiv: 2304.10644.
- [3] P. Alexandersson. "LLT polynomials, elementary symmetric functions and melting lollipops". In: J. Algebraic Combin. 53 (2021).
- [4] P. Alexandersson and G. Panova. "LLT polynomials, chromatic quasisymmetric functions and graphs with cycles". In: *Discrete Math.* 341 (2018).
- [5] P. Alexandersson and R. Sulzgruber. "A combinatorial expansion of vertical-strip LLT polynomials in the basis of elementary symmetric functions". In: *Adv. in Math.* 400 (2022).
- [6] F. Aliniaeifard, S. Li, and S. van Willigenburg. "Generalized chromatic functions". In: Int. Math. Res. Not, IMRN (2023).
- [7] F. Aliniaeifard, V. Wang, and S. van Willigenburg. "The chromatic symmetric function of a graph centred at a vertex". In: (2021). arXiv: 2108.04850.
- [8] J. Aliste-Prieto et al. "A vertex-weighted Tutte symmetric function, and constructing graphs with equal chromatic symmetric function". In: *Electron. J. of Combin.* 28.P2.1 (2021).
- [9] C. Athanasiadis. "Power sum expansion of chromatic quasisymmetric functions". In: Electron. J. of Combin. 22.P2.7 (2015).
- [10] J. Blasiak et al. "Noncommutative Schur functions for posets". In: (2022). arXiv: 2211.03967.
- [11] P. Brosnan and T. Chow. "Unit interval orders and the dot action on the cohomology of regular semisimple Hessenberg varieties". In: *Adv. in Math.* 329.10 (2015).
- [12] E. Carlsson and A. Mellit. "A proof of the shuffle conjecture". In: J. Amer. Math. Soc. 31 (2018).
- [13] S. Cho and J. Hong. "Positivity of chromatic symmetric functions associated with Hessenberg functions of bounce number 3". In: *Electron. J. Combin.* 29.2.19 (2022).
- [14] S. Cho, J. Hong, and E. Lee. "Bases of the equivariant cohomologies of regular semisimple Hessenberg varieties". In: Adv. Math. 423.109018 (2023).
- [15] S. Cho and J. Huh. "On *e*-positivity and *e*-unimodality of chromatic quasisymmetric functions". In: SIAM J. Discrete Math. 33.4 (2019).
- [16] S. Clearman et al. "Evaluation of Hecke algebra traces at Kazhdan–Lusztig basis elements". In: *Electron. J. of Combin.* 23 (2016).
- [17] L. Colmenarejo, A. Morales, and G. Panova. "Chromatic symmetric functions of Dyck paths and q-rook theory". In: *European J. Combin.* 107 (2023).
- [18] L Crew, O. Pechenik, and S. Spirkl. "The Kromatic Symmetric Function: A K-theoretic Analogue of X_G ". In: (2023). arXiv: 2301.02177.
- [19] L. Crew and S. Spirkl. "A Deletion-Contraction Relation for the Chromatic Symmetric Function". In: Eur. J. Comb. 89 (2020).

REFERENCES

- [20] L. Crew and Y. Zhang. "e-basis Coefficients of Chromatic Symmetric Functions". In: (2022). arXiv: 2210.03803.
- [21] M. D'Adderio, R. Riccardi, and V. Siconolfi. "Chromatic functions, interval orders and increasing forests". In: (2023). arXiv: 2311.09685.
- [22] S. Dahlberg. "Triangular ladders $P_{d,2}$ are *e*-positive". In: (2018). arXiv: 1811.04885.
- [23] S. Dahlberg, A. Foley, and S. van Willigenburg. "Resolving Stanley's *e*-positivity of claw-contractible-free graphs". In: *J. Eur. Math. Soc.* 22 (2020).
- [24] S. Dahlberg, A. She, and S. van Willigenburg. "Chromatic posets". In: J. Combin. Theory Ser. A 184 (2021).
- [25] S. Dahlberg, A. She, and S. van Willigenburg. "Schur and *e*-positivity of trees and cut vertices". In: *Electron. J. of Combin.* 27 (2020).
- [26] S. Dahlberg and S. van Willigenburg. "Chromatic symmetric functions in noncommuting variables revisited". In: *Adv. in Appl. Math.* 112 (2020).
- [27] S. Dahlberg and S. van Willigenburg. "Lollipop and lariat symmetric functions". In: SIAM J. Discrete Math. 32 (2018).
- B. Ellzey. "A directed graph generalization of chromatic quasisymmetric functions". In: (2017). arXiv: 1709.00454.
- [29] A. Foley, C. Hoang, and O. Merkel. "Classes of graphs with *e*-positive chromatic symmetric function". In: *Elec. J. of Combin.* 26.P3.51 (2019).
- [30] V. Gasharov. "Incomparability graphs of (3+1)-free posets are *s*-positive". In: *Discrete Math.* 157 (1996).
- [31] D. Gebhard and B. Sagan. "A chromatic symmetric function in noncommuting variables". In: J. Algebr. Comb. 13 (2001).
- [32] M. Guay-Paquet. "A modular law for the chromatic symmetric functions of (3+1)-free posets". In: (2013). arXiv: 1306.2400.
- [33] M. Guay-Paquet. "A second proof of the Shareshian–Wachs conjecture, by way of a new Hopf algebra". In: (2016). arXiv: 1601.05498.
- [34] J. Haglund. The q,t-Catalan numbers and the space of diagonal harmonics. Amer. Math. Soc., 2007.
- [35] M. Harada and M. Precup. "The cohomology of abelian Hessenberg varieties and the Stanley–Stembridge conjecture". In: *Algebr. Comb.* 2 (2019).
- [36] J. Huh, S. Nam, and M. Yoo. "Melting lollipop chromatic quasisymmetric functions and Schur expansion of unicellular LLT polynomials". In: *Discrete Math.* 343 (2020).
- [37] B.-H. Hwang. "Chromatic quasisymmetric functions and noncommutative *P*-symmetric functions". In: (2022). arXiv: 2208.09857.
- [38] A. Lascoux, B. Leclerc, and J-Y. Thibon. "Ribbon tableaux, Hall–Littlewood functions, quantum affine algebras, and unipotent varieties". In: J. Math. Phys. 38.2 (1997).
- [39] G. Li and A. Yang. "On the *e*-Positivity of $(claw, 2K_2)$ -Free Graphs". In: *Electron. J. Combin.* 28.P2.40 (2021).
- [40] J. McDonough, P. Pylyavskyy, and S. Wang. "The Stanley–Stembridge Conjecture for (2+1+1)-avoiding unit interval orders". In: (2024). arXiv: 2404.07280.

REFERENCES

- [41] A. Mendes and J. Remmel. *Counting with symmetric functions*. Vol. 43. Developments in Mathematics, 2015.
- [42] A. Rok and A. Szenes. "Eschers and Stanley's chromatic e-positivity conjecture in length-2". In: (2023). arXiv: 2305.00963.
- [43] B. Sagan and F. Tom. "Chromatic symmetric functions and change of basis". In: In preparation (2024).
- [44] B. Sagan and V. Vatter. "Bijective proofs of proper coloring theorems". In: The American Mathematical Monthly 128 (2021).
- [45] J. Shareshian and M. Wachs. "Chromatic quasisymmetric functions". In: Adv. Math. (2016).
- [46] J. Shareshian and M. Wachs. "Chromatic quasisymmetric functions and Hessenberg varieties". In: *Configuration Spaces* (2012).
- [47] I. Siegl. "Cylindric P-Tableaux for (3+1)-Free Posets". In: (2022). arXiv: 2211.03953.
- [48] M. Skandera. "Characters and chromatic symmetric functions". In: Electron. J. Combin. 28.P2.19 (2021).
- [49] R. Stanley. "A symmetric function generalization of the chromatic polynomial". In: Adv. Math. 111 (1995).
- [50] R. Stanley. *Enumerative Combinatorics. Vol. 2.* Cambridge University Press, 1999.
- [51] R. Stanley. "Graph colorings and related symmetric functions: ideas and applications". In: Discrete Math. 193.1 (1998).
- [52] R. Stanley. "Log-concave and unimodal sequences in Algebra, Combinatorics, and Geometry". In: Annals of the New York Academy of Sciences 576 (1989).
- [53] R. Stanley and J. Stembridge. "On immanants of Jacobi–Trudi matrices and permutations with restricted position". In: J. Combin. Theory Ser. A 62 (1993).
- [54] D. Wang. "All cycle-chords are *e*-positive". In: (2024). arXiv: 2405.01166.
- [55] D. Wang and M. Wang. "The *e*-positivity and Schur positivity of the chromatic symmetric functions of some trees". In: (2021). arXiv: 2112.06619.
- [56] S. Wang. "The *e*-positivity of the chromatic symmetric functions and the inverse Kostka matrix". In: (2022). arXiv: 2210.07567.